Randomized lasso associates freshwater lake-system specific bacterial taxa with

heterotrophic production through flow cytometry

Running Title: Linking taxa with production through flow cytometry

Peter Rubbens[1]*#, Marian L. Schmidt[23]*#, Ruben Props[4], Bopaiah A. Biddanda[5], Nico Boon[4],

Willem Waegeman[1], Vincent J. Denef[2]

*Peter Rubbens and Marian L. Schmidt contributed equally to this work.

#Corresponding author: marschmi@umich.edu, peter.rubbens@ugent.be

[1]KERMIT, Department of Data Analysis and Mathematical Modelling, Ghent University,

Coupure Links 653, B-9000, Gent, Belgium;

[2]Department of Ecology and Evolutionary Biology, University of Michigan, 1105 North

University Ave., Ann Arbor, MI 48109, USA;

[3]Current address: Department of Integrative Biology, University of Texas at Austin, 2506

Speedway, Austin, Texas 78712, USA;

[4]CMET, Center for Microbial Ecology and Technology, Ghent University, Coupure Links 653,

B-9000, Gent, Belgium;

[5]Annis Water Resources Institute, Grand Valley State University, 740 West Shoreline Drive,

Muskegon, MI 49441, USA

Abstract

High- (HNA) and low-nucleic acid (LNA) bacteria are two operational groups identified by flow cytometry (FCM) in aquatic systems. HNA cell density often correlates strongly with heterotrophic production, while LNA cell density does not. However, which taxa are specifically associated with these groups, and by extension, productivity has remained elusive. Here, we addressed this knowledge gap by using a machine learning-based variable selection approach that integrated FCM and 16S rRNA gene sequencing data collected from 14 freshwater lakes spanning a broad range in physicochemical conditions. There was a strong association between bacterial heterotrophic production and HNA absolute cell abundances ($R^2 = 0.65$), but not with the more abundant LNA cells. This solidifies findings, mainly from marine systems, that HNA and LNA could be considered separate functional groups, the former contributing a disproportionately large share of carbon cycling. Taxa selected by the models could predict HNA and LNA absolute cell abundances at all taxonomic levels, with the highest performance at the OTU level. Selected OTUs ranged from low to high relative abundance and were mostly lake system-specific (89.5%-99.2%). A subset of selected OTUs was associated with both LNA and HNA groups (12.5%-33.3%) suggesting either phenotypic plasticity or within-OTU genetic and physiological heterogeneity. These findings may lead to the identification of systems-specific putative ecological indicators for heterotrophic productivity. Generally, our approach allows for the association of OTUs with specific functional groups in diverse ecosystems in order to improve our understanding of (microbial) biodiversity-ecosystem functioning relationships.

Importance

A major goal in microbial ecology is to understand how microbial community structure

influences ecosystem functioning. Research is limited by the ability to readily culture most

bacteria present in the environment and the difference in bacterial physiology *in situ* compared to

in laboratory culture. Various methods to directly associate bacterial taxa to functional groups in

the environment are being developed. In this study, we applied machine learning methods to

relate taxonomic data obtained from marker gene surveys to functional groups identified by flow

cytometry. This allowed us to identify the taxa that are associated with heterotrophic productivity

in freshwater lakes and indicated that the key contributors were highly system-specific, regularly

rare members of the community, and that some could switch between being low and high

contributors. Our approach provides a promising framework to identify taxa that contribute to

ecosystem functioning and can be further developed to explore microbial contributions beyond

heterotrophic production.

Keywords

bacterioplankton, 16S rRNA, flow cytometry, machine learning, variable selection, aquatic

microbiology, heterotrophic productivity

Introduction

A key goal in the field of microbial ecology is to understand the relationship between microbial diversity and ecosystem functioning. However, it is challenging to associate bacterial taxa to specific ecosystem processes. Marker gene surveys have shown that natural bacterial communities are extremely diverse and the presence of a taxon does not imply its activity. The taxa observed in these surveys may have low metabolic potential, be dormant, or have recently died (1, 2). An additional hurdle is that the current standard unit of measure for microbial taxonomic analysis is relative abundance. This results in a negative correlation bias (3), which makes it difficult to quantitatively associate specific microbial taxa with microbial ecosystem functions using traditional correlation measures (4). Therefore, in order to ultimately model and predict bacterial communities, new methodologies, which integrate different data types, are needed to associate bacterial taxa with ecosystem functions (5).

One such advance is the use of flow cytometry (FCM), which has been used extensively to study aquatic microbial communities (6–8). This single-cell technology partitions individual microbial cells into phenotypic groups based on their observable optical characteristics. Most commonly, cells are stained with a nucleic acid stain (*e.g.* SYBR Green I) and upon analysis assigned to either a low nucleic acid (LNA) or a high nucleic acid (HNA) group (9–12). HNA cells differ from LNA cells in both a considerable increase in fluorescence due to cellular nucleic acid content and scatter intensity due to cell morphology. The HNA group is thought to contribute more, whereas the LNA population has been considered to contribute less to productivity of a microbial community (6, 13–15). This is based on positive linear relationships between HNA abundance and (a) bacterial heterotrophic production (BP) (10, 14–17), (b) bacterial activity

measured using the dye 5-cyano-2,3-ditolyl tetrazolium chloride (18, 19), (c) phytoplankton

abundance (20), and (d) dissolved organic carbon concentrations (21). Additionally, growth rates

are higher for HNA than LNA cells (13, 16, 22) and HNA cells accrue cell damage significantly

faster than the LNA cells under temperature (23) and chemical oxidant stress (24). In contrast,

LNA bacterial growth rates are positively correlated with temperature and negatively correlated

with chlorophyll a (25). However, it is important to note that LNA cells are often smaller than

HNA cells (12, 25–27) and therefore LNA cells could have similar amino acid incorporation

rates compared to HNA cells when evaluating biomass-specific production (12).

Here we used a data-driven approach to associate the dynamics of individual taxa with those of

the LNA and HNA groups in freshwater lakes by adopting a machine learning variable selection

strategy. We applied two variable selection methods, the Randomized Lasso (RL) (28) and the

Boruta algorithm (29) to associate individual taxa with HNA and LNA cell abundances. These

methods extend on traditional machine learning algorithms (*i.e.* the Lasso and Random forest

algorithm for the RL and Boruta algorithm, respectively) by making use of resampling and

randomization. These extensions are needed as (a) the Lasso algorithm is not suited for

compositional data because the regression coefficients have an unclear interpretation, and single

variables may be selected when correlated to other variables (30), and (b) Random Forest

algorithms can be biased towards correlated variables (31), which is an intrinsic issue with

relative abundance data (3). The extended methods allow the user to either assign a probability of

selection (RL) or statistically decide which taxa to select (Boruta).

We gathered samples from three types of lake systems (i) a set of oligo- to eutrophic small inland

lakes, (ii) a short residence time mesotrophic freshwater estuary lake (Muskegon Lake), and (iii)

a large oligotrophic Great Lake (Lake Michigan), all located in Michigan, USA. We then used

the RL and Boruta algorithms to associate specific bacterial taxa to HNA and LNA FCM

functional groups, and via the observed HNA-productivity relationship, to functioning. To

validate the RL-based association with the HNA and/or LNA group, we correlated taxon

abundances with specific regions within the FCM fingerprint at finer resolution (*i.e.* bins)

without prior knowledge of the HNA/LNA groups. Furthermore, we tested for phylogenetic

conservation of HNA and LNA functional groups using the probabilities from the RL output and

for the association between the selected taxa and productivity.

Results

Study lakes are dominated by LNA cells

The inland lakes (6.3 x 10^6 cells/mL) and Muskegon Lake (6.0 x 10^6 cell/mL) had significantly

higher total cell abundances than Lake Michigan (1.7 x 10^6 cell/mL; p = 2.7 x 10^{-14}). Across all

lakes, the mean proportion of HNA cell counts (HNAcc) to total cell counts was much lower

(30.4 ± 9%) compared to the mean proportion of LNA cell counts (LNAcc; 69.6 ± 9%). Through

ordinary least squares regression, there was a strong correlation between HNAcc and LNAcc

across all data (R^2 = 0.45, P = 2 x 10^{-24}; **Figure 1A**), however, only Lake Michigan (R^2 = 0.59, P

= 5 x 10^{-11}) and Muskegon Lake (R^2 = 0.44, P= 2 x 10^{-9}) had significant correlations when the

three ecosystems were considered separately.

HNA cell counts and heterotrophic bacterial production are strongly correlated

For mesotrophic Muskegon Lake, there was a strong correlation between total bacterial

heterotrophic production and HNAcc (R^2 = 0.65, P = 1e-05; **Figure 1B**), no correlation between

BP and LNAcc (R^2 = 0.005, P = 0.31; **Figure 1C**), and a weak correlation between heterotrophic

production and total cell counts (R^2 = 0.18, P = 0.03; **Figure 1D**). There was a positive (HNA)

and negative (LNA) correlation between the fraction of HNA or LNA to total cells and

productivity, however, the relationship was weak and not significant (R^2 = 0.14, P = 0.057).

Association of OTUs to HNA and LNA groups by Randomized Lasso

The relevance of specific OTUs for predicting FCM functional group abundance was assessed

using the Randomized Lasso (RL), which assigns a score between 0 (unimportant) to 1 (highly

important) to each taxon in function of the target variable: HNAcc or LNAcc. To assess the

predictive power of a subset of OTUs based on the RL, we iteratively removed the OTUs with

the lowest RL score in a recursive variable elimination scheme. R^2_{CV}, a goodness-of-fit measure

using the R^2 of how well a set of selected OTUs predicts HNAcc or LNAcc compared to true

values using cross-validation, increased when lower-ranked OTUs were removed (moving from

right to left on **Figure 2**). The increase was gradual for the inland lakes (**Figure 2A**) and

Muskegon Lake (**Figure 2C**) but was abrupt for Lake Michigan (**Figure 2B**). The proportion of

taxa that resulted in the highest R^2_{CV} (*see solid (HNA) and dotted (LNA) lines in* **Figure 2**) was

10.2% of all taxa for HNA and 17.7% for LNA for the inland lakes, 4.0% for HNA and 3.0% for

LNA for Lake Michigan, and 21.1% for both HNA and LNA in Muskegon Lake. Lake Michigan

differed the most from other lake systems, having the lowest R^2_{CV}, a sharp increase in R^2_{CV} as

OTUs were eliminated, and a considerably lower number of OTUs that were retained (13 for

HNAcc, 10 for LNAcc). No relationship could be established between rankings of variable

selection methods and the relative abundance of individual OTUs (**Figure S1**). HNAcc and

LNAcc could be predicted with equivalent performance to relative HNA and LNA proportions,

yet the increase between initial and optimal performance was larger (**Figure S2**). The final

predictive performance was higher when relative OTU abundances were transformed using the

CLR-transformation (**Figure S3**).

OTU-level predictions outperform other taxonomic levels

R^2_{CV} values were considerably higher than zero on all taxonomic levels, indicating that our results

were consistent across all taxonomic levels and that different levels can be related to changes in

HNAcc and LNAcc. While the OTU level resulted in the best prediction of HNAcc and LNAcc

(**Figure 3**), each individual OTU had a lower RL score compared to other taxonomic levels,

which on average became lower as the taxonomic level decreased (**Figure S4**). The fraction of

variables (*i.e.* taxa) that could be removed to reach the maximum R^2_{CV} decreased as the

taxonomic level became less resolved.

Validation of RL OTU selection results using the Boruta algorithm and Kendall tau statistic

Venn diagrams were constructed to visualize consistency in the number of OTUs that were

selected according to the RL method, the Boruta algorithm, and individual correlations with

HNAcc and LNAcc via the Kendall rank correlation coefficient (**Figure S5**). The Kendall rank

correlation coefficient selected the most OTUs, followed by the RL, and then the Boruta

algorithm (except for HNAcc in Lake Muskegon; **Figure S5**). The Boruta algorithm selects

relevant variables based on the importance of the most permuted variable as retrieved from

multiple Random Forest models (*see materials and methods*). The Boruta algorithm ranks

selected OTUs as '1', tentative OTUs as '2', and all other OTUs have lower ranks, depending on

the stage in which they were eliminated. The fraction of selected OTUs was always smaller than

1% across lake systems and functional groups (**Figure S6**). All methods agreed on only a small

subset of OTUs.

For each lake system individually, the top RL-scored OTU for HNAcc was also selected by the

Boruta algorithm, whereas both methods only agreed for Lake Michigan LNAcc (**Table 1**).

Across all lake systems, OTU060 (Proteobacteria;Sphingomonadales;alfIV_unclassified) was the

only OTU selected across all lake systems (LNAcc-associated). As Random Forest regressions

are the base method of the Boruta algorithm, we compared the predictive power of Boruta

selected OTUs to those of all OTUs using Random Forest regression. For all lake systems and

functional groups, the performance increased when only Boruta-selected OTUs were included in

the model (**Table S1**). Lasso predictions, in which OTUs were selected according to the RL,

were better as opposed to Random Forest predictions in which OTUs were selected according to

the Boruta algorithm (**Figure S7**).

Although all methods only agreed on a minority of OTUs, we can still formulate a number of

general conclusions across these methods: (1) the selected OTUs were mostly lake systems

specific, (2) a small fraction of OTUs was needed to predict changes in community composition,

(3) selected OTUs were associated with absolute HNA or LNA abundance, (4) top RL-ranked

HNA-associated OTUs were also selected according to the Boruta algorithm and (5) when the

RL and Boruta both agreed on an OTU it was always significantly correlated with both HNAcc

or LNAcc.

HNA- and LNA-associated OTUs differed across lake systems

RL-selected OTUs were mostly assigned to either the HNA or LNA groups and there was

limited correspondence across lake systems between the selected OTUs (**Figure 4**). 1.5%-1.9%

of the OTUs selected for Lake Michigan were also associated with HNAcc or LNAcc for the

inland lakes or Muskegon Lake. This amount was higher for the shared OTUs between the inland

lakes and Lake Muskegon, but still only amounted to 6.0% (HNAcc) or 10.5% (LNAcc) of all

common OTUs. For OTUs selected in all three freshwater environments, RL scores were lake

ecosystem specific, with only a significant similarity between the Inland lakes and Muskegon

lake for HNAcc (r = 0.21, P = 0.0042; **Figure S8**). The Boruta algorithm selected mostly OTUs

that were unique both for the lake system and FCM group (**Figure S9**).

The Bacteroidetes, Betaproteobacteria, Alphaproteobacteria, and Verrucomicrobia contributed

54% of the 258 OTUs selected by the RL (**Figure 5**). Most selected OTUs belonging to these

four phyla were associated with the LNA group (41-52% of selected OTUs), less than one third

with the HNA group (14-30% of selected OTUs), and the remainder were selected as associated

with both the LNA and HNA groups (23-36% of selected OTUs). In Muskegon Lake, OTU173

(Bacteroidetes;Flavobacteriales;bacII-A) was selected as the major HNA-associated taxon while

OTU29 (Bacteroidetes;Cytophagales;bacIII-B) had the highest RL score for LNA OTUs. In Lake

Michigan, OTU25 (Bacteroidetes*;*Cytophagales;bacIII-A), was selected as the major HNA-

associated taxon while OTU168 (Alphaproteobacteria:Rhizobiales:alfVII) was selected as a

major LNA-associated taxon. For the inland lakes, OTU369

(Alphaproteobacterial;Rhodospirillales;alfVIII) was the major HNA-associated OTU while the

OTU555 (Deltaproteobacteria;Bdellovibrionaceae;OM27) was the major LNA-associated taxon.

Most OTUs were selected for Muskegon Lake (153 OTUs; compared to 136 OTUs from the

Inland Lakes and 20 OTUs from Lake Michigan) and 33% of these OTUs were associated with

both FCM groups.

Association with HNA and LNA is not phylogenetically conserved

To evaluate how much evolutionary history explains whether a selected taxon was associated

with the HNA and/or LNA group(s), we calculated Pagel's λ, Blomberg's K, and Moran's I,

which are different measures for testing whether there was a phylogenetic conservation of these

traits. No phylogenetic signal was detected when using Pagel's λ with either using FCM

functional group as a discrete variable (*i.e.* associating an OTU with HNA, LNA, or Both;

Figure 5) or in relation to the HNA RL score, which is a continuous variable (lambda = 0.16; P

= 1; **Figure S10**). However, there was a significant phylogenetic signal for the LNA RL score (p

= 0.003, λ = 0.66), suggesting a stronger phylogenetic structure in the LNA group compared to

the HNA group. This significant result in the LNA group was not found when other measures of

phylogenetic signal were considered (Blomberg's K (HNA: p = 0.63; LNA: p = 0.54), and

Moran's I (HNA: p = 0.88; LNA: p = 0.12)).

Flow cytometry fingerprints confirm associated taxa and reveal more complex relationships

between taxonomy and flow cytometric features

To confirm the association of the final selected OTUs with the HNA and LNA groups, and

resolve how HNA and LNA groups correspond to OTU-level clustering of cells on the FCM

fingerprints, we calculated the correlation between the density of individual small regions (*i.e.*

"bins") in the flow cytometry data with the relative abundances of the OTUs. Note that (i) as

these values denote correlations, they do not indicate actual presence, and (ii) the threshold that

was used to manually make the distinction between HNAcc and LNAcc (*i.e.* dotted line in

Figure 6) lies very close to the border between the two regions of positive and negative

correlation. OTU25 correlated with bins that when aggregated corresponded to almost the entire

HNA region, whereas OTU173 was limited to bins corresponding to the bottom of the HNA

region (**Figure 6**). In contrast, OTU369 was positively correlated to bins situated in both the

LNA and HNA regions of the cytometric fingerprint, highlighting results from **Figure 4** where

OTU369 was selected for both HNA and LNA.

Proteobacteria and rare taxa correlate with productivity measurements

The Kendall rank correlation coefficient was calculated between CLR-transformed abundances of individual OTUs and productivity measurements. OTU481 was the sole OTU that correlated with productivity after a multiple testing correction (Kendall's tau-b = -0.67, P = 0.00003, P_adj = 0.016), but had a low RL score (0.022) for HNAcc and was not selected according to the Boruta algorithm. Of the top 10 OTUs selected for HNAcc according to the RL, three were still significantly associated with productivity (OTU614: P = 0.0064; OTU412, P = 0.044; OTU487, P = 0.014), but not when corrected for multiple hypothesis testing. Some OTUs that had a high RL score also had a positive response to productivity measurements, though they were insignificant after multiple testing correction (**Figure S11**). At the phylum level, only Proteobacteria were significantly correlated to productivity measurements (Kendall's tau = 0.49, P = 0.002, P_adj = 0.05).

Discussion

Our study furthers the integration of functional and genotypic information to determine the complex relationships between microbial diversity and ecosystem functioning. Our results confirmed previous findings that flow cytometric operational groups are distinct functional groups having divergent correlations with heterotrophic productivity. Using two machine learning based variable selection strategies, we could associate bacterial taxa identified by 16S rRNA gene sequencing to these two functional groups in three types of freshwater lake systems in the Great Lakes region. We revealed that (i) HNA and LNA cell abundances were best predicted by a small subset of OTUs that were unique to each lake type, (ii) some OTUs were included in the best model for both HNA and LNA abundance, (iii) there was no phylogenetic conservation of HNA and LNA group association and (iv) freshwater FCM fingerprints display

more complex patterns related to OTUs and productivity compared to the traditional dichotomy

of HNA and LNA.

Although high-nucleic acid cell counts (HNAcc) and low-nucleic acid cell counts (LNAcc) were

correlated with each other, only the association between bacterial heterotrophic production (BP)

and HNAcc was strong and significant. This is in line with previous reports, though past studies

have focused on the proportion of HNA rather than absolute cell abundances and are strongly

biased towards marine systems. For example, Bouvier et al. (11) found a correlation between the

fraction of HNA cells and BP within a large dataset of 640 samples across various freshwater to

marine environments (Pearson's r $= 0.49$), whereas a study off the coast of the Antarctic

Peninsula found a moderate correlation ($R^2 = 0.36$; (17)). Another study in the Bay of Biscay

also found this association ($R^2 = 0.16$; (15)), however, the authors attributed this difference to be

related to cell size and not due to the activity of HNA. Notably, these studies were predominantly

testing the association of marine HNA groups. The high correlation coefficients observed in our

study may indicate a strong coupling between freshwater carbon cycling and HNA group

abundance in freshwater lake systems. Consequently, this suggests an important role of HNA

bacteria in the disproportionately large role that freshwater systems in the global carbon cycle

(32). It has to be noted that our study only evaluated bacterial heterotrophic production using

leucine amino acid incorporation, which biases our analyses against bacterial groups that cannot

import or assimilate this compound (33). Finally, as our correlations with proportional HNA

group abundances also indicated less strong correlations than with absolute HNAcc, we suggest

absolute HNAcc should be used to best predict and study heterotrophic bacterial production.

Similar to other microbiome studies that use machine learning, only a minority of OTUs were

needed to predict the phenotype of interest, with low predictive power of each single OTU, but

strong predictive capacity of the selected group of OTUs (17, 34–36). Both the RL and Boruta

algorithm have been applied to microbiome studies before, for example in the selection of genera

in the human microbiome associated with BMI (37), salivary pH and lysozyme activity (38), and

in relation to multiple sclerosis (39) or with differing diets during primate pregnancy (40). The

Boruta algorithm has also recently been proposed as one of the top-performing variable selection

methods that make use of Random Forests (41). Despite the power of these approaches,

improvements can be made when attempting to integrate different types of data. For example,

16S rRNA gene sequencing still faces the hurdles of DNA extraction (42) and 16S copy number

bias (43). Moreover, detection limits are different for FCM (expressed in the number of cells)

and 16S rRNA gene sequencing (expressed in the number of gene counts or relative abundance),

therefore creating data that may be different in resolution.

The selection of different sets of HNA and LNA OTUs across the three freshwater systems

indicates that different taxa underlie the universally observed HNA and LNA functional groups

across aquatic systems. This is perhaps not surprising as it has been shown that there is strong

species sorting in lake systems (44, 45), shaping community composition through diverging

environmental conditions between the lake systems presented here (46). This high system

specificity also explains the low RL scores for individual OTUs, as the spatial and temporal

dynamics of an OTU diverged strongly across systems. For example, an OTU that has an RL

score of 0.5 implies that on average it will only be chosen one out of two times in a Lasso model.

Some OTUs were associated with both HNAcc and LNAcc. There are multiple possible

explanations for this: (a) In line with scenario 1 from Bouvier et al (11), cells transition from

active growth (primarily HNA) to death or a dormant state (primarily LNA), depending on

variable conditions over the spatiotemporal gradients sampled in this study. A large fraction of

cells (40-95%) in aquatic systems has indeed been inferred to be dormant (47–49), in line with

the predominance of LNA cells. (b) The same OTU may occur in both HNA and LNA groups

due to phenotypic plasticity, which is more in line with scenario 4 from Bouvier et al (11).

Bacterial phenotypic plasticity in size and morphology has been observed (50), and agrees with

suggestions that HNA and LNA groups correspond to cells of differing size (12, 15, 27). (c) The

association of taxa to LNA and HNA can also mean that these taxonomic groups thrive within

either high or low productivity ecosystems and not necessarily that they are responsible for the

change in productivity. (d) Finally, OTU level grouping of bacterial taxa can disguise genomic

and corresponding phenotypic heterogeneity (51–54), which may be an alternate explanation for

inconsistent associations between OTUs and FCM functional groups.

We found no clear phylogenetic conservation of association to HNAcc or LNAcc. This is in

contrast to a recent study that found a clear signal at the phylum level across different aquatic

systems (27). However, lake water samples were an exception to the general trend. In addition, it

is notable that Proctor et al. (27) separated HNA and LNA cells based on cell size (where HNA

cells were defined at approximately >0.4 **μ**m and LNA cells were approximately 0.2-0.4 **μ**m,

based on 50-90% removal of HNA cells after filtering using a 0.4 **μ**m filter), while our study

separated these FCM functional groups on the basis of fluorescence intensity alone. A more

direct estimation of phylogenetic conservation that directly combines cell sorting of HNA or

LNA cells and sequencing, such as the approach of Vila-Costa et al. (55), will be needed to

resolve these contrasting results. Considering the correlations between FCM-based phenotypic

diversity and sequencing-based taxonomic diversity (56, 57), there clearly is a link between

taxonomy and the structure in microbial cytometry data (17). However, the HNA/LNA

dichotomy is too unresolved, as our correlation analysis between smaller regions in the

cytometric fingerprint and the highly-ranked OTUs revealed a more complex relationship. This

agrees with recent research, in which more than two FCM operational groups in aquatic systems

were identified (17, 58, 59).

The Boruta algorithm and RL scores agreed on a small subset of OTUs, including the top-ranked

HNA OTU for all lake systems according to RL, which motivates further investigation of the

ecology of these OTUs. While little detailed information on the identities and ecology of HNA

and LNA freshwater lake bacterial taxa exists, several studies identified Bacteroidetes among the

most prominent HNA taxa, which is in line with our findings. Independent research by Vila-

Costa et al. (55) found that the HNA group was dominated by Bacteroidetes in summer samples

from the Mediterranean Sea, Read et al. (19) showed that HNA abundances correlated with

Bacteroidetes, and Schattenhofer et al. (60) reported that the Bacteroidetes accounted for the

majority of HNA cells in the North Atlantic Ocean. In Muskegon Lake, OTU173 was the

dominant HNA taxon and is a member of the Order *Flavobacteriales* (bacII-A). The bacII group

is a very abundant freshwater bacterial group and has been associated with senescence and

decline of an intense algal bloom (61), suggesting their potential for bacterial production. BacII-

A has also made up ~10% of the total microbial community during cyanobacterial blooms,

reaching its maximum density immediately following the bloom (62). In Lake Michigan,

OTU25, a member of the Bacteroidetes Order *Cytophagales* known as bacIII-A, was the top

HNA OTU. However, much less is known about this specific group of Bacteroidetes. Though,

the bacII-A/bacIII-A group has been strongly associated with more heterotrophically productive

headwater sites (compared to higher order streams) from the River Thames, showing a negative

correlation in rivers with dendritic distance from the headwaters, indicating that these taxa may

contribute more to productivity (19). In the inland lakes, OTU369 was the major HNA taxon and

is associated with the Alphaproteobacteria Order Rhodospirillales (alfVIII), which to our

knowledge is a group with very little information available in the literature. In contrast to our

findings of Bacteroidetes and Alphaproteobacterial HNA selected OTUs, Tada & Suzuki (63)

found that the major HNA taxon from an oceanic algal culture was from the Betaproteobacteria

whereas LNA OTUs were within the Actinobacteria phylum.

Conclusions

We integrated flow cytometry (FCM) and 16S rRNA gene amplicon sequencing data to associate

bacterial taxa with productivity in freshwater lake systems. Our results on a diverse set of

freshwater lake systems indicate that the taxa associated with HNA and LNA functional groups

are lake-specific, and that association with these functional groups is not phylogenetically

conserved. With this study, we show the potential and limitations of integrating flow cytometry-

derived *in situ* functional information with sequencing data using machine learning approaches.

This integration of data enhances our insights into which taxa may contribute to ecosystem

functioning in aquatic bacterial communities. While these data-driven hypotheses will need

further verification, the method is promising considering the wide application of FCM in aquatic

environments, its recent application in other sample matrices (*e.g.*, faeces (64), soils (65), and

wastewater sludge (66)), and the introduction of novel stains to delineate operational groups

based on phenotypic traits (67).

Materials and Methods

Data collection and DNA extraction, sequencing and processing

In this study, we used a total of 173 samples collected from three types of lake systems described

previously (46), including: (a) 49 samples from Lake Michigan (2013 & 2015), (b) 62 samples

from Muskegon Lake (2013-2015; one of Lake Michigan's estuaries), and (c) 62 samples from

twelve inland lakes in Southeastern Michigan (2014-2015). For more details on sampling, please

see **Figure 1** and the *Field Sampling, DNA extraction, and DNA sequencing and processing*

sections within Chiang et al. (46). In all cases, water for microbial biomass samples were

collected and poured through a 210 μm and 20 μm bleach sterilized nitex mesh and sequential in-

line filtration was performed using 47 mm polycarbonate in-line filter holders (Pall Corporation,

Ann Arbor, MI, USA) and an E/S portable peristaltic pump with an easy-load L/S pump head

(Masterflex®, Cole Parmer Instrument Company, Vernon Hills, IL, USA) to filter first through a

3 μm isopore polycarbonate (TSTP, 47 mm diameter, Millipore, Billerica, MA, USA) and

second through a 0.22 μm Express Plus polyethersulfone membrane filters (47 mm diameter,

Millipore, MA, USA). The current study only utilized the 3 - 0.22 μm fraction for analyses.

DNA extractions and sequencing were performed as described in Chiang et al. (46). Fastq files

were submitted to NCBI sequence read archive under BioProject accession number

PRJNA414423 (inland lakes), PRJNA412983 (Lake Michigan), and PRJNA412984 (Muskegon

Lake). We analyzed the sequence data using MOTHUR V.1.38.0 (seed = 777; (Schloss et al.

2009) based on the MiSeq standard operating procedure and put together at the following link:

https://github.com/rprops/Mothur_oligo_batch. A combination of the Silva Database (release

123; (68)) and the freshwater TaxAss 16S rRNA database and pipeline (69) was used for

classification of operational taxonomic units (OTUs).

For the taxonomic analysis, each of the three lake datasets were analyzed separately and treated

with an OTU abundance threshold cutoff of at least 5 sequences in 10% of the samples in the

dataset (similar strategy to (70)). For comparison of taxonomic abundances across samples, each

of the three datasets were then rarefied to an even sequencing depth, which was 4,491 sequences

for Muskegon Lake samples, 5,724 sequences for the Lake Michigan samples, and 9,037

sequences for the inland lake samples. Next, the relative abundance at the OTU level was

calculated using the *transform_sample_counts()* function in the phyloseq R package (71) by

taking the count value and dividing it by the sequencing depth of the sample. For all other

taxonomic levels, the taxonomy was merged at certain taxonomic ranks using the *tax_glom()*

function in phyloseq (71) and the relative abundance was re-calculated.

Heterotrophic bacterial production measurements

Muskegon Lake samples from 2014 and 2015 were processed for heterotrophic bacterial

production using the [^{3}H] leucine incorporation into bacterial protein in the dark method (72, 73).

At the end of the incubation with [^{3}H]-leucine, cold trichloroacetic acid-extracted samples were

filtered onto 0.2 µm filters that represented the leucine incorporation by the bacterial community.

Measured leucine incorporation during the incubation was converted to bacterial carbon

production rate using a standard theoretical conversion factor of 2.3 kg C per mole of leucine

(73).

Flow cytometry, measuring HNA and LNA

In the field, a total of 1 mL of 20 μm filtered lake water were fixed with 5 μL of glutaraldehyde

(20% vol/vol stock), incubated for 10 minutes on the bench (covered with aluminum foil to

protect from light degradation), and then flash frozen in liquid nitrogen to later be stored in -

80°C freezer until later processing with a flow cytometer. Flow cytometry procedures followed

the protocol laid out in Props et al. (56), which also uses the samples presented in the current

study (Michigan and Muskegon samples). Samples were stained with SYBR Green I and

measured in triplicate. The lowest number of cells collected after denoising was 2342. HNA and

LNA groups were selected using the fixed gates introduced in Prest et al. (74) and plotted in

Figure S11. Cell counts were determined per HNA and LNA group and averaged over the three

replicates (giving rise to HNAcc and LNAcc). All cytometry data is available on the

FlowRepository database (75): inland lakes (ID:FR-FCM-ZY9J), Michigan and Muskegon

(ID:FR-FCM-ZYZN).

Data analysis

Processed data and analysis code for the following analyses can be found on the GitHub page for

this project at https://deneflab.github.io/HNA_LNA_productivity/.

HNA-LNA and HNA-Productivity Statistics and Regressions

We tested the difference in absolute number of cells within HNA and LNA functional groups

across running analysis of variance with a post-hoc Tukey HSD test (*aov()* and *TukeyHSD();*

stats R package; (76). In addition, we tested the association of HNA and LNA to each other and

with productivity by running ordinary least squares regression with the *lm()* (*stats* R package;

(76)).

Ranking correlation

Ranking correlation between variables was calculated using the Kendall rank correlation

coefficient, using the *kendalltau()* function in Scipy (v1.0.0) or *cor()* in R (v3.2). The 'tau-b'

implementation was used, which is able to deal with ties. Values range from -1 (strong

disagreement) to 1 (strong agreement). The same statistic was used to assess the similarity

between rankings of variable selection methods.

Centered-log ratio transform

First, following guidelines from Paliy & Shanker (77), Gloor et al. (3) and Quinn et al. (78),

relative abundances of OTUs were transformed using a centered log-ratio (CLR) transformation

before variable selection was applied. This means that the relative abundance x_i of a taxa was

transformed according to the geometric mean of that sample, in which there are p taxa present:

$$x_i' = \log(x_i/(\prod_{j=1}^{p} x_j)^{1/p})$$.

Zero values were replaced by $\delta = 1/p^2$. This was done using the scikit-bio package

(www.scikit-bio.org, v0.4.1).

Lasso & stability selection

Scores were assigned to taxa based on an extension of the Lasso estimator, which is called

stability selection (28). In the case of n samples, the Lasso estimator fits the following regression

model:

$$\hat{\beta}^\lambda = \text{argmin}_{\beta \in \mathbb{R}^p} ||y - X\beta||_2^2 + \lambda \sum_{j=1}^{p} |\beta_j|$$,

in which X denotes the abundance table, y the target to predict, which either is HNA cell

abundances (HNAcc) or LNA cell abundances (LNAcc), β the weight of each variable and λ is a

regularization parameter which controls the complexity of the model and prevents overfitting.

The Lasso performs an intrinsic form of variable selection, as the weights of certain variables

will be put to zero.

Stability selection, when applied to the Lasso, is in essence an extension of the Lasso regression.

It implements two types of randomizations to assign a score to the variables, and is therefore also

called the *Randomized Lasso* (RL). The resulting RL score can be seen as the probability that a

certain variable will be included in a Lasso regression model (*i.e.,* its weight will be non-zero

when fitted). When performing stability selection, the Lasso is fitted to B different subsamples of

the data of fraction $n/2$, denoted as X' and corresponding y'. A second randomization is added by

introducing a weakness parameter α. In each model, the penalty λ changes to a randomly chosen

value in the set $[\lambda, \lambda/\alpha]$, which means that a higher penalty will be assigned to a random subset

of the total amount of variables. The Randomized Lasso therefore becomes:

$$\hat{\beta}^\lambda = \text{argmin}_{\beta \in \mathbb{R}^p} ||y' - X'\beta||_2^2 + \lambda \sum_{j=1}^{p} \frac{|\beta_j|}{w_j}$$,

where w_j is a random variable which is either α or 1. Next, the Randomized Lasso score (RL

score) is determined by counting the number of times the weight of a variable was non-zero for

each of the B models and divided by B. Meinshausen and Bühlmann (28) show that, under

stringent conditions, the number of falsely selected variables is controlled for the Randomized

Lasso when the RL score is higher than 0.5. If λ is varied, one can determine the stability path,

which is the relationship between π and λ for every variable. For our implementation, $B = 500$,

$\alpha = 0.5$ and the highest score was selected in the stability path for which λ ranged from 10^{-3}

until 10^3, logarithmically divided in 100 intervals. The *RandomizedLasso()* function from the

scikit-learn machine learning library was used ((79), v0.19.1).

Random Forests & Boruta

The Boruta algorithm is a *wrapper* algorithm that makes use of Random Forests as a base

classification or regression method in order to select all relevant variables in function of a

response variable (29). Similar to stability selection, the method uses an additional form of

randomness in order to perform variable selection. Random Forests are fitted to the data multiple

times. To remove the correlation to the response variable, each variable gets per iteration a so-

called *shadow variable*, which is a permuted copy of the original variable. Next, the Random

Forest algorithm is run with the extended set of variables, after which variable importances are

calculated for both original and shadow variables. The shadow variable that has the highest

importance score is used as reference, and every variable with significantly lower importance, as

determined by a Bonferroni corrected t-test, is removed. Likewise, variables containing an

importance score that is significantly higher are included in the final list of selected variables.

This procedure can be repeated until all original variables are either discarded or included in the

final set; variables that remain get the label 'tentative' (i.e., after all repetitions it is still not

possible to either select or discard a certain variable). We used the boruta_py package to

implement the Boruta algorithm (https://github.com/scikit-learn-contrib/boruta_py). Random

Forests were implemented using *RandomForestRegressor()* function from scikit-learn (79),

v0.19.1. Random Forests were run with 200 trees, the number of variables considered at every

split of a decision tree was $p/3$ and the minimal number of samples per leaf was set to five. The

latter were based on default values for Random Forests in a regression setting (80). The Boruta

algorithm was run for 300 iterations, variables were selected or discarded at $P < .05$ after

performing Bonferroni correction.

Recursive variable elimination

Scores of the Randomized Lasso were evaluated using a recursive variable elimination strategy

(81). Variables were ranked according to the RL score. Next, the lowest-ranked variables were

eliminated from the dataset, after which the Lasso was applied to predict HNAcc and LNAcc

respectively. This process was repeated until only the highest-scored taxa remained. In this way,

performance of the Randomized Lasso was assessed from a minimal-optimal evaluation

perspective (82). In other words, the lowest amount of variables that resulted in the highest

predictive performance was determined.

Performance evaluation

In order to account for the spatiotemporal structure of the data, a blocked cross-validation

scheme was implemented (83). Samples were grouped according the site and year that they were

collected. This results in 5, 10 and 16 distinctive groups for the Michigan, Muskegon and Inland

lake systems respectively. Predictive models were optimized in function of the R^2 between

predicted and true values of held-out groups using a leave-one-group-out cross-validation

scheme with the *LeaveOneGroupOut()* function. This results in a cross-validated R^2_{CV} value. For

the Lasso, λ was determined using the lassoCV() function, with setting eps=10^{-4} and

n_alphas=400. The Random Forest object was optimized using a grid search where max_features

was chosen in the interval $[1, \sqrt{p}, 2\sqrt{p}, ..., p]$ (all variables) or $[1, ..., p]$ (Boruta selected variables)

and min_samples_leaf in the interval $[1, ..., 5]$, using the *GridSearchCV()* function. The number

of decision trees (n_trees) was set to 200. All functions are part of scikit-learn ((79); v0.19.1)

Stability of the Randomized Lasso

Similarity of RL scores between lake systems and functional groups was quantified using the

Pearson correlation. This was done using the *pearsonr()* function in Scipy (v1.0.0).

Patterns of HNA and LNA OTUs across ecosystems and phylogeny

To visualize patterns of selected HNA and LNA OTUs across the three ecosystems, a heatmap

was created with the RL scores of each OTU from the Randomized Lasso regression that were

higher than specified threshold values. The heatmap was created with the *heatmap.2()* function

(*gplots* R package) using the euclidean distances of the RL scores and a complete linkage

hierarchical clustering algorithm (**Figure 4**).

Correlations between taxa and productivity measurements

The Kendall ranking correlation coefficient or Kendall's tau-b between productivity

measurements and individual abundances were calculated on the phylum and OTU level using

the *kendalltau()* function from Scipy (v1.0.0). P-values were corrected using Benjamini-

Hochberg correction, reported as P_adj. This was done using the *multitest()* function from the

Python module Statsmodels ((84); v0.5.0).

Phylogenetic tree construction and signal calculation

We calculated the best performing maximum likelihood tree using the GTR-CAT model (-gtr -

fastest) model of nucleotide substitution with FastTree (version 2.1.9 No SSE3; (85)).

Phylogenetic signal is a measure of the dependence among a species' trait values on their

phylogenetic history (86). If the phylogenetic signal is very strong, taxa belonging to similar

phylogenetic groups (*e.g.* a Phylum) will share the same trait (*i.e.* association with HNAcc or

LNAcc). Alternatively, if the phylogenetic signal is weak, taxa within a similar phylogenetic

group will have different traits. The phylogenetic signal was measured with both discrete (*i.e.*

HNA, LNA, or both) and continuous traits (*i.e.* the RL score) using the newick tree from

FastTree. For the most part, Pagel's lambda was used (87) to test for phylogenetic signal and was

calculated with the fitDiscrete() function from the geiger R package (discrete trait; (88)) and the

phylosig() function from the phytools R package (continuous trait; (89)). The lambda value

varies between 0 and 1, with 1 indicating complete phylogenetic patterning and 0 representing no

phylogenetic patterning, leading to a tree collapsing into a single polytomy. was then used to

model phylogenetic signal using Pagel's lambda, Blomberg's K (phylosig() function from the

phytools R package (89)), and Moran's I (abouheif.moran() function from the adephylo R

package (90)).

Acknowledgements

PR was supported by Ghent University (BOFSTA2015000501) and MLS was supported by the National Science Foundation Graduate Research Fellowship Program (Grant No. DGE 1256260). Part of the computational resources (Stevin Supercomputer Infrastructure) and services used in this work were provided by the VSC (Flemish Supercomputer Center), funded by Ghent University, the Hercules Foundation and the Flemish Government department EWI. Flow cytometry analysis was supported through a Geconcerteerde Onderzoeksactie (GOA) from Ghent University (BOF15/GOA/006).

Author Contributions

MLS and PR co-wrote the paper with contributions from RP, BB, NB, WW, and VJD. MLS, RP, and BB generated the data. MLS, PR, and RP performed the data analysis. PR, MLS, RP, WW, and VJD designed the study.

References

1. Lennon JT, Jones SE. 2011. Microbial seed banks: the ecological and evolutionary implications of dormancy. Nature Reviews Microbiology 9:119–130.

2. Carini P, Marsden PJ, Leff JW, Morgan EE, Strickland MS, Fierer N. 2017. Relic DNA is abundant in soil and obscures estimates of soil microbial diversity. Nature Microbiology 2:16242.

3. Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ. 2017. Microbiome Datasets Are Compositional: And This Is Not Optional. Frontiers in Microbiology 8.

4. Knight R, Vrbanac A, Taylor BC, Aksenov A, Callewaert C, Debelius J, Gonzalez A, Kosciolek T, McCall L-I, McDonald D, Melnik AV, Morton JT, Navas J, Quinn RA, Sanders JG, Swafford AD, Thompson LR, Tripathi A, Xu ZZ, Zaneveld JR, Zhu Q, Caporaso JG, Dorrestein PC. 2018. Best practices for analysing microbiomes. Nature Reviews Microbiology 16:410–422.

5. Widder S, Allen RJ, Pfeiffer T, Curtis TP, Wiuf C, Sloan WT, Cordero OX, Brown SP, Momeni B, Shou W, Kettle H, Flint HJ, Haas AF, Laroche B, Kreft J-U, Rainey PB, Freilich S, Schuster S, Milferstedt K, van der Meer JR, Groβkopf T, Huisman J, Free A, Picioreanu C, Quince C, Klapper I, Labarthe S, Smets BF, Wang H, Isaac Newton Institute Fellows, Soyer OS. 2016. Challenges in microbial ecology: building predictive understanding of community function and dynamics. The ISME Journal 10:2557–2568.

6. Gasol JM, Del Giorgio PA. 2000. Using flow cytometry for counting natural planktonic bacteria and understanding the structure of planktonic bacterial communities. Scientia Marina 64:197–224.

7. Vives-Rego J, Lebaron P, Nebe-von Caron G. 2000. Current and future applications of flow cytometry in aquatic microbiology. FEMS Microbiology Reviews 24:429–448.

8. Wang Y, Hammes F, De Roy K, Verstraete W, Boon N. 2010. Past, present and future applications of flow cytometry in aquatic microbiology. Trends in Biotechnology 28:416–424.

9. Gasol JM, Zweifel UL, Peters F, Fuhrman JA. 1999. Significance of Size and Nucleic Acid Content Heterogeneity as Measured by Flow Cytometry in Natural Planktonic Bacteria. Applied and Environmental Microbiology 65:4475–4483.

10. Lebaron P, Servais P, Agogue H, Courties C, Joux F. 2001. Does the high nucleic acid content of individual bacterial cells allow us to discriminate between active cells and inactive cells in aquatic systems? Applied and Environmental Microbiology 67:1775–1782.

11. Bouvier T, del Giorgio PA, Gasol JM. 2007. A comparative study of the cytometric characteristics of High and Low nucleic-acid bacterioplankton cells from different aquatic ecosystems. Environmental Microbiology 9:2050–2066.

12. Wang Y, Hammes F, Boon N, Chami M, Egli T. 2009. Isolation and characterization of low nucleic acid (LNA)-content bacteria. The ISME Journal 3:889–902.

13. Lebaron P, Servais P, Baudoux A, Bourrain M, Courties C, Parthuisot N. 2002. Variations
of bacterial-specific activity with cell size and nucleic acid content assessed by flow
cytometry. Aquatic Microbial Ecology 28:131–140.

14. Servais P, Casamayor E, Courties C, Catala P, Parthuisot N, Lebaron P. 2003. Activity and
diversity of bacterial cells with high and low nucleic acid content. Aquatic Microbial
Ecology 33:41–51.

15. Morán X, Bode A, Suárez L, Nogueira E. 2007. Assessing the relevance of nucleic acid
content as an indicator of marine bacterial activity. Aquatic Microbial Ecology 46:141–152.

16. Servais P, Courties C, Lebaron P, Troussellier M. 1999. Coupling Bacterial Activity
Measurements with Cell Sorting by Flow Cytometry. Microbial Ecology 38:180–189.

17. Bowman JS, Amaral-Zettler LA, J Rich J, M Luria C, Ducklow HW. 2017. Bacterial
community segmentation facilitates the prediction of ecosystem function along the coast of
the western Antarctic Peninsula. The ISME Journal 11:1460–1471.

18. Moràn XAG, Ducklow HW, Erickson M. 2011. Single-cell physiological structure and
growth rates of heterotrophic bacteria in a temperate estuary (Waquoit Bay, Massachusetts).
Limnology and Oceanography 56:37–48.

19. Read DS, Gweon HS, Bowes MJ, Newbold LK, Field D, Bailey MJ, Griffiths RI. 2015.
Catchment-scale biogeography of riverine bacterioplankton. The ISME Journal 9:516–526.

20. Sherr EB, Sherr BF, Longnecker K. 2006. Distribution of bacterial abundance and cell-
specific nucleic acid content in the Northeast Pacific Ocean. Deep Sea Research Part I:
Oceanographic Research Papers 53:713–725.

21. García FC, Calleja ML, Al-Otaibi N, Røstad A, Morán XAG. 2018. Diel dynamics and
coupling of heterotrophic prokaryotes and dissolved organic matter in epipelagic and
mesopelagic waters of the central Red Sea. Environmental Microbiology 20:2990–3000.

22. Jochem FJ, Lavrentyev PJ, First MR. 2004. Growth and grazing rates of bacteria groups
with different apparent DNA content in the Gulf of Mexico. Marine Biology 145:1213–
1225.

23. Arnoldini M, Heck T, Blanco-Fernández A, Hammes F. 2013. Monitoring of Dynamic
Microbiological Processes Using Real-Time Flow Cytometry. PLoS ONE 8:e80117.

24. Ramseier MK, von Gunten U, Freihofer P, Hammes F. 2011. Kinetics of membrane
damage to high (HNA) and low (LNA) nucleic acid bacterial clusters in drinking water by
ozone, chlorine, chlorine dioxide, monochloramine, ferrate(VI), and permanganate. Water
Research 45:1490–1500.

25. Huete-Stauffer T, Morán X. 2012. Dynamics of heterotrophic bacteria in temperate coastal
waters: similar net growth but different controls in low and high nucleic acid cells. Aquatic
Microbial Ecology 67:211–223.

26. Morán XAG, Alonso-Sáez L, Nogueira E, Ducklow HW, González N, López-Urrutia Á,
Díaz-Pérez L, Calvo-Díaz A, Arandia-Gorostidi N, Huete-Stauffer TM. 2015. More,
smaller bacteria in response to ocean's warming? Proceedings of the Royal Society B:
Biological Sciences 282:20150371.

27. Proctor CR, Besmer MD, Langenegger T, Beck K, Walser J-C, Ackermann M, Bürgmann
H, Hammes F. 2018. Phylogenetic clustering of small low nucleic acid-content bacteria
across diverse freshwater ecosystems. The ISME Journal 12:1344–1359.

28. Meinshausen N, Bühlmann P. 2010. Stability selection: Stability Selection. Journal of the
Royal Statistical Society: Series B (Statistical Methodology) 72:417–473.

29. Kursa MB, Rudnicki WR. 2010. Feature Selection with the **Boruta** Package. Journal of
Statistical Software 36.

30. Li H. 2015. Microbiome, Metagenomics, and High-Dimensional Compositional Data
Analysis. Annual Review of Statistics and Its Application 2:73–94.

31. Strobl C, Boulesteix A-L, Kneib T, Augustin T, Zeileis A. 2008. Conditional variable
importance for random forests. BMC Bioinformatics 9.

32. Biddanda B. 2017. Global Significance of the Changing Freshwater Carbon Cycle. Eos
98:1–5.

33. Salcher MM, Posch T, Pernthaler J. 2013. *In situ* substrate preferences of abundant
bacterioplankton populations in a prealpine freshwater lake. The ISME Journal 7:896–907.

34. Schubert AM, Rogers MAM, Ring C, Mogle J, Petrosino JP, Young VB, Aronoff DM,
Schloss PD. 2014. Microbiome Data Distinguish Patients with *Clostridium difficile*
Infection and Non-*C. difficile*-Associated Diarrhea from Healthy Controls. mBio 5.

35. Baxter NT, Zackular JP, Chen GY, Schloss PD. 2014. Structure of the gut microbiome
following colonization with human feces determines colonic tumor burden. Microbiome
2:20.

36. Herren CM, McMahon KD. 2018. Keystone taxa predict compositional change in microbial
communities: Keystone microbes predict community turnover. Environmental
Microbiology 20:2207–2217.

37. Lin W, Shi P, Feng R, Li H. 2014. Variable selection in regression with compositional
covariates. Biometrika 101:785–797.

38. Zaura E, Brandt BW, Prodan A, Teixeira de Mattos MJ, Imangaliyev S, Kool J, Buijs MJ,
Jagers FL, Hennequin-Hoenderdos NL, Slot DE, Nicu EA, Lagerweij MD, Janus MM,
Fernandez-Gutierrez MM, Levin E, Krom BP, Brand HS, Veerman EC, Kleerebezem M,
Loos BG, van der Weijden GA, Crielaard W, Keijser BJ. 2017. On the ecosystemic
network of saliva in healthy young adults. The ISME Journal 11:1218–1231.

39. Chen J, Chia N, Kalari KR, Yao JZ, Novotna M, Soldan MMP, Luckey DH, Marietta EV,
Jeraldo PR, Chen X, Weinshenker BG, Rodriguez M, Kantarci OH, Nelson H, Murray JA,
Mangalam AK. 2016. Multiple sclerosis patients have a distinct gut microbiota compared to
healthy controls. Scientific Reports 6:28484.

40. Ma J, Prince AL, Bader D, Hu M, Ganu R, Baquero K, Blundell P, Alan Harris R, Frias
AE, Grove KL, Aagaard KM. 2014. High-fat maternal diet during pregnancy persistently
alters the offspring microbiome in a primate model. Nature Communications 5:3889.

41.   Degenhardt F, Seifert S, Szymczak S. 2017. Evaluation of variable selection methods for
718   random forests and omics data sets. Briefings in Bioinformatics.

719   42.   McCarthy A, Chiang E, Schmidt ML, Denef VJ. 2015. RNA Preservation Agents and
720   Nucleic Acid Extraction Method Bias Perceived Bacterial Community Composition. PLOS
721   ONE 10:e0121659.

722   43.   Louca S, Doebeli M, Parfrey LW. 2018. Correcting for 16S rRNA gene copy numbers in
723   microbiome surveys remains an unsolved problem. Microbiome 6:41.

724   44.   Van der Gucht K, Cottenie K, Muylaert K, Vloemans N, Cousin S, Declerck S, Jeppesen E,
725   Conde-Porcuna J-M, Schwenk K, Zwart G, Degans H, Vyverman W, De Meester L. 2007.
726   The power of species sorting: Local factors drive bacterial community composition over a
727   wide range of spatial scales. Proceedings of the National Academy of Sciences 104:20404–
728   20409.

729   45.   Adams HE, Crump BC, Kling GW. 2014. Metacommunity dynamics of bacteria in an arctic
730   lake: the impact of species sorting and mass effects on bacterial production and
731   biogeography. Frontiers in Microbiology 5.

732   46.   Chiang E, Schmidt ML, Berry MA, Biddanda BA, Burtner A, Johengen TH, Palladino D,
733   Denef VJ. 2018. Verrucomicrobia are prevalent in north-temperate freshwater lakes and
734   display class-level preferences between lake habitats. PLOS ONE 13:e0195112.

735   47.   Aanderud ZT, Vert JC, Lennon JT, Magnusson TW, Breakwell DP, Harker AR. 2016.
736   Bacterial Dormancy Is More Prevalent in Freshwater than Hypersaline Lakes. Frontiers in
737   Microbiology 7.

738   48.   Jones SE, Lennon JT, Karl D. 2010. Dormancy contributes to the maintenance of microbial
739   diversity. Proceedings of the National Academy of Sciences of the United States of
740   America 107:5881–5886.

741   49.   Zimmermann R, Iturriaga R, Becker-Birck J. 1978. Simultaneous Determination of the
742   Total Number of Aquatic Bacteria and the Number Thereof Involved in Respiration.
743   Applied and Environmental Microbiology 36:926–935.

744   50.   Corno G, Jürgens K. 2006. Direct and Indirect Effects of Protist Predation on Population
745   Size Structure of a Bacterial Strain with High Phenotypic Plasticity. Applied and
746   Environmental Microbiology 72:78–86.

747   51.   Coleman ML, Sullivan MB, Martiny AC, Steglich C, Barry K, DeLong EF, Chisholm SW.
748   2006. Genomic islands and the ecology and evolution of *Prochlorococcus*. Science
749   311:1768–1770.

750   52.   Hunt DE, David LA, Gevers D, Preheim SP, Alm EJ, Polz MF. 2008. Resource Partitioning
751   and Sympatric Differentiation Among Closely Related Bacterioplankton. Science
752   320:1081–1085.

753   53.   Denef VJ, Kalnejais LH, Mueller RS, Wilmes P, Baker BJ, Thomas BC, VerBerkmoes NC,
754   Hettich RL, Banfield JF. 2010. Proteogenomic basis for ecological divergence of closely
755   related bacteria in natural acidophilic microbial communities. Proceedings of the National
756   Academy of Sciences 107:2383–2390.

54.   Shapiro BJ, Polz MF. 2014. Ordering microbial diversity into ecologically and genetically
cohesive units. Trends in Microbiology 22:235–247.

55.   Vila-Costa M, Gasol JM, Sharma S, Moran MA. 2012. Community analysis of high- and
low-nucleic acid-containing bacteria in NW Mediterranean coastal waters using 16S rDNA
pyrosequencing: Bacterial composition of different cytometric populations in
Mediterranean waters. Environmental Microbiology 14:1390–1402.

56.   Props R, Schmidt ML, Heyse J, Vanderploeg HA, Boon N, Denef VJ. 2018. Flow
cytometric monitoring of bacterioplankton phenotypic diversity predicts high population-
specific feeding rates by invasive dreissenid mussels: Phenotypic tracking of
bacterioplankton. Environmental Microbiology 20:521–534.

57.   Props R, Kerckhof F-M, Rubbens P, De Vrieze J, Hernandez Sanabria E, Waegeman W,
Monsieurs P, Hammes F, Boon N. 2017. Absolute quantification of microbial taxon
abundances. The ISME Journal 11:584–587.

58.   Amalfitano S, Fazi S, Ejarque E, Freixa A, Romaní AM, Butturini A. 2018. Deconvolution
model to resolve cytometric microbial community patterns in flowing waters:
Deconvolving Cytometric Microbial Subgroups. Cytometry Part A 93:194–200.

59.   Song Y, Wang Y, Mao G, Gao G, Wang Y. 2019. Impact of planktonic low nucleic acid-
content bacteria to bacterial community structure and associated ecological functions in a
shallow lake. Science of The Total Environment 658:868–878.

60.   Schattenhofer M, Wulf J, Kostadinov I, Glöckner FO, Zubkov MV, Fuchs BM. 2011.
Phylogenetic characterisation of picoplanktonic populations with high and low nucleic acid
content in the North Atlantic Ocean. Systematic and Applied Microbiology 34:470–475.

61.   Newton RJ, Jones SE, Eiler A, McMahon KD, Bertilsson S. 2011. A Guide to the Natural
History of Freshwater Lake Bacteria. Microbiology and Molecular Biology Reviews 75:14–
49.

62.   Woodhouse JN, Kinsela AS, Collins RN, Bowling LC, Honeyman GL, Holliday JK, Neilan
BA. 2016. Microbial communities reflect temporal changes in cyanobacterial composition
in a shallow ephemeral freshwater lake. The ISME Journal 10:1337–1351.

63.   Tada Y, Suzuki K. 2016. Changes in the community structure of free-living heterotrophic
bacteria in the open tropical Pacific Ocean in response to microalgal lysate-derived
dissolved organic matter. FEMS Microbiology Ecology 92:fiw099.

64.   Vandeputte D, Kathagen G, D'hoe K, Vieira-Silva S, Valles-Colomer M, Sabino J, Wang J,
Tito RY, De Commer L, Darzi Y, Vermeire S, Falony G, Raes J. 2017. Quantitative
microbiome profiling links gut community variation to microbial load. Nature.

65.   Frossard A, Hammes F, Gessner MO. 2016. Flow Cytometric Assessment of Bacterial
Abundance in Soils, Sediments and Sludge. Frontiers in Microbiology 7.

66.   Foladori P, Bruni L, Tamburini S, Ziglio G. 2010. Direct quantification of bacterial biomass
in influent, effluent and activated sludge of wastewater treatment plants by using flow
cytometry. Water Research 44:3807–3818.

67.   Couradeau E, Sasse J, Goudeau D, Nath N, Hazen TC, Bowen BP, Malmstrom RR,
Northen TR. 2018. Study of Oak Ridge soils using BONCAT-FACS-Seq reveals that a
large fraction of the soil microbiome is active. bioRxiv.

68.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.
2012. The SILVA ribosomal RNA gene database project: improved data processing and
web-based tools. Nucleic Acids Research 41:D590–D596.

69.   Rohwer RR, Hamilton JJ, Newton RJ, McMahon KD. 2018. TaxAss: Leveraging a Custom
Freshwater Database Achieves Fine-Scale Taxonomic Resolution. mSphere 3:e00327-18.

70.   Weiss S, Van Treuren W, Lozupone C, Faust K, Friedman J, Deng Y, Xia LC, Xu ZZ,
Ursell L, Alm EJ, Birmingham A, Cram JA, Fuhrman JA, Raes J, Sun F, Zhou J, Knight R.
2016. Correlation detection strategies in microbial data sets vary widely in sensitivity and
precision. The ISME Journal 10:1669–1681.

71.   McMurdie PJ, Holmes S. 2013. phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. PLoS ONE 8:e61217.

72.   Kirchman D, K'Nees E, Hodson R. 1985. Leucine Incorporation and Its Potential as a
Measure of Protein Synthesis by Bacteria in Natural Aquatic Systemst. Applied and
Environmental Microbiology 49:9.

73.   Simon M, Azam F. 1989. Protein content and protein synthesis rates of planktonic marine
bacteria. Marine Ecology Progress Series 51:201–213.

74.   Prest EI, Hammes F, Kötzsch S, van Loosdrecht MCM, Vrouwenvelder JS. 2013.
Monitoring microbiological changes in drinking water systems using a fast and
reproducible flow cytometric method. Water Research 47:7131–7142.

75.   Spidlen J, Breuer K, Rosenberg C, Kotecha N, Brinkman RR. 2012. FlowRepository: A
resource of annotated flow cytometry datasets associated with peer-reviewed publications.
Cytometry Part A 81A:727–731.

76.   R Core Team. 2018. R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria.

77.   Paliy O, Shankar V. 2016. Application of multivariate statistical techniques in microbial
ecology. Molecular Ecology 25:1032–1057.

78.   Quinn TP, Erb I, Richardson MF, Crowley TM. 2018. Understanding sequencing data as
compositions: an outlook and review. Bioinformatics 34:2870–2878.

79.   Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M,
Prettenhofer P, Weiss R, Dubourg V, Vanderplas J, Passos A, Cournapeau D. Scikit-learn:
Machine Learning in Python. MACHINE LEARNING IN PYTHON 6.

80.   Probst P, Wright M, Boulesteix A-L. 2018. Hyperparameters and Tuning Strategies for
Random Forest.

81.   Guyon I, Weston J, Barnhill S. 2002. Gene Selection for Cancer Classification using
Support Vector Machines. Machine Learning 46:389–422.

82.   Nilsson R, Bjorkegren J, Tegner J. Consistent Feature Selection for Pattern Recognition in
835   Polynomial Time 24.

836   83.   Roberts DR, Bahn V, Ciuti S, Boyce MS, Elith J, Guillera-Arroita G, Hauenstein S, Lahoz-
837   Monfort JJ, Schröder B, Thuiller W, Warton DI, Wintle BA, Hartig F, Dormann CF. 2017.
838   Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic
839   structure. Ecography 40:913–929.

840   84.   Seabold S, Perktold J. 2010. Statsmodels: Econometric and Statistical Modeling with
841   Python. Proceedings of the 9th Python in Science Conference (SciPy 2010) 57–61.

842   85.   Price MN, Dehal PS, Arkin AP. 2010. FastTree 2 – Approximately Maximum-Likelihood
843   Trees for Large Alignments. PLoS ONE 5:e9490.

844   86.   Revell LJ, Harmon LJ, Collar DC. 2008. Phylogenetic Signal, Evolutionary Process, and
845   Rate. Systematic Biology 57:591–601.

846   87.   Pagel M. 1999. Inferring the historical patterns of biological evolution. Nature 401:877–
847   884.

848   88.   Harmon LJ, Weir JT, Brock CD, Glor RE, Challenger W. 2008. GEIGER: investigating
849   evolutionary radiations. Bioinformatics 24:129–131.

850   89.   Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other
851   things). Methods in Ecology and Evolution 3:217–223.

852   90.   Jombart T, Balloux F, Dray S. 2010. adephylo: new tools for investigating the phylogenetic
853   signal in biological traits. Bioinformatics 26:1907–1909.

Figure 1: (A) Correlation between HNA cell counts and LNA cell counts across the three
freshwater lake ecosystems. **(B-D)** Muskegon Lake bacterial heterotrophic production and
its correlation with **(B)** HNA cell counts (HNAcc), **(C)** LNA cell counts, (LNAcc) and **(D)**
total cell counts. The grey area in plots A, B, and D represents the 95% confidence
intervals.

Figure 2: R_{CV}^2 in function of the number of OTUs, which were iteratively removed based on the
RL score and evaluated using the Lasso at every step. The solid (HNA) and dashed (LNA)
vertical lines corresponds to the threshold (i.e., number of OTUs) which resulted in a maximal
R_{CV}^2. **(A)** Inland system ($R_{CV,max}^2 = 0.92$), HNAcc; **(B)** Lake Michigan ($R_{CV,max}^2 = 0.53$),
HNAcc; **(C)** Muskegon lake, HNAcc ($R_{CV,max}^2 = 0.85$); **(D)** Inland system, LNAcc ($R_{CV,max}^2 =$
0.87); **(E)** Lake Michigan, LNAcc ($R_{CV,max}^2 = 0.79$; **(F)** Muskegon lake, LNAcc ($R_{CV,max}^2 =$
0.91).

Figure 3: Evaluation of HNA cell counts (HNAcc) and LNA cell counts (LNAcc) predictions

using the Lasso at all taxonomic levels for the Muskegon lake system, expressed in terms of R^2_{CV},

using different subsets of taxonomic variables. Subsets were determined by iteratively

eliminating the lowest-ranked taxonomic variables based on the RL score.

Figure 4: Hierarchical clustering of the RL score for the top 10 selected OTUs within each lake

system and FCM functional groups with the selected OTU (rows) across HNA and LNA groups

within the three lake systems (columns).

Figure 5: Phylogenetic tree with all HNA and LNA selected OTUs from each of the three lake

systems with their phylum level taxonomic classification and association with HNA, LNA or to

both groups based on the RL score threshold values.

Figure 6: Correlation (Kendall's tau-b) between the relative abundances of the top three OTUs

selected by the RL and the densities in the cytometric fingerprint. The fluorescence threshold

used to manually define HNA and LNA populations is indicated by the dotted line.

Table 1: Top scored OTUs according to the RL per functional population and lake ecosystem.

Selection according to the Boruta algorithm is given in addition to the RL score. Descriptive

statistics by means of the Kendall rank correlation coefficient have been added with level of

significance in function of the HNA/LNA population. Full taxonomy of the OTUs is given in

Table S2.

Lake system	Functional group	OTU	RL score	Boruta selected	Kendall's tau (HNA)	P-value (HNA)	Kendall's tau (LNA)	P-value (LNA)
Inland	HNA	OTU369	0.382	yes	-0.43	<0.001	-0.28	0.0012
	LNA	OTU555	0.384	no	0.089	N.S.	0.22	0.011
Michigan	HNA	OTU025	0.362	yes	0.46	<0.001	0.41	<0.001
	LNA	OTU168	0.428	yes	0.26	0.0092	0.4	<0.001
Muskegon	HNA	OTU173	0.462	yes	0.5	<0.001	0.2	0.019
	LNA	OTU029	0.568	no	0.26	0.0029	0.49	<0.001