Multimodal integration and vividness in the angular gyrus during episodic encoding and retrieval

Roni Tibon*[1], Delia Fuhrmann[1], Daniel A. Levy[2], Jon S. Simons[3], & Richard Henson[1]

[1] MRC Cognition & Brain Sciences Unit, University of Cambridge, UK, CB2 7EF

[2] Baruch Ivcher School of Psychology, The Interdisciplinary Center, Herzliya, Israel, 4610101

[3] Department of Psychology, University of Cambridge, Cambridge, UK, CB2 3EB

*Corresponding author:

Dr. Roni Tibon
MRC Cognition & Brain Sciences Unit
University of Cambridge
15 Chaucer Road
Cambridge, CB2 7EF
UNITED KINGDOM
Telephone: +44 1223 355 294
Email: roni.tibon@mrc-cbu.cam.ac.uk

RUNNING TITLE: ANGULAR GYRUS IN EPISODIC MEMORY

Pages: 25   **Tables**: 2   **Figures**: 3

Word count in **Abstract** : 225   **Introduction** : 650   **Discussion** : 1337

Acknowledgements

RT is supported by a Newton International Fellowship by the Royal Society and the British Academy (grant SUAI/009 RG91715) and by a British Academy Postdoctoral Fellowship (grant SUAI/028 RG94188). JSS is supported by the James S McDonnell Foundation (#220020333). RH is supported by UK Medical Research Council grant (SUAG/010 RG91365). The authors wish to thank Josefina Weinerova for her assistance in collecting the data.

The authors declare no competing financial interests.

Abstract

Much evidence suggests that the angular gyrus (AnG) is involved in episodic memory, but its precise role is yet to be determined. We examined two possible accounts, within the same experimental paradigm: the CoBRA account (Shimamura, 2011), which suggests that the AnG acts as a convergence zone that binds multimodal episodic features; and the Subjectivity account (Yazar et al., 2012), which implicates AnG involvement in subjective mnemonic experience (such as vividness or confidence). fMRI was employed during both encoding and retrieval of paired-associates. During study, participants memorised picture-pairs of common objects (in the unimodal task) or of an object-picture and an environmental sound (in the crossmodal task). At test, they performed a cued-recall task, and further indicated the vividness of their memory. During retrieval, BOLD activation in the AnG was greatest for vividly remembered associates, consistent with the Subjectivity account. During encoding, the same effect of vividness was found, but this was further modulated by task: Greater activations were associated with subsequent recall in the crossmodal than the unimodal task. Thus, encoding data suggests an additional role to the AnG in cross-modal integration, consistent with its role at retrieval proposed by CoBRA. These results resolve some of the puzzles in the literature and indicate that the AnG can play different roles during encoding and retrieval, determined by the cognitive demands posed by different mnemonic tasks.

Significance Statement

We offer new insights into the multiplicity of processes that are associated with angular gyrus (AnG) activation during encoding and retrieval of newly formed memories. We used fMRI while human participants learned and subsequently recalled pairs of objects presented to the same sensory modality or to different modalities. We were able to show that the AnG is involved when vivid memories are created and retrieved, as well as when encoded information is integrated across different sensory modalities. These findings provide novel evidence for the contribution of the AnG to our subjective experience of remembering, alongside its role in integrative processes that promote subsequent memory.

Introduction

The ventral posterior parietal cortex (vPPC), and the angular gyrus (AnG) in particular, have been associated with numerous cognitive functions, including episodic memory. Although the vPPC is considered one of the most active regions during successful episodic retrieval (reviewed by Wagner et al., 2005; Vilberg & Rugg, 2008; Shimamura, 2011; Levy, 2012; Rugg & King, 2017; Sestieri et al., 2017), patients with lateral parietal lesions can often successfully retrieve episodic memories, and are not usually considered to be amnesic.

Two main explanations have been proposed to account for AnG activation during episodic retrieval. The first derives from findings of fewer 'remember' responses, fewer high-confidence responses, and lack of richness, vividness and specificity of retrieved episodic events in patients with parietal lesions (Berryhill et al., 2007; Simons et al., 2010; Hower et al., 2014). Under this account (Yazar et al., 2012), vPPC involvement is viewed in terms of subjective mnemonic abilities, with the AnG involved in one's own experience of episodic memory retrieval, rather than in objective memory performance as expressed in response accuracy. This "Subjectivity account" has been further supported by several studies with healthy individuals (Qin et al., 2011; Yazar et al., 2014; Richter et al., 2016).

The second account is the "Cortical Binding of Relational Activity" (CoBRA) theory (Shimamura, 2011), which suggests that during retrieval, the vPPC (including the AnG) acts as a convergence zone that binds episodic features stored in disparate neocortical regions. This account builds on anatomical features of the vPPC—its central location and connections to many neocortical regions (Seghier, 2013)—making it well situated for binding episodic memories by establishing intermodal links across diverse event features. CoBRA critically predicts that tasks that demand the reinstatement of intermodal features, such as voices with faces, will depend more on the vPPC than tasks that depend on within-modality associations. This account has received support from several studies (Bonnici et al., 2016; Yazar et al., 2017), but other examinations of CoBRA were inconclusive,

with multimodal integration deficits found in parietal lesion patients (Ben-Zvi et al., 2015), but no multimodal vs. unimodal associative memory effects on parietal scalp ERPs recorded in healthy volunteers (Tibon & Levy, 2014).

Even though theoretically these two accounts are distinct, in practice they are not easily dissociated, as the multisensory features comprising episodic representations enable us to experience the rich and vivid details that characterise remembering. To tease apart these accounts, we utilised a pair-associate learning paradigm, based on our prior work (Tibon & Levy, 2014; Ben-Zvi et al., 2015). fMRI of healthy volunteers was employed during two learning tasks. In the unimodal task, stimulus pairs were two semantically-unrelated object-pictures, and in the crossmodal task, stimulus pairs were an object-picture and a nameable but unrelated sound. At study, participants created a mental image of the two stimuli. At test, an object-picture was presented as the cue, and participants recalled the associated target from study, either another picture or a sound, and indicated the vividness of their memory. Thus, crossmodal "events" involved different modalities for separate objects, rather than multimodal representations of the same object. Therefore, even though these events are multimodal, they are not necessarily more vivid.

To test the retrieval-focused predictions of CoBRA and the Subjectivity accounts, we examined BOLD activation during test. The CoBRA account predicts greater parietal activity associated with objective retrieval (the difference between recall success and failure trials) in the crossmodal than the unimodal task, regardless the vividness of retrieved memories. The Subjectivity account, on the other hand, predicts a linear pattern of activity—vivid recall > non-vivid recall > recall failure—that is independent of modality. In addition, as some previous findings from patients' studies might be due to encoding rather than retrieval deficits, we also examined memory effects from the study phase. Finally, given recent claims regarding AnG laterality effects (Bellana et al., 2016), we examined whether the left and right AnG display different memory-related activation patterns.

Methods

Participants

Twenty-four adults participated in the experiment (20 females, *mean age* = 25.83, *SD* = 4.02) and were reimbursed for their time. One participant who was not fully scanned due to a problem with the equipment, and two participants, who did not provide any responses in one of the experimental conditions (one with no success responses and another with no non-vivid responses), were excluded, leaving 21 participants whose data were analysed. Participants were recruited from the volunteer panel of the MRC Cognition and Brain Science Unit in Cambridge, UK. All were fluent English speakers, MR-compatible, had normal hearing and normal or corrected-to-normal vision, and were never diagnosed with ADHD, dyslexia or any other developmental or learning disabilities. They provided informed consent for a protocol approved by a local ethics committee (Cambridge Psychological Research Ethics Committee reference PRE2016.055).

Materials

The materials for the experiment were previously used by Tibon & Levy (2014), although some (~10%) were replaced to improve stimulus quality. Replaced stimuli were verified as recognisable by 6 pilot participants, who did not take part in the main study. Auditory stimuli were 120 environmental sounds (such as animal sounds, tools, vehicles, etc.) downloaded from various internet sources, and edited using Audacity audio editing software, at 44 kHz, 16- bit resolution, mono mode. They were all adjusted to the same amplitude level and edited to last 2 seconds. Twelve additional sounds were used for practice trials and examples. Visual stimuli were 360 colour drawings of common objects obtained from various internet sources, including fruits and vegetables, tools, sporting goods, electrical and electronic devices, animals, food, vehicles, body parts, furniture, and clothing, each approximately 6–8 cm in on-screen size.

To form the various experimental conditions, several stimulus lists were created. Two lists of 120 pictures served as cue lists for recollection. Two additional lists served as target lists, one containing 120 pictures and the other containing 120 sounds. Each entry in the cue lists was pseudorandomly assigned to an entry in the target lists, so that all stimulus-pairs were semantically unrelated. While, unavoidably, the target lists (one list of pictures and another of sounds) remained fixed for the crossmodal and unimodal conditions, two experimental lists were created to counterbalance the cues across participants: half the participants viewed the cues from the first cue list with the auditory target and the cues from the second cue list with the visual target, and vice versa for the other half. Each experimental list was further divided into three sub-lists, to allow 1/3 of the stimuli to be presented three times (the other 2/3 presented only once, see more details below), with the sub-lists used to counter-balance the repeated stimuli across participants.

Procedure

The experiment consisted of two tasks, which took place on different days (*mean days apart* = 7.14, *SD* = 4.8). One was the crossmodal pair-associate learning and cued recall task, and the other was the unimodal pair-associate learning and cued recall task. Task order was counter-balanced across participants. On arrival at the lab, participants were seated in a quiet room, where they signed an informed consent form and performed the unscanned part of the relevant task. Participants were told that they would be presented with pairs of stimuli (sound-picture pairs in the crossmodal task and picture-picture pairs in the unimodal task), and were instructed to study those pairs for subsequent retrieval by forming an association between the stimuli. They were told that after 40 pairs had been presented, a test phase would follow, in which cue pictures alone would be presented. They would then need to recall and say the name of the paired object portrayed in the picture (in the unimodal task) or of the paired object making the sound (in the crossmodal task) that had accompanied that cue. Following these instructions, a headset with a microphone was provided,

together with a practice block of 5 trials. During practice, the experimenter ascertained that the participant understood the nature of the associations that were to be generated for the stimulus pairs. After completion of the practice block, participants were told that the unscanned part of the task would be repeated twice, with the same 40 pairs. They were further informed that following the unscanned part, they would be asked to perform the same task in the fMRI scanner, but for 120 pairs: 40 pairs that had just been studied and recalled at the unscanned part, and 80 new pairs.

During the unscanned part of the task, participants viewed two (identical) study-test blocks of 40 stimulus pairs each. Each study trial started with a fixation cross, displayed for an exposure time jittered across a range of 1-4 s. Stimulus pairs were then presented for 2 s, followed by a 7 s fixation cross. Participants were instructed to use this time to generate an association. Next, a screen with the text "Easy/Hard?" was shown for 2 s. During this time, participants were asked to press a green key if they found it easy to come up with an association, or a red key if they found it difficult. The right index and middle fingers were used for these keypresses, and finger assignment was counterbalanced across participants.

After all 40 study trials had been presented, the test phase started. Test trials also started with a 1-4 s jittered display time fixation cross. The cue picture was then presented alone for 2 s, followed by a 6 s fixation cross. Next, a slide with the text "Pair associate?" was shown for 3 s, and participants were asked to provide their verbal response while the slide was presented. They were encouraged to formulate their response while the fixation cross was presented, so that once the "Pair associate?" slide appeared they could provide their answer immediately. They were further told that if they could not recall the target object, they should not try to guess. Instead, they should say "pass" when the "Pair associate?" slide was displayed. Finally, a slide with the legend "Vivid?" appeared for 2 s, and participants were asked to press a green key if their memory was vivid, or a red key if it was not. Finger assignment was counterbalanced across participants, and matched the

assignment during the study phase (such that the same green key was used for both "Easy" and "Vivid" responses, and the same red key was used for both "Hard" and "Not vivid" responses).

After completing the unscanned part, which included 2 study-test blocks, participants were provided with instructions for the MRI scan. They were told that even though they would need to give verbal responses in the scanner, they should try to minimize head movements. They then performed the task in the MRI scanner, with one block of 120 pairs (40 pairs that were studied during the unscanned part intermixed with 80 new pairs). Finally, after the scan, participants returned to the lab where they performed a debriefing session. In this session, all target stimuli (either sounds in the crossmodal task, or pictures in the unimodal task) were displayed again, and participants were asked to type their names. The data from this session was used to subsequently eliminate (a negligible number of) trials, in which certain participants were unable to provide a name (even if inaccurate) for the sound / picture that was presented.

fMRI Acquisition

The same acquisition and pre-processing protocol was used for both visits (one visit for the crossmodal task, and another for the unimodal task). MRI data were collected using a Siemens 3 T TIM TRIO system (Siemens, Erlangen, Germany). Structural images were acquired with a T1-weighted 3D Magnetization Prepared RApid Gradient Echo (MPRAGE) sequence [repetition time (TR) = 2250 ms; echo time (TE) = 3.02 ms; inversion time (TI) = 900 ms; 230 Hz per pixel; flip angle = 9°; field of view (FOV) 256 x 256 x 192 mm; GRAPPA acceleration factor 2]. Functional images were acquired using an echoplanar imaging (EPI) sequence with multi-band (factor = 4) acquisition. Volumes were acquired over 2 runs, one for the study phase (*mean number of volumes* = 1095.5, *SD* = 11.1) and one for the test phase (*M* = 1263, *SD* = 6.8). Each volume contained 76 slices (acquired in interleaved order within each excitation band), with a slice thickness of 2 mm and no interslice gap

(TR = 1448 ms; TE = 33.4 ms; flip angle = 74°; FOV = 192 mm × 192 mm; voxel-size = 2 mm × 2 mm × 2 mm). Field maps for EPI distortion correction were also collected (TR = 541 ms; TE = 4.92 ms; flip angle = 60°; FOV = 192 × 192 mm).

Data Pre-processing

Data were pre-processed using SPM12 (www.fil.ion.ucl.ac.uk/spm), automated in Matlab (v8.0.0.783 R2012b, The MathWorks) with Automatic Analysis (AA) 5.0 (Cusack, Vicente-Grabovetsky, Mitchell, Wild, Auer, Linke, & Peelle, 2015; https://github.com/rhodricusack/automaticanalysis). T1 anatomical images, for each participant in each visit, were coregistered to the Montreal Neurological Institute (MNI) template using rigid-body transformation, bias-corrected, and segmented by tissue class. Diffeomorphic registration was then applied across participants, separately for each visit, to the grey matter to create a group template using DARTEL (Ashburner, 2007), which was, in turn, affine-transformed to MNI space. EPI distortions in the functional images were corrected using field maps. Next, the images were corrected for motion and then for slice acquisition times by interpolating to the 26th slice in time. The images were rigid-body co-registered to the corresponding T1 image and transformed to MNI space using the diffeomorphic + affine transformations. These normalised images were then smoothed by 6mm FWHM.

Statistical Analyses

Trial Classification. Study and test trials were classified into three response types of interest, based on subsequent response to that trial (for the study phase, to allow examination of subsequent memory effects) or actual response (for the test phase, to allow examination of recall success effects): (1) vivid: trials with correctly recalled targets, marked as "vivid"; (2) non-vivid: correctly

recalled targets, classified as "non-vivid"; (3) failure: all trials with a "pass" response. When the participants' response did not match the name of the object, the data from the debriefing session was used to determine whether the mismatch was due to misidentification (i.e., the response at test matched the response at debriefing) in which case the trial was classified as "correct", or from failure to retrieve the correct answer (i.e., the response at test did not match the response at debriefing) in which case it was marked as a "false alarm". False alarm trials, trials in which no response was collected during the scanning session and trials in which the target was not named during the debriefing session, were marked as "excluded" (see fMRI acquisition).

Behavioural Analyses. All statistical analyses were performed in R v3.4.1 and RStudio v1.0153. After classifying the trials according to their responses, we analysed the behavioural data from study and test. To test for task effects (differences between the unimodal and the crossmodal task) during the study phase, data were analysed using a paired-sample t-test, to compare encoding difficulty (calculated as % of pairs for which an "easy" response was recorded at encoding) between the unimodal and the crossmodal tasks. To test for memory effects at test, accuracy data (number of trials for which vivid, non-vivid and failure responses were obtained) were analysed with a repeated measures ANOVA, which included task (crossmodal, unimodal) and response type (vivid, non-vivid, failure) as within-subject factors, using the ez package (Lawrence, 2016). Whenever sphericity assumptions were violated, Greenhouse-Geisser corrected p-values are reported. Main effects and interactions were decomposed with Bonferroni corrected pairwise comparisons using the lsmeans package (Lenth, 2016). Degrees of freedoms were corrected using the Satterthwaite method, as implemented in the lmerTest package (Kuznetsova et al., 2017).

ROI Analyses. A priori ROIs in the AnG were defined based on the coordinates of peak left AnG activation from a previous univariate meta-analysis of episodic memory by Vilberg and Rugg (2008; [-43, -66, 38]) and its homologous location in the right hemisphere [43, -66, 38]. These coordinates were also used more recently in a study by Bonnici et al. (2016). Timeseries data from

these ROIs were extracted by taking the first eigenvariate across voxels within a 6mm-radius sphere centred on these coordinates, and adjusting the data using an F-contrast on the first-level univariate analysis that identified the regressors of interest (see whole-brain univariate analysis below), thereby adjusting for all other regressors (e.g. movement parameters). The first eigenvariate reflects the first principal component of the timeseries across all voxels in an ROI, i.e., the principal source of variance within a region, without assuming that all voxels contribute equally.

To analyse ROI data, we used mixed-models to accommodate both within- and between-subject variability. This approach is particularly recommended for unbalanced data (an unequal number of trials in each condition, Tibon & Levy, 2015), which we had here due to the post-hoc division of trials into vivid, non-vivid and failure trials (See Tibon et al., 2014; Tibon & Levy, 2014a; Tibon & Levy, 2014b for a similar use of this approach). This procedure allows estimation of the BOLD response for each trial and does not require averaging over trials for each participant. We estimated this response (separately for study and test) using the Least Squares Separate (LSS-N) approach (Mumford et al., 2012; Abdulrahman & Henson, 2016), where N is the number of conditions. LSS-N fits a separate General Linear Model (GLM) for each trial, with one regressor for the trial of interest, and one regressor for all other trials of each condition. This implements a form of temporal smoothness regularisation on the parameter estimation (Abdulrahman & Henson, 2016). The regressors were created by convolving a delta function at the onset of each stimulus with a canonical HRF. The parameters for the regressor of interest for the ROI were then estimated using ordinary least squares, and the whole process repeated for each separate trial.

The resulting Betas were submitted to a linear mixed-model. The model included phase (study, test), task (crossmodal, unimodal), response type (vivid, non-vivid, failure), laterality (left, right) and all possible interactions between these factors as the fixed part of the model, and subject-

specific slopes for each factor[1], plus a subject-specific intercept, as the random part of the model. We fitted the model using restricted maximum likelihood in R package lme4 (Kuznetsova et al., 2014) with the following formula ('*' indicates all possible interactions, and '(…|x)' indicates random effects):

*betas ~ phase * task * response type * laterality + (1 + phase + task + response type + laterality | subject)*

To test the predictions of the CoBRA and the Subjective accounts, we set two pre-defined contrasts, which we applied separately for each phase using the phia R package (De Rosario-Martinez, 2015). The first contrast was set to test CoBRA's prediction of a greater objective recall success effect (success [collapsed across vividness] – failure) for crossmodal than for unimodal memories. The second contrast was set to test the prediction of the Subjectivity account of a linear pattern that is independent of modality (vivid > non-vivid > failure). Reported results are Bonferroni-corrected for two multiple comparisons (across study and test).

Whole-brain Univariate Analyses. We further conducted whole-brain analyses to ensure that we replicated the basic encoding and retrieval mnemonic effects found in previous studies. SPM12 and AA were used to construct General Linear Models (GLMs) for each participant, separately for each run (study, test) and for each task (crossmodal, unimodal; which took place at different visits). These first-level GLMs included 3 separate regressors for each response type of interest (vivid, non-vivid, failure) and a regressor for excluded responses (either false alarms or unnameable targets, see above). These regressors were modelled at the onset of stimulus presentation. For the study run, the GLM further included a regressor for the motor response to the "Easy/Hard" slide (locked to the motor response). For the test run, the GLM further included a

---

[1] We first attempted to estimate the maximal random effects model (Barr et al., 2013), i.e., a model that also included all possible interactions as random slopes. However, this overparameterised model failed to converge (see Bates et al., 2018 for an exhaustive discussion of this problem). We therefore reduced the complexity of the random effects structure to the model described above, which was the maximal model that converged.

regressor for the motor response to the "Vivid?" slide (locked to the motor response) and a regressor for the verbal response (at the onset of the "Pair-associate?" slide). Each of these regressors was generated with a delta function convolved with a canonical hemodynamic response function (HRF). Six subject-specific movement parameters were also included to capture residual movement-related artefacts. The GLM was fit to the data in each voxel. The autocorrelation of the error was estimated using an AR(1)-plus-white-noise model, together with a set of cosines that functioned to high-pass the model and data to 1/128 Hz, fit using Restricted Maximum Likelihood (ReML). The estimated error autocorrelation was then used to "pre-whiten" the model and data, and ordinary least squares used to estimate the model parameters.

The images for the parameter estimates for each of the three response types for each task and each phase were then entered into second-level GLM corresponding to a repeated-measures ANOVA, which treated subject as a random effect, and with phase (study, test), task (unimodal, crossmodal) and response type (vivid, non-vivid, failure) as repeated factors. SPMs were created of the T-statistic for the "recall success" contrast of interest (success [vivid + non-vivid] − failure) at study and test. The statistical threshold was set to $p < .05$ (FWE corrected for multiple comparisons across the whole brain) at the cluster-level, when the voxel-level was thresholded at $p < .0001$ (uncorrected).

Results

Behavioural results

Study Phase. A paired-sample t-test revealed a significant difference in task difficulty during study, $t(20) = 3.63$, $p = .0017$, CI [.03 .13], with greater proportion of easily formed associations in the unimodal task ($M = .69$, $SD = .19$) than in the crossmodal task ($M = .61$, $SD = .23$). This difference was accounted for in our subsequent ROI analyses (see below).

Test Phase. The number of responses for each response type in the crossmodal and unimodal tasks are shown in Figure 1. Analysis of these data showed a significant main effect of task, $F(1, 20) = 31.89$, $p < .0001$, $\eta^2G = .01$, a significant main effect of response type, $F(2, 40) = 46.48$, $p < .0001$, $\eta^2G = .65$, and a significant interaction between these factors, $F(2, 40) = 5.11$, $p = .01$, $\eta^2G = .04$. Bonferroni corrected pairwise comparisons showed greater number of vivid responses in the unimodal than in the crossmodal task, $t(42.67) = 4.05$, $p = .0002$, but no difference between the tasks in the amount of non-vivid or failure responses. These results support our expectation that, when different modalities are associated with separate objects within the event, rather than with the same object, crossmodal events are not necessarily remembered more vividly than unimodal events.

Figure 1. Number of responses for each response type in the crossmodal task (dark grey) and the unimodal task (light grey). Error bars represent SEs for each condition separately (*** $p < .001$).

ROI Mixed-Effects Analyses. For the main hypotheses under investigation, we focused on left and right AnG ROIs, defined a priori from independent data. Figure 2 depicts contrasts of the Beta values, according to our a priori contrasts of interest, averaged across left and right AnG; Table 1 shows the Betas for each condition and each ROI separately.

Figure 2. Contrasts of Beta values of the fitted model, in the AnG (collapsed across left and right AnG). Each plot represents a planned contrast: the CoBRA account (left; greater recall success effect for crossmodal than unimodal) and the Subjective account (right; linear trend of vivid > non-vivid > fail) during study (top panel) and test (bottom panel). Error bars represent Satterthwaite approximation of the pooled SE, and were computed for each condition separately. ** $p < .01$; *** $p < .005$.

The estimated model showed a significant main effect of response type, $\chi^2(2) = 64.2$, $p < .0001$, a significant interaction between phase and response type, $\chi^2(2) = 18.1$, $p = .0001$, a significant interaction between phase and laterality, $\chi^2(1) = 21.33$, $p < .0001$, a significant interaction between task and laterality, $\chi^2(1) = 8.14$, $p = .004$, and a significant 3-way interaction between phase, task, and response type, $\chi^2(2) = 11.27$, $p = .004$. Because laterality did not interact with response type, $\chi^2(2) = 1.76$, $p = .42$, we collapsed across this factor when computing our planned comparisons.

The first planned comparison confirmed a greater recall success effect for crossmodal than unimodal memories during the study phase, $\chi^2(1) = 8.77$, $p = .006$, but not during the test phase, $\chi^2(1) = 0.19$, $p = 1$. The second planned comparison confirmed a linear trend, whereby vivid > non-vivid > fail, during both study, $\chi^2(1) = 29.14$, $p < .0001$, and test, $\chi^2(1) = 72.07$, $p < .0001$.

Table 1. Adjusted mean Beta values in the left and right AnG during study and test for each response type (failure, non-vivid, vivid) in each task (crossmodal, unimodal). SEs calculated for each condition separately, are given in parentheses.

	Left AnG			Right AnG		
	Failure	**Non-vivid**	**Vivid**	**Failure**	**Non-vivid**	**Vivid**
Study						
Crossmodal	-0.043	-0.028	-0.019	-0.037	-0.024	-0.014
	(0.007)	(0.006)	(0.004)	(0.006)	(0.006)	(0.004)
Unimodal	-0.033	-0.034	-0.022	-0.023	-0.018	-0.009
	(0.008)	(0.007)	(0.006)	(0.007)	(0.007)	(0.004)
Test						
Crossmodal	-0.036	-0.014	-0.01	-0.035	-0.032	-0.01
	(0.007)	(0.007)	(0.005)	(0.007)	(0.007)	(0.005)
Unimodal	-0.037	-0.026	-0.008	-0.038	-0.025	-0.005
	(0.008)	(0.007)	(0.006)	(0.008)	(0.007)	(0.005)

To ensure that the interaction between modality and response type at study cannot be explained by difficulty in forming associations (as our behavioural analysis indicated greater difficulty in the crossmodal vs. the unimodal task), we ran an additional analysis, which included the difficulty rating provided for each trial during study as a covariate in our model. Despite the addition of the covariate, the three-way interaction remained significant, $\chi^2(2) = 11.28$, $p = .004$, and the same pattern as before was confirmed by our planned comparisons, i.e., a greater recall success effect for crossmodal than unimodal memories at study phase, $\chi^2(1) = 8.22$, $p = .008$, but not at test, $\chi^2(1) = 0.58$, $p = 09$, as well as a linear trend during both study, $\chi^2(1) = 25.46$, $p < .0001$, and test, $\chi^2(1) = 65.28$, $p < .0001$.

Whole-brain Univariate Analyses. Looking beyond the AnG, to confirm that our paradigm replicated the typical engagement of regions across the brain during episodic memory encoding and retrieval, we conducted a whole-brain analysis (results are shown in Figure 3). We examined the activation to successful memory trials (collapsed across vividness), relative to failure trials, at both study and test. For both phases, this contrast yielded increased BOLD response in multiple brain regions that have previously been linked with episodic memory (see Table 2 for full results).

Figure 3. Brain regions showing stronger BOLD response for successful vs. failure recall at study (left) and test (right). *p* < 0.05 FWE cluster-level corrected, with voxel-level threshold at *p* < 0.0001 (uncorrected). Data are shown on sagittal, coronal, and axial slices of the group-averaged brain (*n* = 21).

Table 2. Regions of increased BOLD activation during study and test in the recall success contrast of interest.

Lat	Region	Peak *x,y,z*	Cluster size	*T* value
Study				
R	Precuneus	6, -68, 46	1242	6.19
R	Angular gyrus	44, -54, 54	515	5.01
R	Middle temporal gyrus	66, -24, -8	80	5.02
R	Middle cingulate gyrus	4, -18, 30	214	4.87
R	Middle frontal gyrus	32, 66, 2	225	4.52
L	Angular gyrus	-38, -62, 42	955	5.9
L	Middle temporal gyrus	-64, -38, -8	348	5.63
L	Middle frontal gyrus	-30, 58, 0	216	4.65
Test				
R	Cerebellum	38, -56, -38	1376	7.13
R	Angular gyrus	42, -50, 20	1648	5.62
R	Superior frontal gyrus	22, 30, 48	482	5.19
R	Caudate	16, 2, 26	172	5.1
R	Postcentral gyrus	50, -10, 26	247	5.05
R	Hippocampus	28, -34, -4	74	4.99
R	Middle frontal gyrus	58, -40, -14	98	4.7
L	Supramarginal gyrus [a]	-50, -40, 42	5356	7.9
L	Angular gyrus	-50, -62, 32		
L	Superior frontal gyrus	-20, 26, 56	5047	7.76
L	Precuneus	-6, -52, 30	4494	7.69
L	Inferior frontal gyrus	-54, 12, 8	1882	6.96
L	Hippocampus	-26, -28, -10	321	6.57
L	Supplementary motor cortex	-4, 2, 62	123	5.95
L	Putamen	-12, 8, -12	135	5.88
L	Thalamus	-8, -28, 8	84	5.04
L	Cerebellum [b]	-18, -64, -26	75	4.99
L		-36, -74, -42	236	4.89

All regions arising from the contrast of interest, at a threshold of *p* < 0.05 FWE cluster-level corrected, with voxel-level threshold at *p* < .0001 (uncorrected). Anatomical labels were provided by the Neuromorphometrics atlas, available via SPM (Neuromorphometrics, Inc. http://www.neuromorphometrics.com/).

[a] Cluster included local maxima in the supramarginal gyrus and the angular gyrus.

[b] Two local maxima were observed in the left cerebellum at test.

Discussion

The current study tested two dominant accounts of the activation in parietal cortex consistently found in neuroimaging studies of episodic memory—the CoBRA account and the Subjectivity account—and revealed a dissociation between encoding and retrieval of memory-related activations in the angular gyrus (AnG). During retrieval, we found a linear trend as predicted by the Subjectivity account: BOLD activation was greatest for vividly remembered associates and least for associates that failed to be recalled. Contrary to the prediction of the CoBRA account however, which predicts greater activation when information from multiple modalities is reinstated at retrieval, the magnitude of the objective recall success effect did not differ between the unimodal and crossmodal tasks. During encoding, activation associated with subsequent memory showed the same linear pattern predicted by the Subjectivity account, but in addition, this pattern was now moderated by task, with greater activations associated with subsequent recall success in the crossmodal than unimodal task. Memory effects at encoding and retrieval did not interact with lateralisation, suggesting that, at least with the current task, the left and right AnG play a similar role. Taken together, these results support the Subjectivity account, but not the CoBRA account as currently specified, which implicates the involvement of the AnG in multimodal reinstatement - specifically at retrieval; rather, in our current data, the involvement of AnG in multimodal processing was restricted to encoding.

Our finding that AnG activity scales with ratings of vividness during retrieval is consistent with prior studies that have found AnG to be sensitive to qualitative characteristics of memory such as vividness, confidence, and precision (e.g., Kuhl & Chun, 2014; Yazar et al., 2014; Richter et al., 2016), supporting the notion that the AnG is involved in the subjective experience of remembering. We show further here that the linear pattern associated with vivid remembering in the AnG also occurs at encoding, suggesting that it is also involved in the construction of representations that enable vivid subsequent memories. We note that this pattern of encoding-related activity, i.e.,

greater AnG activation for subsequently remembered vs. forgotten memories, is inconsistent with some previous studies (e.g., Daselaar et al., 2009; Lee et al., 2017), which reported negative subsequent memory effects in the AnG (that is, greater activation for subsequently forgotten vs. remembered memories). Nevertheless, an extensive meta-analysis (Uncapher & Wagner, 2009) indicated that both positive and negative subsequent memory effects are observed in the vPPC. Although a comprehensive explanation of this discrepancy is not yet available, Uncapher and Wagner (2009) suggest that retention interval is a crucial predictor of the directionality of subsequent memory effects in the vPPC, whereby positive effects are more often associated with relatively short retention times (< 45 minutes), which is consistent with our present findings.

This dissociation between encoding and retrieval potentially resolves a puzzle arising from our prior studies using a very similar paradigm, where we found that patients with parietal lesions did show a deficit in multimodal reinstatement (Ben-Zvi et al., 2015), prima facie supporting the CoBRA account, but we also failed to find any effect of multimodal reinstatement on parietal ERPs recorded during retrieval in healthy volunteers (Tibon & Levy, 2014), contrary to the CoBRA account. The lack of effects of multimodal reinstatement at retrieval in the present fMRI study agree with the prior ERP study, while the effect of multimodal processing that was found at encoding suggests that the deficit in patients might arise when they encode the paired-associates, rather than being a problem at retrieval.

Our finding that AnG activation is not modulated by multimodal reinstatement during retrieval needs to be considered with respect to other findings. In particular, two previous studies have shown retrieval-related involvement of the AnG in tasks that require multimodal reinstatement. In the first study, Yazar et al. (2017) used continuous post-encoding theta burst transcranial magnetic stimulation to interrupt AnG functioning. Participants encoded objects which were presented both visually and auditorily, and were subsequently asked to retrieve unimodal sources (e.g., which side + which location) or crossmodal sources (e.g., which side + male or female

voice) of the studied objects. Following stimulation, participants' ability to retrieve information was reduced for crossmodal but not for unimodal sources. In the second study, by Bonnici et al. (2016), participants memorised audio clips, visual clips, and audio-visual clips, presenting different objects (e.g., a train). At test, they were given a verbal cue for the to-be-recalled clip (e.g., the word "train"), and were asked to recall the associated clip as vividly as possible. This study showed greater AnG activation when crossmodal than unimodal memories were retrieved, coupled with above chance classification accuracy of individual crossmodal (but not unimodal) memories in the AnG. This previous evidence of AnG involvement in multimodal retrieval effects suggest that although our current results seem to dissociate between encoding and retrieval of memory-related activations in AnG, a simple distinction between memory stages might be too simplistic. Instead, we suggest that what determines AnG involvement is not the memory stage per-se (encoding vs. retrieval), but rather the specific combination of cognitive demands posed by the task. Hence, unlike CoBRA, which predicts AnG involvement whenever multimodal elements are reinstated, we suggest that such activation might be further constrained by other task characteristics, such as the type of information that is retrieved and the mnemonic cues that are provided. This suggestion is in line with the component process model of memory (e.g., Moscovitch, 1992; Witherspoon & Moscovitch, 1989), which posits that numerous different processing components, associated with distinct brain regions, are recruited in various combinations by different memory tasks. Thus, different task demands would involve distinct process-specific alliances (that is, transient interactions between several brain regions, Cabeza & Moscovitch, 2013; Moscovitch et al., 2016). Notably, in accord with this view, even though some aspects of our results seem to be inconsistent with previous studies that used different tasks (Bonnici et al., 2016; Yazar et al., 2017), our findings do converge across the three studies that used the same paradigm (i.e., the current study, Ben-Zvi et al., 2015, and Tibon & Levy, 2014), where task demands were kept similar. Importantly, by elucidating the nature of process-specific alliances, future studies can provide a more decisive and fine-grained account of AnG involvement in multimodal processing.

In the current study, unimodal associations were comprised of two visual stimuli, and crossmodal associations were comprised of a visual and an auditory stimulus. We chose these materials to allow simultaneous presentation of the stimuli (given that simultaneous processing of two auditory stimuli, for example, is perceptually challenging). However, this means that a limitation of the current study is that we cannot be certain that our conclusions generalise to other kinds of materials. It could be, for example, that the AnG is selectively involved in multimodal integration of audio-visual associations, but not in the integration of information deriving from other sensory modalities. While we have no reason to assume that audio-visual associations are unique in this sense, future studies can use a similar paradigm to test other forms of within- and between-modality combinations.

In summary, the results of the current study show that the AnG is involved in multiple mnemonic processes, during both encoding and retrieval. They provide a straightforward answer to the puzzle arising from our previous findings (in Ben-Zvi et al., 2015 versus Tibon & Levy, 2014), by suggesting that the AnG is involved in (at least) two memory-related processes: multimodal integration at encoding, and (re)construction of vivid representations, which takes place during both encoding and retrieval. Based on current and prior evidence (Bonnici et al., 2016; Yazar et al., 2017), it is plausible that the extent of AnG activation in mnemonic processes is determined by specific cognitive demands posed by the task (e.g., Moscovitch et al., 2016). Future studies might systematically manipulate such demands within the same experimental paradigm, and examine activation during encoding and retrieval, possibly using a more fine-grained vividness rating (see also Richter et al., 2017). Nonetheless, our current results provide an important step towards clarification of the complexities regarding AnG involvement in episodic memories.

References

Abdulrahman, H., & Henson, R. N. (2016). Effect of trial-to-trial variability on optimal event-related fMRI design: Implications for Beta-series correlation and multi-voxel pattern analysis. *Neuroimage, 125*, 756-766. https://doi.org/10.1016/j.neuroimage.2015.11.009

Ashburner, J. (2007). A fast diffeomorphic image registration algorithm. *Neuroimage, 38*(1), 95-113. https://doi.org/10.1016/j.neuroimage.2007.07.007

Bellana, B., Liu, Z., Anderson, J. A. E., Moscovitch, M., & Grady, C. L. (2016). Laterality effects in functional connectivity of the angular gyrus during rest and episodic retrieval. *Neuropsychologia*, *80*, 24–34. http://doi.org/10.1016/j.neuropsychologia.2015.11.004

Ben-Zvi, S., Soroker, N., & Levy, D. A. (2015). Parietal lesion effects on cued recall following pair associate learning. *Neuropsychologia*, *73*, 176–194. http://doi.org/10.1016/j.neuropsychologia.2015.05.009

Berryhill, M. E., Phuong, L., Picasso, L., Cabeza, R., & Olson, I. R. (2007). Parietal Lobe and Episodic Memory: Bilateral Damage Causes Impaired Free Recall of Autobiographical Memory. *Journal of Neuroscience*, *27*(52), 14415–14423. http://doi.org/10.1523/JNEUROSCI.4163-07.2007

Bonnici, H. M., Richter, F. R., Yazar, Y., & Simons, J. S. (2016). Multimodal Feature Integration in the Angular Gyrus during Episodic and Semantic Retrieval. *Journal of Neuroscience*, *36*(20), 5462–5471. http://doi.org/10.1523/JNEUROSCI.4310-15.2016

Cabeza, R., Ciaramelli, E., & Moscovitch, M. (2012). Cognitive contributions of the ventral parietal cortex: an integrative theoretical account. *Trends in Cognitive Sciences*, *16*(6), 338–52. http://doi.org/10.1016/j.tics.2012.04.008

Cabeza, R., Ciaramelli, E., Olson, I. R., & Moscovitch, M. (2008). The parietal cortex and episodic memory: an attentional account. *Nature Reviews Neuroscience*, 9(8), 613. http://doi.org/10.1038/nrn2459

Cabeza, R., & Moscovitch, M. (2013). Memory systems, processing modes, and components: functional neuroimaging evidence. Perspectives on Psychological Science, 8(1), 49-55. https://doi.org/10.1177/1745691612469033

Cusack, R., Vicente-Grabovetsky, A., Mitchell, D. J., Wild, C. J., Auer, T., Linke, A. C., & Peelle, J. E. (2015). Automatic analysis (aa): efficient neuroimaging workflows and parallel processing using Matlab and XML. *Frontiers in Neuroinformatics*, 8. https://doi.org/10.3389/fninf.2014.00090

Daselaar, S. M. (2009). Posterior midline and ventral parietal activity is associated with retrieval success and encoding failure. *Frontiers in Human Neuroscience*, *3*. http://doi.org/10.3389/neuro.09.013.2009

Gilmore, A. W., Nelson, S. M., & McDermott, K. B. (2015). A parietal memory network revealed by multiple MRI methods. *Trends in Cognitive Sciences*, *19*(9), 534–543. http://doi.org/10.1016/j.tics.2015.07.004

Hower, K. H., Wixted, J., Berryhill, M. E., & Olson, I. R. (2014). Impaired perception of mnemonic oldness, but not mnemonic newness, after parietal lobe damage. *Neuropsychologia*, *56*(1), 409–417. http://doi.org/10.1016/j.neuropsychologia.2014.02.014

Kuhl, B. A., & Chun, M. M. (2014). Successful Remembering Elicits Event-Specific Activity Patterns in Lateral Parietal Cortex. *Journal of Neuroscience*, *34*(23), 8051–8060. http://doi.org/10.1523/JNEUROSCI.4328-13.2014

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software, 82*(13), pp. 1-26. http://doi.org/10.18637/jss.v082.i13

Lee, H., Chun, M. M., & Kuhl, B. A. (2017). Lower parietal encoding activation is associated with sharper information and better memory. *Cerebral Cortex*, *27*(4), 2486–2499. http://doi.org/10.1093/cercor/bhw097

Lenth, R. (2016). Least-Squares Means: The R Package lsmeans. *Journal of Statistical Software, 69*(1), 1-33. doi:10.18637/jss.v069.i01

Levy, D. A. (2012). Towards an understanding of parietal mnemonic processes: some conceptual guideposts. *Frontiers in Integrative Neuroscience*, *6*. http://doi.org/10.3389/fnint.2012.00041

Moscovitch, M. (1992). Memory and working-with-memory: A component process model based on modules and central systems. Journal of cognitive neuroscience, 4(3), 257-267. https://doi.org/10.1162/jocn.1992.4.3.257

Moscovitch, M., Cabeza, R., Winocur, G., & Nadel, L. (2016). Episodic memory and beyond: the hippocampus and neocortex in transformation. Annual review of psychology, 67, 105-134. https://doi.org/10.1146/annurev-psych-113011-143733

Mumford, J. A., Davis, T., & Poldrack, R. A. (2014). The impact of study design on pattern estimation for single-trial multivariate pattern analysis. *Neuroimage, 103*, 130-138. https://doi.org/10.1016/j.neuroimage.2014.09.026

Qin, S., van Marle, H. J., Hermans, E. J., & Fernández, G. (2011). Subjective sense of memory strength and the objective amount of information accurately remembered are related to distinct neural correlates at encoding. *Journal of Neuroscience, 31*(24), 8920-8927. https://doi.org/10.1523/JNEUROSCI.2587-10.2011

Richter, F. R., Cooper, R. A., Bays, P. M., & Simons, J. S. (2016). Distinct neural mechanisms underlie the success, precision, and vividness of episodic memory. *ELife*, *5*. http://doi.org/10.7554/eLife.18260

Rugg, M. D., & King, D. R. (2017). Ventral lateral parietal cortex and episodic memory retrieval. *Cortex*. http://doi.org/10.1016/j.cortex.2017.07.012

Seghier, M. L. (2013). The angular gyrus: Multiple functions and multiple subdivisions. *The Neuroscientist*, *19*(1), 43–61. http://doi.org/10.1177/1073858412440596

Sestieri, C., Shulman, G. L., & Corbetta, M. (2017). The contribution of the human posterior parietal cortex to episodic memory. *Nature Reviews Neuroscience*, *18*(3), 183–192. http://doi.org/10.1038/nrn.2017.6

Shimamura, A. P. (2011). Episodic retrieval and the cortical binding of relational activity. *Cognitive, Affective & Behavioral Neuroscience*, *11*(3), 277–91. http://doi.org/10.3758/s13415-011-0031-4

Simons, J. S., Peers, P. V., Mazuz, Y. S., Berryhill, M. E., & Olson, I. R. (2010). Dissociation between memory accuracy and memory confidence following bilateral parietal lesions. *Cerebral Cortex*, *20*(2), 479–485. http://doi.org/10.1093/cercor/bhp116

Tibon, R., & Levy, D. A. (2014). The time course of episodic associative retrieval: Electrophysiological correlates of cued recall of unimodal and crossmodal pair-associate learning. *Cognitive, Affective and Behavioral Neuroscience*, *14*(1), 220–235. http://doi.org/10.3758/s13415-013-0199-x

Tibon, R., & Levy, D. A. (2015). Striking a balance: analyzing unbalanced event-related potential data. *Frontiers in Psychology, 6*. https://doi.org/10.3389/fpsyg.2015.00555

Uncapher, M. R., & Wagner, A. D. (2009). Neurobiology of Learning and Memory Posterior parietal cortex and episodic encoding : Insights from fMRI subsequent memory effects and dual-attention theory. *Neurobiology of Learning and Memory*, *91*(2), 139–154. http://doi.org/10.1016/j.nlm.2008.10.011

Vilberg, K. L., & Rugg, M. D. (2008). Memory retrieval and the parietal cortex: A review of evidence from a dual-process perspective. *Neuropsychologia*, *46*(7), 1787–1799. http://doi.org/10.1016/j.neuropsychologia.2008.01.004

Wagner, A. D., Shannon, B. J., Kahn, I., & Buckner, R. L. (2005). Parietal lobe contributions to episodic memory retrieval. *Trends in Cognitive Sciences*, *9*(9), 445–53. http://doi.org/10.1016/j.tics.2005.07.001

Witherspoon, D., & Moscovitch, M. (1989). Stochastic independence between two implicit memory tasks. Journal of Experimental Psychology: Learning, Memory, and Cognition, 15(1), 22. http://dx.doi.org/10.1037/0278-7393.15.1.22

Yazar, Y., Bergström, Z. M., & Simons, J. S. (2014). Continuous theta burst stimulation of angular gyrus reduces subjective recollection. *PLoS ONE*, *9*(10). http://doi.org/10.1371/journal.pone.0110414

Yazar, Y., Bergström, Z. M., & Simons, J. S. (2012). What is the parietal lobe contribution to long-term memory? *Cortex*, *48*(10), 1381–1382. http://doi.org/10.1016/j.cortex.2012.05.011

Yazar, Y., Bergström, Z. M., & Simons, J. S. (2017). Reduced multimodal integration of memory features following continuous theta burst stimulation of angular gyrus. *Brain Stimulation*, *10*(3), 624–629. http://doi.org/10.1016/j.brs.2017.02.011