Persistent hotspots of S*chistosoma mansoni* transmission in the Kenyan waters of Lake Victoria:

a search for snail-related answers beneath the waves

Snails and hotspot transmission of *S. mansoni* in Lake Victoria

M. W. MUTUKU[1,5]*, M. R. LAIDEMITT[2], I. N. MWANGI[1]   F. O. OTIATO[3], E. L. AGOLA[1], M. L.

STEINAUER[4], H. OCHANDA[5] , B. KAMEL[2], E. S. LOKER[2], G. M. MKOJI[1]

1. Centre for Biotechnology Research and Development, Kenya Medical Research Institute,

9   Nairobi, Kenya.

2. Center for Evolutionary and Theoretical Immunology, Parasitology Division, Museum of

Southwestern Biology, Department of Biology, University of New Mexico, Albuquerque,

New Mexico, United States of America

3. Influenza Surveillance Program, Centers for Disease Control and Prevention, Nairobi,

Kenya

4. Department of Basic Medical Sciences, Western University of Health Sciences, Lebanon,

Oregon, United States of America

5. School of Biological Sciences, College of Biological and Physical Sciences, University of

Nairobi, Nairobi, Kenya

* Corresponding author

Email: mmutuku@kemri.org

Abstract

Background

Schistosoma mansoni transmission in Lake Victoria is often intense and hard to interrupt with praziquantel (PZQ) treatment alone. Following termination of a five-year annual treatment campaign, we sought to reveal underlying explanations by investigating snail-related aspects of transmission in six PZQ non-responsive villages that lie within a transmission hotspot located along the west-facing Kenyan shore of the lake, and in four PZQ-responding villages facing the more protected waters of Winam Gulf.

Methodology/Principal Findings

We examined on four occasions over a two-year period the relative abundance and *S. mansoni* prevalence for *Biomphalaria sudanica* along the lake shore and for *B. choanomphala* mostly from deeper waters, and acquisition of *S. mansoni* worms by sentinel mice, at 20 sampling sites (two per village). *Biomphalaria sudanica* was found at all sites and abundance did not differ significantly between hotspot and responding villages. By contrast, *B. choanomphala* was found at all six hotspot villages and only one responding village. Overall, more *B. sudanica* (83/10,249, 0.81%) were positive for *S. mansoni* than *B. choanomphala* (3/2,120, 0.14%). *S. mansoni* prevalence among *B. sudanica* in hotspot villages (60/5,346, 1.12%) was higher than in responding villages (23/4,903, 0.47%), and sentinel mice worm recoveries were also higher in hotspot villages, though variability among sampling times and locations precluded statistical significance in either case. The ratio of predatory echinostome to schistosome infections in snails was significantly higher at responding than hotspot sites.

Conclusions/Significance

All ten villages offer considerable opportunities for transmission because of abundant

Biomphalaria populations. Hot spot villages had higher prevalence of snails positive for *S.*

mansoni suggesting transmission trends higher than in responding villages. Although recovery

of *B. choanomphala* including individuals infected with *S. mansoni* was relatively low, these

snails offer a different, potentially underestimated, deeper water avenue for transmission largely

unavailable in the responding villages.

Author summary

Schistosomiasis is a widespread disease of poverty that has been difficult to control

despite mass drug administration (MDA) campaigns throughout affected regions. After

treatment, people become rapidly reinfected because there is a reservoir of schistosome

parasites in freshwater snail vectors. Our study investigated the role of snails in transmission of

S. mansoni at 10 villages near Lake Victoria that were part of a recently-concluded MDA

campaign. The villages were categorized by the campaign as "responding" if prevalence was

substantially reduced over five years, or as "hotspots of transmission" if prevalence diminished

only slightly or increased despite MDA. We measured relative abundance and *S. mansoni*

prevalence in local snail populations, and infection rates of sentinel mice, at four sampling times

over 13 months. Opportunities for transmission were high at all 10 villages, though hotspot

villages yielded a higher prevalence of infected snails and higher worm recoveries from sentinel

mice. Results were suggestive of higher transmission in hotspot villages though variability

among locations and times precluded statistical significance. Hotspot sites also differed from

responding sites in that the deep-water vector, *B. choanomphala*, was primarily present only at

hotspots, and echinostomes (trematodes that prey on *S. mansoni* sporocysts in snails), were

relatively fewer.

Introduction

One of the most prevalent and persistent of the world's neglected tropical diseases is

human schistosomiasis, caused primarily by three members of the digenetic trematode genus

Schistosoma, *S. mansoni* and *S. japonicum* being responsible for intestinal schistosomiasis and

S. haematobium for urogenital schistosomiasis [1]. There is currently considerable momentum

to bring schistosomiasis under control and to proceed to elimination efforts [2] that would

eventually nullify its under-appreciated impact on public health [3, 4]. An essential weapon in the

elimination of schistosomiasis is the drug praziquantel (PZQ) which has been used extensively

in a variety of control programs that have had considerable impact in reducing prevalence and

intensity of infection [5]. There is also a growing appreciation for the need for integrated control

programs taking into account sanitation, provision of safe water, education, and acknowledging

the fundamental role played by freshwater snails as vectors of the disease [6, 7]. Although

snails are not vectors in a conventional sense, much of the success of schistosome parasites is

owed to their molluscan hosts which support the prolific production of human-infective cercariae,

and that often exist in huge populations across immense freshwater habitats such as Lake

Victoria. Here we focus our attention on the relationship of *Biomphalaria* snails to PZQ-based

control efforts for *S. mansoni* in and around the portion of Lake Victoria in Kenya.

Africa harbors 90% of the global burden of *S. mansoni* [8]. Across the continent, 12

species of snails in the genus *Biomphlaria* perpetuate transmission of this parasite. In Kenya,

there are multiple vectors that appear to be adapted to distinctive habitats: 1) *B. pfeifferi*, a

stream inhabitant whose distribution includes tributaries feeding Lake Victoria, canals in the

Mwea irrigation scheme in central Kenya, and in small impoundments and both seasonal and

perennial streams throughout the country, except in the tropical lowland belt along the coast; 2)

B. sudanica, mainly found along shores of Lake Victoria and its surrounding swamps, and in

Lake Jipe; and 3) *B. choanomphala*, a deeper water inhabitant of Lake Victoria [9, 10]. The

latter two taxa are likely members of the same species [11, 12], and although they are

frequently referred to as ecophenotypes or ecomorphs, their distinctive conchological

differences persist upon lab culture (Mutuku, personal observations). Which of the two formal

names, *sudanica* or *choanomphala,* deserves precedence requires further study and molecular

confirmation using specimens from the type localities (see discussion in [11]).

Prevalence of intestinal schistosomiasis in Kenya is highest (> 50%) in the Mwea

irrigation scheme, central Kenya and in the Lake Victoria basin in western Kenya. Recent

studies assessing the prevalence of *S. mansoni* infection in school children in the lake basin [13,

14] reported overall prevalence of 60.5% and 69%, respectively, with island areas having 2-fold

higher prevalence rates than mainland villages. Schools closer to the lake had higher

prevalence rates than those further from the lake. This would seem to be associated with more

frequent contact with lake water containing infected snails, rendering such children more

vulnerable to infection or re-infections after treatment.

Several chemotherapy-based initiatives have been undertaken to control

schistosomiasis in this region. Among them, the Schistosomiasis Consortium for Operational

Research and Evaluation (SCORE) project implemented school-based and village-wide annual

mass drug administration (MDA) in varying treatment levels for 5 consecutive years (2011-2015)

for 150 Kenyan villages situated on or near the eastern shore of Lake Victoria [15]. In schools

that reported significant declines in prevalence, the initial prevalence ranged from 52-81%, and

following treatment declined to 5.5-19%, and were termed "responding" sites (Table 1).

However, in other schools, PZQ treatment resulted in smaller reductions in prevalence, starting

from 60-91% before to 39-83% prevalence following treatment, with one school actually

experiencing an increase in prevalence over the course of the study [15]. These sites were

referred to as "non-responding" sites and generally fell within a persistent hotspot of infection

identified by [15].

In the current study, building upon the results of Wiegand et al. [15], we investigated the

hypothesis that the hotspot villages, relative to the responding villages, are adjacent to habitats

containing more snails and more infected snails, resulting in greater numbers of infectious

cercariae in the environment. These factors would contribute to persistently high prevalence in

the face of control operations. During the two years immediately following the cessation of the

SCORE-sponsored MDA project, on four separate occasions, we surveyed *Biomphalaria*

populations and their schistosome infection rates adjacent to 10 villages with different degrees

of responsiveness to treatment. We examined both shore-associated *B. sudanica* and *B.*

choanomphala dredged from deeper waters. *B. pfeifferi* was not found in the habitats sampled.

Additionally, we concurrently monitored the presence of viable *S. mansoni* cercariae by placing

sentinel mice at water contact sites associated with each village and later perfused the mice for

evidence of infection. Below we describe the results of looking beneath the waves to ascertain

possible snail-related influences on the pattern of persistent hotspots noted for *S. mansoni*

along the Kenyan shore of Lake Victoria.

Materials and methods

Study sites

The study involved ten villages along the shores of Lake Victoria, western Kenya. The

villages had been subjected to annual school based mass drug administration (MDA) for 5

consecutive years. Six of the villages (Minya, Agok, Migiro, Miyandhe, Kanyibok, Usenge,) had

persistent high prevalence of *S. mansoni* (designated hotspot villages) despite the MDA while

four (Kotieno, Seka Dok, St. Douglas Weta, Mumbo) recorded a significant drop in infection

prevalence (responding villages). In each village, two shoreline habitats were identified where

there was evidence of human-water contact activities, and these were established as our

sampling sites. Each of the 20 sampling sites was visited at four times (April 2016, August 2016,

January 2017 and May 2017).

Fig 1: Map showing position of study villages along the shore of Lake Victoria, Kenya.

The map was constructed using QGIS version 2.18.22 and 3.2.1, the basemap layer was

added from the standard terrain tiles provided from http://maps.stamen.com through the

OpenLayers function. The tiles are © Stamen Design, under a creative commons

attribution (CC BY 3.0) license. The data for the locations were obtained manually using

cellular phone GPS functionality according to the WGS1984 mercator projection.

Table 1: *S. mansoni* prevalence in schools within the study villages showing response to

praziquantel treatment following five years of Mass Drug Administration. Data acquired

from and shown with permission of the SCORE project

SCHOOL	Prevalence (%) Year 1 (2011)	Prevalence (%) Year 2 (2012)	Prevalence (%) Year 3 (2013)	Prevalence (%) Year 4 (2014)	Prevalence (%) Year 5 (2015)
Hotspot villages					
MINYA	68.6	77	75	74	82
AGOK	60.5	59.8	60.2	67	39
MIGIRO	69.4	76.7	49.2	72	48
MIYANDHE	78.6	70.5	71.4	73.5	54
KANYIBOK	91.8	93	83.7	84	83
USENGE	69.4	68	65	46.5	65
Responding villages					
KOTIENO	52.9	40.9	19	16	12
SEKA DOK	81.6	73.2	55.8	48	19
ST.DOUGLAS WETA	55.2	27.4	24	19	19
MUMBO	57.7	30	11.5	15.6	5.5

Snail Sampling

Two methods were used to collect snails from each of the 20 designated sampling sites:

scooping from the shore or dredging by boat. From the shore, two experienced collectors

scooped snails for 15 minutes using long-handled scoops (stainless steel sieve with a mesh

size of 2 × 2 mm, supported on an iron frame). Snails were collected offshore by passing a

dredge (0.75 m long and 0.4 m wide with attached sieve, 2 × 2 mm mesh size) along the bottom.

At each sampling site a series of dredge hauls covering a combined length of 150 meters was

made. Material collected was sorted and live biomphalarid snails were retrieved. All snails were

taken to the laboratory in KEMRI, Center for Global Health Research, Kisumu where they were

sorted into species based on shell morphology characteristics, using standard taxonomic

identification keys [16], counted and screened.

Snail screening

Each snail was placed in an individual well of a 24-well plastic culture plate containing 1

ml of aged de-chlorinated tap water. The plate was placed in indirect sunlight for 2 hours

between 10:00-13:00hr. Individual wells were then examined with the aid of a dissecting

microscope for presence of cercariae. Using standard identification keys [17], cercariae were

identified to basic taxonomic groups, and have been further identified with molecular methods

as parts of other studies [18]. The snails were then maintained in the laboratory as described by

Mutuku et al., [19] for four weeks and screened again to enable detection of snails that harbored

pre-patent infections at the time of collection.

Sentinel mice exposures

Mice were exposed to lake water at each near shore sampling site in floating cages that

ensured the mice were exposed to a depth of no more than 5 mm. The cages were made from a

plastic container measuring 20 X 14 X 7 cm with the bottom of the container replaced with wire

mesh with openings measuring 5 x 5 mm. Styrofoam blocks were attached to the plastic

container to insure the cage floated at the desired position. Lids of the containers were

perforated with 5 mm diameter holes to allow air circulation. The abdomen of each mouse was

shaved to facilitate cercariae penetration. For each of the shoreline sampling habitats, five

cages each with 2 mice were tethered adjacent to locations where snails were collected (total of

20 mice per village per sampling time). Mice were in contact with the water for 3 hours (11:00

hr – 14:00 hr), an interval when shedding of cercariae by *S. mansoni*-infected snails is expected

to be at its peak (Steinauer et al., 2008)**.** Mice from each sampling site were then marked with

an identifying ear tag and maintained together until perfused as described by Tamarath et al.

[20] at eight weeks post exposure. Recovered worms were counted and tabulated with respect

to village, sampling site and sampling time.

Ethics statement

All experiments involving mice were approved by institutional animal care and use

committee (IACUC) at the Kenya Medical Research Institute (Protocol KEMRI/ACUC/03.10.15).

All protocols and practices for the handling and manipulation of mice were in accordance with

the guidelines of the American Veterinary Medical Association (AVMA) for humane treatment of

laboratory animals.

Results

Overview of *B. sudanica* and *B. choanomphala* recovered from the 10 villages

A total of 12,369 snails (10,249 *B. sudanica* and 2,120 *B. choanomphala*) were collected

from the study sites during four collection periods (Table 2, Fig 2). No *B. pfeifferi* were found in

the lake habitats sampled. Snails of *B. sudanica* were recovered along the shoreline from all 20

sampling locations while those identified as *B. choanomphala* were dredged from deeper water

at all sampling locations in hotspot villages. *B. choanomphala* was also found in the scooping

samples from shallow water habitats in two hotspot locations, Kanyibok and Usenge. At Usenge

Beach, the shallow water *B. choanomphala* persisted for at least five months with populations of

B. sudanica found nearby. *B. choanomphala* was also occasionally recovered from the shore

intermingled with *B. sudanica* in the Anyanga beach sampling site at Kanyibok. Mumbo was the

only responding village from which *B. choanomphala* was found (from deep water habitats only).

Of note is that Mumbo is located closer to the hotspot villages than the remaining three

responding villages (Fig 1). One hotspot village, Miyandhe, is located adjacent to a rocky shore

with a steep drop-off in water depth, and thus yielded relatively few snails compared to the

remaining nine villages (Table 2).

The number of snails collected varied widely among the sites and among time periods

(Figs 2 and 3). The number of all *Biomphalaria* snails collected among the 20 sampling sites for

the four sampling times was not significantly different between hotspot villages and responding

villages (two-way repeated measures ANOVA; $F_{1,8}$ = 0.346, p = 0.5750); but the number of

snails were significantly different across sampling time periods, with an increasing trend over

time ($F_{3,24}$ = 64.235; p=0.0155) (Fig 3). As the majority of snails collected were *B. sudanica*, their

numbers also did not differ between hotspot villages and responding villages ($F_{1,8}$=0.41,

p=0.5399), and were also significantly different across sampling time periods ($F_{3,24}$ =4.971;

p=0.008). However, as shown in Fig 4, the mean number of *B. choanomphala* (337.7)

recovered from hotspot villages was 14 times higher than from responding villages (23.5) ($F_{1,8}$

=48.44, p=0.0001; time $F_{3,24}$ =0.845; p=0.4826). In fact, *B. choanomphala* was collected at each

of the hotspot villages, but only one of the four responding villages.

Table 2: The 20 sampling sites and their GPS coordinates, total number of *B. sudanica*

and *B. choanomphala* collected from each, and the number of each species positive for *S.*

***mansoni*.**

Village	Habitat	Latitude	Longitude	*Biomphalaria sudanica*	No. of snails (+) for *S. mansoni*	*Biomphalaria choanomphala*	No. of snails (+) for *S. mansoni*
			Hotspot villages				
MINYA	Kochilo beach	$00^0$14'10.96"S	$034^0$14'45.79"E	522	0	96	0
	Luthi beach	$00^0$14'04.58"S	$034^0$13'58.74"E	758	1	282	0
AGOK	Gudwa beach	$00^0$21'26.52"S	$034^0$19'48.37"E	209	1	238	2
	Kogoye beach	$00^0$20'51.94"S	$034^0$20'74.32"E	220	6	85	0
MIGIRO	Uyawi beach	$00^0$12'18.58"S	$034^0$09'18.36"E	728	1	89	0
	Kadede beach	$00^0$12'12.09"S	$034^0$09'11.74"E	132	0	100	0
MIYANDHE	Kamumbo beach	$00^0$11'49.04"S	$034^0$10'28.17"E	48	0	228	0
	Daraja beach	$00^0$11'46.65"S	$034^0$09'44.11"E	103	0	71	0
KANYIBOK	Anyanga beach	$00^0$05'22.49"S	$034^0$05'09.34"E	897	28	380	1
	Ndiara beach	$00^0$05'27.33"S	$034^0$05'06.24"E	407	0	161	0
USENGE	Police beach	$00^0$04'34.80"S	$034^0$03'24.74"E	833	19	4	0
	Usenge beach	$00^0$04'21.49"S	$034^0$03'35.84"E	489	4	292	0
	B. sudanica 60/5,346 (1.12%), *B. choanomphala* 3/2,026 (0.15%)						
			Responding villages				
KOTIENO	Kotieno beach	$00^0$21'09.07"S	$034^0$40'03.52"E	674	2	0	0
	Maguti beach	$00^0$21'17.57"S	$034^0$39'32.50"E	649	16	0	0
SEKA DOK	Seka beach	$00^0$21'18.58"S	$034^0$40'59.95"E	462	2	0	0
	Kagwa beach	$00^0$21'19"S	$034^0$40'52.12"E	834	1	0	0
ST. DOUGLAS WETA	Kobala beach	$00^0$21'05"S	$034^0$45'31.21"E	735	0	0	0
	Alara beach	$00^0$21'08"S	$034^0$45'16.01"E	584	0	0	0
MUMBO	Kabuong beach	$0^0$20'20.06"S	$034^0$21'04.09"E	576	1	93	0
	Mumbo beach	$0^0$20'0.99"S	$034^0$21'33.03"E	389	1	1	0
	B. sudanica 23/4,903 (0.47%), *B. choanomphala* 0/94 (0%)						

Fig 2: Variation in number of *B. sudanica* (blue bars) and *B. choanomphala* (black bars) collected at each of the four sampling times (April and August 2016, January and May 2017) from the 6 hotspot villages on the top versus the 4 responding villages on the bottom. Numbers over bars indicate number of *S. mansoni* shedders/total number of snails found. Numbers shown within mouse Figs represent the number of worms recovered from sentinel mice at that time point.

Fig 3: **Number of all *Biomphalaria* snails collected from the 20 sampling sites for the four**

collection times was not significantly different for hotspot villages (red bars) and

responding villages (green bars), error bars indicate standard error.

Fig 4: Mean number of *B. choanomphala* snails collected over the four times points was

significantly different for hotspot villages (red bars) and responding villages (green bars),

error bars indicate standard error.

Pearson correlation analysis indicated no significant relationship between the total

abundance of *Biomphalaria* snails at a given site and infection prevalence of *S. mansoni* among

school children at the villages (prevalence measured after 5 years of prior MDA) (Pearson's

r=0.3861, p=0.1352) (Fig 5). However, it was noted that one of the hotspot sites, Miyandhe, with

its rocky, wave-swept lakeshore habitat and low abundance of sampled snails, may be an

outlier due to a small number of snails at the sampling habitats. Indeed, if this site is removed

from the analysis, the correlation becomes significant (Pearson r=0.6337, p=0.0334). In this

case, the correlation accounts for 40% of the variation in the data. Alternatively, the responding

sites and hotspot sites appear to have very different linear relationships (Fig 5) and if only the

hotspot sites are considered, the correlation accounts for 69% of the variation in the data, and

Miyandhe appears to be much less of an outlier (Pearson r=0.8329, p=0.0198).

Fig 5 Relationship between the total number of *Biomphalaria* snails collected from

habitats at each of ten villages and prevalence of *S. mansoni* in school children after 5

years of MDA. Red indicates hotspot villages and green indicates responding villages.

The star indicates the village of Miyandhe, a potential outlier in the analysis.

Snails recovered with *S. mansoni* infections

The overall number of positive snails and prevalence for *S. mansoni* recovered among

all 10 villages examined was higher for *B. sudanica* (83/10,249, 0.81%) than for *B.*

choanomphala (3/2,120, 0.14%). A Z-ratio test indicated prevalence was significantly higher in *B.*

sudanica (p = 0.0007). The overall number and prevalence for *S. mansoni* among *B. sudanica*

was also higher in hotspot villages (60/5,346, 1.12%) than from responding villages (23/4,903,

0.47%), with prevalence in hotspot villages being significantly higher (Z-ratio test p=0.0001) (Fig

2). However, when each site is considered separately, and time is a factor, there was no

significant difference in the number of infected snails between hotspot villages and responding

villages (2-way repeated measures ANOVA, $F_{(1,8)}$=0.582, p=0.4766). There was large variance

among the prevalence of infected snails among both hotspot sites and responding villages (Figs

2 and 6). It is noteworthy that two of the responding villages with the highest residual post-

treatment prevalences of *S. mansoni* in children, Kanyibok and Usenge, had the highest overall

prevalence of positive snails (1.57% and 1.42% respectively (Table 2, Figs 2 and 6).

Fig 6: Prevalence of *S. mansoni* infections amongst *Biomphalaria* snails collected from

hotspot villages (red bars) and responding villages (green bars).

Infection of *B. sudanica* with other digenetic trematodes

In all 10 villages, the number of trematode infections other than *S. mansoni* recovered

from *B. sudanica* (Table 3) exceeded those of *S. mansoni* (totals of 241 and 83, respectively). In

hotspot villages 60 of 184 trematode infections were of *S. mansoni* (32%), whereas in

responding villages, 23 of 150 (15%) were of *S. mansoni*. Prevalence of trematode infections

was similar between hotspot and responding villages (3.5 and 3.1%, respectively),

Echinostomes were the most abundant non-*S. mansoni* infections with 112 snails infected from

the ten study villages. Echinostomes are of particular interest because their intramolluscan

stages ingest schistosome sporocysts, a characteristic which could aid in the control of

schistosomiasis. Indeed, the ratio of echinostomes to schistosomes was significantly higher in

responding sites vs. hotspot sites (T-test with Welch's correction for unequal variance, t=2.047,

p=0.0396), consistent with the hypothesized role for echinostomes in control in schistosomiasis.

Overall, there were significant positive correlations between *S. mansoni* infected snails and echinostome infected snails (Pearson r = 0.6438, p = 0.0445) as well as between *S. mansoni* infected snails and amphistome infected snails (Pearson r = 0.9308, p < 0.0001). Relationships between *S. mansoni* and strigeids and also between *S. mansoni* and xiphidiocercariae were non-significant (strigeid Pearson r = 0.603, p = 0.0650; xiphidicercariae Pearson r = -0.1573, p = 0.6643).

Table 3: Number of *B. sudanica* snails collected from each village and number of snails infected with *S. mansoni* or other trematode cercariae. SM = *Schistosoma mansoni*, ECH = echinostomes, AMPH = amphistomes, STRIG = strigeids, XIPH = xiphidiocercariae, SM:ECH = ratio of *S. mansoni* to echinostomes.

VILLAGE	Total Snails	SM	ECH	AMPH	STRIG	XIPH	Total Trematodes	Trematode Prevalence	SM:EC
Hotspot villages									
MINYA	1280	1	5	0	0	19	25	2.0	0.2
AGOK	429	7	4	0	1	7	19	4.4	1.7
MIGIRO	860	1	5	0	1	6	13	1.5	0.2
MIYANDHE	151	0	1	0	1	0	2	1.3	0.0
KANYIBOK	1304	28	22	7	4	1	62	4.8	1.2
USENGE	1322	23	10	6	12	14	65	4.9	2.3
TOTAL	5346	60	47	13	19	47	186	3.5	1.2
%		32	25	7	10	25			
Responding villages									
KOTIENO	1323	18	29	7	17	2	73	5.5	0.6
SEKA DOK	1296	3	10	2	4	1	20	1.5	0.3
WETA	1319	0	13	1	6	15	35	2.7	0.0
MUMBO	965	2	13	1	2	4	22	2.3	0.1
TOTAL	4903	23	65	11	29	22	150	3.1	0.3
%		15	43	7	19	15			

Schistosomes recovered from sentinel mice

In general, adult schistosome recoveries from the sentinel mice were low (53 worms

recovered, or 0.066 worms per mouse, or 0.022 worms/mouse/hour of exposure). The greatest

number of worms acquired by a single mouse was 7. Of the total number of worms recovered,

40 (75.5%) were from hotspot villages (0.083 worms/mouse) while 13 (24.5%) were from

responding villages (0.041 worms/mouse). The mean number of worms recovered per village

was consistently higher in hotspot villages than in responding villages during the 4 sampling

times, however, this difference was not statistically significant (2-way repeated measures

ANOVA $F_{(1,8)}$ =0.5652, p=0.4737) (Fig 8).

The number of infected snails recovered per village was positively correlated with the

number of worms recovered in sentinel mice (Fig 7) indicating that the sentinel mice exposures

and snail screening give similar results in terms of measuring the force of infection in the water

(Pearson r = 0.607, p=0.0326). However, this analysis indicated that at one site, there were

positive sentinel mice with no infected snails and another site that showed the opposite,

negative sentinel mice and infected snails. Therefore, both measures can underrepresent the

force of infection. There was no significant correlation between the total number of snails

(infected + non-infected) recovered per village and the number of worms recovered from the

sentinel mice (Pearson r = 0.03413, p = 0.347) (Fig 7).

Fig 7: The relationship between sampling snails and sentinel mouse exposure for

measuring the force of infection of schistosomes. A. There is a significant positive

relationship between the number of infected snails and number of schistosomes

recovered from sentinel mice. B. There was no significant relationship between the total

number of snails at a site and the number of schistosomes recovered from sentinel mice.

Red indicates hotspot villages and green indicates responding villages.

Fig 8: Mean number of worms recovered from sentinel mice from hotspot villages (red) and responding villages (green) during the four sampling times. Error bars indicate standard error.

Discussion

The SCORE-supported PZQ-based treatment program in and around Lake Victoria in western Kenya recorded significant overall progress in reducing the prevalence and intensity of *S. mansoni* transmission in the 150 villages included in the program [15]. A hotspot comprised of approximately a third of the villages was identified, especially as gauged by 9-12 year old students, who had a relative risk of infection that was nearly 4 times higher than for students outside the hotspot area. The SCORE program is by no means unique in identifying difficulties in reducing prevalence of infection in some villages adjacent to Lake Victoria [21]. The hotspot for *S. mansoni* transmission identified by Wiegand et al. [15] was in Siaya County, Bondo region, of western Kenya, associated with west-facing villages adjacent to the open, unprotected waters of the lake. In contrast, the PZQ responsive villages faced the more protected waters of Winam Gulf. Among the factors postulated by them to be associated with the hotspot of transmission were presence of superspreaders, larger snail populations or presence of different species of vector snails, and worms more refractory to PZQ.

As expected, *B. sudanica* was abundant along the shoreline in nearly all the sampling sites except in the hotspot village Miyandhe where the shoreline was rocky and wave-swept. By contrast, *B. choanomphala,* was most commonly dredged from waters offshore from the west-facing hotspot villages, and was on average 14 times more common in hotspot than responding villages. This species was found only in one responding village, Mumbo, lying close to the mouth of the gulf. *B. choanomphala* was recovered from shallow water in Usenge where it co-occurred with another typically deeper water gastropod species *Gabbiella humerosa*,

suggesting they had both been transported there, possibly by a storm. Previous studies have

also shown *B. choanomphala* to be rare or absent within Winam Gulf [9], possibly because the

bottom substrate is too muddy, whereas, the substrate found off the shore of the hotspot

villages was more a combination of sand and mud, known to be favourable to *B. choanomphala*

[22]. Gouvras et al. [23] found *B. choanomphala* to be focally distributed in the Tanzanian

waters of the lake and interestingly noted, as did we, a general trend for this species to be more

common in western than eastern locations. They considered a possible gradual change in

substratum conditions from west to east might be responsible. In contrast to our results for

Kenya, Gouvras et al. [23] did note high *B. choanomphala* populations in some more protected

inlets and, in general, found more *B. choanomphala* positive for *S. mansoni* than we did: 12.2%

of all *S. mansoni* shedders were *B. choanomphala* in their study vs. only 3.5% in our study. In

both studies, *B. sudanica* was identified as the major vector for *S. mansoni* in the lake. Many

other factors like levels of dissolved oxygen, pollution and seasonal effects like rainfall may also

influence *B. choanomphala* populations [23, 24]. There was no indication that *B. pfeifferi* played

a role in *S. mansoni* transmission in any of the study villages, as no snails of this species were

found in the streams or small impoundments within the study localities. Overall, abundance of

all *Biomphalaria* snails did not differ between hotspot and responding villages in spite of the

disparities noted for *B. choanomphala*.

With respect to snails positive for *S. mansoni*, the overall prevalence of *S. mansoni*

infections in all the *Biomphalaria* snails collected was almost 2-fold higher for the hotspot

villages (63/7,732, 0.81%) than for the responding villages (23/4,997, 0.46%), with most of all *S.*

mansoni infections coming from *B. sudanica* (96.5%). It was similarly observed that the mean

number of worms recovered per sentinel mouse was also 2-fold higher in hotspot (0.0833) as

compared to responding villages (0.0406). Although relatively few worms were recovered from

the mice so caution is warranted, they do provide a second independent suggestion that the

force of infection from snail to mammal is higher in the hotspot villages. A consequence of this is

that the rate of re-infections would be higher in the hotspot villages than in the responding

villages, having the ultimate effect of re-establishing a high prevalence rate by one year post-

treatment.

We by no means wish to suggest that the differences between hotspot and responding

villages were always clear-cut. For instance, we recovered relatively few snails and no *S.*

mansoni positive snails from Miyandhe, and only one positive snail each from Migiro or Minya,

all hotspot villages. Gouvras et al. [23] also noted localities where human infection rates were

high yet no positive snails were retrieved. Conversely, in one of four responding villages

(Kotieno) no worms were recovered from sentinel mice, even though infected snails were found

on multiple occasions. At St. Douglas Weta, worms were recovered from sentinel mice

exposure even though no infected snails were ever collected from the habitats sampled. Once

cercariae are shed from infected snails, they can swim or move away under the influence of

water currents or waves, and come into contact with a susceptible vertebrate host located away

from the shedding snail [25, 26].

Although *S. mansoni* infection rates were generally low among the *B. choanomphala*

collected (0.14%), our dredge hauls across the extensive bottom surface of the lake can only be

expected to retrieve a small proportion of the snails present and likely grossly underestimate the

role of this snail species in perpetuating transmission. Given the vastness of the lake, it seems

somewhat improbable that *any S. mansoni* miracidia would find *B. choanomphala* in deep water,

especially given there are so many *B. sudanica* close to shore where most human feces would

be deposited. At least two factors might facilitate infections in *B. choanomphala*: currents

moving from offshore to deeper water, and fishermen who may defecate directly into deeper

water from their boats. Another factor potentially favouring involvement of *B. choanomphala* in

transmission is that experimental infection studies of Kenya-derived specimens suggest this

species, like *B. pfeifferi*, is inherently more susceptible to *S. mansoni* than *B. sudanica* [27].

With respect to issues of snail susceptibility, including for *B. sudanica*, given the nearly 2-fold

higher prevalence of infection in the hotspot villages noted, we cannot rule out the possibility

that *B. sudanica* from these localities are simply more susceptible to *S. mansoni* than snails

from the responsive villages.

Not unexpectedly, and as reported from previous studies [28-30], *Biomphalaria* snails

shedding other trematode cercariae were also frequently collected. Similar to Rowel et al. [28]

who found 2.1% of all *Biomphalaria* snails from Ugandan waters of Lake Victoria to be

trematode positive, we found an overall prevalence of 2.6% in our Kenyan study. Non-

schistosome infections exceeded *S. mansoni* infections by a factor of 2.9 in our study. Rowel et

al. [28] also noted excesses of non-schistosome trematodes in Lake Albert (5.33 fold) and Lake

Victoria (6.17 fold). As an aside, numbers like this provide an interesting snapshot of the extent

of *Biomphalaria* usage by trematodes of human vs. animal origin.

Included among the trematodes found from *B. sudanica* were strigeids, xiphidiocercariae,

amphistomes and echinostomes that collectively number approximately 20 species [18]. Such

infections are potentially of interest because larvae of other trematode species, particularly the

rediae of echnostomes, can actively displace schistosome sporocysts in snails via predation, or

may otherwise compete with and either pre-empt or delay cercariae production for *S. mansoni*

should they co-occur in the same snail [31]. Consequently, snails infected with other trematode

species pose hazards for *S. mansoni* and in many ways are effectively removed from the

population of available hosts to infect. Although the preponderance of non-schistosome

infections over *S. mansoni* infections was greater in the responding than hotspot villages, it is

hard to ascribe a primary effect of these infections on limiting *S. mansoni* infections or

contributing to disparities between responding and hotspot villages. This is because the overall

prevalence of such infections is relatively low in the lake. There could be as yet poorly

understood secondary effects of their presence however, that bear consideration. One

possibility is that some trematodes like echinostome produce cercariae that encyst in other

snails as metacercariae. Metacercariae are very commonly recovered from lake-dwelling snails.

It is possible that *S. mansoni* miracidia can somehow detect their presence and avoid such

snails, further diminishing the pool of available snails to infect. Also, if a proportion of the snails

in a population were refractory to all trematode infections, then the competitive interactions

among all trematode species - including *S. mansoni* - for the remaining compatible snails might

be more intense than currently estimated.

In conclusion, the results of the present study suggest that snail-related factors may be

partly responsible for the differences observed in *S. mansoni* prevalence between the hotspot

and responding villages. We estimate that the force of transmission in hotspot villages was

nearly 2 times higher as compared to responding villages as determined by the mean number of

worms recovered per sentinel mouse and number of schistosome-infected snails recovered

from the same hotspot villages. However, caution is required as there is variability amongst

villages in the magnitude of these effects, and numbers of positive snails recovered at different

times were variable even within a village. We noted that *B. sudanica* was present in all the

villages we studied and based on the number of positive snails of this species recovered we

conclude it is also the major vector for *S. mansoni* in all villages. However *B. choanomphala*

was much more frequently found near hotspot villages and in our view play an under-

appreciated and more diffuse and difficult to measure role in transmission. Many other

trematode species infect *Biomphalaria* snails in Lake Victoria but their overall prevalence is low

suggested they do not have a dramatic primary effect on *S. mansoni* prevalence rates in snails,

though secondary effects may be greater than appreciated. Control of *S. mansoni* in and around

the lake will remain a daunting challenge and argues more persuasively than ever for

implementation of improved sanitation and provision of safe water supplies as both parts of an

integrated control approach and as fundamental rights for the people living in the area.

Acknowledgements

The authors acknowledge, with thanks, Dr. Stephen Munga, Director, Center for Global

Health Research, Kenya Medical Research Institute (KEMRI) for providing laboratory space to

carry out the study and to Joseph Kinuthia, Geoffrey Maina, Polycarp Oraro and Boaz Oduor for

their technical assistance in the laboratory and field sampling. This research was undertaken

with approval of National Commission for Science, Technology and Innovations permit number

P/16/9609/12754 and also with approval from National Environmental Management Authority,

permit number NEMA/AGR/46/2014. Field sampling of snails in the habitats was done under

authorization from Kenya Wildlife Service, permit number KWS/BRM/5001. This paper is

published with the approval of the Director, KEMRI.

References

1.   Colley DG, Bustinduy AL, Secor E, King CH. Human schistosomiasis. Lancet.
2014;383(9936):2253-64. doi: 10.1016/S0140-6736(13)61949-2. PubMed PMID: ISI:000338657600030.
2.   Rollinson D, Knopp S, Levitz S, Stothard JR, Tchuente LAT, Garba A, et al. Time to set the agenda
for schistosomiasis elimination. Acta Trop. 2013;128(2):423-40. doi: 10.1016/j.actatropica.2012.04.013.
PubMed PMID: ISI:000326771900027.
3.   King CH. Health metrics for helminth infections. Acta Trop. 2015;141:150-60.
4.   King CH, Dangerfield-Cha M. The unacknowledged impact of schistosomiasis. Chronic Illness.
2008;4:65-79.
5.   Webster JP, Molyneux DH, Hotez PJ, Fenwick A. The contribution of mass drug administration to
global health: past, present and future. Philosophical Transactions of the R oyal Society of London B
Biological Sciences. 2014;369(1645):20130434. doi: doi: 10.1098/rstb.2013.0434. PubMed Central
PMCID: PMC PMC4024227
6.   Sokolow SH, Wood CL, Jones IJ, Swartz SJ, Lopez M, Hsieh MH, et al. Global assessment of
schistosomiasis control over the past century shows targeting the snail intermediate host works best.
PLoS Negl Trop Dis. 2016;10(7):e0004794. doi: 10.1371/journal.pntd.0004794.
7.   Ross AG, Chau TN, Inobaya M, Olveda RM, Li Y, Harn DA. A new global strategy for the
elimination of schistosomiasis. Int J Infect Dis. 54:130-7.
8.   Hotez PJ, Kamath A. Neglected tropical diseases in Sub-Saharan Africa: Review of their
prevalence, distribution, and disease burden. Plos Neglected Tropical Diseases. 2009;3(8):e412. doi:
ARTN e412

DOI 10.1371/journal.pntd.0000412. PubMed PMID: ISI:000269220900003; PubMed Central PMCID: PMCPMC2727001.

9. Loker ES, Hofkin BV, Mkoji GM, Mungai B, Kihara J, Koech DK. Distributions of freshwater snails in southern Kenya with implications for the biological control of schistosomiasis and other snail-mediated parasites. Journal of Medical and Applied Malacology. 1993;5:1-20.

10. Brown DS. Freshwater Snails of Africa and Their Medical Importance. London: Taylor and Francis Limited; 1994.

11. Zhang SM, Bu L, Laidemitt MR, Lu L, Mutuku MW, Mkoji GM, et al. Complete mitochondrial and rDNA complex sequences of important vector species of *Biomphalaria*, obligatory hosts of the human-infecting blood fluke, *Schistosoma mansoni*. Sci Rep-Uk. 2018;8:7341. doi: doi: 10.1038/s41598-018-25463-z. PubMed Central PMCID: PMCPMCID: PMC5943310.

12. Standley CJ, Wade CM, Stothard JR. A fresh insight into transmission of schistosomiasis: a misleading tale of *Biomphalaria* in Lake Victoria. PLoS ONE. 2011;6:e26563. doi: doi:10.1371/journal.pone.0026563.

13. Odiere MR, Rawago FO, Ombok, M., Secor WE, Karanja DM, Mwinzi PN, et al. High prevalence of schistosomiasis in Mbita and its adjacent islands of Lake Victoria, western Kenya. . Parasites & Vectors. 2012;5:278.

14. Samuels AM, Matey E, Mwinzi PN, Wiegand RE, Muchiri G, Ireri E, et al. *Schistosoma mansoni* Morbidity among School-Aged Children: A SCORE Project in Kenya. American Journal of Tropical Medicine & Hygiene. 2012;87:874-82.

15. Wiegand RE, Mwinzi PNM, Montgomery SP, Chan YL, Andiego K, Omedo M, et al. A persistent hotspot of *Schistosoma mansoni* infection in a five-year randomized trial of praziquantel preventative chemotherapy strategies. J Infect Dis. 2017;216(11):1425-33. PubMed PMID: WOS:000417960300014.

16. Kristensen TK. A field guide to African freshwater snails. East African species. Second Edition ed: Danish Bilharziasis Laboratory; 1987. 51 p.

17. Frandsen F, Christensen NO. An introductory guide to the identification of cercariae from African freshwater snails with special reference to cercariae of trematode species of medical and veterinary importance. Acta Trop. 1984:181-202.

18. Laidemitt MR, Zawadzki ET, Brant SV, Mutuku MW, Mkoji GM, Loker ES. Loads of trematodes: discovering hidden diversity of paramphistomoids in Kenyan ruminants. Parasitology. 2017;144(2):131-47. PubMed PMID: WOS:000395068400003.

19. Mutuku MW, Dweni CK, Mwangi M, Kinuthia JM, Mwangi IN, Maina GM, et al. Field-derived *Schistosoma mansoni* and *Biomphalaria pfeifferi* in Kenya: a compatible association characterized by lack of strong local adaptation, and presence of some snails able to persistently produce cercariae for over a year. Parasite Vector. 2014;7. PubMed PMID: WOS:000345896700001.

20. K.Y. T, Moore DV, DeGiusti DL, Ripsom CA, Meleney HE. A technique for the perfusion of laboratory animals for the recovery of schistosomes. J Parasitol. 1947;33:419-26.

21. Crellen T, Walker M, Lamberton PHL, Kabatereine NB, Tukahebwa EM, Cotton JA, et al. Reduced efficacy of praziquantel against *Schistosoma mansoni* is associated with multiple rounds of mass drug administration. Clin Infect Dis. 2016;63(9):1151-9. PubMed PMID: WOS:000387986200006.

22. Magendantz M. Biology of *Biomphalaria choanomphala* and *B. sudanica* in relation to their role in transmission of *Schistosoma mansoni* in Lake Victoria at Mwanza, Tanzania. B World Health Organ. 1972;47(3):331-42. PubMed PMID: WOS:A1972O655800009.

23. Gouvras AN, Allan F, Kinung'hi S, Rabone M, Aidan E, Teckla A, et al. Longitudinal survey on the distribution of *Biomphalaria sudanica* and *B. choanomophala* in Mwanza region, on the shores of Lake Victoria, Tanzania: implications for schistosomiasis transmission and control. Parasite Vector. 2017;10:316.

24.   Kazibwe F, Makanga B, Rubaire-Akiiki C, Ouma J, Kariuki C, Kabatereine NB, et al. Ecology of
Biomphalaria (Gastropoda : Planorbidae) in Lake Albert, Western Uganda: snail distributions, infection
with schistosomes and temporal associations with environmental dynamics. Hydrobiologia.
2006;568:433-44. PubMed PMID: ISI:000239393800036.
25.   Radke MR, Ritchie LS, Rowan WB. Effects of water velocities on worm burdens of animals
exposed to *Schistosoma mansoni* cercariae released under laboratory and field conditions. Experimental
Parasitology. 1961;11:323-31.
26.   Upatham ES. Dispersion of St. Lucian *Schistosoma mansoni* cercariae in natural standing and
running waters determined by cercaria counts and mouse exposure. Ann Trop Med Parasit.
1974;68:343-52.
27.   Mutuku MW, Lu L, Otiato FO, Mwangi IN, Kinuthia JM, Maina GM, et al. A comparison of Kenyan
Biomphalaria pfeifferi and *B. sudanica* as vectors for *Schistosoma mansoni*, including a discussion of the
need to better understand the effects of snail breeding systems on transmission. J Parasitol. 2017:669-
76.
28.   Rowel C, Fred B, Betson M, Sousa-Figueiredo JC, Kabatereine NB. Environmental epidemiology
of intestinal schistosomiasis in Uganda : population dynamics of Biomphalaria (Gastropoda :
Planorbidae) in Lake Albert and Lake Victoria with observations on natural infections with digenetic
trematodes. Biomed Res Int. 2015;2015:Article ID 717261, 11 pages.
29.   Okeke OC, Ubachukwu PO. Trematode infections of the freshwater snail *Biomphalaria pfeifferi*
from a south-east Nigerian community with emphasis on cercariae of *Schistosoma*. J Helminthol.
2017;91:295-301.
30.   Nidal A, Mohammed I, Madsen H, Ahmed AAARM. Types of trematodes infecting freshwater
snails found in irrigation canals in the East Nile locality, Khartoum, Sudan. Infect Dis Poverty. 2016;5:16.
31.   Lim HK, Heyneman D. Intramolluscan inter-trematode antagonism: a review of factors
influencing the host-parasite system and its possible role in biological control. Advances in Parasitology.
1972;10:191-268.

Usenge beach
Police beach
KANYIBOK
USENGE
Anyanga beach
Ndiara beach

Daraja beach
MIYANDHE
MIGIRO
Kamumbo beach
Kadede beach
Kochilo beach
Uyawi beach
MINYA
Luthi beach

Mumbo beach
Kabuong beach
Kotieno beach
MUMBO
Maguti beach
Seka beach
Kogoye beach
Gudwa beach
SEKA DOK
AGOK
KOTIENO GUMBA
Alara beach
Kagwa beach
Kobala beach
ST. DOUGLAS WETA

🏠 Non Responding

🏠 Responding

10   0   10   20   30   40 km