Single-cell analysis reveals the *KIT* D816V mutation in hematopoietic stem and progenitor cells in systemic mastocytosis

Short title: Single-cell mutation analysis in mastocytosis

Jennine Grootens[1], Johanna S. Ungerstedt[2,3], Maria Ekoff[1], Elin Rönnberg Höckerlind[1], Monika Klimkowska[4], Rose-Marie Amini[5], Michel Arock[6], Stina Söderlund[7,8], Mattias Mattsson[5,7], Gunnar Nilsson[1,7,8], and Joakim S. Dahlin[1]

Author affiliations:
[1]Department of Medicine Solna, Karolinska Institutet, and Karolinska University Hospital, Stockholm, Sweden;
[2]Department of Medicine Huddinge, Karolinska Institutet, Stockholm, Sweden;
[3]Hematology Center, Karolinska University Hospital, Stockholm, Sweden;
[4]Department of Clinical Pathology and Cytology, Karolinska University Hospital Huddinge, Stockholm, Sweden;
[5]Department of Immunology, Genetics and Pathology, Uppsala University, Uppsala, Sweden;
[6]Molecular and Cellular Oncology, Ecole Normale Supérieure de Cachan, Cachan, France and Clinical Hematology Laboratory, Pitié-Salpêtrière Hospital, Paris, France;
[7]Division of Hematology, Uppsala University Hospital, Uppsala, Sweden;
[8]Department of Medical Sciences, Uppsala University, Uppsala, Sweden

Address correspondence to:
Gunnar Nilsson, Department of Medicine, Immunology and Allergy unit, Karolinska Institutet, SE-171 76 Stockholm, Sweden; e-mail: gunnar.p.nilsson@ki.se
or
Joakim Dahlin, Department of Medicine, Immunology and Allergy unit, Karolinska Institutet, SE-171 76 Stockholm, Sweden; e-mail: joakim.dahlin@ki.se

Abstract word count: 239

Text word count: 3604

Number of figures: 6

Number of tables: 1

35   Number of references: 37

37   Key points:

38   • Single-cell analysis reveals the *KIT* mutation in hematopoietic stem and

39   progenitor cells and its accumulation in mast cells of SM patients

40   • CD45RA is a specific marker for aberrant mast cells in SM

Abstract

Systemic mastocytosis (SM) is a hematological disease characterized by organ

infiltration by neoplastic mast cells. Almost all SM patients have a mutation in the

gene encoding the tyrosine kinase receptor KIT causing a D816V substitution and

autoactivation of the receptor. In this study, we analyzed the occurrence of the D816V

mutation during hematopoiesis to evaluate whether this mutation is an early or late

event in SM. Using a panel of 9 fluorescently labeled antibodies, 10 different

hematopoietic progenitor subsets and mature mast cells were identified. The

frequencies of the progenitor subsets did not differ between controls and SM patients.

Single-cell index sorting and D816V mutation assessment were applied to analyze

mast cells and more than 10,000 $CD34^+$ bone marrow progenitors from SM patients.

We found that in SM 60-99% of the mast cells harbored the mutation, and the

mutation could be detected throughout the hematopoietic landscape, from stem cells

to more lineage-primed progenitors. However, the occurrence of the mutation in stem

and progenitor cells was heterogeneous and varied between patients. Thus, the D816V

mutation appears to be an early event in the development of SM but accumulates in

the mast cell lineage. In addition, we demonstrate that mast cells originate from an

$Fc\varepsilon RI^+$ bone marrow progenitor, and we describe aberrant CD45RA expression on

SM mast cells for the first time. Taken together, our findings provide novel insight

into hematopoiesis and the occurrence of the *KIT* D816V mutation in SM.

Introduction

Hematopoietic stem cells (HSCs) in the bone marrow give rise to all blood cells, including mast cells. Differentiating HSCs progress through a number of intermediate progenitors with multilineage-forming capacity before commitment to the mast cell lineage.[1,2] The binding of stem cell factor (SCF) to its receptor, KIT, promotes the maturation and proliferation of mast cells.[3-6] It is therefore not surprising that mutations in the *KIT* gene coincide with the mast cell-driven disease systemic mastocytosis (SM).[7]

SM is a hematologic neoplasm in which infiltrates of neoplastic mast cells occur in various tissues.[8,9] The majority of SM patients carry a mutation in *KIT,* most commonly affecting codon 816.[10] The *KIT* D816V mutation makes receptor signaling constitutively active, independent of binding to its ligand SCF. The detection of *KIT* D816V in either bone marrow or peripheral blood samples is one of the criteria for the clinical diagnosis of SM using a standardized qPCR assay.[11] When the *KIT* D816V mutation is present in mast cells, it may also be detected in mature bone marrow or peripheral blood cells, such as basophils, eosinophils, neutrophils, and B and T lymphocytes, which depends on the patient.[12-17] Furthermore, the precursors of erythroid and myeloid cells as well as CD34$^+$ progenitors may carry the *KIT* D816V mutation.[12,18,19] Taken together, these findings suggest that the mutation may arise in early hematopoietic stem and progenitor cells (HSPCs), rather than in a lineage-restricted precursor.

The CD34$^+$ cell compartment is highly heterogeneous, spanning HSCs to late lineage-committed progenitors. In the present study, we delineated the cellular origin of the

KIT D816V mutation in single bone marrow cells in SM by combining FACS index

sorting of CD34[+] HSPCs with a multiplex qPCR assay. The data revealed that the

D816V mutation in *KIT* arises in early HSPCs. The mutation burden is low but

variable in multipotent and lineage-restricted progenitor populations and increases in

mature mast cells. Furthermore, the present study provides more insight into

hematopoiesis in SM subjects and reveals a novel marker for aberrant mast cells.

Patients, materials, and methods

Patients and sample preparation

Bone marrow and occasional peripheral blood samples were collected from patients

under evaluation for SM and from control subjects. The study was approved by the

Stockholm and Uppsala Regional Ethics committee. Oral and written informed

consent was obtained from each patient and control subject. The study was conducted

in accordance with the World Medical Association Declaration of Helsinki. The

patients included in our experimental studies were classified according to the World

Health Organization criteria[20] as having indolent SM (ISM, n = 19), aggressive SM

(ASM, n = 3), SM with associated hematological neoplasm (SM-AHN, n = 3), or

other malignancies, including myelodysplastic syndrome or myeloproliferative

neoplasm (MDS or MPN, n = 2). Table S1 shows the patient characteristics. For

analytic purposes, ASM and SM-AHN patients were combined in an advanced SM

group (AdvSM). The controls (Ctrl) included patients with cutaneous mastocytosis

(CM, n = 2) and subjects without SM (n = 7). All samples were processed on the day

of collection. Red blood cells were lysed with PharmLyse buffer (BD Biosciences,

Franklin Lakes, NJ) and washed with PBS containing 2% fetal calf serum (FCS,

Thermo Fisher Scientific, Waltham, MA).

Cell lines

Two human mast cell lines, HMC-1.2 and ROSA[KIT WT], with and without the *KIT*

D816V mutation, respectively, were cultured as previously described.[21,22]

Flow cytometry and cell sorting

Bone marrow cells were stained with fluorophore-conjugated monoclonal antibodies

in Brilliant Stain Buffer (BD Biosciences, Franklin Lakes, NJ). The antibodies used

for staining included CD10 (HI10a), CD14 (M5E2), CD34 (581), CD38 (Hb7),

CD45RA (HI100), CD90 (5E10), CD117 (104D2), CD123 (6H6), CD133 (AC133),

and FcεRIα (CRA-1) (all from BD Biosciences, Biolegend, San Diego, CA, or

Miltenyi Biotec, Bergisch Gladbach, Germany). The live/dead cell marker DAPI (BD

Bioscience) was used for the analysis of cultured cells. Cell sorting and analysis were

performed using the FACSAria III and LSRFortessa systems (BD Biosciences). For

cell culture experiments, cells were sorted into sterile-filtered PBS with 2% FCS

following a two-step sorting protocol, using the yield followed by the purity precision

mode. The analysis of 20-100 sorted cells verified the sorting purity. Single CD34[+]

cells were sorted into 2 μl of lysis buffer containing 20 μg/ml PCR-grade proteinase K

(Thermo Fisher Scientific) in EB buffer (Qiagen, Hilden, Germany). These cells were

sorted either immediately from the sample or following an enrichment sort. The

single-cell precision mode ensured that only one cell was deposited per well. The

plates were then centrifuged and stored at -80°C until analysis. The sorting data from

single cells were saved using the index sort module of the software FACSDIVA

software version 8 (BD Biosciences). Flow cytometry data analysis was performed

using FlowJo software version 10 (TreeStar, Ashland, OR).

Cell culture and colony-forming assays

Colony-forming potential was evaluated by plating 500 cells in 35 x 10 mm tissue

culture dishes (Thermo Fisher Scientific) in duplicate. The cells were cultured in

Methocult H4434 (STEMCELL Technologies Inc., Vancouver, Canada). Colonies

were counted after 12-13 days, distinguishing erythroid, granulocyte/monocyte, and

mixed colonies following the manufacturer's recommendations. The mast cell-

forming potential was assessed via liquid culture as described previously.[23] Briefly,

sorted cells were cultured in medium with recombinant human (rh) IL-3 (10 ng/mL;

PeproTech, Rocky Hill, NJ) and rhIL-6 (10 ng/mL; PeproTech). In some experiments,

the medium was changed on day 5, and the cells were subsequently cultured with

rhIL-6 (10 ng/mL) and SCF (100 ng/mL, r-metHuSCF; Swedish Orphan Biovitrum,

Stockholm, Sweden), as indicated in the figure legends. Enzyme cytochemical

staining of trypsin-like activity was used to assess tryptase expression.[24,25] Cell

morphology was evaluated by May-Grünwald Giemsa staining (MGG; Sigma-

Aldrich, St. Louis, MO).

Single-cell mutation analysis

Before the addition of lysis buffer, 96-well plates (Bio-Rad Laboratories, Hercules,

CA) were UV-treated for 30 minutes using a UVT-B-AR UV Cabinet (Grant

Instruments, Cambridge, UK). Tubes and pipettes were treated with the DNA AWAY

Surface Decontaminant (Thermo Fisher Scientific). After sorting, the cells were lysed

for 45 minutes at 55°C, followed by a 10-min heat-inactivating step at 95°C.

Multiplex qPCR analysis was performed via a Taqman gene expression assay

(Thermo Fisher Scientific) in lysed single cells using 96-well CFX Manager (Bio-Rad

Laboratories). A mutation assay was designed based on a previously published

method[11] using Primer3 (v.0.4.0) which specifically detected *KIT* D816V using the

forward primer 5'-AGAGACTTGGCAGCCAGAAA-3'; a reverse primer matching

the mutation at the 3' position, 5'-

TTAACCACATAATTAGAATCATTCTTGATGA-3'; and the probe 5'-FAM-

TCCTCCTTACTCATGGTCGGATCACA-BHQ1-3'. A control assay was designed

to target a flanking exon and intron eight in GAPDH using the forward primer 5'-

CTGACTTCAACAGCGACACC-3'; the reverse primer 5'-

AGAGTTGTCAGGGCCCTTTT-3'; and the probe 5'-HEX-

TCAAGCTCATTTCCTGGTATGTG-BHQ1-3' (all primers came from TAG

Copenhagen, Denmark). The control and mutation assays were tested and optimized

using the cell lines ROSA[KIT WT] and HMC-1.2. The primer concentrations, optimized

for single cells, were as follows; control assay 150 nM forward/reverse primers, 100

nM probe; mutation assay 100 nM forward/reverse primers, 67 nM probe. Each plate,

containing 86 CD34[+] cells, 5 mature mast cells, 3 empty wells, and 2 wells with bulk

CD34[+] cells following sorting, was subjected to the multiplex qPCR assay. Cells were

excluded from analysis when amplification failed in the control assay, or when

amplification of the control assay resulted in a CT value higher than 40. In the

mutation assay, cells were regarded as unmutated when there was no amplification or

when the CT value was higher than 45. Outliers from each plate were then detected

using the ROUT outlier test in Prism software and removed from further analyses

(Figure S5B). Plates and cells that passed the quality control procedure were linked

with the index data and gated similarly to the whole sample for analysis.

Statistical analysis

Statistical analysis was performed using Prism software (Version 6.0h, GraphPad

Software, La Jolla, CA). The unpaired two-tailed Student's t-test was employed when

comparing two groups, and the unpaired one-way ANOVA with Tukey's multiple

comparison test was employed when comparing more than two groups. Statistical

analysis of log-transformed data was performed for some experiments as indicated in

the figure legends. Differences were considered significant when $P < .05$.

For original data, please contact joakim.dahlin@ki.se

Results

Systemic mastocytosis patients have a normal CD34$^+$ bone marrow stem and

progenitor cell composition

To assess the composition of the hematopoietic cells in the bone marrow of SM

patients, we designed a multicolor flow cytometry panel that identifies ten distinct

HSPC subtypes and mature mast cells (Table 1; Figures 1A-B). Colony formation

assays confirmed the validity of the GMP, CMP, and MEP gating strategies in SM

subjects (Figure S1A-C). The flow cytometry data provided a specific HSPC profile

for 34 bone marrow samples: 9 control, 19 ISM and 6 AdvSM (Table S1; Figure 1C).

The percentages of HSPCs in SM samples did not significantly differ from those in

control samples (Figures 1D and S2A). In addition, the ISM and AdvSM samples

exhibited percentages of total CD34$^+$ cells similar to the control samples (Figure

S2B). Thus, the overall hematopoiesis profile did not differ between the control and

SM subjects.

The bone marrow CMP$^{Fc\varepsilon RI+}$ fraction presents an enhanced mast cell-forming

potential

Mast cell progenitors circulate in the blood as Lin$^-$ CD34^{hi} CD117$^{int/hi}$ FcεRI$^+$ cells.[33]

These cells express CD123 and lack CD45RA,[23,33] overlaying the CMP$^{Fc\varepsilon RI+}$ gate in

bone marrow (Figure S3A). We therefore analyzed the mast cell-forming potential of

the bone marrow CMP$^{Fc\varepsilon RI+}$ fraction by culturing the cells with IL-3 and IL-6. Flow

cytometry analysis revealed that the CMP$^{Fc\varepsilon RI+}$ cultures gave rise to a higher

frequency of CD117^{hi} FcεRI$^+$ mast cells than the CMPs$^{Fc\varepsilon RI-}$ cells (Figures 2A-C),

despite overall lower CD117 expression (Figure S4A). May-Grünwald Giemsa and

enzymatic tryptase staining showed that the CMP$^{Fc\varepsilon RI+}$ progeny generated a high

frequency of granulated tryptase-positive mast cells when cultured with SCF (Figures

2D-E and S3B). Cultured GMPs did not give rise to any cells with a mast cell

phenotype (Figure S3C).

The emerging model of hematopoiesis proposes that MEPs, basophils, and

eosinophils develop separately from GMPs following an asymmetric cell division at

the MPP stage. This process can be traced by the segregation of CD133, which is

expressed in GMPs but not in progenitors with erythromyeloid output.[34] Indeed, we

found that CMP$^{Fc\varepsilon RI+}$ cells lacked CD133 staining, similar to erythromyeloid cells

(Figure S4B).

CD45RA is expressed on aberrant mast cells in systemic mastocytosis

The multicolor flow cytometry panel used for the gating of HSPCs was also employed

to identify mast cells in bone marrow (Figure 1B). May-Grünwald Giemsa and

tryptase staining confirmed that CD14⁻ CD34⁻ CD117^{hi} FcεRI^{hi} cells constitute mast

cells (Figure 3A). SM samples presented higher percentages of mast cells than

controls (Figure 3B). Notably, mast cells from almost all SM samples exhibited

aberrant CD45RA expression, as verified with fluorescence minus one control and

internal control staining of CD34^{+} cells (Figures 3C-D). The cutaneous (non-

systemic) mastocytosis samples and MDS/MPN samples lacked expression of

CD45RA on bone marrow mast cells (Figures 3C-D). In addition, mast cell

frequencies were correlated with the clinical percentage of BMMC infiltrate and

serum tryptase levels (Figures 4A-C; Table S2). Taken together, the results indicate

that SM bone marrow samples contain more mast cells than control samples and these

mast cells express the aberrant marker CD45RA.

Single-cell index sorting reveals the KIT D816V mutation in early HSPCs in systemic

mastocytosis patients

The cellular origin of the *KIT* D816V mutation in SM has not been established, and

the relative abundance of this mutation in different CD34^{+} HSPC populations and

mast cells has not been determined. We therefore developed a single-cell method to

trace the *KIT* D816V mutation using two cell lines: HMC-1.2 cells carrying the

mutation and ROSA cells carrying wild-type *KIT*. The cells were labeled with two

different fluorochrome-labeled antibodies, mixed, and randomly FACS sorted with

the index sort function. A multiplex qPCR assay was used to detect an irrelevant gene,

for quality control purposes, and the *KIT* D816V mutation (Figure S5A-B). Among all

single cells, more than 96% passed the quality control procedure (Figure S5C).

Linking the results with the index data revealed that the mutation assay was negative

for the wild-type ROSA cells. The assay detected the mutation in 98.1% of all HMC-

1.2 cells. Thus, the assay did not detect false positive cells in the mutation assay and

only a few false negatives, demonstrating the high sensitivity and specificity of the

method (Figure S5D).

The high success rate with cell lines allowed us to use the assay for HSPCs and

mature mast cells from patient samples. Single cells from 12 samples (6 ISM and 6

AdvSM) were analyzed for mutation assessment. Specifically, CD34$^+$ HSPCs were

index sorted from ten patients, whereas specific HSPC subsets were sorted from two

patients. At the time of SM diagnosis, the presence of the *KIT* D816V mutation was

confirmed in either peripheral blood or bone marrow (Table S1). We analyzed 40-194

single mast cells and 351-1583 single CD34$^+$ cells from each bone marrow sample,

providing mutation data for 12,859 single cells in total (Figure 5A; Table S3). All SM

samples carried the *KIT* D816V mutation in the mast cell compartment at a high

frequency in ISM (mean, 80.9%) and AdvSM (mean, 93.2%) (Figure 5B). The

percentage of mutated mast cells was correlated with the total percentage of mast cells

in the sample and with the serum tryptase levels (Figure 6A-B; Table S2). The

mutation rate in CD34$^+$ HSPCs was less than 2.7% in all samples (Figure 5C), except

for one advanced SM sample with a high mutation burden of 37.0% (Table S4).

Nevertheless, the percentage of mutated CD34$^+$ HSPCs was correlated with serum

tryptase levels (Figure 6C; Table S2).

Coupling the index data with the mutation analysis of 10 bone marrow samples

revealed that the *KIT* D816V mutation appeared throughout the hematopoietic

landscape (Figure 5D; Table S3-4). Specifically, different HSPCs presented similar

mutation rates in all samples when the median percentage of mutation was compared

between different HSPCs (Figure 5E). Overall, the method of index sorting provided a

representative set of single cells from the whole-bone-marrow aspirates (Figure S6A).

However, the CMP$^{Fc\varepsilon RI+}$ population with an enriched mast cell potential was rare. To

determine the mutation frequency in the CMP$^{Fc\varepsilon RI+}$ fraction, we selectively sorted 86

and 278 single CMPs$^{Fc\varepsilon RI+}$ from 2 SM samples. One patient exhibited less than 2%

mutated CMPs$^{Fc\varepsilon RI+}$, HSCs and GMPs, whereas the fraction of mutated mast cells was

98.9% (Figure 5F; Table S4). The second patient presented 95.7% mutated

CMPs$^{Fc\varepsilon RI+}$ and 96.4 % mutated mast cells. (Figure 5G; Table S4). Peripheral blood

MCPs (CD14$^-$ CD34$^+$ CD117$^+$ FcεRI$^+$) showed mutation rates comparable to that of

bone marrow CMPs$^{Fc\varepsilon RI+}$ in one patient (Figure S6B), but the mutation could not be

detected in the MCPs from 4 other SM samples, likely due to the low mutation

frequency and the small number of cells analyzed due to the rarity of the cells (14-23

cells analyzed per sample). Taken together, we detected the *KIT* D816V mutation

throughout the hematopoietic landscape, starting with the HSCs.

Discussion

The cellular origin of the aberrant clone in SM is still under debate. Here, we show

that SM patients exhibit an increased frequency of mast cells in bone marrow. Almost

all these mast cells carry the *KIT* D816V mutation. Despite normal HSPC frequencies

in bone marrow, the *KIT* D816V mutation can be traced all the way back to the

CD34$^+$ CD38$^-$ CD10$^-$ CD45RA$^-$ CD90$^+$ HSC fraction.

The *KIT* D816V mutation causes receptor autoactivation, which signals survival,

proliferation, and differentiation, likely explaining the competitive growth advantage

of the aberrant mast cell clone and the high frequency of mutated mast cells.

However, the frequency of mutated HSPCs was low in most patients. This

observation is consistent with the overall normal HSPC composition of SM patients.

There are two main potential explanations for the low D816V mutation burden

observed in HSPCs and its accumulation in mast cells: HSPCs rapidly differentiate

into mast cells when the mutation occurs, or mutated HSPCs do not present a strong

growth advantage over non-mutated HSPCs. In fact, we have previously shown that

KIT signaling is dispensable for mast cell progenitor development, and that other

stimuli can induce cell proliferation.[23] However, the aberrant progenitor clone may in

some cases proliferate to reach a high frequency, as demonstrated in two patients with

advanced SM who presented more than 20% mutated cells in the HSC fraction.

Advanced forms of SM have previously been associated with detection of the *KIT*

mutation in bulk-sorted CD34[+] progenitors and various mature cell lineages, whereas

in indolent disease, the mutation is mainly restricted to mast cells.[13]

It is of particular interest to study the mutation burden in progenitors with mast cell-

forming potential. We have previously reported that CD34[+] CD117[int/hi] FcεRI[+] cells in

peripheral blood are committed or nearly committed MCPs.[33] Considering the CD34[+]

CD117[+] FcεRI[+] phenotype of CMPs[FcεRI+] cells, these cells might be expected to

present a similar mast cell-forming capacity as to blood MCPs. In the current study,

we showed that CMPs[FcεRI+] gave rise to a higher frequency of mast cells than

CMPs[FcεRI-] and GMPs. However, the mast cell frequency in cultured CMPs[FcεRI+] was

less than 10% and a substantial fraction were found to be CD117[-/lo] FcεRI[+] cells after

culture, consistent with a basophil-like phenotype. Thus, CMPs$^{Fc\epsilon RI+}$ exhibit a

considerable mast cell-forming capacity but also include constitute progenitors with

non-mast cell output. Basophils mature in bone marrow, whereas mast cells mature in

peripheral tissues, which likely explains why bone marrow CMPs$^{Fc\epsilon RI+}$ form CD117$^-$

FcϵRI$^+$ basophil-like cells, whereas CD34$^+$ CD117$^{int/hi}$ FcϵRI$^+$ blood progenitors do

not. The high frequency of CD117$^-$ FcϵRI$^+$ basophil-like cells that develop from

CMPs$^{Fc\epsilon RI+}$ is also in agreement with the reduced CD117 expression of CMPs$^{Fc\epsilon RI+}$

compared with that of CMPs$^{Fc\epsilon RI-}$. CD117 is upregulated upon mast cell maturation,

and blood mast cell progenitors express intermediate to high levels of CD117.

CMPs$^{Fc\epsilon RI+}$ express CD117 at low levels overall, even though there are individual

CMPs$^{Fc\epsilon RI+}$ cells that express high CD117 levels, which might correspond to the cells

with mast cell-forming capacity. Taken together, our results indicate that the low *KIT*

mutation prevalence in the CMP$^{Fc\epsilon RI+}$ fraction is likely explained by the capacity of

these cells' to form not only mast cells but also CD117$^-$ FcϵRI$^+$ basophil-like cells.

In SM, mast cells are known to aberrantly express multiple surface proteins, among

CD2 and CD25 are currently used for diagnosis.[35] These markers are expressed

independently of the SM subtype. Other markers, such as CD30 and CD52, are

associated with advanced SM and mast cell leukemia.[36,37] In the present study, we

demonstrate that CD45RA is a novel marker for aberrant mast cells in SM.

Remarkably, bone marrow mast cells from two patients with cutaneous mastocytosis

did not express CD45RA, suggesting that CD45RA is a specific marker for systemic

mast cell disease. Taken together, our results indicate that CD45RA expression should

be considered a novel marker for diagnosis of SM.

CD45RA is present on aberrant mast cells in SM, but at what stage during mast cell

differentiation is the marker upregulated? CMPs$^{Fc\epsilon RI+}$, which lack CD45RA, present

mast cell-forming potential in SM patients. In contrast, we show that cells in the

classic CD45RA$^+$ GMP gate lack mast cell potential in SM patients. An alternative

explanation is that the mast cell progenitors in SM fall into the immature CD38$^-$ gate.

In fact, examples of rare mutated CD117$^{int/hi}$ FcϵRI$^{int/hi}$ cells in the CD45RA$^+$ LMPP

gate were seen (Figure S6C). It is also tempting to speculate that CD45RA is

upregulated once aberrant mast cells are mature.

The CD34$^+$ HSPC population is highly heterogeneous and a single-cell resolution is

therefore necessary to delineate the cellular distribution of the mutation. Here, we

used single-cell index sorting to track cellular identity when performing mutation

analysis. Notably, the mutation was detected throughout differentiation, from HSCs to

the MEP, GMP, and LMPP fractions, depending on the patient. The detection of the

mutation in early HSPCs explains previous findings of the mutation being present in

bulk-sorted mature cell lineages of some patients.

Multiplex TaqMan qPCR assays detected the *KIT* mutation in index-sorted

progenitors in the present investigation. The qPCR method was originally developed

for sensitive mutation detection in DNA extracted from bone marrow or blood cells.

GAPDH primers and probes were added to the assay to confirm the presence or

absence of a sorted cell. This method allowed the detection of mutated cells with

close to 100% efficiency, with no observed false positives. The single-cell index

sorting and DNA mutation analysis approach can similarly be applied to other

mutations and hematologic diseases to reveal the origin of aberrant clones. One

limitation of the method is the number of cells that can be analyzed, meaning that

mutated cells in very rare subpopulations might go undetected. However, more than

10,000 HSPCs were analyzed in the present investigation, and single-cell index

sorting coupled with qPCR allowed unbiased and prospective analysis of mutated

HSPC populations in SM. Whole-genome amplification and nested PCR methods for

amplifying and sequencing DNA mutations from single cells are associated with the

phenomenon known as allelic dropout, which is the amplification of only a single

allele, leading to an overestimation of the frequency of non-mutated cells. Mutational

profiling of single colonies, as performed by Jawhar *et al*[18], avoids this problem.

However, such analysis can be performed only on progenitors that form sufficiently

large colonies. Mast cell progenitors present a poor proliferative capacity, and this

technique was therefore not applied in the present investigation.

In summary, we have explored the composition and mutation profile of $CD34^+$ HSPCs

from SM patients. The results show that the *KIT* mutation can be found in cells

throughout the hematopoietic landscape and exhibits accumulation in the mast cell

lineage, supporting the notion that the aberrant clone may arise in the hematopoietic

stem cell compartment. Furthermore, the mutated mast cells express CD45RA, a

potential novel clinical biomarker for aberrant mast cells in SM.

Acknowledgements

We wish to thank Kerstin Hamberg Levedahl for help with the patients. Stem cell

factor was a kind gift from SOBI, Stockholm, Sweden. This study was supported by

the Swedish Research Council, the Swedish Cancer Society, the Stockholm Cancer

Society (Radiumhemmets Forskningsfonder), Magnus Bergvall's Foundation, Tore

Nilsson's Foundation for Medical Research, the Karolinska Institutet, and through the

regional agreement on medical training and clinical research (ALF) between

Stockholm County Council and the Karolinska Institutet, and ALF-funding from

Uppsala University Hospital.

415   Authorship

J.G., J.S.D. and G.N. designed the experiments; J.G., E.R.H. and J.S.D. performed the

experiments; J.G. and M.E. processed the patient samples. J.S.U., M.K., R.M.A., S.S.

and M.M. provided patient samples and diagnosed patients; M.A. provided the ROSA

cell line. J.G. and J.S.D. analyzed the data and wrote the first draft of the paper; and

all authors reviewed the manuscript; and J.S.D. and G.N. designed and supervised the

study and secured funding.

Conflict-of-interest disclosure: The authors have declared that no conflict of interest

exists.

References

1.   Dahlin JS, Hallgren J. Mast cell progenitors: origin, development and
migration to tissues. *Mol Immunol*. 2015;63(1):9-17.

2.   Grootens J, Ungerstedt JS, Nilsson G, Dahlin JS. Deciphering the
differentiation trajectory from hematopoietic stem cells to mast cells. *Blood Advances*.
In press.

3.   Irani AM, Nilsson G, Miettinen U, et al. Recombinant human stem cell factor
stimulates differentiation of mast cells from dispersed human fetal liver cells. *Blood*.
1992;80(12):3009-3021.

4.   Kirshenbaum AS, Goff JP, Kessler SW, Mican JM, Zsebo KM, Metcalfe DD.
Effect of IL-3 and stem cell factor on the appearance of human basophils and mast
cells from CD34+ pluripotent progenitor cells. *J Immunol*. 1992;148(3):772-777.

5.   Valent P, Spanblochl E, Sperr WR, et al. Induction of differentiation of human
mast cells from bone marrow and peripheral blood mononuclear cells by recombinant
human stem cell factor/kit-ligand in long-term culture. *Blood*. 1992;80(9):2237-2245.

6.   Mitsui H, Furitsu T, Dvorak AM, et al. Development of human mast cells
from umbilical cord blood cells by recombinant human and murine c-kit ligand. *Proc*
Natl Acad Sci U S A. 1993;90(2):735-739.

7.   Nagata H, Worobec AS, Oh CK, et al. Identification of a Point Mutation in the
Catalytic Domain of the Protooncogene C-Kit in Peripheral-Blood Mononuclear-Cells
of Patients Who Have Mastocytosis with an Associated Hematologic Disorder. *Proc*
Natl Acad Sci U S A. 1995;92(23):10560-10564.

8.   Valent P, Akin C, Escribano L, et al. Standards and standardization in

mastocytosis: consensus statements on diagnostics, treatment recommendations and

response criteria. *Eur J Clin Invest*. 2007;37(6):435-453.

9.   Valent P, Akin C, Metcalfe DD. Mastocytosis: 2016 updated WHO

classification and novel emerging treatment concepts. *Blood*. 2017;129(11):1420-

1427.

10.   Arock M, Sotlar K, Akin C, et al. KIT mutation analysis in mast cell

neoplasms: recommendations of the European Competence Network on Mastocytosis.

Leukemia. 2015;29(6):1223-1232.

11.   Kristensen T, Vestergaard H, Moller MB. Improved detection of the KIT

D816V mutation in patients with systemic mastocytosis using a quantitative and

highly sensitive real-time qPCR assay. *J Mol Diagn*. 2011;13(2):180-188.

12.   Afonja O, Amorosi E, Ashman L, Takeshita K. Multilineage involvement and

erythropoietin-"independent" erythroid progenitor cells in a patient with systemic

mastocytosis. *Ann Hematol*. 1998;77(4):183-186.

13.   Garcia-Montero AC, Jara-Acevedo M, Teodosio C, et al. KIT mutation in

mast cells and other bone marrow hematopoietic cell lineages in systemic mast cell

disorders: a prospective study of the Spanish Network on Mastocytosis (REMA) in a

series of 113 patients. *Blood*. 2006;108(7):2366-2372.

14.   Kocabas CN, Yavuz AS, Lipsky PE, Metcalfe DD, Akin C. Analysis of the

lineage relationship between mast cells and basophils using the c-kit D816V mutation

as a biologic signature. *J Allergy Clin Immunol*. 2005;115(6):1155-1161.

15.   Yavuz AS, Lipsky PE, Yavuz S, Metcalfe DD, Akin C. Evidence for the

involvement of a hematopoietic progenitor cell in systemic mastocytosis from single-

cell analysis of mutations in the c-kit gene. *Blood*. 2002;100(2):661-665.

16.   Pardanani A, Reeder T, Li CY, Tefferi A. Eosinophils are derived from the

neoplastic clone in patients with systemic mastocytosis and eosinophilia. *Leuk Res*.

2003;27(10):883-885.

17.   Mayado A, Teodosio C, Dasilva-Freire N, et al. Characterization of CD34(+)

hematopoietic cells in systemic mastocytosis: Potential role in disease dissemination.

Allergy. 2018;73(6):1294-1304.

18.   Jawhar M, Schwaab J, Schnittger S, et al. Molecular profiling of myeloid

progenitor cells in multi-mutated advanced systemic mastocytosis identifies KIT

D816V as a distinct and late event. *Leukemia*. 2015;29(5):1115-1122.

19.   Jara-Acevedo M, Teodosio C, Sanchez-Munoz L, et al. Detection of the KIT

D816V mutation in peripheral blood of systemic mastocytosis: diagnostic

implications. *Mod Pathol*. 2015;28(8):1138-1149.

20.   Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health

Organization classification of myeloid neoplasms and acute leukemia. *Blood*.

2016;127(20):2391-2405.

21.   Saleh R, Wedeh G, Herrmann H, et al. A new human mast cell line expressing

a functional IgE receptor converts to tumorigenic growth by KIT D816V transfection.

Blood. 2014;124(1):111-120.

22.   Sundstrom M, Vliagoftis H, Karlberg P, et al. Functional and phenotypic

studies of two variants of a human mast cell line with a distinct set of mutations in the

c-kit proto-oncogene. *Immunology*. 2003;108(1):89-97.

23.   Dahlin JS, Ekoff M, Grootens J, et al. KIT signaling is dispensable for human

mast cell progenitor development. *Blood*. 2017;130(16):1785-1794.

24.   Harvima IT, Naukkarinen A, Harvima RJ, Fraki JE. Immunoperoxidase and

enzyme-histochemical demonstration of human skin tryptase in cutaneous mast cells

in normal and mastocytoma skin. *Arch Dermatol Res*. 1988;280(6):363-370.

25.   Dahlin JS, Ungerstedt JS, Grootens J, et al. Detection of circulating mast cells

in advanced systemic mastocytosis. *Leukemia*. 2016;30(9):1953-1956.

26.   Notta F, Doulatov S, Laurenti E, Poeppl A, Jurisica I, Dick JE. Isolation of

Single Human Hematopoietic Stem Cells Capable of Long-Term Multilineage

Engraftment. *Science*. 2011;333(6039):218-221.

27.   Majeti R, Park CY, Weissman IL. Identification of a hierarchy of multipotent

hematopoietic progenitors in human cord blood. *Cell Stem Cell*. 2007;1(6):635-645.

28.   Manz MG, Miyamoto T, Akashi K, Weissman IL. Prospective isolation of

human clonogenic common myeloid progenitors. *Proc Natl Acad Sci U S A*.

2002;99(18):11872-11877.

29.   Goardon N, Marchi E, Atzberger A, et al. Coexistence of LMPP-like and

GMP-like leukemia stem cells in acute myeloid leukemia. *Cancer Cell*.

2011;19(1):138-152.

30.   Lee J, Breton G, Oliveira TY, et al. Restricted dendritic cell and monocyte

progenitors in human cord blood and bone marrow. *J Exp Med*. 2015;212(3):385-399.

31.   Olweus J, BitMansour A, Warnke R, et al. Dendritic cell ontogeny: a human

dendritic cell lineage of myeloid origin. *Proc Natl Acad Sci U S A*.

1997;94(23):12551-12556.

32.   Doulatov S, Notta F, Eppert K, Nguyen LT, Ohashi PS, Dick JE. Revised map

of the human progenitor hierarchy shows the origin of macrophages and dendritic

cells in early lymphoid development. *Nat Immunol*. 2010;11(7):585-593.

33.   Dahlin JS, Malinovschi A, Ohrvik H, et al. Lin- CD34hi CD117int/hi

FcepsilonRI+ cells in human blood constitute a rare population of mast cell

progenitors. *Blood*. 2016;127(4):383-391.

34.   Gorgens A, Radtke S, Mollmann M, et al. Revision of the human

hematopoietic tree: granulocyte subtypes derive from distinct hematopoietic lineages.

Cell Rep. 2013;3(5):1539-1552.

35.   Teodosio C, Mayado A, Sanchez-Munoz L, et al. The immunophenotype of

mast cells and its utility in the diagnostic work-up of systemic mastocytosis. *J Leukoc*

Biol. 2015;97(1):49-59.

36.   Sotlar K, Cerny-Reiterer S, Petat-Dutter K, et al. Aberrant expression of CD30

in neoplastic mast cells in high-grade mastocytosis. *Mod Pathol*. 2011;24(4):585-595.

37.   Hoermann G, Blatt K, Greiner G, et al. CD52 is a molecular target in

advanced systemic mastocytosis. *FASEB J*. 2014;28(8):3540-3551.

Table legend

Table 1. Immunophenotypes of HSPCs

Figure legends

Figure 1. HSPC gating and profiling shows a normal course of hematopoiesis in patients with systemic mastocytosis. (A) A landscape model of the hematopoiesis. (B) Gating strategy for bone marrow HSPCs and mature mast cells, showing representative plots with population frequencies as percentages. (C) The hematopoietic progenitor profile of 34 bone marrow samples. Cutaneous mastocytosis samples are indicated in italics. (D) SM samples were compared with control samples without SM, including CM (open circles), and the mean expression + SEM of each HSPC is presented as a percentage of the total $CD34^+$ cells. The unpaired t-test was used in panel D to compare the control and SM groups for each HSPC subset.

Figure 2. The CMP$^{Fc\varepsilon RI+}$ fraction in the bone marrow has mast cell potential. (A-C) Flow cytometry analysis of cells cultured with rhIL-3 and rhIL-6 up to 21 days. (A) Flow cytometry plots of representative cultures, showing the percentages of $CD117^-$, $CD117^+$, and $CD117^{hi}$ cells in each plot. (B) Percentage of $CD117^{hi}$ $Fc\varepsilon RI^+$ cells at days 5-6 (n = 5). (C) Two of the samples in panel B were followed for up to 21 days under culture with rhIL-3 and rhIL-6. The percentage of $CD117^{hi}$ cells was analyzed on days 5-6, 13-14, and 19-21. (D-E) Cells stained for enzymatic tryptase and MGG after culturing with rhIL-3 and rhIL-6 for 5 days, followed by rhIL-6 and SCF up to day 21 (n = 5). (D) Quantification of tryptase-positive cells at days 13-14 and 17-21 (n = 5). (E) Representative images of cultured CMP$^{Fc\varepsilon RI-}$ and CMP$^{Fc\varepsilon RI+}$ cells after 13-14 days, showing tryptase staining (red) and granules by May-Grünwald

Giemsa (MGG) staining. Samples Ctrl29, SM28, SM32, SM33, and MDS30 were

analyzed in panels B, D, and E. Samples SM32 and SM33 were analyzed in panels A

and C. The bars and lines in panel B, C and D represent the means ± SEM. The paired

t-test was used in panel B. **$P < .01$. Images were captured using an Olympus XC10

camera (Olympus, Tokyo, Japan). The image width corresponds to 29 μm.

Figure 3. SM mast cells aberrantly express CD45RA. (A) Enzymatic staining

demonstrates tryptase expression, and MGG staining reveals granules in SM mast

cells. The images are representative of two experiments (samples SM32 and SM33).

Images were captured using an Olympus XC10 camera (Olympus, Tokyo, Japan). The

image width corresponds to 29 μm. (B) Mast cell frequencies in Ctrl, ISM, AdvSM,

and MDS/MPN patients, showing the median of log-transformed values. (C)

Histograms showing CD45RA expression in mast cells (red line), FMO controls (grey

histogram) and CD34$^+$ cells (dotted line). (D) Frequency of CD45RA-expressing mast

cells, showing the mean and SEM. Control subjects included two subjects with CM

(open circles). One-way ANOVA with Tukey's multiple comparison was used in

panels B and D, *$P < .05$, **$P < .01$, ****$P < .0001$.

Figure 4. The percentage of mast cells in bone marrow correlates with clinical

features. The percentage of mast cells was correlated with (A) serum tryptase levels

and (B) the percentage of BMMC infiltrate. (C) Serum tryptase levels were correlated

with the percentage of BMMC infiltrate. The samples include ISM (closed dots) and

AdvSM (open dots). Correlation values and significance are indicated in the figure

and in Table S2. Linear regression was calculated based on log-transformed data: in

panel A, n = 25, and in panels B and C, n = 23. The percentage of BMMC infiltrate

was not determined for two patients, accounting for the difference in numbers

between panels. For analytical purposes, BMMC infiltrate presented as <1% or <5%

in Table S1 were set at 1% and 5% respectively.

Figure 5. Mutation analysis of KIT D816V in single cells identifies mutated

HSPCs and mast cells. (A) Method of single-cell mutation analysis using index

sorting and multiplex qPCR. (B-C) The bars indicate the means + SEM percentage of

mutated cells for (B) mast cells and (C) HSPCs. (D) The cellular distribution of the

mutation in 10 different HSPCs, showing the total numbers of mutated and sorted

cells for each sample that passed the quality control procedure. (E) The bars indicate

the median percentage of mutated cells per HSPC. (F-G) Mutation frequencies of

selectively isolated HSPCs following a 2-step sorting protocol in two samples (SM28,

ISM; and SM31, AdvSM). These two samples are indicated with open circles in

panels B and E.

Figure 6. The mutation rate in mast cells and HSPCs correlates with clinical

features. (A-C) Linear regression of the percentage of mutated mast cells, showing

significant correlations with (A) the percentage of mast cells and (B) serum tryptase

levels. (C) The percentage of mutated HSPCs was correlated with serum tryptase

levels. The samples included ISM (closed dots) and AdvSM (open dots). Correlation

and significance are indicated in the figure and in Table S2. Linear regression was

calculated based on log-transformed data in panels A-C, n = 12 and D, n = 10.

Table 1. Immunophenotypes of hematopoietic stem and progenitor cells

Short	Cell type	Gating strategy	Reference
HSC	Hematopoietic stem cell	$CD14^-$ $CD34^+$ $CD38^-$ $CD45RA^-$ $CD90^+$	26, 27
MPP	Multipotent progenitor	$CD14^-$ $CD34^+$ $CD38^-$ $CD45RA^-$ $CD90^-$	26, 27
$CMP^{Fc\varepsilon RI-}$	Common myeloid progenitor $Fc\varepsilon RI^-$	$CD14^-$ $CD34^+$ $CD38^+$ $CD45RA^-$ $CD10^-$ $CD123^+$ $Fc\varepsilon RI^-$	28 *
$CMP^{Fc\varepsilon RI+}$	Common myeloid progenitor $Fc\varepsilon RI^+$	$CD14^-$ $CD34^+$ $CD38^+$ $CD45RA^-$ $CD10^-$ $CD123^+$ $Fc\varepsilon RI^+$ $CD117^+$	28 *
MEP	Megakaryocyte-erythroid progenitor	$CD14^-$ $CD34^+$ $CD38^+$ $CD45RA^-$ $CD10^-$ $CD123^-$	28
LMPP	Lymphoid-primed multipotent progenitor	$CD14^-$ $CD34^+$ $CD38^-$ $CD45RA^+$	29
GMP	Granulocyte-monocyte progenitor	$CD14^-$ $CD34^+$ $CD38^+$ $CD45RA^+$ $CD10^-$ $CD123^+$	28
CDP	Common dendritic progenitor	$CD14^-$ $CD34^+$ $CD38^+$ $CD45RA^+$ $CD10^-$ $CD123^{hi}$	30, 31
MLP	Multilymphoid progenitor	$CD14^-$ $CD34^+$ $CD38^-$ $CD45RA^+$ $CD10^+$	32
BNKP	B- and NK cell progenitor	$CD14^-$ $CD34^+$ $CD38^+$ $CD45RA^+$ $CD10^+$	32
MC	Mast cell	$CD14^-$ $CD34^-$ $CD117^{hi}$ $Fc\varepsilon RI^+$	

* Refers to the original CMP population without the inclusion of the FcεRI marker.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6