Single-cell analysis reveals the *KIT* D816V mutation in hematopoietic stem and

progenitor cells in systemic mastocytosis

Brief title: Single-cell analysis in systemic mastocytosis

Jennine Grootens, MSc[a], Johanna S. Ungerstedt, MD, PhD[b,c], Maria Ekoff, PhD[a], Elin

Rönnberg, PhD[a], Monika Klimkowska, MD, PhD[d], Rose-Marie Amini, MD, PhD[e],

Michel Arock, PhD[f], Stina Söderlund, MD, PhD[g,h], Mattias Mattsson, MD[e,g], Gunnar

Nilsson, PhD[a,g,h,*], and Joakim S. Dahlin, PhD[a,*]

Affiliations:

[a]Department of Medicine Solna, Karolinska Institutet, and Karolinska University

Hospital, Stockholm, 17164, Sweden;

[b]Department of Medicine Huddinge, Karolinska Institutet, Stockholm, 14186,

Sweden;

[c]Hematology Center, Karolinska University Hospital, Stockholm, 17176, Sweden;

[d]Department of Clinical Pathology and Cytology, Karolinska University Hospital

Huddinge, Stockholm, 14186, Sweden;

[e]Department of Immunology, Genetics and Pathology, Uppsala University, Uppsala,

75185, Sweden;

[f]Molecular and Cellular Oncology, Ecole Normale Supérieure de Cachan, Cachan,

94235, France and Clinical Hematology Laboratory, Pitié-Salpêtrière Hospital, Paris,

75013, France;

[g]Division of Hematology, Uppsala University Hospital, Uppsala, 75185, Sweden;

[h]Department of Medical Sciences, Uppsala University, Uppsala, 75185, Sweden;

[*]Corresponding authors at: Department of Medicine, Division of Immuology and

Allergy, Karolinska Institutet, 17164 Stockholm, Sweden.

Email addresses: gunnar.p.nilsson@ki.se (G.N. Nilsson), joakim.dahlin@ki.se (J.S.

Dahlin).

Abstract

Background: Systemic mastocytosis (SM) is a hematological disease characterized

by organ infiltration by neoplastic mast cells. Almost all SM patients have a mutation

in the gene encoding the tyrosine kinase receptor KIT causing a D816V substitution

and autoactivation of the receptor. Mast cells and CD34$^+$ hematopoietic progenitors

can carry the mutation, however, in which progenitor cell subset the mutation arises is

unknown. We aimed to investigate the distribution of the D816V mutation in single

mast cells and single hematopoietic stem and progenitor cells.

Methods: Fluorescence-activated single-cell index sorting and D816V mutation

assessment were applied to analyze mast cells and more than 10,000 CD34$^+$ bone

marrow progenitors across 10 hematopoietic progenitor subsets. In vitro assays

verified cell-forming potential.

Findings: We found that in SM 60-99% of the mast cells harbored the D816V

mutation. Despite increased frequencies of mast cells in SM patients compared with

control subjects, the hematopoietic progenitor subset frequencies were comparable.

Nevertheless, the mutation could be detected throughout the hematopoietic landscape

of SM patients, from hematopoietic stem cells to more lineage-primed progenitors. In

addition, we demonstrate that FcεRI$^+$ bone marrow progenitors exhibit mast cell-

forming potential, and we describe aberrant CD45RA expression on SM mast cells for

the first time.

Interpretation: The *KIT* D816V mutation arises in early hematopoietic stem and

progenitor cells and the mutation frequency is approaching 100% in mature mast

cells, which express the aberrant marker CD45RA.

Key words (4-6)

Systemic Mastocytosis; Single-Cell; Mast Cell; Mast Cell Progenitor; CD45RA;

Hematopoiesis; Hematopoietic Stem Cells; KIT D816V

Research in Context

Evidence before this study: Systemic mastocytosis is a myeloid neoplasm associated

with a mutation in the receptor KIT. The mutation is found in multiple mature cell

lineages and can be detected in bulk-sorted CD34$^+$ hematopoietic stem and progenitor

cells.

Added value of this study: We use single-cell analysis to study the distribution of

the common *KIT* mutation within the heterogenous CD34$^+$ progenitor landscape and

in mast cells of systemic mastocytosis patients. Isolation of progenitor subsets

followed by cell culture identifies a population of progenitors in bone marrow with

increased mast cell-forming potential. Furthermore, a novel marker for aberrant mast

cells in systemic mastocytosis is described.

Implications of all the available evidence: Our study provides novel insights into

the cellular origin and propagation of the common *KIT* mutation in systemic

mastocytosis. The identification of a novel marker for aberrant mast cells shows

potential to improve disease diagnosis.

Introduction

Hematopoietic stem cells (HSCs) in the bone marrow give rise to blood cells and mast cells. Differentiating HSCs progress through a number of intermediate progenitors with multilineage-forming capacity before commitment to the mast cell lineage.[1,2] The binding of stem cell factor (SCF) to its receptor, KIT, promotes the maturation and proliferation of mast cells.[3-6] It is therefore not surprising that mutations in the *KIT* gene coincide with the mast cell-driven disease systemic mastocytosis (SM).[7]

SM is a hematologic neoplasm in which infiltrates of neoplastic mast cells occur in various tissues.[8,9] The majority of SM patients carry a mutation in *KIT*, most commonly affecting codon 816.[10] The *KIT* D816V mutation makes receptor signaling constitutively active, independent of binding to its ligand SCF. The detection of *KIT* D816V in either bone marrow or peripheral blood samples is one of the criteria for the clinical diagnosis of SM using a standardized qPCR assay.[11] When the *KIT* D816V mutation is present in mast cells, it may also be detected in mature bone marrow or peripheral blood cells, such as basophils, eosinophils, neutrophils, and B and T lymphocytes, which depends on the patient.[12-17] Furthermore, the precursors of erythroid and myeloid cells as well as CD34$^+$ progenitors may carry the *KIT* D816V mutation.[12,18,19] Taken together, these findings suggest that the mutation may arise in early hematopoietic stem and progenitor cells (HSPCs), rather than in a lineage-restricted precursor.

The CD34$^+$ cell compartment is highly heterogeneous, spanning HSCs to late lineage-committed progenitors. In the present study, we delineated the cellular origin of the *KIT* D816V mutation in single bone marrow cells in SM by combining FACS index

sorting of CD34$^+$ HSPCs with a multiplex qPCR assay. The data revealed that the

D816V mutation in *KIT* arises in early HSPCs. The mutation burden is low but

variable in multipotent and lineage-restricted progenitor populations and increases in

mature mast cells. Furthermore, the present study provides more insight into

hematopoiesis in SM subjects and identifies high CD45RA expression in aberrant

mast cells.

Materials and Methods

Patients and sample preparation

Bone marrow and occasional peripheral blood samples were collected from patients

under evaluation for SM and from control subjects. The study was approved by the

Stockholm and Uppsala Regional Ethics committee. Oral and written informed

consent was obtained from each patient and control subject. The study was conducted

in accordance with the World Medical Association Declaration of Helsinki. The

patients included in our experimental studies were classified according to the World

Health Organization criteria[20] as having indolent SM (ISM, n = 19), aggressive SM

(ASM, n = 3), SM with associated hematological neoplasm (SM-AHN, n = 3), or

other malignancies, including myelodysplastic syndrome or myeloproliferative

neoplasm (MDS or MPN, n = 2). Supplementary Table 1 shows the patient

characteristics. For analytic purposes, ASM and SM-AHN patients were combined in

an advanced SM group (AdvSM). The controls (Ctrl) included patients with

cutaneous mastocytosis (CM, n = 2) and subjects without SM (n = 7). All samples

were processed on the day of collection. Red blood cells were lysed with PharmLyse

buffer (BD Biosciences, Franklin Lakes, NJ) and washed with PBS containing 2%

fetal calf serum (FCS, Thermo Fisher Scientific, Waltham, MA).

Cell lines

Two human mast cell lines, HMC-1.2 and ROSA$^{KIT WT}$, with and without the *KIT*

D816V mutation, respectively, were cultured as previously described.[21,22]

Flow cytometry and cell sorting

Bone marrow cells were stained with fluorophore-conjugated monoclonal antibodies

in Brilliant Stain Buffer (BD Biosciences, Franklin Lakes, NJ). The antibodies used

for staining included CD10 (HI10a), CD14 (M5E2), CD34 (581), CD38 (Hb7),

CD45RA (HI100), CD90 (5E10), CD117 (104D2), CD123 (6H6), CD133 (AC133),

and FcεRIα (CRA-1) (all from BD Biosciences, Biolegend, San Diego, CA, or

Miltenyi Biotec, Bergisch Gladbach, Germany). The live/dead cell marker DAPI (BD

Bioscience) was used for the analysis of cultured cells. Cell sorting and analysis were

performed using the FACSAria III and LSRFortessa systems (BD Biosciences). For

cell culture experiments, cells were sorted into sterile-filtered PBS with 2% FCS

following a two-step sorting protocol, using the yield followed by the purity precision

mode. The analysis of 20-100 sorted cells verified the sorting purity. Single $CD34^+$

cells were sorted into 2 μl of lysis buffer containing 20 μg/ml PCR-grade proteinase K

(Thermo Fisher Scientific) in EB buffer (Qiagen, Hilden, Germany). These cells were

sorted either immediately from the sample or following an enrichment sort. The

single-cell precision mode ensured that only one cell was deposited per well. The

plates were then centrifuged and stored at -80°C until analysis. The sorting data from

single cells were saved using the index sort module of the software FACSDIVA

software version 8 (BD Biosciences). Flow cytometry data analysis was performed

using FlowJo software version 10 (TreeStar, Ashland, OR).

Cell culture and colony-forming assays

Colony-forming potential was evaluated by plating 500 cells in 35 x 10 mm tissue

culture dishes (Thermo Fisher Scientific) in duplicate. The cells were cultured in

Methocult H4434, including SCF, IL-3, EPO, and GM-CSF (STEMCELL

Technologies Inc., Vancouver, Canada). Colonies were counted after 12-13 days,

158 distinguishing erythroid, granulocyte/monocyte, and mixed colonies following the

159 manufacturer's recommendations. The mast cell-forming potential was assessed via

160 liquid culture as described previously.[23] Briefly, sorted cells were cultured in medium

161 with recombinant human (rh) IL-3 (10 ng/mL; PeproTech, Rocky Hill, NJ) and rhIL-6

162 (10 ng/mL; PeproTech). In some experiments, the medium was changed on day 5, and

163 the cells were subsequently cultured with rhIL-6 (10 ng/mL) and SCF (100 ng/mL, r-

164 metHuSCF; Swedish Orphan Biovitrum, Stockholm, Sweden), as indicated in the

165 figure legends. Enzyme cytochemical staining of trypsin-like activity was used to

166 assess tryptase expression.[24,25] Cell morphology was evaluated by May-Grünwald

167 Giemsa staining (MGG; Sigma-Aldrich, St. Louis, MO).

169 *Single-cell mutation analysis*

170 Before the addition of lysis buffer, 96-well plates (Bio-Rad Laboratories, Hercules,

171 CA) were UV-treated for 30 minutes using a UVT-B-AR UV Cabinet (Grant

172 Instruments, Cambridge, UK). Tubes and pipettes were treated with the DNA AWAY

173 Surface Decontaminant (Thermo Fisher Scientific). After sorting, the cells were lysed

174 for 45 minutes at 55°C, followed by a 10-min heat-inactivating step at 95°C.

175 Multiplex qPCR analysis was performed via a Taqman gene expression assay

176 (Thermo Fisher Scientific) in lysed single cells using 96-well CFX Manager (Bio-Rad

177 Laboratories). A mutation assay was designed based on a previously published

178 method[11] using Primer3 (v.0.4.0) which specifically detected *KIT* c.2447A>T

179 (D816V) using the forward primer 5'-AGAGACTTGGCAGCCAGAAA-3'; a reverse

180 primer matching the mutation at the 3' position, 5'-

181 TTAACCACATAATTAGAATCATTCTTGATGA-3'; and the probe 5'-FAM-

182 TCCTCCTTACTCATGGTCGGATCACA-BHQ1-3'. A control assay was designed

to target a flanking exon and intron eight in GAPDH using the forward primer 5'-

CTGACTTCAACAGCGACACC-3'; the reverse primer 5'-

AGAGTTGTCAGGGCCCTTTT-3'; and the probe 5'-HEX-

TCAAGCTCATTTCCTGGTATGTG-BHQ1-3' (all primers came from TAG

Copenhagen, Denmark). The control and mutation assays were tested and optimized

using the cell lines ROSA$^{KIT\ WT}$ and HMC-1.2. The primer concentrations, optimized

for single cells, were as follows: control assay 150 nM forward/reverse primers, 100

nM probe; mutation assay 100 nM forward/reverse primers, 67 nM probe. Each plate,

containing 86 CD34$^+$ cells, 5 mature mast cells, 3 empty wells, and 2 wells with bulk

CD34$^+$ cells following sorting, was subjected to the multiplex qPCR assay. Cells were

excluded from analysis when amplification failed in the control assay, or when

amplification of the control assay resulted in a CT value higher than 40. In the

mutation assay, cells were regarded as unmutated when there was no amplification or

when the CT value was higher than 45. Outliers from each plate were then detected

using the ROUT outlier test in Prism software and removed from further analyses

(Supplementary Fig. 6B). Plates and cells that passed the quality control procedure

were linked with the index data and gated similarly to the whole sample for analysis.

Statistical analysis

Statistical analysis was performed using Prism software (Version 6.0h, GraphPad

Software, La Jolla, CA). The unpaired two-tailed Student's t-test was employed when

comparing two groups, and the unpaired one-way ANOVA with Tukey's multiple

comparison test was employed when comparing more than two groups. Statistical

analysis of log-transformed data was performed for some experiments as indicated in

the figure legends. Differences were considered significant when $P < 0.05$.

Results

Systemic mastocytosis patients have a normal CD34$^+$ bone marrow stem and

progenitor cell composition

To assess the composition of the hematopoietic cells in the bone marrow of SM

patients, we designed a multicolor flow cytometry panel that identifies ten distinct

HSPC subtypes and mature mast cells (Table 1; Fig. 1A-B). Colony formation assays

confirmed the validity of the GMP, CMP, and MEP gating strategies in SM subjects

(Supplementary Fig. 1A-C). The flow cytometry data provided a specific HSPC

profile for 34 bone marrow samples: 9 control, 19 ISM and 6 AdvSM (Fig. 1C;

Supplementary Table 1). The percentages of HSPCs in SM samples did not

significantly differ from those in control samples (Fig. 1D and Supplementary Fig.

2A). In addition, the ISM and AdvSM samples exhibited percentages of total CD34$^+$

cells similar to the control samples (Supplementary Fig. 2B). Thus, the overall

hematopoiesis profile did not differ between the control and SM subjects.

The bone marrow CMP$^{Fc\varepsilon RI+}$ fraction presents an enhanced mast cell-forming

potential

Mast cell progenitors circulate in the blood as Lin$^-$ CD34^{hi} CD117$^{int/hi}$ FcεRI$^+$ cells.[33]

These cells express CD123 and lack CD45RA,[23,33] overlaying the CMP$^{Fc\varepsilon RI+}$ gate in

bone marrow (Supplementary Fig. 3A) that exhibit a distinct expression profile

compared with mature mast cells (Supplementary Fig. 3B). We therefore analyzed the

mast cell-forming potential of the bone marrow CMP$^{Fc\varepsilon RI+}$ fraction by culturing the

cells with IL-3 and IL-6. Flow cytometry analysis revealed that the CMP$^{Fc\varepsilon RI+}$ cultures

gave rise to a higher frequency of CD117^{hi} FcεRI$^+$ mast cells than the CMPs$^{Fc\varepsilon RI-}$ cells

(Fig. 2A-C), despite overall lower CD117 expression (Supplementary Fig. 4A). May-

Grünwald Giemsa and enzymatic tryptase staining showed that the CMP$^{Fc\epsilon RI+}$ progeny

generated a high frequency of granulated tryptase-positive mast cells when cultured

with SCF (Fig. 2D-E and Supplementary Fig. 5). Cultured GMPs did not give rise to

any cells with a mast cell phenotype (Fig. 2A-E and Supplementary Fig. 5).

The emerging model of hematopoiesis proposes that MEPs, basophils, and

eosinophils develop separately from GMPs following an asymmetric cell division at

the MPP stage. This process can be traced by the segregation of CD133, which is

expressed in GMPs but not in progenitors with erythromyeloid output.[34] Indeed, we

found that CMP$^{Fc\epsilon RI+}$ cells exhibited only low levels of CD133 staining, similar to

erythromyeloid cells (Supplementary Fig. 4B).

CD45RA is expressed on aberrant mast cells in systemic mastocytosis

The multicolor flow cytometry panel used for the gating of HSPCs was also employed

to identify mast cells in bone marrow (Fig. 1B). May-Grünwald Giemsa and tryptase

staining confirmed that CD14$^-$ CD34$^-$ CD117^{hi} FcϵRI^{hi} cells constitute mast cells (Fig.

3A). SM samples presented higher percentages of mast cells than controls (Fig. 3B).

Notably, mast cells from almost all SM samples exhibited aberrant CD45RA

expression, as verified with fluorescence minus one control and internal control

staining of CD34$^+$ cells (Fig. 3C-D). The cutaneous (non-systemic) mastocytosis

samples and MDS/MPN samples expressed low levels of CD45RA on bone marrow

mast cells (Fig. 3C-D). In addition, mast cell frequencies were correlated with the

clinical percentage of BMMC infiltrate and serum tryptase levels (Fig. 4A-C;

Supplementary Table 2). Taken together, the results indicate that SM bone marrow

samples contain more mast cells than control samples and these mast cells express

high levels of the aberrant marker CD45RA.

Single-cell index sorting reveals the KIT D816V mutation in early HSPCs in systemic

mastocytosis patients

The cellular origin of the *KIT* D816V mutation in SM has not been established, and

the relative abundance of this mutation in different CD34$^+$ HSPC populations and

mast cells has not been determined. We therefore developed a single-cell method to

trace the *KIT* D816V mutation using two cell lines: HMC-1.2 cells carrying the

mutation and ROSA cells carrying wild-type *KIT*. The cells were labeled with two

different fluorochrome-labeled antibodies, mixed, and randomly FACS sorted with

the index sort function. A multiplex qPCR assay was used to detect an irrelevant gene,

for quality control purposes, and the *KIT* D816V mutation (Supplementary Fig. 6A-

B). Among all single cells, more than 96% passed the quality control procedure

(Supplementary Fig. 6C). Linking the results with the index data revealed that the

mutation assay was negative for the wild-type ROSA cells. The assay detected the

mutation in 98.1% of all HMC-1.2 cells. Thus, the assay did not detect false positive

cells in the mutation assay and only a few false negatives, demonstrating the high

sensitivity and specificity of the method (Supplementary Fig. 6D).

The high success rate with cell lines allowed us to use the assay for HSPCs and

mature mast cells from patient samples. Single cells from 12 samples (6 ISM and 6

AdvSM) were analyzed for mutation assessment. Specifically, CD34$^+$ HSPCs were

index sorted from ten patients, whereas specific HSPC subsets were sorted from two

patients. At the time of SM diagnosis, the presence of the *KIT* D816V mutation was

confirmed in either peripheral blood or bone marrow (Supplementary Table 1). We

analyzed 40-194 single mast cells and 351-1583 single CD34$^+$ cells from each bone

marrow sample, providing mutation data for 12,859 single cells in total (Fig. 5A;

Supplementary Table 3). All SM samples carried the *KIT* D816V mutation in the mast

cell compartment at a high frequency in ISM (mean, 80.9%) and AdvSM (mean,

93.2%) (Fig. 5B). Index data analysis of individual mast cells demonstrated that

CD45RA expression was associated with presence of the *KIT* mutation (Fig. 5C). The

percentage of mutated mast cells was correlated with the total percentage of mast cells

in the sample and with the serum tryptase levels (Fig. 6A-B; Supplementary Table 2).

The mutation rate in CD34$^+$ HSPCs was less than 3% in all samples (Fig. 5D), except

for one AdvSM sample with a high mutation burden of 37.0% (Supplementary Table

4). Nevertheless, the percentage of mutated CD34$^+$ HSPCs was correlated with serum

tryptase levels (Fig. 6C; Supplementary Table 2).

Coupling the index data with the mutation analysis of 10 bone marrow samples

revealed that the *KIT* D816V mutation appeared throughout the hematopoietic

landscape (Fig. 5E; Supplementary Table 3-4). Specifically, different HSPCs

presented similar mutation rates in all samples when the median percentage of

mutation was compared between different HSPCs (Fig. 5F). Overall, the method of

index sorting provided a representative set of single cells from the whole-bone-

marrow aspirates (Supplementary Fig. 7A). However, the CMP$^{FcεRI+}$ population with

an enriched mast cell potential was rare. To determine the mutation frequency in the

CMP$^{FcεRI+}$ fraction, we selectively sorted 86 and 278 single CMPs$^{FcεRI+}$ from 2 SM

samples. One patient exhibited less than 2% mutated CMPs$^{FcεRI+}$, HSCs and GMPs,

whereas the fraction of mutated mast cells was 98.9% (Fig. 5G; Supplementary Table

4). The second patient presented 95.7% mutated CMPs$^{Fc\varepsilon RI+}$ and 96.4 % mutated mast

cells (Fig. 5H; Supplementary Table 4). Peripheral blood MCPs (CD14$^-$ CD34$^+$

CD117$^+$ FcεRI$^+$) showed mutation rates comparable to that of bone marrow

CMPs$^{Fc\varepsilon RI+}$ in one patient (Supplementary Fig. 7B), but the mutation could not be

detected in the MCPs from 4 other SM samples, likely due to the low mutation

frequency and the small number of cells analyzed due to the rarity of the cells (14-23

cells analyzed per sample). Taken together, we detected the *KIT* D816V mutation

throughout the hematopoietic landscape, starting with the HSCs.

Discussion

The cellular origin of the aberrant clone in SM is still under debate. Here, we show

that SM patients exhibit an increased frequency of mast cells in bone marrow. Almost

all these mast cells carry the *KIT* D816V mutation. Despite normal HSPC frequencies

in bone marrow, the *KIT* D816V mutation can be traced all the way back to the

$CD34^+$ $CD38^-$ $CD10^-$ $CD45RA^-$ $CD90^+$ HSC fraction.

The *KIT* D816V mutation causes receptor autoactivation, which signals survival,

proliferation, and differentiation, likely explaining the competitive growth advantage

of the aberrant mast cell clone and the high frequency of mutated mast cells.

However, the frequency of mutated HSPCs was low in most patients. This

observation is consistent with the overall normal HSPC composition of SM patients.

There are two main potential explanations for the low D816V mutation burden

observed in HSPCs relative to mast cells: HSPCs rapidly differentiate into mast cells

when the mutation occurs, or mutated HSPCs do not present a strong growth

advantage over non-mutated HSPCs. In fact, we have previously shown that KIT

signaling is dispensable for mast cell progenitor development in vitro and in vivo, and

that other stimuli can induce cell proliferation.[23] However, the aberrant progenitor

clone may in some cases proliferate to reach a high frequency, as demonstrated in two

patients with AdvSM who presented more than 20% mutated cells in the HSC

fraction. Advanced forms of SM have previously been associated with detection of

the *KIT* mutation in bulk-sorted $CD34^+$ progenitors and various mature cell lineages,

whereas in indolent disease, the mutation is mainly restricted to mast cells.[13]

It is of particular interest to study the mutation burden in progenitors with mast cell-

forming potential. We have previously reported that $CD34^+$ $CD117^{int/hi}$ $Fc\epsilon RI^+$ cells in

peripheral blood are committed or nearly committed MCPs.[33] Considering the $CD34^+$

$CD117^+$ $Fc\epsilon RI^+$ phenotype of $CMPs^{Fc\epsilon RI+}$ cells, these cells might be expected to

present a similar mast cell-forming capacity as to blood MCPs. In the current study,

we showed that $CMPs^{Fc\epsilon RI+}$ gave rise to a higher frequency of mast cells than

$CMPs^{Fc\epsilon RI-}$ and GMPs. However, the mast cell frequency in cultured $CMPs^{Fc\epsilon RI+}$ was

less than 10% and a substantial fraction were found to be $CD117^{-/lo}$ $Fc\epsilon RI^+$ cells after

culture, consistent with a basophil-like phenotype. Thus, $CMPs^{Fc\epsilon RI+}$ exhibit a

considerable mast cell-forming capacity but also include progenitors with non-mast

cell output. Basophils mature in bone marrow, whereas mast cells mature in

peripheral tissues, which likely explains why bone marrow $CMPs^{Fc\epsilon RI+}$ form $CD117^-$

$Fc\epsilon RI^+$ basophil-like cells, whereas $CD34^+$ $CD117^{int/hi}$ $Fc\epsilon RI^+$ blood progenitors do

not. The high frequency of $CD117^-$ $Fc\epsilon RI^+$ basophil-like cells that develop from

$CMPs^{Fc\epsilon RI+}$ is also in agreement with the reduced CD117 expression of $CMPs^{Fc\epsilon RI+}$

compared with that of $CMPs^{Fc\epsilon RI-}$. CD117 is upregulated upon mast cell maturation,

and blood mast cell progenitors express intermediate to high levels of CD117.

$CMPs^{Fc\epsilon RI+}$ express CD117 at low levels overall, even though there are individual

$CMPs^{Fc\epsilon RI+}$ cells that express high CD117 levels, which might correspond to the cells

with mast cell-forming capacity. Taken together, our results indicate that the low *KIT*

mutation prevalence in the $CMP^{Fc\epsilon RI+}$ fraction is likely explained by the capacity of

these cells' to form not only mast cells but also $CD117^-$ $Fc\epsilon RI^+$ basophil-like cells.

In SM, mast cells are known to aberrantly express multiple surface proteins, among

CD2 and CD25 are currently used for diagnosis.[35] These markers are expressed

independently of the SM subtype. Other markers, such as CD30 and CD52, are

associated with AdvSM and mast cell leukemia.[36,37] In the present study, we

demonstrate that CD45RA is a novel marker for aberrant mast cells in SM.

Remarkably, bone marrow mast cells from two patients with cutaneous mastocytosis

expressed low CD45RA levels, suggesting that CD45RA is a marker for systemic

mast cell disease. The increase in the CD45RA isoform of CD45 likely explains the

high levels of CD45 observed in systemic mastocytosis patients.[38-41] Taken together,

our results indicate that high CD45RA expression should be considered a novel

marker for diagnosis of SM.

CD45RA is present on aberrant mast cells in SM, but at what stage during mast cell

differentiation is the marker upregulated? CMPs$^{Fc\epsilon RI+}$, which lack CD45RA, present

mast cell-forming potential in SM patients. In contrast, we show that cells in the

classic CD45RA$^+$ GMP gate lack mast cell potential in SM patients. An alternative

explanation is that the mast cell progenitors in SM fall into the immature CD38$^-$ gate.

In fact, examples of rare mutated CD117$^{int/hi}$ FcϵRI$^{int/hi}$ cells in the CD45RA$^+$ LMPP

gate were seen (Supplementary Fig. 7C). It is also tempting to speculate that CD45RA

is upregulated once aberrant mast cells are mature.

The CD34$^+$ HSPC population is highly heterogeneous and a single-cell resolution is

therefore necessary to delineate the cellular distribution of the mutation. Here, we

used single-cell index sorting to track cellular identity when performing mutation

analysis. Notably, the mutation was detected throughout differentiation, from HSCs to

the MEP, GMP, and LMPP fractions, depending on the patient. The detection of the

mutation in early HSPCs explains previous findings of the mutation being present in

bulk-sorted mature cell lineages of some patients.

Multiplex TaqMan qPCR assays detected the *KIT* mutation in index-sorted

progenitors in the present investigation. The qPCR method was originally developed

for sensitive mutation detection in DNA extracted from bone marrow or blood cells.

GAPDH primers and probes were added to the assay to confirm the presence or

absence of a sorted cell. This method allowed the detection of mutated cells with

close to 100% efficiency, with no observed false positives. The single-cell index

sorting and DNA mutation analysis approach can similarly be applied to other

mutations and hematologic diseases to reveal the origin of aberrant clones. One

limitation of the method is the number of cells that can be analyzed, meaning that

mutated cells in very rare subpopulations might go undetected. However, more than

10,000 HSPCs were analyzed in the present investigation, and single-cell index

sorting coupled with qPCR allowed unbiased and prospective analysis of mutated

HSPC populations in SM. Whole-genome amplification and nested PCR methods for

amplifying and sequencing DNA mutations from single cells are associated with the

phenomenon known as allelic dropout, which is the amplification of only a single

allele, leading to an overestimation of the frequency of non-mutated cells. Mutational

profiling of single colonies, as performed by Jawhar *et al.*[18] avoids this problem.

However, such analysis can be performed only on progenitors that form sufficiently

large colonies. Mast cell progenitors present a poor proliferative capacity, and this

technique was therefore not applied in the present investigation.

In summary, we have explored the composition and mutation profile of CD34$^+$ HSPCs

from SM patients. The results show that the *KIT* mutation can be found in cells

throughout the hematopoietic landscape with increased burden in the mast cell

lineage, supporting the notion that the aberrant clone may arise in the hematopoietic

stem cell compartment. Furthermore, the mutated mast cells express CD45RA, a

potential novel clinical biomarker for aberrant mast cells in SM.

Acknowledgements

We wish to thank Kerstin Hamberg Levedahl for help with the patients and Petter

Woll for helpful discussions. Stem cell factor was a kind gift from SOBI, Stockholm,

Sweden.

Funding Sources

This study was supported by the Swedish Research Council, the Swedish Cancer

Society, the Stockholm Cancer Society (Radiumhemmets Forskningsfonder), Magnus

Bergvall's Foundation, Tore Nilsson's Foundation for Medical Research, the

Karolinska Institutet, and through the regional agreement on medical training and

clinical research (ALF) between Stockholm County Council and the Karolinska

Institutet, and ALF-funding from Uppsala University Hospital.

Declaration of Interest

The authors declare no competing interests

Author Contributions

Conceptualization and Methodology, J.G., G.N. and J.S.D.; Investigation, J.G., M.E.

and E.H.; Formal analysis, J.G.; Visualization and Validation, J.G. and J.S.D.;

Resources, J.S.U., M.K., R.A., S.S., M.M. and M.A.; Writing – original draft, J.G.

and J.S.D.; Writing – review & editing, all authors; Funding acquisition and

Supervision, J.U., G.N. and J.S.D.

References

1.   Dahlin JS, Hallgren J. Mast cell progenitors: origin, development and

migration to tissues. *Mol Immunol* 2015; **63**(1): 9-17.

2.   Grootens J, Ungerstedt JS, Nilsson G, Dahlin JS. Deciphering the

differentiation trajectory from hematopoietic stem cells to mast cells. *Blood Adv* 2018;

2(17): 2273-81.

3.   Irani AM, Nilsson G, Miettinen U, et al. Recombinant human stem cell factor

stimulates differentiation of mast cells from dispersed human fetal liver cells. *Blood*

1992; **80**(12): 3009-21.

4.   Kirshenbaum AS, Goff JP, Kessler SW, Mican JM, Zsebo KM, Metcalfe DD.

Effect of IL-3 and stem cell factor on the appearance of human basophils and mast

cells from CD34+ pluripotent progenitor cells. *J Immunol* 1992; **148**(3): 772-7.

5.   Valent P, Spanblochl E, Sperr WR, et al. Induction of differentiation of human

mast cells from bone marrow and peripheral blood mononuclear cells by recombinant

human stem cell factor/kit-ligand in long-term culture. *Blood* 1992; **80**(9): 2237-45.

6.   Mitsui H, Furitsu T, Dvorak AM, et al. Development of human mast cells

from umbilical cord blood cells by recombinant human and murine c-kit ligand. *Proc*

Natl Acad Sci U S A 1993; **90**(2): 735-9.

7.   Nagata H, Worobec AS, Oh CK, et al. Identification of a Point Mutation in the

Catalytic Domain of the Protooncogene C-Kit in Peripheral-Blood Mononuclear-Cells

of Patients Who Have Mastocytosis with an Associated Hematologic Disorder. *Proc*

Natl Acad Sci U S A 1995; **92**(23): 10560-4.

8.   Valent P, Akin C, Escribano L, et al. Standards and standardization in

mastocytosis: consensus statements on diagnostics, treatment recommendations and

response criteria. *Eur J Clin Invest* 2007; **37**(6): 435-53.

9.   Valent P, Akin C, Metcalfe DD. Mastocytosis: 2016 updated WHO

classification and novel emerging treatment concepts. *Blood* 2017; **129**(11): 1420-7.

10.   Arock M, Sotlar K, Akin C, et al. KIT mutation analysis in mast cell

neoplasms: recommendations of the European Competence Network on Mastocytosis.

Leukemia 2015; **29**(6): 1223-32.

11.   Kristensen T, Vestergaard H, Moller MB. Improved detection of the KIT

D816V mutation in patients with systemic mastocytosis using a quantitative and

highly sensitive real-time qPCR assay. *J Mol Diagn* 2011; **13**(2): 180-8.

12.   Afonja O, Amorosi E, Ashman L, Takeshita K. Multilineage involvement and

erythropoietin-"independent" erythroid progenitor cells in a patient with systemic

mastocytosis. *Ann Hematol* 1998; **77**(4): 183-6.

13.   Garcia-Montero AC, Jara-Acevedo M, Teodosio C, et al. KIT mutation in

mast cells and other bone marrow hematopoietic cell lineages in systemic mast cell

disorders: a prospective study of the Spanish Network on Mastocytosis (REMA) in a

series of 113 patients. *Blood* 2006; **108**(7): 2366-72.

14.   Kocabas CN, Yavuz AS, Lipsky PE, Metcalfe DD, Akin C. Analysis of the

lineage relationship between mast cells and basophils using the c-kit D816V mutation

as a biologic signature. *J Allergy Clin Immunol* 2005; **115**(6): 1155-61.

15.   Yavuz AS, Lipsky PE, Yavuz S, Metcalfe DD, Akin C. Evidence for the

involvement of a hematopoietic progenitor cell in systemic mastocytosis from single-

cell analysis of mutations in the c-kit gene. *Blood* 2002; **100**(2): 661-5.

16.   Pardanani A, Reeder T, Li CY, Tefferi A. Eosinophils are derived from the

neoplastic clone in patients with systemic mastocytosis and eosinophilia. *Leuk Res*

2003; **27**(10): 883-5.

17.   Mayado A, Teodosio C, Dasilva-Freire N, et al. Characterization of CD34(+)

hematopoietic cells in systemic mastocytosis: Potential role in disease dissemination.

Allergy 2018; **73**(6): 1294-304.

18.   Jawhar M, Schwaab J, Schnittger S, et al. Molecular profiling of myeloid

progenitor cells in multi-mutated advanced systemic mastocytosis identifies KIT

D816V as a distinct and late event. *Leukemia* 2015; **29**(5): 1115-22.

19.   Jara-Acevedo M, Teodosio C, Sanchez-Munoz L, et al. Detection of the KIT

D816V mutation in peripheral blood of systemic mastocytosis: diagnostic

implications. *Mod Pathol* 2015; **28**(8): 1138-49.

20.   Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health

Organization classification of myeloid neoplasms and acute leukemia. *Blood* 2016;

127(20): 2391-405.

21.   Saleh R, Wedeh G, Herrmann H, et al. A new human mast cell line expressing

a functional IgE receptor converts to tumorigenic growth by KIT D816V transfection.

Blood 2014; **124**(1): 111-20.

22.   Sundstrom M, Vliagoftis H, Karlberg P, et al. Functional and phenotypic

studies of two variants of a human mast cell line with a distinct set of mutations in the

c-kit proto-oncogene. *Immunology* 2003; **108**(1): 89-97.

23.   Dahlin JS, Ekoff M, Grootens J, et al. KIT signaling is dispensable for human

mast cell progenitor development. *Blood* 2017; **130**(16): 1785-94.

24.   Harvima IT, Naukkarinen A, Harvima RJ, Fraki JE. Immunoperoxidase and

enzyme-histochemical demonstration of human skin tryptase in cutaneous mast cells

in normal and mastocytoma skin. *Arch Dermatol Res* 1988; **280**(6): 363-70.

25.   Dahlin JS, Ungerstedt JS, Grootens J, et al. Detection of circulating mast cells

in advanced systemic mastocytosis. *Leukemia* 2016; **30**(9): 1953-6.

26.   Notta F, Doulatov S, Laurenti E, Poeppl A, Jurisica I, Dick JE. Isolation of

Single Human Hematopoietic Stem Cells Capable of Long-Term Multilineage

Engraftment. *Science* 2011; **333**(6039): 218-21.

27.   Majeti R, Park CY, Weissman IL. Identification of a hierarchy of multipotent

hematopoietic progenitors in human cord blood. *Cell Stem Cell* 2007; **1**(6): 635-45.

28.   Manz MG, Miyamoto T, Akashi K, Weissman IL. Prospective isolation of

human clonogenic common myeloid progenitors. *Proc Natl Acad Sci U S A* 2002;

99(18): 11872-7.

29.   Goardon N, Marchi E, Atzberger A, et al. Coexistence of LMPP-like and

GMP-like leukemia stem cells in acute myeloid leukemia. *Cancer Cell* 2011; **19**(1):

138-52.

30.   Lee J, Breton G, Oliveira TY, et al. Restricted dendritic cell and monocyte

progenitors in human cord blood and bone marrow. *J Exp Med* 2015; **212**(3): 385-99.

31.   Olweus J, BitMansour A, Warnke R, et al. Dendritic cell ontogeny: a human

dendritic cell lineage of myeloid origin. *Proc Natl Acad Sci U S A* 1997; **94**(23):

12551-6.

32.   Doulatov S, Notta F, Eppert K, Nguyen LT, Ohashi PS, Dick JE. Revised map

of the human progenitor hierarchy shows the origin of macrophages and dendritic

cells in early lymphoid development. *Nat Immunol* 2010; **11**(7): 585-93.

33.   Dahlin JS, Malinovschi A, Ohrvik H, et al. Lin- CD34hi CD117int/hi

FcepsilonRI+ cells in human blood constitute a rare population of mast cell

progenitors. *Blood* 2016; **127**(4): 383-91.

34.   Gorgens A, Radtke S, Mollmann M, et al. Revision of the human

hematopoietic tree: granulocyte subtypes derive from distinct hematopoietic lineages.

Cell Rep 2013; **3**(5): 1539-52.

35. Teodosio C, Mayado A, Sanchez-Munoz L, et al. The immunophenotype of

mast cells and its utility in the diagnostic work-up of systemic mastocytosis. *J Leukoc*

Biol 2015; **97**(1): 49-59.

36. Sotlar K, Cerny-Reiterer S, Petat-Dutter K, et al. Aberrant expression of CD30

in neoplastic mast cells in high-grade mastocytosis. *Mod Pathol* 2011; **24**(4): 585-95.

37. Hoermann G, Blatt K, Greiner G, et al. CD52 is a molecular target in

advanced systemic mastocytosis. *FASEB J* 2014; **28**(8): 3540-51.

38. Chisholm KM, Merker JD, Gotlib JR, et al. Mast cells in systemic

mastocytosis have distinctly brighter CD45 expression by flow cytometry. *Am J Clin*

Pathol 2015; **143**(4): 527-34.

39. Escribano L, Diaz-Agustin B, Lopez A, et al. Immunophenotypic analysis of

mast cells in mastocytosis: When and how to do it. Proposals of the Spanish Network

on Mastocytosis (REMA). *Cytometry B Clin Cytom* 2004; **58**(1): 1-8.

40. Morgado JM, Sanchez-Munoz L, Teodosio CG, et al. Immunophenotyping in

systemic mastocytosis diagnosis: 'CD25 positive' alone is more informative than the

'CD25 and/or CD2' WHO criterion. *Mod Pathol* 2012; **25**(4): 516-21.

41. Teodosio C, Garcia-Montero AC, Jara-Acevedo M, et al. Mast cells from

different molecular and prognostic subtypes of systemic mastocytosis display distinct

immunophenotypes. *J Allergy Clin Immunol* 2010; **125**(3): 719-26, 26 e1-26 e4.

Figure legends

Fig. 1

HSPC gating and profiling shows a normal course of hematopoiesis in patients

with systemic mastocytosis. (A) A landscape model of the hematopoiesis. (B) Gating

strategy for bone marrow HSPCs and mature mast cells, showing representative plots

with population frequencies as percentages. (C) The hematopoietic progenitor profile

of 34 bone marrow samples. Cutaneous mastocytosis samples are indicated in italics.

(D) SM samples were compared with control samples without SM, including CM

(open circles), and the mean expression + SEM of each HSPC is presented as a

percentage of the total $CD34^+$ cells. The unpaired *t*-test was used in panel D to

compare the control and SM groups for each HSPC subset.

Fig. 2

The $CMP^{Fc\epsilon RI+}$ fraction in the bone marrow has mast cell potential. (A-C) Flow

cytometry analysis of of $CMP^{Fc\epsilon RI-}$, $CMP^{Fc\epsilon RI+}$ and GMP cells cultured with rhIL-3

and rhIL-6 up to 21 days. (A) Flow cytometry plots of representative cultures,

showing the percentages of $CD117^-$, $CD117^+$, and $CD117^{hi}$ cells in each plot. (B)

Percentage of $CD117^{hi}$ $Fc\epsilon RI^+$ cells at days 5-6. (C) Two of the samples in panel B

were followed for up to 21 days under culture with rhIL-3 and rhIL-6. The percentage

of $CD117^{hi}$ cells was analyzed on days 5-6, 13-14, and 19-21. (D) Representative

images of $CMP^{Fc\epsilon RI-}$, $CMP^{Fc\epsilon RI+}$ and GMP cells cultured with rhIL-3 and rhIL-6 for 5

days, followed by rhIL-6 and SCF until day 13-14, showing tryptase staining (red)

and granules by May-Grünwald Giemsa (MGG) staining. (E) Quantification of

tryptase-positive cells at days 13-14 and 17-21. $CMP^{Fc\epsilon RI-}$, $CMP^{Fc\epsilon RI+}$ cells were

analyzed from Ctrl29, SM28, SM32, SM33, and MDS30. GMPs were analyzed from

SM32 and SM33. The bars and lines in panel B, C and D represent the means ± SEM.

One-way ANOVA with Tukey's multiple comparison was used in panel B, **$P <$

0.01, ***$P < 0.001$. Images were captured using an Olympus XC10 camera

(Olympus, Tokyo, Japan). The image width corresponds to 29 μm.

Fig. 3

SM mast cells aberrantly express CD45RA. (A) Enzymatic staining demonstrates

tryptase expression, and MGG staining reveals granules in SM mast cells. The images

are representative of two experiments (samples SM32 and SM33). Images were

captured using an Olympus XC10 camera (Olympus, Tokyo, Japan). The image width

corresponds to 29 μm. (B) Mast cell frequencies in Ctrl, ISM, AdvSM, and

MDS/MPN patients, showing the median of log-transformed values. (C) Histograms

showing CD45RA expression in mast cells (red line), FMO controls (grey histogram)

and CD34$^+$ cells (dotted line). (D) Frequency of CD45RA-expressing mast cells,

showing the mean and SEM. Control subjects included two subjects with CM (open

circles). One-way ANOVA with Tukey's multiple comparison was used in panels B

and D, *$P < 0.05$, **$P < 0.01$, **** $P < 0.0001$.

Fig. 4

The percentage of mast cells in bone marrow correlates with clinical features.

The percentage of mast cells was correlated with (A) serum tryptase levels and (B) the

percentage of BMMC infiltrate in the biopsy section. (C) Serum tryptase levels were

correlated with the percentage of BMMC infiltrate in the biopsy section. The samples

include ISM (closed dots) and AdvSM (open dots). Correlation values and

significance are indicated in the figure and in Supplementary Table 2. Linear

regression was calculated based on log-transformed data: in panel A, n = 25, and in

panels B and C, n = 23. The percentage of BMMC infiltrate was not determined for

two patients, accounting for the difference in numbers between panels. For analytical

purposes, BMMC infiltrate presented as <1% or <5% in Supplementary Table 1 were

set at 1% and 5% respectively.

Fig. 5

Mutation analysis of KIT D816V in single cells identifies mutated HSPCs and

mast cells. (A) Method of single-cell mutation analysis using index sorting and

multiplex qPCR. (B) The bars indicate the means + SEM percentage of mutated mast

cells. (C) The ratio between the median CD45RA expression of mutated and wild type

(WT) mast cells was determined for each patient. Each data point represents one

patient. The numbers of cells analyzed from each patient are presented in

Supplementary Table 3. Eleven patients are shown, as index data was not available

from 1 of the 12 patients. The two-tailed Wilcoxon signed rank test determined

whether the ratio was different from 1, indicated with a dotted line. *$P < 0.05$. (D)

The bars indicate the means + SEM percentage of mutated HSPCs. (E) The cellular

distribution of the mutation in 10 different HSPCs, showing the total numbers of

mutated and sorted cells for each sample that passed the quality control procedure. (F)

The bars indicate the median percentage of mutated cells per HSPC. (G-H) Mutation

frequencies of selectively isolated HSPCs following a 2-step sorting protocol in two

samples (SM28, ISM; and SM31, AdvSM). These two samples are indicated with

open circles in panels B and F.

Fig. 6

The mutation rate in mast cells and HSPCs correlates with clinical features.

(A-C) Linear regression of the percentage of mutated mast cells, showing significant

correlations with (A) the percentage of mast cells and (B) serum tryptase levels. (C)

The percentage of mutated HSPCs was correlated with serum tryptase levels. The

samples included ISM (closed dots) and AdvSM (open dots). Correlation and

significance are indicated in the figure and in Supplementary Table 2. Linear

regression was calculated based on log-transformed data in panels A-C, n = 12 and D,

n = 10.

Table legend

Table 1. Immunophenotypes of HSPCs

Fig. 1

Fig. 2

Fig. 3

A

Tryptase   MGG

B

% mast cells

*
**

Ctrl   ISM   AdvSM   MDS/MPN

C

% of maximum

Ctrl37

CM38

SM39

CD45RA

D

% CD45RA⁺ mast cells

****
****   ****

Ctrl   ISM   AdvSM   MDS/MPN

Mast cells
FMO
CD34⁺

Fig. 4

Fig. 5

Fig. 6

Table 1. Immunophenotypes of hematopoietic stem and progenitor cells

Short	Cell type	Gating strategy	References
HSC	Hematopoietic stem cell	$CD14^-$ $CD34^+$ $CD38^-$ $CD45RA^-$ $CD90^+$	26, 27
MPP	Multipotent progenitor	$CD14^-$ $CD34^+$ $CD38^-$ $CD45RA^-$ $CD90^-$	26, 27
$CMP^{Fc\varepsilon RI-}$	Common myeloid progenitor $Fc\varepsilon RI^-$	$CD14^-$ $CD34^+$ $CD38^+$ $CD45RA^-$ $CD10^-$ $CD123^+$ $Fc\varepsilon RI^-$	28 *
$CMP^{Fc\varepsilon RI+}$	Common myeloid progenitor $Fc\varepsilon RI^+$	$CD14^-$ $CD34^+$ $CD38^+$ $CD45RA^-$ $CD10^-$ $CD123^+$ $Fc\varepsilon RI^+$ $CD117^+$	28 *
MEP	Megakaryocyte-erythroid progenitor	$CD14^-$ $CD34^+$ $CD38^+$ $CD45RA^-$ $CD10^-$ $CD123^-$	28
LMPP	Lymphoid-primed multipotent progenitor	$CD14^-$ $CD34^+$ $CD38^-$ $CD45RA^+$	29
GMP	Granulocyte-monocyte progenitor	$CD14^-$ $CD34^+$ $CD38^+$ $CD45RA^+$ $CD10^-$ $CD123^+$	28
CDP	Common dendritic progenitor	$CD14^-$ $CD34^+$ $CD38^+$ $CD45RA^+$ $CD10^-$ $CD123^{hi}$	30, 31
MLP	Multilymphoid progenitor	$CD14^-$ $CD34^+$ $CD38^-$ $CD45RA^+$ $CD10^+$	32
BNKP	B- and NK cell progenitor	$CD14^-$ $CD34^+$ $CD38^+$ $CD45RA^+$ $CD10^+$	32
MC	Mast cell	$CD14^-$ $CD34^-$ $CD117^{hi}$ $Fc\varepsilon RI^+$	

*Refers to the original CMP population without the inclusion of the FcεRI marker.