Bacterial community dynamics during embryonic and

larval development of three confamilial echinoids

Tyler J. Carrier[1,*] and Adam M. Reitzel[1]

[1] Department of Biological Sciences, University of North Carolina at Charlotte, Charlotte,

NC USA, 28223

[*] Corresponding author: tcarrie1@uncc.edu

Running title: Bacterial dynamics during development

Keywords: microbiome; *Strongylocentrotus*; *Mesocentrotus*; host-microbe; symbiosis

ABSTRACT

16
17
       Development of some animals is influenced by and, in some cases, dependent on

18
the associated microbiota. The timing of when associated bacterial communities are

19
established during the development of marine invertebrates and their subsequent dynamics

20
across stages are known for only a few species. Here, we compare the bacterial

21
communities of three confamilial echinoids from egg to juvenile using sequence-based

22
approaches. Bacterial communities are established on unfertilized eggs and change

23
gradually during embryonic and larval development. Despite the differences amongst these

24
pre-metamorphic stages, approximately thirty-percent of OTUs identified in association

25
with unfertilized egg were present in the juveniles. During embryonic development, host-

26
associated communities diverged from the environmental microbiota but later converged

27
following the onset of larval feeding. Taken together, the data presented here support the

28
hypothesis that bacterial communities are established prior to fertilization and community

29
composition shifts gradually thereafter, all while remaining distinct from the environment.

30
Future work will need to determine the relative influence of the host and bacteria-bacteria

31
interactions in shaping the associated bacterial community to determine the potential

32
functional importance of bacteria during the development of larval sea urchins and benthic

33
marine invertebrates more broadly.

INTRODUCTION

The journey from fertilized egg to adulthood has fascinated developmental biologists for centuries. A growing consensus in the field of animal-microbe symbioses suggests that developmental processes are, at least in part, affected by the presence and activity of associated microbiota (McFall-Ngai & Ruby 2000, McFall-Ngai 2002, Fraune & Bosch 2010, Nyholm & McFall-Ngai 2014). The degree of dependency on these communities is highly variable: the removal of bacterial associates inhibits the differentiation of gut tissues in zebrafish (Bates et al. 2006) while other taxa are seemingly unaffected (Manahan et al. 1993). Nevertheless, several patterns have emerged from comparative studies of animal-bacteria interactions. First, animals tend to associate with developmental stage-specific bacterial communities (Apprill et al. 2012, Mortzfeld et al. 2015, Carrier & Reitzel 2018). Second, changes in morphology and life history stage correlate with shifts in the composition and structure of this community (Apprill et al. 2012, Sharp et al. 2012, Mortzfeld et al. 2015, Fieth et al. 2016). Lastly, during ontogeny, animal-associated bacterial communities can be highly specific and diverge from the environmental microbiota through host-mediated selection (Davidson & Stahl 2008, Fraune et al. 2010, Pradea 2011, Franzenburg et al. 2013, McFall-Ngai et al. 2013).

Development of many marine invertebrates includes several morphologically diverse life-history stages. Regardless of taxonomy, reproduction in most species involves an external fertilization, embryogenesis, and larval development before settling and metamorphosing into a benthic juvenile (*e.g.*, Thorson 1950, Mileikovsky 1971). When microbial communities are established during early development for marine invertebrates appears to vary among species. Larvae of deep-sea vestimentiferans, for example, are

symbiont free until settlement while several species from various taxa are known to

vertically inherit microbial symbionts and diversify this community later in development

by horizontal (environmental) acquisitions (Nussbaumer et al. 2006, Stewart et al. 2008,

Bright & Bulgheresi 2010, Apprill et al. 2012, Sharp et al. 2012, Sipkema et al. 2015).

Once established, the location of the bacterial symbionts can also vary. The embryos and

larvae of eight echinoderm species have, for example, symbiotic bacteria in the out-pockets

of the extracellular matrix surrounding the larval body as well as near or in gut epidermal

cells (Cameron & Holland 1983, Bosch 1992, Cerra et al. 1997, Galac et al. 2016, Carrier

& Reitzel 2018, Carrier et al. 2018b).

The composition and structure of the bacterial communities associated with

echinoderm larvae is highly variable and has been shown to be influenced by food

availability (Carrier & Reitzel 2018, Carrier et al. 2018b), phenotype (Carrier & Reitzel

2018), asexual reproduction (Galac et al. 2016), and, potentially, disease (Carrier et al.

2018a). Three confamilial echinoderms (*Strongylocentrotus purpuratus*, *Mesocentrotus*

franciscanus, and *S. droebachiensis*), in particular, associate with species-specific bacterial

communities that shift according to larval stage (Carrier & Reitzel 2018). When the diverse

and dynamic bacterial community associating with these echinoid larvae are established

and how this community varies throughout the entirety of pelagic development is unknown.

By sampling each developmental stage from the unfertilized egg to benthic juvenile

and assaying the associated bacterial communities using next-generation sequencing, we

test two explicit hypotheses. First, we test if stage-specific bacterial communities during

larval development result from associations established in earlier embryonic stages and

continue after metamorphosis. Second, we test that three confamilial Strongylocentrotids

exhibit relatively similar developmental patterning in the bacterial communities due to their

phylogenetic relatedness. Implications from these hypotheses are predicted to complement

previous studies showing that the biology of echinoid larvae is inseparable from their

microbial community and is an informative system to understand the ecology of animal-

microbe symbiosis.

MATERIALS AND METHODS

Specimen collection and larval rearing

Adult urchin for each species were collected from populations in the Salish Sea in

April 2016 (Carrier & Reitzel 2018). *Strongylocentrotus purpuratus* was hand-collected at

low tide at Slip Point, Clallam Bay, WA (48°15'39" N, 124°15'03" W; Figure S1) and

transferred overnight to the Friday Harbor Laboratories (FHL; University of Washington;

Friday Harbor, WA, USA; Figure S1). Similarly, *S. droebachiensis* were hand-collected at

low tide at Cattle Point, San Juan Island, WA (48°27'00" N, 122°57'43" W; Figure S1) and

were transferred to FHL within one hour. *M. franciscanus*, on the other hand, were

collected by SCUBA off Bell Island, WA (48°35'49" N, 122°58'55" W; Figure S1) and

transferred to FHL within two hours. All urchins were suspended in separate sub-tidal

cages off the dock at FHL and fed *Nereocystis* spp. (sugar kelp) *ad libitum* until spawning

two weeks later.

Urchins were spawned by a one to two mL intracoelomic injection of 0.50 M KCl.

Gametes from *S. purpuratus* (n=3 for males and females), *M. franciscanus* (n=1 for males

and n=3 for females), and *S. droebachiensis* (n=3 for males and females) were collected,

rinsed, and pooled separately for each species. Fertilization of eggs and larval rearing

followed Strathmann (1987), with the exception that embryos and larvae were reared using

5.0-µm filtered seawater (FSW) to include the environmental microbiota.

Embryos were incubated in one-liter of FSW at ambient conditions (~10ºC and

12ºC; Carrier & Reitzel 2018). Two hours post-fertilization, embryos were transferred to

triplicate one-liter jars at a density of two individuals•mL^{-1}. Beginning at the prism stage,

larvae were provided *Rhodomonas lens* at 10,000 cells•mL^{-1}, for which was made growth

medium-free by centrifugation and resuspension in FSW prior to the introduction to larval

cultures. Larval cultures were subsequently diluted to one larvae per two mL at advanced

developmental stages. Larval cultures were given 90 to 95% water changes every other day

and *R. lens* was replenished at 10,000 cells•mL^{-1}.

Monocultures of *R. lens* were grown in f/2 media at room temperature with a

combination of ambient and artificial lighting for 24 hours per day (Guillard 1975).

Sampling across development

Each urchin species was cultured from egg to juvenile. We sampled ten stages: (i)

unfertilized eggs, (ii) 16-cell embryo (referred to hereafter as 'cleavage'), (iii) pre-hatched

blastula, (iv) hatched blastula, (v) gastrula, (vi) pre-feeding prism, (vii) 4-, (viii) 6-, and (ix)

8-armed larvae, and (x) post-metamorphic juvenile (see, Table S1 for biological replication

for each species at each developmental stage). Developmental stages were identified using

a compound microscope (Nikon Eclipse E600; camera: QImaging MicroPublisher 5.0

RTV). At each developmental stage, 100 individuals per jar (n=3) were sampled at random

for each species. Immediately after collections, samples were concentrated into a pellet

using a microcentrifuge, the FSW was removed with a sterile glass pipette, and pelleted

embryos, larvae, or juveniles were then preserved in RNAlater (Thermo Scientific,

Massachusetts, USA) and stored at -20°C before DNA extractions.

Complementary to sampling of the developmental stages, the environmental

microbiota from the seawater was also sampled. When cultures were sampled, triplicate ~1

L of 5.0-μm FSW was filtered onto a 0.22-μm Millipore filter to retain the environmental

microbiota. Full filter disks were then preserved in RNAlater and stored at -20°C before

DNA extractions.

Assaying bacterial communities

Total DNA was extracted from larval samples using the GeneJet Genomic DNA

Purification Kit (Thermo Scientific, Massachusetts, USA). For FSW samples, we extracted

eDNA using the FastDNA Spin Kit for Soil (MP Biomedical, Illkirch, France). DNA was

then quantified using the NanoDrop 2000 UV-Vis Spectrophotometer (Thermo Scientific,

Massachusetts, USA) and diluted to 5 ng•μL^{-1} using RNase/DNase-free water.

Bacterial sequences were amplified using universal primers for the V3/V4 regions

of the 16S rRNA gene (Forward: 5′ CTACGGGNGGCWGCAG, Reverse: 5′

GACTACHVGGGTATCTAATCC; Klindworth et al. 2013). Products were purified using

the Axygen AxyPrep Mag PCR Clean-up Kit (Axygen Scientific, New York, USA),

indexed via PCR using the Nextera XT Index Kit V2 (Illumina, California, USA), and then

purified again. At each of these three steps, fluorometric quantitation was performed using

a Qubit (Life Technologies, California, USA) and libraries were validated using a

Bioanalyzer High Sensitivity DNA chip (Agilent Technologies, California, USA). Illumina

MiSeq sequencing (v3, 2x300 bp paired-end reads) was performed at the University of

North Carolina at Charlotte. PCR recipe and thermal profiles are available in Table S2.

Forward and reverse sequences were paired and trimmed using PEAR (Zhang et al.

2014) and Trimmomatic (Bolger et al. 2014), respectively, converted from fastq to fasta

using a custom script, and chimeric sequences were detected using USEARCH (Edgar et

al. 2011) and removed using filter_fasta.py prior to the analysis of bacterial 16S rRNA

sequences. Using QIIME 1.9.1 (or Quantitative Insights Into Microbial Ecology; Caporaso

et al. 2010), bacterial 16S rRNA sequences were analyzed and grouped into operational

taxonomic units (OTUs) based on a minimum 97% similarity. The biom table generated

by the pick_open_reference_otus.py script was filtered of OTUs with ten or less reads as

well as sequences matching the chloroplast of cryptophytes (*i.e.*, *R. lens*; Carrier & Reitzel

2018).

Using the filtered biom table and "biom summarize-table" function to count total

sequences per sample, the rarefaction depth of 25,396 was determined and applied to all

subsequent analyses (Figure S2). Alpha diversity estimates were calculated using the

alpha_diversity.py script and compared statistically by a two-way analysis of variance

(ANOVA) in JMP (Pro v. 13). Beta diversity was calculated using unweighted and

weighted UniFrac (Lozupone & Knight 2005) as part of jackknifed_beta_diversity.py, and

compared using principal coordinate analyses (PCoA) that were then visualized in

EMPeror (Vazquez-Baeza et al. 2013) and stylized in Adobe Illustrator CS6. UPGMA-

based microbial dendrograms were generated using jackknifed_beta_diversity.py as part

of the PCoA analyses. Community similarity across species and developmental stages were

compared statistically using an analysis of similarity (ANOSIM) in QIIME as part of the

compare_categories.py script, with p<0.05 considered significant. Community

composition was generated using summarize_taxa_through_plots.py script and visualized

using Prism 7 (GraphPad Software, California, USA). Lastly, observed OTUs were

calculated using the alpha_diversity.py script, shared OTUs were calculated using the

shared_phylotypes.py script, and the Bray-Curtis index was calculated using the

beta_diversity.py script, and compared statistically using ANOVA in JMP (Pro v. 13).

A step-by-step listing of QIIME scripts used to convert raw reads to OTUs for

visualization of the data is located in Supplemental Note 1. The raw sequence reads as part

of this dataset are available on Dryad.

RESULTS

Community-level shifts across development

Urchin-specific bacterial communities (ANOSIM, unweighted: p=0.0011; weighted:

p<0.001; Figure 1) shifted significantly in OTU membership (ANOSIM, p=0.0013; Figure

1A; Figures S3-4) and composition (ANOSIM, p=0.0011; Figure 1B; Figures S2-3) during

embryonic and larval development. Moreover, community evenness (Shannon and

Robbins), diversity (Simpson), richness (Chao1), phylogenetic distance (Faith's), and

Fishers alpha varied significantly across developmental stages (p<0.0001 for all tests) and

amongst urchin species (p<0.0001 for all tests) (Figure S5; Table S2).

Unweighted and weighted UniFrac comparisons of the associated bacterial

communities for each urchin species suggests that community similarity largely group by

developmental stage (Figure 2). Bacterial communities were grouped largely by sequential

developmental stages with some specific clusters for: (i) early embryological stages, (ii)

late embryogenesis and early larval development, and (iii) later larval stages. The benthic

juveniles had more variable positioning on the microbial dendrograms: the bacterial

community of *S. purpuratus* juveniles were intermediate to early- and late-stage larvae, *M.*

franciscanus juvenile was most similar to eggs, and the *S. droebachiensis* juvenile were

most similar to late-stage larvae (Figure 2).

A total of nine classes of bacteria represented at least 1% of the community (Figure

2). Of these, the relative abundance of the γ-proteobacteria (Phylum: Proteobacteria), α-

proteobacteria (Phylum: Proteobacteria), and Flavobacteriia (Phylum: Bacteroidetes) were,

on average, each more than 10% of the community (Figure 2). For the γ-proteobacteria, the

relative abundance ranged from 26.6% to 61.0% for *S. purpuratus*, 12.2% to 62.9% for *M.*

franciscanus, and 39.6% to 88.7% for *S. droebachiensis* (Figure 2). Next, for the α-

proteobacteria, the relative abundance ranged from 7.7% to 42.4% for *S. purpuratus*,

11.0% to 31.1% for *M. franciscanus*, and 5.1% to 22.7% for *S. droebachiensis* (Figure 2).

Lastly, for the Flavobacteriia, the relative abundance ranged from 4.9% to 30.5% for *S.*

purpuratus, 5.1% to 36.1% for *M. franciscanus*, and 1.5% to 34.2% for *S. droebachiensis*

(Figure 2).

Dynamics of the associated bacterial species

Total OTUs in association with these confamilial echinoids differed significantly

between developmental stages (ANOVA, $p<0.0001$) for each species (ANOVA, *S.*

purpuratus: $p<0.0001$; *M. franciscanus*: $p<0.0001$; *S. droebachiensis*: $p<0.0001$; Figure

3A; Table S1). Unfertilized eggs, on average, associated with a similar number of OTUs

amongst species (~1,400 OTUs; ANOVA, $p=0.508$), which increased during embryonic

development (Figure 3A). Following the onset of feeding and larval development, the

number of OTUs subsequently decreased until metamorphosis (Figure 3A). The maximum

number of associated OTUs during development varied amongst species (*S. purpuratus*:

1,839; *M. franciscanus*: 1,723; *S. droebachiensis*: 1,864) and at which stages this occurred

(*S. purpuratus*: gastrula; *M. franciscanus*: 4-armed larva; *S. droebachiensis*: pre-hatched

blastula (Figure 3A).

A significant proportion of the OTUs associated with echinoid eggs did not remain

in association throughout embryonic and larval development for each urchin species

(ANOVA, *S. purpuratus*: p<0.0001; *M. franciscanus*: p<0.0001; *S. droebachiensis*:

p<0.0001; Figure 3B; Table S1). Following fertilization and early embryogenesis, ~52%

of OTUs from the egg remained in association while ~48% were lost (Figure 3B). In the

subsequent larval and juvenile stages, egg-associated OTUs were lost gradually, resulting

in juvenile urchins with ~29% of the egg-associated OTUs (Figure 3B). Moreover, when

transitioning between any two sequential developmental stages, ~48% of OTUs retained

while the rest were specific to either stage (Figure 3C). This result was not simply due to a

high diversity of OTUs in the egg samples because OTU number increased after the egg

stage for all species (Figure 3A).

Community dissimilarity (Bray-Curtis) between the sea urchin host and

environmental microbiota showed significant differences during embryonic and larval

developmental for each species (ANOVA, *S. purpuratus*: p<0.0001; *M. franciscanus*:

p<0.0001; *S. droebachiensis*: p<0.0001; Figure 3D, Figure S6, Table S1). During

development from egg to hatched blastula, the associated bacterial community became

more dissimilar from the environmental microbiota (Figure 3D), with the magnitude of

dissimilarity varying amongst species (Figure 3D; ANOVA, egg through hatched blastula,

each p≤0.0001). Furthermore, following development of the mouth and the onset of

feeding, late embryos and early stage larvae became more similar to the environmental

bacterial community (Figure 3D). The relative dissimilarity converged in the early larval

stages (Figure 3D; ANOVA, 4-armed larvae: p=0.406), which was the last sampled time

point for which we determined the bacterial community of the FSW. During divergence

and convergence events between the larval host and the environmental microbiota, the

environmental bacterial community was consistent throughout the experiment and did not

cluster with bacterial communities associated with the urchin (Figure S6).

Psychromonas and echinoid eggs

Bacteria in the genus _Psychromonas_ showed the highest abundance for any

bacterial genus for each urchin species at any one of the sampled stages, in contrast to their

relative rarity (0.7%) in the seawater (Figure 4; Tables S6-7). For each urchin species, the

abundance and diversity (Figure S7; ANOVA, p<0.0001 for each) of _Psychromonas_ was

highest in the unfertilized eggs and declined significantly following fertilization (ANOVA,

p<0.0001 for each), until being in very low abundance in 4-armed larvae and the

subsequent developmental stages (Figure 4). The magnitude of relative abundance varied

amongst urchin species (ANOVA, p<0.0001) and was directly proportion to egg diameter

(R^2=0.833). The OTUs within _Psychromonas_ totaled 96 for _S. purpuratus_, 100 for _M._

franciscanus, and 116 for _S. droebachiensis_, while the FSW had 81 (Figure 4). Of those,

nine of the ten most abundant _Psychromonas_ OTUs for each urchin species were shared

amongst urchin species.

DISCUSSION

Like many marine invertebrates, echinoid life cycles involve a distinct embryonic, larval, and juvenile stage. Our assay of the bacterial communities of three confamilial echinoids supports the hypothesis that the bacterial consortium is highly dynamic across developmental stages, with taxonomic membership and the relative abundance of those taxa shifting gradually from egg to juvenile. Furthermore, over the course of development, the bacterial community of urchin embryos remains distinct and diverges from the environmental microbiota until larvae begin feeding. Lastly, eggs of these echinoids appear to have a specific association with diverse OTUs that match the genus *Psychromonas*.

The bacterial communities associated with echinoplutei are highly dynamic when the larval holobiont faces variation in the abiotic and biotic environment (Carrier & Reitzel 2018, Carrier et al. 2018b). For example, following diet-restriction the bacterial communities of *S. purpuratus*, *M. franciscanus*, and *S. droebachiensis* each shift considerably from their well-fed counterparts, to potentially cope with reduced exogenous resources (Carrier & Reitzel 2018). When diet-restriction is prolonged and the larval host expresses phenotypic plasticity, the composition and structure of the bacterial community is specific to phenotype, a response correlated with the magnitude in which phenotypic plasticity is expressed (Carrier & Reitzel 2018).

Phenotypic plasticity for echinoid larvae, more broadly, is the interaction between phenotype, nutritional state, and the environment. Characterization of bacterial communities across developmental stages for these three confamilial echinoids extends the understanding of the complex interplay between developmental processes and the

surrounding environment in which these species develop. Particularly, after unfertilized

eggs establish a bacterial community, the composition of these prokaryotic species shift

gradually during embryogenesis and during transitions towards a feeding larva. In these

gradual community transitions, it was surprising that more coarse shifts were not observed

between major developmental milestones, such as when blastula hatch and shed the

fertilization envelope, larvae begin feeding, and/or when larvae metamorphose into benthic

juveniles. The latter may be because echinoid larvae undergoing metamorphosis reabsorb

some of the larval digestive tract (Chia & Burke 1978). The juveniles therefore may have

retained, instead of shed (Fieth et al. 2016), a portion of their bacterial community present

in late larval stages.

The influence of particular bacteria and bacterial communities on the development

of aquatic and terrestrial invertebrates is well-studied in few species (McFall-Ngai & Ruby

2000, Fraune & Bosch 2010, McFall-Ngai 2014, Nyholm & McFall-Ngai 2014).

Established invertebrate systems where bacteria play a significant functional role include

Vibrio fisheri-mediated maturation of the light organ in the bobtail squid *Euprymna*

scolopes (McFall-Ngai & Ruby 2000, McFall-Ngai 2014), immune regulation relating to

the epithelial bacteria associated with the freshwater cnidarian *Hydra* spp. (Fraune & Bosch

2010, Bosch 2013, Franzenburg et al. 2013), and the nutritional endosymbionts (*e.g.*,

Buchnera) of aphids (Douglas 1998). Provided the vast genomic and molecular foundation

established over the past several decades for the echinoid clade used in this study

(Kudtarkar & Cameron 2017), these confamilial urchins have the potential to be established

as a comparative system for beneficial animal-microbe interactions during development

(Buckley & Rast 2017, Heyland et al. 2018) and in bi-phasic life-histories. One route for

testing the impact of microbial symbionts on echinoid development may be through the egg-associated *Psychromonas*.

Eggs of echinoids have been used as a metric in larval biology, ecology, and evolution (Vance 1973, Wray & Raff 1991, Jaeckle 1995, Moran & McAlister 2009, Marshall et al. 2012) to make inferences about life history mode and transitions (Vance 1973, Collins & Moran 2018), fertilization ecology (Pennington 1985, Levitan 1995), and maternal inheritance (Byrne et al. 1999, Villinski et al. 2002). The high abundance of egg-associated *Psychromonas* and their loss and replacement may imply that this network of bacterial symbionts are functionally connected to early developmental processes.

Interestingly, egg size of echinoids is strongly correlated with total lipid content (Jaeckle 1995, Moran & McAlister 2009) and data here support that egg size is also highly correlated with the abundance of *Psychromonas*. Maternally-inherited lipids for *Dendraster excentricus*, a related echinoid found in the same geographical region of this study, are depleted by the same developmental stage that *Psychromonas* was undetectable here (Rendleman et al. 2018). Moreover, symbiotic bacteria associated with developing embryos of the deep-sea clam *Calyptogena okutanii* is spatiotemporally regulated and has the genetic components to process lipopolysaccharides (Ikuta et al. 2016). *Psychromonas* may, therefore, aid in lipid metabolism (Riley et al. 2008) during embryonic development or serve as a commensal symbiont that relies on the lipids leaching out maturing embryos.

An additional pattern we observed during embryonic development was that urchin-associated bacterial communities diverged from the environmental microbiota. Divergence of host-associated microbiota from the environmental community suggests that urchin embryos and larvae may select, at least, part of their microbial symbiont community.

Maintenance of host-specific symbionts may further suggest that these communities are

functionally important. To explicitly test how urchin larvae maintain a specific bacterial

community and their functional relevance during embryonic and larval development, a

multifaceted molecular and -omic (*e.g.*, meta-genomic and -transcriptomic) approach

would be needed (Williams & Carrier 2018). Approaches for characterizing the

mechanisms that establish and regulate host specificity may be adopted from well-

established biological systems, such as *Hydra* (Fraune et al. 2010, Fraune & Bosch 2010,

Franzenburg et al. 2013) or the winnowing of *Euprymna scolopes* to the luminescent

mutualist *Vibrio fisheri* (McFall-Ngai & Ruby 2000, McFall-Ngai 2002, Nyholm &

Mcfall-Ngai 2004).

Taken together, data presented here support the hypothesis that the bacterial

community associated with the developmental stages of three sea urchin species are

established prior to fertilization and shift gradually thereafter. The functional importance

and evolutionary significance of this bacterial community and the mechanisms for these

corresponding shifts remains unknown. Future experiments involving germ-free culturing

as well as manipulations of the host, members of the microbial community, and variable

culturing conditions will be needed to determine roles of associated microbiota. Results

from these experiments would serve as a key piece towards establishing echinoids as a

comparative system for understanding the interdependence of animal development and the

associated bacterial communities.

Acknowledgements. We thank Jason Hodin and Billie Swalla for providing laboratory

space; Morgan Eisenlord, Gustav Pauley, Katie Dobkowski, Joe Gaydos, and Richard

Strathmann for collecting urchins; Colette Feehan and Richard Strathmann for larval

rearing advise; Daniel Janies for sequencing resources; Karen Lopez for technical

assistance with sequencing; and the reviewers for improving the specificity and clarity of

this manuscript. TJC was supported by an NSF Graduate Research Fellowship, Charles

Lambert Memorial Endowment fellowship from the Friday Harbor Laboratories, and a

Sigma Xi Grants-in-Aid of Research grant, and AMR was supported by Human Frontier

Science Program Award RGY0079/2016.

 FIGURE LEGEND

Figure 1. Similarity amongst the associated bacterial community across development for

three species of echinoids. Community similarity for *Strongylocentrotus purpuratus*

(triangle), *Mesocentrotus franciscanus* (square), and *S. droebachiensis* (circle) for ten

developmental stages when considering the presence/absence of taxa (A) and their relative

abundance (B).

Figure 2. Relatedness of bacterial communities during development. Weighted UniFrac

dendrograms and corresponding class-level taxonomic plots for the bacterial communities

of *Strongylocentrotus purpuratus* (left), *Mesocentrotus franciscanus* (center), and *S.*

droebachiensis (right) across ten developmental stages.

Figure 3. Measures of OTU shifts during development. OTU dynamics across embryonic

and larval development for *Strongylocentrotus purpuratus* (purple), *Mesocentrotus*

franciscanus (red), and *S. droebachiensis* (green), and the average of the urchin species

(black), when considering (A) total observed OTUs, (B) percent of OTUs shared with

unfertilized eggs, (C) proportion of OTUs shared between developmental stages, and (D)

similarity with the environmental microbiota.

Figure 4. Decay in abundance of *Psychromonas* during development. Relative abundance

of the bacterial genus *Psychromonas* across embryonic and larval development for

Strongylocentrotus purpuratus (purple), *Mesocentrotus franciscanus* (red), *S.*

droebachiensis (green), and the seawater (blue) with the number of total OTUs.

LITERATURE CITED

Apprill A, Marlow HQ, Martindale MQ, Rappe MS (2012) Specificity of associations

between bacteria and the coral *Pocillopora meandrina* during early development.

Applied and Environmental Microbiology 78:7467-7475

Bates JM, Mittge E, Kuhlman J, Baden KN, Cheesman SE, Guillemin K (2006) Distinct

signals from the microbiota promote different aspects of zebrafish gut

differentiation. Dev Biol 297:374-386

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: A flexible trimmer for Illumina

sequence data. Bioinformatics 30:2114-2120

Bosch I (1992) Symbiosis between bacteria and oceanic clonal sea star larvae in the

western North Atlantic Ocean. Marine Biology 114:495-502

Bosch TCG (2013) Cnidarian-microbe interactions and the origin of innate immunity in

metazoans. Annual Review of Microbiology 67:499-518

Bright M, Bulgheresi S (2010) A complex journey: transmission of microbial symbionts.

Nat Rev Microbiol 8:218-230

Buckley KM, Rast JP (2017) An organismal model for gene expression networks in the

gut-associated immune response. Frontiers in Immunology 8:1297

Byrne M, Villinski JT, Cisternas P, Siegel RK, Popodi E, Raff RA (1999) Maternal factors

and the evolution of developmental mode: evolution of oogenesis in *Heliocidaris*

erythrogramma. Development Genes and Evolution 209:275-283

Cameron RA, Holland ND (1983) Electron microscopy of extracellular materials during

the development of a seastar, *Patiria miniata* (Echinoderm: Asteroidea). Cell

Tissue Research 243:193-200

Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer

N, Pena AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig

JE, Ley RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J,

Sevinsky JR, Turnbaugh PJ, Walters WA, Widmann J, Yatsunenko T, Zaneveld J,

Knight R (2010) QIIME allows analysis of high-throughput community sequencing

data. Nat Methods 7:335-336

Carrier TJ, Macrander J, Reitzel AM (2018a) A microbial perspective on the life-history

evolution of marine invertebrate larvae: if, where, and when to feed. Marine

Ecology 39:e12490

Carrier TJ, Reitzel AM (2018) Convergent shifts in host-associated microbial communities

across environmentally elicited phenotypes. Nature Communications 9:952

Carrier TJ, Wolfe K, Lopez K, Gall M, Janies DA, Byrne M, A.M. R (2018b) Diet-induced

shifts in the crown-of-thorns (*Acanthaster* sp.) larval microbiome. Marine Biology

165:157

Cerra A, Byrne M, Hoegh-Guldberg O (1997) Developments of the hyaline layer around

the planktonic embryos nad larvae of the asteroid *Patiriella calcar* and the presence

of associated bacteria. Invertebrate Reproduction and Development 31:337-343

Chia F-S, Burke RD (1978) Echinoderm metamorphosis: fate of larval structures. In: Chia

F-S, Rice ME (eds) Settlement and metamorphosis of marine invertebrate larvae:

proceedings of the Symposium on settlement and metamorphosis of marine

Invertebrate larvae, American Zoological Society Meeting. Elsevier

Collins R, Moran AL (2018) Evolutionary transitions in mode of development. In: Carrier

TJ, Reitzel AM, Heyland A (eds) Evolutionary Ecology of Marine Invertebrate

Larvae. Oxford University Press

Davidson SK, Stahl DA (2008) Selective recruitment of bacteria during embryogenesis of

an earthworm. The ISME Journal 2:510-518

Douglas AE (1998) Nutritional interactions in insect-microbial symbioses: aphids and their

symbiotic bacteria *Buchnera*. Annu Rev Entomol 43:17-37

Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R (2011) UCHIME improves

sensitivity and speed of chimera detection. Bioinformatics 27:2194-2200

Fieth RA, Gauthier M-EA, Bayes J, Green KM, Degnan SM (2016) Ontogenetic changes

in the bacterial symbiont community of the tropical demosponge Amphimedon

queenslandica: metamorphosis is a new beginning. Frontiers in Marine Science

3:228

Franzenburg S, Fraune S, Altrock PM, Künzel S, Baines JF, Traulsen A, Bosch TCG

(2013) Bacterial colonization of Hydra hatchlings follows a robust temporal

pattern. The ISME Journal 7:781

Fraune S, Augustin R, Anton-Erxleben F, Wittlieb J, Gelhaus C, Klimovich VB,

Samoilovich MP, Bosch TCG (2010) In an early branching metazoan, bacterial

colonization of the embryo is controlled by maternal antimicrobial peptides. P Natl

Acad Sci USA 107:18067-18072

Fraune S, Bosch TCG (2010) Why bacteria matter in animal development and evolution.

Bioessays 32:571-580

Galac MR, Bosch I, Janies DA (2016) Bacterial communities of oceanic sea star

(Asteroidea: Echinodermata) larvae. Marine Biology 163:162

Guillard RRL (ed) (1975) Culture of phytoplankton for feeding marine invertebrates.

Plenum Press, New York, USA

Heyland A, Schuh NW, Rast JP (2018) Sea urchin larvae as a model for postembryonic

development. In: Kloc M, Kubiak JZ (eds) Marine Organisms as Model Systems in

Biology and Medicine. Springer

Ikuta T, Igawa K, Tame A, Kuroiwa T, Kuroiwa H, Aoki Y, Takaki Y, Nagai Y, Ozawa

G, Yamamoto M, Deguchi R, Fujikura K, Maruyama T, Yoshida T (2016) Surfing

the vegetal pole in a small population: extracellular vertical transmission of an

'intracellular' deep-sea clam symbiont. Royal Society Open Science 3:160130

Jaeckle WB (1995) Variation in size, energy content, and biochemical composition of

invertebrate eggs: correlates to the mode of larval development. In: McEdward LR

(ed) Ecology of Marine Invertebrate Larvae. CRC Press

Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, Glockner FO (2013)

Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-

generation sequencing-based diversity studies. Nucleic Acids Research 41:e1

Kudtarkar P, Cameron RA (2017) Echinobase: an expanding resource for echinoderm

genomic information. Database 2017:bax074

Levitan DR (1995) The ecology of fertilization in free-spawning invertebrates. In:

McEdward LR (ed) Ecology of Marine Invertebrate Larvae. CRC Press

Lozupone C, Knight R (2005) UniFrac: a new phylogenetic method for comparing

microbial communities. Applied and Environmental Microbiology 71:8228-8235

Manahan DT, Davis JP, Stephens GC (1993) Bacteria-free sea urchin larvae: selective

uptake of neutral amino acids from seawater. Science 220:204-206

Marshall DJ, Krug PJ, Kupriyanova EK, Byrne M, Emlet RB (2012) The biogeography of

marine invertebrate life histories. Annual Review of Ecology, Evolution, and

Systematics 43:97-114

McFall-Ngai M (2014) The importance of microbes in animal development: lessons from

the squid-*Vibrio* symbiosis. Annual Review of Microbiology 68:177-194

McFall-Ngai M, Hadfield MG, Bosch TCG, Carey HV, Domazet-Loso T, Douglas AE,

Dubilier N, Eberl G, Fukami T, Gilbert SF, Hentschel U, King N, Kjelleberg S,

Knoll AH, Kremer N, Mazmanian SK, Metcalf JL, Nealson K, Pierce NE, Rawls

JF, Reid A, Ruby EG, Rumpho M, Sanders JG, Tautz D, Wernegreen JJ (2013)

Animals in a bacterial world, a new imperative for the life sciences. P Natl Acad

Sci USA 110:3229-3236

McFall-Ngai M, Ruby EG (2000) Developmental biology and marine invertebrate

symbioses. Current Opinion in Microbiology 3:603-607

McFall-Ngai MJ (2002) Unseen forces: the influence of bacteria on animal development.

Dev Biol 242:1-14

Mileikovsky SA (1971) Types of larval development in marine bottom invertebrates, their

distribution and ecological significance: a re-evalution. Marine Biology 10:193-

213

Moran AL, McAlister JS (2009) Egg size as a life history character of marine invertebrates:

Is it all it's cracked up to be? Biological Bulletin 216:226-242

Mortzfeld BM, Urbanski S, Reitzel AM, Kunzel S, Technau U, Fraune S (2015) Response
of bacterial colonization in *Nematostella vectensis* to development, environment
and biogeography. Environmental Microbiology 18:1764-1781

Nussbaumer AD, Fisher CR, Bright M (2006) Horizontal endosymbiont transmission in
hydrothermal vent tubeworms. Nature 441:345-348

Nyholm SV, McFall-Ngai M (2014) Animal development in a microbial world. In: Minelli
A, Pradeu T (eds) Towards a Theory of Development. Oxford University Press

Nyholm SV, Mcfall-Ngai MJ (2004) The winnowing: establishing the squid-*Vibrio*
symbiosis. Nat Rev Microbiol 2:632-642

Pennington JT (1985) The ecology of fertilization of echinoid eggs: the consequences of
sperm dilution, adult aggregation, and synchronous spawning. Biological Bulletin
169:417-430

Pradea T (2011) A mixed self: the role of symbionts in development. Biological Theory
6:80-88

Rendleman A, Rodriguez J, Ohanian A, Pace D (2018) More than morphology: differences
in food ration drive physiological plasticity in echinoid larvae. Journal of
Experimental Marine Biology and Ecology 501:1-15

Riley M, Staley JT, Danchin A, Wang TZ, Brettin TS, Hauser LJ, Land ML, Thompson
LS (2008) Genomics of an extreme psychrophile, *Psychromonas ingrahamii*. BMC
Genomics 9:210

Sharp KH, Distel D, Paul VJ (2012) Diversity and dynamics of bacterial communities in
early life stages of the Caribbean coral *Porites astreoides*. The ISME Journal 6:790-
801

Sipkema D, de Caralt S, Morillo JA, Al-Soud WA, Sørensen SJ, Smidt H, Uriz MJ (2015)

Similar sponge-associated bacteria can be acquired via both vertical and horizontal

transmission. Environmental Microbiology 17:3807-3821

Stewart FJ, Young CR, Cavanaugh CM (2008) Lateral symbiont acquisition in a maternally

transmitted chemosynthetic clam endosymbiosis. Mol Biol Evol 25:673-687

Strathmann MF (1987) Reproduction and development of marine invertebrates of the

northern Pacific coast: data and methods for the study of eggs, embryos, and larvae.

University of Washington Press

Thorson G (1950) Reproductive and larval ecology of marine bottom invertebrates.

Biological Reviews 25:1-45

Vance RR (1973) On reproductive strategies in marine benthic invertebrates. The

American Naturalist 107:339-352

Vazquez-Baeza Y, Pirrung M, Gonzalez A, Knight R (2013) EMPeror: a tool for

visualizing high-throughput microbial community data. Gigascience 2:16

Villinski JT, Villinski JC, Byrne M, Raff RA (2002) Convergent maternal provisioning

and life-history evolution in echinoderms. Evolution 56:1764-1775

Williams EA, Carrier TJ (2018) An -omics perspective on marine invertebrate larvae. In:

Carrier TJ, Reitzel AM, Heyland A (eds) Evolutionary Ecology of Marine

Invertebrate Larvae. Oxford University Press, Oxford, UK

Wray GA, Raff RA (1991) The evolution of developmental strategy in marine

invertebrates. Trends in Ecology & Evolution 6:45-50

Zhang J, Kobert K, Flouri T, Stamatakis A (2014) PEAR: A fast and accurate Illumina

Paired-End reAd mergeR. Bioinformatics 30:614-620

Figure 1
2

3
4
5

6 **Figure 2**
7

Strongylocentrotus purpuratus   *Mesocentrotus franciscanus*   *Strongylocentrotus droebachiensis*

Legend:
- Bacteriodetes: Saprospirae
- Bacteriodetes: Flavobacteriia
- Cyanobacteria: Chloroplast
- Planctomycetes: Planctomycetia
- Proteobacteria: Alphaproteobacteria
- Proteobacteria: Betaproteobacteria
- Proteobacteria: Deltaproteobacteria
- Proteobacteria: Gammaproteobacteria
- Verrucomicrobia: Verrucomicrobiae
- Unassigned
- Other

8
9
10

16   **Figure 4**
17

18
19
20

Figure S1. Locations of adult urchin collections throughout the Salish Sea with geographic reference points. Specifically, *Strongylocentrotus purpuratus* were hand-collected at Slip Point, Clallam Bay, WA; *S. droebachiensis* were hand-collected at low tide at Cattle Point, San Juan Island, WA; and, *M. franciscanus* were collected by SCUBA off Bell Island, WA.

Figure S2. Alpha rarefaction curves for three species of echinoids and of the seawater. Alpha rarefaction curves for *Strongylocentrotus purpuratus* (purple), *Mesocentrotus franciscanus* (red), and *S. droebachiensis* (green) and seawater (blue) based on rarefaction depth of 25,396.

16
17
18 **Figure S3.** Similarity between the associated bacterial community across development for
19 three species of echinoids. Community similarity for *Strongylocentrotus purpuratus* (top),
20 *Mesocentrotus franciscanus* (middle), and *S. droebachiensis* (bottom) for ten
21 developmental stages when considering the presence/absence of taxa (*i*) and their relative
22 abundance (*ii*).
23
24

Figure S4. Bacterial community dendrograms across development for three echinoid species. Unweighted (*i*) and weighted (*ii*) bacterial community dendrograms for *Strongylocentrotus purpuratus* (top), *Mesocentrotus franciscanus* (middle), and *S. droebachiensis* (bottom).

33
34
35 **Figure S5.** Various alpha diversity indices for three echinoid species across development.
36 Shannon (A), Simpson (B), Robbins (C), Chao1 (D), Fisher (E), and Faiths (F) indices
37 for *Strongylocentrotus purpuratus* (purple), *Mesocentrotus franciscanus* (red), *S.*
38 *droebachiensis* (green), and the average (black) at ten developmental stages.
39
40

41
42
43 **Figure S6.** Microbial dendrogram of developmental stages and seawater. Dendrogram of
44 the bacterial communities based on weighted UniFrac for the developmental stages of
45 *Strongylocentrotus purpuratus*, *Mesocentrotus franciscanus*, and *S. droebachiensis* as well
46 as the respective environmental bacterial communities.

47
48

49
50
51 **Figure S7.** Diversity of *Psychromonas* during development. Total operational taxonomic
52 units (OTUs) within the bacterial genus *Psychromonas* across embryonic and larval
53 development for *Strongylocentrotus purpuratus* (purple), *Mesocentrotus franciscanus*
54 (red), *S. droebachiensis* (green) as well as the average for the three sea urchin species and
55 the in the seawater.