Title

Geography, not food availability, reflects compositional differences in the bacterial

communities associated with larval sea urchins

Running title

Population-specific microbiome of urchin larvae

Authors

Tyler J. Carrier[1,*], Sam Dupont[2], and Adam M. Reitzel[1]

Affiliations

[1] Department of Biological Sciences, University of North Carolina at Charlotte, 9201

University City Blvd., Charlotte, NC 28223 USA

[2] Department of Biological and Environmental Sciences, University of Gothenburg, The

Sven Lovén Centre for Marine Infrastructure, Kristineberg, Sweden

[*] Corresponding author: tcarrie1@uncc.edu

Keywords

Microbiome; host-microbe; ecology; plasticity; echinopluteus; symbiosis

Abstract

Determining the principles underlying the assembly, structure, and diversity of symbiont communities remains a focal point of animal-microbiome research. Much of these efforts focus on taxonomic variation within or between animal populations, but rarely test the proportional impacts of ecological components that may affect animal-associated microbiota. Using the sea urchin *Strongylocentrotus droebachiensis* from the Atlantic and Pacific Oceans and the ability of their larvae to acclimate to heterogeneous feeding environments, we test the hypothesis that inter-population differences in the composition of animal-associated bacterial communities are more pronounced than intra-population variation. Despite significant differences in bacterial community structure within each *S. droebachiensis* larval population (based on food availability, time, development, and phenotype), variation in OTU membership and community composition correlated more strongly with geography. Moreover, across these populations in the Atlantic and Pacific Oceans, 20-30% of bacterial OTUs were specific to a single location while less than 10% were shared, and more than half of this community is predicted to contribute to metabolism of the larval holobiont. Taken together, these results suggest that inter-populational variation in symbiont communities is more pronounced than intra-populational variation. This difference may suggest that ecological variables over broad geographic scales may mask smaller scale ecological variables; however, explicit testing of this potential principle requires common garden experiments as well as molecular and functional manipulations.

Importance

A central question in animal-microbiome research that remains largely unanswered concerns what factors govern taxonomic composition of symbiotic communities and what drives these differences. Efforts to answer this question primarily focus on variation within a single population or differences between geographically distant populations. Using sea urchin larvae, we compare whether variation in bacterial communities is larger between locations than within when larvae are cultured at different food concentrations. While each population of sea urchin larvae associated with specific bacterial communities when cultured at different food concentration, these differences were relatively minor compared to variation based on geography. Our comparisons suggest that variation between populations d is more pronounced than variation within populations, and that larger scale ecological components may mask factors at smaller scales. Future experiments explicitly testing for the potential of host genotype using common garden experiments and functional manipulations will help determine mechanisms driving this variation.

Introduction

Acclimating to environmental variability through morphological, developmental, and/or physiological plasticity is a common trait of animals (1-7). Over the past decade, the appreciation for the role that animal-associated microbiota play in responding to environmental stressors has grown profoundly (8-14). When experiencing heterogeneous environments, an animal may recruit, expel, and/or shuffle the relative proportion of associated microbiota (15, 16) to assemble a community with particular molecular pathways for the environmental conditions (17-19).

Microbial communities associated with animals often vary in response to a multitude of abiotic and biotic factors, including temperature, salinity, diet quality and quantity, season, and habitat-type (see reviews by 8, 12). Of these, dietary responses are best studied and shown to have a major impact on the composition and functions of this community (20-23). When faced with prolonged diet restriction, the composition and diversity of microbiota associated with both invertebrate and vertebrate hosts shift considerably (13, 24), a response hypothesized to buffer against reduced exogenous nutrients.

Microbial communities associated with animals are also species-specific (*e.g.*, 13, 25, 26) and taxonomically variable across the geographical distribution of the host (27-31). Population-specific microbial communities are partially dependent on environmental conditions (32) and dispersal limitations (33). Despite this taxonomic variation, microbial communities can remain functionally similar due to shared genes across bacterial taxa (18, 34). The microbial communities of the bromeliad *Aechmea nudicaulis*, for example, are taxonomically dynamic but functionally consistent between individuals (19). Similarly, the bacterial community associated with the green alga *Ulva* spp. are too spatially and

temporally variable to define a 'core' community; however, nearly 70% of the microbial

genes are conserved across this broad geography (18).

Efforts to classify the principles underlying the assembly of host-associated microbial

communities have primarily been within (*e.g.*, 13, 20, 35-38) or between (*e.g.*, 27-31, 39)

populations (but see 40). This conceptual discontinuity highlights a gap in understanding

the patterns and principles underlying community assembly. To test for differences in

assembly structure, direct comparisons of animal-associated microbial communities within

and between populations are needed to determine the relative strength of these factors. The

fields of sensory, behavioral, and circadian biology have shown that organismal responses

to ecological variables are unequal and may be masked based on spatial significance (41-

45). Factors within and between populations, therefore, may have different proportional

impacts on the composition of host-associated microbiota.

Planktotrophic (feeding) larvae are one biological system to compare ecological

components and how they influence the composition of animal-associated microbiota.

Many planktotrophic larvae (*e.g.*, the pluteus of sea urchins) inhabit heterogeneous

nutritional environments and are morphologically and physiologically plastic to food

availability (2, 46-53). Larvae of the echinoid *Strongylocentrotus droebachiensis*, for

example, associate with phenotype-, diet-, and developmental stage-specific bacterial

communities (13, 54). *S. droebachiensis*, in particular, has an Arctic-boreal distribution

composed of a multiple ocean population with limited gene flow (55-57) and signs of

potential local adaption (58, 59).

Using the circumpolar distribution of *S. droebachiensis* and the ability of their larvae

to acclimate to heterogeneous feeding environments, we test the hypothesis that inter-

population differences in the composition of animal-associated bacterial communities are

more pronounced than intra-population variation. To test this, we performed differential

feeding experiments on *S. droebachiensis* larvae from populations in the Pacific and

Atlantic Oceans (Fig. 1) and used amplicon sequencing to profile the larval-associated

bacterial communities. We then used these sequences to predict the functional gene profiles

of the larval-associated bacterial communities to preliminarily test for similar community

function between urchin populations.

Results

Morphological plasticity of larval sea urchin

Diet-induced morphological plasticity was observed for *S. droebachiensis* larvae from

each population, with the pattern of expression being location-specific (ANOVA,

$p<0.0001$; Figs. S2-3). Diet-restricted larvae from the Salish Sea and Gulf of Maine, in

general, exhibited a higher post-oral arm to mid-body line ratio than *ad libitum* counterparts

(ANOVA, $p<0.0001$; Figs. S1-2; Tables S3-4), even though analyses of Gulf of Maine

larvae were partially confounded by developmental stage (Table S1). Larvae from the

North Sea, however, exhibited the opposite response: *ad libitum* feeding induced a higher

post-oral arm to mid-body line ratio than diet-restricted individuals (ANOVA, $p<0.0001$;

Fig. S1-2; Tables S3-4).

Structural comparisons of bacterial communities within populations

S. droebachiensis larvae from each population associated with a diet-specific bacterial

community (Figs. 2A-C, S4-7; Table S6). Larvae from the Salish Sea and Gulf of Maine

exhibited similar diet-specific community-level patterns in both alpha- and beta-diversity

(Figs. 2A-B, S4-6; Table S6), where the bacterial consortium of food-restricted larvae were

more similar to each other than to well-fed counterparts. Larvae from the North Sea, on the

other hand, exhibited the opposite response (Figs. 2C, S4, S7; Table S6), where diet-

specific bacterial communities were observed (Table S6) except that all food

concentrations were more similar to each other than to starved larvae (Fig. S7; Table S6).

The bacterial communities associated with larvae from each sea urchin population also

varied temporally in both alpha- and beta-diversity (Figs. 2D-F, S4, S8-10; Table S6).

Temporal changes in community diversity was reflective of culturing environment: larval

populations from the Atlantic Ocean increased in weeks one through three and declined in

week four while the larval population from the Pacific Ocean decreased in the first three

weeks and increased in week four. Compositional changes in the associated bacterial

communities for each larval population, on the other hand, experienced similar time-

dependent shifts from week one though four (Figs. 2D-F, S4).

Complementary to dietary and temporal shifts in community structure and

composition, larval-associated bacterial communities for *S. droebachiensis* varied

significantly with larval morphology (Fig. S11; Table S7) and developmental stage (Figs.

S8-10; Table S7). Variation in the larval-associated bacterial communities across food

availability, time, morphology, and development, however, appeared to not co-depend on

the bacterial taxa observed in the seawater, as the composition of larval-associated and

environmental microbiota were significantly different (Fig. S12; Table S7).

Differences in the bacterial communities of urchin populations

Despite differences in bacterial community structure within each *S. droebachiensis*

larval population, variation in OTU membership and composition correlated best with

geography (ANOSIM, unweighted UniFrac, $p < 0.001$; ANOSIM, weighted UniFrac,

p<0.001; Figs. 3, S13-15; Table S8). Relatedness of these communities supports that the

larval-associated microbiota in the Western and Eastern Atlantic Ocean were more similar

to each other than to the Pacific Ocean population (Figs. 3, S13). Moreover, bacterial

community diversity and structure were significantly different between these three

populations (Fisher's alpha: F-ratio = 142.2, $p<0.001$; Shannon equitability: F-ratio =

110.1, $p<0.001$; Faith's phylogenetic distance: F-ratio = 159.2, $p<0.001$; observed OTUs:

F-ratio = 124.2, $p<0.001$; Table S8). For each of the tested indices, bacterial communities

of Gulf of Maine larvae were taxonomically richest and most diverse while Salish Sea

larvae were the least rich and diverse, leaving North Sea larvae as intermediate (Tukey's

test, $p<0.05$ for each; Table S8).

S. droebachiensis larvae from these populations primarily associated with seven

bacterial classes from three phyla: Saprospirae and Flavobacteriia (Bacteroidetes), Alpha-

, Beta-, Delta-, and Gammaproteobacteria (Proteobacteria), and Verrucomicrobiae

(Verrucomicrobia) (Fig. 4A). Of these classes, Flavobacteriia, Alpha- and

Gammaproteobacteria represented at least 10% of the relative community for each

population, where Gammaproteobacteria was the most abundant clade at 20-50% (Fig.

4A). Of the hundreds of OTUs associated with *S. droebachiensis* larvae (Fig. S1), less than

10% were shared across populations, with 20-30% being specific to a single population

and 3-11% being shared between two populations (Fig. 4B). This shared community was

similar to the full communities for each population, in that it was composed primarily by

Flavobacteriia (24.7%), Alpha- (28.1%) and Gammaproteobacteria (34.9%) (Fig. 4A).

Predicted community function and representative taxa

Similar to the 16S rRNA sequence diversity, PICRUSt-generated (predicted)

metagenomic gene profiles for *S. droebachiensis* larvae were specific to urchin population

(ANOVA, p<0.0001; Fig. 5A). Predicted gene contents of PICRUSt-generated

metagenomes were, on average, ~54.9% metabolism, ~17.5% genetic information

processing, ~13.0% environmental information processing, and ~7.3% cellular processes

and signaling (Figs. 5B, S16). Total gene content was significantly different between

locations for metabolism (ANOVA, p<0.0001), cellular processes and signaling (ANOVA,

p<0.0001), and organismal systems (ANOVA, p<0.0001) (Fig. S16).

More than half of the genes of the predicted *S. droebachiensis* larval metagenome

related   to   metabolism   (Figs.   5B,   S15).   Gammaproteobacteria   (42.6%),

Alphaproteobacteria (31.4%), and Flavobacteriia (13.3%) were the major taxa presumed

to have metabolic functionality (Fig. 5C). Moreover, Betaproteobacteria (2.9%),

Verrucomicrobiae (1.7%), Deltaproteobacteria (1.2%), and Planctomycetia (1.2%) were

also minor taxa in this metabolism-prediction subset of the community (Fig. 5C).

Discussion

Principles underlying the assembly and diversity of host-associated microbial

communities has remained a focal point of host-microbiome research over the last few

decades (9, 10, 12, 15, 16, 28, 30, 60-67). Efforts to partition and categorize variation in

community composition have suggested that evolutionary history (10, 16, 61, 64, 65),

physiology (8, 68, 69), and environmental conditions (8, 12, 66, 70) primarily drive these

differences. The majority of this work, however, focuses on two spatiotemporal scales:

within (13, 20, 35-38) or between (27-31) host populations (but see 40). The potential to

distinguish and discern intra- and inter-location processes highlights a conceptual gap in

understanding the patterns and principles underlying microbiome assembly.

Comparing the bacterial communities associated with *S. droebachiensis* larvae across

feeding environments and geographical location supports three primary findings. First,

larvae from each population associate with bacterial communities correlated with food

availability, time, development, and phenotype. Second, despite variation in the microbiota

within each population of urchin larvae, the composition of *S. droebachiensis* larval-

associated bacterial communities best correlated with geography. Third, predicted

metagenomic profiles for *S. droebachiensis* larval-associated bacterial communities were

population-specific, with more than half the predicted gene pool relating to metabolism.

Larval marine invertebrates associate with a microbial community that is diverse and

dynamic yet specific to host species and distinct from the environmental microbiota (13,

35, 38, 54, 71). The bacterial community associated with larval echinoderms, in particular,

shifts with food availability (13, 35), embryonic and larval development (13, 54),

phenotype (13), disease (71), and physiology (68). Common practice for investigating

larval-microbe interactions involve single location studies of a widely distributed taxa (see

13, 35, 38, 54). The data presented here suggest that community-level shifts in response to

biotic (development and phenotype) and abiotic (food availability and time) factors occur

at multiple locations across a population with limited gene flow. Three widely-distributed

sea urchin species (including *S. droebachiensis*) have previously been shown to associate

with food-elicited phenotype-specific bacterial communities (13). It may, therefore, be

hypothesized that phenotype-specific microbiota for these and, potentially, other species

of larval echinoderms (*e.g.*, *Acanthaster* sp. in the Pacific Ocean, including on the Great

Barrier Reef; 35, 72-74) occur throughout the geographical distribution of the species.

Marine invertebrate larvae are not unique in their ability to respond to environmental

variability by exhibiting shifts in the associated bacterial community (8, 12). Several clades

of marine and terrestrial vertebrates (*e.g.*, mammals, reptiles, and fish) respond to diet

variability and abiotic stress by restructuring their gut microbiota that, in turn, are likely to

differ in community function (8, 20, 21, 69, 75-77). Similar to larval marine invertebrates,

marine and terrestrial vertebrates can be widely-distributed and, thus, multiple genetically

distinct populations may exhibit similar community-levels shifts in microbiota at multiple

locations across these populations. To our knowledge, however, parallel responses between

populations for vertebrates and other taxa have not yet been documented.

Inter-populational variation in the composition of *S. droebachiensis* larval-associated

bacterial communities was more pronounced than intra-populational differences. While the

mechanisms driving this result is unknown, genetic and ecological differences between *S.*

droebachiensis populations are likely contributing factors (32, 33). *S. droebachiensis* from

the Salish Sea, Gulf of Maine, and North Sea diverged following larval-mediated restricted

gene flow (55, 56, 78, 79). Moreover, each urchin population inhabit environments that

differ in physical (80-82) and biological (83-85) factors. Inter-populational variation in the

bacterial communities associated with *S. droebachiensis* larvae may, therefore, have been

due to differences in host genetics and the oceanographic environment, a result that

requires future explicit testing using common garden experiments and molecular

manipulations (86).

Population-specific bacterial communities for *S. droebachiensis* larvae is consistent

with the hypothesis that host-associated microbial communities are taxonomically variable

across host geography (27-31). Studies using diverse animal taxa (*e.g.*, sponges, corals,

fish, and humans) support that the environment, not genotype, principally drives

community differences (87-90). Genetics of *S. droebachiensis* indicates that individuals from the Salish Sea and Gulf of Maine sub-populations are more related to each other than to the North Sea sub-population (55-57). Microbial communities from these populations, however, suggest the contrary: larvae from the Gulf of Maine and North Sea are more similar to each other than to Salish Sea larvae. This reinforces that environment (or, here, geography) is a larger influence on community assembly than genetic similarity. The mechanistic linkage between host genetics, the environment, and microbial community is unknown, but based on the predicted larval metagenomics, we suspect it is related to the metabolic activity of and between the larval host and bacterial symbionts.

Metagenomes predicted from 16S rRNA profiles for *S. droebachiensis* larvae also suggest population specificity. From these data, what was perhaps most notable is that the predicted metagenomes primarily include genes relating to metabolism. Planktotrophic (feeding) larvae, such as the echinopluteus of *S. droebachiensis*, are traditionally thought to acquire the energy for larval development by capturing exogenous particulates (*e.g.*, phytoplankton and bacteria; 91). A growing body of research, however, has suggested that planktotrophic larvae also utilize diverse 'alternative' nutritional resources, such as neutral amino acids (92, 93), kelp detritus (94), and, perhaps, phototrophic symbionts (35, 95, 96). Our predicted metagenomic gene profiles suggest the modes of nutrient acquisition for larvae may relate to the bacterial symbiont community. This observation may be of particular importance because planktotrophic larvae often inhabit food-limited environments (97-99). To decrease mortality due to starvation and prolonged exposure to pelagic predators (100-102), a metabolic input from the symbiont community may aid in accelerating development and serve as a physiological buffer that complements metabolic depression induced by diet-restriction (48).

Metabolic activity by animal-associated microbial communities benefiting the host is

widespread. The growth and reproduction of aphids, for example, is reliant on the amino

acids provisioned by the obligate endosymbiont *Buchnera* (103) while autotrophic and

heterotrophic bacteria drive carbon metabolism in sponges from oligotrophic seas (26).

Several bacterial phyla associated with the woodrat *Neotoma* gut (21, 104) and *Daphnia*

(105), on the other hand, are responsible for toxin degradation, allowing the host to access

and subsequently absorb nutrients from otherwise inaccessible diet. While a microbe-based

metabolism aiding in the digestion of diverse diets is not supported functionally for marine

invertebrate larvae, the wealth of data from a diversity of animal system and our 16S rRNA-

derived metagenomes merit testing.

By comparing host-associated bacterial communities across feeding environments and

biogeography in larval sea urchins, we propose that inter-populational variation in

symbiont communities is more biologically significant than intra-populational variation.

This difference suggests that larger scale ecological variables that correspond to different

geographic locations may mask smaller scale ecological variables (*e.g.*, feeding, time,

development, and phenotype). Explicit testing with diverse animal taxa by experimental

manipulations and functional characterization are merited to understand the priority of

exogenous factors in the composition of host-associated microbial communities.

Experimental Procedures

Adult urchin collection and larval rearing

Adult *S. droebachiensis* were collected from populations in the North Sea in March

2015, the Salish Sea in April 2016, and the Gulf of Maine in February 2017 (Fig. 1).

Individuals from the North Sea were collected by divers in Drøbak, Norway (59°39' N,

10°37' E) and transported in cold, aerated seawater to the Sven Lovén Centre for Marine

Infrastructure (Kristineberg, Sweden). Urchins were maintained in natural seawater and

fed *ad libitum* on a live mix of *Ulva lactuca* and *Laminaria* spp. collected from the

Kristineberg shoreline. Urchins from the Salish Sea were hand-collected at low tide at

Cattle Point, San Juan Island, USA (48°27' N, 122°57' W), transferred to the Friday Harbor

Laboratories (FHL; Friday Harbor, WA, USA) within one hour, suspended in sub-tidal

cages off the dock at FHL, and fed *Nereocystis* spp. *ad libitum*. Lastly, individuals from

the Gulf of Maine were collected by divers in Frenchman Bay, Maine (44°25' N 68°12'

W), shipped overnight to the Darling Marine Center (Walpole, ME, USA), and were

maintained in flow-through aquaria and fed *Saccharina latissima ad libitum*.

Within two weeks of collections, adult urchins were spawned by a one to two mL

intracoelomic injection of 0.50 M potassium chloride. For each population, gametes from

three males and females were separately pooled. Fertilization of eggs and larval rearing

followed Strathmann (106), with the exception that embryos and larvae were reared using

5.0-μm filtered seawater (FSW) to include the environmental microbiota. Briefly, embryos

were incubated in 1 L of FSW at ambient temperature and salinity. Two hours post-

fertilization, seawater was decanted to remove excess sperm. Embryos were then

transferred to triplicate 3 L jars and diluted to a density of two individuals•mL^{-1}. Beginning

at the prism stage, larvae were provided *Rhodomonas lens* at 10,000 cells•mL^{-1}, for which

was made growth medium-free by centrifugation and resuspension in FSW prior to the

introduction to larval cultures. Larval cultures were subsequently diluted to one larva per

two mL at advanced developmental stages. Larval cultures were given 90-95% water

changes every other day and *R. lens* was replenished at 10,000 cells•mL^{-1}.

320 Monocultures of *R. lens* were grown in f/2 media at room temperature with a

321 combination of ambient and artificial lighting for 24 hours per day (107).

323 *Experimental feeding and larval morphometrics*

324 At 48 hours post-fertilization, prism-stage larvae were divided into twelve replicate jars

325 that were sub-divided into four experimental feeding treatments: 10,000, 1,000, 100, or 0

326 cells•mL^{-1} of *R. lens*. For each experiment, larvae fed 10,000 cells•mL^{-1} were reared

327 through metamorphosis while diet-restricted (1,000 and 100 cells•mL^{-1}) and starved larvae

328 were cultured until developmental stasis (Table S1). Larvae (n=100) from each replicate

329 for each treatment were sampled weekly. Immediately after sampling, larval samples were

330 concentrated into a pellet using a microcentrifuge, the FSW was removed with a sterile

331 glass pipette, and pelleted larvae were then preserved in RNAlater (Thermo Scientific,

332 Massachusetts, USA) and stored at -20 °C before DNA extractions.

333 Complementary to sampling *S. droebachiensis* larvae, the environmental microbiota

334 from the seawater was also sampled. When larval cultures were sampled, triplicate ~1-L

335 of 5.0-µm FSW was filtered onto a 0.22-µm Millipore filter to retain the environmental

336 microbiota. Full filter disks were then preserved in RNAlater and stored at -20 °C before

337 DNA extractions.

338 In addition to sampling larvae to assay the associated bacterial communities, twenty

339 larvae (n=20) from a single replicate of each dietary treatment were sampled for

340 morphometric analysis. Larvae were imaged using a compound microscope (Salish Sea:

341 Nikon Eclipse E600; camera: QImaging MicroPublisher 5.0 RTV; Gulf of Maine: Zeiss

342 Primo Star HD digital microscope; North Sea: Leica stereomicroscope) and morphometrics

343 (length of larval body, post-oral arms, and stomach area; Figs. S1-2) were performed using

ImageJ, v. 1.9.2 (108). Statistical differences in larval size and stomach volume were

compared using a two-way analysis of variance (ANOVA; JMP Pro v. 13) to test for

differences over time and across feeding conditions. Moreover, differences between sites

were compared with a one-way ANOVA. Where statistical differences were observed

($p<0.05$), a Tukey's post-hoc test were used to determine the affect at each time point and

for each diet.

Assaying microbial communities

Total DNA was extracted from larval samples using the GeneJet Genomic DNA

Purification Kit (Thermo Scientific, Massachusetts, USA). For FSW samples, eDNA was

extracted using the FastDNA Spin Kit for Soil (MP Biomedical, Illkirch, France). DNA

was then quantified using the NanoDrop 2000 UV-Vis Spectrophotometer (Thermo

Scientific, Massachusetts, USA) and diluted to 5 ng•µL^{-1} using RNase/DNase-free water.

Bacterial sequences were amplified using universal primers for the V3/V4 regions of

the 16S rRNA gene (Forward: 5′ CTACGGGNGGCWGCAG, Reverse: 5′

GACTACHVGGGTATCTAATCC; 109). Products were purified using the Axygen

AxyPrep Mag PCR Clean-up Kit (Axygen Scientific, New York, USA), indexed via PCR

using the Nextera XT Index Kit V2 (Illumina, California, USA), and then purified again.

At each of these three steps, fluorometric quantitation was performed using a Qubit (Life

Technologies, California, USA) and libraries were validated using a Bioanalyzer High

Sensitivity DNA chip (Agilent Technologies, California, USA). Illumina MiSeq

sequencing (v3, 2x300 bp paired-end reads) was performed at the University of North

Carolina at Charlotte. See, Table S2 for PCR recipe and thermal profiles.

Forward and reverse sequences were paired and trimmed using PEAR (110) and

Trimmomatic (111), respectively, and converted from fastq to fasta using custom script.

Chimeric sequences were then detected using USEARCH (112) and removed using

filter_fasta.py prior to analysis of bacterial 16S rRNA sequences. Using QIIME 1.9.1 (113)

and Greengenes (v. 13.5), bacterial 16S rRNA sequences were analyzed and grouped into

operational taxonomic units (OTUs) based on a minimum 97% similarity. The biom table

generated by pick_open_reference_otus.py was filtered of OTUs with ten or less sequences

as well as sequences matching the chloroplast of cryptophytes (*i.e.*, *R. lens*; as per 13).

Using the filtered biom table and "biom summarize-table" to count total sequences per

sample, the rarefaction depth of 3,193 reads was determined and applied to all subsequent

analyses (Fig. S3). Alpha diversity (*i.e.*, Fisher's alpha, Shannon equitability, Faith's

phylogenetic distance, and observed OTUs) was calculated using alpha_diversity.py and

compared statistically using one-way ANOVAs in JMP. Beta diversity was calculated

using the unweighted and weighted UniFrac (114) as part of jackknifed_beta_diversity.py.

These values were compared using principal coordinate analyses (PCoA) and recreated

using make_2d_plots.py and stylized for presentation in Adobe Illustrator CS6.

Community similarity across phenotypes, dietary states, developmental stages, and

biogeography were compared statistically using an analysis of similarity (ANOSIM),

permutational multivariate analysis of variance using distance matrices (ADONIS), and

permutational multivariate analysis of variance (PERMANOVA) using

compare_categories.py. The shared OTUs between urchin populations was determined

using compute_core_microbiome.py and shared_phylotypes.py. Average bacterial

community for each urchin population as well as across populations and the shared

community was generated using summarize_taxa_through_plots.py, visualized with Prism

7 (GraphPad Software), and stylized for presentation in Adobe Illustrator CS6.

A step-by-step listing of QIIME scripts used to convert raw reads to OTUs for

visualization of these data is located in Note S1.

Functional predictions using PICRUSt

For the QIIME-generated OTU (*i.e.*, biom) table to be compatible with PICRUSt (115),

all *de novo* OTUs were filtered according to Greengenes (v. 13.5). This 'closed' OTU table

(that retained 50.9% of original OTUs) was normalized by copy number, upon which

metagenomic gene profiles were predicted and categorized by biological function (*i.e.*,

KEGG categories). The PICRUSt output was made compatible for Statistical Analysis of

Metagenomic Profiles (STAMP; 116) using biom_to_stamp.py in Microbiome Helper

(117) and analyzed for population-level differences. A STAMP-generated principle

coordinate analysis was compared statistically with an ANOVA within the STAMP

program. Functional profiles were visualized with Prism 7 (GraphPad Software) and

stylized for presentation in Adobe Illustrator CS6. Lastly, metagenome_contributions.py

and custom scripts to extract taxonomic information were used to determine which

bacterial taxa contributed to predicted genes categories of interest.

A step-by-step listing of PICRUSt scripts used to convert from QIIME and the

subsequent data analysis is located in Note S1.

Acknowledgements

We thank the staffs of the Friday Harbor Laboratories, Darling Marine Center, and

Sven Lovén Centre for Marine Infrastructure for facility access and logistical assistance;

Jason Hodin and Billie Swalla for providing laboratory space; Richard Strathmann for

collection assistance; Daniel Janies for sequencing resources; Karen Lopez for technical

assistance with sequencing; and Jason Macrander for bioinformatics assistance.

T.J.C. was supported by an NSF Graduate Research Fellowship, Charles Lambert

Memorial Endowment fellowship from the Friday Harbor Laboratories, and a Sigma Xi

Grants-in-Aid of Research grant; S.D. was funded by the Centre for Marine Evolutionary

Biology and supported by a Linnaeus grant from the Swedish Research Councils VR and

Formas; and A.M.R. was supported by Human Frontier Science Program Award

RGY0079/2016.

Fig. Legends

Fig. 1. Location of *Strongylocentrotus droebachiensis* experiments. Cartoon representation

of adult populations and the geographical distribution for where differential feeding

experiments were conducted.

Fig. 2. Dietary and temporal shifts in the bacterial communities associated with

Strongylocentrotus droebachiensis larvae. Community similarity based on *S.*

droebachiensis larval-associated bacterial communities based on food availability (A-C)

over a multi-week exposure (D-F) for populations from the Salish Sea (A, D), Gulf of

Maine (B, E), and North Sea (C, F). Comparisons between food availability (A-C) and over

time (D-F) are based on weighted UniFrac comparisons.

Fig. 3. Similarity of the bacterial communities associated with *Strongylocentrotus*

droebachiensis larvae from three populations. Community similarity of *S. droebachiensis*

larval-associated bacterial communities between three geographic locations, with variation

due to diet, time and development, and phenotype collapsed by site. Comparisons between

populations are based on (A) unweighted and (B) weighted UniFrac comparisons.

Fig. 4. Bacterial taxa associated with *Strongylocentrotus droebachiensis* larvae. (A) Class-

level profiles of the bacterial communities associated with *S. droebachiensis* larvae from

the Salish Sea, Gulf of Maine, and North Sea as well as the mean community across

populations and the shared taxa between each population. (B) Distribution of OTUs

associated with *S. droebachiensis* larvae uniquely found in each population, common

between two populations, and shared between all populations.

Fig. 5. Predicted *Strongylocentrotus droebachiensis* metagenome. (A) Community

similarity of level-1 KEGG categories for PICRUSt-generated *S. droebachiensis* larval

metagenomes. (B) Profiles of level-1 KEGG categories from the Salish Sea, Gulf of Maine,

and North Sea as well as the mean community across populations and the shared taxa

between each population. (C) Relative proportions of bacterial classes linked to the

predicted metabolic functionality of the PICRUSt-generated *S. droebachiensis* larval

metagenome.

References

1. Bradshaw A. 1965. Evolutionary significance of phenotypic plasticity in plants. Advances in Genetics 13:115-155.

2. Boidron-Metairon IF. 1988. Morphological plasticity in laboratory-reared echinoplutei of *Dendraster excentricus* (Eschscholtz) and *Lytechinus variegatus* (Lamarck) in response to food conditions. Journal of Experimental Marine Biology and Ecology 119:31-41.

3. Sterns S. 1989. The evolutionary significance of phenotypic plasticity. BioScience 39:436-445.

4. DeWitt TJ, Sih A, Wilson DS. 1998. Costs and limits of phenotypic plasticity. Trends in Ecology & Evolution 13:77-81.

5. Schlichting CD, Smith H. 2002. Phenotypic plasticity: linking molecular mechanisms with evolutionary outcomes. Evolutionary Ecology 16:189-211.

6. West-Eberhard M. 2003. Developmental Plasticity and Evolution. Oxford University Pres, Oxford, UK.

7. Miner BG, Sultan SE, Morgan SG, Padilla DK, Relyea RA. 2005. Ecological consequences of phenotypic plasticity. Trends in Ecology & Evolution 20:685-692.

8. Kohl KD, Carey HV. 2016. A place for host-microbe symbiosis in the comparative physiologist's toolbox. Journal of Experimental Biology 219:3496-3504.

9. Macke E, Tasiemski A, Massol F, Callens M, Decaestecker E. 2017. Life history and eco-evolutionary dynamics in light of the gut microbiota. Oikos 126:508-531.

10. Shapira M. 2016. Gut microbiotas and host evolution: scaling up symbiosis. Trends in Ecology & Evolution 31:539-549.

483   11.   Theis KR, Dheilly NM, Klassen JL, Brucker RM, Baines JF, Bosch TCG, Cryan JF,

484   Gilbert SF, Goodnight CJ, Lloyd EA, Sapp J, Vandenkoornhuyse P, Zilber-

485   Rosenberg I, Rosenberg E, Bordenstein SR. 2016. Getting the hologenome concept

486   right: an eco-evolutionary framework for hosts and their microbiomes. mSystems

487   1:e00028-16.

488   12.   Carrier TJ, Reitzel AM. 2017. The hologenome across environments and the

489   implications of a host-associated microbial repertoire. Frontiers in Microbiology

490   8:802.

491   13.   Carrier TJ, Reitzel AM. 2018. Convergent shifts in host-associated microbial

492   communities across environmentally elicited phenotypes. Nature Communications

493   9:952.

494   14.   Apprill A. 2017. Marine animal microbiomes: toward understanding host-

495   microbiome interactions in a changing ocean. Frontiers in Marine Science 4:222.

496   15.   Bordenstein SR, Theis KR. 2015. Host biology in light of the microbiome: ten

497   principles of holobionts and hologenomes. PLoS Biology 13:e1002226.

498   16.   Zilber-Rosenberg I, Rosenberg E. 2008. Role of microorganisms in the evolution of

499   animals and plants: the hologenome theory of evolution. FEMS Microbiology

500   Reviews 32:723-735.

501   17.   Burke B, Steinberg P, Rusch D, Kjelleberg S, Thomas T. 2011. Bacterial community

502   assembly based on functional genes rather than species. Proceedings of the National

503   Academy of Sciences of the United States of America 108:14288-14293.

504   18.   Roth-Schulze AJ, Pintado J, Zozaya-Valdés E, Cremades J, Ruiz P, Kjelleberg S,

505   Thomas T. 2018. Functional biogeography and host specificity of bacterial

communities associated with the marine green alga *Ulva* spp. Molecular Ecology

27:1952-1965.

19. Louca S, Jacques SMS, Pires APF, Leal JS, Srivastava DS, Parfrey LW, Farjalla VF,

Doebeli M. 2016. High taxonomic variability despite stable functional structure

across microbial communities. Nature Ecology & Evolution 1:15.

20. David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, Ling

AV, Devlin AS, Varma Y, Fischbach MA, Biddinger SB, Dutton RJ, Turnbaugh PJ.

2014. Diet rapidly and reproducibly alters the human gut microbiome. Nature

505:559-564.

21. Kohl KD, Dearing MD. 2012. Experience matters: prior exposure to plant toxins

enhances diversity of gut microbes in herbivores. Ecology Letters 15:1008-1015.

22. Rosenberg E, Zilber-Rosenberg I. 2016. Interaction between the microbiome and

diet: the hologenome concept. Journal of Nutrition & Food Sciences 6:5.

23. Sonnenburg ED, Smits SA, Tikhonov M, Higginbottom SK, Wingreen NS,

Sonnenburg JL. 2016. Diet-induced extinctions in the gut microbiota compound over

generations. Nature 529:212-215.

24. Kohl KD, Amaya J, Passement CA, Dearing MD, McCue MD. 2014. Unique and

shared responses of the gut microbiota to prolonged fasting: a comparative study

across five classes of vertebrate hosts. FEMS Microbiology Ecology 90:883-894.

25. Fraune S, Bosch TCG. 2007. Long-term maintenance of species-specific bacterial

microbiota in the basal metazoan *Hydra*. Proceedings of the National Academy of

Sciences of the United States of America 104:13146-13151.

26. Schmitt S, Tsai P, Bell J, Fromont J, Ilan M, Lindquist N, Perez T, Rodrigo A, Schupp

PJ, Vacelet J, Webster N, Hentschel U, Taylor MW. 2012. Assessing the complex

sponge microbiota: core, variable and species-specific bacterial communities in

marine sponges. ISME Journal 6:564-576.

27. Dishaw LJ, Flores-Torres J, Lax S, Gemayel K, Leigh B, Melillo D, Mueller MG,

Natale L, Zucchetti I, Santis RS, Pinto MR, Litman GW, Gilbert JA. 2014. The gut

of geographically disparate *Ciona intestinalis* harbors a core microbiota. PLoS One

9:e93386.

28. Huang B-H, Chang C-W, Huang C-W, Gao J, Liao P-C. 2018. Composition and

functional specialists of the gut microbiota of frogs reflect habitat differences and

agricultural activity. Frontiers in Microbiology 8:2670.

29. Marino CM, Pawlik JR, López-Legentil S, Erwin PM. 2017. Latitudinal variation in

the microbiome of the sponge *Ircinia campana* correlates with host haplotype but not

anti-predatory chemical defense. Marine Ecology Progress Series 565:53-66.

30. Marzinelli EM, Campbell AH, Valdes EZ, Vergés A, S. N, Wernberg T, Bettignies

Td, Bennett S, Caporaso JG, Thomas T, Steinberg PD. 2015. Continental-scale

variation in seaweed host-associated bacterial communities is a function of host

condition, not geography. Environmental Microbiology 17:4078-4088.

31. Mortzfeld BM, Urbanski S, Reitzel AM, Kunzel S, Technau U, Fraune S. 2015.

Response of bacterial colonization in *Nematostella vectensis* to development,

environment and biogeography. Environmental Microbiology 18:1764-1781.

32. Pantos O, Bongaerts P, Dennis PG, Tyson GW, Hoegh-Guldberg O. 2015. Habitat-

specific environmental conditions primarily control the microbiomes of the coral

Seriatopora hystrix. ISME Journal 9:1916-1927.

33. Moeller AH, Suzuki TA, Lin D, Lacey EA, Wasser SK, Nachman MW. 2017.

Dispersal limitation promotes the diversification of the mammalian gut microbiota.

Proceedings of the National Academy of Sciences of the United States of America

114:13768-13773.

34. Louca S, Polz MF, Mazel F, Albright MBN, Huber JA, O'Connor MI, Ackermann

M, Hahn AS, Srivastava DS, Crowe SA, Doebeli M, Parfrey LW. 2018. Function and

functional redundancy in microbial systems. Nature Ecology & Evolution 2:936-943.

35. Carrier TJ, Wolfe K, Lopez K, Gall M, Janies DA, Byrne M, A.M. R. 2018. Diet-

induced shifts in the crown-of-thorns (*Acanthaster* sp.) larval microbiome. Marine

Biology 165:157.

36. Lokmer A, Wegner KM. 2015. Hemolymph microbiome of Pacific oysters in

response to temperature, temperature stress and infection. The ISME Journal 9:670-

682.

37. Webster NS, Botte ES, Soo RM, Whalan S. 2011. The larval sponge holobiont

exhibits high thermal tolerance. Environmental Microbiology Reports 3:756-762.

38. Webster NS, Soo R, Cobb R, Negri AP. 2011. Elevated seawater temperature causes

a microbial shift on crustose coralline algae with implications for the recruitment of

coral larvae. ISME Journal 5:759-770.

39. Yatsunenko T, Rey FE, Manary MJ, Trehan I, Dominguez-Bello MG, Contreras M,

Magris M, Hidalgo G, Baldassano RN, Anokhin AP, Heath AC, Warner B, Reeder J,

Kuczynski J, Caporaso JG, Lozupone CA, Lauber C, Clemente JC, Knights D, Knight

R, Gordon JI. 2012. Human gut microbiome viewed across age and geography.

Nature 486:222-227.

40. Webster TMU, Consuegra S, Hitchings M, de Leaniz CG. 2018. Inter-population

variation in the Atlantic salmon microbiome reflects environmental and genetic

diversity. Applied and Environmental Microbiology 84:e00691-18.

41.   Hodin J, Ferner MC, Heyland A, Gaylord B. 2017. I feel that! Fluid dynamics and

sensory aspects of larval settlement across scales. *In* Carrier TJ, Reitzel AM, Heyland

A (ed), Evolutionary Ecology of Marine Invertebrate Larvae. Oxford University

Press, Oxford, UK.

42.   Beauchard O, Veríssimo H, Queirós AM, Herman PMJ. 2017. The use of multiple

biological traits in marine community ecology and its potential in ecological indicator

development. Ecological Indicators 76:81-96.

43.   Kerr JR, Kemp PS. In Press. Masking a fish's detection of environmental stimuli:

application to improving downstream migration at river infrastructure. Journal of

Fish Biology:1-10.

44.   Li Y, He N, Hou J, Xu L, Liu C, Zhang J, Wang Q, Zhang X, Wu XB. 2018. Factors

influencing leaf chlorophyll content in natural forests at the biome scale. Frontiers in

Ecology and the Environment 6:64.

45.   Yang Y, Niu H, Xiao L, Lin Q, Han B-P, Naselli-Flores L. 2018. Spatial

heterogeneity of spring phytoplankton in a large tropical reservoir: could mass effect

homogenize the heterogeneity by species sorting? Hydrobiologia 819:109-122.

46.   Adams DK, Sewell MA, Angerer RC, Angerer LM. 2011. Rapid adaptation to food

availability by a dopamine-mediated morphogenetic response. Nature

Communications 2:592.

47.   Byrne M, Sewell MA, Prowse TAA. 2008. Nutritional ecology of sea urchin larvae:

influence of endogenous and exogenous nutrition on echinopluteal growth and

phenotypic plasticity in *Tripneustes gratilla*. Functional Ecology 22:643-648.

48.   Carrier TJ, King BL, Coffman JA. 2015. Gene expression changes associated with

the developmental plasticity of sea urchin larvae in response to food availability.

Biological Bulletin 228:171-180.

49.   Hart MW, Strathmann RR. 1994. Functional consequences of phenotypic plasticity

in echinoid larvae. Biological Bulletin 186:291-299.

50.   McAlister JS, Miner BG. 2018. Phenotypic plasticity of feeding structures in marine

invertebrate larvae. *In* Carrier TJ, Reitzel AM, Heyland A (ed), Evolutionary Ecology

of Marine Invertebrate Larvae. Oxford University Press, Oxford, UK.

51.   Miner BG. 2004. Evolution of feeding structure plasticity in marine invertebrate

larvae: a possible trade-off between arm length and stomach size. Journal of

Experimental Marine Biology and Ecology 315:117-125.

52.   Miner BG. 2011. Mechanisms underlying feeding structure plasticity in echinoderm

larvae, p 221-229. *In* Flatt T, Heyland A (ed), Mechanisms of Life History Evolution.

Oxford University Press.

53.   Soars NA, Prowse TAA, Byrne M. 2009. Overview of phenotypic plasticity in

echinoid larvae, '*Echinopluteus transversus*' type vs. typical echinoplutei. Marine

Ecology Progress Series 383:113-125.

54.   Carrier TJ, Reitzel AM. Accepted. Bacterial community dynamics during embryonic

and larval development of three confamilial echinoids. Marine Ecology Progress

Series.

55.   Addison JA, Hart MW. 2004. Analysis of population genetic structure of the green

sea urchin (*Strongylocentrotus droebachiensis*) using microsatellites. Marine

Biology 144:243-251.

56.   Addison JA, Hart MW. 2005. Colonization, dispersal, and hybridization influence

phylogeography of North Atlantic sea urchins (*Strongylocentrotus droebachiensis*).

Evolution 59:532-543.

57.   Scheibling RE, Hatcher BG. 2013. *Strongylocentrotus droebachiensis*. *In* Lawrence

JM (ed), Sea Urchins: Biology and Ecology. Acedemic Press, San Diego, CA.

58.   Marks JA, Biermann CH, Eanes WF, Kryvi H. 2008. Sperm polymorphism within

the sea urchin *Strongylocentrotus droebachiensis*: divergence between Pacific and

Atlantic Oceans. Biological Bulletin 215:115-125.

59.   Manier MK, Palumbi SR. 2008. Intraspecific divergence in sperm morphology of the

green sea urchin, *Strongylocentrotus droebachiensis*: implications for selection in

broadcast spawners. BMC Evolutionary Biology 8:283.

60.   Bright M, Bulgheresi S. 2010. A complex journey: transmission of microbial

symbionts. Nature Reviews Microbiology 8:218-230.

61.   Brooks AW, Kohl KD, Brucker RM, van Opstal EJ, Bordenstein SR. 2016.

Phylosymbiosis: relationships and functional effects of microbial communities across

host evolutionary history. PLoS Biology 14:e2000225.

62.   Coyte KZ, Schluter J, Foster KR. 2015. The ecology of the microbiome: networks,

competition, and stability. Science 350:663-666.

63.   Gilbert SF, Bosch TCG, Ledon-Rettig C. 2015. Eco-Evo-Devo: developmental

symbiosis and developmental plasticity as evolutionary agents. Nature Reviews

Genetics 16:611-622.

64.   Ley RE, Hamady M, Lozupone C, Turnbaugh PJ, Ramey RR, Bircher JS, Schlegel

ML, Tucker TA, Schrenzel MD, Knight R, Gordon JI. 2008. Evolution of mammals

and their gut microbes. Science 320:1647-1651.

65.   Ley RE, Petersen DA, Gordon JI. 2006. Ecological and evolutionary forces shaping

microbial diversity in the human intestine. Cell 124:837-848.

66.   Miller ET, Svanbäck R, Bohannan BJM. 2018. Microbiomes as metacommunities:

understanding host-associated microbes through metacommunity ecology. Trends in

Ecology & Evolution 33:926-935.

67.   Rosenberg E, Sharon G, Zilber-Rosenberg I. 2009. The hologenome theory of

evolution contains Lamarckian aspects within a Darwinian framework.

Environmental Microbiology 11:2959-2962.

68.   Buckley KM, Rast JP. 2017. An organismal model for gene expression networks in

the gut-associated immune response. Frontiers in Immunology 8:1297.

69.   Kohl KD, Brun A, Magallanes M, Brinkerhoff J, Laspiur A, Acosta JC, Bordenstein

SR, Caviedes-Vidal E. 2016. Physiological and microbial adjustments to diet quality

permit facultative herbivory in an omnivorous lizard. Journal of Experimental

Biology 219:1903-1912.

70.   Alberdi A, Aizpurau O, Bohmann K, Zepeda-Mendoze ML, Gilbert MTP. 2016. Do

vertebrate gut metagenomes confer rapid ecological adaptation? Trends in Ecology

& Evolution 311:689-699.

71.   Carrier TJ, Macrander J, Reitzel AM. 2018. A microbial perspective on the life-

history evolution of marine invertebrate larvae: if, where, and when to feed. Marine

Ecology 39:e12490.

72.   Wolfe K, Graba-Landry A, Dworjanyn S, Byrne M. 2015. Larval phenotypic

plasticity in the boom-and-bust crown-of-thorns seastar, *Acanthaster planci*. Marine

Ecology Progress Series 539:179-189

73.   Wolfe K, Graba-Landry A, Dworjanyn S, Byrne M. 2015. Larval starvation to

satiation: influence of nutrient regime on the success of *Acanthaster planci*. PLoS

ONE:e0122010.

74.   Wolfe K, Graba-Landry A, Dworjanyn S, Byrne M. 2017. Superstars: Assessing

nutrient thresholds for enhanced larval success of *Acanthaster planci*, a review of the

evidence. Marine Pollution Bulletin 116:307-314.

75.   Schmidt VT, Smith KF, Melvin DW, Amaral-Zettler LA. 2015. Community

assembly of a euryhaline fish microbiome during salinity acclimation. Molecular

Ecology 24:2537-2550.

76.   Kohl KD, Oakeson KF, Orr TJ, Miller AW, Forbey JS, Phillips CD, Dale C, Weiss

RB, Dearing MD. 2018. Metagenomic sequencing provides insights into microbial

detoxification in the guts of small mammalian herbivores (*Neotoma* spp.). FEMS

Microbiology Ecology 94:fiy184.

77.   Muegge BD, Kuczynski J, Knights D, Clemente JC, Gonzalez A, Fontana L,

Henrissat B, Knight R, Gordon JI. 2011. Diet drives convergence in gut microbiome

functions across mammalian phylogeny and within humans. Science 332:970-974.

78.   Norderhaug KM, Anglès d'Auriac MB, Fagerli CW, Gundersen H, Christie H, Dahl

K, Hobæk A. 2016. Genetic diversity of the NE Atlantic sea urchin

Strongylocentrotus droebachiensis unveils chaotic genetic patchiness possibly linked

to local selective pressure. Marine Biology 163:36.

79.   Cowen RK, Sponaugle S. 2009. Larval dispersal and marine population connectivity.

Annual Review of Marine Science 1:443-466.

80.   Otto L, Zimmerman JTF, Furnes GK, Mork M, Saetre R, Becker G. 1990. Review of

693   the physical oceanography of the North Sea. Netherlands Journal of Sea Research

694   26:161-238.

695   81.   Sutherland DA, MacCready P. 2011. A model study of the Salish Sea estuarine

696   circulation. Journal of Physical Oceanography 41:1125-1143.

697   82.   Pettigrew NR, Churchill JH, Janzen CD, Mangum LJ, Signell RP, Thomas AC,

698   Townsend DW, Wallinga JP, Xue H. 2005. The kinematic and hydrographic structure

699   of the Gulf of Maine coastal current. Deep Sea Research Part II: Topical Studies in

700   Oceanography 52:2369-2391.

701   83.   McQuatters-Gollop A, Raitsos DE, Edwards M, Pradhan Y, Mee LD, Lavender SJ,

702   Attrill MJ. 2007. A long-term chlorophyll dataset reveals regime shift in North Sea

703   phytoplankton   biomass   unconnected   to   nutrient   levels.   Limnology   and

704   Oceanograhpy 52:635-648.

705   84.   Thomas AC, Townsend DW, Weatherbee R. 2003. Satellite-measured phytoplankton

706   variability in the Gulf of Maine. Continental Shelf Research 23:971-989.

707   85.   Winter DF, Banse K, Anderson GC. 1975. The dynamics of phytoplankton blooms

708   in puget sound a fjord in the Northwestern United States. Marine Biology 29:139-

709   176.

710   86.   Williams EA, Carrier TJ. 2018. An -omics perspective on marine invertebrate larvae,

711   p 288-304. *In* Carrier TJ, Reitzel AM, Heyland A (ed), Evolutionary Ecology of

712   Marine Invertebrate Larvae. Oxford University Press, Oxford, UK.

713   87.   Burgsdorf I, Erwin PM, López-Legentil S, Cerrano C, Haber M, Frenk S, Steindler

714   L. 2014. Biogeography rather than association with cyanobacteria structures

symbiotic microbial communities in the marine sponge *Petrosia ficiformis*. Frontiers

in Microbiology 5:529.

88. Pantos O, Bongaerts P, Dennis PG, Tyson GW, Hoegh-Guldberg O. 2015. Habitat-

specific environmental conditions primarily control the microbiomes of the coral

Seriatopora hystrix. The ISME Journal 9:1916-1927.

89. Rothschild D, Weissbrod O, Barkan E, Kurilshikov A, Korem T, Zeevi D, Costea PI,

Godneva A, Kalka IN, Bar N, Shilo S, Lador D, Vila AV, Zmora N, Pevsner-Fischer

M, Israeli D, Kosower N, Malka G, Wolf BC, Avnit-Sagi T, Lotan-Pompan M,

Weinberger A, Halpern Z, Carmi S, Fu J, Wijmenga C, Zhernakova A, Elinav E,

Segal E. 2018. Environment dominates over host genetics in shaping human gut

microbiota. Nature 555:210-215.

90. Sullam KE, Essinger SD, Lozupone CA, O'Connor MP, Rosen GL, Knight R,

Kilham SS, Russell JA. 2012. Environmental and ecological factors that shape the

gut bacterial communities of fish: a meta-analysis. Molecular Ecology 21:3363-3378.

91. Rivkin RB, Bosch I, Pearse JS, Lessard EJ. 1986. Bacterivory: a novel feeding mode

for asteroid larvae. Science 233:1311-1314.

92. Manahan DT. 1990. Adaptations by invertebrate arvae for nutrient acquisition from

seawater. Inegrative and Comparative Biology 30:147-160.

93. Manahan DT, Davis JP, Stephens GC. 1993. Bacteria-free sea urchin larvae: selective

uptake of neutral amino acids from seawater. Science 220:204-206.

94. Feehan CJ, Grauman-Boss BC, Strathmann RR, Dethier MN, Duggins DO. 2018.

Kelp detritus provides high-quality food for sea urchin larvae. Limnology and

Oceanograhpy 63:5299-5306.

95.   Mies M, Sumida MYG, Rädecker N, Voolstra CR. 2017. Marine invertebrate larvae

associated with *Symbiodinium*: a mutualism from the start? Frontiers in Ecology and

the Environment 5:56.

96.   Bosch I. 1992. Symbiosis between bacteria and oceanic clonal sea star larvae in the

western North Atlantic Ocean. Marine Biology 114:495-502.

97.   Fenaux L, Strathmann MF, Strathmann RR. 1994. Five tests of food-limited growth

of larvae in coastal waters by comparison of rates of development and form of

echinoplutei. Limnology and Oceanography 39:84-98.

98.   Olson RR, Olson MH. 1989. Food limitation of planktotrophic marine invertebrate

larvae: does it control recruitment success? Annual Review of Ecology and

Systematics 20:225-247.

99.   Pauley G, Boring L, Strathmann RR. 1985. Food limited growth and development of

larvae: experiments with natural sea water. Journal of Experimental Marine Biology

and Ecology 93:1-10.

100. Morgan SG. 1995. Life and death in the plankton: larval mortality and adaptation, p

279-321. *In* McEdwards L (ed), Ecology of Marine Invertebrate Larvae. CRC Press.

101. Rumrill SS. 1990. Natural mortality of marine invertebrate larvae. Ophelia 32:163-

198.

102. Young CM, Chia F-S. 1987. Abundance and distribution of pelagic larvae as

influenced by predation, behavior, and hydrographic factors. *In* Giese AC, Pearse JS,

Pearse VB (ed), Reproduction of Marine Invertebrates IX General aspects: seeking

unity in diveristy. Blackwell Scienfic Publications, Palo Alto, California.

103. Douglas AE. 1998. Nutritional interactions in insect-microbial symbioses: aphids and

their symbiotic bacteria *Buchnera*. Annual Review of Entomology 43:17-37.

104. Kohl KD, Weiss RB, Cox J, Dale C, Dearing MD. 2014. Gut microbes of mammalian
herbivores facilitate intake of plant toxins. Ecology Letters 17:1238-1246.

105. Macke E, Callens M, Meester LD, Decaestecker E. 2017. Host-genotype dependent
gut microbiota drives zooplankton tolerance to toxic cyanobacteria. Nature
Communications 8:1608.

106. Strathmann MF. 1987. Reproduction and development of marine invertebrates of the
northern Pacific coast: data and methods for the study of eggs, embryos, and larvae.
University of Washington Press.

107. Guillard RRL (ed). 1975. Culture of phytoplankton for feeding marine invertebrates.
Plenum Press, New York, USA.

108. Schneider C, Rasband W, Eliceiri K. 2012. NIH Image to ImageJ: 25 years of image
analysis. Nature Methods 9:671-675.

109. Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, Glockner FO.
2013. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and
next-generation sequencing-based diversity studies. Nucleic Acids Research 41:e1.

110. Zhang J, Kobert K, Flouri T, Stamatakis A. 2014. PEAR: a fast and accurate Illumina
Paired-End reAd mergeR. Bioinformatics 30:614-620.

111. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: A flexible trimmer for Illumina
sequence data. Bioinformatics 30:2114-2120.

112. Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. 2011. UCHIME improves
sensitivity and speed of chimera detection. Bioinformatics 27:2194-2200.

113. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK,
Fierer N, Pena AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D,
Koenig JE, Ley RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J,

Sevinsky JR, Turnbaugh PJ, Walters WA, Widmann J, Yatsunenko T, Zaneveld J,

Knight R. 2010. QIIME allows analysis of high-throughput community sequencing

data. Nature Methods 7:335-336.

114. Lozupone C, Knight R. 2005. UniFrac: a new phylogenetic method for comparing

microbial communities. Applied and Environmental Microbiology 71:8228-8235.

115. Langille MGI, Zaneveld J, Caporaso JG, McDonald D, Knights D, Reyes JA,

Clemente JC, Burkepile DE, Vega Thurber RL, Knight R, Beiko RG, Huttenhower

C. 2013. Predictive functional profiling of microbial communities using 16S rRNA

marker gene sequences. Nature Biotechnology 31:814-821.

116. Parks DH, Tyson GW, Hugenholtz P, Beiko RG. 2014. STAMP: statistical analysis

of taxonomic and functional profiles. Bioinformatics 30:3123-3124.

117. Comeau AM, Douglas GM, Langille MGI. 2017. Microbiome helper: a custom and

streamlined workflow for microbiome research. mSystems 2:e00127-16.

A

Proportional abundance

Salish Sea | Gulf of Maine | North Sea | Average | Shared

- Bacteroidetes: Saprospirae
- Bacteroidetes: Flavobacteriia
- Proteobacteria: Alphaproteobacteria
- Proteobacteria: Betaproteobacteria
- Proteobacteria: Deltaproteobacteria
- Proteobacteria: Gammaproteobacteria
- Verrucomicrobia: Verrucomicrobiae
- Other

B

Salish Sea   Gulf of Maine

22.5%   6.8%   26.3%

9.7%

3.4%   10.9%

20.4%

North Sea

Figure S1. *Strongylocentrotus droebachiensis* larvae alter phenotype to feeding environment. Ratio between the post-oral arm and mid body line (mean ± standard error; n=20) for *S. droebachiensis* from the Salish Sea (A), Gulf of Maine (B), and North Sea (C) larvae having been fed either 10,000 (black), 1,000 (dark grey), 100 (grey), and 0 cells•mL^{-1} (white).

Figure S2. *Strongylocentrotus droebachiensis* larvae alter gut volume to feeding environment. Gut volume (mean ± standard error; n=20) for larval *S. droebachiensis* from the Salish Sea (A), Gulf of Maine (B), and North Sea (C) having been fed either 10,000 (black), 1,000 (dark grey), 100 (grey), and 0 cells•mL^{-1} (white).

Figure S3. Alpha rarefaction curves for *Strongylocentrotus droebachiensis* larvae from three geographic locations and of the seawater. Alpha rarefaction curves for the associated microbiota for *S. droebachiensis* larvae (circles) from the Salish Sea (blue), Gulf of Maine (red), and North Sea (yellow) and seawater (squares) based on the rarefaction depth (3,193 sequences) used for all analyses.

Figure S4. Dietary and temporal shifts in the bacterial communities associated with *Strongylocentrotus droebachiensis* larvae. Community similarity based on *S. droebachiensis* larval-associated bacterial communities based on food availability (A-C) over a multi-week exposure (D-F) for populations from the Salish Sea (A, D), Gulf of Maine (B, E), and North Sea (C, F). Comparisons between food availability (A-C) and over time (D-F) are based on unweighted UniFrac comparisons. (Note: this is the unweighted complement to Fig. 2.)

Figure S5. Community similarity of *Strongylocentrotus droebachiensis* larvae from the Salish Sea across time. Similarity of bacterial communities associated with *S. droebachiensis* larvae from the Salish Sea for one (A; p < 0.002), two (B; p < 0.001), three (C; p < 0.001), or four (D; p < 0.001) weeks and having been fed 10,000 (orange), 1,000 (blue), 100 (purple), or 0 (yellow) cells•mL^{-1}.

42
43
44 **Figure S6.** Community similarity of *Strongylocentrotus droebachiensis* larvae from the Gulf of
45 Maine across time. Similarity of bacterial communities associated with *S. droebachiensis* larvae
46 from the Gulf of Maine for one (A; p < 0.001), two (B; p < 0.005), three (C; p < 0.007), or four
47 (D; p = 0.430) weeks and having been fed 10,000 (orange), 1,000 (blue), 100 (purple), or 0 (yellow)
48 cells•mL^{-1}.
49
50

51
52
53 **Figure S7.** Community similarity of *Strongylocentrotus droebachiensis* larvae from the North Sea
54 across time. Similarity of bacterial communities associated with *S. droebachiensis* larvae from the
55 North Sea for one (A; $p < 0.002$), two (B; $p < 0.001$), three (C; $p < 0.001$), or four (D; $p < 0.020$)
56 weeks and having been fed 10,000 (orange), 1,000 (blue), 100 (purple), or 0 (yellow) cells•mL^{-1}.
57
58

59
60
61 **Figure S8.** Community similarity of *Strongylocentrotus droebachiensis* larvae from the Salish Sea
62 across diets. Similarity of bacterial communities associated with *S. droebachiensis* larvae from the
63 Salish Sea having been fed 10,000 (A; $p < 0.001$), 1,000 (B; $p < 0.001$), 100 (C; $p < 0.001$), or 0
64 (D; $p < 0.001$) cells•mL^{-1} for one (orange), two (blue), three (purple), or four (yellow) weeks.
65
66

67
68
69 **Figure S9.** Community similarity of *Strongylocentrotus droebachiensis* larvae from the Gulf of
70 Maine across diets. Similarity of bacterial communities associated with *S. droebachiensis* larvae
71 from the Gulf of Maine having been fed 10,000 (A; $p < 0.005$), 100 (B; $p < 0.001$), or 0 (C; $p <$
72 0.002) cells•mL^{-1} for one (orange), two (blue), three (purple), or four (yellow) weeks.
73
74

75
76
77 **Figure S10.** Community similarity of *Strongylocentrotus droebachiensis* larvae from the North
78 Sea across diets. Similarity of bacterial communities associated with *S. droebachiensis* larvae from
79 the North Sea having been fed 10,000 (A; p < 0.002), 1,000 (B; p < 0.002), 100 (C; p < 0.002), or
80 0 (D; p < 0.006) cells•mL^{-1} for one (orange), two (blue), three (purple), or four (yellow) weeks.
81
82

83
84
85 **Figure S11.** Community similarity of the associated microbial community along a phenotypic
86 continuum for *Strongylocentrotus droebachiensis* from three geographic locations. Similarity of
87 bacterial communities associated with *S. droebachiensis* larvae from the Salish Sea (A; p = 0.046),
88 Gulf of Maine (B; p < 0.005), and North Sea (C; p < 0.004) pre (orange) and post (blue and purple)
89 expression of plasticity.
90
91

94 **Figure S12.** Similarity of the associated microbial community for *Strongylocentrotus*
95 *droebachiensis* larvae from three geographic locations. Community similarity (p < 0.001) of the
96 associated microbiota between *S. droebachiensis* larvae (circles) from the Salish Sea (blue), Gulf
97 of Maine (red), and North Sea (yellow) and seawater (squares).

100
101
102 **Figure S13.** Community similarity of *Strongylocentrotus droebachiensis* larvae across diets.
103 Similarity of bacterial communities associated with *S. droebachiensis* larvae from the Salish Sea
104 (purple), Gulf of Maine (blue), or North Sea (orange) and having been fed 10,000 (A; $p < 0.001$),
105 1,000 (B; $p < 0.001$), 100 (C; $p < 0.001$), and 0 (D; $p < 0.001$) cells•mL^{-1}.
106
107

Figure S14. Community similarity of *Strongylocentrotus droebachiensis* larvae across time. Similarity of bacterial communities associated with *S. droebachiensis* larvae from the Salish Sea (purple), Gulf of Maine (blue), or North Sea (orange) and having been differentially fed for one (A; p < 0.001), two (B; p < 0.001), three (C; p < 0.001), or four (D; p < 0.001) weeks.

116
117
Figure S15. Similarity between the associated bacterial communities of *Strongylocentrotus*
droebachiensis larvae from three geographic locations. Dendrograms of *S. droebachiensis* larvae
representing the total community for the Salish Sea, Gulf of Maine, and North Sea, with variation
due to phenotypes, dietary options, and developmental stages being collapsed. Dendrograms were
generated as part of the "jackknifed_beta_diversity.py" command for Figure 3 using weighted
UniFrac.
124

125
126
127 **Figure S16.** Predicted functional profiles of bacterial communities associated with
128 *Strongylocentrotus droebachiensis* larvae between geographical locations. Bar charts of PICRUSt-
129 generated metagenomics profiles group at level 1 KEGG categories for the bacterial communities
130 associated with *S. droebachiensis* larvae from the Salish Sea (red), Gulf of Maine (blue), and North
131 Sea (orange).
132
133

Note S1

QIIME analysis

Starting with raw read files from MiSeq:

1) Pair forward and reverse files using PEAR (pear-0.9.10-bin-64)
2) Trim paired files using Trimmomatic (trimmomatic-0.36.jar)
3) Convert paired and trimmed .fastq to .fasta using the custom code: cat [input .fastq] | paste –
    - - - | cut -f 1,2 | sed 's/^/>/'| tr "\t" "\n" >[output .fasta]
4) Validate tab-delimited mapping file using "validate_mapping_file.py"
5) Generate meta-.fasta files using "add_qiime_labels.py"
6) Detect chimeras from meta-.fasta (called, "combined_seqs.fna")
7) Filter chimeras using "filter_fasta.py"
8) Pick OTUs using "pick_open_reference_otus.py"
9) Filter OTUs with >10 reads using "filter_otus_from_otu_table.py"
10) Filter 'o__Cryptophyta' using "filter_taxa_from_otu_table.py"
11) Determine rarefaction depth using "biom summarize-table"
12) Filtered .biom table was rarified using "single_rarefaction.py"
13) Filtered filtered .biom table was split using "split_otu_table.py" to test specific hypotheses
14) Alpha diversity was compared using "alpha_diversity.py"
15) Beta diversity was compared using "jackknifed_beta_diversity.py" and 2D PCoA plots of unweighted and weighted UniFrac values were generated using "make_2d_plots.py"
16) Statistical comparisons of unweighted and weighted UniFrac were performed with "compare_categories.py" using the '--method' "anosim," "adonis," and "permanova."
17) Taxonomic plots were generated using "summarize_taxa_through_plots.py"
18) Unique and shared OTUs were determined both using "compute_core_microbiome.py"and "shared_phylotypes.py"

PICRUSt analysis

Starting with the biom table following 'Step 12' from 'QIIME analysis':

1) OTUs picked *de novo* were filtered using "filter_otus_from_otu_table.py" from QIIME
2) The closed OTU table was normalized using "normalized normalize_by_copy_number.py"
3) Metagenomes were predicted using "predict_metagenomes.py"
4) Predicted functions were categorized using "categorize_by_function.py"
5) The output from 'Step 4' was made STAMP-compatible using biom_to_stamp.py from Microbiome Helper
6) Resultant principles coordinates were extracted and analyzed using JMP

Taxa relating to metabolism

Starting with the 'closed' and 'normalized' biom table following 'Step 3' from 'PICRUSt analysis':

1) Determine which taxa contributed to 'Metabolism' using "metagenome_contributions.py" from PICRUSt for all related KEEG ID:
    "K00010,K00020,K00030,K00040,K00051,K00052,K00053,K00061,K00062,K00071,K00072,K00073,K00100,K00120,K00121,K00130,K00140,K00190,K00195,K00196,K00220,K00230,K00231,K00232,K00240,K00250,K00253,K00254,K00260,K00261,K00270,K00280,K00281,K00290,K00300,K00310,K00311,K00330,K00331,K00332,K00333,K00340,

K00350,K00360,K00361,K00362,K00363,K00364,K00365,K00380,K00400,K00401,K00402,K00403,K00404,K00405,K00410,K00430,K00440,K00450,K00460,K00471,K00472,K00473,K00480,K00500,K00510,K00511,K00512,K00513,K00514,K00515,K00520,K00521,K00522,K00523,K00524,K00525,K00531,K00532,K00533,K00534,K00540,K00550,K00561,K00562,K00563,K00564,K00565,K00571,K00572,K00590,K00591,K00592,K00600,K00601,K00603,K00604,K00620,K00621,K00622,K00623,K00624,K00625,K00626,K00627,K00630,K00633,K00640,K00642,K00643,K00650,K00660,K00670,K00680,K00710,K00720,K00730,K00740,K00750,K00760,K00770,K00780,K00785,K00790,K00791,K00830,K00860,K00900,K00901,K00902,K00903,K00904,K00905,K00906,K00908,K00909,K00910,K00920,K00930,K00940,K00941,K00942,K00943,K00944,K00945,K00950,K00960,K00965,K00966,K00980,K00981,K00982,K00983,K00984,K00999,K01010,K01040,K01051,K01052,K01053,K01054,K01055,K01056,K01057,K01058,K01059,K01060,K01061,K01062,K01063,K01064,K01065,K01066,K01070,K01100,K01110,K01120,K01130,K01200,K01210,K01212,K01220,K01230"

2) Determine class-level composition using custom code: awk '{print $c__*}' [output txt file from '1' of this section]