Development of an environmental DNA method for monitoring fish communities: ground truthing in diverse lakes with characterised fish faunas

Running title: eDNA method for lake fish monitoring

Jianlong Li[1, 5]*, Tristan W. Hatton-Ellis[2], Lori-Jayne Lawson Handley[1], Helen S. Kimbell[1, 3], Marco Benucci[1], Graeme Peirson[4], Bernd Hänfling[1]

1 Evolutionary and Environmental Genomics Group (@EvoHull), School of Environmental Sciences, University of Hull, Hull, UK

2 Natural Resources Wales, Bangor, UK

3 Frontiers, Lausanne, Switzerland

4 Environment Agency, Kidderminster, UK

5 Institute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Sciences, Haikou, China

*Corresponding author:

haikuilee@gmail.com (JL)

Abstract

1. Accurate, cost-effective monitoring of fish is required to assess the quality of lakes under the European Water Framework Directive (WFD). Recent studies have shown that environmental DNA (eDNA) metabarcoding is an effective and non-invasive method, which can provide semi-quantitative information on fish communities in large lakes.

2. This study further investigated the potential of eDNA metabarcoding as a tool for WFD status assessment by collecting and analysing water samples from eight Welsh lakes and six meres in Cheshire, England, with well described fish faunas. Water samples ($N = 252$) were assayed using two mitochondrial DNA regions (Cytb and 12S rRNA).

3. eDNA sampling indicated the presence of very similar species in the lakes compared to those expected on the basis of existing and historical information. In total, 24 species were detected with a total of 111 species occurrences in the lakes studied using eDNA. Secondly, there was a significant positive correlation between expected faunas and eDNA data in terms of confidence of species occurrence (Spearman's $r = 0.74$, $df = 109$, $p < 0.001$). Thirdly, eDNA data can estimate relative abundance with the standard five-level classification scale ("DAFOR"). Lastly, four ecological fish communities were characterised using eDNA data which agrees with the pre-defined lake types according to environmental characteristics.

4. *Synthesis and applications*. This study provides further evidence that eDNA metabarcoding could be a powerful and non-invasive monitoring tool for WFD purpose in a wide range of lake types, considerably outperforming other methods for community level analysis.

Key-words: eDNA, metabarcoding, fish monitoring, community ecology, EC Water Framework Directive, lakes, lentic systems

Introduction

The impact of anthropogenic pressures on aquatic ecosystems is ubiquitous and usually leads to alterations of the environment (Scheffer et al. 2001). To mitigate negative human effects, the first step in the improvement of the ecological quality of degraded water bodies is the development of an assessment system to evaluate the current situation. The main European initiative for water quality assessment and improvements, the Water Framework Directive 2000/60/EC (WFD), requires all member states to reach "good" ecological status of lakes, rivers and ground waters based on biological elements including phytoplankton, macrophytes and phytobenthos, benthic invertebrates and fish (CEC 2000).

Fish are widely considered as relevant for detecting and quantifying impacts of anthropogenic pressures on lakes and reservoirs (Argillier et al. 2013). Although most UK WFD ecological tools have been developed and deployed, an effective fish-based assessment tool for lakes remains problematic (Winfield 2002; Kelly et al. 2012). The current fish-based indices (e.g. Argillier et al. 2013) or tools (e.g. "FIL2" in Kelly et al. 2012) rely on semi-quantitative capture-based methods such as electro-fishing or gill-netting to provide information on fish composition and abundance, as well as age-structure of fish populations. Furthermore, the choice of survey methods is heavily dependent on the depth and size of lake. However, both of these sampling methods are relatively laborious and thus expensive, sometimes destructive, taxonomically biased, cannot be deployed in all situations (e.g. in or near dense vegetation, or entire water column in a very deep lake) and have poor sampling accuracy and precision. These drawbacks restrict their ability to meet WFD requirements (Kubecka et al. 2009; Winfield et al. 2009). A future strategy for WFD purposes thus requires a highly cost-effective approach, with a minimum amount of destructive sampling.

A significant "game-changer" in biodiversity monitoring in recent years is the analysis of environmental DNA (eDNA) which is a non-invasive genetic method that takes advantage of

intracellular or extra-organismal DNA in the environment to detect the presence of organisms

(Lawson Handley 2015; Thomsen & Willerslev 2015; Taberlet et al. 2018). A particularly

promising approach is to simultaneously screen whole communities of organisms using

eDNA metabarcoding. Several studies have shown that eDNA metabarcoding using High-

Throughput Sequencing offers tremendous potential as a complementary method tool to

established monitoring methods for ecology and conservation of aquatic species (e.g.

Hänfling et al. 2016; Port et al. 2016; Valentini et al. 2016). After reviewing and discussing

the potential of eDNA metabarcoding as a tool for WFD status assessment, Hering et al.

(2018) suggested that this approach is well-suited for fish biodiversity assessment

considering representativeness, sensitivity, precision, comparability, cost-effectiveness, and

environmental impact. The mitochondrial COI gene is widely, but not unanimously accepted

as the standard metabarcode for metazoans (e.g. Riaz et al. 2011; Clarke et al. 2014).

However, with environmental samples, most primers targeting the COI region amplify large

proportions of microbial species (e.g. Stat et al. 2017). This fact remains the strongest reason

for the use of other genes such as mitochondrial cytochrome b (Cytb) and 12S rRNA (12S)

regions. A prototype eDNA tool assayed by Cytb and 12S regions for fish biodiversity

assessment was tested in three lakes in the English Lake District, and initial results from this

are promising (Hänfling et al. 2016). However, in order to understand factors such as

responses to ecological pressures, taxon specific biases, sampling requirements in different

lake types and management of low confident occurrence, a much larger dataset from a range

of lakes with different ecological characteristics is required. Thus, the objectives of this study

were (a) to broaden the lake fish dataset using eDNA-based metabarcoding analysis by

collecting and analysing water samples from 14 UK lakes with well described fish faunas; (b)

to explore a possible approach to evaluate the confidence of species presence based on site

occupancy and read counts and (c) to use a relative abundance scale to estimate species

abundance by comparing the eDNA metabarcoding results to historical data.

Materials and methods

Study sites

The distribution of the 14 lakes including eight Welsh lakes and six Cheshire meres are

shown in Figure 1, and the general characteristics are outlined in Appendix S1 Table S1.1

based on data from the UK Lakes Portal (Hughes et al. 2004). In brief, the 14 lakes can be

divided into three types according to environmental characteristics including Type 1: three

low alkalinity lakes that are broadly upland in character ("CWE" Llyn Cwellyn, "PAD" Llyn

Padarn and "OGW" Llyn Ogwen); Type 2: two high alkalinity but shallow lakes ("PEN"

Llyn Penrhyn and "TRA" Llyn Traffwll) on the west coast of Anglesey (North Wales) which

are close to sea and accessible for migratory fish, and Type 3: nine high alkalinity but

shallow lakes that are dominated by coarse fish ("KEN" Kenfig Pool, "LLB" Llan Bwch-llyn,

"LLG" Llangorse Lake, "MAP" Maer Pool, "CAM" Chapel Mere, "OSS" Oss Mere, "FEN"

Fenemere, "WLF" Watch Lane Flash and "BET" Betley Mere) (Appendix S2).

Figure 1 Distribution of sampling lakes. Sampling lake codes: (CWE) Llyn Cwellyn; (PAD) Llyn Padarn; (OGW) Llyn Ogwen; (PEN) Llyn Penrhyn; (TRA) Llyn Traffwll; (KEN) Kenfig Pool; (LLB) Llan Bwch-llyn; (LLG) Llangorse Lake; (MAP) Maer Pool; (CAM) Chapel Mere; (OSS) Oss Mere; (FEN) Fenemere; (WLF) Watch Lane Flash; (BET) Betley Mere. The GPS coordinates of sampling locations in each lake are listed in Appendices S3 & S4.

eDNA capture, extraction, library preparation and sequencing

eDNA capture and extraction

In total, 252 samples were collected from the 14 lakes between 01/12/2015 to 12/07/2016 (Appendix S1 Table S1.2). The detail of sampling strategy can be found in Appendix S5.1. All samples were filtered within 24 hrs of collection. Two litres of each water sample were filtered through a 47 mm diameter 0.45-μm mixed cellulose acetate and nitrate filter (Whatman, UK) using Nalgene filtration units in combination with a vacuum pump (15–20 in. Hg; Pall Corporation, USA). Our previous study demonstrated that the 0.45-μm filters are suitable for fish metabarcoding, with low variation and high repeatability between the filtration replicates (Li et al. 2018a) compared to other filtration methods. The filtration units were cleaned with 10% v/v commercial bleach solution (containing ~3% sodium hypochlorite) and 5% v/v microsol detergent (Anachem, UK), and then rinsed thoroughly with de-ionised water after each filtration run to prevent cross-contamination. For each sampling lake, a filtration blank (2 L deionised water) was filtered before sample filtration in order to test for possible contamination at the filtration stage. After filtration, all filters were placed into 50 mm sterile petri dishes using sterile tweezers, sealed with parafilm and then immediately stored at –20°C until extraction. DNA extraction was carried out using the DNeasy PowerWater Kit (QIAGEN, Germany) following the manufacturer's protocol. The DNA was eluted in 100 μL 10 mM Tris (Solution PW6) and stored at –20 °C freezer.

Library preparation and sequencing

Sequencing libraries were generated from PCR amplicons targeting Cytb and 12S. We previously tested both fragments *in vitro* on 22 UK freshwater fish species and *in situ* on

three deep lakes in the English Lake District, and demonstrated their suitability for eDNA

metabarcoding of UK lake fish communities (Hänfling et al. 2016).

To enable the detection of possible PCR contamination, we included no-template controls

(NTCs, i.e. negative control) of molecular grade water (Fisher Scientific, UK) and single-

template controls (STCs, i.e. positive control) of cichlid fish DNA (the Eastern happy,

Astatotilapia calliptera, a cichlid from Lake Malawi, which is not present in natural waters in

the UK) within each library. All lake samples (N = 252), together with 14 sampling blanks,

14 filtration blanks, 14 NTCs, and 14 STCs were included in each library construction (N =

308) for sequencing on an Illumina MiSeq. Three PCR technical replicates were performed

for each sample then pooled to minimise bias in individual PCRs. All PCRs were set up using

eight-strip PCR tubes in a PCR workstation with UV hood and HEPA filter in the eDNA

laboratory at University of Hull (UoH) to minimise the risk of contamination.

The Cytb locus, targeting a 414-bp vertebrate-specific fragment, was amplified with the

L14912 and H15149 primer pair (Kocher et al. 1989) using a one-step library preparation

protocol (Kozich et al. 2013) (see more detail in Appendix S5.2).. To improve clustering

during initial sequencing, the final 10 pM denatured Cytb library mixed with 30% PhiX

control was sequenced with the MiSeq reagent kit v3 (2×300 cycles) at UoH. The 12S locus

targets a 106-bp vertebrate-specific fragment amplified with the 12S_V5_F and 12S_V5_R

primer pair (Riaz et al. 2011). Following the consistently lower sequencing yield of the one-

step protocol for the 12S compared to the Cytb in previous studies, we decided to switch to a

two-step library preparation protocol using nested tagging (Kitson et al. 2018) (see more

detail in Appendix S5.2). Since the heterogeneity spacers included in the two-step library

preparation protocol will improve the nucleotide diversity, the denatured 12S library (13 pM)

was mixed with lower amount of PhiX control (i.e. 10%) compared to the Cytb library, then

sequenced with the MiSeq reagent kit v2 (2×250 cycles) at UoH.

Data analysis

Collation of fish data

Existing fish data for the 14 lakes was collated using a range of data sources (see the detail in Appendix S2). These included site-specific surveys carried out by Natural Resources Wales or the Environment Agency and their predecessor bodies, third party fishery surveys, and data published in the literature and *ad hoc* records stored on databases such as the National Biodiversity Network Atlas (https://nbnatlas.org). Based on existence of survey data and expert opinion (TWH-E and GP), fish species were placed into four general categories with confidence of species presence/absence for each lake (Table 1), reflecting both the available data and the uncertainty around it. Assessing relative abundance *a priori* was more challenging. For the Welsh lakes, an abundance category approach was used, with each species being placed on DAFOR scale (D = Dominant; A = Abundant; F = Frequent; O = Occasional; R = Rare) (Tansley 1993) using available data where possible and/or TWH-E's expert opinion. All records where no abundance was recorded were assumed to be rare. For Cheshire meres, summaries of fish community composition and fish density per unit area (ind. ha^{-1} and kg ha^{-1}) were produced from the Ecological Consultancy Ltd (ECON) fishery survey using point abundance sampling by electro-fishing (PASE) and seine netting in the same period as water samples collection with this study in 2016 (Appendix S1 Tables S1.3 & S1.4). The ECON fishery survey in 2016 was undertaken by ECON and commissioned by Natural England as part of an investigation of the impacts of fisheries on Sites of Special Scientific Interests (SSSIs) condition.

190   Table 1 Criteria for assessing the confidence of species presence/absence using existing data.

Category	Confidence in Presence	Description
Probably absent (Pa)	Very low	No records and at least one of outside biogeographic range/habitat unsuitable.
Possibly present (Po)	Low	Either not recorded since 1977 or not recorded from the lake, but present in connected water body or diadromous species with direct access to the sea.
Probably present (Pr)	Moderate	Either multiple records but not recorded since 1997 or only one record since 1997. For Welsh lakes, this category has also been used where the current status of a population is uncertain, even if records data meet the criteria for Established.
Established (E)	High	For Welsh lakes, multiple records including at least one since 2000. For Cheshire meres, records in the Ecological Consultancy Ltd (ECON) survey.

Bioinformatics analysis

Raw read data from Illumina MiSeq sequencing have been submitted to NCBI (BioProject:

PRJNA454866). Bioinformatics analysis was implemented following a custom reproducible

metabarcoding pipeline (metaBEAT v0.97.9) with custom-made UK freshwater fish

reference databases (Cytb and 12S) as described in our previous study (Hänfling et al. 2016).

Sequences which the best BLAST hit had a bit score below 80 or had less than 95%/100%

identity (Cytb/12S) identity to any sequence in the curated databases were considered non-

target sequences. The different level of similarity for each locus depended on the length of

metabarcode and the knowledge of intraspecific diversity of the studied taxon. To assure full

reproducibility of our bioinformatics analysis, the custom reference databases and the Jupyter

notebooks for data processing have been deposited in an additional dedicated GitHub

repository (https://github.com/HullUni-

bioinformatics/Li_et_al_2018_eDNA_fish_monitoring). The Jupyter notebook also performs

demultiplexing of the indexed barcodes added in the first PCR reactions.

Low-frequency noise threshold

Filtered data were summarised into the number of sequence reads per species/sample for

downstream analyses (Appendixes S3 & S4). After bioinformatics analysis, the low-

frequency noise threshold (proportion of STC species read counts in the lake sample) was set

to 0.07% and 0.3% for Cytb and 12S respectively to filter high-quality annotated reads

passing the previous filtering steps that have high-confidence BLAST matches but may result

from contamination during the library construction process or sequencing (De Barba et al.

2014; Hänfling et al. 2016; Port et al. 2016). The low-frequency noise threshold was guided

by the analysis of sequence data from STCs under several thresholds. Filtered data were

summarised in two ways for downstream analyses: (a) the number of sequence reads per

species at each lake (hereafter referred to as read counts) and (b) the proportion of sampling

locations in which a given species was detected (hereafter referred to as site occupancy).

Estimating abundance with eDNA

221   Based on our previous results (Hänfling et al. 2016), site occupancy is a better proxy for

222   estimates of abundance than read counts. However, read counts should not be completely

223   ignored because they still contain important abundance information. The maximum site

224   occupancy across both loci was used as a score for species abundance (Table 2). In addition

to site occupancy score, the maximum relative read count (i.e. proportion of read counts per species at each sampling lake) across both loci, and the number of loci with which the species was detected were used as confidence indicators to assign fish species into the same general categories which were used for non-eDNA survey data (Table 3). Those species that were assigned the lowest confidence scores (probably absent) were excluded from downstream analysis (i.e. roach *Rutilus rutilus* in CWE, bream *Abramis brama* and bullhead *Cottus gobio* in OGW, pike *Esox lucius* in PEN, bullhead in TRA, gudgeon *Gobio gobio* in BET). Each retained species was assigned a corrected abundance score by multiplying site occupancy score × confidence score. The corrected abundance score was used to assign each species to a relative abundance DAFOR scale (Table 4) (For instance, bream in LLB: $2 \times 5 = 10$ resulting in "Frequent" category in DAFOR scale).

Table 2 Site occupancy score based on maximum site occupancy across Cytb and 12S.

Site occupancy score	Maximum site occupancy (across Cytb and 12S)
1	> 0 and ≤ 0.1
2	> 0.1 and ≤ 0.3
3	> 0.3 and ≤ 0.6
4	> 0.6 and ≤ 0.8
5	> 0.8

Table 3 Criteria for assessing confidence of species occurrence using eDNA data.

Category	Confidence score	Description		
		Locus	Site occupancy score	Maximum relative read count (across Cytb and 12S)
Probably absent (PrA)	Very low (-1)	Any	= 1	> 0 and ≤ 0.005

Possibly present (PoP)	Low (1)	Any	> 1	> 0 and ≤ 0.005
		One	≥ 1	> 0.005 and ≤ 0.05
Probably present (PrP)	Moderate (3)	Both	≥ 1	> 0.005 and ≤ 0.01
		Both	= 1	> 0.01 and ≤ 0.1
		One	≥ 1	> 0.05 and ≤ 0.1
		One	= 1	> 0.1 and ≤ 0.5
Established (E)	High (5)	Both	> 1	> 0.01
		Both	= 1	> 0.1
		One	> 1	> 0.1
		One	= 1	> 0.5

241 Table 4 The relative abundance DAFOR scale based on eDNA data.

Abundance rank number	DAFOR scale	Corrected abundance score (site occupancy score × confidence score)
0	None	= 0
1	Rare	≥ 1 and < 4
2	Occasional	≥ 4 and < 9
3	Frequent	≥ 9 and < 15
4	Abundant	≥ 15 and < 25
5	Dominant	= 25

Ecological and statistical analyses

All downstream analyses were performed in R v3.3.2 (R_Core_Team 2016) and graphs

plotted using GGPLOT2 v2.2.1 (Wickham & Chang 2016). Before investigating species

detection and abundance estimate with eDNA, we first evaluated whether Cytb and 12S

datasets produced consistent results by calculating the Pearson's product-moment correlation

coefficient for both read counts and site occupancy. Spearman's rank correlations were

performed between historical data and eDNA data – firstly in terms of confidence of species

presence and secondly in terms of relative abundance. To investigate differences in fish

communities between sampling lakes, hierarchical clustering dendrogram was used to assess

the existence of distinct community types using the function *fviz_dend* in FACTOEXTRA

v1.0.4 (Kassambara & Mundt 2017) to extract and visualize the results calculated from the

function *hclust* using Canberra distances method based on site occupancy. Furthermore, non-

metric multidimensional scaling (NMDS) based on read counts in individual sampling

location of lakes, allied with analysis of similarities (ANOSIM), were performed using the

abundance-based Bray-Curtis dissimilarity index with the function *metaMDS* and *anosim*

respectively in VEGAN v2.4-4 (Oksanen et al. 2017). The sequencing depth and sample-

based rarefaction was performed using the *rarecurve* and *rarc* function in VEGAN v2.4-4

(Oksanen et al. 2017), respectively. The full R script is available on the GitHub repository

(https://github.com/HullUni-

bioinformatics/Li_et_al_2018_eDNA_fish_monitoring/tree/master/R_script)

Results

The Cytb and 12S libraries generated 17.15 and 15.97 million reads, respectively, with

15.91 and 13.4 million reads passing filters (including 49.01% or 11.07% PhiX). After

quality filtering and removal of chimeric sequences, the average read count of lake sample

was 12,503 for Cytb and 21,703 for 12S. After BLAST searches for taxonomic assignment,

$59.47\% \pm 35.56\%$ and $26.46\% \pm 19.60\%$ reads in each lake sample were assigned to fish for

Cytb and 12S, respectively. Rarefaction analysis revealed our sequencing depth to be

sufficient to identify all of the amplified fish species present in the 14 lakes for both loci

(Appendix S1 Figure S1.1).

Significant correlations were found between site occupancy and average read counts, for

both loci and all sampling lakes apart from OGW (Pearson's $r = 0.81$, $df = 4$, $p = 0.05$) and

LLB (Pearson's $r = 0.94$, $df = 2$, $p = 0.06$) for Cytb (Appendix S1 Figure S1.2a, b). Data from

the two loci were significantly correlated (Pearson's r consistently $p < 0.05$) for site

occupancy for each lake (Appendix S1 Figure S1.2c), and for average read counts apart from

in three lakes: PAD (Pearson's $r = 0.55$, $df = 7$, $p = 0.12$), TRA (Pearson's $r = 0.59$, $df = 5$, p

$= 0.16$) and LLB (Pearson's $r = 0.89$, $df = 2$, $p = 0.11$) (Appendix S1 Figure S1.2d). The more

detail of comparisons between these two loci in detection probabilities can be found in

Appendix S5.3.

Comparison of eDNA data and expected results

In total, 24 species with 111 species occurrences in 14 lakes were detected across Cytb

and 12S. Perch *Perca fluviatilis* was the most common species which were detected in all

lakes. The second most frequently occurring species was roach with detections in 10 lakes. In

addition to these two species, pike were found in all nine coarse fish lakes (Table 5; Figure 2;

Appendix S1 Figure S1.3). There was a significant positive correlation between expected

faunas and eDNA data in terms of confidence of species occurrence (Spearman's $r = 0.74$, df

$= 109$, $p < 0.001$). A total of 73 "established" species occurrences have been recorded across

all lakes according to historical data. Of these, 72 (98.6%) were detected with eDNA (Table

5). The only false negative in the eDNA datasets was tench *Tinca tinca* in PEN. Tench were

detected by 12S in PEN but at a read count less than the low-frequency noise threshold

(Appendix S4; site occupancy $= 0.3$, average read counts $= 3.9$). Tench are likely to be very

rare in PEN: only two tench were recorded by the 2016 Royal Society for the Protection of

Birds survey and small numbers have also been detected in previous years (Appendix S2.4).

Twenty-five of the 111 occurrences were "probably or possibly present", and 13 were "very

low confidence present" (i.e. probably absent records based on historical data) (Table 5).

There were consistent positive correlations between DAFOR scale based on eDNA data

and expected DAFOR scale based on both historical data for Welsh lakes and fish density per

unit area from the ECON survey of Cheshire meres (Figure 3). The correlations were

significant in three out of eight Welsh lakes (PAD, OGW and KEN). Significant correlations

were observed in three (MAP, CAM and BET) and four (MAP, CAM, FEN and WLF)

Cheshire meres based on individual density (ind. ha^{-1}) and biomass density (kg ha^{-1}),

respectively (Figure 3; Appendix S1 Figure S1.4).

Furthermore, the rigorous sampling of shore and offshore locations is carried out to

compare the suitability of the different spatial sampling location (Appendix S1 Table S1.2).

The more detail of comparisons between shore and offshore samples can be found in

Appendix S5.4.

Characterisation of fish assemblages using eDNA data

The hierarchical clustering dendrograms based on site occupancy for two loci indicated

that there were four distinct community types compared to three pre-defined lake types

according to environmental characteristics. Specifically, the clustering dendrograms accorded

with the pre-defined lake Type 1 (CWE, OGW and PAD) and Type 2 (TRA and PEN), but

indicated that WLF and BET can be divided from the pre-defined lake Type 3 (Figure 4a1,

b1). The NMDS ordination allied with ANOSIM based on read counts for two loci confirmed

the clustering dendrograms results that there were four distinct community types according to

the predominant groups of fish including Community 1: salmonids and minnow *Phoxinus*

phoxinus; Community 2: mixed diadromous fish; Community 3: coarse fish, and Community

4: bream-dominated coarse fish (Figure 4a2, b2). The *R* statistic in the ANOSIM global test

with these four different communities were high (Appendix S1 Table S1.5, 0.75 ± 0.02)

supporting statistical differences between communities (Appendix S1 Table S1.5, p = 0.001).

The overall distance pattern of these four fish communities was that both coarse fish

communities (Communities 3 & 4) were close to each other; the mixed diadromous fish

community (Community 2) was between the coarse fish communities and the salmonids and

minnow community (Community 1) (Figure 4; Appendix S1 Table S1.5).

The salmonids and minnow community (Community 1) is characteristic of three low

alkalinity upland lakes (CWE, OGW and PAD). These sites were generally dominated by

brown trout *Salmo trutta*, rainbow trout *Oncorhynchus mykiss* or Arctic charr *Salvelinus*

alpinus, but also included minnow and Atlantic salmon *Salmo salar* (Figure 2; Appendix S1

Figure S1.3). CWE and PAD are designated as SSSIs under the Wildlife and Countryside Act

1981 (as amended) to conserve Arctic charr, and CWE, a oligotrophic lake, is also protected

as a Special Areas of Conservation under the EU Habitats Directive. The eDNA data showed

that PAD contained a small number of Arctic charr, whereas this species was dominant in

CWE (Figure 2; Appendix S1 Figure S1.3). Rainbow trout were only detected in OGW,

where they are regularly stocked by the local angling club (Appendix S2.3). Some of these

sites may also contain low density of other species; usually diadromous species were detected

using eDNA as well such as European eel *Anguilla anguilla* in all three upland lakes and

three-spined stickleback *Gasterosteus aculeatus* in PAD. Perch have recently been recorded

from PAD for the first time using captured-based survey methods (Appendix S2.2).

Surprisingly, this species was detected using eDNA (though shown to be rare), in all of these

three lakes (Figure 2; Table 5). The diadromous fish community (Community 2) is

characteristic TRA and PEN which were dominated by European eel and three-spined

stickleback. Other species included brown trout, nine-spined stickleback *Pungitius pungitius*,

roach, perch and rudd *Scardinius erythrophthalmus* (Figure 2). The coarse fish communities

consisted of bream, common carp *Cyprinus carpio*, pike, perch, roach, rudd and tench, plus

some additional species (e.g. European eel, bullhead, three-spined stickleback and gudgeon).

Pike, perch, roach and tench were dominant in KEN, MAP, CAM, LLB, LLG, OSS and FEN

(Community 3); however, the dominant species in WLF and BET (Community 4) was bream

(Figure 2; Appendix S1 Figure S1.3).

Table 5 Correspondence of confidence of species occurrence between predicted and eDNA data in 14 lakes.

Community type			1		2		3							4		Number of lakes
Scientific name	Code	CWE	PAD	OGW	PEN	TRA	KEN	LLB	LLG	MAP	CAM	OSS	FEN	WLF	BET	
Abramis brama	BRE							PrP/E	E/E	PoP/PoP	PoP/PoP	E/E	E/E	E/E	E/E	8
Alburnus alburnus	BLE			PrA/PoP			PrA/PoP									2
Anguilla anguilla	EEL	E/PrP	E/E	PoP/PrP	E/E	E/E	E/E		E/E				E/E	PoP/PoP		9
Carassius auratus	GOF		PrA/PrP													1
Carassius carassius	CRU														Po/Pr	1
Cottus gobio	BUL						PrA/PoP		PoP/PrP						Po/Po	3
Cyprinus carpio	CAR									E/E	PoP/PoP	E/E	E/E	E/E	E/E	6
Esox lucius	PIK						E/E	E/E	E/E	E/E	E/E	E/E	E/E	E/E	E/E	9
Gasterosteus aculeatus	3SS		E/E		E/E	E/E					PoP/E		E/PrP	PoP/E	Po/Pr	7
Gobio gobio	GUD		PrA/PoP									PoP/PoP		E/PoP		3
Leucaspius deliniatus	SUN													E/E		1
Oncorhynchus mykiss	RTR			E/E												1
Perca fluviatilis	PER	PrA/E	PrP/PoP	PrA/PrP	E/E	PrA/E	E/E	E/E	E/E	E/E	E/E	E/E	E/E	E/E	E/E	14
Phoxinus phoxinus	MIN	E/E	E/E	E/E								PrA/PoP				4
Pseudorasbora parva	TMG														Pa/Po	1
Pungitius pungitius	9SS				PoP/PrP	PoP/PoP	PoP/PoP									3
Rhodeus amarus	BIT													PoP/PoP		1
Rutilus rutilus	ROA				E/E	E/E	PoP/PoP	E/E	E/E	PrA/PoP		E/E	E/E	E/E	E/E	10
Salmo salar	SAL	E/PoP	E/E													2
Salmo trutta	BTR	E/E	E/E	E/E	PoP/PoP	PoP/E									Po/Po	6
Salvelinus alpinus	CHA	E/E	E/PrP													2
Scardinius erythrophthalmus	RUD				E/E	PoP/PrP	E/E		E/E		PrA/PoP	E/E	E/E			7
Squalius cephalus	CHU													PrA/PoP		1
Tinca tinca	TEN				E/PrA		E/E		E/E	E/E	E/E	E/E	PoP/PoP	PoP/PrP	E/E	9

Number of species		6	9	6	8	7	9	4	8	6	7	9	9	12	11	111 in total

353 Notes: Categories of presence confidence ("PrA" probably absent, "PoP" possibly present, "PrP" probably present and "E" established) are described in

354 Table 1 showing as based on predicted/eDNA data. The "Established" species occurrences based on historical data are shown in blue background ($N =$

355 73). Solid vertical lines indicate community types (Community 1–3) with similar faunas. Dashed line indicates division between whether bream *Abramis*

356 *brama* is dominant species in the lake (Community 4). Sampling lake codes are given in Figure 1.

Figure 2 Species composition of site occupancy for Cytb and 12S. Species three letter codes and sampling lake codes are given in Table 5 and Figure 1, respectively.

Figure 3 Correlations between DAFOR scale based on eDNA data and expected DAFOR scale based on existing data of Welsh lakes or individual density from the ECON survey of Cheshire meres. Sampling lake codes, abundance rank numbers to DAFOR scale and species three letter codes are given in Figure 1, Table 4 and Table 5, respectively.

362
363
364
365

Figure 4 Hierarchical clustering dendrograms and non-metric multidimensional scaling (NMDS) ordination of the sampling lakes. Dendrograms using Canberra distances method based on site occupancy for (a1) Cytb and (b1) 12S. The dashed frames are drawn around each cluster in dendrograms. The three pre-defined lake types according to environmental characteristics. NMDS in individual sampling location of lakes using Bray-Curtis dissimilarity index based on read counts for (a2) Cytb and (b2) 12S. Each symbol corresponds to a sampling lake, with circles corresponding to shore samples ("SL") and triangles corresponding to offshore samples ("OSL") in NMDS ordination. The ellipse indicates 70% similarity level within each community type in ordinations. Scores for species taxa are plotted on the same axes to better visualise the ordination in space between species and samples. Species three letter codes and community type (Com 1–4) with individual lake codes are given in Table 5 and Figure 1, respectively.

Discussion

This study confirmed key results from our previous study in that there was (a) a consistent, strong correlation between Cytb and 12S in terms of read counts and site occupancy; (b) a consistent, strong correlation between site occupancy and read counts and (c) site occupancy was consistently better than read counts for estimating relative abundance, for both Cytb and 12S datasets (Hänfling et al. 2016).

Comparison of eDNA and existing data for species detection

A total of 73 "established" species occurrences were recorded across the 14 lakes according to existing and historical fish data. Of these, 72 (98.6%) were detected with eDNA, which demonstrated that eDNA metabarcoding outperformed the conventional surveys and observations using a range of sampling methods over date ranges spanning several decades in species richness estimates. This result is consistent with previous studies that have demonstrated that eDNA metabarcoding produced a more comprehensive species list than alternative survey techniques with a similar effort in both marine and freshwater ecosystems (Hänfling et al. 2016; Port et al. 2016; Valentini et al. 2016).

Moreover, there was a significant positive correlation between historical data and eDNA data in terms of confidence of species presence (Spearman's $r = 0.74$, $df = 109$, $p < 0.001$). Species occurrences that were assessed as "probably or possibly present" (25 of the 111 species occurrences) and "probably absent" (13 of the 111 species occurrences) based on distribution data or anecdotal evidence also fitted the eDNA criteria for lowered confidence levels. In most cases, the "probably absent" occurrences were at very low site occupancy ($\leq$ 0.1) and read counts (~0.5%), just above our threshold for accepting a positive record. These records could be genuine detections of species that have previously been missed. For example,

perch were recorded with both Cytb and 12S in CWE, OGW and TRA. The confidence of

species present was either "established present" or "probably present" according to eDNA

data. The recent appearance of this species in nearby lakes PAD and PEN (see detail in

Appendix S2) suggest that the species is spreading in North Wales either by natural means or

as a result of illegal introductions. In addition, the other "probably absent" occurrences could

be either false positives from sequencing error, laboratory or environmental contamination.

For instance, bleak *Alburnus alburnus* in OGW, goldfish *Carassius auratus* and Gudgeon in

PAD, rudd in CAM, roach in MAP, topmouth gudgeon *Pseudorasbora parva* in BET and

chub *Squalius cephalus* in WLF are most likely explained by low-level cross contamination

or sequencing barcode misassignment since these species were only detected by either Cytb

or 12S in single sample of the lake.

Barcode misassignment and tag jumps have been proved in metabarcoding studies (e.g.

Deakin et al. 2014; Schnell et al. 2015). To minimise contamination or false assignments,

further work needs to be done to optimise each step, as well as the incorporation of robust

controls to identify contamination when it occurs. Together with a growing number of studies

(Stat et al. 2017; Li et al. 2018b), the results demonstrate the importance of using two (or

more) markers for metabarcoding to avoid problems due to primer bias (Clarke et al. 2014),

gene copy number, PCR or sequencing artefacts (Schloss et al. 2011) and/or contamination.

Other strategies such as using consistency of presence across technical replicates as used by

Port et al. (2016) might be a more suitable approach to control for false positives if rare

species are of particular interest. Where verification of rare species detections is considered a

high priority, additional confirmation from targeted species approaches (e.g. quantitative

PCR) and/or field surveys may be required.

Use of eDNA for assessing relative abundance of fish

WFD specifies that abundance should be considered when determining ecological status;

hence, current WFD approaches include estimates of abundance (often as abundance classes).

For instance, a five-level scale, adapted from DAFOR scale, is accepted for Austrian standard

and national monitoring techniques applied under WFD for surveying aquatic macrophytes

(Pall & Moser 2009). Here, we used DAFOR scale to estimate fish relative abundance in the

present study based on DNA-based identification to facilitate integration into current WFD

approaches.

There were consistently positive correlations in terms of relative abundance between the

eDNA data and historical data. However, these correlations were not always statistically

significant. The correlations were not significant in TRA and PEN probably because three-

spined stickleback were more abundant based on eDNA data than expected. This species is

often under-represented or overlooked in surveys using established fish capture methods

(Hänfling et al. 2016). Moreover, tench were not detected by eDNA in PEN. LLB is a

species-poor lake with only four species detected, which could reduce the statistical power.

The non-significant correlation in WLF could be attributed to under-representation of three-

spined stickleback in previous fish surveys and reduced detection probability of gudgeon with

Cytb. Although these results are generally encouraging, further work is critical to obtain

enough statistical power to directly test the relationship between abundance estimates from

eDNA and surveys using other methods. It is also important to investigate taxon-specific

detection probabilities and abundance estimates so that a pressure-sensitive tool can be

developed.

Using eDNA to describe ecological communities

According to environmental characteristics, there were three pre-defined lake types.

Encouragingly, four distinct community types could be identified based on clustering

dendrograms and NMDS ordinations using eDNA data. Basically, the Community 1 and

Community 2 agreed with the pre-defined lake Type 1 and Type 2, respectively. The pre-

defined coarse fish lakes (Type 3) can be further divided into Community 3 and Community

4 based on whether bream was dominant species in the lake. These findings indicated that

eDNA metabarcoding has great potential as fish-based assessment tool for WFD lake status

assessment.

Specifically, our results showed that the low alkalinity lakes within a predominantly

upland catchment were mainly dominated by salmonids reflecting their relatively deep and

oligotrophic nature. Except from salmonids, all these sites contained minnow, most likely

introduced by anglers using them as live bait (Hatton-Ellis 2005). Compared to the salmonids

and minnow lakes (Community 1), perch, pike and cyprinids (such as bream, common carp,

roach and tench) were prevalent in the coarse fish lakes (Communities 3 & 4) reflecting their

relatively shallow and eutrophic nature. These findings support that eDNA profiles are

suitable to reflect the eutrophic conditions of Lake Windermere in which species that prefer

less eutrophic conditions (Atlantic salmon, brown trout, Arctic charr, minnow and bullhead)

were more abundant in the mesotrophic North Basin, and species associated with eutrophic

conditions (roach, tench, rudd, bream and European eel) were more common in the eutrophic

South Basin (Hänfling et al. 2016). Furthermore, both of the diadromous fish lakes

(Community 2) contained a different fauna including various combinations of mostly

diadromous species such as European eel, three-spined stickleback, nine-spined stickleback

and brown trout. Alongside these, several coarse fish species such as roach, perch, and rudd

also occurred in these lakes. However, of greater conservation interest is the distribution of

nine-spined stickleback. This species occurs only locally in Wales, mainly at lowland sites

close to the sea (Davies et al. 2004; Hatton-Ellis 2005).

Conclusions

In this study, we have extended the geographical, ecological and taxonomic extent of the

lake fish eDNA-based metabarcoding dataset, including 14 UK lakes with well described fish

faunas. Four eDNA communities are characterised, which is consistent with earlier

assessments and ecological interpretations. Moreover, eDNA metabarcoding outperformed

established survey techniques in terms of species detection, relative abundance using the

standard five-level classification scale and characterisation of ecological fish communities,

suggesting eDNA metabarcoding has great potential as fish-based assessment tool for WFD

lake status assessment. Further optimisation and development such as understanding taxon-

specific biases, abundance estimates and management of low confident occurrence are

strongly recommended with a larger dataset, to improve the accuracy, effectiveness and

applicability of WFD monitoring tool.

Authors' contributions

494 JL wrote the manuscript; BH, TWH-E, and GP conceived and designed the study; TWH-E

495 and GP prepared historical fish data; HSK, MB, and JL carried out the fieldwork; MB

496 prepared the sampling map; JL conducted laboratory work and performed bioinformatics

497 analyses; JL, LLH, and BH performed the statistical analyses. All authors commented the

498 final manuscript.

Acknowledgments

This work was part of PhD project of JL, who was supported by the University of Hull and China Scholarship Council. This work was funded by the UK Environment Agency and Natural Resources Wales. We are particularly grateful to all the site owners and managers for granting access and the Bangor University for providing access to their laboratory facilities. Ecological Consultancy Ltd provided fishery survey data and cooperation in data collation of Cheshire meres. Robert Jaques, Hayley Watson, Mags Cousins, Peter Clabburn, Rob Evans, Sophie Gott, Emma Keenan, Ian Sims, Emma Brown, Jason Jones, Dawn Parry, Peter Shum, Harriet Johnson, Rob Donnelly and Christoph Hahn provided invaluable help during fieldwork and lab work.

Data accessibility

Data is available from the GitHub repository (https://github.com/HullUni-bioinformatics/ Li_et_al_2018_eDNA_fish_monitoring). The repository is archived with Zenodo (https://doi.org/10.5281/zenodo.1462898).

References

Argillier, C., Causse, S., Gevrey, M., Pedron, S., De Bortoli, J., Brucet, S., ... Holmgren, K. (2013) Development of a fish-based index to assess the eutrophication status of European lakes. *Hydrobiologia,* **704,** 193-211.

CEC (2000) Directive 2000/60/EC of the European Parliament and of the Council: establishing a framework for Community action in the field of water policy. Official Journal of the European Communities, Luxembourg.

Clarke, L.J., Soubrier, J., Weyrich, L.S. & Cooper, A. (2014) Environmental metabarcodes for insects: *in silico* PCR reveals potential for taxonomic bias. *Molecular Ecology Resources,* **14,** 1160-1170.

Davies, C.E., Shelley, J., Harding, P.T., McLean, I.F.G., Gardiner, R. & Peirson, G. (2004) Freshwater Fishes in Britain - the species and their distribution. pp. 176. Harley Books, Colchester.

De Barba, M., Miquel, C., Boyer, F., Mercier, C., Rioux, D., Coissac, E. & Taberlet, P. (2014) DNA metabarcoding multiplexing and validation of data accuracy for diet assessment: application to omnivorous diet. *Molecular Ecology Resources,* **14,** 306-323.

Deakin, C.T., Deakin, J.J., Ginn, S.L., Young, P., Humphreys, D., Suter, C.M., ... Hallwirth, C.V. (2014) Impact of next-generation sequencing error on analysis of barcoded plasmid libraries of known complexity and sequence. *Nucleic Acids Research,* **42,** e129.

Hänfling, B., Lawson Handley, L., Read, D.S., Hahn, C., Li, J., Nichols, P., ... Winfield, I.J. (2016) Environmental DNA metabarcoding of lake fish communities reflects long-term data from established survey methods. *Molecular Ecology,* **25,** 3101-3119.

Hatton-Ellis, T. (2005) Fish communities and fisheries in Wales's National Nature Reserves: a review. *Freshwater Forum,* **24,** 82-104.

Hering, D., Borja, A., Jones, J.I., Pont, D., Boets, P., Bouchez, A., ... Kelly, M. (2018) Implementation options for DNA-based identification into ecological status assessment under the European Water Framework Directive. *Water Research,* **138,** 192-205.

Hughes, M., Hornby, D.D., Bennion, H., Kernan, M., Hilton, J., Phillips, G. & Thomas, R. (2004) The development of a GIS-based inventory of standing waters in Great Britain together with a risk-based prioritisation protocol. *Water, Air and Soil Pollution: Focus,* **4,** 73-84.

Kassambara, A. & Mundt, F. (2017) factoextra: extract and visualize the results of multivariate data analyses: an R package in CRAN. Available from https://CRAN.R-project.org/package=factoextra.

Kelly, F.L., Harrison, A.J., Allen, M., Connor, L. & Rosell, R. (2012) Development and application of an ecological classification tool for fish in lakes in Ireland. *Ecological Indicators,* **18,** 608-619.

Kitson, J.J.N., Hahn, C., Sands, R.J., Straw, N.A., Evans, D.M. & Lunt, D.H. (2018) Detecting host-parasitoid interactions in an invasive Lepidopteran using nested tagging DNA-metabarcoding. *Molecular Ecology***,** https:/doi.org/10.1111/mec.14518.

Kocher, T.D., Thomas, W.K., Meyer, A., Edwards, S.V., Pääbo, S., Villablanca, F.X. & Wilson, A.C. (1989) Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers. *Proceedings of the National Academy of Sciences,* **86,** 6196-6200.

Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K. & Schloss, P.D. (2013) Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Applied and Environmental Microbiology,* **79,** 5112-5120.

Kubecka, J., Hohausova, E., Matena, J., Peterka, J., Amarasinghe, U.S., Bonar, S.A., ... Winfield, I.J. (2009) The true picture of a lake or reservoir fish stock: a review of needs and progress. *Fisheries Research,* **96,** 1-5.

Lawson Handley, L. (2015) How will the 'molecular revolution' contribute to biological recording? *Biological Journal of the Linnean Society,* **115,** 750-766.

Li, J., Lawson Handley, L.J., Read, D.S. & Hänfling, B. (2018a) The effect of filtration method on the efficiency of environmental DNA capture and quantification via metabarcoding. *Molecular Ecology Resources,* **18,** 1102-1114.

Li, Y., Evans, N.T., Renshaw, M.A., Jerde, C.L., Olds, B.P., Shogren, A.J., ... Pfrender, M.E. (2018b) Estimating fish alpha- and beta-diversity along a small stream with environmental DNA metabarcoding. *Metabarcoding and Metagenomics,* **2,** e24262.

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., ... Wagner, H. (2017) Vegan: community ecology package. Retrieved from https://CRAN.R-project.org/package=vegan.

Pall, K. & Moser, V. (2009) Austrian Index Macrophytes (AIM-Module 1) for lakes: a Water
Framework Directive compliant assessment system for lakes using aquatic macrophytes.
Hydrobiologia, **633,** 83-104.
Port, J.A., O'Donnell, J.L., Romero-Maraccini, O.C., Leary, P.R., Litvin, S.Y., Nickols,
K.J., ... Kelly, R.P. (2016) Assessing vertebrate biodiversity in a kelp forest ecosystem
using environmental DNA. *Molecular Ecology,* **25,** 527-541.
R_Core_Team (2016) R: a language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. Retrieved from http://www.R-project.org/.
Riaz, T., Shehzad, W., Viari, A., Pompanon, F., Taberlet, P. & Coissac, E. (2011) ecoPrimers:
inference of new DNA barcode markers from whole genome sequence analysis. *Nucleic*
Acids Research, **39,** e145.
Scheffer, M., Carpenter, S., Foley, J.A., Folke, C. & Walker, B. (2001) Catastrophic shifts in
ecosystems. *Nature,* **413,** 591-596.
Schloss, P.D., Gevers, D. & Westcott, S.L. (2011) Reducing the effects of PCR amplification
and sequencing artifacts on 16S rRNA-based studies. *PloS ONE,* **6,** e27310.
Schnell, I.B., Bohmann, K. & Gilbert, M.T.P. (2015) Tag jumps illuminated – reducing
sequence‐to‐sample misidentifications in metabarcoding studies. *Molecular Ecology*
Resources, **15,** 1289-1303.
Stat, M., Huggett, M.J., Bernasconi, R., DiBattista, J.D., Berry, T.E., Newman, S.J., ... Bunce,
M. (2017) Ecosystem biomonitoring with eDNA: metabarcoding across the tree of life in
a tropical marine environment. *Scientific Reports,* **7,** 12240.
Taberlet, P., Bonin, A., Zinger, L. & Coissac, E. (2018) Environmental DNA: for biodiversity
research and monitoring. pp. 253. Oxford University Press, Oxford.
Tansley, A.G. (1993) An introduction to plant ecology. pp. 228. Discovery Publishing House,
New Delhi.
Thomsen, P.F. & Willerslev, E. (2015) Environmental DNA–an emerging tool in
conservation for monitoring past and present biodiversity. *Biological Conservation,* **183,**
4-18.
Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J., Thomsen, P.F., ... Dejean, T.
(2016) Next-generation monitoring of aquatic biodiversity using environmental DNA
metabarcoding. *Molecular Ecology,* **25,** 929-942.
Wickham, H. & Chang, W. (2016) ggplot2: create elegant data visualisations using the
grammar of graphics. Retrieved from https://CRAN.R-project.org/package=ggplot2.
Winfield, I.J. (2002) Monitoring lake fish communities for the Water Framework Directive: a
UK perspective. *TemaNord,* **566,** 69-72.
Winfield, I.J., Fletcher, J.M., James, J.B. & Bean, C.W. (2009) Assessment of fish
populations in still waters using hydroacoustics and survey gill netting: experiences with
Arctic charr (*Salvelinus alpinus*) in the UK. *Fisheries Research,* **96,** 30-38.
616