1

Centromeric CENP-A loading requires accurate mitotic timing, which is linked to checkpoint proteins

Anne Laure Pauleau[1,2,3], Andrea Bergner[1,2,3], Janko Kajtez[1,4], and Sylvia Erhardt[1,2,3]

[1] Center for Molecular Biology of Heidelberg University (ZMBH), Im Neuenheimer Feld 282, 69120 Heidelberg, Germany.

[2] DKFZ-ZMBH-Alliance, 69120 Heidelberg, Germany.

[3] CellNetworks Excellence Cluster, Heidelberg University, 69120 Heidelberg, Germany.

[4] current address: Department of Micro- and Nanotechnology, Technical University of Denmark, Produktionstorvet 423, DK-2800 Kgs. Lyngby, Denmark.

Correspondence to Sylvia Erhardt: s.erhardt@zmbh.uni-heidelberg.de

Abstract

A defining feature of centromeres is the presence of the histone H3 variant CENP-A that

replaces H3 in a subset of centromeric nucleosomes. In *Drosophila* cultured cells CENP-A

deposition at centromeres takes place during the metaphase stage of the cell cycle and

strictly depends on the presence of its specific chaperone CAL1. How CENP-A loading is

restricted to mitosis is unknown. We found that CAL1 interacts with the spindle assembly

checkpoint protein and RZZ complex component Zw10 and thus constitutes the anchor for

the recruitment of RZZ. We identify Zw10 as determinant of a mitotic duration and show

that it is inversely correlated with CENP-A levels at centromeres. Moreover, overexpression

of CAL1 is associated with increased CENP-A levels at centromeres and completely

uncouples CENP-A loading from mitosis. Therefore CAL1 controls CENP-A incorporation at

centromeres both quantitatively and temporally, connecting it to the spindle assembly

checkpoint to further ensure mitotic fidelity.

Introduction

The formation of two genetically identical daughter cells with a correct and stable genome is of utmost importance during mitosis. Condensed chromosomes are attached and segregated to the opposing poles of the dividing cell at anaphase by the mitotic spindle. At the interface between the chromosomes and the spindle microtubules lies the kinetochore. This multi-protein complex assembles during mitosis on a specialized domain of the chromosome called centromere, which is defined by an enrichment of nucleosomes containing the histone H3 variant CENP-A, also called CID in *Drosophila*. In many organisms, CENP-A recruits the components of the inner kinetochore components collectively known as CCAN (Constitutive Centromere-Associated Network), which in turn recruit the outer kinetochore components of the KMN network (formed by the Knl1 complex, the Mis12 complex and the Ndc80 complex) (Joglekar and Kukreja, 2017). The Ndc80 complex is mainly responsible for connecting microtubules with kinetochores while the Knl1 complex mostly functions to coordinate the Spindle Assembly Checkpoint (SAC) (Musacchio and Desai, 2017). The SAC delays entry into anaphase until all chromosomes are properly attached and aligned at the metaphase plate. The metaphase to anaphase transition is controlled by the activation by Cdc20 of the APC/C, a multisubunit ubiquitin ligase that triggers the degradation of cell cycle regulators by the proteasome (Kapanidou et al., 2017). The SAC proteins Mad2, BubR1, and Bub3 sequester Cdc20 forming the Mitotic Checkpoint Complex (MCC) thereby preventing the activation of the APC/C. Finally, the Mis12 complex serves as a hub interacting with all kinetochore complexes as well as with the inner centromere CCAN (Musacchio and Desai, 2017).

In some organisms, including *Drosophila melanogaster,* most CCAN proteins cannot be

identified with the exception of CENP-C (Westermann and Schleiffer, 2013). CENP-C has

been shown to act as a linker between CENP-A nucleosomes and the Mis12 complex

(Przewloka et al., 2011). CENP-C is also implicated in CENP-A replenishment at centromeres

during mitosis, serving as a platform to recruit the CENP-A chaperone CAL1 (Erhardt et al.,

2008). The *Drosophila* outer kinetochore is less complex than in other organisms as several

subunits of the KMN network including Mis12 complex subunit Dsn1, Knl1 complex subunit

Zwint, and Ndc80 complex subunit Spc24, are absent (Przewloka et al., 2007; Schittenhelm

et al., 2007). These changes affect the recruitment of the SAC proteins to the kinetochore

and their regulation, and it still remains to be analyzed how the *Drosophila* kinetochore has

solved this. In addition to the checkpoint proteins mentioned above, other proteins have

been implicated in SAC activity including Bub1, Mad1, the Mps1 and Aurora B kinases and

the RZZ complex (formed by the three proteins Rough Deal (ROD), Zw10 and Zwilch)

(Musacchio, 2015). In *Drosophila*, BubR1 and Bub1 localization to the kinetochore partly

depends on Spc105R (Schittenhelm et al., 2009), while Mad1 and Mad2 rely on the previous

recruitment of the RZZ complex (Buffin et al., 2005; Defachelles et al., 2015b). The RZZ

complex is conserved in all metazoans (Karess, 2005; Schmitt, 2010; Vleugel et al., 2012).

Besides Mad1-Mad2, RZZ also recruits Spindly and the motor protein Dynein, thereby being

essential for SAC silencing (Gassmann et al., 2008; Griffis et al., 2007; Starr et al., 1998;

Wojcik et al., 2001). Moreover, a recent study has shown that kinetochore size regulation

and maturation depends on RZZ and Spindly (Sacristan et al., 2018). RZZ recruitment to the

kinetochore at prometaphase has been shown to be mediated by KNL1 and Zwint in higher

eukaryotes (Caldas et al., 2015; Kasuboski et al., 2011) but no such dependency exist in

Drosophila. Therefore, we set out to identify the RZZ receptor in *Drosophila*. In the course of

these studies we found a surprising interaction of the RZZ complex with constitutive inner

kinetochore components and the loading machinery of CENP-A. Strikingly, we discovered a

co-dependence of mitotic duration and accurate CENP-A loading that seems mediated by

the interaction with these checkpoint proteins. Importantly, experimentally changing the

duration of mitosis affects CENP-A loading during mitosis, whereas increasing the pool of its

loading factor CAL1 uncouples CENP-A loading from mitosis. These data suggest an intricate

coordination of SAC, CENP-A loading, and mitotic duration.

Results

Zw10 mediates the recruitment of the RZZ to the kinetochore independently of Spc105R.

How RZZ is recruited to the kinetochore at mitosis onset remains elusive in *Drosophila*. We

first sought to determine if one of the RZZ components is responsible for targeting the

whole complex to the kinetochore. The RZZ complex is a conserved complex composed of

the three proteins, Rough Deal (ROD), Zwilch and Zw10 (Barisic and Geley, 2011; Karess,

2005; Schmitt, 2010). We established S2 cell lines stably expressing GFP-tagged ROD, Zwilch,

or Zw10 and found that the localization of these tagged proteins is in accordance with

endogenous proteins (Scaerou et al., 1999; Williams et al., 1992; Williams et al., 2003) as

well as transgenes expression (Basto et al., 2004; Buffin et al., 2005; Defachelles et al.,

2015b) reported in Drosophila tissues: GFP-Zw10 located in the cytoplasm during interphase

whereas low levels of GFP-Zwilch and GFP-ROD could be detected in the cytoplasm during

interphase (Fig. S1A-C). Following the disassembly of the nuclear envelope at the onset of

mitosis all three proteins were recruited to the kinetochore where they remained until

complete chromosome biorientation. RZZ proteins were then transported along the spindle

microtubules (Fig. S1A-C) in a process mediated by Spindly and Dynein required to silence

the SAC (Gassmann et al., 2008; Griffis et al., 2007; Starr et al., 1998; Wojcik et al., 2001).

Knockdown of all RZZ complex components (Fig. S1D-E) led to the known spindle

morphology defects in all cell lines (Fig. 1A-B) (Defachelles et al., 2015a; Karess and Glover,

1989; Williams et al., 1992; Williams et al., 2003). Importantly, depletion of any complex

component led to a reduction of the other GFP-RZZ proteins at kinetochores (Fig. 1A-B)

suggesting that the RZZ components are mutually dependent on each other similar to what

has been reported in Drosophila tissues (Defachelles et al., 2015a; Scaerou et al., 2001;

Wainman et al., 2012; Williams and Goldberg, 1994; Williams et al., 2003)(Fig. 1A).

However, levels of GFP-Zw10 at kinetochores were usually the least affected indicating that

some GFP-Zw10 is still recruited to kinetochores in absence of its complex partners ROD or

Zwilch (Fig. 1A-B). These data suggest that Zw10 may be the most proximal component of

the complex at kinetochores in Drosophila cells.

In mammalian cells, the recruitment of the RZZ complex depends on the presence of KNL1

and Zwint at kinetochores (Kops et al., 2005; Starr et al., 2000). As no ortholog of Zwint has

been identified in Drosophila so far, we tested if the Drosophila KNL1 ortholog Spc105R is

required for RZZ recruitment to the kinetochore. Depletion of Spc105R (Fig. S1D) did not

affect Zw10 localization (Fig. 1C-D). Reciprocally, depletion of Zw10 (Fig.S1D) did not

prevent Spc105R localization at the kinetochore (Fig. S2A-B). We then tested Mis12 (Fig.

S1D), which has been shown to localize to kinetochores throughout the cell cycle (Przewloka

et al., 2007) and found that it is also not involved in Zw10 recruitment to the kinetochore

(Fig. 1E-F). Therefore, we propose that the Drosophila checkpoint proteins and outer

kinetochore components are assembled through two independent branches: on the one

hand the RZZ complex, which is necessary to localize the MAD proteins to the kinetochore

(Buffin et al., 2005), and on the other hand Mis12-Spc105R, which are required to recruit all

others components of the KMN network as well as the BUB proteins (Basu et al., 1998; Orr

and Sunkel, 2010; Przewloka et al., 2007; Schittenhelm et al., 2009).

RZZ recruitment to the kinetochore requires direct interaction with inner centromere

components.

Like all kinetochore proteins tested so far (Przewloka et al., 2007), the localization of Zw10

to the kinetochore is abolished when the inner centromere protein CENP-A (Fig. 2A-B)

and/or its loading factor CAL1 (Fig. S2C) are depleted even though Zw10 protein levels

remained largely unchanged in Western blot analysis (Fig. S1E). Importantly, CENP-A and

CAL1 co-immunoprecipitated with GFP-Zw10 (Fig. 2C). Although we could observe GFP-ROD

by immunofluorescence, we were unable to detect it by immunoblotting precluding analysis

of ROD immunoprecipitate complexes. Regarding GFP-Zwilch we obtained inconsistent

results: CAL1 and CENP-A were immunoprecipitated in some but not all experiments. Zw10

is also known to form a complex with the *Drosophila* RINT1 at the Golgi during interphase

(Wainman et al., 2012) (Fig. S2D). We observed that GFP-RINT1 immunoprecipitated with

Zw10 but neither with CAL1 nor CENP-A indicating that there are, indeed, two Zw10-

containing complexes and that Zw10 interacts with CENP-A and CAL1 when it is in a complex

with Zwilch and ROD in mitosis (Fig. 2C). To confirm these results, we performed Yeast-two-

hybrid (YTH) experiments that repeatedly showed that Zw10 weakly but consistently binds

to full length CAL1 and CENP-A (Fig. 2D). Since YTH interaction are often challenging with

bigger proteins, we repeated the YTH analysis with yeasts expressing fragments of CAL1 and

Zw10 and identified strong interactions of the C-terminus (amino acids 680-979) of CAL1

and the C-terminus of Zw10 (amino acids 482-721) (Fig.S2E-F). The C-terminus of CAL1 also

binds CENP-C and Roadkill (Bade et al., 2014; Schittenhelm et al., 2010). No further

interactions were detected between the other components of the complex and the

centromeric proteins supporting our previous observation that Zw10 might be responsible

for targeting the RZZ complex to the kinetochore.

Zw10 affects the loading of newly synthesized CENP-A during mitosis.

A prominent function of the RZZ complex is to recruit the MAD proteins to the kinetochore,

thereby making the RZZ essential for the SAC (Buffin et al., 2005). SAC activity peaks in

prometaphase when CENP-A incorporation into centromeric chromatin takes place in

Drosophila cells (Mellone et al., 2011). Therefore, we sought to identify a possible link

between SAC activity and CENP-A loading. We used the SNAP-tag technology (Jansen et al.,

2007) in a quench-chase-pulse experiment to label newly synthesized and incorporated

CENP-A. At day 3 of Zw10 RNAi, we quenched existing SNAP-CENP-A molecules with a non-

fluorescent dye, cultured the cells for 24 hours to allow approximately one additional cell

division, then marked newly synthesized SNAP-CENP-A molecules with a fluorophore and

measure fluorescent intensity of newly incorporated SNAP-CENP-A at centromeres (Fig. 3A).

We found that Zw10 depletion led to a 30% reduction of newly synthesized SNAP-CENP-A

(Fig. 3A-B) and SNAP-CAL1 (Fig. S3A-B) at centromeres compared to control-depleted cells.

This significant decrease of SNAP-CENP-A centromeric levels may be caused either by a

decreased stability or by decreased loading. Since stability of centromeric SNAP-CENP-A was

not affected by Zw10 depletion as determined by SNAP-tag pulse chase experiment (Fig.

S3C-D), we hypothesized that Zw10 depletion interferes with SNAP-CENP-A loading directly.

To address a potential role of Zw10 in mitotic loading of CENP-A we measured Fluorescence

Recovery After Photobleaching (FRAP) of CENP-A, on the assumption that fast recovery of

the signal at centromeres corresponds to active loading of CENP-A. We partially bleached

GFP-CENP-A signal during prophase and followed recovery of the signal at centromeres until

anaphase. We observed a partial recovery of GFP-CENP-A at centromeres in control cells

confirming previous reports that loading of new CENP-A takes place during mitosis (Mellone

et al., 2011) (Fig. 3C). Importantly, the recovery rate of GFP-CENP-A at centromeres was

comparable in control and Zw10-depleted cells (Fig. 3D-E). Thus, the CENP-A mitotic loading

machinery seemed functional in Zw10-depleted cells and does not explain the observed

impaired loading of CENP-A.

Besides mitotic loading, additional CENP-A loading has been reported in G1 in S2 cells

(Lidsky et al., 2013). We, therefore, followed cells through mitosis, partially bleached GFP-

CENP-A in early G1 and measured GFP-CENP-A signal intensity for 3 hours but did not detect

any recovery of the bleached GFP-CENP-A signal (Fig. S3E-F). We explained these results

with a likely absence of any active and measurable CENP-A loading in early G1. Strikingly,

however, during the course of these live cell analyses we noticed that Zw10-depleted cells

displayed a shorter mitosis compared to control cells (Fig. 3C). This result is in accordance

with the Premature Sister Chromatids Separation (PSCS) phenotype observed in Zw10

mutant embryos and larval tissues (Basto et al., 2000; Scaerou et al., 1999; Williams et al.,

1992).

CENP-A loading correlates with mitosis duration.

To confirm that Zw10-depleted cells proceed faster through mitosis, we performed time-

lapse microscopy analysis of cells stably expressing H2B-GFP and mCherry-Tubulin. Indeed,

mitotic timing (defined as the time that elapses between late prophase (determined by the

partial nuclear envelop breakdown (NEB) concomitant with the formation of two spindle

poles) and anaphase entry) was significantly shorter in Zw10-depleted cells than in control

cells (Fig. 4A-B; videos 1-2). This raised the possibility that decreased SNAP-CENP-A

incorporation in Zw10-depleted cells could be a consequence of a shortened mitosis. To test

this, we depleted two other mitotic checkpoint proteins, Mad2 and BubR1, known to reduce

mitosis duration (Lince-Faria et al., 2009; Orr et al., 2007; Rahmani et al., 2009). These

depletions (Fig. S4A) also led to a decrease of newly synthesized centromeric SNAP-CENP-A

incorporation similar to Zw10 depletion (Fig. 4C-D). On the contrary, depletion of Mis12 (Fig.

S4A), which has been previously shown not to affect duration of mitosis in Drosophila

cultured cells (Feijao et al., 2013), did not alter SNAP-CENP-A amount at centromeres (Fig.

4C-D). Therefore, we hypothesized that the amount of newly synthesized CENP-A that can

be incorporated into centromeric chromatin depends on mitosis duration. Yet, extending

metaphase duration by preventing entry into anaphase through depletion the APC/C

cofactor Cdc27 (Huang and Raff, 2002) or Spindly (required for silencing of the SAC) (Griffis

et al., 2007) (Fig. S4A) did not cause an increase of CENP-A levels at centromeres when

compared to control (Fig. 4C-D). Arrest in prometaphase by the microtubule-depolymerizing

drug Nocodazole or in metaphase by the proteasome inhibitor MG132 also did not cause

changes in loading of newly synthesized SNAP-CENP-A (Fig. S4B-C). SNAP-CAL1 centromeric

loading was similarly affected: an acceleration of mitotic timing by BubR1 or Mad2 RNAi let

to a decrease of loading, whereas mitotic arrest by Cdc27 or Spindly RNAi did not affect

loading (Fig. S4D). This set of experiments suggests that the amount of new CENP-A

molecules that are incorporated into centromeric chromatin is limited at each round of

mitosis. Importantly, FRAP analysis revealed an inverse correlation between CENP-A

incorporation rate and mitosis duration in control cells (Fig. 4E), which suggested that either

CENP-A loading or CENP-A centromeric levels directly controls the duration of mitosis.

CENP-A levels correlate with mitosis duration.

To test if there is a direct correlation between centromeric CENP-A levels and mitosis length

we needed an experimental system that reduces CENP-A levels without arresting the cells in

mitosis due to the known phenotypes associated with CENP-A depletion (e.g. chromosome

segregation defects) that are impedimental to study mitosis progression (Blower and

Karpen, 2001). Therefore, we performed CENP-A RNAi depletion in GFP-CENP-A

overexpressing cells. Interestingly, depletion of CENP-A in GFP-CENP-A cells led to

undetectable levels of endogenous CENP-A but left sufficient GFP-CENP-A to allow some

cells to go through mitosis (Fig. 5A-B). As hypothesized, mitosis duration was extended in

CENP-A-reduced cells in comparison to control cells (Fig. 5B-C; videos 3-4) as observed in

CENP-A mutant fly embryos (Blower et al., 2006), supporting our hypothesis that CENP-A

levels might control mitotic length.

We next sought to determine if elevated CENP-A levels could also influence mitosis

duration. However, upregulation of CENP-A to high levels is associated with ectopic CENP-A

incorporation into chromosome arms and subsequent defects in mitosis (Heun et al., 2006).

In order to avoid this secondary effect, we induced CENP-A overexpression from pMT-CENP-

A-GFP plasmid with a short and very low $CuSO_4$ pulse. Under those conditions, CENP-A-GFP

was over-expressed while endogenous CENP-A levels were unaffected (Fig. 5D). However,

22 hours after the low $CuSO_4$ pulse endogenous CENP-A was almost undetectable, indicating

that the cell compensate ectopic CENP-A overexpression by decreasing its endogenous

counterpart (Fig. 5D). We also observed a reduction of endogenous CENP-A in cells

expressing constitutively GFP-CENP-A and SNAP-CENP-A (Fig. 5D). Under these conditions,

non-centromeric CENP-A was undetectable and CENP-A-GFP localized to centromeres (Fig.

5E). Using time-lapse analysis we found that the cells with a mild increase of CENP-A

progressed significantly faster through mitosis (Fig. 5G-H; videos 5-6). Importantly we

observed no increase in mitotic defects usually associated with CENP-A overexpression such

as chromosomes misalignment and lagging chromosomes (Fig 5I). Together these results

confirm our hypothesis that centromeric CENP-A levels influence mitosis duration.

CAL1 levels regulate the amount of CENP-A at centromere.

To confirm our observations that CENP-A levels influences mitosis duration, we sought to

manipulate CENP-A indirectly. Centromeric CENP-A loading is mediated by its loading factor

CAL1 (Chen et al., 2014; Erhardt et al., 2008). In addition, CAL1 has been shown to be the

limiting factor of CENP-A incorporation in flies (Schittenhelm et al., 2010). We

overexpressed CAL1 in S2 cells from a $CuSO_4$-inducible pMT promoter and observed an

increase of both, endogenous and overexpressed CENP-A (GFP-CENP-A, SNAP-CENP-A)

levels (Fig. 6A). Overexpressed CAL1 primarily localized to centromeres and the nucleolus

similarly to the endogenous protein (84 % of cells showing CAL1-V5). Cytoplasmic and/or

nucleoplasmic CAL1 were visible in about 15 % of the cells but CENP-A was only localized to

centromeres (Fig. 6B) and was not detectable at any other chromatin regions on metaphase

chromosomes (Fig. 6C). This result indicates that CAL1 overexpression alone does not lead

to ectopic incorporation of CENP-A. Importantly, however, and in contrast to previous

observations in embryos (Schittenhelm et al., 2010), centromeric CENP-A levels were

massively increased, up to 200%, after 24 hours of CAL1 overexpression (Fig. 6D).

Surprisingly, CENP-C levels remained largely unaffected by CAL1 overexpression (Fig. 6E-F).

These results show that the centromeric amount of CENP-A is regulated by the available

amount of CAL1. To test whether the increased CENP-A level is due to more CENP-A loading

we overexpressed CAL1 in SNAP-CENP-A expressing cells and measured incorporation of

newly synthesized CENP-A. As expected, significantly more SNAP-CENP-A incorporated into

centromeres in CAL1 overexpressing cells (Fig. 6G-H). These data confirm that CENP-A

centromeric levels are regulated by CAL1 and that CAL1 overexpression specifically

increases incorporation of newly synthesized CENP-A in centromeric chromatin.

Overexpression of CAL1 leads to centromeric CENP-A loading outside of mitosis.

Since CAL1 overexpression is suitable to study the effect of CENP-A levels on mitosis

duration we performed time-lapse microscopy experiments on CAL1-overexpressing cells

and, indeed, found that they proceed faster through mitosis compared to control cells (Fig.

7A-B; videos 7-8). Of note, CAL1 overexpressing cells did neither exhibit more mitotic

defects (Fig. S5A) nor did they recruit more endogenous kinetochore proteins (CENP-C,

Spc105R, BubR1) during prometaphase (Fig. S5B). CAL1-overexpressing cells also display a

functional SAC because cells treated with the microtubule interfering drug Taxol triggered a

cell cycle arrest in prometaphase in control cells as well as in CAL1-overexpressing cells (Fig.

S5C-D). We concluded that CAL1-overexpressing cells not only incorporated more CENP-A at

centromeres, they also progressed faster through mitosis.

To determine if CAL1 overexpression accelerates CENP-A loading during mitosis, we

performed FRAP experiments and measured the recovery rate of GFP-CENP-A after

photobleaching in prophase. Surprisingly, GFP-CENP-A recovery rate decreased in CAL1-

overexpressing cells as compared to control cells (Fig. 7C-D). In addition, the inverse

correlation that we observed in control cells between CENP-A incorporation rate at

centromeres and mitosis duration was not preserved in CAL1-overexpressing cells. This

suggests that rather than the rate of incorporation, the amount of CENP-A itself correlates

with mitosis duration in control cells.

Since the availability of CAL1 seems to be the crucial aspect for CENP-A incorporation, we

tested if CAL1-misexpressing cells incorporate CENP-A into centromeric chromatin outside

mitosis. Thus, we arrested cells at the metaphase to anaphase transition using MG132

treatment for 2 hours, blocked existing SNAP-CENP-A molecules using SNAP-Block and

cultured the cells for an additional 4 hours in the presence of MG132 (Fig. S6A) similar to a

previously published protocol (Mellone et al., 2011). Whereas we observed about 10%

SNAP-CENP-A loading in uninduced pMT-CAL1 cells, the induction of pMT-CAL1 and

resulting overexpression of CAL1 let to an incorporation of newly synthesized SNAP-CENP-A

in about 40% of the cells pointing to loading outside of mitosis (Fig. S6A). To further confirm

that overexpressed CAL1, indeed, loads CENP-A to centromeres outside of mitosis, we

followed newly synthesized SNAP-CENP-A live by time-lapse microscopy. We blocked

existing SNAP-CENP-A with SNAP-Block, washed and added SNAP-640 fluorophore

(Butkevich et al., 2017) to the culture medium in order to stain newly synthesized SNAP-

CENP-A molecules. Despite some aggregation of the dye, SNAP-CENP-A loading at

centromeres was discernable in both, CAL1 overexpressing and control cells. We only

analyzed cycling cells that eventually entered mitosis during the course of imaging and determined whether and when they incorporated SNAP-CENP-A. We found that in the uninduced control cells 38% had SNAP-CENP-A signals at centromeres by the end of the imaging (Fig. 7E-F; videos 9). Of those, 91 % loaded SNAP-CENP-A during mitosis and 9% after the cell exited mitosis. In contrast, 81% of cycling CAL1-overexpressing cells had SNAP-CENP-A signals at centromeres (Fig. 7E-F; videos 10), and of those 31% loaded before mitosis and 51% after the cells exited of mitosis. The specificity of the signal was verified by co-staining with anti-CENP-A antibody immediately after block (0 min), and 20 hours of incubation with SNAP-640 (Fig. S6B). No SNAP-CENP-A was detected directly after SNAP-block treatment, confirming the efficient quenching of old SNAP-CENP-A molecules (Fig. S6B). After 20 hours newly synthesized SNAP-CENP-A perfectly co-localized with total CENP-A signals at centromeres. Interestingly, when we counted the percentage of cells that displayed SNAP-CENP-A at centromeres in these fixed samples we did not observe any differences between control and CAL1-overexpressing cells after 20 h (Fig. S6C). However, SNAP-CENP-A centromeric intensity was four times higher in CAL1-overexpressing cells than in control cells (Fig. S6D). This suggests that low SNAP-CENP-A protein levels were masked in the presence of excess SNAP dye in live cell analysis. Nevertheless, the analysis of SNAP-CENP-A loading in cycling cells revealed that cells overexpressing CAL1 loaded CENP-A throughout the cell cycle while loading outside mitosis did not occur in control cells (Fig. 7G). In conclusion, overexpression of the CENP-A-specific chaperone CAL1 uncouples CENP-A centromeric loading from mitosis and leads to an uncontrolled overload of CENP-A at centromeres that is usually prevented by a crosstalk with the SAC and the length of mitosis.

Discussion

Many central components of the spindle assembly checkpoint require outer kinetochore

components for their localization to centromeric regions. In yeast and worms, for instance,

Mad1/2 recruitment depends on KNL1 (London and Biggins, 2014; Moyle et al., 2014), in

other eukaryotes, Mad1/Mad2 recruitment to the kinetochore also requires the RZZ

complex (Caldas et al., 2015; Kops et al., 2005; Silio et al., 2015). In mammalian cells, RZZ

recruitment to the kinetochore depends on Bub1, which itself requires KNL1/Zwint (Caldas

et al., 2015; Varma et al., 2013; Zhang et al., 2015) although experimental results suggest

the existence of a KNL1-independent mechanism (Caldas et al., 2015; Petsalaki et al., 2018;

Silio et al., 2015). In *Drosophila*, many outer kinetochore components are missing

(Przewloka et al., 2007; Schittenhelm et al., 2007) and even though Mad1/2 recruitment to

the kinetochore also depends on the RZZ complex (Buffin et al., 2005) the factors necessary

for the localization of the RZZ to kinetochores and whether RZZ and spindly are required for

kinetochore expansion remain elusive (Sacristan et al., 2018). In this paper we show that

RZZ localization to the kinetochores does not require KNL1[Spc105R] but depends on the inner

kinetochore proteins CAL1 and CENP-A. Therefore, we hypothesize that the *Drosophila*

outer kinetochore and components of the SAC assemble through two independent

pathways: the CENP-C-KMN-Bub1-Bub3/BubR1 branch (Basu et al., 1998; Przewloka et al.,

2011; Schittenhelm et al., 2009) or the here described CAL1-RZZ-Mad1/2 branch. How those

two pathways interconnect for the formation of MCC complexes remains to be determined.

One link may be the KMN complex, since Mad2 has been shown to be diminished in the

absence of KMN proteins (Feijao et al., 2013). Interestingly, Spc105R mutation does not

affect SAC function in fly embryos suggesting that flies rely more on the RZZ-Mad1/2 branch

to engage the SAC (Schittenhelm et al., 2009).

We show here that CAL1 binds to Zw10 through its C-terminus, the same region required for

its interaction with CENP-C (Schittenhelm et al., 2010) and the E3 ligase adaptor Roadkill

(Bade et al., 2014). On the other hand, no binding partner of CAL1 N-terminal region other

than CENP-A has been identified so far. This suggests that, while the interaction between

CAL1 and CENP-A is very stable, the C-terminal interacting partners of CAL1 are potentially

of regulatory nature and more versatile, as one would expect for component of the SAC.

In *Drosophila* cells, CENP-A loading primarily takes place during prometaphase-metaphase

(Mellone et al., 2011). Additional turnover of CENP-A in G1 has been reported leading to the

hypothesis that CENP-A could be further incorporated at this stage (Lidsky et al., 2013). We

did not observe any CENP-A turnover during early G1 in our FRAP experiments when we

bleached centromeric CENP-A at the end of cytokinesis. However, our FRAP experiments

and our live staining of newly synthesized SNAP-CENP-A confirmed the observations

reported by Mellone et al., that the majority of CENP-A loading takes place during mitosis in

Drosophila cultured cells (Mellone et al., 2011).

With SNAP-tagged CENP-A we showed that centromeric loading of newly synthesized CENP-

A was decreased in Zw10-deficient cells even though the loading machinery for CENP-A

mitotic incorporation was functional in these cells. Importantly, we found that Zw10 affects

the length of mitosis allowing sufficient time for chromosomes to attach and align. Since

CENP-A loading occurs during this time window, we investigated a possible link between

CENP-A loading and mitotic timing. Indeed, shortening mitosis duration through the

depletion of the two proteins Mad2 and BubR1 previously shown to affect mitosis duration

(Meraldi et al., 2004) recapitulated the decreased CENP-A loading observed after Zw10

depletion strongly arguing that mitosis duration affects CENP-A loading. Elongating the

mitotic time window during which CENP-A is usually loaded (Spindly, Cdc27 depletion or by

drug treatment) did not increase the amount of CENP-A incorporated at centromeres,

showing that only a defined amount of CENP-A can be incorporated at each mitosis. How

CENP-A levels at centromeres are sensed is unclear; one model is that BubR1 and Mad2

function as mitotic timer (Buffin et al., 2007; Lince-Faria et al., 2009; Meraldi et al., 2004;

Rahmani et al., 2009) and we could envision a role of Zw10 in this process. Mad2 but not

Mad1 has been shown to be a central determinant of the mitotic timing (Emre et al., 2011;

Meraldi et al., 2004), leading to the hypothesis that cytosolic Mad2-Cdc20 complexes form

independently of Mad1 and determine the timing of mitosis (Emre et al., 2011; Li et al.,

2017). It has previously been proposed that low CENP-A levels at centromeres lead to more

cytosolic Mad2 thereby keeping the timer active longer (Blower et al., 2006). We

hypothesized that higher CENP-A levels at centromeres during early mitosis would

accelerate the recruitment of RZZ and consequently Mad2 to the kinetochores, therefore,

releasing the timer and shortening mitosis duration in cells where kinetochores attach

properly to the spindle microtubules. Interestingly, Nocodazole treatment did not affect

CENP-A loading confirming previous observations that kinetochores attachment to the

microtubule spindle does not play a role in CENP-A loading (Mellone et al., 2011). These

results are pointing further to an additional function of the SAC independent of the control

of microtubule attachment. However, recent evidence show that RZZ together with Spindly

plays a central role in kinetochore expansion during early mitosis to form a fibrous corona

that then compacts upon microtubule capture (Sacristan et al., 2018). How this plays a role

in CENP-A loading needs to be investigating.

In flies, CENP-A incorporation is controlled by its chaperone CAL1 (Chen et al., 2014; Erhardt

et al., 2008; Schittenhelm et al., 2010). We found that overexpression of CAL1 leads to

increased CENP-A protein levels but Chen et al., previously reported that CAL1-GFP-LacI

overexpression alone did not affect CENP-A when targeted to an exogenous LacO array

(Chen et al., 2014). Under our conditions, CAL1-V5 was strongly overexpressed and caused

an intense centromeric and nucleolar localization of CAL1 and a centromere-specific

overload of CENP-A. Importantly, however, ectopic incorporation of CENP-A was never

observed further enhancing the evidence that CAL1 loads CENP-A exclusively to

centromeres. Important for our study is that the increased centromeric CENP-A levels

following CAL1 overexpression correlated with faster mitosis. A similar acceleration of

mitotic timing was observed when CENP-A was overexpressed in a controlled manner that

did not lead to massive incorporation at ectopic sites.

The live analysis of newly synthesized CENP-A allowed us to visualize CENP-A incorporation

during mitosis. CAL1-overexpressing cells loaded CENP-A in all stages of the cell cycle. This

strongly suggests that CAL1's availability and protein amount control CENP-A incorporation

into centromeric chromatin both quantitatively and temporally. CENP-A expression (Ahmad

and Henikoff, 2001) and its stability (Bade et al., 2014; Moreno-Moreno et al., 2006)

together with its dependence on the low abundant and highly specific loading factor CAL1

and the here shown connection to mitotic events are likely interconnected cellular

surveillance mechanisms to avoid misincorporation of CENP-A and, therefore, securing

genome stability. How CAL1 itself is regulated to obtain such specificity is currently

unknown. In conclusion, we show that there is a direct crosstalk between the SAC and the

maintenance of centromeric chromatin, ensuring mitotic fidelity not only by controlling

microtubule attachment but also by regulating accurate composition of centromeres.

Materials and methods

Gene constructs

Plasmid encoding CENP-A tagged with a N-terminal EGFP has been described previously (Erhardt et al., 2008) as well as those encoding CENP-A and CAL1 with N-terminal SNAP tag (Mellone et al., 2011). Constructs created for this study (GFP-ROD, GFP-Zwilch, GFP-Zw10, GFP-RINT1) were cloned into AscI and PacI sites of the pCopia–localization and purification (LAP) vector with a basal expression Copia promoter and an N-terminal mEGFP tag (Erhardt et al., 2008). For overexpression studies, CAL1-V5 and CENP-A-GFP were cloned into pMT vector (Invitrogen). For yeast two hybrid studies, pMM6 plasmids containing full length CAL1, CAL1 truncations and CENP-A have been described in (Bade et al., 2014). Zw10 (full length, aa1-240, aa241-481, aa482-721), Zwilch and Rod (full length, aa1-696, aa697-1392, aa1393-2090) were cloned into pMM5 plasmids.

Cell culture

Cell maintenance

Drosophila S2 cells were maintained under sterile conditions at 25 °C in Schneider medium containing 10 % heat-inactivated fetal bovine serum (FBS) and 100 µg/ml Penicillin/Streptomycin. Expression of genes under the metallothionein promoter (pMT vector) was induced by supplementing the medium with $CuSO_4$ (CENP-A-GFP: 10 µM; CAL1-V5: 100 µM).

RNAi and Transfection Experiments

Double-stranded RNA (dsRNA) was produced using the MEGAscript (Ambion) kit according to the manufacturer's protocol. RNAi was carried out as described previously (Erhardt et al.,

2008). In short, 10^6 S2 cells were incubated for 1 h with 15 μg dsRNA in serum-free medium.

Subsequently, 2 volumes of 15 % serum containing medium were added. Analysis was

usually performed 96 h after treatment.

Stable cell lines were generated by co-transfecting the plasmid of interest together with a

plasmid carrying the resistance gene for hygromycin with Cellfectin II (Life Technologies),

and the selection process was started 2 d after transfection by supplementing the media

with 250 μg/ml Hygromycin B.

SNAP quench chase pulse experiments

Cells overexpressing either SNAP-tagged CENP-A or CAL1 proteins were treated with dsRNA

as described in RNAi section. After 3 days, cells were collected and incubated in fresh SM

containing 10 μM SNAP-Cell Block (New England Biolabs S9106), for 30 min at 25 °C, 400

rpm. Cells were then pelleted, washed 3 times (1st wash for 30 min, at 25 °C, 400 rpm) and

plated in conditional medium. After 24 h, cells were collected and incubated with 4 μM

SNAP-Cell TMR-Star (New England Biolabs S9105) or SNAP-SiR647 (New England Biolabs

S9102) for 15 min at 25 °C, 400 rpm, protected from light. The cells were then pelleted,

washed 3 times with SM (1st wash for 30 min, at 25 °C, 400 rpm), one time in PBS and

prepared for imaging. Cells were settled on a glass coverslip for 10 min, fixed with 4 % PFA

for 10 min, washed three times in PBS, and counterstained with DAPI before mounting.

SNAP pulse-chase

Cells overexpressing either SNAP-tagged CENP-A or CAL1 proteins were treated with dsRNA

as described in RNAi section. After 3 days, cells were collected and incubated in fresh

medium containing 4 μM SNAP-Cell TMR-Star (New England Biolabs, Inc.) for 15 min at 25

°C, 400 rpm, protected from light. Cells were then washed 3 times with SM (1st wash for 30

min, at 25 °C, 400 rpm) and plated in conditional medium. After 24 h, cells were collected

and prepared for imaging as previously described.

SNAP live imaging

To follow incorporation of newly synthesized SNAP-CENP-A, SNAP-CENP-A was blocked as

previously described (see section SNAP quench chase pulse). Cells were then resuspended in

conditional medium containing 0.5 µM SNAP-640 dye (Butkevich et al., 2017) and placed

into an Ibidi imaging chamber before imaging.

Immunofluorescence

IF was essentially performed as described previously (Erhardt et al., 2008). Cells were

harvested, washed once with PBS and placed to settle on a glass coverslip for 10 min before

fixation with 4 % PFA for 10 min. Cells were washed three times in PBS and permeabilized

for 5 min with PBS containing with 0.5 % Triton X-100. Unspecific binding was prevented by

blocking the cells for 30 min with 1 % BSA in PBS. Primary antibodies diluted in blocking

solution were incubated for 30 min at RT. After three washes with PBS, cells were incubated

with the corresponding, fluorescently labeled secondary antibodies (diluted in blocking

solution) for 30 min at RT. The secondary antibody incubation and all the following steps

were performed while protected from light. After three washes in PBS, DNA was stained for

5 min with DAPI (1 µg/ml in PBS). The cells were then washed two more times with PBS

before mounting in Aqua/Polymount medium (Polysciences, Inc.) on a glass slide.

Preparation of mitotic chromosome spreads

Mitotic chromosome spreads were essentially performed as described previously (Mathew

et al., 2014). To obtain mitotic chromosomes, 2×10^5 exponentially growing cells were

arrested in mitosis with 2.5 µg/µl Colcemid for 1 h, centrifuged for 3 min at 800 *g*,

resuspended in 0.5 ml hypotonic sodium citrate solution (0.5 % Na-citrate in ddH2O), and

incubated for 8–10 min. Cells were spun on positively charged slides in a cytocentrifuge

(Shandon 4 Cytospin; Thermo Fisher Scientific) at 900 rpm for 10 min. Mitotic chromosome

spreads were fixed with 4 % PFA for 10 min and subsequently treated for IF.

Microscopy and Image analysis

All Images were acquired on a DeltaVision Core system (Applied Precision) equipped with an

Olympus UPlanSApo 100X oil immersion objective (n.a. 1.4) at binning 1x1 (mitotic figures)

or 2x2 (interphase). If not stated otherwise, images were taken as z stacks of 0.3 µm

increments. All images were deconvolved (20 cycles additive with noise filtering) and

maximum-projected using the Applied Precisions soft-WoRx 3.7.1 suite. Image

quantification was done using ImageJ software. For determining the number and intensities

of centromeric dots of CAL1 and CENP-A, a set of plugins developed by the Nikon Imaging

Centre, University Heidelberg was used. After marking the nuclear boundaries in the DAPI

channel and saving them with the ROI tool, the spots in the channel of interest were

enhanced using the DoG spot enhancer plugin. The threshold was adjusted using ''Li''

instead of default settings. The number of spots with their mean intensities relative to the

nuclear area was determined using the ROI particle analyzer plugin.

Time-lapse imaging was performed on a DeltaVision Core system (Applied Precision)

equipped either with an Olympus 60X/1.42, N Plan Apo oil immersion objective or an

Olympus 40X/1.35, UApo/340 oil objective, at binning 2x2 with 3 min (mitosis duration and

fidelity measurements) or 15 min (SNAP-CENP-A live imaging) time-lapse series, for 16 h,

with z stacks of 1 µm increment.

FRAP was performed on a Zeiss LSM 780 confocal microscope equipped with a 63x/1.4 oil

immersion objective. Number of focal planes per time point, as well as time intervals and

total duration were adjusted according to experimental needs. The spacing of focal planes

was 1 μm in all experiments. Recovery of EGFP-CENP-A signal was analyzed using FRAP

analysis plugin developed by ZMBH imaging facility. After marking the boundaries of the cell

of interest, the threshold allowing visualization of centromeric signal exclusively was defined

on pre-bleach image. Total intensity of the thresholded signal was then measured for each

time point as well as the area. Similar measurements were performed on non-bleached cells

from the same area to determine the photobleaching level in each time-series. After

correction, EGFP-CENP-A signal intensity was plotted against the time, a linear regression

curve was fitted and the rate of recovery extracted.

Immunoprecipitation

GFP-tagged proteins were isolated using a GFP-specific single chain antibody (GBP) coupled

to NHS-activated Sepharose (GE) (Rothbauer et al., 2008). A total of 1.0–2.0 x 10^8 cells were

lysed in two pellet volumes of 20 mM Tris (pH 7.5), 1 % NP-40, 250 mM NaCl, 2 mM PMSF, 2

mg/ml aprotinin, 5 mg/ml leupeptin, 1 mg/ml pepstatin and Roche anti-proteases cocktail

tablet for 15 min at 4 °C. After clearing the lysate by centrifugation, the supernatant was

incubated for 2 h at 4 °C with 50 μl pre-equilibated GBP-beads. The beads were washed six

times with lysis buffer and eluted with 2x Laemmli buffer 5 min at 95 °C.

Antibodies

The following primary antibodies were used: rabbit H3 (Abcam AB 1791), rabbit CAL1 (Bade

et al., 2014), chicken CENP-A (from Dr. Heun), rabbit CENP-A (Active Motif 39713), rabbit

YFP (from Pr. Bukau), mouse tubulin (Sigma T9026), mouse V5 (Invitrogen V8012), sheep

Scp105R (from Dr. Glover), rabbit Zw10 (from Dr. Goldberg), guinea pig CENP-C (Erhardt et

al., 2008), rabbit phospho-S10 histone H3 (Abcam ab5176), BubR1 (from Dr. Glover).

Secondary antibodies coupled to Alexa Fluor 488, Alexa Fluor 546 and Alexa Fluor 647

fluorophores (Molecular Probes) were used for IF, and horseradish peroxidase-conjugated

secondary antibodies (Sigma) were used for WB analysis.

Reverse transcription and Quantitative qPCR

RNA was isolated using Trizol(R) according to the manufacturer's procedures. Whole

genome cDNA synthesis was performed using the Quantitect Reverse Transcription kit

(Invitrogen), with a combination of oligo (dT) and random hexamer primers in equal

proportions. A control reaction with no reverse transcription was always performed in

parallel.

qPCR was performed after cDNA synthesis on a LightCycler 480 (Roche) using LightCycler

480 SYBR Green I Master (Roche). All reactions were run in triplicate in a LightCycler 480

multiwell plate. Actin was used as a reference. The level of each targeted gene in the control

mock-treated sample was normalized to 1, and compared with the corresponding–depleted

samples.

Yeast two hybrid

Protein-protein interactions were tested using pMM5 and pMM6 fusion constructs and the

yeast strain SGY37VIII. The YTH was performed as described previously (Schramm et al.,

2000). In short, interactions were judged based on the activity of β-galactosidase that

results in the conversion of X-Gal (5-bromo-4-chloro-3-indolyl-b-D-galactosidase) into a blue

dye.

Acknowledgements:

We thank B. Bukau, D. Glover, M. Przewloka, M. Goldberg, R. Karess for antibodies and

constructs, V. Belov, A. Butkevich and the Facility of synthetic chemistry of the Max Planck

Institute for Biophysical Chemistry (MPIBPC) for the gift of the compound 640SiRH-BG. We

thank the ZMBH Imaging Facility and A. Jafar Pour for ImageJ FRAP analysis plugins; H.

Lorenz for help with imaging, R. Vlijm and the Erhardt laboratory members for discussions;

E. Schiebel and S. Corless for critical comments on the manuscript. We acknowledge funding

from the Deutsche Forschungsgemeinschaft through the grant EXC81 (CellNetworks),

ER576/2-2, SFB1036, the European Research Council through the grant ERC-CoG-682496

(cenRNA) to SE. ALP is an alumna of the CellNetworks postdoc program.

Competing interests: No competing interests declared

      Figures legends

Figure 1: RZZ recruitment to kinetochores does not require Spc105R in *Drosophila*.

A. Immunofluorescence of cells expressing GFP-Rod, GFP-Zw10 or GFP-Zwilch in

combination with mCherry-Tubulin after depletion of RZZ components as indicated. After 96

h dsRNA treatment, cells were fixed and stained with anti-GFP antibody (green) and anti-

CENP-A (red), DNA was counterstained with DAPI. **B. *Upper panel.*** Prometaphase cells from

A. were counted for the presence or absence of GFP signal at kinetochores. ***Lower panel.***

Quantification of A. showing the total GFP-RZZ kinetochore intensity per cell. GFP

fluorescence intensity at kinetochores was measured for each cell and then normalized

within one experiment allowing pooling of the measurements for at least 3 experiments per

condition. **C.** Immunofluorescence of GFP-Zw10/mCherry-Tubulin expressing cells after

depletion of Spc105R. At day 4 of RNAi, cells were fixed and stained with anti-GFP (green)

and anti-Spc105R antibodies (red), DNA was counterstained with DAPI. **D.** Quantifications

of C. ***Left.*** Prometaphase cells were counted for the presence or absence of GFP-Zw10 signal

at kinetochores. ***Right.*** Total GFP-Zw10 kinetochore intensity per cell measured as in B. **E.**

Immunofluorescence of GFP-Zw10/mCherry-Tubulin expressing cells after depletion of

Mis12. After 96 h dsRNA treatment, cells were fixed and stained with anti-GFP (green) and

anti-Spc105R (red) antibodies, DNA was counterstained with DAPI. **F.** Quantifications of E.

Left. Prometaphase cells were counted for the presence or absence of GFP-Zw10 signal at

kinetochores. ***Right.*** Total GFP-Zw10 kinetochore intensity per cell measured as in B. All

graphs show Mean +/- SEM of 3 experiments (n>90 cells), Student t-test (n.s.: non-

significant; *: $p<0.05$; **: $p<0.01$; ***: $p<0.001$). Scale bar= 2 μm.

Figure 2: RZZ recruitment to the kinetochore depends on a direct interaction with inner

centromere proteins.

A. Immunofluorescence of GFP-Zw10/mCherry-Tubulin expressing cells after depletion of

CENP-A. At day 4 of RNAi, cells were fixed and stained with anti-GFP (green), anti-CENP-A

(red) antibodies, DNA was counterstained with DAPI. Scale bar= 2 μm. **B.** Quantifications of

A. *Left.* Prometaphase cells were counted for the presence or absence of GFP-Zw10 signal at

kinetochores. *Right.* Total GFP-Zw10 kinetochore intensity was measured in each cell,

normalized and plotted. Mean +/- SEM of 3 experiments (n>90 cells), Student t-test (*:

p<0.05; ***: p<0.001). **C.** Co-immunoprecipitation of RZZ components with centromeric

proteins. GFP (Ctrl), GFP-Rod, GFP-Zw10, GFP-Zwilch and GFP-RINT1 proteins were

immunoprecipitated using the GFP-binding protein (GBP). Pulled down fractions were

analyzed by immunoblotting for the presence of CAL1, CENP-A and Zw10. **D.** Yeast two

hybrid interaction tests using the RZZ components Zw10, Zwilch and three domains of ROD

with CAL1 and CENP-A. Blue color reflects the interaction between the 2 proteins tested.

Figure 3: Zw10 affects the loading of CENP-A during mitosis.

A. Time line of RNAi experiment followed by a Quench-Chase-Pulse SNAP-tag experiment.

After 72 h dsRNA treatment, cells were incubated with SNAP-Block to quench existing SNAP-

CENP-A molecules (Q), washed and cultured for 24 h. After 24 h chase, cells were incubated

with TMR-Star (red) to stain newly synthesized SNAP-CENP-A molecules (P). The images

show representative immunofluorescence staining of SNAP-CENP-A expressing cells after

Zw10 or control depletion. DNA was counterstained with DAPI. Scale bar= 2 μm. **B.**

Quantification of A. showing the total SNAP-CENP-A centromeric intensity per nucleus as %

of control. Mean +/- SEM of 3 experiments (n>300 cells), Student t-test (***: p<0.001). **C.**

FRAP of GFP-CENP-A in mitosis after Zw10 depletion. After 72 h dsRNA treatment, GFP-

CENP-A signal was partially (about 50-60 %) bleached in prophase and cells were imaged

until telophase. Time lapse: 90 s. Scale bar= 2 μm. **D.** Quantification of C. The total GFP-

CENP-A centromeric signal of the cells shown in C is displayed. **E.** Recovery rate of GFP-

CENP-A in control and Zw10-depleted cells. Mean +/- SEM, n≥10 cells, Student t-test (n.s=

non-significant).

Figure 4: CENP-A loading is correlated with mitosis duration.

A. Time-lapse imaging of H2B-GFP/mCherry-tubulin expressing cells after Zw10 depletion.

After 72 h dsRNA treatment, cells were imaged for 16 h. Time lapse 3 min. Scale bar= 2 μm.

B. Quantification of A. The mitosis duration from nuclear envelope breakdown (determined

by mCherry-Tubulin nuclear diffusion concomitant to DNA condensation) to anaphase entry

is shown. Mean +/- SEM, n>200 cells. Student t-test (***: p<0.001). **C.** SNAP-CENP-A

incorporation in BubR1, Mad2, Mis12, Spindly or Cdc27 depleted cells. After 72 h dsRNA

treatment, a Quench-Chase-Pulse experiment (scheme in Fig. 3A) was performed to stain

newly synthesized SNAP-CENP-A molecules. DNA was counterstained with DAPI. Scale bar: 2

μm. **D.** Quantification of C. The total SNAP-CENP-A centromeric signal intensity per nucleus

is shown as % of control dsRNA-treated cells. Mean +/- SEM of 3 experiments (n>300 cells),

Student t-test (*n.s.*: non-significant; ***: p<0.001). **E.** The recovery rate of GFP-CENP-A after

photobleaching in mitosis is plotted against the mitosis duration for each analyzed cell.

Pearson's correlation test, p<0.01.

Figure 5: CENP-A levels affects mitosis duration.

A. Immunoblot showing CENP-A knockdown efficiency in GFP-CENP-A/mCherry-Tubulin expressing cells. An anti-CENP-A antibody was used to detect endogenous CENP-A and overexpressed GFP-CENP-A. **B.** Time-lapse imaging of GFP-CENP-A/mCherry-tubulin expressing cells after CENP-A depletion. After 72 h dsRNA treatment, cells were imaged for 16 h. Time lapse 3 min. Scale bar= 5 μm. **C.** Quantification of B. showing the mitosis duration of control and CENP-A depleted cells. Mean +/- SEM, n>80 cells. Student t-test (***: p<0.001). **D.** Immunoblot showing CENP-A levels in the different cell lines used. CENP-A antibody detects endogenous and tagged versions of CENP-A. GFP-CENP-A and SNAP-CENP-A are under the constitutive *copia* promoter. CENP-A-GFP expression can be induced by activating the *metallothionein* promoter with $CuSO_4$. H3 serves as loading control. **E.** Immunofluorescence of CENP-A-GFP cells. CENP-A-GFP expression was induced with 10 μM $CuSO_4$ for 2 h. Cells were washed and cultured in conditional medium for 22 h before fixation. DNA was counterstained with DAPI. Scale bar= 2 μm. **F.** Quantification of E. showing the total CENP-A-GFP centromeric intensity per nucleus as % of control. Mean +/- SEM of 3 experiments (n>300 cells), Student t-test (n.s= non-significant). **G.** Time lapse imaging of cells expressing mCherry-tubulin and inducible CENP-A-GFP. GFP-CENP-A expression was induced by addition of 10 μM $CuSO_4$ to the medium for 2 h. Cells were washed before imaging (16 h). Time lapse 3 min. Scale bar= 5 μm. The intensity levels of CENP-A-GFP in control cells are shown enhanced for visualization purposes **H.** Quantification of G. showing mitosis duration of control and CENP-A-GFP expressing cells. Mean +/- SEM, n>300 cells. Student t-test (***: p<0.001). **I.** Mitotic phenotypes of CENP-A-GFP/mCherry-Tubulin cells imaged in G. Cells were scored for the accuracy of mitosis: normal or defective

(presence of lagging chromosomes in anaphase, formation of tripolar spindles). Mean +/-

SEM n > 200 cells. Student t-test (n.s.: non-significant).

Figure 6: CAL1 overexpression leads to increased CENP-A levels at centromeres.

A. Immunoblot of CENP-A (either GFP or SNAP N-terminally tagged) expressing cells with or

without concomitant CAL1 overexpression. CAL1-V5 expression was induced by addition of

100 µM $CuSO_4$ for 24 h. Anti-CAL1 (endogenous CAL1 and CAL1-V5), anti-CENP-A

(endogenous CENP-A and tagged-CENP-A) and anti-H3 antibodies were used. **B.**

Immunofluorescence of CAL1-V5/GFP-CENP-A expressing cells. CAL1-V5 expression was

induced as in A. Cells were fixed and stained with anti-V5 antibody (red). DNA was

counterstained with DAPI. **C.** Metaphase chromosomes of CAL1-V5 expressing cells. CAL1-V5

expression was induced as in A. Metaphase chromosomes were prepared and stained with

anti-CENP-A antibody. DNA was counterstained with DAPI. **D.** Quantification of B. GFP-

CENP-A signal intensity per centromere was measured and is shown as % of non-induced

control cells. **E.** Immunofluorescence of CAL1-V5/GFP-CENP-A expressing cells. CAL1-V5

expression was induced as in A. Then the cells were fixed and stained with anti-CENP-C

antibody (red). DNA was counterstained with DAPI. **F.** Quantification of E. CENP-C signal

intensity per centromere was measured and is shown as % of non-induced control cells. **G.**

Immunofluorescence of newly synthesized SNAP-CENP-A in CAL1 overexpressing cells. CAL1-

V5 expression was induced as in A. SNAP-CENP-A molecules were quenched with SNAP-

Block. A 24 h chase was performed in presence of $CuSO_4$, before labeling with SNAP-SiR647

and fixation. DNA was counterstained with DAPI. **H.** Quantification of G. SNAP-CENP-A signal

intensity per centromere was measured and is shown as % of non-induced control. All

graphs show Mean +/- SEM of 3 experiments (n>300 cells), Student t-test (n.s.: non-

significant; *: p<0.05; **: p<0.01). Scale bar= 2 μm.

Figure 7: Overexpression of CAL1 leads to CENP-A centromeric loading outside of mitosis.

A. Time-lapse imaging of H2B-GFP/mCherry Tubulin cells with or without concomitant

overexpression of CAL1. CAL1-V5 expression was induced for 24 h. The cells were imaged for

16 h. Time lapse 3 min. Scale bar= 2 μm. **B.** Quantification of A showing the mitosis duration.

Mean +/- SEM, n > 200 cells. Student t-test (***: p<0.001). **C.** FRAP experiments of GFP-

CENP-A in mitosis in CAL1-overexpressing cells. CAL1-V5 expression was induced for 24 h.

GFP-CENP-A signal was partially bleached in prophase and cells were imaged until

telophase. Time lapse: 90 s. Scale bar= 2 μm. **D.** Recovery rate of GFP-CENP-A in control and

CAL1-overexpressing cells. Mean +/- SEM, n≥ 12 cells, Student t-test (*: p<0.05; **: p<0.01).

E. Time-lapse imaging of SNAP-CENP-A/mCherry-Tubulin cells with or without concomitant

overexpression of CAL1. CAL1-V5 expression was induced for 24 h. Then the cells were

incubated with SNAP-Block to quench existing SNAP-CENP-A molecules, washed and

resuspended in conditional medium containing 0.5 μM SNAP-640 dye. Cells were imaged for

16 h to visualize newly synthesized SNAP-CENP-A. Time lapse 15 min. Scale bar= 2 μm.

Intensity levels have been adjusted separately for each condition. **F-G.** Quantifications of E.

F. Table showing the % of SNAP-CENP-A positive cells (+ cells) and the timing of loading. **G.**

For each cycling cell, the earliest detection time point of SNAP-CENP-A (Y-axis) is plotted

versus the time of cytokinesis (X-axis). t0 on both axis corresponds to start of imaging after

SNAP-Block.

Online supplemental materials include 6 figures and 10 videos. Fig. S1 shows localization of

the GFP-tagged RZZ-proteins during cell cycle, qPCR and WB results for RNAi efficiency. Fig.

S2 shows Spc105R localization in Zw10-depleted cells, Zw10 localization in CAL1-depleted

cells, localization of GFP-RINT during cell cycle and yeast two hybrid with CAL1 and Zw10

fragments. Fig. S3 shows SNAP-CAL1 quench-chase-pulse experiment in Zw10-depleted

cells, centromeric SNAP-CENP-A stability assay and GFP-CENP-A FRAP in early G1. Fig. S4

shows qPCR results for RNAi efficiency, SNAP-CENP-A and SNAP-CAL1 quench-chase-pulse

experiments in different RNAi and drug treatment. Fig. S5 shows SAC activity and

kinetochore components recruitment in CAL1-overexpressing cells. Fig. S6 shows SNAP-

CENP-A centromeric loading outside mitosis.

Supplemental figures legends

Figure S1 related to fig. 1.

Localization of GFP-ROD (**A.**), GFP-Zwilch (**B**.), GFP-Zw10 (**C**.) during cell cycle. Cells

expressing each GFP-RZZ component concomitant with mCherry-Tubulin were fixed and

stained with anti-GFP (green) and anti-CENP-A (red) antibodies, DNA was counterstained

with DAPI. Scale bar= 2 μm. **D**. qPCR results showing mRNA levels after indicated

knockdown in percent of control. **E.** Immunoblot showing protein levels after dsRNA

treatment.

Figure S2 related to fig. 1-2.

A. Immunofluorescence of S2 cells after depletion of Zw10. At day 4 of RNAi, cells were fixed

and stained with anti-Zw10 (green), anti-Spc105R (red) and anti-tubulin (grey) antibodies,

DNA was counterstained with DAPI. Scale bar= 2 μm. **B.** Quantification of A. Prometaphase cells were counted for the presence or absence of Spc105R signal at kinetochores. **C.** Immunofluorescence of GFP-Zw10/mCherry-Tubulin expressing cells after depletion of CAL1. After 96 h dsRNA treatment, cells were fixed and stained with anti-GFP (green) and anti-CENP-A (red) antibodies, DNA was counterstained with DAPI. Scale bar= 2 μm. **D.** Localization of GFP-RINT1, during cell cycle. Cells expressing each GFP-RINT1 concomitant with mCherry-Tubulin were fixed and stained with anti-GFP (green) and anti-CENP-A (red) antibodies, DNA was counterstained with DAPI. Scale bar= 2 μm. **F.** Yeast two hybrid interaction tests. Blue color reflects the interaction between the 2 proteins tested. **G.** Scheme showing CAL1 functional domains and its binding to its known partners. BD: binding domain.

Figure S3 related to fig. 3

A. Immunofluorescence of SNAP-CAL1 expressing cells after Zw10 depletion. After 72 h dsRNA treatment, a Quench-Chase-Pulse experiment (scheme in Fig. 3A) was performed to stain newly synthesized SNAP-CAL1 molecules. DNA was counterstained with DAPI. Scale bar= 2 μm. **B.** Quantification of A. showing the total SNAP-CAL1 centromeric intensity per nucleus as % of control. Mean +/- SEM of 3 experiments (n>300 cells), Student t-test (**: $p<0.01$). **C.** Immunofluorescence of SNAP-CENP-A expressing cells after Zw10 depletion. After 72 h dsRNA treatment, cells were incubated with TMR-Star (P) to stain existing SNAP-CENP-A molecules, washed and put back in culture (t0). After 24 h chase, cells were fixed (t24). Note that no SNAP-Block was performed for this experiment. DNA was counterstained with DAPI. Scale bar= 2 μm. **D.** Quantification of C. showing the total SNAP-CENP-A

centromeric intensity per nucleus as % of t0. Mean +/- SEM of 3 experiments (n>300 cells),

Student t-test (n.s= non-significant). **E.** FRAP of GFP-CENP-A in G1 phase. Cells expressing

GFP-CENP-A and mCherry-Tubulin were followed through mitosis, GFP-CENP-A signal was

partially bleached in early G1 and cells were further imaged for > 2 h. Time lapse: 6 min.

Scale bar: 2 μm. **F.** Quantification of E. The total GFP-CENP-A centromeric signal of 4

representative cells is shown.

Figure S4 related to fig. 4

A. qPCR results showing mRNA levels after indicated knockdown in SNAP-CENP-A cells as

percent of control. **B.** Immunofluorescence of SNAP-CENP-A expressing cells after MG132 or

Nocodazole treatment. A Quench-Chase-Pulse experiment was performed to stain newly

synthesized SNAP-CENP-A molecules. The 24 h chase was performed in presence or absence

of the indicated drug. Cells were stained with an anti-S10-phospho-H3 antibody to identify

mitotic cells. DNA was counterstained with DAPI. Scale bar= 2 μm. **C.** Quantification of B.

showing the SNAP-CENP-A centromeric intensity as % of control. Only Phospho-H3 positive

cells were analyzed. Mean+/- SEM of 3 experiments (n>300 cells), Student t-test (n.s= non-

significant). **D.** Quantification showing the total SNAP-CAL1 centromeric intensity per

nucleus as % of control after depletion of the indicated proteins. Mean+/- SEM of 3

experiments (n>300 cells), Student t-test (**: $p<0.01$; n.s.: non-significant).

Figure S5 related to fig. 7

A. Mitotic phenotypes of H2B-GFP/mCherry-Tubulin expressing cells with or without

concomitant overexpression of CAL1. CAL1-V5 expression was induced for 24 h with 100 μM

CuSO$_4$ then the cells were imaged for 16 h. Cells were scored for the accuracy of mitosis:

normal or defective (presence of lagging chromosomes in anaphase, formation of tripolar

spindles). Mean +/- SEM n > 200 cells. Student t-test (n.s.: non-significant). **B.** Amount of

kinetochore proteins recruited during mitosis in presence or absence of CAL1

overexpression. CAL1-V5 expression was induced for 24 h in GFP-CENP-A/mCherry-Tubulin

cells. Cells were then fixed and stained with anti-CENP-C (upper panel, red), anti-Spc105R

(middle panel, red) or anti-BubR1 antibody (lower panel, red). DNA was counterstained with

DAPI. Scale bar= 2 μm. Kinetochore signal intensity of the indicated proteins is shown as %

of control. Only prometaphase cells were analyzed. Mean +/- SEM of 3 experiments (n > 300

cells). Student t-test (n.s.: non-significant). **C.** Stills from time-lapse imaging experiments

showing GFP-CENP-A/mCherry-Tubulin cells with or without CAL1 overexpression imaged

immediately after addition of 1 μM Taxol and after 5 h. CAL1-V5 expression was induced for

24 h. Then 1 μM taxol was added and the cells were imaged for 16 h. **D.** Quantification of C.

showing the percentage of cells that arrest (arrest) in response to Taxol treatment or arrest

and restart (finish).

Figure S6 related to fig. 7

A. Quantification of cells showing SNAP-CENP-A signal at centromeres in control and CAL1-

overexpressing cells after 4 h MG132 treatment. CAL1-V5 expression was induced for 24 h.

then the cells were incubated for 2 h with MG132 to arrest cells in mitosis prior to SNAP-

CENP-A block. A 4-h chase was performed in presence of MG132 to allow synthesis and

incorporation of new SNAP-CENP-A into centromeres before incubation with SNAP-Si647

and fixation. **B.** Immunofluorescence of SNAP-CENP-A in control and CAL1 overexpressing

cells. CAL1-V5 expression was induced for 24 h. The cells were then incubated with SNAP-

Block, washed and resuspended in conditional medium containing SNAP-640 dye. After 20

h, the cells were washed 3 times before fixation and immunostaining with an anti-CENP-A

antibody (green). Cells were also taken after block, stained with 0.5 µM SNAP-640 dye for 15

min to check the efficiency of the block. DNA was counterstained with DAPI. Scale bar= 2

µm. **C.** Quantification of B. showing the percentage of cells positive for centromeric SNAP-

CENP-A staining. **D.** Quantification of B. showing the total SNAP-CENP-A centromeric

intensity per nucleus as % of control. All graphs show Mean +/- SEM of 3 experiments

(n>300 cells), Student t-test (n.s.: non-significant; *: $p < 0.05$; **: $p < 0.01$).

Supplemental Videos.

Video 1 (related to Figure 4A).

H2B-GFP/mCherry-tubulin expressing cells after Ctrl dsRNA treatment (72 h). Imaging was

performed for 16 h on a Deltavision microscope (60X, Bin 2), at 25 °C. Time lapse 3 min.

Scale bar= 5 µm.

Video 2 (related to Figure 4A):

H2B-GFP/mCherry-tubulin expressing cells after Zw10 dsRNA treatment (72 h). Imaging was

performed for 16 h on a Deltavision microscope (60X, Bin 2), at 25 °C. Time lapse 3 min.

Scale bar= 5 µm.

Video 3 (related to Figure 5B):

GFP-CENP-A/mCherry-Tubulin expressing cells after Ctrl dsRNA treatment (72 h). Imaging

was performed for 16 h on a Deltavision microscope (40X, Bin 2), at 25 °C. Time lapse 3 min.

Scale bar= 5 µm.

Video 4 (related to Figure 5B):

GFP-CENP-A/mCherry-Tubulin expressing cells after CENP-A dsRNA treatment (72 h).

Imaging was performed for 16 h on a Deltavision microscope (40X, Bin 2), at 25 °C. Time

lapse 3 min. Scale bar= 5 µm.

Video 5 (related to Figure 5G):

CENP-A-GFP/mCherry-tubulin expressing cells. Visible CENP-A-GFP results from leaky

metallothionein promoter. Imaging was performed for 16 h on a Deltavision microscope

(40X, Bin 2), at 25 °C. Time lapse 3 min. Scale bar= 5 µm.

Video 6 (related to Figure 5G):

CENP-A-GFP/mCherry-tubulin expressing cells. CENP-A-GFP expression was induced for 2 h

with 10 µM $CuSO_4$ then washed 3 times before being placed in Ibidi chamber. Imaging was

performed for 16 h on a Deltavision microscope (40X, Bin 2), at 25 °C. Time lapse 3 min.

Scale bar= 5 µm.

Video 7 (related to Figure 7A):

H2B-GFP/mCherry-tubulin expressing cells containing inducible CAL1-V5. No CAL1-V5

induction. Imaging was performed for 16 h on a Deltavision microscope (60X, Bin 2), at 25

°C. Time lapse 3 min. Scale bar= 5 µm.

Video 8 (related to Figure 7A):

H2B-GFP/mCherry-tubulin expressing cells containing inducible CAL1-V5. CAL1-V5

expression was induced for 24 h prior to imaging with 100 µM $CuSO_4$. Imaging was

performed for 16 h on a Deltavision microscope (60X, Bin 2), at 25 °C. Time lapse 3 min.

Scale bar= 5 µm.

Video 9 (related to Figure 7E):

SNAP-CENP-A/mCherry-tubulin expressing cells containing inducible CAL1-V5. No CAL1-V5

induction. SNAP-CENP-A molecules were quenched before addition of 0.5 µM SNAP-640

dye. Imaging was performed for 16 h on a Deltavision microscope (60X, Bin 2), at 25 °C.

Time lapse 15 min. Scale bar= 5 µm.

Video 10 (related Figure 7E):

SNAP-CENP-A/mCherry-tubulin expressing cells containing inducible CAL1-V5. CAL1-V5

expression was induced for 24 h with 100 µM $CuSO_4$. SNAP-CENP-A molecules were

quenched before addition of 0.5 µM SNAP-640 dye. Imaging was performed for 16 h on a

Deltavision microscope (60X, Bin 2), at 25 °C. Time lapse 15 min. Scale bar= 5 µm.

References

Ahmad, K., and S. Henikoff. 2001. Centromeres are specialized replication domains in
heterochromatin. *J Cell Biol*. 153:101-110.
Bade, D., A.L. Pauleau, A. Wendler, and S. Erhardt. 2014. The E3 ligase CUL3/RDX controls
centromere maintenance by ubiquitylating and stabilizing CENP-A in a CAL1-dependent manner. *Dev*
Cell. 28:508-519.
Barisic, M., and S. Geley. 2011. Spindly switch controls anaphase: spindly and RZZ functions in
chromosome attachment and mitotic checkpoint control. *Cell Cycle*. 10:449-456.
Basto, R., R. Gomes, and R.E. Karess. 2000. Rough deal and Zw10 are required for the metaphase
checkpoint in Drosophila. *Nat Cell Biol*. 2:939-943.
Basto, R., F. Scaerou, S. Mische, E. Wojcik, C. Lefebvre, R. Gomes, T. Hays, and R. Karess. 2004. In
vivo dynamics of the rough deal checkpoint protein during Drosophila mitosis. *Curr Biol*. 14:56-61.
Basu, J., E. Logarinho, S. Herrmann, H. Bousbaa, Z. Li, G.K. Chan, T.J. Yen, C.E. Sunkel, and M.L.
Goldberg. 1998. Localization of the Drosophila checkpoint control protein Bub3 to the kinetochore
requires Bub1 but not Zw10 or Rod. *Chromosoma*. 107:376-385.
Blower, M.D., T. Daigle, T. Kaufman, and G.H. Karpen. 2006. Drosophila CENP-A mutations cause a
BubR1-dependent early mitotic delay without normal localization of kinetochore components. *PLoS*
Genet. 2:e110.
Blower, M.D., and G.H. Karpen. 2001. The role of Drosophila CID in kinetochore formation, cell-cycle
progression and heterochromatin interactions. *Nat Cell Biol*. 3:730-739.

Buffin, E., D. Emre, and R.E. Karess. 2007. Flies without a spindle checkpoint. *Nat Cell Biol*. 9:565-572.

Buffin, E., C. Lefebvre, J. Huang, M.E. Gagou, and R.E. Karess. 2005. Recruitment of Mad2 to the
kinetochore requires the Rod/Zw10 complex. *Curr Biol*. 15:856-861.

Butkevich, A.N., V.N. Belov, K. Kolmakov, V.V. Sokolov, H. Shojaei, S.C. Sidenstein, D. Kamin, J.
Matthias, R. Vlijm, J. Engelhardt, and S.W. Hell. 2017. Hydroxylated Fluorescent Dyes for Live-Cell
Labeling: Synthesis, Spectra and Super-Resolution STED. *Chemistry*. 23:12114-12119.

Caldas, G.V., T.R. Lynch, R. Anderson, S. Afreen, D. Varma, and J.G. DeLuca. 2015. The RZZ complex
requires the N-terminus of KNL1 to mediate optimal Mad1 kinetochore localization in human cells.
Open Biol. 5.

Chen, C.C., M.L. Dechassa, E. Bettini, M.B. Ledoux, C. Belisario, P. Heun, K. Luger, and B.G. Mellone.
2014. CAL1 is the Drosophila CENP-A assembly factor. *J Cell Biol*. 204:313-329.

Defachelles, L., S.G. Hainline, A. Menant, L.A. Lee, and R.E. Karess. 2015a. A maternal effect rough
deal mutation suggests that multiple pathways regulate Drosophila RZZ kinetochore recruitment. *J
Cell Sci*. 128:2952.

Defachelles, L., N. Raich, R. Terracol, X. Baudin, B. Williams, M. Goldberg, and R.E. Karess. 2015b. RZZ
and Mad1 dynamics in Drosophila mitosis. *Chromosome Res*. 23:333-342.

Emre, D., R. Terracol, A. Poncet, Z. Rahmani, and R.E. Karess. 2011. A mitotic role for Mad1 beyond
the spindle checkpoint. *J Cell Sci*. 124:1664-1671.

Erhardt, S., B.G. Mellone, C.M. Betts, W. Zhang, G.H. Karpen, and A.F. Straight. 2008. Genome-wide
analysis reveals a cell cycle-dependent mechanism controlling centromere propagation. *J Cell Biol*.
183:805-818.

Feijao, T., O. Afonso, A.F. Maia, and C.E. Sunkel. 2013. Stability of kinetochore-microtubule
attachment and the role of different KMN network components in Drosophila. *Cytoskeleton
(Hoboken)*. 70:661-675.

Gassmann, R., A. Essex, J.S. Hu, P.S. Maddox, F. Motegi, A. Sugimoto, S.M. O'Rourke, B. Bowerman, I.
McLeod, J.R. Yates, 3rd, K. Oegema, I.M. Cheeseman, and A. Desai. 2008. A new mechanism
controlling kinetochore-microtubule interactions revealed by comparison of two dynein-targeting
components: SPDL-1 and the Rod/Zwilch/Zw10 complex. *Genes Dev*. 22:2385-2399.

Griffis, E.R., N. Stuurman, and R.D. Vale. 2007. Spindly, a novel protein essential for silencing the
spindle assembly checkpoint, recruits dynein to the kinetochore. *J Cell Biol*. 177:1005-1015.

Heun, P., S. Erhardt, M.D. Blower, S. Weiss, A.D. Skora, and G.H. Karpen. 2006. Mislocalization of the
Drosophila centromere-specific histone CID promotes formation of functional ectopic kinetochores.
Dev Cell. 10:303-315.

Huang, J.Y., and J.W. Raff. 2002. The dynamic localisation of the Drosophila APC/C: evidence for the
existence of multiple complexes that perform distinct functions and are differentially localised. *J Cell
Sci*. 115:2847-2856.

Jansen, L.E., B.E. Black, D.R. Foltz, and D.W. Cleveland. 2007. Propagation of centromeric chromatin
requires exit from mitosis. *J Cell Biol*. 176:795-805.

Joglekar, A.P., and A.A. Kukreja. 2017. How Kinetochore Architecture Shapes the Mechanisms of Its
Function. *Curr Biol*. 27:R816-R824.

Kapanidou, M., N.L. Curtis, and V.M. Bolanos-Garcia. 2017. Cdc20: At the Crossroads between
Chromosome Segregation and Mitotic Exit. *Trends Biochem Sci*. 42:193-205.

Karess, R. 2005. Rod-Zw10-Zwilch: a key player in the spindle checkpoint. *Trends Cell Biol*. 15:386-
392.

Karess, R.E., and D.M. Glover. 1989. rough deal: a gene required for proper mitotic segregation in
Drosophila. *J Cell Biol*. 109:2951-2961.

Kasuboski, J.M., J.R. Bader, P.S. Vaughan, S.B. Tauhata, M. Winding, M.A. Morrissey, M.V. Joyce, W.
Boggess, L. Vos, G.K. Chan, E.H. Hinchcliffe, and K.T. Vaughan. 2011. Zwint-1 is a novel Aurora B
substrate required for the assembly of a dynein-binding platform on kinetochores. *Mol Biol Cell*.
22:3318-3330.

Kops, G.J., Y. Kim, B.A. Weaver, Y. Mao, I. McLeod, J.R. Yates, 3rd, M. Tagaya, and D.W. Cleveland. 2005. ZW10 links mitotic checkpoint signaling to the structural kinetochore. *J Cell Biol*. 169:49-60.

Li, J., N. Dang, D.J. Wood, and J.Y. Huang. 2017. The kinetochore-dependent and -independent formation of the CDC20-MAD2 complex and its functions in HeLa cells. *Sci Rep*. 7:41072.

Lidsky, P.V., F. Sprenger, and C.F. Lehner. 2013. Distinct modes of centromere protein dynamics during cell cycle progression in Drosophila S2R+ cells. *J Cell Sci*. 126:4782-4793.

Lince-Faria, M., S. Maffini, B. Orr, Y. Ding, F. Claudia, C.E. Sunkel, A. Tavares, J. Johansen, K.M. Johansen, and H. Maiato. 2009. Spatiotemporal control of mitosis by the conserved spindle matrix protein Megator. *J Cell Biol*. 184:647-657.

London, N., and S. Biggins. 2014. Mad1 kinetochore recruitment by Mps1-mediated phosphorylation of Bub1 signals the spindle checkpoint. *Genes Dev*. 28:140-152.

Mathew, V., A.L. Pauleau, N. Steffen, A. Bergner, P.B. Becker, and S. Erhardt. 2014. The histone-fold protein CHRAC14 influences chromatin composition in response to DNA damage. *Cell Rep*. 7:321-330.

Mellone, B.G., K.J. Grive, V. Shteyn, S.R. Bowers, I. Oderberg, and G.H. Karpen. 2011. Assembly of Drosophila centromeric chromatin proteins during mitosis. *PLoS Genet*. 7:e1002068.

Meraldi, P., V.M. Draviam, and P.K. Sorger. 2004. Timing and checkpoints in the regulation of mitotic progression. *Dev Cell*. 7:45-60.

Moreno-Moreno, O., M. Torras-Llort, and F. Azorin. 2006. Proteolysis restricts localization of CID, the centromere-specific histone H3 variant of Drosophila, to centromeres. *Nucleic Acids Res*. 34:6247-6255.

Moyle, M.W., T. Kim, N. Hattersley, J. Espeut, D.K. Cheerambathur, K. Oegema, and A. Desai. 2014. A Bub1-Mad1 interaction targets the Mad1-Mad2 complex to unattached kinetochores to initiate the spindle checkpoint. *J Cell Biol*. 204:647-657.

Musacchio, A. 2015. The Molecular Biology of Spindle Assembly Checkpoint Signaling Dynamics. *Curr Biol*. 25:R1002-1018.

Musacchio, A., and A. Desai. 2017. A Molecular View of Kinetochore Assembly and Function. *Biology (Basel)*. 6.

Orr, B., H. Bousbaa, and C.E. Sunkel. 2007. Mad2-independent spindle assembly checkpoint activation and controlled metaphase-anaphase transition in Drosophila S2 cells. *Mol Biol Cell*. 18:850-863.

Orr, B., and C.E. Sunkel. 2010. Drosophila CENP-C is essential for centromere identity. *Chromosoma*. 120:83-96.

Petsalaki, E., M. Dandoulaki, and G. Zachos. 2018. The ESCRT protein Chmp4c regulates mitotic spindle checkpoint signaling. *J Cell Biol*. 217:861-876.

Przewloka, M.R., Z. Venkei, V.M. Bolanos-Garcia, J. Debski, M. Dadlez, and D.M. Glover. 2011. CENP-C Is a Structural Platform for Kinetochore Assembly. *Curr Biol*. 21:399-405.

Przewloka, M.R., W. Zhang, P. Costa, V. Archambault, P.P. D'Avino, K.S. Lilley, E.D. Laue, A.D. McAinsh, and D.M. Glover. 2007. Molecular analysis of core kinetochore composition and assembly in Drosophila melanogaster. *PLoS One*. 2:e478.

Rahmani, Z., M.E. Gagou, C. Lefebvre, D. Emre, and R.E. Karess. 2009. Separating the spindle, checkpoint, and timer functions of BubR1. *J Cell Biol*. 187:597-605.

Rothbauer, U., K. Zolghadr, S. Muyldermans, A. Schepers, M.C. Cardoso, and H. Leonhardt. 2008. A versatile nanotrap for biochemical and functional studies with fluorescent fusion proteins. *Mol Cell Proteomics*. 7:282-289.

Sacristan, C., M.U.D. Ahmad, J. Keller, J. Fermie, V. Groenewold, E. Tromer, A. Fish, R. Melero, J.M. Carazo, J. Klumperman, A. Musacchio, A. Perrakis, and G.J. Kops. 2018. Dynamic kinetochore size regulation promotes microtubule capture and chromosome biorientation in mitosis. *Nat Cell Biol*. 20:800-810.

Scaerou, F., I. Aguilera, R. Saunders, N. Kane, L. Blottiere, and R. Karess. 1999. The rough deal protein
is a new kinetochore component required for accurate chromosome segregation in Drosophila. *J Cell*
Sci. 112 (Pt 21):3757-3768.
Scaerou, F., D.A. Starr, F. Piano, O. Papoulas, R.E. Karess, and M.L. Goldberg. 2001. The ZW10 and
Rough Deal checkpoint proteins function together in a large, evolutionarily conserved complex
targeted to the kinetochore. *J Cell Sci*. 114:3103-3114.
Schittenhelm, R.B., F. Althoff, S. Heidmann, and C.F. Lehner. 2010. Detrimental incorporation of
excess Cenp-A/Cid and Cenp-C into Drosophila centromeres is prevented by limiting amounts of the
bridging factor Cal1. *J Cell Sci*. 123:3768-3779.
Schittenhelm, R.B., R. Chaleckis, and C.F. Lehner. 2009. Intrakinetochore localization and essential
functional domains of Drosophila Spc105. *EMBO J*. 28:2374-2386.
Schittenhelm, R.B., S. Heeger, F. Althoff, A. Walter, S. Heidmann, K. Mechtler, and C.F. Lehner. 2007.
Spatial organization of a ubiquitous eukaryotic kinetochore protein network in Drosophila
chromosomes. *Chromosoma*. 116:385-402.
Schmitt, H.D. 2010. Dsl1p/Zw10: common mechanisms behind tethering vesicles and microtubules.
Trends Cell Biol. 20:257-268.
Schramm, C., S. Elliott, A. Shevchenko, and E. Schiebel. 2000. The Bbp1p-Mps2p complex connects
the SPB to the nuclear envelope and is essential for SPB duplication. *EMBO J*. 19:421-433.
Silio, V., A.D. McAinsh, and J.B. Millar. 2015. KNL1-Bubs and RZZ Provide Two Separable Pathways for
Checkpoint Activation at Human Kinetochores. *Dev Cell*. 35:600-613.
Starr, D.A., R. Saffery, Z. Li, A.E. Simpson, K.H. Choo, T.J. Yen, and M.L. Goldberg. 2000. HZwint-1, a
novel human kinetochore component that interacts with HZW10. *J Cell Sci*. 113 (Pt 11):1939-1950.
Starr, D.A., B.C. Williams, T.S. Hays, and M.L. Goldberg. 1998. ZW10 helps recruit dynactin and
dynein to the kinetochore. *J Cell Biol*. 142:763-774.
Varma, D., X. Wan, D. Cheerambathur, R. Gassmann, A. Suzuki, J. Lawrimore, A. Desai, and E.D.
Salmon. 2013. Spindle assembly checkpoint proteins are positioned close to core microtubule
attachment sites at kinetochores. *J Cell Biol*. 202:735-746.
Vleugel, M., E. Hoogendoorn, B. Snel, and G.J. Kops. 2012. Evolution and function of the mitotic
checkpoint. *Dev Cell*. 23:239-250.
Wainman, A., M.G. Giansanti, M.L. Goldberg, and M. Gatti. 2012. The Drosophila RZZ complex - roles
in membrane trafficking and cytokinesis. *J Cell Sci*. 125:4014-4025.
Westermann, S., and A. Schleiffer. 2013. Family matters: structural and functional conservation of
centromere-associated proteins from yeast to humans. *Trends Cell Biol*. 23:260-269.
Williams, B.C., and M.L. Goldberg. 1994. Determinants of Drosophila zw10 protein localization and
function. *J Cell Sci*. 107 (Pt 4):785-798.
Williams, B.C., T.L. Karr, J.M. Montgomery, and M.L. Goldberg. 1992. The Drosophila l(1)zw10 gene
product, required for accurate mitotic chromosome segregation, is redistributed at anaphase onset.
J Cell Biol. 118:759-773.
Williams, B.C., Z. Li, S. Liu, E.V. Williams, G. Leung, T.J. Yen, and M.L. Goldberg. 2003. Zwilch, a new
component of the ZW10/ROD complex required for kinetochore functions. *Mol Biol Cell*. 14:1379-
1391.
Wojcik, E., R. Basto, M. Serr, F. Scaerou, R. Karess, and T. Hays. 2001. Kinetochore dynein: its
dynamics and role in the transport of the Rough deal checkpoint protein. *Nat Cell Biol*. 3:1001-1007.
Zhang, G., T. Lischetti, D.G. Hayward, and J. Nilsson. 2015. Distinct domains in Bub1 localize RZZ and
BubR1 to kinetochores to regulate the checkpoint. *Nat Commun*. 6:7162.

1034

Fig.1

A

B

C

D

Fig.2

A

B

SNAP-CENP-A centromeric signal (as % of Ctrl)

***

dsRNA: Ctrl, Zw10

C

GFP-CENP-A / mCherry-Tubulin

D

GFP-CENP-A centromeric signal vs Time (Min)

Ctrl dsRNA / Zw10 dsRNA

E

Recovery rate of GFP-CENP-A (a.u.)

n.s.

n=10, n=8

dsRNA: Ctrl, Zw10

Fig.3

Fig.4

Fig.5

Fig.6

A

Prophase Prophase (NEB) Metaphase Anaphase

Ctrl

CAL1 Ind.

GFP-H2B / mCherry-Tubulin

B

n=228   n= 270

CAL1 induction

C

Pre-bleach Bleach Prometaphase Metaphase Anaphase Telophase

Ctrl

CAL1 Ind.

GFP-CENP-A / mCherry-Tubulin

D

n=16   n=21   n=12

CAL1 induction

E

Ctrl

CAL1 Ind.

SNAP-CENP-A / mCherry-Tubulin

G

No CAL1 induction

24h CAL1 induction

F

	Ctrl		CAL1 induction	
SNAP-CENP-A + cells	38 %	(167/452)	91 %	(371/458)
SNAP-CENP-A signal:				
Before Mitosis	0		31 %	(114/371)
During mitosis	91 %	(152/167)	18 %	(67/371)
After cytokinesis	9 %	(15/167)	51 %	(191/371)

Fig.7

A

GFP-ROD / CENP-A / Tubulin / DAPI

B

GFP-Zwilch / CENP-A / Tubulin / DAPI

C

GFP-Zw10 / CENP-A / Tubulin / DAPI

D

E

Fig.S1

GFP-RINT1 / CENP-A / Tubulin / DAPI

Fig.S2

Fig.S3

Fig.S4

Fig.S5

A

B

SNAP-CENP-A   CENP-A   SNAP-CENP-A / CENP-A
DAPI

SNAP-CENP-A   CENP-A   SNAP-CENP-A / CENP-A
DAPI

Fig.S6