Whole brain imaging reveals distinct spatial patterns of amyloid beta deposition in three mouse
models of Alzheimer's disease

Running Title: Whole brain imaging in mouse models of Alzheimer's disease

Jennifer D. Whitesell[1], Alex R. Buckley[2], Joseph E. Knox[1], Leonard Kuan[1], Nile Graddis[1], Andrew
Pelos[1,3], Alice Mukora[1], Wayne Wakeman[1], Phillip Bohn[1], Anh Ho[1], Karla E. Hirokawa[1], Julie A. Harris[1*]
[1]Allen Institute for Brain Science, Seattle WA. [2]Washington University in St. Louis. [3]Pomona College,
Claremont, CA

*Correspondence should be addressed to J.A.H. (julieha@alleninstitute.org)

Acknowledgments

We thank Dr. Stefan Mihalas for helpful discussion and advice on statistical methods, Josh Royall for
graphic design work, Dr. Song-Lin Ding for assistance correlating human and mouse anatomical regions,
and the Animal Care, Transgenic Colony Management, Lab Animal Services, and imaging teams for
mouse husbandry, tissue preparation, and imaging. Tg2576 mice were a generous gift from Dr. Xavier
Figueroa at Nativis. Research reported in this publication was supported by the National Institute on
Aging of the National Institutes of Health under Award Number R01AG047589 to J.A.H. The content is
solely the responsibility of the authors and does not necessarily represent the official views of the
National Institutes of Health. We thank the Allen Institute founder, Paul G. Allen, for his vision,
encouragement, and support.

Abstract (250 words)

A variety of Alzheimer's disease (AD) mouse models overexpress mutant forms of human amyloid
precursor protein (APP), producing high levels of Aβ and forming plaques. However, the degree to which
these models mimic spatiotemporal patterns of Aβ deposition in brains of AD patients is unknown. Here,
we mapped the spatial distribution of Aβ plaques across ages in three APP-overexpression mouse lines
(APP/PS1, Tg2576, hAPP-J20) using *in vivo* labeling with methoxy-X04, high throughput whole brain
imaging, and an automated informatics pipeline. Images were acquired with high resolution serial 2-
photon tomography and labeled plaques were detected using custom-built segmentation algorithms.
Image series were registered to the Allen Mouse Brain Common Coordinate Framework, a 3D reference
atlas, enabling automated brain-wide quantification of plaque density, number, and location. In both
APP/PS1 and Tg2576 mice, plaques were identified first in isocortex, followed by olfactory, hippocampal,
and cortical subplate areas. In hAPP-J20 mice, plaque density was highest in hippocampal areas,
followed by isocortex, with little to no involvement of olfactory or cortical subplate areas. Within the major
brain divisions, distinct regions were identified with high (or low) plaque accumulation; *e.g.*, the lateral
visual area within the isocortex of APP/PS1 mice had relatively higher plaque density compared with
other cortical areas, while in hAPP-J20 mice, plaques were densest in the ventral retrosplenial cortex. In
summary, we show how whole brain imaging of amyloid pathology in mice reveals the extent to which a
given model recapitulates the regional Aβ deposition patterns described in AD.

Keywords

Alzheimer's mouse model, Amyloid beta, plaque deposition, whole brain imaging

1. Introduction

Alzheimer's disease (AD) is classically defined after death by the presence of two neuropathologies, amyloid β (Aβ) plaques and tau tangles. Recently, widespread advances in the use of biomarkers as rigorous measures of these pathologies in living people has led to a coordinated proposal for new definitions and staging of AD, incorporating biomarker profiles specific for Aβ and tau (Jack et al., 2018). One major benefit of using biomarkers to define disease stages is the ability to then design experiments testing novel therapeutics with the goal of intervening at early, presymptomatic, stages. This will likely accelerate translational efforts, but does not replace the need for animal models early in the discovery and testing process. Thus, an important goal for the preclinical research field is to systematically characterize whether existing and novel mouse models adequately mimic the distribution and progression of these pathologies as mapped in human patients.

Pathological alterations in Aβ are currently the earliest detectable biomarker change, occurring before changes in tau pathologies, and sometimes decades before clinical symptoms (Jack et al., 2013). Most commonly used mouse models were engineered to express mutant forms of the amyloid precursor protein (APP), and/or presenilin 1 (PS1) which cause early onset forms of AD. These mutant APP mouse lines develop amyloid pathology, but little to no tau pathology or frank neurodegeneration, suggesting that they might best model early stages of AD (Sasaguri et al., 2017). Although mouse models have received much of the blame for the lack of success in translating preclinical research findings to approved therapeutics for clinical use, one possibility may be that these failures are due to misalignments in the disease stage modeled by these mice compared to the stage at which interventions are made in the clinical trials.

The large variety, accessibility, and sheer number of Alzheimer's disease mouse models is an immensely important resource, revealing many novel basic science insights into Aβ pathologies. However, characterization of these lines with respect to molecular, circuit, network, and cognitive alterations is still very much incomplete, although at least one good resource based on post-hoc compilation of results exists (e.g., https://www.alzforum.org/research-models). As these data come from many independent labs using only a single line, often with different experimental focus on selected brain areas, and different methods and techniques, it is difficult to compare results across lines or interpret reported differences. Systematic characterization of mouse models, and a centralized database of results, would be a large asset for the AD community, assisting researchers in selecting the most appropriate lines based on experimental needs. New large-scale collaborative efforts promise to make progress toward this goal, at least for newly developed mouse models (e.g., MODEL-AD, https://model-ad.org/).

Here, our goal was to demonstrate the use of systematic whole brain imaging to characterize key pathological features in multiple mouse lines. We modified a high-throughput imaging and informatics pipeline, first developed for our Allen Mouse Connectivity Atlas project (Oh et al., 2014; http://connectivity.brain-map.org/projection), to label and map regional progression patterns of Aβ plaques across the entire brain in three frequently used APP mouse models: APP/PS1 (Jankowsky et al., 2001), hAPP-J20 (Mucke et al., 2000) and Tg2576 (Hsiao et al., 1996). Plaques were labeled across the entire brain via systemic injections of Methoxy-X04, a fluorescent Congo red derivative that crosses the blood-brain barrier (Klunk et al., 2002). Previous reports of plaque density in these AD mouse models report different ages of onset (Hall & Roberson, 2012; J.-E. Lee & Han, 2013), with numerous reports of plaque density in cortex and hippocampus (Garcia-Alloza et al., 2006; H. Huang et al., 2016; Jährling et al., 2015; E. B. Lee et al., 2006; Liu et al., 2017; Mucke et al., 2000; Samaroo et al., 2012; Wright et al., 2013; Zhang et al., 2017). Some recent studies have reported plaque density for large subdivisions the cortex (Kim et al., 2012) or a subset of structures (Liebmann et al., 2016) but they do not comprehensively describe plaque loads within subregions of these major brain divisions or across the rest of the brain.

In humans, progression and regional patterns of amyloid pathology have been described based on autopsy cases (Thal, Rüb, Orantes, & Braak, 2002), but also more increasingly with amyloid PET imaging (Buckner, 2005; Grothe et al., 2017; Rice & Bisdas, 2017). From these studies, we know that Aβ deposition occurs selectively first in the cortex, followed by hippocampal regions, including entorhinal and CA1, then other subcortical regions; e.g., striatum, basal forebrain, thalamus, and finally brainstem nuclei

and cerebellum. Within the cortex, Aβ aggregates appear first, and are heaviest, in associational cortical
areas, and specifically in the default mode network (DMN), which includes the precuneus, posterior
cingulate cortex (PCC), retrosplenial cortex (RSP), medial prefrontal cortex, and lateral posterior parietal
cortex (Buckner, Andrews-Hanna, & Schacter, 2008; Raichle et al., 2001).

Our results show plaque load is densest and appears earliest in the isocortex in both APP/PS1 and
Tg2576 mice, like the early amyloid phases described by Thal et al., (2002). In contrast, plaque density
was highest in hippocampal areas first, followed by isocortex in hAPP-J20 mice. We also identified
plaques in select subcortical structures, mostly in the APP/PS1 line, in areas homologous to those
described in the later amyloid deposition phases. Within the isocortex, the hAPP-J20 mice appeared to
more closely mimic early stage human AD regional amyloid deposition; plaque load was higher in
associational cortical areas as opposed to sensory and motor regions. Thus, systematic whole brain
imaging of amyloid pathology in mice reveals line-specific regional deposition patterns. These data can
be used together with characterization of other pathologies to identify the most suitable mouse models for
testing early interventions in the progression of Alzheimer's disease.

2. Materials and Methods

Animals. All experimental procedures related to the use of mice were approved by the Institutional Animal
Care and Use Committee of the Allen Institute for Brain Science, in accordance with NIH guidelines. We
used heterozygous APP$^{+/-}$ mice from the following transgenic lines: ***APP/PS1*** (B6.Cg-
Tg(APPswe,PSEN1dE9)85Dbo/Mmjax, MMRRC Stock No: 34832-JAX) (Jankowsky et al., 2001)***, hAPP-***
J20 (B6.Cg-Zbtb20Tg(PDGFB-APPSwInd)20Lms/2Mmjax, MMRRC Stock No: 34836-JAX) (Mucke et al.,
2000), ***Tg2576*** (B6;SJL-Tg(APPSWE)2576Kha) (Hsiao et al., 1996). All animals were group-housed with
a 10/14 light cycle (lights on from 6 AM to 8 PM, temperature = 68-72 degrees, humidity = 30-70%).
APP/PS1 and hAPP-J20 mice were on the C57Bl/6J background and Tg2576 mice were on an FVB
background. Mice were separated into six groups by age at perfusion: 5 months (P141-P156), 7 months
(P202-P218), 9 months (P263-P307), 13 months (P386-P423), 19 months (P529-P589). The number of
mice from each sex in each age group/transgenic line combination is listed in **Table 1**. We only very minor
differences between the sexes in one region that had very low plaque density (the thalamus in hAPP-J20
mice), so we pooled male and female brains for all analysis (however, the two sexes were not equally
distributed in our dataset, see **Table 1**). Our control dataset contained 35 APP$^{-/-}$ littermates from 6 - 18
months old from the APP/PS1 and hAPP-J20 lines (15 APP/PS1, 20 hAPP-J20; details in **Table 1**). All
mice used in this study received a stereotaxic injection of AAV2/1.pCAG.FLEX.EGFP in the left
hemisphere 20-25 days before perfusion; analyses of these data are not included in the current study.
Informatics processing including segmentation and registration were performed on whole brain images,
but all quantification was performed in the right hemisphere to minimize potential interference from the
stereotaxic injection or EGFP fluorescence on plaque measurements.

Two-photon serial imaging of methoxy-X04 labeled plaques. To label plaques, mice received an
intraperitoneal (i.p.) injection of 3.3 mg/kg methoxy-X04 in 3.3% DMSO, 6.7% Kolliphor-EL (Millipore
Sigma) in PBS. Twenty to twenty-four hours after injection, mice were perfused with 4%
paraformaldehyde (PFA, 4°C), then brains were dissected and post-fixed in 4% PFA at room temperature
for 3-6 hours, followed by overnight at 4°C. Whole brain fluorescence imaging was performed as
described in (Oh et al., 2014) with serial two-photon (STP) tomography (Ragan et al., 2012; TissueCyte
1000, TissueVision Inc. Somerville, MA), using 925 nm excitation, a 500 nm dichroic mirror, and a 447/60
bandpass emission filter on the blue channel. Serial block-face images were acquired at 0.35 μm/pixel
lateral resolution with a 100 μm sectioning interval. We acquired 140 serial sections through each brain
from cerebellum through olfactory bulb.

Segmentation and registration. Automated image segmentation was performed as previously described
(Kuan et al., 2015) with the following modifications. Candidate plaque areas were identified by performing
adaptive thresholding on band-passed blue channel pixel strength in relation to the relative signal
strength in green channel. This additional step was implemented because many artifacts with detectable
blue signal tended to have relatively lower green signals than that of true plaques identified by expert
annotation. Since artifacts were more prevalent around tissue borders and in ventricles, the candidate

plaques are then further probabilistically filtered by a simple morphometric classifier which measures and
tests the object shape elongation, spatial location/distance to tissue border, and its relative signal strength
to tissue background autofluorescence in both green and red channels. Thirty-five of one hundred eleven
image series were acquired with a lower PMT voltage (below 750 V) initially and were then processed
with a higher sensitivity initial classifier.

Automated 3D registration was also performed as previously described (Kuan et al., 2015). Briefly,
segmented fluorescence output is a full resolution mask that classifies each 0.35 µm × 0.35 µm pixel
as either signal or background. An isotropic 3D summary of each brain is constructed by dividing each
image series into 10 µm × 10 µm x 10 µm grid voxels. Total signal is computed for each voxel by
summing the number of signal positive pixels in that voxel. Each image stack is registered in a multi-step
process using both global affine and local deformable registration to the 3D Allen Mouse Brain Common
Coordinate Framework, v3 (CCFv3). Plaque density for each structure in the reference atlas ontology was
calculated by summing voxels from the same structure. We also used a standard feature labeling
algorithm to obtain plaque counts within each structure. The algorithm labels the connected components
of the registered volume in which any two voxels of segmented plaque areas are connected if the voxels
are touching or diagonally adjacent. Due to the 100 µm z-sampling interval, our resolution limit for
detecting separate plaques in the z axis was 100 µm.

Quality Control. All image series were subjected to manual QC checks for completeness and uniformity of
raw fluorescence images, minimum fluorescence intensity, and artifacts. Severe artifacts such as missing
tissue or sections, poor orientation, edge cutoff, tessellation, and low signal strength caused image series
failure. Brains that contained slices or other damage from the stereotaxic injection, dissection, and
imaging/sectioning process were failed if the damage was in the right hemisphere. APP$^{-/-}$ brains did not
undergo additional QC checking beyond the raw image series. For APP$^{+/-}$ brains, automatic segmentation
results were checked for overall quality and false positive signals using a two-step process. First, every
image series was manually scored by an expert annotator by overlaying segmentation results for 3-5
single coronal sections with raw fluorescent images from STP imaging. A qualitative score (from 1-7) was
assigned to each image series based on the perceived overlap of the segmentation and raw image and
absence of artifacts. Second, 3D gridded plaque images for every brain were loaded in ITK-SNAP
(Yushkevich et al., 2006) and checked for obvious artifacts at tissue edges. Artifacts identified in 3D
images were subsequently checked by overlaying the corresponding single sections. In some cases,
(~25), the quality control process extended to identification and masking of areas of high intensity/high
frequency artifacts and areas of signal dropout. The edge of the cerebral aqueduct, the cerebellum, and
the medial border of the orbital cortex were particularly prone to bright tissue edge effects (**Figure 2f-k**),
so these regions were manually checked in 1-5 coronal sections and the 3D grid file for every image
series. In some cases, false positives in these regions could not be masked due to overlapping true
positive signal and manual annotators made a judgment call for inclusion based on the overall quality of
the image series. In general, APP$^{+/-}$ segmentations were failed for false positives in rostral cortex and
severe false positives in ventricles, but not for minor false positive signal in the ventricles, cerebellum and
olfactory bulb.

Antibody Characterization **6E10** (Covance #B228658, RRID: AB_1977025, 1:1000) This mouse
monoclonal IgG1 antibody is reactive to amino acid residues 1-16 of human β amyloid, specifically
recognizing the epitope of amino acids 3-8 of the sequence (EFRHDS). This antibody has been
previously shown to label amyloid plaques in the brains of humans (Patton et al., 2006) and all three AD
transgenic mice used in this study (Y. Huang et al., 2015; Pozueta et al., 2013; Thakker et al., 2009). No
staining was observed when the antibody was used to stain tissue from APP$^{-/-}$ mice. ***Goat Anti-Mouse***
IgG1(Alexa Fluor® 568). This goat polyclonal IgG1 antiserum (Invitrogen #1964384, RRID:
AB_2535766, 1:1000) is reactive against mouse IgG1. No staining was observed in tissue that was left
unexposed to 6E10 primary antibody but incubated with the IgG1 secondary, for mice of all genotypes.

Immunohistochemistry. Coronal sections retrieved after two-photon serial imaging (100 µm thickness)
were immunostained to assess Aβ load and spatial distribution. For antigen retrieval, sections were
placed in 70% formic acid for 15 minutes, followed by a PBS rinse. All sections were then incubated in
blocking solution (4% normal goat serum + 0.5% Triton X-100 in PBS) for 2 hours. After blocking,

sections were stained with primary antibody overnight (6E10, Covance, Princeton, NJ, Lot
#B2286581:1000). After three 2-hour washes in PBS + 0.1% Triton X-100, sections were then incubated
in secondary antibody overnight (Alexa Fluor goat anti-mouse 568). After three additional 2-hour washes
in PBS + 0.1% Triton X-100, sections were counterstained with DAPI (Invitrogen, Carlsbad, CA, Lot
#1874814) and coverslipped with Fluoromount G medium (Southern Biotechnology, Birmingham, AL;
catalog #J3017-XE67B). All slides were imaged on the VS120 multichannel epifluorescence microscope
system (Olympus, Center Valley, PA) with a 10X objective. Selected ROIs were subsequently imaged on
the confocal laser scanning system FV3000 (Olympus, Center Valley, PA).

Image Quantification. To quantify the fraction of methoxy-X04 labeled plaques detected by the automated
segmentation algorithm following STP imaging, we manually counted the number of plaques in 1.4 mm x
1.4 mm ROIs in raw STP coronal images and their associated segmentation mask files (n=15 ROIs total,
3 ROIs per experiment in 5 experiments). All ROIs were drawn in the isocortex and hippocampus in 12-
month-old or 18-month-old brains (n=3 APP/PS1, n=1 Tg2576, n=1 hAPP-J20). We also quantified the
fraction of antibody-labeled plaque area detected by STP imaging and automated segmentation of
methoxy-X04 labeling. We drew ROIs corresponding to major brain divisions on two sections from a
single brain for each mouse line, then applied a threshold to the 6E10 antibody labeling using Fiji (**Figure
4c**). The area in the antibody-labeled image and the corresponding segmentation of the STP image were
measured for each ROI, and the fraction detected was quantified as area in segmentation of the STP
image / area in thresholded IHC image. To quantify the ratio of dense core to diffuse plaques, we drew
polygons around the edge of the methoxy-XO4 labeling and the Aβ antibody fluorescence in maximum
intensity projections of confocal images (n=6 plaques per brain, 3 cortical and 3 hippocampal) and
measured the area inside each polygon using Fiji (Schindelin et al., 2012). We used sections from one
13-month-old APP/PS1 mouse, one 13-month-old hAPP-J20 mouse, and one 19-month-old Tg2576
mouse.

Statistics. Previous studies have reported that plaque density is not normally distributed (Liu et al., 2017).
We performed the Shapiro-Wilk test for normality on the 13-month-old APP/PS1 group since it was the
only group of mouse line and age having at least 30 samples (Razali & Wah, 2011). The test rejects the
normality of the brain-wide plaque density distribution with a p-value of 0.001. Therefore, unless
otherwise specified in the text, we used a Kruskal-Wallis one-way ANOVA with a significance level of 0.05
and Wilcoxon rank-sum test for post hoc comparisons for all hypothesis testing herein and we report
plaque densities as median ± interquartile range (IQR).

3. Results

We measured the brain-wide distribution of plaque pathology in three AD mouse models that express
mutant forms of APP using a high-throughput, high resolution imaging and analysis pipeline (**Figure 1a**).
First, plaques were labeled *in vivo* with i.p. injection of methoxy-X04 one day before transcardial perfusion
(Klunk et al., 2002). Methoxy-X04 crosses the blood brain barrier, labeling amyloid brain-wide and
producing bright fluorescence that is natively detectable; properties critical for use in our STP pipeline
(Amato, Pan, Schwartz, & Ragan, 2016). Methoxy-X04 primarily fluoresces in the blue channel, but, at
least given our acquisition parameters, some signal was also detected in green and red channels. Some
of this signal appears to be autofluorescence, as low, but still detectable, plaque signal was also
observed in APP+/- mice that did not receive a methoxy-X04 injection as has been previously reported
(Diez, Koistinaho, Kahn, Games, & Hökfelt, 2000; Dowson, 1981; Kwan, Duff, Gouras, & Webb, 2009; D
R Thal, Ghebremedhin, Haass, & Schultz, 2002; Zipfel et al., 2003, **Figure 1e**). To establish a time
course for plaque deposition in each transgenic mouse line, methoxy-X04 injections were performed in
multiple age groups between 5 and 19 months (**Table 1**). Second, following plaque labeling, brains were
imaged with serial two-photon tomography (STP) at high x,y resolution (0.35 x 0.35 μm) every 100 μm
throughout the entire rostral-caudal extent of the brain in coronal planes (**Figure 1b**). Next, each whole
brain image series was processed using the informatics pipeline adapted from (Kuan et al., 2015). This
step consists of two parts; signal detection (segmentation) and registration. We developed a custom-built
signal detection algorithm to automatically segment the methoxy-X04 labeled plaques in each serial
section (**Figure 2,** see methods for details). Segmented image stacks were deformably registered to the
Allen 3D Common Coordinate Framework v3 (CCFv3) and resampled to 10 μm voxel resolution ("3D grid

files"). Finally, we quantified the segmented methoxy-X04 labeling in terms of density and/or plaque
number for every region annotated in the Allen CCFv3 reference atlas across the entire brain.

3.1 Evaluation of automated plaque segmentation

The performance of the segmentation algorithm was analyzed in two ways. First, automatic
segmentation results were manually scored for overall quality by an expert annotator (see methods for
details). Second, we compared the number of methoxy-X04 labeled plaques detected by the automated
segmentation algorithm with the number of plaques manually identified on the raw images for a subset of
experiments and regions of interest (ROIs). On average, the automated segmentation algorithm produced
slightly higher plaque counts compared to manual counts (115+/-33%). This discrepancy is likely due in
part to the observation that some brain areas were particularly prone to false positives in the
segmentation, most often caused by bright tissue edges. These artifacts were commonly seen in the
cerebellum, around the edges of ventricles, particularly the cerebral aqueduct (**Figure 2d)**, in the rostral
cortex, particularly layer 1 of orbital cortex, and in the glomerular layer of the olfactory bulb where
lipofuscin deposits could not be distinguished from methoxy-X04-labeled plaques. The segmentation and
3D grid files were manually checked for every brain, with particular attention paid to these artifact-prone
areas.

Since some regions of the brain were prone to segmentation artifacts, confidence in automatically-
generated quantification of plaque densities is lower for some regions than others. To determine which
regions were the most problematic (and therefore had the lowest confidence), we subjected a set of 35
STP image series from wild type, non-transgenic APP$^{-/-}$ littermates with and without methoxy-X04
injections, to the same analysis pipeline as our APP$^{+/-}$ brains, *except for the segmentation QC*, so that we
could directly measure the occurrence of artifacts contributing to false positives in our data. Following the
application of the plaque segmentation algorithm to these brains, we quantified the percentage of voxels
containing false positive signal for structures in the isocortex, hippocampus, midbrain, olfactory areas,
and cortical subplate (**Figure 2e-i).** To estimate the magnitude of false positive signals across the whole
brain, we calculated the relative error per structure by subtracting the mean signal in the control dataset
from the mean signal in the full plaque dataset for every structure in annotated in the Allen CCFv3
reference atlas (840 structures). The mean false positive signal for every structure is plotted in **Figure 2j**
and included in **Appendix 1.** Fifty-two percent of structures had an error between -0.1 and 0.1 indicating
that less than 0.1% of their signal was affected by false positives. There was a second, smaller peak in
the histogram around 0.5% composed of structures that had some false positive signal. Sixteen percent
of structures had more than 0.5% difference between their plaque density and their false positive density,
and no structure had more than a 0.9% difference (**Figure 2j**). Sixteen percent of the 840 structures had
small negative values for the difference between signal and false positive percent coverage, indicating
higher false positives than true plaque signal. However, it is important to note that the control dataset did
not undergo the same segmentation QC process as the plaque dataset, so the reported false positive
values are an upper bound on the error. To make it easier to visualize the anatomical locations of regions
with high false positive signal, we generated a false positive heat map showing the spatial distribution of
regions with high segmentation background (**Figure 2k,l**). Most of these regions are located along the
midline or near ventricles and are subject to false positives from bright tissue edges (arrowheads in
Figure 1).

3.2 Metrics for quantification of plaque load using methoxy-X04 and comparison to Aβ antibody labeling

We explored several features for quantifying plaque pathology. Most frequently, we report plaque density
per structure, defined as the percent of each structure's total volume that contains segmented plaque
signal. We also automatically calculated plaque counts based on the segmentation results. To accomplish
this, we computationally identified voxels with segmented plaque that were touching and/or diagonally
adjacent to each other in the 3D grid images (**Figure 3**). Our 100 μm z-sampling interval caused plaques
to have an elongated appearance along the anterior-posterior axis (**Figure 3c,d**). However, we also
identified individual plaques that extended for several millimeters across serial sections, much further
than our sampling interval would explain. This was particularly apparent along the midline where many

plaques appeared to follow the path of the vasculature. This was more obvious in hAPP-J20 and Tg2576
mice (**Figure 3e, f**) than in APP/PS1 mice (**Figure 3d**), but was observed in all three APP mouse lines.
Because we could not easily differentiate between these extended length vascular-associated plaques
and the non-vascular plaques based on the automated counting results, in most of our subsequent
analyses we report plaque density to describe pathological load. However, the number of plaques per
cubic mm are reported in **Figure 5a** and included in **Appendix 2**, and it should be noted that these values
were not corrected for the presence of vascular-associated plaques.

Methoxy-X04, a congo red derivative, was reported to label dense-core amyloid but not diffuse plaques
(Condello, Schain, & Grutzendler, 2011). Thus, we wanted to characterize the relative fraction of *total*
plaque detectable with methoxy-X04 compared to labeling sections with an Aβ antibody. We retrieved
individual sections from a subset of experiments following methoxy-X04 labeling and STP imaging, and
immunostained them with the 6E10 anti-Aβ antibody (**Figure 4a,b**). We quantified the area covered by
plaque in thresholded, antibody-labeled sections compared with the automated segmentation of the same
sections **(Figure 4c**, *e.g.,* compare red to white masks). In all sections, the overall plaque density in the
antibody-labeled sections was higher than the automated segmentation of methoxy-X04 fluorescence, but
the densities for the two methods of plaque labeling differed between mouse lines (APP/PS1: 10.4±7.8%
IHC, 1.2±0.80% segmentation (13-months); Tg2576: 2.8±2.5% IHC, 0.44±0.37% segmentation (19-
months); hAPP-J20: 7.2±7.2% IHC, 0.08±0.07% segmentation (13-months)). To more directly measure
our detection level, we calculated the fraction of Aβ antibody-labeled area that was detected by automatic
segmentation of methoxy-X04 fluorescence in the same section for all ROIs in each image. Surprisingly,
we found a significant effect of mouse line on the fraction of antibody-labeled plaque area detected
(**Figure 4f,** p=0.04, one-way ANOVA). In APP/PS1 and Tg2576 mouse lines, the fraction detected by
segmentation was similar (0.15±0.10 for APP/PS1, 0.15±0.11 for Tg2576), but a lower proportion of
antibody labeled plaques were detected in hAPP-J20 (0.03±0.03, p=0.04 compared to APP/PS1, Tukey's
post hoc comparison). This line-specific difference may be due to the significantly lower ratio of dense
core to diffuse plaque area that we measured in hAPP-J20 mice compared to the other two lines
(APP/PS1: 0.09±0.04, Tg2576: 0.1±0.03, hAPP-J20: 0.03±0.02, p=0.001 one-way ANOVA, **Figure 4g**).
We did not observe any antibody-labeled plaques in regions without methoxy-X04 labeled plaques, and
critically, plaque densities measured by both methods are very highly and significantly correlated for all
regions tested in all three mouse lines (r=0.95, **Figure 4e**), supporting the use of the methoxy-X04 label in
our systematic pipeline approach to describe brain-wide distribution of plaques. Thus, we find that
methoxy-X04 does underestimate total, or absolute, plaque density (including diffuse plaques) by ~7- to
10-fold for APP/PS1 and Tg2576 mice, and ~30-fold for hAPP-J20 mice. Due to this difference in the
fraction of plaques labeled with methoxy-X04, plus known differences in the rate of plaque accumulation
in each mouse line (Jankowsky & Zheng, 2017), we focused our analysis on *patterns* of plaque deposition
rather than comparing absolute levels between lines. Where we do compare plaque levels (**Section 3.3**),
it is important to note that these reported values refer only to dense-core, not diffuse, plaques.

3.3 Comparison of Methoxy-X04-labeled plaque levels between three APP mouse models.

We measured methoxy-X04 labeled plaque density and count for each mouse line across multiple ages in
the whole brain (**Figure 5a,b**). Of the three lines, APP/PS1 mice showed the most aggressive rates of
dense-core plaque accumulation, as expected from previous reports (Garcia-Alloza et al., 2006;
Jankowsky et al., 2004). Whole-brain dense core-plaque densities were higher in APP/PS1 mice
compared to the other two mouse lines in every age group, and whole brain plaque counts were higher in
APP/PS1 mice compared to the other two lines for every age except 5 months. At 5 months-old, plaques
were readily observable in APP/PS1 mice. Occasionally, but still quite infrequently, plaques were
detected in 5 and 7-month-old hAPP-J20 mice. At 13 months, hAPP-J20 mice had significantly higher
plaque density than Tg2576 mice (**Figure 5b**, p=0.04). Very sporadic plaques were observed in Tg2576
mice at 13 months, and they then accumulated more plaques rapidly between 13 and 19 months.

We also measured methoxy-X04-labeled plaque densities within each of the twelve major brain divisions,
and fiber tracts, annotated in the Allen CCFv3 reference atlas. In **Figure 5c,d**, we show plaque density
across all age groups for the isocortex (**Figure 5c**) and hippocampal formation (**Figure 5d**), As seen at
the whole brain level, methoxy-X04 labeled dense-core plaque densities in these two major regions were

also higher in APP/PS1 mice compared to Tg2576 or hAPP-J20 mice at every age. Tg2576 and hAPP-
J20 mice did not differ in isocortex plaque density, but hAPP-J20 mice had significantly higher plaque
density in the hippocampal formation compared to Tg2576 mice at 9 and 13-months-old (p=0.02, p=0.04,
Figure 5d).

3.4 Temporal and spatial progression patterns of Methoxy-X04 labeled plaques in three APP mouse models.

Plaque densities across all the major brain divisions are shown for all three mouse lines in 13- and 19-
month-old brains in **Figure 5e-g**. APP/PS1 mice had the highest plaque density in the isocortex but also
had dense plaques in olfactory areas, hippocampal formation, cortical subplate, and cerebellum at both
13- and 19-months-old. At 13 months, APP/PS1 mice were also beginning to show some plaque
accumulation in the thalamus and striatum although the plaque density in these two structures was not
significantly higher than the median for the whole brain at either 13 or 19 months. Tg2576 mice had only
very sporadic plaques at 13 months, but by 19 months-old their plaque distribution across major regions
looked similar to APP/PS1 mice; densest plaques in isocortex, followed by olfactory regions, hippocampal
formation, cortical subplate, and cerebellum. However, even though the plaque distribution was broadly
similar, Tg2576 still had much lower plaque densities than APP/PS1 mice (note the different scales for
the APP/PS1 mouse line plot compared with Tg2576 and hAPP-J20 in **Figures 5e** and **5g**). Although
there was a significant effect of structure on plaque density in major brain divisions in Tg2576 mice
(p=10^{-5} at 13 months, p=0.001 at 19 months), no individual brain division had a significantly higher plaque
density than the median in post hoc testing. Notably, hAPP-J20 mice had a different brain-wide plaque
distribution from the other two lines. Plaque levels in the isocortex and hippocampal formation were
significantly higher than the whole brain median at 13 months, and plaque density in the cortical subplate
was significantly lower than the median at the same time point (**Figure 5e**). By 19 months, hAPP-J20
mice still had more plaques in the hippocampal formation than any other brain division and there was a
significant effect of structure on plaque density (p=0.02), but none of the plaque densities in individual
brain divisions were significantly different from the median at 19-months.

To examine the regional distribution of methoxy-X04 labeled plaques at a finer scale, we quantified
densities for each of 316 subdivisions of the 12 major brain regions (aka, "summary structures") in the
Allen CCFv3. Plaque densities are plotted in **Figure 6** for summary structures in the 4 major brain
divisions with the most plaque (isocortex, hippocampal formation, olfactory areas, and cortical subplate)**.**
Very small summary structures are not shown here (*i.e.,* total volume < 0.5% of its major division), but the
plaque densities for all structures in all mouse lines and age groups are reported in **Appendix 2**.

The isocortex is parcellated into 43 regions (or, summary structures) in the Allen CCFv3. We observed
widespread distribution of plaques across these regions in APP/PS1 mice at 13 and 19 months (**Figure
6a)**. However, there was still some regional specificity within the cortex for this APP line. Plaque densities
in the lateral visual area (VISl) were significantly higher than the median plaque density across the entire
isocortex at 13 months (**Figure 6a,b**). Prefrontal areas prelimbic (PL), infralimbic (ILA), and medial orbital
cortex (ORBm), had lower plaque density compared to the whole isocortex median (**Figure 6a**). hAPP-
J20 mice showed a very striking and specific regional distribution of plaque in the isocortex at both 13 and
19 months, but we only performed statistical testing at 13 months since the isocortex was not identified as
significant with post-hoc testing across all major divisions at 19 months. Plaque density in the isocortex of
13-month-old hAPP-J20 mice was extremely variable, with 23 summary structures having median plaque
densities that were significantly higher or lower than the median for the whole structure (**Figure 6a**).
Specifically, we identified levels that were 5-13 times higher than the median in three subdivisions of
retrosplenial cortex (RSPagl, RSPd, RSPv) at 13 months and 2-4 times higher than the median at 19
months, an observation consistent with previous reports (Harris et al, 2010). Interestingly, plaque density
in RSPd and RSPv was significantly *lower* than the isocortex median for APP/PS1 mice (**Figure 6a,b**). In
addition to RSP subdivisions, hAPP-J20 mice had high plaque density in ventral anterior cingulate cortex
(ACAv), parietal association cortex (anterior visual area, VISa, anteromedial visual area, VISam, and
posteromedial visual area, VISpm), and surprisingly, primary visual cortex (VISp). Tg2576 mice also
appeared to have heavier plaque accumulation in dorsal anterior cingulate (ACAd) and parietal (VISam,

VISa, and VISrl) cortex than in other cortical areas at 19 months (**Figure 6a**) but there was no significant
effect of structure on plaque density in isocortex for this mouse line at any age.

The hippocampal formation (HPF) contains 14 summary structures (**Figure 6c**), including the
hippocampus itself (CA1, CA3, DG, SUB) as well as associated regions (e.g., entorhinal cortex, ENT).
Within the HPF, structure had a significant effect on plaque density for every mouse line/age group
combination. In the hippocampus proper, all three lines had higher plaque density in the DG than in CA1,
CA2, or CA3 (**Figure 6c,d**). APP/PS1 and Tg2576 mice also had high plaque density in lateral and
medial entorhinal cortex (ENTl and ENTm). hAPP-J20 mice had plaques in ENTm but plaques in ENTl
were rarely observed. All three mouse lines had relatively heavy plaque accumulation in the subiculum
(SUB) and prosubiculum (ProS), but the Tg2576 mouse line had a strong preferential accumulation of
plaques specifically in the dorsal subiculum (**Figure 6d,** white arrow in **Figure 1**). Notably, every single
individual Tg2576 mouse that we imaged had plaques in the dorsal subiculum, even at 9 months-old.

Two other brain divisions with differences in plaque density between mouse lines were the olfactory areas
which had dense plaques in many regions for APP/PS1 and Tg2576 mice, but virtually no plaques in
hAPP-J20 mice (**Figure 1, Figure 6e,f**) and the cortical subplate which contains several amygdalar
structures, where plaques were similarly high in APP/PS1 and Tg2576 but low or absent in hAPP-J20
mice (**Figure 6g,h**). Importantly, we also did not observe Aβ antibody-labeled plaques in olfactory regions
or the amygdala in hAPP-J20 mice. Also of note, both APP/PS1 and Tg2576 mice had plaques in the
cerebellum, whereas we did not identify any in hAPP-J20 mice (**Figure 7**). APP/PS1 mice developed
cerebellar plaques as early as 4 months-old, but plaques were not observed in the cerebellum of Tg2576
mice until 19 months-old, and then many were associated with vasculature.

We also explored the relationship between regional plaque distribution patterns in the isocortex with the
structural connectivity-based isocortical modules found from a network analysis of modeled connectivity
weights in wild type C57BL/6J mice (Harris et al., 2018, Knox et al., 2018). These modules are comprised
of cortical regions that are more strongly connected to each other than to regions in other modules. The
medial and prefrontal modules contain most of the regions reported to be part of the rodent DMN (Lu et
al., 2012; Sforazzini, Schwarz, Galbusera, Bifone, & Gozzi, 2014; Stafford et al., 2014; Zerbi, Grandjean,
Rudin, & Wenderoth, 2015); including RSP, ACA, posterior parietal (PTLp aka VISa and VISrl), ORB, and
PL. The plaque density in each of the six cortical modules for each mouse line compared to the expected
density if plaques were evenly distributed across the entire cortex is plotted in **Figure 8a**. hAPP-J20 mice
had three times as many plaques in the medial module as would be expected with an even distribution
and one and one-half times the expected density in the prefrontal module, indicating a distribution that
overlaps with the rodent DMN. The relative plaque density in the medial module for hAPP-J20 mice was
significantly higher than any other module (p<0.0001, 2-way ANOVA with Sidak's multiple comparisons
test), and relative plaque density in the prefrontal module was significantly higher than all other modules
except visual (p<0.0001 for all other modules, p=0.5 for visual, 2-way ANOVA with Sidak's multiple
comparisons test). hAPP-J20 mice also had lower plaque density in the anterolateral, somatomotor, and
temporal modules compared to all other modules, but these three modules did not differ from each other.
Tg2576 mice had significantly fewer plaques in the temporal module compared to the prefrontal and
visual modules (p=0.02, p=0.003, 2-way ANOVA with Sidak's multiple comparisons test) and APP/PS1
mice had significantly fewer plaques in the medial compared to the visual module (p=0.003, 2-way
ANOVA with Sidak's multiple comparisons test). To visualize these differences in plaque distribution
across modules, we created images of the Allen CCFv3 reference atlas structures showing the positions
of the modules with a colormap corresponding to relative plaque levels for each of the three mouse lines
(**Figure 8b**).

To assist in visualizing the whole brain, 3D, regional distribution patterns of methoxy-X04 labeled plaques
for the different ages and mouse lines, we created 3D heatmaps (**Figure 9, Figure 10**). These maps are
based on the Allen CCFv3 with each annotated brain structure colored by plaque density. **Figures 9 and**
10 show example images from different 3D locations (coronal, sagittal, and horizontal slices) in 13-month-
old APP/PS1 and hAPP-J20 maps, and 19-month-old Tg2576 maps (since plaque density was very low
at 13 months in this line). **Movie 1** shows a flythrough of these three maps in the coronal plane. Maps for

all mouse line/age group combinations are available for download and can be viewed in image software
such as ITK-SNAP (http://www.itksnap.org/) [link to be provided].

One spatial feature of amyloid deposition that is easily visualized using the plaque heatmaps is the
distribution of plaques by layer in the isocortex. The high density of plaques in layer 5 for APP/PS1 mice
and layer 1 for Tg2576 mice are prominent features in the 3D maps (i.e. compare **Figure 9b** with **Figure**
9d). To quantify differences in plaque density across cortical layers, we calculated the plaque density per
layer across all isocortical structures and compared it to the plaque density that would be expected with
even plaque distribution based on the relative volume of each layer in the Allen CCFv3. The relative
plaque density in each layer for the three mouse lines is plotted in **Figure 11**. Plaque density in APP/PS1
mice was more uniform across layers than the other two lines, but was higher in layer 5 (p<0.0001 than in
layers 4, 6a, and 6b; p=0.03 vs. layer 1; 2-way ANOVA with Sidak's multiple comparisons test) (**Figure**
11a,b). There appeared to be some regions such as VISrl where plaques were as dense in layer 2/3 as in
layer 5 (**Figure 9b, Figure 10b**), and indeed the density in layer 5 was not different from layer 2/3 (p=0.8,
2-way ANOVA with Sidak's multiple comparisons test). In the retrosplenial cortex of hAPP-J20 mice,
plaques density was highest in layer 5 **(Figure 9c, Figure 10c)**, but in other cortical regions, hAPP-J20
and Tg2576 mice both had the highest plaque density in layer 1 (**Figure 11c-f, Figure 9c,d; Figure**
10c,d). Across all regions, both Tg2576 and hAPP-J20 mice had higher plaque density in layer 1 than in
any other layer (p<0.0001, 2-way ANOVA with Sidak's multiple comparisons test). The strong preference
for layer 1 plaques in Tg2576 and hAPP-J20 mice appeared to come partly from the surface vasculature
which had heavy methoxy-X04 labeling in Tg2576 mice at all ages and 19-month-old hAPP-J20 mice
(**Figure 4**). In Tg2576 mice, there was no difference in the relative plaque density between layer 2/3 and
layer 5, but hAPP-J20 mice had significantly lower plaque density in layer 5 than in layer 2/3 across the
whole cortex (p=0.007, two-way ANOVA with Sidak's multiple comparisons).

In summary, we quantified patterns of plaque deposition across brain regions at the major division and
summary structure level for three APP transgenic mouse lines. We found line-specific differences in
plaque density for different brain divisions and for structures within major brain divisions. The pattern of
plaque deposition in the isocortex differed between mouse lines at both the regional and layer levels.

3.5 Comparison of plaque deposition patterns between mouse lines and human

In AD patients, plaque accumulation follows a predictable pattern starting in the isocortex, followed by
hippocampal regions, then other subcortical regions such as striatum, basal forebrain, and thalamus, and
finally brainstem nuclei and cerebellum. These stages of Aβ deposition were described by (Thal, Rüb, et
al., 2002) based on quantifying the percent of human tissue samples with plaques in a list of regions
across the whole brain. To compare the brain-wide patterns of plaque deposition described here with the
pattern of Aβ deposits observed in autopsy tissue from patients in different phases of AD, we examined
plaque density in rodent homologs to the human brain regions used to quantify AD staging in (Thal, Rüb,
et al., 2002). For each region reported to contain Aβ deposits (left column, **Figure 12a**), the closest
mouse homolog from the Allen CCFv3 reference atlas is listed in the second column. In cases where
there was more than one mouse structure that could correspond to the human brain region listed, we
used the median plaque density across all the corresponding structures. We calculated the median
plaque density per structure across ages for each of the three APP mouse lines and, for APP/PS1 and
hAPP-J20 mouse lines, we subtracted the median false positive value for each structure in control mice of
the same line. We applied a color map spanning from the 10th to 90th percentile of plaque densities for **all**
annotated structures (840) for each group to create a table similar to the one in (Thal, Rüb, et al., 2002)
(**Figure 12a**). The temporal and regional pattern of plaque deposition for APP/PS1 and Tg2576 mice was
remarkably similar to the pattern reported by Thal et al. with a few notable exceptions. Most importantly,
plaques accumulated in the cerebellum (CBX) very early for APP/PS1 mice and were also seen in
Tg2576 mice, but plaques were not regularly seen in the cerebellum until the last phase of Aβ deposition,
Phase 5, in human brain tissue. Plaque density was high in central gray for all three mouse lines and in
the substantia nigra for APP/PS1 and hAPP-J20 mouse lines, but these were both regions in which we
observed false positive signal in our automated segmentation, so these values may overestimate the true
plaque signal even though we subtracted the control data values. In the top portion of the chart in **Figure**
12a, the plaque distribution in APP/PS1 mice and Tg2576 mice follows a similar step-wise pattern to that

described in autopsy tissue for Phase 1, Phase 2, and the beginning of Phase 3. The thalamus,
hypothalamus, and striatum were key regions that defined Phase 3 for human brains. While APP/PS1
mice began to accumulate plaques in the thalamus and striatum at 13 months, at 19-months-old they did
not have plaques in the hypothalamus and still had low plaque density in the basal forebrain and
brainstem nuclei. This implies that even after aging for 1.5 years, APP/PS1 mice, which had the most
aggressive plaque deposition of the three tested, had only partially entered Phase 3. Tg2576 mice had
low but measurable plaque density in the thalamus, hypothalamus, and striatum, but we did not observe
plaques in these regions so these values were likely false positives.

Counting the number of cases in which plaques were observed in all regions from the table would be
difficult to perform computationally with our data because of the need to choose a threshold for calling a
region "positive" for the presence of plaques. We instead made the assumption that plaque density and
percent prevalence are related, and compared the plaque density values from our quantification with the
percent prevalence values reported in (Thal, Rüb, et al., 2002). To determine which Phase of Aβ
deposition was most closely matched by each experimental mouse model, we created a correlation matrix
by calculating the Pearson correlation between the plaque density in all the structures in the table for
each mouse line/age combination and the percent prevalence of plaques in each structure as reported by
(Thal, Rüb, et al., 2002) (**Figure 12c**). For the APP/PS1 mouse line, plaque distribution was most highly
correlated with Phase 2 of Aβ deposition at all ages, but the correlation between Phase 2 and Phase 3
was approximately equal for the 19-month age group confirming that 19-month-old APP/PS1 mice were
transitioning to the equivalent of Phase 3. Thirteen and nineteen-month-old Tg2576 mice also had the
highest correlation with Phase 2. hAPP-J20 mice, however, did not have a high correlation with any of the
phases described in human cases. At 19-months-old, hAPP-J20 mice had a moderate correlation with
Phased 1 in human tissue (which is defined only by plaques in the neocortex), but this correlation
coefficient was still less than 0.5. In the cases used to develop the phasing system, nondemented
patients had Aβ pathology in Phases 1, 2, and 3, while clinically proven Aβ cases exhibited Aβ-Phases 3,
4, and 5 (Thal, Rüb, et al., 2002). Based on this metric, 19-month-old APP/PS1 and Tg2576 mice most
closely resemble patients around the time of first diagnosis.

4. Discussion

We adapted the imaging, automated detection, and registration pipelines previously used for brain-wide
quantification of axonal projections (Oh et al., 2014) to measure whole-brain amyloid pathology in three
commonly used mouse models of AD. Automated segmentation of methoxy-X04 labeled plaques,
combined with the 3D registration of whole brain image series to the Allen CCF volumetric reference atlas
allow us to systematically quantify brain-wide distribution of dense-core plaque pathology at high
resolution for a large sample size. The processing of non-transgenic APP$^{-/-}$ littermates with the same
pipeline facilitated interpretation of results by providing a confidence metric for the automated brain-wide
plaque densities reported here. We found that plaque deposition patterns were heterogeneous across
structures in all three transgenic mouse lines examined, pointing to the need for robust whole brain
characterization to optimize the usage of existing and novel mouse models.

Comparing estimates of plaque density across different studies and brain regions is challenging, as it
requires considering the labeling method, staining conditions, and imaging conditions used in each study
in addition to animal age and transgenic line. Our pipeline approach provided consistency in labeling
method and imaging conditions, and our use of automatic segmentation ensured consistent detection of
plaque density across many images and samples, allowing us to quantitatively compare plaque density
across the whole brain with unprecedented anatomical specificity. However, quantitatively comparing
results obtained with different methods of plaque labeling is still an important challenge. We found that
methoxy-X04 labeling is inconsistent between mouse lines and that a lower fraction of total plaque area
was detected with methoxy-X04 in hAPP-J20 mice compared with APP/PS1 and Tg2576 mouse lines.
We measured the magnitude of this difference by comparing automatically-detected methoxy-X04
plaques with antibody labeled plaques in the same sections, but it is important to note that our automatic
segmentation is based on single confocal section images while the same sections were imaged with a
widefield microscope after antibody labeling. However, the ratio between the two labeling methods would
not be expected to differ between mouse lines. Importantly, the fraction of individual plaque area covered

by methoxy-X04 vs. Aβ antibody measured from confocal sections is similar to the ratio between
methoxy-X04 and antibody labeled areas measured in large ROIs (**Figure 4f,g**), indicating that the
difference in detection likely reflects a biological difference in the composition of plaques for the different
mouse lines. To our knowledge, this difference in the ratio of dense-core/diffuse plaques in hAPP-J20
mice compared with other APP models has not been previously reported. It will be important to confirm
this result using other methods of labeling dense-core and diffuse plaques. Critically, although our
experiments underestimate total plaque density, they do represent brain-wide plaque location more
precisely than any other study to date in mouse models of AD.

Despite the challenges in directly comparing results between labs, we did compare our quantified plaque
density values for the isocortex and hippocampus to published measurements when possible to both
establish general trends in our data as compared to others, and to evaluate our automated segmentation
technique. One recent comparable study quantified Thioflavin S-labeled plaque load in the cortex and
hippocampus for both APP/PS1 and Tg2576 mice (Liu et al., 2017). Our measured plaque densities are
on average 35% of those reported by Liu et al. (e.g., for our 13-month-old APP/PS1 mice and Liu et al.'s
10-13-month age group: cortex median = 0.80% vs. 2.1% and hippocampus median = 0.64% vs. 1.4%;
for our 19-month-old Tg2576 and Liu et al.'s 14-17-month age group: cortex median = 0.20% vs. 1.9%
and hippocampal median = 0.19% vs. 0.54%). On the other hand, our plaque densities are slightly higher
than another report (Garcia-Alloza et al., 2006) that also used Thioflavin S labeling to quantify plaque
density in the cortex of APP/PS1 mice (for our 9-month-old age group and Garcia-Alloza et al.'s 10-
month-old age group: cortex median = 0.96 (IQR 0.67-1.8) vs. ~0.13-0.22%). We also report lower
densities (0.2% (7 mo), 0.5% (9 mo) vs. ~2% (7-8.5 mo)) than a group using light sheet microscopy to
image methoxy-X04-labeled plaques in cleared APP/PS1 mouse brains (Jährling et al., 2015). The
plaque density we measured in 9-month-old hAPP-J20 hippocampus is nearly identical to a recent report
using Thioflavin S staining (0.03%, Zhang et al., 2017). Overall, it seems that the densities we measured
are within reported range of previous studies, albeit on the lower end of ranges.

Previous studies have reported plaque density in the cortex and hippocampus for different AD mouse
models across ages, but generally regional distribution of plaque burden is not resolved beyond these
broad divisions, much less across the entire brain. Recently, a 3D survey of Alzheimer's pathologies in
transgenic mouse models including APP/PS1 and Tg2576 was reported using the iDISCO clearing
method (Liebmann et al., 2016). The authors imaged plaques brain-wide using tissue clearing followed by
light sheet microscopy and registration of the whole brain images to the Allen CCFv3 reference atlas. The
values reported by Liebmann et al. for plaque count (plaques per mm^3) are higher than ours by a factor of
~10, which could potentially reflect the higher axial resolution of light sheet microscopy (2.5 μm vs. our
100 μm sectioning interval), but may also be caused by our automated detection algorithm merging
individual plaques (see **Figure 4**). Liebmann et al. did not report plaque densities for all structures across
the brain, but the twelve structures with highest plaque burden in APP/PS1 mice in their study generally
agreed with our findings. In the isocortex, they found the highest plaque load in layers 2/3 and 5 of
primary motor cortex (MOp), layers 2/3 and 5 of barrel cortex (SSp-bfd), supplementary somatosensory
cortex (SSs), and dorsal auditory cortex (AUDd). While none of these structures had significantly higher
density compared to the median for the isocortex in our data, all regions were in the top 5% of densities
for both our 13-month-old and 19-month-old mice. Our results were similarly in agreement with the
structures in the hippocampal formation found to have high plaque load (i.e., subiculum, lateral entorhinal
cortex, CA1, and the molecular layer of the dentate gyrus, **Figure 6**). There were, however, notable
differences between our datasets, as Liebmann et al. reported high plaque density in fiber tracts,
specifically in the piriform area and the corpus callosum. While the piriform cortex and corpus callosum
both had measurable plaque densities in our data, neither was particularly high. Liebmann et al. focused
more on individual plaques than on brain-wide plaque distribution, and they also reported variability in the
shape of individual amyloid deposits, including the presence of elongated structures and vascular
deposits.

Our method has several advantages over the clearing method used by Liebmann et al. Most importantly,
we collected image data using the STP imaging system that was also used to construct the Allen CCFv3,
so the variability in our 3D registration to this reference atlas was as low as is technically possible. Our
sample size is also very high for some groups, increasing confidence in our results. We put a great deal

of effort into developing and validating our custom segmentation algorithm as described in the results,
although we did not similarly optimize our plaque counting algorithm, which could potentially be improved
with future iteration. Finally, we labeled plaques with methoxy-X04 *in vivo*, rather than post-hoc, so
penetration of the plaque label into the tissue was not a concern. Tissue clearing, however, has the
distinct advantage of being amenable to more types of labels, and continues to be further optimized to
best combine antibody labeling and high resolution whole brain imaging methods (like STP and light
sheet microscopy). It will be useful to compare whole brain quantification of Aβ antibody-labeled plaque
density with the methoxy-X04 density reported here, particularly for the hAPP-J20 line since its fraction of
dense core plaque labeling is lower than the other APP lines.

The advent of PiB-PET imaging as a biomarker in AD has been a major breakthrough for the field. Not
only does it allow non-invasive estimation of whole-brain plaque density and deposition patterns in
humans, but patients can also be diagnosed with probable AD much earlier in the disease process and
clinical trials can be more closely controlled based on biomarker measurements (Counts, Ikonomovic,
Mercado, Vega, & Mufson, 2017). To best align pre-clinical research in mouse models with human
biomarker data, the same PET imaging methods should be applied to mice. While STP imaging of
methoxy-X04 labeling gives much higher spatial resolution than possible with PET imaging, ideally one
would like to seamlessly move across scales in mice to enable better experimental testing of potential
mechanistic underpinnings of disease pathology that could be extensible to humans. Unfortunately, PiB
does not label Aβ deposits in some mouse models, including both APP/PS1 and Tg2576 (Klunk et al.,
2005; Snellman et al., 2013), but at least one quantitative comparison between amyloid PET and
methoxy-X04 labeling in different mouse lines has been performed (Brendel et al., 2015). The
significance of the differential affinity of imaging probes for various forms of Aβ is still an active area of
research (Schilling et al., 2016).

For many years, regional distribution patterns have been studied in relationship to tau pathology and its
progression with disease stage (H Braak & Braak, 1991; Heiko Braak, Alafuzoff, Arzberger, Kretzschmar,
& Del Tredici, 2006). Tau progression through brain networks has more recently been interpreted as
prion-like propagation between cells (Lewis & Dickson, 2016). Not to be left behind, there has been
increasing recognition of the occurrence of specific patterns of regional amyloid pathology as well in AD,
starting with the discovery that Aβ plaques preferentially form in regions of the brain that are part of the
resting state default mode network (DMN, R. L. Buckner, 2005). Not only do AD patients have early and
heavy deposition of plaques in the DMN (Mormino et al., 2011; Palmqvist et al., 2017), the activity of this
brain-wide network is also increasingly impaired as disease progresses (Brier et al., 2012; Hafkemeijer,
van der Grond, & Rombouts, 2012; Jones et al., 2015; Mormino et al., 2011). These functional
connectivity deficits related to the DMN are robust enough that they have been proposed as a biomarker
to track disease progress, although several limitations still need to be addressed before this could be
implemented (Jones, 2016). The mechanisms underlying this link between Aβ pathology and functional
impairment of large-scale networks have been elusive at the cellular and behavioral level, where plaque
pathology, neurodegeneration, and cognitive impairment have not been well-correlated (Nelson et al.,
2012). One intriguing hypothesis is centered around the findings that Aβ deposition is related to neuronal
activity (Bero et al., 2011; Cirrito et al., 2008; Kamenetz et al., 2003; Li et al., 2013). A recent study
directly tested this hypothesis in Tg2576 mice by depriving the barrel cortex of sensory input, which
decreased neuronal activity, and measured the effect regional plaque load (Bero et al., 2011). These
authors wisely began their study by quantifying plaque load across cortical areas before using sensory
deprivation to perturb the distribution of plaque levels. An increased focus on brain-wide distribution of
pathology, in addition to pathology levels, should be an important component of ongoing research in
rodent models of AD to best align with data from patients.

In AD patients, plaque deposition expands from the DMN and other cortical structures to hippocampus
and subcortical regions. Plaque pathology in the APP transgenic lines tested here mimics these patterns
of regional vulnerability in different ways. hAPP-J20 mice have a cortical plaque distribution pattern that
most resembles the recent characterization of a rodent DMN homolog (Lu et al., 2012; Sforazzini et al.,
2014; Stafford et al., 2014; Zerbi et al., 2015). The preferential accumulation of plaques in retrosplenial
and prefrontal cortex in these mice suggests that they could be useful for testing models of regional
vulnerability to amyloid deposition in the cortex. In contrast, the brain-wide sequence of plaque deposition

in APP/PS1 and Tg2576 mice is more similar to the pattern seen in patients (Thal, Rüb, et al., 2002),
suggesting these mouse lines may be more appropriate for studies focusing on brain-wide pathology or
disease progression.

There are several possible reasons for the differences we observed in spatial patterns of Aβ deposition
between APP-overexpressing transgenic mouse lines. One obvious possibility is the different promotors
driving transgene expression (**Table 1**) and/or their different genomic insertion sites. A recent study found
differences in expression pattern for some AD mouse models using two newly developed antibodies that
can differentiate between mouse and human APP (Höfling et al., 2016). Tg2576 mice were included in
their study but unfortunately APP/PS1 and hAPP-J20 mice were not. However, this approach could be
applied to additional mouse lines to characterize the interaction between APP expression and Aβ
deposition. Interpreting the relationship between APP expression and plaque density may be complicated
by the fact that measurements of APP expression patterns are dominated by signal in cell bodies, but
APP levels in axons may be more relevant to plaque deposition. Testing this hypothesis by correlating
APP and plaque levels in the same region or in synaptically connected regions requires quantification of
both APP expression and plaque deposition patterns with high spatial resolution, but this type of study
could potentially reveal interesting mechanistic details of plaque formation.

Another possibility for the differences in spatial pattern of Aβ deposition is that each pattern reflects a
separate aspect of cellular and network vulnerability. Selective vulnerability could occur at the regional
level (e.g. from differences in activity across brain regions) or at the cellular level (e.g. from damage to
specific neuronal projection types). Since the time course and type (dense-core vs. diffuse) of Aβ
deposition differs across lines, each mouse lines may be susceptible to damage by different mechanisms.
Unraveling the differences between mouse lines and cell types in their vulnerability to AD pathology is a
challenging problem but these experiments have great potential for unlocking mechanisms of network
neurodegeneration.

The age- and line-specific average brain-wide plaque maps described here are available for download as
nearly raw raster data (nrrd) files that can be viewed with the free software ITK-SNAP
(http://www.itksnap.org/) or loaded into Python or Matlab (The MathWorks, Inc.) as 3D arrays. We
encourage researchers to use these maps for comparing amyloid distribution patterns between these
transgenic models and choosing appropriate brain regions and mouse lines for hypothesis testing. These
significant differences in amyloid deposition patterns between similar transgenic mouse lines underline
the importance of careful consideration when choosing an AD mouse model (Jankowsky & Zheng, 2017)
and importantly, indicate that appropriately chosen APP mouse models can be useful for modeling
amyloid related brain-wide network degeneration.

Author Contributions
Conceptualization: JAH, JDW. Supervision: JAH, JDW. Data Acquisition: JDW, AP, PB, AH. Data
Curation: JDW, ARB, AM, KEH. Informatics Pipeline Development: LK, NG, WW. Data Analysis: JDW,
ARB, JEK, NG, AP, AM. Visualization: JDW, ARB, AP, AM, KEH. The original draft was written by JDW
and JAH, with input from JEK, LK, AP. All co-authors reviewed the manuscript.

8. References

Amato, S. P., Pan, F., Schwartz, J., & Ragan, T. M. (2016). Whole Brain Imaging with Serial Two-Photon
Tomography. *Frontiers in Neuroanatomy*, *10*, 31. https://doi.org/10.3389/fnana.2016.00031

Bero, A. W., Yan, P., Roh, J. H., Cirrito, J. R., Stewart, F. R., Raichle, M. E., … Holtzman, D. M. (2011).
Neuronal activity regulates the regional vulnerability to amyloid-β deposition. *Nature Neuroscience*,
14(6), 750–756. https://doi.org/10.1038/nn.2801

Braak, H., Alafuzoff, I., Arzberger, T., Kretzschmar, H., & Del Tredici, K. (2006). Staging of Alzheimer
disease-associated neurofibrillary pathology using paraffin sections and immunocytochemistry. *Acta
Neuropathologica*, *112*(4), 389–404. https://doi.org/10.1007/s00401-006-0127-z

Braak, H., & Braak, E. (1991). Neuropathological stageing of Alzheimer-related changes. *Acta

Neuropathologica, *82*(4), 239–259. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/1759558

Brendel, M., Jaworska, A., Grießinger, E., Rötzer, C., Burgold, S., Gildehaus, F.-J., … Rominger, A.
(2015). Cross-sectional comparison of small animal [18F]-florbetaben amyloid-PET between
transgenic AD mouse models. *PloS One*, *10*(2), e0116678.
https://doi.org/10.1371/journal.pone.0116678

Brier, M. R., Thomas, J. B., Snyder, A. Z., Benzinger, T. L., Zhang, D., Raichle, M. E., … Ances, B. M.
(2012). Loss of Intranetwork and Internetwork Resting State Functional Connections with
Alzheimer's Disease Progression. *Journal of Neuroscience*, *32*(26), 8890–8899.
https://doi.org/10.1523/JNEUROSCI.5698-11.2012

Buckner, R. L. (2005). Molecular, Structural, and Functional Characterization of Alzheimer's Disease:
Evidence for a Relationship between Default Activity, Amyloid, and Memory. *Journal of*
Neuroscience, *25*(34), 7709–7717. https://doi.org/10.1523/JNEUROSCI.2177-05.2005

Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L. (2008). The brain's default network: Anatomy,
function, and relevance to disease. *Annals of the New York Academy of Sciences*, *1124*, 1–38.
https://doi.org/10.1196/annals.1440.011

Cirrito, J. R., Kang, J.-E., Lee, J., Stewart, F. R., Verges, D. K., Silverio, L. M., … Holtzman, D. M. (2008).
Endocytosis is required for synaptic activity-dependent release of amyloid-beta in vivo. *Neuron*,
58(1), 42–51. https://doi.org/10.1016/j.neuron.2008.02.003

Condello, C., Schain, A., & Grutzendler, J. (2011). Multicolor time-stamp reveals the dynamics and
toxicity of amyloid deposition. *Scientific Reports*, *1*(1), 19. https://doi.org/10.1038/srep00019

Counts, S. E., Ikonomovic, M. D., Mercado, N., Vega, I. E., & Mufson, E. J. (2017). Biomarkers for the
Early Detection and Progression of Alzheimer's Disease. *Neurotherapeutics⃞: The Journal of the*
American Society for Experimental NeuroTherapeutics, *14*(1), 35–53.
https://doi.org/10.1007/s13311-016-0481-z

Diez, M., Koistinaho, J., Kahn, K., Games, D., & Hökfelt, T. (2000). Neuropeptides in hippocampus and
cortex in transgenic mice overexpressing V717F β-amyloid precursor protein — initial observations.
Neuroscience, *100*(2), 259–286. https://doi.org/10.1016/S0306-4522(00)00261-X

Dowson, J. H. (1981). A sensitive method for the demonstration of senile plaques in the dementing brain.
Histopathology, *5*(3), 305–310. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/7016712

Garcia-Alloza, M., Robbins, E. M., Zhang-Nunes, S. X., Purcell, S. M., Betensky, R. A., Raju, S., …
Frosch, M. P. (2006). Characterization of amyloid deposition in the APPswe/PS1dE9 mouse model
of Alzheimer disease. *Neurobiology of Disease*, *24*(3), 516–524.
https://doi.org/10.1016/j.nbd.2006.08.017

Grothe, M. J., Barthel, H., Sepulcre, J., Dyrba, M., Sabri, O., Teipel, S. J., & Alzheimer's Disease
Neuroimaging Initiative, F. the A. D. N. (2017). In vivo staging of regional amyloid deposition.
Neurology, *89*(20), 2031–2038. https://doi.org/10.1212/WNL.0000000000004643

Hafkemeijer, A., van der Grond, J., & Rombouts, S. A. R. B. (2012). Imaging the default mode network in
aging and dementia. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*, *1822*(3),
431–441. https://doi.org/10.1016/j.bbadis.2011.07.008

Hall, A. M., & Roberson, E. D. (2012). Mouse models of Alzheimer's disease. *Brain Research Bulletin*,
88(1), 3–12. https://doi.org/10.1016/j.brainresbull.2011.11.017

Harris, J. A., Mihalas, S., Hirokawa, K. E., Whitesell, J. D., Knox, J., Bernard, A., … Zeng, H. (2018). The
organization of intracortical connections by layer and cell class in the mouse brain. *BioRxiv*.
Retrieved from http://biorxiv.org/content/early/2018/04/18/292961.abstract

Höfling, C., Morawski, M., Zeitschel, U., Zanier, E. R., Moschke, K., Serdaroglu, A., … Kuhn, P.-H.
(2016). Differential transgene expression patterns in Alzheimer mouse models revealed by novel
human amyloid precursor protein-specific antibodies. *Aging Cell*, *15*(5), 953–963.
https://doi.org/10.1111/acel.12508

Hsiao, K., Chapman, P., Nilsen, S., Eckman, C., Harigaya, Y., Younkin, S., … Cole, G. (1996). Correlative
memory deficits, Aβ elevation, and amyloid plaques in transgenic mice. *Science.*
https://doi.org/10.1126/science.274.5284.99

Huang, H., Nie, S., Cao, M., Marshall, C., Gao, J., Xiao, N., … Xiao, M. (2016). Characterization of AD-
like phenotype in aged APPSwe/PS1dE9 mice. *Age*, *38*(4), 303–322.
https://doi.org/10.1007/s11357-016-9929-7

Huang, Y., Skwarek-Maruszewska, A., Horré, K., Vandewyer, E., Wolfs, L., Snellinx, A., … Thathiah, A.
(2015). Loss of GPR3 reduces the amyloid plaque burden and improves memory in Alzheimer's
disease mouse models. *Science Translational Medicine*, *7*(309), 309ra164-309ra164.
https://doi.org/10.1126/scitranslmed.aab3492

Jack, C. R., Bennett, D. A., Blennow, K., Carrillo, M. C., Dunn, B., Haeberlein, S. B., … Contributors, R.
(2018). NIA-AA Research Framework: Toward a biological definition of Alzheimer's disease.
Alzheimer's & Dementia: The Journal of the Alzheimer's Association, *14*(4), 535–562.
https://doi.org/10.1016/j.jalz.2018.02.018

Jack, C. R., Knopman, D. S., Jagust, W. J., Petersen, R. C., Weiner, M. W., Aisen, P. S., … Trojanowski,
J. Q. (2013). Tracking pathophysiological processes in Alzheimer's disease: an updated
hypothetical model of dynamic biomarkers. *The Lancet. Neurology*, *12*(2), 207–216.
https://doi.org/10.1016/S1474-4422(12)70291-0

Jährling, N., Becker, K., Wegenast-Braun, B. M., Grathwohl, S. A., Jucker, M., & Dodt, H.-U. (2015).
Cerebral β-Amyloidosis in Mice Investigated by Ultramicroscopy. *PloS One*, *10*(5), e0125418.
https://doi.org/10.1371/journal.pone.0125418

Jankowsky, J. L., Fadale, D. J., Anderson, J., Xu, G. M., Gonzales, V., Jenkins, N. A., … Borchelt, D. R.
(2004). Mutant presenilins specifically elevate the levels of the 42 residue ??-amyloid peptide in
vivo: Evidence for augmentation of a 42-specific ?? secretase. *Human Molecular Genetics*, *13*(2),
159–170. https://doi.org/10.1093/hmg/ddh019

Jankowsky, J. L., Slunt, H. H., Ratovitski, T., Jenkins, N. A., Copeland, N. G., & Borchelt, D. R. (2001).
Co-expression of multiple transgenes in mouse CNS: A comparison of strategies. *Biomolecular
Engineering*, *17*(6), 157–165. https://doi.org/10.1016/S1389-0344(01)00067-3

Jankowsky, J. L., & Zheng, H. (2017). Practical considerations for choosing a mouse model of
Alzheimer's disease. *Molecular Neurodegeneration*, *12*(1), 89. https://doi.org/10.1186/s13024-017-
0231-7

Jones, D. T., Knopman, D. S., Gunter, J. L., Graff-radford, J., Vemuri, P., Boeve, B. F., … Jack, C. R.
(2015). Cascading network failure across the Alzheimer's disease spectrum. *Brain*, 1–16.
https://doi.org/10.1093/brain/awv338

Kamenetz, F., Tomita, T., Hsieh, H., Seabrook, G., Borchelt, D., Iwatsubo, T., … Malinow, R. (2003). APP
processing and synaptic function. *Neuron*, *37*(6), 925–937. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/12670422

Kim, T.-K., Lee, J.-K. J.-Y. J.-E., Park, S.-K., Lee, K.-W., Seo, J.-S., Im, J.-Y., … Han, P.-L. (2012).
Analysis of differential plaque depositions in the brains of Tg2576 and Tg-APPswe/PS1dE9
transgenic mouse models of Alzheimer disease. *Experimental & Molecular Medicine*, *44*(8), 492–
502. https://doi.org/10.3858/emm.2012.44.8.056

Klunk, W. E., Bacskai, B. J., Mathis, C. A., Kajdasz, S. T., McLellan, M. E., Frosch, M. P., … Yman, B. R.
T. H. (2002). Imaging A β Plaques in Living Transgenic Mice with Multiphoton Microscopy and
Methoxy-X04 , a Systemically Administered Congo Red Derivative. *Journal of Neuropathology &*
Experimental Neurology, *61*(9), 797–805. https://doi.org/10.1093/jnen/61.9.797

Klunk, W. E., Lopresti, B. J., Ikonomovic, M. D., Lefterov, I. M., Koldamova, R. P., Abrahamson, E. E., …
Mathis, C. A. (2005). Binding of the Positron Emission Tomography Tracer Pittsburgh Compound-B
Reflects the Amount of Amyloid-in Alzheimer's Disease Brain But Not in Transgenic Mouse Brain.
https://doi.org/10.1523/JNEUROSCI.2990-05.2005

Kuan, L., Li, Y., Lau, C., Feng, D., Bernard, A., Sunkin, S. M., … Ng, L. (2015). Neuroinformatics of the
Allen Mouse Brain Connectivity Atlas. *Methods*, *73*, 4–17.
https://doi.org/10.1016/j.ymeth.2014.12.013

Kwan, A. C., Duff, K., Gouras, G. K., & Webb, W. W. (2009). Optical visualization of Alzheimer's
pathology via multiphoton-excited intrinsic fluorescence and second harmonic generation. *Optics*
Express, *17*(5), 3679–3689. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/19259208

Lee, E. B., Leng, L. Z., Zhang, B., Kwong, L., Trojanowski, J. Q., Abel, T., & Lee, V. M. Y. (2006).
Targeting amyloid-β peptide (Aβ) oligomers by passive immunization with a conformation-selective
monoclonal antibody improves learning and memory in Aβ precursor protein (APP) transgenic mice.
Journal of Biological Chemistry, *281*(7), 4292–4299. https://doi.org/10.1074/jbc.M511018200

Lee, J.-E., & Han, P.-L. (2013). An update of animal models of Alzheimer disease with a reevaluation of
plaque depositions. *Experimental Neurobiology*, *22*(2), 84–95.
https://doi.org/10.5607/en.2013.22.2.84

Lewis, J., & Dickson, D. W. (2016). Propagation of tau pathology: hypotheses, discoveries, and yet
unresolved questions from experimental and human brain studies. *Acta Neuropathologica*, *131*(1),
27–48. https://doi.org/10.1007/s00401-015-1507-z

Li, X., Uemura, K., Hashimoto, T., Nasser-Ghodsi, N., Arimon, M., Lill, C. M., … Berezovska, O. (2013).
Neuronal activity and secreted amyloid β lead to altered amyloid β precursor protein and presenilin 1
interactions. *Neurobiology of Disease*, *50*, 127–134. https://doi.org/10.1016/j.nbd.2012.10.002

Liebmann, T., Renier, N., Bettayeb, K., Greengard, P., Tessier-Lavigne, M., & Flajolet, M. (2016). Three-
Dimensional Study of Alzheimer's Disease Hallmarks Using the iDISCO Clearing Method. *Cell*
Reports, *16*(4), 1138–1152. https://doi.org/10.1016/J.CELREP.2016.06.060

Liu, P., Reichl, J. H., Rao, E. R., McNellis, B. M., Huang, E. S., Hemmy, L. S., … Zahs, K. R. (2017).
Quantitative Comparison of Dense-Core Amyloid Plaque Accumulation in Amyloid-β Protein
Precursor Transgenic Mice. *Journal of Alzheimer's Disease*, *56*(2), 743–761.
https://doi.org/10.3233/JAD-161027

Lu, H., Zou, Q., Gu, H., Raichle, M. E., Stein, E. A., & Yang, Y. (2012). Rat brains also have a default
mode network. *Proc Natl Acad Sci U S A*, *109*(10), 3979–3984.
https://doi.org/10.1073/pnas.1200506109

Mormino, E. C., Smiljic, A., Hayenga, A. O., Onami, S. H., Greicius, M. D., Rabinovici, G. D., … Jagust,
W. J. (2011). Relationships between β-amyloid and functional connectivity in different components
of the default mode network in aging. *Cerebral Cortex (New York, N.Y. : 1991)*, *21*(10), 2399–2407.
https://doi.org/10.1093/cercor/bhr025

Mucke, L., Masliah, E., Yu, G. Q., Mallory, M., Rockenstein, E. M., Tatsuno, G., … McConlogue, L.
(2000). High-level neuronal expression of abeta 1-42 in wild-type human amyloid protein precursor
transgenic mice: synaptotoxicity without plaque formation. *The Journal of Neuroscience : The*
Official Journal of the Society for Neuroscience, *20*(11), 4050–4058.

842    https://doi.org/10.1523/JNEUROSCI.20-11-04050.2000

843 Nelson, P. T., Alafuzoff, I., Bigio, E. H., Bouras, C., Braak, H., Cairns, N. J., … Beach, T. G. (2012).
844    Correlation of Alzheimer Disease Neuropathologic Changes With Cognitive Status: A Review of the
845    Literature. *Journal of Neuropathology & Experimental Neurology*, *71*(5), 362–381.
846    https://doi.org/10.1097/NEN.0b013e31825018f7

847 Oh, S. W., Harris, J. A., Ng, L., Winslow, B., Cain, N., Mihalas, S., … Zeng, H. (2014). A mesoscale
848    connectome of the mouse brain. *Nature*, *508*(7495), 207–214. https://doi.org/10.1038/nature13186

849 Palmqvist, S., Schöll, M., Strandberg, O., Mattsson, N., Stomrud, E., Zetterberg, H., … Hansson, O.
850    (2017). Earliest accumulation of β-amyloid occurs within the default-mode network and concurrently
851    affects brain connectivity. *Nature Communications*, *8*(1), 1214. https://doi.org/10.1038/s41467-017-
852    01150-x

853 Patton, R. L., Kalback, W. M., Esh, C. L., Kokjohn, T. A., Van Vickle, G. D., Luehrs, D. C., … Roher, A. E.
854    (2006). Amyloid-beta peptide remnants in AN-1792-immunized Alzheimer's disease patients: a
855    biochemical analysis. *The American Journal of Pathology*, *169*(3), 1048–1063.
856    https://doi.org/10.2353/AJPATH.2006.060269

857 Pozueta, J., Lefort, R., Ribe, E. M., Troy, C. M., Arancio, O., & Shelanski, M. (2013). Caspase-2 is
858    required for dendritic spine and behavioural alterations in J20 APP transgenic mice. *Nature
859    Communications*, *4*(1), 1939. https://doi.org/10.1038/ncomms2927

860 Ragan, T. M., Kadiri, L. R., Venkataraju, K. U., Bahlmann, K., Sutin, J., Taranda, J., … Osten, P. (2012).
861    Serial two-photon tomography for automated ex vivo mouse brain imaging. *Nature Methods*, *9*(3),
862    255–258. https://doi.org/10.1038/nmeth.1854

863 Raichle, M. E., MacLeod, A. M., Snyder, A. Z., Powers, W. J., Gusnard, D. A., & Shulman, G. L. (2001). A
864    default mode of brain function. *Proceedings of the National Academy of Sciences of the United
865    States of America*, *98*(2), 676–682. https://doi.org/10.1073/pnas.98.2.676

866 Razali, N. M., & Wah, Y. B. (2011). Power comparisons of shapiro-wilk, kolmogorov-smirnov, lilliefors and
867    anderson-darling tests. *Journal of Statistical Modeling and Analytics*. Retrieved from
868    https://www.researchgate.net/profile/Bee_Yap/publication/267205556_Power_Comparisons_of_Sha
869    piro-Wilk_Kolmogorov-Smirnov_Lilliefors_and_Anderson-
870    Darling_Tests/links/5477245b0cf29afed61446e1/Power-Comparisons-of-Shapiro-Wilk-Kolmogorov-
871    Smirnov-Lilliefors-an

872 Rice, L., & Bisdas, S. (2017). The diagnostic value of FDG and amyloid PET in Alzheimer's disease—A
873    systematic review. *European Journal of Radiology*, *94*, 16–24.
874    https://doi.org/10.1016/J.EJRAD.2017.07.014

875 Samaroo, H. D., Opsahl, A. C., Schreiber, J., O'Neill, S. M., Marconi, M., Qian, J., … Stephenson, D. T.
876    (2012). High throughput object-based image analysis of β-amyloid plaques in human and transgenic
877    mouse brain. *Journal of Neuroscience Methods*, *204*(1), 179–188.
878    https://doi.org/10.1016/J.JNEUMETH.2011.10.003

879 Sasaguri, H., Nilsson, P., Hashimoto, S., Nagata, K., Saito, T., De Strooper, B., … Saido, T. C. (2017).
880    APP mouse models for Alzheimer's disease preclinical studies. *The EMBO Journal*, *36*(17), 2473–
881    2487. https://doi.org/10.15252/embj.201797397

882 Schilling, L. P., Zimmer, E. R., Shin, M., Leuzy, A., Pascoal, T. A., Benedet, A. L., … Rosa-Neto, P.
883    (2016). Imaging Alzheimer's disease pathophysiology with PET. *Dementia & Neuropsychologia*,
884    *10*(2), 79–90. https://doi.org/10.1590/S1980-5764-2016DN1002003

885 Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., … Cardona, A.
886    (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682.

https://doi.org/10.1038/nmeth.2019

Sforazzini, F., Schwarz, A. J., Galbusera, A., Bifone, A., & Gozzi, A. (2014). Distributed BOLD and CBV-
weighted resting-state networks in the mouse brain. *NeuroImage*, *87*, 403–415.
https://doi.org/10.1016/j.neuroimage.2013.09.050

Snellman, A., López-Picón, F. R., Rokka, J., Salmona, M., Forloni, G., Scheinin, M., … Haaparanta-Solin,
M. (2013). Longitudinal amyloid imaging in mouse brain with 11C-PIB: comparison of APP23,
Tg2576, and APPswe-PS1dE9 mouse models of Alzheimer disease. *Journal of Nuclear Medicine⍰:*
Official Publication, Society of Nuclear Medicine, *54*(8), 1434–1441.
https://doi.org/10.2967/jnumed.112.110163

Stafford, J. M., Jarrett, B. R., Miranda-Dominguez, O., Mills, B. D., Cain, N., Mihalas, S., … Fair, D. A.
(2014). Large-scale topology and the default mode network in the mouse connectome. *Proceedings*
of the National Academy of Sciences, *111*(52), 201404346.
https://doi.org/10.1073/pnas.1404346111

Thakker, D. R., Weatherspoon, M. R., Harrison, J., Keene, T. E., Lane, D. S., Kaemmerer, W. F., …
Shafer, L. L. (2009). Intracerebroventricular amyloid-beta antibodies reduce cerebral amyloid
angiopathy and associated micro-hemorrhages in aged Tg2576 mice. *Proceedings of the National*
Academy of Sciences of the United States of America, *106*(11), 4501–4506.
https://doi.org/10.1073/pnas.0813404106

Thal, D. R., Ghebremedhin, E., Haass, C., & Schultz, C. (2002). UV light-induced autofluorescence of full-
length Abeta-protein deposits in the human brain. *Clinical Neuropathology*, *21*(1), 35–40. Retrieved
from http://www.ncbi.nlm.nih.gov/pubmed/11846043

Thal, D. R., Rüb, U., Orantes, M., & Braak, H. (2002). Phases of A beta-deposition in the human brain
and its relevance for the development of AD. *Neurology*, *58*(12), 1791–1800.
https://doi.org/10.1212/WNL.58.12.1791

Wright, A. L., Zinn, R., Hohensinn, B., Konen, L. M., Beynon, S. B., Tan, R. P., … Vissel, B. (2013).
Neuroinflammation and Neuronal Loss Precede Aβ Plaque Deposition in the hAPP-J20 Mouse
Model of Alzheimer's Disease. *PLoS ONE*, *8*(4), e59586.
https://doi.org/10.1371/journal.pone.0059586

Yushkevich, P. A., Piven, J., Hazlett, H. C., Smith, R. G., Ho, S., Gee, J. C., & Gerig, G. (2006). User-
guided 3D active contour segmentation of anatomical structures: Significantly improved efficiency
and reliability. *NeuroImage*, *31*(3), 1116–1128. https://doi.org/10.1016/j.neuroimage.2006.01.015

Zerbi, V., Grandjean, J., Rudin, M., & Wenderoth, N. (2015). Mapping the mouse brain with rs-fMRI: An
optimized pipeline for functional network identification. *NeuroImage*, *123*, 11–21.
https://doi.org/10.1016/j.neuroimage.2015.07.090

Zhang, H., Zhang, L., Zhou, D., He, X., Wang, D., Pan, H., … Sun, B. (2017). Ablating ErbB4 in PV
neurons attenuates synaptic and cognitive deficits in an animal model of Alzheimer's disease.
Neurobiology of Disease, *106*, 171–180. https://doi.org/10.1016/J.NBD.2017.07.001

Zipfel, W. R., Williams, R. M., Christie, R., Nikitin, A. Y., Hyman, B. T., & Webb, W. W. (2003). Live tissue
intrinsic emission microscopy using multiphoton-excited native fluorescence and second harmonic
generation. *Proceedings of the National Academy of Sciences of the United States of America*,
100(12), 7075–7080. https://doi.org/10.1073/pnas.0832308100

Tables

Table 1. Experimental Dataset

Mouse Line	Promoter	5 mo (M/F)	7 mo (M/F)	9 mo (M/F)	13 mo (M/F)	19 mo (M/F)
APP/PS1 (APP$^{+/-}$)	moPrP	6 (4/2)	13 (9/4)	6 (3/3)	39 (36/3)	9 (7/2)
hAPP-J20 (APP$^{+/-}$)	PDGF-β	4 (2/2)	3 (2/1)	5 (2/3)	11 (6/5)	3 (0/3)
Tg2576 (APP$^{+/-}$)	haPrP			3 (0/3)	5 (0/5)	4 (0/4)
Controls						
APP/PS1 (APP$^{-/-}$)				2 (2/0)	11 (7/4)	2 (0/2)
hAPP-J20 (APP$^{-/-}$)			2 (1/1)		17 (9/8)	1 (0/1)

930
Transgenic mouse lines used in this study and the number of experimental animals in each group.
Numbers in parentheses indicate the number of males and females in each mouse line/age group
combination. The promoter used to drive APP expression for each transgenic line is indicated in the table.
Control mice were APP$^{-/-}$ littermates of experimental mice.

Table 2. Abbreviations and full structure names for all annotated isocortical, hippocampal, cortical
subplate, and olfactory areas in the Allen Mouse Brain Common Coordinate Framework (CCF) at the
summary structure level.

Abbreviation	Full structure name
Isocortex	
FRP	Frontal pole, cerebral cortex
MOs	Secondary motor area
ACAd	Anterior cingulate area, dorsal part
ACAv	Anterior cingulate area, ventral part
PL	Prelimbic area
ILA	Infralimbic area
ORBl	Orbital area, lateral part
ORBm	Orbital area, medial part
ORBvl	Orbital area, ventrolateral part
AId	Agranular insular area, dorsal part
AIv	Agranular insular area, ventral part
AIp	Agranular insular area, posterior part
GU	Gustatory areas
VISC	Visceral area
SSs	Supplemental somatosensory area
SSp-bfd	Primary somatosensory area, barrel field
SSp-tr	Primary somatosensory area, trunk
SSp-ll	Primary somatosensory area, lower limb
SSp-ul	Primary somatosensory area, upper limb
SSp-un	Primary somatosensory area, unassigned
SSp-n	Primary somatosensory area, nose
SSp-m	Primary somatosensory area, mouth
MOp	Primary motor area
VISal	Anterolateral visual area
VISl	Lateral visual area
VISp	Primary visual area
VISpl	Posterolateral visual area
VISli	Laterointermediate area
VISpor	Postrhinal area
VISrl	Rostrolateral visual area
VISa	Anterior area
VISam	Anteromedial visual area
VISpm	posteromedial visual area

RSPagl	Retrosplenial area, lateral agranular part
RSPd	Retrosplenial area, dorsal part
RSPv	Retrosplenial area, ventral part
AUDd	Dorsal auditory area
AUDp	Primary auditory area
AUDpo	Posterior auditory area
AUDv	Ventral auditory area
TEa	Temporal association areas
ECT	Ectorhinal area
Hippocampal Formation	
CA1	Field CA1
CA2	Field CA2
CA3	Field CA3
DG	Dentate gyrus
FC	Fasciola cinerea
IG	Induseum griseum
ENTl	Entorhinal area, lateral part
ENTm	Entorhinal area, medial part, dorsal zone
PAR	Parasubiculum
POST	Postsubiculum
PRE	Presubiculum
SUB	Subiculum
ProS	Prosubiculum
HATA	Hippocampo-amygdalar transition area
APr	Area prostriata
Olfactory Areas	
MOB	Main olfactory bulb
AOB	Accessory olfactory bulb
AON	Anterior olfactory nucleus
TT	Taenia tecta
DP	Dorsal peduncular area
PIR	Piriform area
NLOT	Nucleus of the lateral olfactory tract
COAa	Cortical amygdalar area, anterior part
COAp	Cortical amygdalar area, posterior part
PAA	Piriform-amygdalar area
TR	Postpiriform transition area
Cortical Subplate	
CLA	Claustrum
EPd	Endopiriform nucleus, dorsal part
EPv	Endopiriform nucleus, ventral part
LA	Lateral amygdalar nucleus
BLA	Basolateral amygdalar nucleus
BMA	Basomedial amygdalar nucleus
PA	Posterior amygdalar nucleus

Figure Legends

Figure 1. Pipeline workflow for labeling and quantification of brain-wide plaque distribution. (a)
Mice were injected with methoxy-X04 to label plaques *in vivo* 24 hours before perfusion. Brains were
imaged with serial 2-photon tomography which involves embedding the brain in agar (1), acquisition of 2-
photon images in the coronal plane (2), and sectioning with an integrated vibratome (3). Images were
processed in an automated informatics pipeline that included automated detection of plaques from

section images and deformable 3D registration to the Allen CCFv3. Plaque density and count were then
quantified for all structures. **(b)** Selected images from coronal STP image series spanning the brain from
cerebellum to olfactory bulb are shown for a 19-month-old mouse with methoxy-X04 labeling from each of
the three APP mouse lines used in this study: APP/PS1, Tg2576, and hAPP-J20. Images in the furthest
right column show similar sections from a 6-month-old APP/PS1$^{+/-}$ mouse that was not injected with
methoxy-X04. Some lower intensity plaque fluorescence is visible without methoxy-X04 labeling. Arrow
indicates methoxy-X04 labeled plaques in the subiculum of Tg2576 mice. Arrowheads show examples of
bright tissue edges that can cause false positives in automated segmentation.

Figure 2. Automated segmentation of plaque fluorescence. (a) An example ROI from a single section
image in somatosensory cortex, barrel field area (SSp-bfd) showing methoxy-X04 fluorescence. A portion
of the hippocampus is also visible in the bottom left corner of the image. **(b)** Segmented plaque signal
(red) overlaid on the image section shown in (a). **(c)** An ROI from a different section in the same brain
showing the cerebral aqueduct (AQ) and surrounding periaqueductal gray (PAG). **(d)** Segmented plaque
signal for the image section shown in c (red) showing false positive signal along the bright tissue edge at
the border of the cerebral aqueduct. Similar signal was observed in APP$^{-/-}$ mice. Scale = 1 mm. **(e-i)**
Quantification of false positives as percent of structure volume for 35 image series collected from APP$^{-/-}$
mice without plaques to identify regions prone to segmentation artifacts. False positives are plotted for
summary structures in hippocampus **(e)**, selected midbrain regions including PAG **(f)**, olfactory areas **(g)**,
cortical subplate **(h)**, and isocortex **(i)**. Red points indicate APP$^{-/-}$ mice that received a methoxy-X04
injection one day before perfusion. Abbreviations for summary structures in isocortex, hippocampus,
olfactory areas, and cortical subplate can be found in **Table 2**. Abbreviations in (f) SNr: Substantia nigra,
reticular part; SCm: Superior colliculus, motor related; PAG: Periaqueductal gray; RN: Red nucleus. **(j)**
Histogram showing the difference between signal and false positive levels (% coverage in APP$^{+/-}$ mice - %
coverage in APP$^{-/-}$ mice) for every structure in the Allen CCFv3 reference atlas. **(k)** False positive
heatmap showing regions with the highest segmentation artifacts. Abbreviations: BLAa: Basolateral
amygdalar nucleus, anterior part; IG: Induseum griseum; AD: Anterodorsal nucleus of the thalamus; CEA:
Central amygdalar nucleus; ORBm: Medial orbital cortex; SFO: Subfornical organ; MH: Medial habenula.
(l) False positive heatmap for a different anatomical location. Abbreviations: CEAm: Central amygdalar
nucleus, medial part; TTv: Taenia tecta, ventral part; CA2: Hippocampal field CA2.

Figure 3. Automatic plaque counts reveal large, vascular-associated amyloid deposits. Portion of a
single coronal **(a,b)** and sagittal **(c,d)** image from the CCFv3 template with gridded, automatically
detected methoxy-X04 labeled plaques from a 19-month-old APP/PS1 mouse brain overlaid in white. In
(b) and **(d)**, the same images are shown with automatically counted plaques in labeled with different
colors. Plaques in the same area that are the same color had diagonally adjacent voxels in at least one
image section and were counted as one single object. The sagittal view in (d) shows that plaques are
elongated on the z-axis. **(e)** Sagittal image from the CCFv3 template with gridded plaques from a 19-
month-old hAPP-J20 mouse with automatic plaque count results overlaid. Nearly all of the plaques along
the midline in this brain were counted as a single plaque (blue). **(f)** Sagittal CCFv3 template image with
gridded plaques and counts overlaid for a 19-month-old Tg2576 mouse reveals large, vascular-
associated deposits.

Figure 4. Methoxy-X04 accurately labels the location of plaques but underestimates plaque
density. (a-c) Automated segmentation results **(a)**, IHC for Aβ (6E10 antibody) **(b)**, and an overlay of the
automated segmentation and thresholded IHC image from the same section **(c)** are shown for one
section from each mouse line. APP/PS1 and hAPP-J20 brains are from 13-month-old mice. The Tg2576
brain is from a 19-month-old mouse. In the segmentation images in (a), red indicates segmented plaques,
and gray the putative plaque signal that was removed by the adaptive filtering step in the segmentation
algorithm. Automatically detected tissue edges are also colored gray. Image alignment in the overlay
images **(c)** is imperfect due to tissue deformation during antibody labeling. **(d)** maximum intensity
projection of a confocal image stack through the center of one plaque for each mouse line. Upper
grayscale image shows 6E10 labeling; bottom color image shows 6E10 labeling overlaid with methoxy-
X04 and DAPI fluorescence (both in cyan). Arrowheads indicate the methoxy-X04 positive core of each
plaque. Scale = 500 μm (a-c), 50 μm (d). **(e)** Aβ antibody-labeled plaque density and segmented
methoxy-X04-labeled plaque density plotted for each ROI measured (log scale). Points are colored by the

Allen CCFv3 reference atlas region in which each ROI was drawn (isocortex, hippocampal formation,
cortical subplate, olfactory areas, striatum, and thalamus). Inset shows a coronal section from the Allen
CCFv3 Reference Atlas with region colors. The shape of each point indicates which mouse line it belongs
to: circles for APP/PS1, squares for Tg2576, and triangles for hAPP-J20. Methoxy-X04 density was lower
than $A\beta$ antibody-labeled density for every point, but the two measurements were highly correlated
(Pearson's r = 0.95, p<0.0001). **(f)** The fraction of $A\beta$ antibody-labeled plaque density that was detected
by automated segmentation was significantly lower for hAPP-J20 mice compared to APP/PS1 mice. **(g)**
For individual plaques, the ratio of methoxy-X04 labeling to $A\beta$ antibody labeling is plotted for each mouse
line. hAPP-J20 mice had significantly lower fractions of methoxy-X04/$A\beta$ antibody labeling.

Figure 5. Automated quantification of plaque density and count across whole brain and major
divisions for three APP mouse lines at various ages. (a,b) Plaque density measured by automated
segmentation and registration plotted as count (plaques per mm^3; **a**) or density (% of structure volume; **b-**
g). Plaque density is plotted by age for the whole brain **(a,b)**, isocortex **(c)**, and hippocampal formation **(d)**
in each of the three APP lines characterized. Dark circles indicate the median values and light points
individual animals. Lines connect points at the median. Plaque density at 13 months **(e)** and 19 months
(g) divided by major brain division is plotted separately for each mouse line. Box plots show the median
and IQR, with whiskers extending to 1.5 times the IQR. Outliers are plotted as individual points beyond
the whiskers. Dashed vertical line in the APP/PS1 plots in (e) and (g) show the maximum value from the
hAPP-J20 and Tg2576 mouse line plots below at the same age. Plots in (e) and (g) are colored by major
brain division, and the cartoons in (c), (d), and (f) show the anatomical locations of the major brain
divisions in the Allen CCFv3 reference atlas. Abbreviations: OLF: Olfactory areas, HPF: Hippocampal
formation, CTXsp: Cortical subplate, STR: Striatum, PAL: Pallidum, TH: Thalamus, HY: Hypothalamus,
MB: Midbrain, P: Pons, MY: Medulla, CB: Cerbellum.

Figure 6. Quantification of plaque density for selected summary structures in the isocortex,
hippocampus, olfactory areas, and cortical subplate. Plaque density (percent of structure volume) for
summary structures in **(a)** isocortex, **(c)** hippocampal formation**,** **(e)** olfactory areas, and **(g)** cortical
subplate in 13-month-old (left) and 19-month-old (right) mice. Darker lines connect points at the median
and lighter error bars indicate the interquartile range. Dashed vertical lines indicate the median value for
each mouse line for all graphed structures in each plot. Plots for plaque density at 13 months have split y-
axes to allow all three datasets to be plotted on one graph. Asterisks in (a) indicate structures that were
significantly higher or lower than the median plaque density in isocortex for APP/PS1 and hAPP-J20
mice. **(b)** Representative images showing plaque deposition patterns in each of the three APP mouse
lines for two isocortex structures: lateral visual cortex (VISl, top) and retrosplenial cortex (RSP, bottom).
(d) Images showing plaque deposition patterns in the hippocampus proper for the three mouse lines. The
location of subiculum (SUB), CA1, CA3, and dentate gyrus (DG) are indicated on the hAPP-J20 image. **(f)**
Images showing plaque deposition patterns in the piriform (olfactory) cortex for the three mouse lines.
The hAPP-J20 image shows one of only a few plaques observed in this region in hAPP-J20 mice. **(h)**
Images showing plaque deposition patterns in different amygdalar structures in the cortical subplate for
the three mouse lines. The approximate location of the central amygdala (CEA), medial amygdala (MEA),
basolateral amygdala (BLA), and piriform cortex (PIR) is indicated on the hAPP-J20 image. All
abbreviations used in the graphs can be found in **Table 2**. Scale = 500 µm.

Figure 7. APP/PS1 and Tg2576 mice have plaques in the cerebellum but hAPP-J20 mice do not.
Single STP image planes in the cerebellum (top row) and higher magnification views of the region
indicated with the white box (bottom row). All images are from 19-month-old mouse brains. Scale = 1 mm.

Figure 8. Plaques are unequally distributed across structural modules in the isocortex. (a). Plaque
density in the isocortex modules from (Harris et al., 2018) relative to the density that would be expected
with even plaque distribution across all cortical regions (dashed gray line). Error bars indicate the 95%
confidence interval. **(b).** Illustrations of the Allen CCFv3 with isocortex structures colored by the relative
plaque density of their corresponding module in each mouse line.

Figure 9. Whole brain 3D heatmaps showing the anatomical distribution of plaques. Single coronal
(left), sagittal (middle), and horizontal (right) planar views from the associated full 3D map for each age-

mouse line combination. **(a)** Sections from the Allen CCFv3 reference atlas showing structure annotation
at the summary structure level. **(b-d)** Single sections corresponding to the reference atlas plates in the top
row colored by plaque density. APP/PS1 **(b)** and hAPP-J20 **(c)** maps for 13-month-old animals are
shown, and the Tg2576 **(d)** map is for 19-month-old animals. Each plaque map has its own scaled
colormap (indicated in color bar on right) since absolute plaque levels are variable between ages and
mouse lines. The full 3D plaque maps for all ages and mouse lines are available to download as nrrd files
online at [link to be provided]. Abbreviations: LA: Lateral amygdalar nuclus, DG: dentate gyrus, VISrl:
rostrolateral visual area, BLAv: Basolateral amygdalar nucleus, ventral part, ENTl: lateral entorhinal
cortex, RSPv: ventral retrosplenial cortex, ENTm: medial entorhinal cortex, CEA: Central amygdalar
nucleus, SUB: subiculum, BLA: Basolateral amygdala, PIR: Piriform cortex.

Figure 10. Whole brain 3D heatmaps showing the anatomical distribution of plaques at other
locations. (Single coronal (left), sagittal (middle), and horizontal (right) planar views from the associated
full 3D map for each age-mouse line combination. **(a)** Sections from the Allen CCFv3 reference atlas
showing structure annotation at the summary structure level. **(b-d)** Single sections corresponding to the
reference atlas plates in the top row colored by plaque density. APP/PS1 **(b)** and hAPP-J20 **(c)** maps for
13-month-old animals are shown, and the Tg2576 **(d)** map is for 19-month-old animals. Each plaque map
has its own scaled colormap (indicated in color bar on right) since absolute plaque levels are variable
between ages and mouse lines. The full 3D plaque maps for all ages and mouse lines are available to
download as nrrd files online at [link to be provided]. Abbreviations: VISrl: rostrolateral visual area, ACA:
anterior cingulate cortex, ENTm: medial entorhinal cortex, RSPv: ventral retrosplenial cortex, DG: dentate
gyrus, SUBd: dorsal subiculum, TR: Postpiriform transition area.

Figure 11. Plaque distribution across cortical layers differs between mouse lines. Images showing
plaques in the parietal cortex of 19-month-old APP/PS1 **(a)**, Tg2576 **(b)**, and hAPP-J20 **(c)** mice.
Approximate layer boundaries are indicated in text to the right of each image (wm = white matter). Scale
= 500 µm. The relative plaque density in each cortical layer across the entire isocortex is plotted
separately for APP/PS1 **(b)**, Tg2576 **(d)**, and hAPP-J20 **(f)** mouse lines. Box plots show median and IQR
with whiskers extending up to 1.5 times the IQR. Outliers are plotted as individual points.

Figure 12. Brain-wide spatial-temporal patterns of plaque deposition in APP/PS1 and Tg2576 mice
is most similar to the pattern seen in human patients with AD progression. (a) Comparison of
relative plaque density in similar structures between human autopsy tissue and APP mouse models.
Human brain regions where Aβ pathology was quantified by Thal et al. (2012) are listed in the left column
and the corresponding region(s) from the Allen CCFv3 reference atlas are listed in the second column.
Plaque density (or median plaque density where there are multiple regions) for each age group and
mouse line is indicated by the heatmap in the columns to the right. The colormap spans from the 10th
percentile to the 90th percentile of the plaque density for **all** structures at all ages in that mouse line. **(b)**
10%, 50%, and 95% values for each mouse line. **(c)** Similarity measured using the Pearson correlation
coefficient for comparisons between plaque density in each mouse line and age group with the fraction of
patients showing plaques in each region during the five Phases of Aβ deposition in patients.
Abbreviations: ENTl: Entorhinal area, lateral part; ENTm: Entorhinal area, medial part, dorsal zone; ACA:
Anterior cingulate area; RSP: Retrosplenial area; LA: Lateral Amygdalar nucleus; BLA: Basolateral
amygdalar nucleus; BMA: Basomedial amygdalar nucleus; PA: Posterior amygdalar nucleus; CEA:
Central amygdalar nucleus; MEA: Medial amygdalar nucleus; COA: Cortical amygdalar area; DG: Dentate
gyrus; PRE: Presubiculum; POST: Postsubiculum; TH: Thalamus; CP: Caudoputamen; ACB: Nucleus
accumbens; HY: Hypothalamus; SI: Substantia innominata; MA: Magnocellular nucleus; NDB: Diagonal
band nucleus; PAG: Periaqueductal gray; SCs: Superior colliculus, sensory related; SCm: Superior
colliculus, motor related, RN: Red nucleus; IO: Inferior olivary complex; Substantia nigra, reticular part;
SNc: Substantia nigra, compact part; GRN: Gigantocellular reticular nucleus; IRN: Intermediate reticular
nucleus; PARN: Parvicellular reticular nucleus; CBX: Cerebellar cortex; PRNr: Pontine reticular nucleus;
PRNc: Pontine reticular nucleus, caudal part; RAmb: Midbrian raphe nuclei; LC: Locus ceruleus; PB:
Parabrachial nucleus; DN: Dentate nucleus.

Movie 1. Plaque maps for 13-month-old APP/PS1, 19-month-old Tg2576, and 13-month-old hAPP-J20
mouse lines. Each frame of the movie shows a coronal section from the Allen CCFv3 reference atlas with
structures colored by plaque density (percent of structure volume occupied by plaque).

Appendix 1. False positive values for all structures in the Allen CCFv3 reference atlas. False
positive plaque signal detected in APP$^{-/-}$ control brains by our automated informatics pipeline. For each
structure annotated in the Allen CCFv3 reference atlas, we list the structure abbreviation, unique structure
id, plaque volume, structure volume, plaque density (plaque volume/structure volume), and plaque count.

Appendix 2. Plaque volume, density, and count for all structures in all animals. For each animal in
our dataset, this table lists the unique image series id, mouse line, age at death, age group, sex. For all
840 structures in the Allen CCFv3, we report the structure acronym, plaque volume, structure volume,
plaque density (plaque volume/structure volume), and plaque count.

1110

Figure 1. Pipeline workflow for labeling and quantification of brain-wide plaque distribution

a.

methoxy-X04 injection

hAPP-J20
Tg2576
APP/PS1

Serial 2-Photon Imaging

Automated Informatics Processing

Quantification by Structure

b.

APP/PS1   Tg2576   hAPP-J20   APP/PS1 No Methoxy-X04

Figure 2. Automated Segmentation of plaque fluorescence

Figure 3. Automatic plaque counts reveal large, vascular-associated amyloid deposits.

Figure 4. Methoxy-X04 accurately labels the location of plaques but underestimates plaque density

a. Automated Segmentation   b. Anti Aβ IHC (6E10)   c. Segmentation /Thresh IHC   d. Confocal MIP

APP/PS1

Tg2576

hAPP-J20

Anti-Aβ (6E10)
Anti-Aβ (6E10) Methoxy-X04/DAPI
Anti-Ab (6E10)
Anti-Aβ (6E10) Methoxy-X04/DAPI
Anti-Aβ (6E10)
Anti-Aβ (6E10) Methoxy-X04/DAPI

e. Segmented methoxy-X04 plaque density is correlated with antibody-labeled plaque density

Log10 methoxy-X04 plaque density

10-fold
30-fold
r = 0.95

○ APP/PS1
□ Tg2576
△ hAPP-J20

Log10 Aβ antibody-labeled plaque density

f. Fraction of antibody-stained area detected with automated segmentation

methoxy-X04 / Aβ antibody area detected (whole ROI)

p=0.04
ns

APP/PS1   Tg2576   hAPP-J20

g. Ratio of dense core / diffuse plaques for each APP mouse line

methoxy-X04 / Aβ antibody ratio per plaque

p=0.002
p=0.006

APP/PS1   Tg2576   hAPP-J20

Figure 5. Automated quantification of plaque density and count whole brain and major divisions for three APP mouse lines at various ages.

Figure 6. Quantification of plaque density for selected summary structures in the isocortex, hippocampus, olfactory areas, and cortical subplate.

Figure 7. APP/PS1 and Tg2576 mice have plaques in the cerebellum but hAPP-J20 mice do not.

Figure 8. Plaques are unequally distributed across structural modules in the isocortex.

Figure 9. Heatmaps showing the anatomical distribution of plaques in three APP mouse lines.

Figure 10. Heatmaps showing the anatomical distribution of plaques in three APP mouse lines.

Figure 11. Plaque distribution across cortical layers differs between mouse lines

Figure 12. Brain-wide temporal patterns of plaque distribution in APP/PS1 and Tg2576 mice are similar to the pattern of Aβ deposition in human AD brains.

a.

Human region from Thal et al. 2002	Allen CCFv3 region
Phase 1 Neocortex	Isocortex
Phase 2 CA1	CA1
Entorhinal Region	ENTl, ENTm
Gyrus Cinguli	ACA, RSP
Amygdala	LA, BLA, BMA, PA, CEA, MEA, COA
Fascia Dentata	DG
Phase 3 Presubiculum	PRE, POST
Thalamus	TH
Striatum	CP, ACB
Hypothalamus	HY
Basal Forebrain Nuclei (Meynert)	SI, MA, NDB
CA4	Not annotated
Central Gray	PAG
Phase 4 Superior Collicle	SCs, SCm
Red Nucleus	RN
Inferior Olivary Nucleus	IO
Substantia Nigra	SNr, SNc
Reticular Formation of the Medulla Oblongata	GRN, IRN, PARN
Cerebellar Molecular Layer	CBX
Reticular Formation of the Pons	PRNr, PRNc
Anterior and Central Raphe nuclei	RAmb
Phase 5 Locus coeruleus	LC
Parabrachial Nuclei	PB
Reticulo Tegmental Nucleus (Bechterew)	TRN
Dorsal Tegmental Nucleus (Gudden)	DTN
Nuclei Pontis	PG
Cerebellar Granule Cell Layer	Not annotated
no deposits Dentate Nucleus	DN

b.

Plaque density in all structures			
percentile	APP/PS1	Tg2576	J20
90%	0.523	0.056	0.029
50%	0.001	0.001	0.000
10%	0.000	0.000	0.000

c.

	APP/PS1					Tg2576			hAPP-J20				
	5	7	9	13	19	9	13	19	5	7	9	13	19
Phase 1	0.45	0.40	0.44	0.42	0.34	0.03	0.34	0.50	0.04	0.08	0.39	0.33	0.46
Phase 2	0.87	0.81	0.80	0.81	0.75	0.02	0.71	0.86	0.12	0.02	0.09	0.08	0.18
Phase 3	0.63	0.63	0.69	0.71	0.72	0.11	0.60	0.68	0.29	0.01	-0.12	-0.07	0.04
Phase 4	0.46	0.42	0.48	0.50	0.51	0.00	0.38	0.50	0.38	0.07	0.01	0.17	0.22
Phase 5	0.33	0.37	0.40	0.42	0.47	0.24	0.46	0.41	0.18	0.09	0.10	0.25	0.28

-0.2 -0.1 0 0.1 0.2 0.3 0.4 0.5 0.6 0.7 0.8 0.9 1

Pearson r