1 **Compression-Assisted Arthrocentesis of the Knee as a Quality Improvement**
2 **Intervention**
3

James F. Bennett[1], Wilmer L. Sibbitt, Jr. [1], Philip A. Band[2], Sabeen Yaqub[1], N.

Suzanne Emil[1], Monthida Fangtham[1], Roderick A. Fields[1], William A. Hayward[3],

Selma D. Kettwich[1], Arthur D. Bankhurst[1].

From

[1]Department of Internal Medicine, Division of Rheumatology and School of Medicine,

MSC 10 5550, 5[th] FL ACC, University of New Mexico Health Sciences Center,

Albuquerque, NM, USA 87131. WSibbitt@salud.unm.edu

[2]Department of Orthopaedic Surgery, Biochemistry & Molecular Pharmacology

NYU School of Medicine, New York City, NY, USA. Email: Philip.Band@nyumc.org

[3] The Department of Exercise and Sport Sciences, New Mexico Highlands University,

Las Vegas, NM, USA hayward1962@hotmail.com

[4]338 Amherst Dr. NE, Albuquerque, NM, 87106 Email: selmakettwich@gmail.com

[5]Medical Student, School of Medicine, University of New Mexico Health Sciences

Center, Albuquerque, NM, USA 87131. Email: LRoldan@salud.unm.edu

Corresponding Author:

Wilmer L. Sibbitt, Jr., MD, MSC 10 5550, 5[th] FL ACC, Department of Internal Medicine,

Division of Rheumatology, University of New Mexico Health Sciences Center,

Albuquerque, New Mexico, USA 87131, tele 505 272-4762, email:

WSibbitt@salud.unm.edu

Compliance with Ethical Standards:

Disclosure of Potential Conflicts of Interest: None of authors have a conflict of

interest in this manuscript. All products used are commercially available and the authors

have no interest in any of the manufacturers, products, or distributors.

Research involving human participants and/or animals: The institutional review

board (the Human Research Review Committee) has approved the above study and has

permitted us to perform data analysis and manuscript submission. No animal studies were

performed.

Informed Consent: The subjects gave informed consent to participate prior to all

interventions and study was approved by the institutional review board. All studies were

carried out in accordance with the World Medical Association Declaration of Helsinki

(JBJS 79A:1089-98,1997). Patient confidentiality was protected according to the U.S.

Health Insurance Portability and Accountability Act (HIPAA) and all data has been de-

identified.

Funding and Grants: There was no funding for this study.

Clinical Trial: This was not a clinical trial. This was a report of continuous quality

improvement using bioengineering advances.

ABSTRACT

Objective: The present study reports the introduction of mechanical compression of the

knee for arthrocentesis as quality improvement intervention in a procedure clinic.

Methods: 430 consecutive symptomatic osteoarthritic knees underwent arthrocentesis

followed by corticosteroid injection (1mg/kg of triamcinolone acetonide). The first 215

consecutive knees underwent conventional arthrocentesis and injection; the quality

intervention of a mechanical compression brace was introduced, and the next 215

consecutive knees underwent mechanical compression-assisted arthrocentesis follow by

injection. Pain scores, arthrocentesis success, fluid yield, time-to-next-intervention,

injections/year, and medical costs were measured.

Results: No serious adverse events occurred in 430 subjects. Diagnostic synovial fluid

($\geq$2 ml) was obtained in 9.3% (20/215) without compression and 40.9% (88/215) with

compression (p=0.00001, z for 95% CI= 1.96, Pierson). Mechanical compression was

associated with a 231% increase in mean arthrocentesis volume: compression 5.3$\pm$11.2

ml, conventional 1.6$\pm$6.4 ml (CI of difference 2.0 <3.7< 5.4; p=0.00001). Time-to-next-

intervention after compression-assisted arthrocentesis was longer: 6.9$\pm$3.5 months

compared to conventional: 5.1$\pm$2.7 months (p<0.00001, 95% CI of difference 1.2 <1.8<

2.3). Mechanical compression was associated with a reduction in the number of

corticosteroid injections administered per year: mechanical compression: 1.7$\pm$0.9

injections/year; conventional: 2.4$\pm$0.5 injections/year (p<0.00001, 95% CI of difference -

0.83 <-0.70< -0.56). Mechanical compression did not increase overall yearly costs

associated with management of the symptomatic knee (mechanical compression:

$293.30/year/knee, conventional: $373.29/year/knee) (p<0.0001, 95% CI of difference 47

<80< 112).

Conclusions: Routine mechanical compression of the knee for arthrocentesis and

injection is an effective bioengineering quality improvement intervention in a procedure

clinic.

Key words: arthrocentesis, intraarticular, injections, quality, knee

77
78

INTRODUCTION

After non-pharmacologic interventions and topical and oral medications, complete arthrocentesis followed intraarticular injection of corticosteroids is often recommended for symptomatic flares and short-term relief of pain of arthritis of the knee [1-5]. Klocke and colleagues have recently reported that corticosteroid injection of the osteoarthritic knee may actually decrease cartilage degradation in the short-term as measured by biomarkers, providing indirect support for the above recommendations [1-6]. Recently, Meehan et al, Bhavsar et al, and Yaqub et al have demonstrated that mechanical compression of the knee provides more complete arthrocentesis before injection in both the effusive and non-effusive knee and suggest that mechanical compression might be a reasonable quality improvement intervention for arthrocentesis [6-9]. In the present study we determined knee arthrocentesis and injection quality measures before and after the introduction of mechanical compression as a quality improvement intervention.

MATERIALS AND METHODS

This Quality Improvement program was formalized in our medical center in a formal effort to improve rheumatology outcomes and services with a commitment to continuous quality improvement in outpatient rheumatology as recently described recently by Chow et al [10]. The study was approved by the institutional review board (IRB) and in compliance with the Helsinki Declaration and subsequent revisions. The sequential study design was typical of quality improvement prospective cohort study with 1) measurement of baseline quality factors in consecutive traditionally treated patients, 2) introduction of the quality intervention, and 3) re-measurement of quality factors in

patients after the intervention. A total of 430 knees with grade II-III osteoarthritis were

included in this study.   In the primary corticosteroid arm, the first consecutive 215 knees

underwent conventional arthrocentesis and injection, then the quality intervention was

introduced, and the second 215 consecutive knees underwent mechanical compression-

assisted arthrocentesis and injection. Inclusion criteria for the study consisted of: 1)

painful symptomatic grade II-III osteoarthritis knee with the patient requesting a knee

injection, 2) indications for therapeutic-diagnostic arthrocentesis and/or injection, 3)

indication for corticosteroid injection, and 4) formal signed consent of the patient to

undergo the procedure [11]. Prior to the procedure, the presence or absence of clinical

effusion was confirmed by physical examination by palpation of the extended knee for

suprapatellar bursa distention, ballottement of a floating patella while applying pressure

superiorly and inferiorly, and visible-palpable fluid shift with asymmetric compression.

The knee was then classified as "effusive" (swollen) or "non-effusive" (dry).

Arthrocentesis prior to injection was attempted in all subjects prior to injection regardless

of the presence or absence of a clinical effusion.

Arthrocentesis and Joint Injection Technique

Chlorhexidine 2% was used for antisepsis. Knee procedures were performed

using the conventional lateral approaches with the one-needle multiple syringe technique

[12-20]. A 22 gauge 2 inch needle (4710007050 – 22 GX2" (0.7X50 mm), FINE-JECT,

Henke Sass Wolf, Kettenstrasse 1 D-78532 Tuttlingen, Germany) on a 3 ml syringe (3 ml

Luer Lok syringe, BD, 1 Becton Drive, Franklin Lakes, NJ 07417, website:

http://www.bd.com) filled with 3 ml of 1% lidocaine (Xylocaine® 1%, AstraZeneca

Pharmaceuticals LP, 1800 Concord Pike, P.O. Box 15437, Wilmington, DE 19850-5437)

was introduced into the lateral parapatellar recess of the suprapatellar bursa and if no

fluid returned, into the patellofemoral joint. The knee was then compressed or "milked"

by the operator's free hand to optimize fluid return [8,9,12-17]. If fluid return was

obtained and the syringe filled with synovial fluid, the intraarticular needle was left in

place, the 3 ml syringe was removed, and a syringe exchange was performed with

additional 20 ml syringes until fluid return ceased.   Pain scores, arthrocentesis success,

and fluid yield were recorded. After the fluid return ceased, a final syringe exchange was

performed, and the joint was then injected through the already-placed intraarticular

needle using a 3 ml syringe with 1 mg/kg triamcinolone acetonide suspension (maximum

80 mg) (Kenalog® 40, Westwood-Squibb Pharmaceuticals, Inc (Bristol-Myers Squibb),

345 Park Ave, New York, NY 10154-0004, USA). In the quality intervention group, an

elastomeric knee brace (YooSoo Adjustable Knee brace, Shenzhen Shi Hai Xun Yun Wei

Co., Ltd. No.203, 69 Dong, Liyuan Xin Cun, Bantian Street, Longgang, 518000,

Shenzhen, China, Amazon, https://www.amazon.com) was placed and modified so the

lateral suprapatellar bursa and patellofemoral joint could expand with fluid (Figures 1 and

2]. The brace applies constant mechanical compression to the medial suprapatellar bursa

and the synovial compartments of the inferior knee, collapsing these compartments, and

forcing fluid to the lateral suprapatellar bursa and the patellofemoral joint where the fluid

could be accessed (Figures 1 and 2). Before the brace was applied, the skin surfaces of

the knee were cleaned with 2% chlorhexidine. As the brace itself was clean but non-

sterile, after the compressive brace was placed, the antisepsis with 2% chlorhexidine was

again applied to the skin of the operative portal overlying the lateral recess of the

suprapatellar bursa, and arthrocentesis and injection were performed identically as

described above.

Outcome Measures:

Patients were observed for serious adverse events. Diagnostic fluid was defined as

$\geq$ to 2.0 milliliters (ml) (1 ml for culture, and 1 ml for cell counts and crystal examination

[8]. Patient pain was measured with the 0-10 cm Visual Analogue Pain Scale (VAS Pain

Scale), where 0 cm = no pain and 10 cm = unbearable pain [18-20]. The total number of

corticosteroid injections per year were recorded in each individual including those

administered in different clinics leveraging the electronic medical record to tract

comprehensive quality of care as described by Schmajuk and colleagues [21]. Weitoft et

al has demonstrated that complete arthrocentesis extends the time-to-next-symptomatic-

flare and thus the time-to-next-intervention [5]. Time-to-next-intervention also

corresponds to the time to the next corticosteroid or hyaluronan injection, surgical

intervention, physical therapy or splint referral, joint imaging, or referral to another

specialist, all of which increase costs. Thus the time-to-next-intervention permits

preliminary baseline cost determinations [5,18,22-24].

Costs associated with the Quality Intervention

Costs of the procedure in US dollars ($) were defined as those costs reimbursed

by 2017 Medicare (United States) national rates for HCPC/CPT 20610 code for a large

joint arthrocentesis for a physician office ($68.97/procedure), 15 minute outpatient

encounter ($76.69), 60 mg triamcinolone acetonide ($13.00/procedure), and compressive

brace ($10.00/brace) [25]. Yearly costs were calculated by multiplying the

costs/procedure x 12 months divided by the months to-the-next-intervention (time to

reinjection, surgical intervention, physical therapy or splint referral, joint imaging, or

referral to another specialist) – this calculation is an actual underestimate of true medical

costs but provides a standardization of the lowest level of costs from an injection

intervention as previously described [5,20,24,25].

*Statistical analysis***:** Data were entered into Excel (Version 5, Microsoft, Seattle, WA),

and analyzed in SISA (Simple Interactive Statistical Analysis,

http://www.quantitativeskills.com/sisa/). A power calculation was made using

preliminary data at this level where $\alpha=0.0001$, power = 0.9, and allocation ratio = 1.0

indicated that n=100 in each group would provide statistical power at the $p<0.001$ level

and n = 200 in each group at the $p<0.0001$ level. Pierson Chi square two by two table

analysis was performed on categorical data calculating both p values and confidence

intervals with significance reported at the $P <0.05$ level. Measurement data was analyzed

using the t-test calculating both p values and confidence intervals.

RESULTS

The two study cohorts (215 conventional and 215 compression-assisted) were

similar in demographics and baseline pain measures. The mean age of the conventional

cohort was 63.2 ± 11.6 years and the compression-assisted cohort was 60.1 ± 12.1 years

($p=0.007$, 95% CI of difference -5.3 <-3.1< -0.9). Male:female ratio was 17:198 (92%

female) in the conventional cohort, and 19:196 (88% female) in the compression cohort

($p=0.73$, z for 95% CI= 1.96, Pierson), typical of studies of osteoarthritis of the knee

demonstrating a female gender bias. Pre-procedural pain according to the 10 cm VAS

was 7.2 ± 1.9 cm in the conventional cohort and 7.2 ± 1.8 cm in the compression cohort (p

$=0.87$, 95% CI of difference:-0.4 <0.0< 0.3 (Wald)). Procedural pain was 2.6 ± 1.6 cm in

the conventional cohort and 2.4 ± 1.8 cm in the compression cohort (p = 0.22, 95% CI of

difference -0.5 <-0.2< 0.1 (Wald)). Post-Procedural pain was 1.1 ± 1.5 cm in the

conventional cohort and 1.0 ± 1.6 cm, in the compression cohort (p = 0.5, 95% CI of

difference -0.4 <-0.1< 0.2 (Wald)).

There were no serious adverse events encountered by the 430 patients in the

cohort including but not limited to reaction to local anesthesia, needle-stick, infection,

septic joint, hemarthrosis, deep venous thrombosis, pseudoseptic arthritis, dermal

atrophy, significant bruising, hemorrhage or post-injection visits to emergency facilities.

Diagnostic synovial fluid ($\geq$2 ml) was obtained in 9.3% (20/215) without

compression and 40.9% (88/215) with compression (p=0.00001, z for 95% CI= 1.96,

Pierson). Absolute volume of arthrocentesis fluid yield without compression was 1.6 ± 6.4

ml versus 5.3 ± 11.2 ml with compression (231% increase, CI of difference 2.0 <3.7< 5.4,

p=0.00001). These data are shown in Figure 3 in graphic form.

In the palpably non-effusive (dry) knees (n =177) the absolute volume of

arthrocentesis fluid obtained without compression was 0.0 ± 0.00 ml versus 1.5 ± 2.4 ml

with compression (n = 178) (>100 % increase, 95% CI of difference 0.995 <1.5< 2.005,

p=0.0001).

In the palpably effusive knee absolute volume of arthrocentesis without

compression (n=38) was 14.7 ± 13.8 ml versus 25.3 ± 15.5 ml with compression (n = 37)

(72.1% increase, 95% CI of mean difference: -3.0 <-1.7< -0.3, p=0.02).

Mechanical compression was associated with fewer subsequent corticosteroid

injections per year: mechanical compression: 1.7 ± 0.9 injections/year as opposed to

conventional: 2.4±0.5 injections/year (p<0.00001, 95% CI of difference -0.83<-0.70<-

0.56).

Time-to-next-intervention after intraarticular corticosteroid injection was longer

with constant compression 6.9±3.5 months as opposed to conventional treatment 5.1±2.7

months (p<0.00001, 95% CI of difference 1.2<1.8< 2.3). These data are shown in Figure

4 in graphic form.

Time-to-next-intervention was also longer in the effusive knees treated with

mechanical compression, 7.3±3.0 months as opposed to conventional 5.6±3.0 months

(p<0.016, 95% CI of difference-3.1 <-1.7< -0.3).

Time-to-next-intervention was also longer in the 178 non-effusive knees treated

with constant compression, 6.9±3.5 months as opposed to 177 non-effusive knees who

receive conventional treatment 5.0±2.6 months (p<0.01, 95% CI of difference -3.3 <-1.9<

-0.5).

Lowest level costs per year for conventional knee aspiration/injection was

$373±189/year/knee; and for mechanical compression was $293±152/year/knee

(p<0.0001, 95% CI of difference 47 <80< 112), thus, mechanical compression did not

increase yearly costs.

DISCUSSION:

In an effort to improve the arthrocentesis process, Meehan et al have recently

demonstrated that an external compressive brace shifts fluid in the knee where it can be

more easily accessed [7]. Similarly, Bhavsar et al demonstrated that external mechanical

compression of the knee collapsed non-target synovial spaces and dilated and expanded

target spaces and thus improved arthrocentesis success and yield in both the effusive and

non-effusive knee [8]. Yaqub et al have also demonstrated that mechanical compression

permits improved arthrocentesis success with the knee in different positioning [9].

Because of low cost and improved arthrocentesis success it has been proposed that

mechanical compression could be used as a low cost quality improvement intervention in

procedure clinics [7-9].

As part of an institutional quality improvement program to reduce occupational

needle-sticks and optimize needle procedures of the knee, we introduced mechanical

compression of the osteoarthritic knee before arthrocentesis and corticosteroid injection.

Meehan et al, Bhavsar et al, and Yaqub et al used a repeated measure paired study design

in the same subject that was sensitive to aspiration success, but not to other outcomes,

costs, or changes in adverse events [7-9]. The present pragmatic study used a different

design to study the quality effects of mechanical compression, specifically a two-cohort

design typical of quality improvement interventions [7-10,21-26].

In the present two-cohort study mechanical compression was associated with an

increased diagnostic joint aspiration success from 9% to 44%, and the volume of fluid

collected by 231% (Figure 3), similar to the findings of Bhavsar et al who used a repeated

measure in the same individual study design [8]. The use of the mechanical compression

brace did not appear to have a negative effect on subsequent injection outcomes. Time-

to-next-intervention after compression-assisted arthrocentesis was 6.9±3.5 months

compared to conventional arthrocentesis: 5.1±2.7 months (p<0.00001). Mechanical

compression was associated with in a reduction in the total number of corticosteroid

injections administered per year: mechanical compression: 1.7±0.9 injections/year;

conventional: 2.4±0.5 injections/year (p<0.00001). Mechanical compression did not

increase overall costs associated with management of the knee (mechanical compression:

$293.30/year/knee, conventional: $373.29/year/knee, p<0.0001). Thus, mechanical

compression of the knee for arthrocentesis before injection appears to be a reasonable

quality improvement intervention for knee procedures that improves quality measures

and does not increase overall costs.

A number of prior studies have focused on the need to accurately place the needle

intraarticularly and completely aspirate as much effusion as possible prior to injection,

and all have generally demonstrated improved joint injection outcomes [5,18,27-29].

In 2003 Weitoft et al demonstrated that complete aspiration of the knee prior to

corticosteroid injection prolonged the time-to-flare, and thus, reduced the need for

repetitive corticosteroid injection [5]. If the data in Figure 4 of the present study are

directly compared to the published data in Figure 1 of Weitoft et al, the present

independent data show a similar relationship in regards to improvement in outcome after

complete aspiration [5].

Extraction of joint fluid for biomarker analysis even in the dry (non-effusive)

knee is presently an important area in arthritis research and may also be integral to future

joint preservation strategies and therapies [6,22,29,30]. Similar to Bhavsar et al, Yaqub

et al, and Meehan et al, but using a different study design, we found that mechanical

compression permitted fluid extraction in a much greater proportion of osteoarthritic

knees that presented clinically as dry, non-effusive knees (Figure 3) [7-9]. The failure of

full synovial fluid extraction during conventional aspiration of both the effusive and non-

effusive knee may be due to a combination of mistargeting of the needle, the complex

intraarticular synovial compartments that trap viscous fluid, ineffective manual

compression, resistance to movement of fluid due to the semi-solid gel-like properties of

synovial fluid, and its non-Newtonian elasticity and viscosity [8,29-31]. A constant

circumferential pressure as provided by a mechanical compressive brace dilates the joint

space to be targeted (Figures 1 and 2) and takes advantage of the classic rheological

properties of synovial fluid to permit linear non-turbulent fluid flow to the needle access

point, enhancing fluid aspiration (Figure 3).

One limitation of this study is that medial approaches to aspiration and injection

were not used; it is possible that medial approaches might be just as successful by placing

the compressive brace in a reverse orientation. Another potential limitation was the

sequential rather than randomized study design that potentially could be a cause of

consistent bias; however, sequential methodology is a standard quality improvement

vehicle in all hospitals [10,22,23,26]. In the present study, an elastomeric brace was

used, it is anticipated that commercially available pneumatic braces would have similar

mechanical and clinical effects [32].

Conclusion.

Mechanical compression of the knee prior to corticosteroid injection permits more

complete synovial fluid extraction, and does not appear to increase costs or adversely

affect outcomes. Routine mechanical compression of the knee for arthrocentesis and

injection is a reasonable quality improvement intervention in a procedure clinic.

Conflict of Interest: None of authors have a conflict of interest in this manuscript. All

products used are commercially available and the authors have no interest in any of the

manufacturers, products, or distributors.

Funding: There was no funding for this study.

References

1.   Hochberg MC, Altman RD, April KT, Benkhalti M, Guyatt G, McGowan J,

Towheed T, Welch V, Wells G, Tugwell P; American College of Rheumatology.

American College of Rheumatology 2012 recommendations for the use of

nonpharmacologic and pharmacologic therapies in osteoarthritis of the hand, hip, and

knee (2012) Arthritis Care Res (Hoboken) 64:465-74.

2.   Brown GA. AAOS clinical practice guideline: treatment of osteoarthritis of the

knee: evidence-based guideline, 2nd edition (2013) J Am Acad Orthop Surg 21:577-9.

3.   Bruyère O, Cooper C, Pelletier JP, Maheu E, Rannou F, Branco J, Luisa Brandi

M, Kanis JA, Altman RD, Hochberg MC, Martel-Pelletier J, Reginster JY. A consensus

statement on the European Society for Clinical and Economic Aspects of Osteoporosis

and Osteoarthritis (ESCEO) algorithm for the management of knee osteoarthritis-From

evidence-based medicine to the real-life setting (2016) Semin Arthritis Rheum 45(4

Suppl):S3-11.

4.   Jüni P, Hari R, Rutjes AW, Fischer R, Silletta MG, Reichenbach S, da Costa BR.

Intra-articular corticosteroid for knee osteoarthritis (2015) Cochrane Database Syst Rev

Oct 22;(10):CD005328.

5.   Weitoft T, Uddenfeldt P. Importance of synovial fluid aspiration when injecting

intra-articular corticosteroids (20000 Ann Rheum Dis 59:233-5.

6.   Klocke R, Levasseur K, Kitas GD, Smith JP, Hirsch G.

Cartilage turnover and intra-articular corticosteroid injections in kneeosteoarthritis

(2018) Rheumatol Int 38:455-459

7.   Meehan R, Wilson C, Hoffman E, Regan E, Altimier L. Ultrasound

quantification of fluid shifts in the knees of arthritis patients before and after inflation of

a pneumatic compressive device (2015) Arthritis Rheum;67 (Absract Supplement):180.

8.   Bhavsar TB, Sibbitt WL. Jr, Band PA, Cabacungan RJ, Moore TS, Salayandia LC,

Fields RA, Kettwich SK., Roldan LP, Emil NS, Fangtham M, Bankhurst AD.

Improvement in diagnostic and therapeutic arthrocentesis via constant compression.

(2017) Clin Rheumatol Sep 14. doi: 10.1007/s10067-017-3836-x. [Epub ahead of print]

PMID: 28913649

9.   Yaqub S, Sibbitt WL Jr, Band PA, Bennett JF, Emil NS, Fangtham M, Fields RA,

Hayward WA, Kettwich SK, Roldan LP, Bankhurst AD. Can diagnostic and therapeutic

arthrocentesis be successfully performed in the flexed knee? (2018) J Clin Rheumatol

Feb 7. doi: 10.1097/RHU.0000000000000707. [Epub ahead of print] PMID:29424762

Jul 1.

10.   Chow SL, Shojania KG."Rheum to Improve": Quality improvement in

outpatient rheumatology (2017) J Rheumatol 44:1304-1310.

11.   Brandt KD, Fife RS, Braunstein EM, Katz B. Radiographic grading of the

severity of knee osteoarthritis: relation of the Kellgren and Lawrence grade to a grade

based on joint space narrowing, and correlation with arthroscopic evidence of articular

cartilage degeneration (1991) Arthritis Rheum 34:1381-6.

12.   Chen B, Lai LP, Putcha N, Stitik TP, Foye PM, et al. Optimal needle placement

for ultrasound-guided knee joint injections or aspirations (2014) J Trauma Treat 3:216.

doi:10.4172/2167-1222.1000216; http://dx.doi.org/10.4172/2167-1222.1000216

Accessed August 27, 2017.

13.   Dabke HV. Accuracy of needle placement into the intra-articular space of the

knee (2004) J Bone Joint Surg Am 86-A:433-4.

14.   Gardner GC. Teaching arthrocentesis and injection techniques: What is the best

way to get our point across? (2007) J Rheumatol 34:1448-1450.

15.   Jackson DW, Evans NA, Thomas BM. Accuracy of needle placement into the

intra-articular space of the knee (2002) J Bone Joint Surg Am 84-A:1522-7.

16.   Roberts WN. Primer: pitfalls of aspiration and injection (2007) Nat Clin Pract

Rheumatol 3:464-72.

17.   Roberts WN, Hayes CW, Breitbach SA, Owen DS Jr. Dry taps and what to do

about them: a pictorial essay on failed arthrocentesis of the knee (1996) Am J Med

100:461-4.

18.   Sibbitt WL Jr, Kettwich LG, Band PA, Chavez-Chiang NR, DeLea SL, Haseler

LJ, Bankhurst AD (2012) Does ultrasound guidance improve the outcomes of

arthrocentesis and corticosteroid injection of the knee? Scand J Rheumatol 41:66-72.

19.   Katz J, Melzack R. Measurement of pain (1999) Surg Clin North Am 79:231-252.

20.   Sibbitt WL Jr, Band PA, Chavez-Chiang NR, Delea SL, Norton HE, Bankhurst

AD (2011) A randomized controlled trial of the cost-effectiveness of ultrasound-guided

intraarticular injection of inflammatory arthritis. J Rheumatol 38:252-63.

21. Schmajuk G, Yazdany J (2017) Leveraging the electronic health record to

improve quality and safety in rheumatology.   Rheumatol Int 37:1603-1610.

22.   Band PA. Using Medicare data to understand health care value: measures of

incremental cost and effectiveness are both needed to estimate value (2015) JAMA

Intern Med 175:462.

23.   Grosse SD, Teutsch SM, Haddix AC. Lessons from cost-effectiveness research

for the United States public health policy (2007) Annu Rev Public Health 28:365-91.

24.   Sibbitt WL Jr, Band PA, Kettwich LG, Chavez-Chiang NR, Delea SL, Bankhurst

AD (2011) A randomized controlled trial evaluating the cost-effectiveness of sonographic

guidance for intra-articular injection of the osteoarthritic knee. J Clin Rheumatol 17:409-

15.

25.   U.S. Department of Health & Human Services. Centers for Medicare &

Medicaide Services. Physician Fee Schedule Look-up.

http://www.cms.hhs.gov/PFSlookup/ assessed 25 March, 2018.

26.   Molina CS, Thakore RV, Blumer A, Obremskey WT, Sethi MK. Use of the

National Surgical Quality Improvement Program in orthopaedic surgery (2015) Clin

Orthop Relat Res 473:1574-81

27.   Zhang Q, Zhang T. Effect on pain and symptoms of aspiration

before hyaluronan injection for knee osteoarthritis: A prospective, randomized, single-

blind study (2016) Am J Phys Med Rehabil 95:366-71.

28.   Tanaka N, Sakahashi H, Sato E, Hirose K, Ishima T, Ishii S. Intra-articular

injection of high molecular weight hyaluronan after arthrocentesis as treatment for

rheumatoid knees with joint effusion (2002) Rheumatol Int 22:151-4.

29.   Band PA, Heeter J, Wisniewski HG, Liublinska V, Pattanayak CW, Karia RJ,

Stabler T, Balazs EA, Kraus VB (2015) Hyaluronan molecular weight distribution is

associated with the risk of knee osteoarthritis progression.   Osteoarthritis Cartilage

23:70-6.

30.   Wisniewski HG, Colón E, Liublinska V, Karia RJ, Stabler TV, Attur M,

Abramson SB, Band PA, Kraus VB. TSG-6 activity as a novel biomarker of progression

in knee osteoarthritis (2014) Osteoarthritis Cartilage 22:235-41.

31.   Chaudhry H, Bukiet B, Roman M, Stecco A, Findley T (2013) Squeeze film

lubrication for non-Newtonian fluids with application to manual medicine. Biorheology.

50:191-202.

32.   Arthroventions: KneeTap Needle Guide.

http://www.arthroventions.com/kneetap-product-overview Accessed Jan 5, 2018.

FIGURES.

FIGURE 1. The arthrocentesis is performed through the superiolateral approach with an

elastomeric brace applying circumferential (radial) constant mechanical compression to

the knee and forcing residual fluid where it can be accessed at the superiolateral portal.

An absorbent impermeable drape can be placed in the access portal to protect the brace.

FIGURE 2. This ultrasound image demonstrates the lateral recess of the suprapatellar

bursa with manual compression versus mechanical constant compression with the

elastomeric brace showing substantial shift of intraarticular synovial fluid toward the

access point at the lateral recess of the suprapatellar bursa. The aspiration/injection

needle can be seen in the effusion on the right hand side after the constant compression

brace has been applied.

FIGURE 3. This graph shows the markedly increased synovial fluid yield with

mechanical constant compression (upper line) (n=215) and conventional (lower line)

(n=215), demonstrating a mean 230% increase in synovial fluid yield (p<0.0001).

FIGURE 4.   This graph shows the increased time-to-next intervention with mechanical

constant compression (upper line) (n=215) and conventional (lower line) (n=215) in 430

knees demonstrating a mean 35% increase in time-to-next-intervention (p<0.00001).

Manual Compression   Mechanical Compression