Bariatric surgery changes gut microbiota composition independent of operation

type

*[1]Fernanda L. Paganelli, *[2]Misha Luyer, [3]C. Marijn Hazelbag, [3]Hae-Won Uh,

[1]Malbert R.C. Rogers, [2]Danielle Adriaans, [1, 4]Roos-Marijn Berbers, [1]Antoni P.A.

Hendrickx, [1]Marco C. Viveen, [1]James A. Groot, [1]Marc J. M. Bonten, [1]Ad C. Fluit,

[1]Rob J.L. Willems, [4]Helen L. Leavis.

1. *Department of Medical Microbiology, University Medical Center Utrecht,*

Utrecht University, The Netherlands.

2. *Department of Surgery, Catharina Hospital Eindhoven, Eindhoven the*

Netherlands

3. *Department of Biostatistics and Research Support, Julius Center, University*

Medical Center Utrecht, Utrecht University, The Netherlands.

4. *Department of Rheumatology and Clinical Immunology, University Medical*

Center Utrecht, Utrecht University, The Netherlands.

* these authors equally contributed to the study

The authors declare no conflict of interest.

Corresponding Author: Helen L. Leavis. Department of Rheumatology and Clinical

Immunology, University Medical Center Utrecht, Utrecht University, The

Netherlands. Heidelberglaan 100. 3584 CX Utrecht. T +31 0887557357.

h.leavis@umcutrecht.nl.

Short running head (not more than 50 characters (count letters and spaces):

Microbiota changes in bariatric surgery patients

Abbreviations list and their definitions for all abbreviations used in the text if

there are 3 or more:

RYGB: Roux-Y Gastric bypass

SG: Sleeve Gastrectomy

OTU: Operational Taxonomic Units

bp: base pairs

PCA: Principle Component Analysis

FDR: False Discovery Rate

HbA1c: glycated hemoglobin

PCoA: Principal coordinate analysis

VitD: vitamin D;

VitB6: vitamin B6.

Abstract

Bariatric surgery in patients with morbid obesity, either through gastric sleeve

gastrectomy or Roux-Y gastric bypass surgery, leads to sustainable weight loss,

improvement of metabolic disorders and changes in the intestinal microbiota. Yet, the

relationship between changes in gut microbiota, weight loss and the surgical

procedure is unknown. We, therefore, determined temporal changes in microbiota

composition in 45 obese patients undergoing a crash diet followed by gastric sleeve

gastrectomy ($n= 22$) or Roux-Y gastric bypass ($n= 23$). Intestinal microbiota

composition was determined before intervention (baseline, S1), 2 weeks after a crash

diet (S2), and 1 week (S3), 3 months (S4) and 6 months (S5) after surgery. Relative to

S1, the microbial diversity index declined at S2 and S3 ($p< 0.05$), and gradually

returned to baseline levels at S5. The crash diet was associated with an increased

abundance of Rikenellaceae and decreased abundances of Ruminococcaceae and

Streptococcaceae ($p< 0.05$). After surgery, at S3, the relative abundance of

Bifidobacteriaceae had decreased (compared to the moment directly after the crash

diet), whereas those of Streptococcaceae and Enterobacteriaceae had increased ($p<$

0.05). Increased weight loss during the next 6 months was not associated without

major changes in microbiota composition. Significant differences between both

surgical procedures were not observed at any of the time points. In conclusion,

undergoing a crash diet and bariatric surgery were associated with an immediate but

temporary decline in the microbial diversity, with immediate and permanent changes

in microbiota composition, with no differences between patients undergoing gastric

sleeve gastrectomy or Roux-Y gastric bypass surgery.

Introduction

Bariatric surgery is the only sustainable effective treatment for obesity (1). Surgical

procedures such as the Roux-Y Gastric bypass (RYGB) and the sleeve gastrectomy

(SG) facilitate a 50-70% decrease in excess body weight and fat mass (1). In addition,

surgery leads to decreased caloric intake or malabsorption and to metabolic changes,

such as an improved glucose metabolism, and is associated with a changed intestinal

microbiota (2-4). The role of altered host-microbial interactions in this process is not

well understood (1). Studies on the composition of the distal gut microbiota in obesity

and after RYGB in humans and rodents yielded long lasting changes in types and ratio

of enteric bacteria (3, 5-8). Furthermore, transfer of the gut microbiota from RYGB-

treated mice to non-operated, germ-free mice resulted in weight loss and decreased fat

mass in the recipient animals (5). These findings support a direct effect of the

microbiota on weight and adiposity.

To further elucidate the results of the entire bariatric surgery procedure on the

intestinal microbiota composition we investigated sequentially collected stool samples

from 45 morbid obese patients undergoing either RYGB or SG at five different time

points before and after surgery.

Subjects and Methods

Ethics Statement

The study protocol was in accordance with the regulations of the Ethics Committee of

Catharina Hospital Eindhoven.

Study Design

In this observational study patients with morbid obesity were recruited from the

Catharina Hospital Eindhoven out-patient obesity clinic between September 2014 and

November 2014. All 45 patients fulfilled the criteria for bariatric surgery and were

screened before surgery for eligibility by a team including a surgeon, dietician and a

psychologist. Two weeks before the planned surgery, patients were subjected to a

crash diet consisting of 500 calories a day for 2 weeks. Type of surgery was

determined based on clinical criteria and shared decision making between surgeon

and patient. Reasons to refrain from RYGB were medication dependence, increased

operation risk or super obesity. A reason to refrain from SG was gastroesophageal

reflux disease. During surgery, patients received 1 g cefazolin antibiotic prophylaxis

intravenously. After hospitalization, general practitioners managed adjustments of

insulin, oral diabetics and other medication in the home setting. Patients visited out-

patient clinic at 3, 6 and 12 months for evaluation and will remain in follow up for 5

years. In this study (flow chart in supplementary figure 1), we used STROBE cohort

reporting guidelines (9).

Sample collection and DNA extraction.

Stool samples (Sterilin specimen container, Thermo-Fisher) were gathered at the out-

patient clinic or at patient homes. Samples were always stored in the freezer and

collected at the homes of the patients using dry-ice and stored at the hospital at -80°C.

Sample were collected at 5 different time points; before the start of the crash diet (S1),

2 weeks after the crash diet (S2), and 1 week (S3), 3 months (S4) and 5 to 6 months

after surgery (S5).

Total bacterial DNA from feces samples was isolated according to Godon *et al.* (10).

When isolated DNA contained PCR inhibitors (20% of the samples random

distributed over the time points), samples were submitted to an extra step of

isopropanol precipitation and column purification with QiAamp stool mini kit

(Qiagen). DNA was stored at −20°C prior to further analysis.

16S rRNA gene sequencing strategy and analysis

A 469 bp encompassing the V3 and V4 hypervariable regions of the 16S rRNA gene

was amplified and sequenced using the Illumina MiSeq Reagent Kit v3 (600-cycle) on

an Illumina MiSeq instrument according to Fadrosh *et al.* (11). A total of 3 amplicon

pools were sequenced, generating 8.9, 7.8 and 14.4 (mean of 10.3) million total reads.

These 2x300 bp paired-end reads were pre-processed as follows. The first 12 bp of

each paired-end containing the index sequences were extracted and afterwards

concatenated to dual-index barcodes of 24 bp specific for each read-pair and sample.

Paired reads were merged, as an overlap of about 90 bp was expected, using FLASH

(version 1.2.11) (12). Subsequently, these merged reads were de-multiplexed using

the split_libraries_fastq.py script rom and analyzed by the QIIME microbial

community analysis pipeline (version 1.9.1) (13). Quality filtering was also performed

during this step, truncating reads with an average PHRED quality score of 20 or less.

After removal of the barcodes, heterogeneity spacers, and primer sequences about

19.8 million sequences were left with a mean length of 410 bp (median length of

405). The obtained sequences with a minimum of 97% similarity were assigned to

operational taxonomic units (OTUs) using QIIME's open-reference OTU picking

workflow (pick_open_reference_otus.py). This workflow was carried out using

USEARCH (version 6.1.544) (14) for OTU picking, in addition to detection and

removal of chimeric sequences. The obtained OTU sequences were aligned to the

Greengenes 16S rRNA gene database (gg_13_8_otus), followed by removal of OTUs

represented by less than 0.005% of the total number of sequences. The generated

OTU table and phylogenetic tree were used for assessing alpha- and beta-diversity

using QIIME's core_diversity_analyses.py workflow with a rarefaction depth of

20001 sequences. The weighted UniFrac distance was used to calculate beta-diversity

of the samples, while the shannon index was used for the alpha-diversity. For

Principal Component Analysis (PCA) R 3.5.0 in an environment of RStudio 1.1.383

(RStudio Team, Boston, MA) (15) was employed, using zCompositions, clr

transformation and ggpplot R packages (16-18).

Statistical analysis

Microbiota changes between time points and operation types were investigated using

ANCOM (19) in R 3.3.3 (15). Changes in the clinical parameters (BMI, vitamin D,

vitamin B6, cholesterol, bilirubin, glycated hemoglobin (HbA1c), iron, ferritin and

folate) between baseline and 6 months after surgery was analyzed by applying t-test in

Prism GraphPad (version 7.0). Associations between changes in total read counts at

family level (at baseline versus 6 months after surgery) and changes in patient

characteristics (at baseline versus 6 months after surgery) were investigated using a

linear regression model. To eliminate possible confounding effects, age and sex were

included as covariates. For these analyses, changes in total read counts were used as

outcome, whereas changes in patient characteristics were used as predictor (model:

change_in_read_counts ∼ β1·age + β2·sex + β3·change_in_patient_characteristic).

For association analysis R 3.5.0 in an environment of RStudio 1.1.383 (RStudio

Team, Boston, MA) was employed (15). Results are presented using pheatmap

package (https://CRAN.R-project.org/package=pheatmap). For statistical testing we

used false discovery rates (FDRs) correct for multiple comparisons, and an FDR-

adjusted p-value < 0.05 was considered as significant (20).

Results

RYGB and SG resulted in significant decrease of BMI in all patients.

In this study, 45 Caucasian Dutch patients were included with an average age of 43

years, 36 (84%) being female, 11 (24%) using proton pump inhibitors and 4 (9%)

having type 2 Diabetes Mellitus at baseline (Table 1). After a crash diet, 22 patients

underwent SG and 23 underwent RYGB. At baseline the mean BMI was 42.9 (+/-

6.56) and 43 (+/-4.13) for patients undergoing RYGB and SG, respectively. At 6

months after the procedure BMI declined to 30.81 (+/-5.35) and 31.52 (+/-3.86),

respectively, with no significant difference based on surgery type (Table 1).

Crash diet reduces microbial alpha diversity, which is restored to baseline levels 6

months after surgery, irrespective of surgery type.

In total 221 fecal samples were collected, with 4 samples missing from 4 unique times

points from 4 different patients. Using a pre-defined cut-off value of 20001 reads, 220

samples could be analyzed.

The initial crash diet had a stronger effect on total microbiota diversity as the Shannon

diversity index declined from 4.5 at baseline (S1) to 4.0 after the crash diet ($p< 0.05$)

(S2) and then gradually returned to 4.5 at 3 (S4) and 6 months (S5) after surgery

(**Figure 1a**). Differences in diversity are reflected by an initial decrease and

subsequent rise in numbers of distinct microbial OTUs. At baseline, 3 months and 6

months after surgery more than 500 OTUs were identified, whereas after the crash

diet and at 1 week after surgery this number was reduced to below 400 OTUs (**Figure**

1b).

In Principle Component Analysis (PCA) (**Figure 1c**), patients at baseline (S1) and

after crash diet (S2) are more similar to each other when compared to the times points

after the surgery (S3, S4, S5), which cluster together further apart.

Distinct microbial changes appear directly after crash diet, but are replaced by

persistent distinct changes shortly after surgery.

Significant changes in total relative abundance of specific families in the different

time points were observed (**Figure 2h**). After the crash diet (S2) there was a

significant reduction in relative abundance of 2 microbial families, Streptococcaceae

and Ruminococcaceae (**Figure 2a and d**), and a significant increase in 1 family,

Rikenellaceae (**Figure 2e**). Subsequent comparison of the microbial composition pre-

surgery (S2) and 1-week post-surgery (S3) revealed a significant increase in the

relative abundance of Streptococcaceae and Enterobacteriaceae families (**Figure 2a**

and b) and a decrease in Bifidobacteriaceae, which persisted until 6 months post-

surgery (S5) (**Figure 2c**). In these 6 months (at S5) microbiota complexity was

restored (Figure 1a and b), which when compared with 1 week after surgery (S3)

coincided with increased relative abundance of low abundance families

Veillonellaceae and the Clostridiales order with no further family classification

(**Figure 2f and g**).

When RYGB and SG surgeries were analyzed separated, no significant differences in

microbiota composition based on beta diversity and relative abundance was observed

at baseline (S1) (**Figure 3a**), 1 week (S3) (**Figure 3b**) or 6 months after surgery (S5)

(**Figure 3c**) between patients that underwent either SG and RYGB (**Figure 3d**).

Significant associations between microbiota changes and clinical markers

Clinical parameters in patients were analyzed at baseline and 6 months after the

surgery. Besides weight loss, serum levels of vitamin D, B6, cholesterol, bilirubin,

HbA1c, iron, ferritin and folate improved 6 months after surgery when compared to

baseline (FDR-adjusted, $p < 0.05$, Table 1).

These changes were associated to overall differences in microbial abundance in

relation to the changes in clinical parameters at S5 versus S1, which are highlighted in

Figure 4. Significant associations were only found in low abundance families.

Decreased bilirubin level was associated with and Prevotellaceae, Bacteroidales and

Peptococcaceae taxa; and increased iron level was associated with Pasteurellaceae. In

addition, a decreased HbA1c was associated with Coriobacteriaceae and Clostridiales

taxa The most pronounced measured effects in the dataset was a negative association

between Prevotellaceae, Veillonellaceae, Streptococcaceae, Bifidobacteriaceae and

Enterobacteriaceae taxa in relation to decreased serum cholesterol levels, whereas

most pronounced positive associations were found between Lachnospiraceae and

Coriobacteriaceae taxa in relation to decreased cholesterol levels (Figure 4). Yet,

these associations were not statistically significant after FDR adjustment. (Figure 4).

Discussion

This study is novel in the fact that it compares microbiota profiles before and shortly

after bariatric surgery with subsequent follow-up profiles. We describe the sequential

impact of a crash diet followed by either RYGB or SG surgery, resulting in

progressive weight loss and changes in the gut microbiota composition. In addition to

substantial weight loss and improvements in clinical parameters, bariatric surgery also

induces long-lasting changes in microbiota composition in most patients. The most

apparent immediate change in microbiota composition occurred after the crash diet,

with a concurrent reduction in alpha diversity, whereas surgery was associated with

early and sustained replacement of distinct bacterial taxa and restoration of the

diversity.

Although significant microbial changes are identified in the gut microbiota of

bariatric surgery patients, at 6 months after surgery the total microbial diversity was

similar to microbial diversity measured at baseline. This probably reflects a severe

stress on the human microbiota by a crash diet, with a significant change in catabolic

state (21). Persisting post-surgery microbiome changes suggest adaptation to anatomic

and physiologic changes induced by the surgery. Previously reported effects of

bariatric surgery on microbiota diversity have ranged from an increase in total

diversity (22, 23) to absence of change and even a decrease in alpha diversity (24).

We suspect that baseline sampling in relation to crash diet may vary between studies,

and might contribute to the reported differences between studies. Moreover, not all

studies describe whether crash diet is part of the entire procedure and, if so, the exact

timing of the diet in relation to baseline patient sampling.

Besides a stable bacterial alpha diversity after surgery, we observed profound

differences after each consecutive intervention on bacterial taxa composition.

The crash diet immediately resulted in an increase in the relative abundance of

Bifidobacteriaceae and decrease of Streptococcaceae, whereas the opposite effect was

observed after surgery; an increase in abundance of Streptococcaceae and decline in

Bifidobacteriaceae that persisted for at least 6 months. Observed persistent increase in

Enterobacteriaceae after surgery, confirms previous sustained changes reported in

humans and animal models (rats) (7, 21, 25, 26). One of the main differences is

exposure to undigested nutrients and biliopancreatic enzymes, which may play

important roles in the microbial composition, intestinal permeability and intestinal

adaptation (27). Since increased intestinal permeability is associated with

inflammation and reductions in alpha diversity, which is also associated with obesity,

it is questionable whether restoration of alpha diversity to baseline level may also

reflect persistent inflammation in the post-surgery state at 6 months, which has been

previous related to increase in Enterobacteriaceae (16).

Although others observed microbiota changes only after RYGB (5), here we observed

this in both surgery types. This suggests that despite the 2 procedures result in distinct

anatomic differences, this did not seem to influence the post-surgery changes in

relative abundance of Bifidobacteriaceae, Streptococcaceae and Enterobacteriaceae

observed amongst both patient groups and which were similar for both types of

surgery. Moreover, bariatric surgery in itself, unlike crash diet, results in an altered

long-lasting composition of the microbiota.

Although a significant association with changed clinical parameters between baseline

and 6 months after surgery was lacking, the relative abundance of Bifidobacteriaceae,

Streptococcaceae and Enterobacteriaceae taxa changed significantly shortly after

surgery. Whether, these changes reflect consequences of surgery induced anatomic

differences rather than improvement in clinical parameters remains speculative. We

observed a significant association between increased serum bilirubin level and

decreased relative abundance of Bacteroidales, Peptococcaceae and Prevotellaceae

taxa in this dataset. The abundance of Bacteroidales in the gut microbiota could

contribute to the increase in bilirubin level, since *Bacteroides fragilis*, which is part of

Bacteroidales taxa, is one of the bacterial species described to be able to metabolize

bilirubin in the gut (28, 29). In addition, a decreased HbA1c was found significantly

associated with decreased Coriobacteriaceae and increased Clostridiales taxa.

Nevertheless, the exact meaning of changes of these low abundance taxa is unknown.

This study failed to confirm the suggested relationship between increased abundance

of Firmicutes and Bacteroidetes and obesity (30, 31), as the relative abundance of the

family members of these phyla remained stable before and after surgery, despite

significant weight loss. In addition, other studies described that Faecalibacterium (*F.*

prausnitzii) was assumed to play a role in inflammation and glucose homeostasis in

obesity with a reduced relative abundance after RYGB surgery (3, 8, 32, 33). In our

study, a decreased abundance of the Ruminococcaceae family, to which *F. prausnitzii*

belongs, was observed after the crash diet, yet this change did not sustain after

surgery.

In conclusion, here we illustrate that temporal sampling of bariatric surgery patients

with subsequent microbiome analysis can lead to increased insights into the relative

contribution of interventions on stability and composition of the microbiota. We show

that a crash diet invoked profound temporary changes in total microbiota diversity and

composition, yet surgery precluded early fixed changes of microbial composition and

restoration of the microbial diversity that likely contribute to weight loss.

Acknowledgments

The authors declare that there are no conflicts of interest. Authors contribution:

conceived and designed the experiments: ML, DA, MJMB, ACF, HLL. Performed the

experiments: ML, DA, APAH, JG, MCV. Analyzed the data: FLP, MRCR, CMH,

HWU, RMB. Wrote the paper: FLP, ML, CMH, HWU, MRCR, DA, RMB, APAH,

MCV, JG, MJMB, ACF, RJLW, HLL.

References

1.   Tilg H. Obesity, metabolic syndrome, and microbiota: multiple interactions. J
Clin Gastroenterol 2010;44 Suppl 1:S16-8. doi:
10.1097/MCG.0b013e3181dd8b64.
2.   Kobyliak N, Conte C, Cammarota G, Haley AP, Styriak I, Gaspar L, Fusek J,
Rodrigo L, Kruzliak P. Probiotics in prevention and treatment of obesity: a
critical view. Nutr Metab (Lond) 2016;13:14. doi: 10.1186/s12986-016-0067-
0.
3.   Furet JP, Kong LC, Tap J, Poitou C, Basdevant A, Bouillot JL, Mariat D,
Corthier G, Dore J, Henegar C, et al. Differential adaptation of human gut
microbiota to bariatric surgery-induced weight loss: links with metabolic and
low-grade inflammation markers. Diabetes 2010;59(12):3049-57. doi:
10.2337/db10-0253.
4.   Kong LC, Tap J, Aron-Wisnewsky J, Pelloux V, Basdevant A, Bouillot JL,
Zucker JD, Dore J, Clement K. Gut microbiota after gastric bypass in human
obesity: increased richness and associations of bacterial genera with adipose
tissue genes. Am J Clin Nutr 2013;98(1):16-24. doi: 10.3945/ajcn.113.058743.
5.   Liou AP, Paziuk M, Luevano JM, Jr., Machineni S, Turnbaugh PJ, Kaplan
LM. Conserved shifts in the gut microbiota due to gastric bypass reduce host
weight and adiposity. Sci Transl Med 2013;5(178):178ra41. doi:
10.1126/scitranslmed.3005687.
6.   Sweeney TE, Morton JM. The human gut microbiome: a review of the effect
of obesity and surgically induced weight loss. JAMA Surg 2013;148(6):563-9.
doi: 10.1001/jamasurg.2013.5.
7.   Zhang H, DiBaise JK, Zuccolo A, Kudrna D, Braidotti M, Yu Y,
Parameswaran P, Crowell MD, Wing R, Rittmann BE, et al. Human gut
microbiota in obesity and after gastric bypass. Proc Natl Acad Sci U S A
2009;106(7):2365-70. doi: 10.1073/pnas.0812600106.
8.   Graessler J, Qin Y, Zhong H, Zhang J, Licinio J, Wong ML, Xu A, Chavakis
T, Bornstein AB, Ehrhart-Bornstein M, et al. Metagenomic sequencing of the
human gut microbiome before and after bariatric surgery in obese patients
with type 2 diabetes: correlation with inflammatory and metabolic parameters.
Pharmacogenomics J 2013;13(6):514-22. doi: 10.1038/tpj.2012.43.
9.   von Elm E AD, Egger M, Pocock SJ, Gotzsche PC, Vandenbroucke JP. . The
Strengthening the Reporting of Observational Studies in Epidemiology
(STROBE) Statement: guidelines for reporting observational studies.
10. Godon JJ, Zumstein E, Dabert P, Habouzit F, Moletta R. Molecular microbial
diversity of an anaerobic digestor as determined by small-subunit rDNA
sequence analysis. Appl Environ Microbiol 1997;63(7):2802-13.
11. Fadrosh DW, Ma B, Gajer P, Sengamalay N, Ott S, Brotman RM, Ravel J. An
improved dual-indexing approach for multiplexed 16S rRNA gene sequencing
on the Illumina MiSeq platform. Microbiome 2014;2(1):6. doi: 10.1186/2049-
2618-2-6.
12. Magoc T, Salzberg SL. FLASH: fast length adjustment of short reads to
improve genome assemblies. Bioinformatics 2011;27(21):2957-63. doi:
10.1093/bioinformatics/btr507.
13. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello
EK, Fierer N, Pena AG, Goodrich JK, Gordon JI, et al. QIIME allows analysis
of high-throughput community sequencing data. Nat Methods 2010;7(5):335-
6. doi: 10.1038/nmeth.f.303.

14.   Edgar RC. Search and clustering orders of magnitude faster than BLAST.
Bioinformatics 2010;26(19):2460-1. doi: 10.1093/bioinformatics/btq461.
15.   R Foundation for Statistical Computing V, Austria. R: A language and
environment for statistical computing., 2018.
16.   Palleja A, Kashani A, Allin KH, Nielsen T, Zhang C, Li Y, Brach T, Liang S,
Feng Q, Jorgensen NB, et al. Roux-en-Y gastric bypass surgery of morbidly
obese patients induces swift and persistent changes of the individual gut
microbiota. Genome Med 2016;8(1):67. doi: 10.1186/s13073-016-0312-1.
17.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag
New York 2016.
18.   Gregory B. Gloor GR. Compositional analysis: a valid approach to analyze
microbiome high-throughput sequencing data. Canadian Journal of
Microbiology 2016;62(8). doi: https://doi.org/10.1139/cjm-2015-0821.
19.   Mandal S, Van Treuren W, White RA, Eggesbo M, Knight R, Peddada SD.
Analysis of composition of microbiomes: a novel method for studying
microbial composition. Microb Ecol Health Dis 2015;26:27663. doi:
10.3402/mehd.v26.27663.
20.   Storey JDaT, R. Statistical significance for genomewide studies
Proc Nat Acad Sci USA 2003;100.
21.   Li JV, Ashrafian H, Bueter M, Kinross J, Sands C, le Roux CW, Bloom SR,
Darzi A, Athanasiou T, Marchesi JR, et al. Metabolic surgery profoundly
influences gut microbial-host metabolic cross-talk. Gut 2011;60(9):1214-23.
doi: 10.1136/gut.2010.234708.
22.   Guo Y, Liu CQ, Shan CX, Chen Y, Li HH, Huang ZP, Zou DJ. Gut
microbiota after Roux-en-Y gastric bypass and sleeve gastrectomy in a
diabetic rat model: Increased diversity and associations of discriminant genera
with metabolic changes. Diabetes Metab Res Rev 2017;33(3). doi:
10.1002/dmrr.2857.
23.   Basso N, Soricelli E, Castagneto-Gissey L, Casella G, Albanese D, Fava F,
Donati C, Tuohy K, Angelini G, La Neve F, et al. Insulin Resistance,
Microbiota, and Fat Distribution Changes by a New Model of Vertical Sleeve
Gastrectomy in Obese Rats. Diabetes 2016;65(10):2990-3001. doi:
10.2337/db16-0039.
24.   Ward EK, Schuster DP, Stowers KH, Royse AK, Ir D, Robertson CE, Frank
DN, Austin GL. The effect of PPI use on human gut microbiota and weight
loss in patients undergoing laparoscopic Roux-en-Y gastric bypass. Obes Surg
2014;24(9):1567-71. doi: 10.1007/s11695-014-1275-1.
25.   Patrone V, Vajana E, Minuti A, Callegari ML, Federico A, Loguercio C,
Dallio M, Tolone S, Docimo L, Morelli L. Postoperative Changes in Fecal
Bacterial Communities and Fermentation Products in Obese Patients
Undergoing Bilio-Intestinal Bypass. Front Microbiol 2016;7:200. doi:
10.3389/fmicb.2016.00200.
26.   Shao Y, Ding R, Xu B, Hua R, Shen Q, He K, Yao Q. Alterations of Gut
Microbiota After Roux-en-Y Gastric Bypass and Sleeve Gastrectomy in
Sprague-Dawley Rats. Obes Surg 2017;27(2):295-302. doi: 10.1007/s11695-
016-2297-7.
27.   Guo Y, Huang ZP, Liu CQ, Qi L, Sheng Y, Zou DJ. Modulation of the gut
microbiome: a systematic review of the effect of bariatric surgery. Eur J
Endocrinol 2018;178(1):43-56. doi: 10.1530/EJE-17-0403.

28. Fahmy K. GCH, Nicholson D.C. . The reduction of bile pigments by faecal
and intestinal bacteria. Biochimica et Biophysica Acta (BBA) - General
Subjects 1972;264(1). doi: https://doi.org/10.1016/0304-4165(72)90119-5.
29. Hamoud A.R WL, Stec D. E., and Hinds Jr T. D. Bilirubin in the Liver–Gut
Signaling Axis. Trends in Endocrinology & Metabolism 2018;29(3). doi:
https://doi.org/10.1016/j.tem.2018.01.002.
30. Turnbaugh PJ, Backhed F, Fulton L, Gordon JI. Diet-induced obesity is linked
to marked but reversible alterations in the mouse distal gut microbiome. Cell
Host Microbe 2008;3(4):213-23. doi: 10.1016/j.chom.2008.02.015.
31. le Roux CW, Bueter M, Theis N, Werling M, Ashrafian H, Lowenstein C,
Athanasiou T, Bloom SR, Spector AC, Olbers T, et al. Gastric bypass reduces
fat intake and preference. Am J Physiol Regul Integr Comp Physiol
2011;301(4):R1057-66. doi: 10.1152/ajpregu.00139.2011.
32. Damms-Machado A, Mitra S, Schollenberger AE, Kramer KM, Meile T,
Konigsrainer A, Huson DH, Bischoff SC. Effects of surgical and dietary
weight loss therapy for obesity on gut microbiota composition and nutrient
absorption. Biomed Res Int 2015;2015:806248. doi: 10.1155/2015/806248.
33. Sokol H, Pigneur B, Watterlot L, Lakhdari O, Bermudez-Humaran LG,
Gratadoux JJ, Blugeon S, Bridonneau C, Furet JP, Corthier G, et al.
Faecalibacterium prausnitzii is an anti-inflammatory commensal bacterium
identified by gut microbiota analysis of Crohn disease patients. Proc Natl
Acad Sci U S A 2008;105(43):16731-6. doi: 10.1073/pnas.0804812105.
433

452   **Table**

453   **Table 1.** Clinical parameters analyzed at baseline (S1) and 6 months after surgery

454   (S5). Asterisk indicates significant differences between S1 and S5 ($p < 0.05$)

455   calculated by t-test with false discovery rate (FDR) multiple testing correction.

	SG	RYGB
Gender	77.27% (female)	91.30% (female)
Age (years)	43.5 ± 12.03	44 ± 9.29
Type of surgery	48.80%	51.10%
Proton pump inhibitors	23.80%	26.08%
Diabetes Mellitus	9.09%	8.69%
BMI (kg/m2) S1	42.9 ± 6.56	43 ± 4.13
BMI (kg/m2) S2	38.91 ± 6.18	40.57 ± 4.19
BMI (kg/m2) S4	33.69 ± 5.51	35.07 ± 4.03
BMI (kg/m2) S5	30.81 ± 5.35*	31.52 ± 3.86*
vitD (nmol/l) S1	40 ± 16.39	42 ± 15.24
vitD (nmol/l) S5	73 ± 14.39*	73 ± 23.16*
vitB6 (nmol/l) S1	74 ± 33.34	66 ± 37.53
vitB6 (nmol/l) S5	104 ± 54.66	105 ± 46.99*
Cholesterol (mmol/l) S1	5.40 ± 0.88	5.47 ± 1.08
Cholesterol (mmol/l) S5	4.93 ± 0.68	4.05 ± 0.78*
Bilirubin (umol/ml) S1	5.70 ± 6.27	5.60 ± 2.44
Bilirubin (umol/ml) S5	9.30 ± 6.53	7.25 ± 4.08
HbA1c (mmol/mol) S1	39 ± 11.8	38 ± 5.48
HbA1c (mmol/mol) S5	32.50 ± 3.91	34 ± 4.35*
Iron (umol/l) S1	10 ± 6.17	12 ± 5.34
Iron (umol/l) S5	17 ± 8.27	16.50 ± 4.88
Ferritin (ug/l) S1	73 ± 232.83	84 ± 119.09
Ferritin (ug/l) S5	114 ± 197.36	108 ± 114.65
Folate (mmol/l) S1	15 ± 6.71	15 ± 5.94
Folate (mmol/l) S5	16 ± 6.11	31.50 ± 10.06*

1. Reference values. BMI: <18.5 (underweight), 18.5-24.9 (normal weight), 25-29.9

(overweight), >30 (obese). VitD: 50-100 nmol/l. VitB6: 34-156 nmol/l. Cholesterol:

3.5-6.5 nmol/l. Bilirubin: 3-21 umol/l. HbA1c: 20-42 mmol/l. Iron: 8-33 umol/l

(male), 5-30 umol/l (female). Ferritin: 25-250 ug/l (male), 20-150 ug/l (female).

Folate: > 6.8 mmol/l.

Figures Legends

Figure 1. Observed and estimated richness of gut microbiota at different time points

during the bariatric surgery procedure. (A). Shannon diversity index estimated a

decrease in bacterial richness at S2 and S3. (B). Rarefication curves showed a

reduction in bacterial richness at S2 and S3. (C). Principal component analysis (PCA)

plot of similarity between the samples; each dot represents 1 sample, each color a

different time point. S1. before surgery (red); S2. after 2 weeks of crash diet (orange);

S3. 1 week after surgery (yellow); S4. 3 months after surgery (light blue); S5. 6

months after surgery (dark blue).

Figure 2. Relative abundance of bacterial families in the gut microbiota at the five

time points analyzed. (A-G). Boxplots show the average relative abundance of 7

families that significantly changed between 2 different time points. (A).

Streptococcaceae. (B). Enterobacteriaceae. (C). Bifidobacteriaceae. (D).

Ruminococcaceae. (E). Rikenellaceae. (F). Veillonellaceae. (G).

O_Clostridiales_f_others. H. Relative abundance of all families identified at the

different time points. Significant families are represented in the same color. Asterisk

(in red) indicates significant fold change differences ($p < 0,05$) analyzed by ANCOM.

Figure 3. (A-C). Principal coordinate analysis (PCoA) plots comparing beta diversity

of Sleeve Gastrectomy (SG) versus Roux-Y Gastric bypass (RYGB) surgery at

baseline (S1) (A) 1 week after surgery (S3) (B) and 6 months after surgery (S5) (C).

SG is indicated in red, RYGB is indicated in blue (D). Relative abundance of bacterial

families in the gut microbiota at the five time points analyzed in SG versus RYGB

surgery.

Figure 4. Association between clinical parameters and family taxa calculated based

on the difference between 6 months after surgery (S6) and baseline (S1). The

significant association (false discovery rate (FDR) adjusted $p<0.05$) is indicated

with a "x". The red color indicates positive effect and the blue color negative effect.

HbA1c, glycated hemoglobin; VitD, vitamin D; VitB6, vitamin B6.