Neural Correlates of Causal Confounding

Mimi Liljeholm

Department of Cognitive Sciences, University of California, Irvine

<u>Corresponding author:</u>

Mimi Liljeholm

Department of Cognitive Sciences

2312 Social and Behavioral Sciences Gateway

University of California

Irvine, CA, 92697-5100

E-mail: m.liljeholm@uci.edu

Keywords: Causal induction, Uncertainty, Bayesian inference, prediction-errors, fMRI

Abstract

As scientists, we are keenly aware that if putative causes perfectly co-vary, the independent influence of neither can be discerned – a "no confounding" constraint on inference, fundamental to philosophical and statistical perspectives on causation. Intriguingly, a substantial behavioral literature suggests that naïve human reasoners, adults and children, are tacitly sensitive to causal confounding. Here, we used a combination of cognitive modeling and fMRI to investigate neural substrates mediating such sensitivity. While being scanned, participants observed and judged the influences of various fictitious allergy medicines on headache. Formal comparisons of a Bayesian causal model, sensitive to both confounding and stochasticity, and a purely error-driven algorithm, sensitive only to stochasticity, revealed dissociable neural representations of predictive uncertainty across several regions, including the insula, dlPFC and hippocampus. Implications for the detection and estimation of distinct forms of uncertainty, and for a neural mediation of domain general constraints on causal induction, are discussed.

Introduction

Consider two scenarios involving a target cause, C, an alternative cause, A, and the presence (+) or absence (-) of some relevant effect. In the first scenario, the effect occurs when C and A are presented in combination, but not when A is presented alone [A-, AC+]. In the second scenario, the A- trials are removed, so that both C and A occur only in combination with each other [AC+]. How would your judgment about the influence of C differ across these two scenarios? Philosophers, scientists and statisticians alike recognize that perfect co-variation of C and A in the second scenario occludes the independent influences of each putative cause, rendering any judgment about their respective effects fraught with uncertainty. Importantly, naïve human reasoners also appear to, tacitly, apply this domain-general constraint on causal induction, as evidenced by behavioral data showing sensitivity to confounding in a wide range of causal and predictive judgments, by children as well as adults (Spellman, 1996; Kushnir and Gopnik, 2005; Meder et al., 2006; Schulz and Bonawitz, 2007; Schulz et al., 2007; Liljeholm, 2015). In contrast, very little is known about the neural substrates mediating sensitivity to, and uncertainty associated with, causal confounding. The current study aims to identify such neural computations.

A second source of uncertainty in causal and predictive inference that has been thoroughly investigated both behaviorally and neurally is the stochasticity, or variance, of the outcome variable, which is greatest when the probability distribution over possible outcome states is uniform. As with causal confounding, behavioral sensitivity to outcome stochasticity is well established (e.g., Holt & Laury, 2002). Moreover, several neuroimaging studies have implicated the anterior insula, dorsomedial

prefrontal cortex (DMPFC) and dorsolateral prefrontal cortex (DLPFC) in the neural representation of outcome stochasticity (Volz et al., 2003; Huettel et al., 2005; Grinband et al., 2006; Abler et al., 2009). An important open question is whether, or which of, these neural regions also encode uncertainty associated with causal confounding.

Formally, two dominant approaches to predictive uncertainty can be discerned in the psychological literature. First, in associative learning theory, an error-driven representation of uncertainty about a cues predictive strength is captured by the "Pearce-Hall" (PH) algorithm, which relates the "associability", α, of a cue to a weighted average of the absolute prediction error on previous trials involving that cue:

$$\alpha_n = \gamma \left| \lambda_{n-1} - \sum V_{n-1} \right| + \left(1 - \gamma \right) \alpha_{n-1} \tag{1}$$

where ΣV_{n-1} is the expectation about the outcome summed over all present cues, λ_{n-1} is the observed state of the outcome on that trial, and γ is a free parameter accounting for the weighting of the previous trial relative to preceding ones. The estimated quantity, α, is proportional to the stochasticity of the outcome.

Conversely, in Bayesian Causal Models (BCM), uncertainty about the predictive strength of a cause is reflected in the entropy of the posterior distribution over its possible strengths, which depends on the variance in the effect variable, but also on the independent occurrence of putative causes. Specifically, for a scenario involving two candidate causes (e.g., C and A in the example above), reasoners make

judgments based on possible causal graphs (see Figure 1A) in which neither, either or

both C and A cause the effect, with all graphs assuming a constantly present

background cause of the effect, B.   The likelihood of a particular graph given the

observed data is derived using a particular parameterization that specifies how causes

combine to influence their effects.   The most commonly used parameterization for

generative causes is the noisy-OR (Cheng, 1997; Glymour, 1998).   For the graph in

which both C and A exert a generative influence on the effect, e, the probability of e is:

$$P\left(e^{+}|b,c,a;w_{B},w_{C},w_{A}\right)=1-\left(1-w_{B}^{b}\right)\times\left(1-w_{C}^{c}\right)\times\left(1-w_{A}^{a}\right) \qquad (2)$$

where c and a are 0 or 1 depending on the absence vs. presence of the candidate and

alternative cause, b, the background cause, is always 1 (i.e., present), and $w_{B,}$ w_{C} and

w_{A} are the respective strengths of putative.

While both the associative and Bayesian causal model predicts sensitivity to

stochasticity, only the causal model accounts for uncertainty due to causal

confounding.   A critical aspect of causal confounding is that, provided a sufficient

sample of trials with a particular confounded configuration, there is no logical or

empirical reason to expect a prediction about the outcome given that stimulus

configuration to be inaccurate.   This property uniquely distinguishes causal

confounding, not just from stochasticity, which scales linearly with outcome prediction

errors, but also from other sources of uncertainty, such as ambiguity and volatility. Of

particular interest, therefore, is whether uncertainty due to causal confounding is

discernable during feedback-based learning, as participants make predictions, and

receive feedback, about the occurrence of the effect given a confounded cause. Here,

a combination of neuroimaging and cognitive modeling is used to dissociate neural

signals scaling with error-driven uncertainty from those encoding uncertainty

associated with causal confounding.

Materials & Methods

Participants. 20 healthy normal volunteers (mean age = 20.9 ± 2.4, range: 18-27, 12

females) participated in the study. In our previous work (e.g., Liljeholm et al., 2011;

Liljeholm et al., 2013), this sample size has proven sufficient to generate effects based

on a univariate model-based fMRI analysis at appropriate corrected thresholds. One

participant was excluded prior to any analyses due to excessive head movement (>

5mm), leaving a sample size of 19. The volunteers were pre-assessed to exclude

those with a history of neurological or psychiatric illness. All subjects gave informed

consent and the study was approved by the Institutional Review Board of the

University of California, Irvine.

Task & procedure. Participants were scanned with functional MRI while performing

a causal induction task in which they assumed the role of a research scientist

assessing the influence of various allergy medicines on headache, a potential side

effect. At the beginning of the study, participants were instructed that each medicine

could either produce headache or have no influence on headache (i.e., there were

no preventive causes) and, further, that the influence of a given medicine might be

stochastic, so that even if that medicine was indeed a cause of headache, it may still

not produce headache every time it was administered. The study focused on three

target medicines, C, D and S, mnemonically labeled here to indicate "confounded",

"deterministic" and "stochastic" influences respectively, with each target medicine

occurring only in combination with some alternative medicine during feedback-based

learning. Target conditions, relevant events and model predictions are listed in

Table 1. With +, - and ± respectively indicating a 1.0, 0 and 0.5 probability of the

effect, and with lower case letters indicating non-target causes, there were 5

different types of medicine treatments: e-, s'±, De+, Se± and Cc'+. None of the

allergy patients had headache in the absence of any medicine, which was explicitly

stated in the initial instructions as well as apparent on several "no medicine" (n-)

trials.

Table 1: Conditions, relevant events and model predictions about uncertainty during feedback-based learning. C, D and S respectively indicate confounded, deterministic and stochastic target causes; e and c' indicate non-target compound elements; +, - and ± respectively indicate a 1.0, 0.0 and 0.5 probability of the effect.

Condition	Relevant events	Target compounds	Uncertainty	
			Causal	Error-driven
Confounded	n-, Cc'+	Cc'	6.97	0.19
Deterministic	n-, e-, De+	De	4.26	0.24
Stochastic	n-, e-, Se±	Se	6.29	0.48

Note that medicines C and c' both occurred *only* in combination with one another, so

that target medicine C was perfectly confounded with its compound counterpart.

Medicine D also occurred only in combination with another medicine, e, however,

"elemental" medicine e also occurred by itself, allowing for an estimation of the

independent causal influence of medicine D across the constant presence of e (i.e.,

across e- and De+ trials). Medicine S was identical to medicine D, except that the

probability of headache given the Se compound was 0.5 rather than 1.0; a

comparison of medicines S and D, therefore, identifies a difference between

deterministic and stochastic causation. Finally, medicine s' also produced headache

with a probability of 0.5 but unlike medicine S, never occurred in combination with

any other medicine; contrasting S with s' identifies differences due to compound

presentation.

On each trial of feedback-based learning (see Figure 1A), participants were

presented with an individual allergy patient and were told either that the patient had

not received any medicine or that a particular medicine, or combination of medicines,

had been administered. Medicines were color coded and labeled accordingly (i.e.,

Medicine "B" was blue) with color assignments counterbalanced across participants.

On the first screen, the allergy patient's state was obscured by a question mark and

participants were asked to indicate whether or not the patient had a headache,

pressing the ''Y'' key for ''yes'' and the ''N'' key for ''no''.   After a prediction was

made, there was a brief (2000ms) pause during which the word "wait" was displayed

on the screen, followed by a screen revealing the allergy patient's state (i.e., with or

without headache). Each of 6 distinct trial types was presented 8 times in each of 4

sessions, for a total of 192 trials, with sessions separated by 2-minute breaks, during

which the scanner was turned off. Each session was further divided into 4, non-

delineated, blocks, with each type of trial occurring once in random order in each

block. All trials were separated by a jittered 4-second ITI. In every other block,

starting with the 2^{nd} block of the 1^{st} session, each trial (except for n- trials) was

followed by a query regarding the causal influence of the target medicine presented

on that trial (i.e., C, D or S), or of medicine e or s' on non-target trials. Specifically,

participants were asked to choose between the following three options regarding the

relevant medicine: "is a cause of headache", "is not a cause headache" and "may or

may not be a cause of headache" (see Figure 1B).

Figure 1. Task illustration. A. Prediction and outcome screens from a feedback-based learning trial. Screens were separated by a 2 sec. pause during which the word "Wait" (not shown in figure) was displayed. B. Causal query regarding the individual influence of a particular medicine on headache.

Computational models. Two formal accounts of predictive strength and uncertainty

were implemented. First, an error-driven, associative learning algorithm updates

predictive strength, on each trial, using the difference between the observed state of

the outcome and the expected state based on all present cues:

$$V_{n+1} \leftarrow V_n + \alpha_n(\lambda_n - \sum V_n) \qquad\qquad (3)$$

where V is the predictive strength of a cue, and all other terms, including α_n, are

defined as in Equation 1 above. In addition to scaling the influence of the prediction

error on predictive strength, α is taken to reflect current levels of uncertainty about the

cues influence. Note that, because of the Σ term in Equations 1 and 3, both the

uncertainty (α), and strength (V) of a cue depend on the effects of other cues with

which it is paired. Specifically, in the current study, the variance in the occurrence of

headache across e- and De+ trials, and on Se± trials, results in a constantly

fluctuating strength of Medicine e and, consequently, in persistent prediction errors,

and corollary changes to strength and uncertainty, on e-, as well as De+, trials (see

model predictions in Table 1).

The error-driven approach further assumes that, during feedback-based learning,

participants select "Yes"/"No" responses to questions about whether the outcome will

be present using probabilities generated by a softmax distribution, in which a free

parameter controls the influence of predictive strength (i.e., the strength with which a

cue predicts the outcome) on choice behavior. Free parameters were fit to behavioral

data by minimizing the negative log likelihood of observed responses for each

individual. On feedback-based trials with target causes, which always occurred in

compound with an alternative cause, the α of the target cause (always greater than or

equal to that of its compound counterpart) was used to indicate uncertainty, while the

summed strength, ΣV, was used to predict the outcome, and to compute the prediction

error.

A second formal account is provided by a Bayesian Causal Model (BCM), in which

reasoners make inferences about which of several causal structures (*G1-G4*) is

responsible for the observed data (Griffiths & Tenenbaum, 2005; Liljeholm, 2015).

Support for a given target cause, C, is defined as the log likelihood ratio in favor of

graphs in which there is a link between that cause and the effect (e.g., *G1* and *G3*),

over graphs in which there is no such link (e.g., *G0* and *G2*).

$$\sup_{C \to E} = \log \frac{P(D|G1) + P(D|G3)}{P(D|G0) + P(D|G2)}.$$

(4)

In the current study, sequential inferences were modeled using the smallest possible

focal set (Novick & Cheng, 1990) – that is, the smallest set of events relevant for

making the inference. Thus, inferences about elemental (non-target) candidate

causes s' and e were modeled based on trials with those respective causes and the

"no causes" trials (i.e., n- and s'± for candidate s', and n- and e- for candidate e), using

only *G0* and *G1*. For each target candidate cause, which always occurred in

compound with some alternative cause (see Table 1), inferences were modeled over

all four graphs (as in Equation 4) using the trials relevant for the particular target

cause (i.e., n- and Cc' for target C; n-, e- and De+ for target D, and n-, e- and Se± for

target S). Each graph has a set of parameters θ, which are the causal strengths w of

the background (w_B), target candidate cause (w_C) and alternative candidate cause (w_A)

respectively. The likelihoods, $P(D|G_i)$, are obtained by averaging out the relevant

unknown parameters, such that, for *G0*

$$P(D|G0) = \int_0^1 P(D|w_B, G0) P(w_B|G0) dw_B,$$

(5)

210 for *G1* (or *G2*, replacing *C* with *A*)

211 $$P(D|G1) = \int_0^1 \int_0^1 P(D|w_B, w_C, G1) P(w_B, w_C|G1) dw_B \, dw_C,$$

(6)

212 and for G3

213 $$P(D|G3) = \int_0^1 \int_0^1 \int_0^1 P(D|w_B, w_C, w_A, G3) P(w_B, w_C, w_A|G3) dw_B \, dw_C \, dw_A,$$

(7)

214   where $P(\theta|G_i)$ denotes the priors on causal strength parameters.

215   The likelihoods $P(D|\theta, G_i)$ are computed using the noisy-OR parameterization (Cheng,

216   1997). Specifically, summarizing data D by contingencies $N(e,c,a)$, the frequencies of

217   each combination of the presence versus absence of the effect, target cause and

218   alternative cause, the likelihood term for $G3$ is:

$$P(D|w_B,w_C,w_A,G3) = \begin{pmatrix} N(c^-,a^-) \\ N(e^+,c^-,a^-) \end{pmatrix} \times \begin{pmatrix} N(c^+,a^-) \\ N(e^+,c^+,a^-) \end{pmatrix} \times \begin{pmatrix} N(c^-,a^+) \\ N(e^+,c^-,a^+) \end{pmatrix} \times \begin{pmatrix} N(c^+,a^+) \\ N(e^+,c^+,a^+) \end{pmatrix} w_B^{N(e^+,c^-,a^-)}$$

219
$$(1-w_B)^{N(e^-,c^-,a^-)}[w_B+w_C-w_Bw_C]^{N(e^+,c^+,a^-)}[1-w_B-w_C+w_Bw_C]^{N(e^-,c^+,a^-)}[w_B+w_A-w_Bw_A]^{N(e^+,c^-,a^+)}[1-w_B-w_A+w_Bw_A]^{N(e^-,c^-,a^+)}$$ (8)
$$[w_B+w_C+w_A-w_Bw_C-w_Bw_A-w_Cw_A+w_Bw_Cw_A]^{N(e^+,c^+,a^+)}[1-w_B-w_C-w_A+w_Bw_C+w_Bw_A+w_Cw_A-w_Bw_Cw_A]^{N(e^-,c^+,a^+)}.$$

where $\binom{n}{k}$ denotes the number of ways of picking k unordered outcomes from n

possibilities. $N(c^+)$ indicates the frequency of events in which the target cause is

present, with analogous definitions for the other $N(.)$ terms. The likelihood terms for

$G0, G1$ and $G2$ are similarly specified, where frequencies $N(.)$ are summed across the

presence and absence of relevant events, and $w_i=0$ for any cause that does not have

link to the effect in the relevant graph.

Causal strength estimates for each elemental and target cause were modeled as the

mean of the marginal posterior distribution $P(w_C|D)$ from the most likely graph in which

a link existed between the cause and the effect. The marginal posterior distribution is

obtained by applying Bayes' rule, such that, if for example $G3$ had the highest

likelihood, then

$$P(w_C|D,G3) = \int_0^1\int_0^1 \frac{P(D|w_B,w_C,w_A,G3)P(w_B,w_C,w_A|G3)}{P(D)}dw_B\,dw_A,$$ (9)

where $P(D|w_B, w_C, w_A, G3)$ is the likelihood term, $P(w_B, w_C, w_A|G3)$ refers, again, to the prior

probabilities of causal strength parameters, and $P(D)$ is the normalizing term, denoting

the probability of the observed data. The mean of w_C is given by

235
$$\bar{w}_C = \int_0^1 w_C P(w_C \mid D) dw_C.$$
(10)

236 Uncertainty about the strength of a particular cause, the focus of the study, was

237 modeled as the Shannon entropy $H(w_c)$ of the marginal posterior distribution $P(w_C|D)$:

238
$$H(w_C) = -\int_0^1 P(w_C \mid D) \log P(w_C \mid D) dw_C.$$
(11)

The model was sequentially implemented on a trial-by-trial basis, such that, for each

cause, likelihoods were computed on each trial of feedback-based learning that

yielded information relevant to that cause, using only the data point provided on that

trial, and using the posterior on the previous trial as the prior on the current trial. On

the first trial in which data relevant to a particular candidate cause was presented, the

priors were assigned independent uniform, $Beta(1,1)$, distributions. Finally, as an

analog to the model-free prediction error, the Kullback-Leibler (KL) divergence

between the prior (Q) and posterior (P) on each trial was computed as:

247
$$KL(P\|Q) = \int_0^1 p(w_C \mid D) \log \frac{p(w_C \mid D)}{q(w_C \mid D)} dw_C.$$
(12)

As noted, target causes only occurred in compound with alternative causes during

feedback-based learning. On such trials, the entropy and KL divergence of the

marginal distribution for the target cause was used to indicate uncertainty and surprise

respectively, while predictions regarding the occurrence of the effect were generated

using the noisy-OR integration of strengths specified in Equation 2.

Imaging procedure and analysis. A 3 Tesla scanner (Phillips Achieva) was used to

acquire structural T1-weighted images and T2*-weighted echoplanar images

(repetition time = 2.65 s; echo time = 30 ms; flip angle = 90°; 45 transverse slices;

matrix = 64 × 64; field of view = 192 mm; thickness = 3 mm; slice gap = 0 mm) with

BOLD contrast. The first four volumes of images were discarded to avoid T1

equilibrium effects. All remaining volumes were corrected for differences in the time of

slice acquisition, realigned to the first volume, spatially normalized to the Montreal

Neurological Institute (MNI) echoplanar imaging template, and spatially smoothed with

a Gaussian kernel (8 mm, full width at half-maximum). High-pass filtering with a cutoff

of 128 s was used.

For each subject, a general linear model (GLM) was specified, with regressors in each

session modeling 1) the onset of the prediction screen on each trial during feedback-

based learning, 2) the onset of the outcome screen on each trial during feedback-

based learning, and 3) the onsets of the screens soliciting judgments about the

influences of individual medicines. Prediction-screen onsets and causal judgment

onsets were modulated, in order, by trial-by-trial error-driven and causal estimates of

uncertainty and strength. Outcome-screen onsets were modulated by trial-by-trial

error-driven and causal estimates of surprise. No orthogonalization was applied. Two

separate regressors modeled the motor responses on each feedback-based and

causal query trial, respectively, and six additional regressors accounted for the

residual effects of head motion. All regressors were convolved with a canonical

hemodynamic response function. To contrast error-driven and causal accounts,

disjunction tests were performed, assessing neural effects that were significant, based

on whole brain correction, given one (e.g., the causal) account but that did not

emerge, in either a positive or negative direction, given the alternative (e.g., error-

driven) account even at an uncorrected threshold of *p<0.1*.

To address loss of power due to multicollinearity, Bayesian model selection (BMS)

analyses were used to further arbitrate between error-driven and causal accounts.

Here, for each subject, two separate GLMs were specified, one for the error-driven

approach and another for the causal approach. Thus, these GLMs were identical to

that described above, except that only error-driven variables were entered as

parametric modulators in the first GLM, and only "causal" variables in the second

GLM. A first-level Bayesian estimation procedure (SPM 12) was used to compute a

log-model evidence map for every subject and each GLM (Penny et al., 2005).

Inferences at the group level were modeled by applying a random effects approach

(Rose et al., 2010) at every voxel of the log evidence data falling within a particular

functional mask, specified based on the results of classical estimation (e.g., inferences

about uncertainty during prediction screens were constrained by a mask of all voxels

that reached significance using whole-brain (CST) correction with either the error-

driven or causal GLM). Group-level exceedance probability maps, reflecting the

probability that one model is more likely than the others, were thresholded to identify

regions in which the exceedance probability was greater than 0.95.

While it effectively addresses colinearity between causal and error-driven models, the

BMS pertains to *all predictions* made by a particular model, discriminating neither

between different computational variables (i.e., uncertainty, strength and surprise) nor

between different experimental events (i.e., feedback-based learning vs. explicit

causal judgments). Given this limitation, we constrain our reporting of BMS results in

the text to whether they confirm the disjunction results (for completeness, a full lists of

all BMS results are provided in Supplementary Tables 1-4), and focus our discussion

on regions that are identified across both disjunction and BMS tests. To further

facilitate the interpretation of disjunction results, we plot neural responses to each

target cause, as well as to increasing values of computational variables, during

feedback-based learning, extracting effect sizes from 8mm spheres centered on peak

MNI coordinates, using rfxplot (Glascher, 2009).

Results

Behavioral results

All statistical tests of behavioral data were planned comparisons, employing two-tailed

t-tests, and were calculated using n=19. Confidence intervals (*CI*) and effect sizes

(Cohen's *d*) are reported for all comparisons.

Figure 2. Mean proportions of "Is cause", "Is not a cause" and ambivalent "May or may not be cause" judgments about the individual influence of each medicine on headache. Target causes (C, D & S) occurred only in compound with alternative causes during feedback-based learning, while non-target causes occurred only individually (s'), or individually as well as in compound (e).

The mean proportions of "is a cause", "is not a cause" and, ambivalent, "may or may

not be a cause" judgments, averaged across multiple probes for each probed

medicine, are shown in Figure 2. Planned comparisons of target causes revealed that

the mean proportion of ambivalent judgments were significantly lower for the

deterministic (D) medicine than for both confounded (C), $t(18)=6.78$, $p<0.001$,

CI=[0.35 0.67], d=1.79, and stochastic (S), $t(18)=4.09$, $p<0.001$, *CI*=[0.14 0.44],

d=1.11, medicines. The difference between the confounded and stochastic medicine

was also significant, with the mean proportion of ambivalent judgments being greater

for C than for S $t(18)=2.65$, $p<0.05$, *CI*=[0.0.05 0.40], d=0.68. Recall that all target

medicines, including S, occurred only in compound with alternative medicines,

controlling for cue complexity across target comparisons. The proportion of ambivalent

judgments for the two stochastic causes, one always occurring on its own (s') and the

other always in a compound (S) during feedback based learning, was only marginally

significant, *p=0.08*, suggesting that cue complexity did not strongly influence

perceived causal ambiguity.

328 Neuroimaging results

Figure 3. Disjunction results favoring the causal model of uncertainty during the prediction period of feedback-based learning trials, showing effects in the anterior insula (top) and DMPFC (bottom). Plotted values show effect sizes (y-axis) extracted from spheres centered on peak coordinates at low (L), medium (M) and high (H) levels of causal (blue) and error-driven (red) models (middle column), and for confounded (C), deterministic (D) and stochastic (S) compound causes (right column). Error bars=SEM.

Uncertainty in causal prediction and judgment. During feedback-based learning

(Figure 3 and Table 2), as participants generated a prediction about whether or not

headache would occur given a particular medicine or combination of medicines,

disjunction analyses revealed significant effects of the causal, but not the error-driven,

measure of uncertainty in the bilateral anterior insula, bilateral fusiform gyrus, anterior

cingulate and rostrolateral prefrontal cortex.

Table 2: Results of disjunction tests in favor of the causal model. Peak MNI coordinates and cluster sizes (>= 142) at p<0.005. R/L = Right/Left, (-) = negative correlation, ' ' = continuous with previous cluster.

Contrast	Area	x, y, z	Cluster size at p<0.005
Prediction Uncertainty	L. Fusiform Gyrus	-34, -50, -20	858
	L. Anterior Insula	-40, 16, -8	672
	L. Rostrolateral Prefrontal Cortex	-34, 52, 12	' '
	R. Fusiform Gyrus	-38, -50, -14	202
	R. Anterior Insula	38, 30, -6	166
	Anterior Cingulate Cortex	8, 44, 22	233
Prediction Strength	Calcarine/Lingual	14, -92, -4	1109
Surprise	R. Angular Gyrus	54, -60, 32	1564
	R. Middle Temporal Gyrus	64, -46, 10	' '
	R. Anterior Insula	38, 18, 0	1294
	R. Inferior Frontal Gyrus	34, 14, 30	' '
	L. Anterior Insula	-30, 22, 0	952
	Anterior Cingulate Cortex	8, 44, 26	733
	Ventral Tegmental Area	4, -22, -26	622
Surprise (-)	L. Caudate	-14, 28, -2	846
	R. Superior Parietal Lobule	20, -58, 60	1086
	R. Hippocampus	32, -38, 4	241
	L. Parahippocampus	-34, -42, 0	258
	R. Supramarginal Gyrus	58, -20, 38	243
Judgment Uncertainty	Dorsomedial Prefrontal Cortex	16, 58, 32	2507
	R. Dorsolateral Prefrontal Cortex	32, 22, 28	143

Conversely, the error-driven, but not causal, measure of predictive uncertainty (Figure

4 & Table 3) was significantly positively correlated with activity in large aspects of the

lingual gyrus, the left inferior parietal lobule, and the dorsolateral prefrontal cortex, as

well as negatively correlated with activity in the bilateral middle temporal gyrus (MTG),

the right temporal-parietal junction (TPJ) and ventromedial prefrontal cortex (vmPFC).

Note, however, that the negative correlation in the vmPFC reflects increased activity in

response to the confounded cause (bottom right of Figure 4), suggesting that this

region is in fact selectively responding to causal confounding. The BMS analysis

(Supplemental Table 1) confirmed the selective effects of the causal model in the

bilateral insula, anterior cingulate and fusiform gyri, as well as the selective effects of

the error-driven algorithm in the vmPFC, lingual gyrus and dorsolateral prefrontal

cortex.

Figure 4. Disjunction results favoring the error-driven model of uncertainty during the prediction period of feedback-based learning trials, showing effects in the DLPFC (top), lingual gyrus (middle) and vmPFC (bottom). Plotted values show effect sizes (y-axis) extracted from spheres centered on peak coordinates at low (L), medium (M) and high (H) levels of causal (blue) and error-driven (red) models (middle column), and for confounded (C), deterministic (D) and stochastic (S) compound causes (right column). Error bars=SEM.

During causal queries, as participants made judgments about the causal influences of

individual medicines, disjunction analyses revealed extensive effects of the causal

measure of uncertainty throughout the DMPFC, as well as more modest effects in the

right DLPFC, while a selective negative correlation with the error-driven measure

again emerged in the vmPFC. BMS analyses (Supplemental Table 2) confirmed that

the causal model provided a better account of activity throughout the DMPFC.

Table 3: Results of disjunction tests in favor of the error-driven algorithm. Peak MNI coordinates and cluster sizes (>= 142) at p<0.005. R/L = Right/Left, (-) = negative correlation.

Contrast	Area	x, y, z	Cluster size at p<0.005
Prediction Uncertainty	Lingual Gyrus	0, -72, 2	1556
	L. Inferior Parietal Lobule	-32, -60, 44	177
	R. Dorsolateral Prefrontal Cortex	44, 26, 26	143
Prediction Uncertainty (-)	Ventromedial Prefrontal Cortex	2, 26, -10	471
	L. Middle Temporal Gyrus	-52, -12, -22	450
	L. Middle Temporal Gyrus	-44, -66, 22	503
	R. Middle Temporal Gyrus	44, -62, 20	340
	R. Middle Temporal Gyrus	60, -54, 2	177
	R. Temporal Parietal Junction	50, -38, 24	175
Surprise	Posterior Cingulate Cortex	-10, -32, 24	149
	R. Inferior Temporal Gyrus	40, -54, -8	151
Judgment Uncertainty (-)	Ventromedial Prefrontal Cortex	-10, 36, -10	154

Measures of surprise. While uncertainty in causal inferences, due to confounding or

stochasticity, is of primary interest here, estimates of the strength of the relationship

between a cause and effect, and of the surprise at the outcome on each trial, were

also modeled from both a causal and error-driven perspective. We focus our reporting

and discussion of results on the measures of surprise, rather than strength: First,

because causal and error-driven measures of strength were highly correlated (mean r

= 0.87) and, second, because, to the extent that they differed, the divergence is

largely incidental – using a linear, rather than noisy-OR, integration function in the

Bayesian causal model would render strength measures virtually identical across

causal and error-driven models, while preserving differences in uncertainty due to

stochasticity vs. confounding (see discussion). For completeness, significant results

of all comparisons of strength measures are listed in relevant tables (i.e., Table 2 for

disjunction tests and Supplemental Table 3 for model selection).

Figure 5. Disjunction results favoring the causal model of surprise, in the dorsomedial prefrontal cortex (top) and hippocampus (bottom). Line graphs show effect sizes (y-axis) extracted from spheres centered on peak coordinates at low (L), medium (M) and high (H) levels of causal (blue) and error-driven (red) models of surprise.

For measures of surprise, disjunction tests identified activity positively correlated with

the causal model (i.e., the information theoretic distance between the prior and

posterior strength distribution on each trial) in the bilateral anterior insula, anterior

cingulate cortex and right middle temporal, inferior frontal and angular gyri, and the

ventral tegmental area.   Activity negatively correlated with the causal measure of

surprise was revealed in the bilateral hippocampus, caudate and right superior and

inferior parietal lobules. In contrast, selective effects of the model-free prediction error

emerged in the posterior cingulate cortex and right inferior temporal gyrus.   BMS

analyses (Supplemental Table 4) confirmed the selective effects of causal surprise in

the bilateral insula, anterior cingulate cortex, bilateral hippocampus and ventral

tegmental area.

Discussion

Sensitivity to the "no confounding" constraint – a requirement that putative causes do

not co-vary, fundamental to philosophical and statistical perspectives on causation –

has been reliably demonstrated in a range of behavioral judgments by children as well

as adults (Spellman, 1996; Kushnir and Gopnik, 2005; Meder et al., 2006; Schulz and

Bonawitz, 2007; Schulz et al., 2007; Liljeholm, 2015).   The current study combined

computational cognitive modeling with fMRI to investigate neural substrates encoding

uncertainty associated with causal confounding. While being scanned, participants

studied the influences of various fictitious allergy medicines on headache – a side

effect. Three target medicines with respectively confounded (C), deterministic (D) and

stochastic (S) influences each occurred only in compound with an alternative medicine

during feedback-based learning, ruling out contingency and cue complexity as sources

of any effects of causal confounding.   Moreover, unlike stochasticity, uncertainty

associated with confounding during feedback-based learning could not be attributed to

experienced or anticipated errors.   Formal model comparisons indicated that a

Bayesian causal model, sensitive both to stochasticity and confounding, provided a

better account of activity in the anterior insula and dorsomedial prefrontal cortex

(DMPFC) than did an error-driven, model-free, algorithm sensitive only to

stochasticity; conversely, the error-driven model selectively elicited activity in the right

DLPFC and posterior lingual gyrus.

The anterior insula has been implicated in encoding outcome stochasticity across a

variety of tasks (e.g., Volz et al., 2003; Huettel et al., 2005; d'Acremont et al., 2013;

Preuschoff et al., 2008; Critchley et al., 2001; Grinband et al., 2006). For example,

scanning participants with fMRI as they classified stimuli according to a variable

category boundary, Grinband et al. (2006) found that activity in the anterior insula, as

well as the DMPFC, scaled with categorization uncertainty – the proximity of a

stimulus to a category boundary. An important aspect of this and other

demonstrations of the role of the anterior insula in encoding outcome stochasticity is

that the uncertainty is directly related to experienced or inferred errors: that is, the

greater the outcome stochasticity on a trial, the more likely it is that the participants

prediction, or decision, on that trial will be incorrect. In contrast, during feedback-

based learning in the current task, the outcome on confounded trials was

deterministic. Consequently, unlike previous work, the increase in anterior insular and

DMPFC activity as participants generated predictions about the outcome on trials with

the confounded cause cannot be attributed to the experience or anticipation of errors.

This strongly suggests that uncertainty representations in these regions may

implement generative models of predictive knowledge, dissociable from the evaluation

of immediate consequences of choice.

Activity in the vmPFC selectively responded to the confounded compound during

feedback-based learning. As with the insula and DMPFC, comparisons with other

target causes rule out stimulus complexity (since all target causes occurred only in

compound with alternative causes during feedback-based learning) as well as the

deterministic nature of the outcome (since that was also the case for target cause D)

as sources of the effect. Moreover, the lack of a response to the stochastic target

cause suggests that this region is sensitive to the presence of confounding, rather

than to uncertainty per se. Notably, causal confounding is an abstract concept,

irreducible to either elemental contingency or contiguity. In a study by Kumaran et al

(2009), in which the conjunction of two stimuli, or of a stimulus and a location,

predicted a weather (sun vs. rain) outcome, the vmPFC, and its interaction with the

hippocampus, was implicated in the acquisition and transfer of conceptual stimulus-

outcome associations (see Bowman and Zeithamova, 2018, for related results). Here,

particularly in light of our hippocampal findings discussed below, we conjecture that

the vmPFC and hippocampus are similarly, albeit in this case futilely, attempting to

extract generalizable knowledge from confounded compound trials.

In addition to scaling with causal uncertainty, activity in the anterior insula and

anterior cingulate cortex, together with the bilateral hippocampus, correlated with the

Kullback-Liebler divergence between the posterior and prior distributions, a variable

termed "Bayesian surprise" (Itti & Baldi, 2006). Notably, surprise at the outcome of a

particular trial during feedback-based learning is fundamentally related to how the two

modeling approaches treat uncertainty. In the error-driven model, the uncertainty (or

associability) of a cue is proportional to the history of prediction errors associated with

that cue; in other words, the greater the long run average surprise, the greater the

uncertainty.   In contrast, in the causal model, while a high level of surprise (i.e., large

KL divergence between the posterior and prior distribution) on a given trial, due either

to outcome stochasticity or to a change in the underlying statistics of the environment,

may result in greater uncertainty (i.e., greater entropy of the posterior distribution) a

confounded deterministic cause generates very low levels of surprise, after a few

initial acquisition trials, but is still associated with very high levels of uncertainty.

Another notable distinction between the models contrasted here is that surprise in the

causal model is unsigned; conversely, in the error-driven model, the unexpected

presence and absence of the effect respectively yield a positive and negative

prediction error. Thus, in the current study, selective effects of the causal measure of

surprise may reflect a neural implementation of a full generative model, consistent with

the effects observed for uncertainty, or simply an unsigned prediction error.   Further

work is needed to arbitrate between these possibilities.

Several previous studies have implicated the DLPFC, together with the anterior insula

and DMPFC, in encoding uncertainty related to outcome stochasticity (Huettel et al.,

2005; Abler et al., 2009; Causse et al., 2013). The specificity of this area with respect

to our comparison of error-driven and causal sources of uncertainty is somewhat

unequivocal; In particular, based on the disjunction tests, the right DLPFC was

selectively recruited by the error-driven measure of uncertainty during feedback-based

learning, but then significantly modulated by the causal measure of uncertainty during

judgments about the existence of causal relationships.   A possible explanation for

this pattern of results is that the right DLPFC reflects the influence of uncertainty on

behavior – recall that uncertainty about the causal influence of the confounded target

cause is only behaviorally relevant during explicit judgments about its causal status;

confounded trials during feedback-based learning deterministically and unambiguously

predict the presence of the effect. Indeed, several previous studies have used neural

disruption techniques to demonstrate that the right DLPFC modulates behavioral

responses to uncertainty (Knoch et al., 2006; Fectau et al., 2007; Ye et al., 2015).

Nonetheless, we caution the reader that only the selective effects of the error-driven

model during feedback-based learning were observed across disjunction and model

selection analyses.

Selective effects of the error-driven account of predictive uncertainty were also

identified throughout large portions of the right posterior lingual gyrus. In a recent

neuroimaging study, Causse et al., (2013) assed decision making under uncertainty

using an aviation task in which participants had to either land a plane or abort landing,

given certain (100% or 0%) vs. uncertain (50%) information regarding landing

conditions. Consistent with the current results, they found greater activity in the right

posterior lingual gyrus, as well as in the DLPFC, during high than during low

uncertainty conditions.   Our results are also consistent with those of Payzan-

LeNestour et al. (2016), who found that the lingual gyrus scaled specifically with

uncertainty due to outcome stochasticity, as opposed to uncertainty due to changes in

the statistical structure of the environment or due to small samples. Payzan-

LeNestour et al. argued that the lingual gyrus might not typically emerge in studies

assessing outcome stochasticity because those studies did not dissociate stochasticity

from other uncertainty components. A similar case can be made here, that the

inclusion of an alternative, generative, model that captures additional aspects of

uncertainty may have increased sensitivity to detect stochasticity-specific signals.

Several neuroeconomic studies have contrasted gambles involving "risk" vs.

"ambiguity", corresponding respectively to stochastic and unknown probabilities (Hsu

et al., 2005; Huettel et al., 2006; Bach et al., 2009; Levy et al., 2010). For example,

when flipping a coin, you might know that the probability of heads is 0.5 (risk) or, in the

case of a biased coin, you may have no knowledge at all about the probability of

heads (ambiguity) – in either case, subjectively, heads and tails are equally likely.

One might argue that, in the current study, the influence of the confounded medicine is

unknown, just as the outcome probabilities of ambiguous gambles are unknown. But

what does it mean to "know" something? Critically, all target medicines occurred only

in   compound   with   alternative   medicines   during   feedback-based   learning:

consequently, when presented individually during causal queries, each was equally

novel and, presumably, equally unknown. In other words, differences in ambiguity

between confounded and independently occurring causes cannot be attributed to a

generalization decrement based on changes in stimulus features. Moreover, as with

stochasticity, ambiguity, but not confounding, should elicit high levels of error

expectation during outcome prediction. Finally, ambiguity and confounding call for

very different plans of action: in the case of ambiguity, but not confounding, a simple

increase in sample size (i.e., repeated observation of the same contingency) can

resolve the uncertainty. Further work is needed to assess the overlap between neural

representations of confounding and ambiguity.

As the mantra "covariation does not equal causation" implies, requirements for causal

induction extend far beyond mere statistical regularity. Countless studies have

demonstrated neural and behavioral effects of error-driven uncertainty, whether due to

stochasticity, insufficient samples or changes in the statistical structure of the

environment. The current study reveals that, given violation of a basic boundary

condition on causal inference, predictive uncertainty can emerge in the presence of a

deterministic, unchanging and well-sampled contingency. These results contribute to

a growing literature on the role of generative models in neural representations of

uncertainty, and shed light on a possible neural implementation of domain general, a

priori, constraints on causal induction.

References

Abler, B., Herrnberger, B., Grön, G., & Spitzer, M. (2009). From uncertainty to reward:

BOLD characteristics differentiate signaling pathways. BMC neuroscience, *10*(1),

154.

Bach, D. R., Seymour, B., & Dolan, R. J. (2009). Neural activity associated with the

passive prediction of ambiguity and risk for aversive events. Journal of

Neuroscience, *29*(6), 1648-1656.

Barthelmé, S., & Mamassian, P. (2009). Evaluation of objective uncertainty in the

visual system. *PLoS computational biology*, *5*(9), e1000504.

Beckers T, De Houwer J, Pineno O, Miller RR (2005) Outcome additivity and outcome

maximality influence cue competition in human causal learning. Journal of

experimental psychology Learning, memory, and cognition 31:238-249.

Been SL, Mitchell, C. J., Bouton, M. E., & Frohardt, R. (2003) Forward and backward

blocking of causal judgment is enhanced by additivity of effect magnitude. Memory

& Cognition 31:133-142.

Bowman, C. R., & Zeithamova, D. (2018). Abstract memory representations in the

ventromedial prefrontal cortex and hippocampus support concept

generalization. *Journal of Neuroscience*, 2811-17.

Causse, M., Péran, P., Dehais, F., Caravasso, C. F., Zeffiro, T., Sabatini, U., & Pastor,

J. (2013). Affective decision making under uncertainty during a plausible aviation

task: An fMRI study. *NeuroImage*, *71*, 19-29.

Cheng P (1997) From covariation to causation: A causal power theory. Psycholgoical

Review:367-405.

Cheng, P., Liljeholm, M., & Sandhofer, C. (2017) Analytic causal knowledge for

constructing useable empirical causal knowledge: Two experiments on

preschoolers. In G. Gunzelmann, A. Howes, T. Tendbrink, & E.J. Davelaar (Eds.),

Proceedings of the 39th Annual Conference of the Cognitive Science Society (pp.

1758-1763). Austin, TX: Cognitive Science Society.

Corlett, P. R., Aitken, M. R., Dickinson, A., Shanks, D. R., Honey, G. D., Honey, R. A.,

... & Fletcher, P. C. (2004). Prediction error during retrospective revaluation of

causal associations in humans: fMRI evidence in favor of an associative model of

learning. *Neuron*, *44*(5), 877-888.

Critchley, H. D., Mathias, C. J., & Dolan, R. J. (2001). Neural activity in the human

brain relating to uncertainty and arousal during anticipation. *Neuron*, *29*(2), 537-

545.

Chumbley, J. R., Flandin, G., Bach, D. R., Daunizeau, J., Fehr, E., Dolan, R. J., &

Friston, K. J. (2012). Learning and generalization under ambiguity: an fMRI

study. *PLoS computational biology*, *8*(1), e1002346.

d'Acremont, M., Fornari, E., & Bossaerts, P. (2013). Activity in inferior parietal and

medial prefrontal cortex signals the accumulation of evidence in a probability

learning task. *PLoS computational biology*, *9*(1), e1002895.

Fecteau, S., Knoch, D., Fregni, F., Sultani, N., Boggio, P., & Pascual-Leone, A.

(2007). Diminishing risk-taking behavior by modulating activity in the prefrontal

cortex: a direct current stimulation study. *Journal of Neuroscience*, *27*(46), 12500-

12505.

Fletcher, P. C., Anderson, J. M., Shanks, D. R., Honey, R., Carpenter, T. A.,

Donovan, T., & Bullmore, E. T. (2001). Responses of human frontal cortex to

surprising events are predicted by formal associative learning theory. Nature

neuroscience, *4*(10), 1043.

Forman, S. D., Cohen, J. D., Fitzgerald, M., Eddy, W. F., Mintun, M. A., & Noll, D. C.

(1995). Improved assessment of significant activation in functional magnetic

resonance imaging (fMRI): use of a cluster-size threshold. Magnetic Resonance in

medicine, *33*(5), 636-647.

Gläscher, J. (2009) Visualization of group inference data in functional neuroimaging.

Neuroinformatics, *7*(1), 73-82.

Gläscher, J., Daw, N., Dayan, P., & O'Doherty, J. P. (2010). States versus rewards:

dissociable neural prediction error signals underlying model-based and error-driven

reinforcement learning. Neuron, *66*(4), 585-595.

Griffiths, T. L., & Tenenbaum, J. B. (2005). Structure and strength in causal

induction. *Cognitive psychology*, *51*(4), 334-384.

Grinband, J., Hirsch, J., & Ferrera, V. P. (2006). A neural representation of

categorization uncertainty in the human brain. *Neuron*, *49*(5), 757-763.

Hare, T. A., O'Doherty, J., Camerer, C. F., Schultz, W., & Rangel, A. (2008).

Dissociating the role of the orbitofrontal cortex and the striatum in the computation

of goal values and prediction errors. Journal of neuroscience, *28*(22), 5623-5630.

Harrison, L. M., Duggins, A., & Friston, K. J. (2006). Encoding uncertainty in the

hippocampus. Neural Networks, *19*(5), 535-546.

Holt, C. A., & Laury, S. K. (2002). Risk aversion and incentive effects. *American*

economic review, *92*(5), 1644-1655.

Huettel, S. A., Stowe, C. J., Gordon, E. M., Warner, B. T., & Platt, M. L. (2006). Neural

signatures of economic preferences for risk and ambiguity. Neuron, *49*(5), 765-

775.

Huettel SA, Song AW, McCarthy G. (2005) Decisions under uncertainty: probabilistic

context influences activity of prefrontal and parietal cortices. J Neurosci. 25:3304–

3311

Hsu, M., Bhatt, M., Adolphs, R., Tranel, D., & Camerer, C. F. (2005). Neural systems

responding   to   degrees   of   uncertainty   in   human   decision-

making. Science, *310*(5754), 1680-1683.

Itti, Laurent, and Pierre F. Baldi. "Bayesian surprise attracts human

attention." *Advances in neural information processing systems*. 2006.

Knoch, D., Gianotti, L. R., Pascual-Leone, A., Treyer, V., Regard, M., Hohmann, M., &

Brugger, P. (2006). Disruption of right prefrontal cortex by low-frequency repetitive

transcranial magnetic stimulation induces risk-taking behavior. *Journal of*

Neuroscience, *26*(24), 6469-6472.

Krain, A. L., Wilson, A. M., Arbuckle, R., Castellanos, F. X., & Milham, M. P. (2006).

Distinct neural mechanisms of risk and ambiguity: a meta-analysis of decision-

making. Neuroimage, *32*(1), 477-484.

Kumaran, D., Summerfield, J. J., Hassabis, D., & Maguire, E. A. (2009). Tracking the

emergence of conceptual knowledge during human decision

making. Neuron, *63*(6), 889-901.

Kushnir T, Gopnik A (2005) Young children infer causal strength from probabilities and

interventions. Psychological science 16:678-683.

Le Pelley ME, Oakeshott SM, McLaren IP (2005) Blocking and unblocking in human

causal learning. Journal of experimental psychology Animal behavior processes

31:56-70.

Levy, I., Snell, J., Nelson, A. J., Rustichini, A., & Glimcher, P. W. (2010). Neural

representation of subjective value under risk and ambiguity. Journal of

neurophysiology, *103*(2), 1036-1047.

Liljeholm, M. (2015). How multiple causes combine: independence constraints on

causal inference. Frontiers in psychology, 6.

Liljeholm, M. & Cheng, P.W. (2007) When is a cause the "same"? Coherent

generalization across contexts. Psychological science 18:1014-1021.

McClure, S. M., Berns, G. S., & Montague, P. R. (2003). Temporal prediction errors in

a passive learning task activate human striatum. Neuron, *38*(2), 339-346.

Meder B, Hagmayer Y, Waldmann M (2006) Understanding the Causal Logic of

Confounds. In: Twenty-Eighth Annual Conference of the Cognitive Science

Society, pp 579-584: Erlbaum.

Morris, R. W., Vercammen, A., Lenroot, R., Moore, L., Langton, J. M., Short, B., ... &

Weickert, T. W. (2012). Disambiguating ventral striatum fMRI-related bold signal

during reward prediction in schizophrenia. Molecular psychiatry, *17*(3), 280.

Cheng, P. W., & Novick, L. R. (1990). A probabilistic contrast model of causal

induction. *Journal of personality and social psychology*, *58*(4), 545.

O'Doherty, J., Dayan, P., Schultz, J., Deichmann, R., Friston, K., & Dolan, R. J.

(2004). Dissociable roles of ventral and dorsal striatum in instrumental

conditioning.   Science,   *304*(5669),   452–454.

http://doi.org/10.1126/science.1094285

Payzan-LeNestour, E., Dunne, S., Bossaerts, P., & O'Doherty, J. P. (2013). The

neural representation of unexpected uncertainty during value-based decision

making. *Neuron*, *79*(1), 191-201.

Preuschoff, K., Quartz, S. R., & Bossaerts, P. (2008). Human insula activation reflects

risk prediction errors as well as risk. *Journal of Neuroscience*, *28*(11), 2745-2752.

Pushkarskaya, H., Smithson, M., Joseph, J. E., Corbly, C., & Levy, I. (2015). Neural

correlates of decision-making under ambiguity and conflict. *Frontiers in behavioral*

neuroscience, *9*, 325.

Schulz L, Bonawitz E (2007) Serious fun: preschoolers engage in more exploratory

play when evidence is confounded. Developmental psychology 43:1045-1050.

Schulz L, Gopnik A, Glymour C (2007) Preschool children learn about causal structure

from conditional interventions. Dev Sci 10:322-332.

Smith, K., Dickhaut, J., McCabe, K., & Pardo, J. V. (2002). Neuronal substrates for

choice under ambiguity, risk, gains, and losses. *Management science*, *48*(6), 711-

718.

Spellman B (1996) Acting as Intuitive Scientists: Contingency Judgments Are Made

While Controlling for Alternative Potential Causes. Psychological science 7:337-

342.

Turner, D. C., Aitken, M. R., Shanks, D. R., Sahakian, B. J., Robbins, T. W.,

Schwarzbauer, C., & Fletcher, P. C. (2004). The role of the lateral frontal cortex in

causal associative learning: exploring preventative and super-learning. Cerebral

Cortex, *14*(8), 872-880.

Volz KG, Schubotz RI, von Cramon DY. (2003) Predicting events of varying

probability: uncertainty investigated by fMRI. Neuroimage. ;19:271–280.

Waldmann MR (2001) Predictive versus diagnostic causal learning: evidence from an

overshadowing paradigm. Psychonomic bulletin & review 8:600-608.

Wasserman EA, Berglan LR (1998) Backward blocking and recovery from

overshadowing in human causal judgement: the role of within-compound

associations. The Quarterly journal of experimental psychology B, Comparative

and physiological psychology 51:121-138.

Ye, H., Chen, S., Huang, D., Wang, S., Jia, Y., & Luo, J. (2015). Transcranial direct

current stimulation over prefrontal cortex diminishes degree of risk

aversion. *Neuroscience letters*, *598*, 18-22.

673   Figure Captions

Figure 1. Task illustration. A. Prediction and outcome screens from a feedback-

based learning trial. Screens were separated by a 2 sec. pause during which the word

"Wait" (not shown in figure) was displayed. B. Causal query regarding the individual

influence of a particular medicine on headache.

Figure 2. Mean proportions of "Is cause", "Is not a cause" and ambivalent "May or may

not be cause" judgments about the individual influence of each medicine on

headache. Target causes (C, D & S) occurred only in compound with alternative

causes during feedback-based learning, while non-target causes occurred only

individually (s'), or individually as well as in compound (e).

Figure 3. Disjunction results favoring the causal model of uncertainty during the

prediction period of feedback-based learning trials, showing effects in the anterior

insula (top) and DMPFC (bottom). Plotted values show effect sizes (y-axis) extracted

from spheres centered on peak coordinates at low (L), medium (M) and high (H) levels

of causal (blue) and error-driven (red) models (middle column), and for confounded

(C), deterministic (D) and stochastic (S) compound causes (right column). Error

bars=SEM.

Figure 4. Disjunction results favoring the error-driven model of uncertainty during the

prediction period of feedback-based learning trials, showing effects in the DLPFC

(top), lingual gyrus (middle) and vmPFC (bottom). Plotted values show effect sizes (y-

axis) extracted from spheres centered on peak coordinates at low (L), medium (M)

and high (H) levels of causal (blue) and error-driven (red) models (middle column),

and for confounded (C), deterministic (D) and stochastic (S) compound causes (right

column). Error bars=SEM.

Figure 5. Disjunction results favoring the causal model of surprise during the outcome

period of feedback-based learning trials, showing effects throughout the DMPFC (top)

and bilateral hippocampus (bottom). Plotted values show effect sizes (y-axis)

extracted from spheres centered on peak coordinates at low (L), medium (M) and high

(H) levels of causal (blue) and error-driven (red) models (middle column), and for

confounded (C), deterministic (D) and stochastic (S) compound causes (right column).

Error bars=SEM.

705   Source data legends

Figure 2—source data 1

This spreadsheet contains the source data for the mean causal judgments plotted in

Figure 2. The spreadsheet can be opened with Microsoft Excel or with open-source

alternatives such as OpenOffice.

Figure 3—source data 1

This spreadsheet contains the source data for the mean effect sizes plotted in the

middle and right columns of Figure 3. The spreadsheet can be opened with Microsoft

Excel or with open-source alternatives such as OpenOffice.

Figure 4—source data 1

This spreadsheet contains the source data for the mean effect sizes plotted in the

middle and right columns of Figure 4. The spreadsheet can be opened with Microsoft

Excel or with open-source alternatives such as OpenOffice.

Figure 5—source data 1

This spreadsheet contains the source data for the mean effect sizes plotted in the right

column of Figure 5. The spreadsheet can be opened with Microsoft Excel or with

open-source alternatives such as OpenOffice.

Supplemental Table 1: Results of Bayesian Model Selection. Peak log-odds MNI coordinates
and cluster sizes (>= 40) for group-level exceedance probability maps, thresholded at a
probability (γ) > 0.95, assessing which model provided the best account of uncertainty during
feedback-based learning. R/L = Right/Left, ' ' = continuous with previous cluster.

Winning model	Area	x, y, z	Cluster size at γ > 0.95
Uncertainty during Prediction			
Causal	Anterior Cingulate Cortex	4, 48, 24	873
	Cerebellum	-36, -52, -30	677
	R. Anterior Insula	34, 18, -10	540
	L. Anterior Insula	-26, 16,8	534
	R. Calcarine	14, -86, 14	359
	R. Fusiform Gyrus	32, -40, -22	253
	L. Precentral Gyrus	-54, 10, 34	139
	Vermis	0, -62, -40	107
	Superior Frontal Gyrus	-12, 14, 58	55
	L. Dorsolateral Prefrontal Cortex	-36, 26, 22	51
Error-driven	Precuneus	-4 -44, 10	1097
	R. Lingual Gyrus	22, -84, -10	' '
	Middle Occipital	-18, -92, 18	165
	Ventral Striatum	-8, 2, -2	90
	Ventromedial Prefrontal Cortex	-2, 22, -8	86
	Posterior Cingulate	-2, -28, 22	71
	L. Angular Gyrus	-46, -72, 28	69
	Ventral Tegmental Area	-8, -20, -10	59
	L. Thalamus	-16, -10, 4	56
	R. Dorsolateral Prefrontal Cortex	44, 44, 22	43

734 Supplemental Table 2: Results of Bayesian Model Selection. Peak log-odds MNI coordinates
735 and cluster sizes (>= 40) for group-level exceedance probability maps, thresholded at a
736 probability (γ) > 0.95, assessing which model provided the best account of uncertainty during
737 causal judgments. R/L = Right/Left, ' ' = continuous with previous cluster.

Winning model	Area	x, y, z	Cluster size at γ > 0.95
Uncertainty during Judgments			
Causal	Anterior Cingulate Cortex	4, 48, 24	1166
	Cerebellum	-24, -80, -36	219
	L. Anterior Insula	-36, 18, -8	164
	L. Putamen	-20, 0, -10	143
	L. Middle Temporal Gyrus	-52, -40, 0	76
	Supplementary Motor Cortex	-12, 14, 59	81
	R. Orbitofrontal Cortex	46, 44, -8	66
	Vermis	2, -62, -40	44
	Ventral Tegmental Area	0, -20, -20	53
	Premotor Cortex	32, 10, 52	51
Error-driven	Rostrolateral Prefrontal Cortex	24, 56, 8	163
	Precuneus	-6, -56, 20	118
	L. Orbitofrontal Cortex	-46, 44, -10	67
	R. Angular Gyrus	46, -66, 38	65
	Cerebellum	-20, -84, -26	57

740 Supplemental Table 3: Results of Bayesian Model Selection. Peak MNI coordinates and cluster
741 sizes (>= 40) for group-level exceedance probability maps, thresholded at a probability (γ) > 0.95,
742 assessing causal and error-driven accounts of predictive strength during feedback-based
743 learning. R/L = Right/Left

Winning model	Area	x, y, z	Cluster size at γ > 0.95
Prediction Strength			
Causal	L. Precentral Gyrus	-26, -10, 58	1068
	L. Postcentral Gyrus	-36, -26, 46	842
	R. Superior Occipital Cortex	16, -76, 28	168
	L. Superior Occipital Cortex	-14, -80, 22	167
	Cerebellum	24, -52, -24	161
	R. Lingual Gyrus	12, -50, 6	144
	L. Middle Temporal Gyrus	-44, -52, 6	369
	Dorsomedial Prefrontal Cortex	0, 34, 36	173
	L. Putamen	-20, 2, -12	224
	R. Precentral Gyrus	32, -10, 62	93
	R. Superior Temporal Gyrus	62, -42, 20	56
	R. Postcentral Gyrus	38, -24, 38	56
Error-driven	R. Lingual Gyrus	6, -80, -12	509
	L. Middle Occipital Cortex	-18, -92, 18	159
	R. Supramarginal Gyrus	56, -36, 36	109
	R. Middle Temporal Gyrus	56, -62, 4	74
Judgment Strength			
Error-driven	R. Middle Frontal Gyrus	26, 28, 30	99

745 Supplemental Table 4: Results of Bayesian Model Selection. Peak MNI coordinates and cluster
746 sizes (>= 40) for group-level exceedance probability maps, thresholded at a probability (γ) > 0.95,
747 assessing causal and error-driven accounts of surprise during feedback-based learning. R/L =
748 Right/Left

Winning model	Area	x, y, z	Cluster size at $\gamma > 0.95$
Surprise			
Causal	R. Anterior Insula	34, 18, -12	821
	L. Anterior Insula	-28, 10, -20	678
	R. Middle Temporal Gyrus	64, -24, -4	510
	Anterior Cingulate Cortex	4, 48, 24	490
	R. Hippocampus	26, -36, 8	348
	L. Precentral Gyrus	-24, -10, 68	267
	L. Postcentral Gyrus	-50, -22, 38	144
	L. Hippocampus	-22, 34, 2	96
	Ventral Tegmental Area	-2, -20, -22	79
	L. Cerebellum	-20, -78, -36	70
	L. Middle Occipital Gyrus	-32, -82, 22	46
Error-driven	R. Inferior Frontal Gyrus	34, 8, 30	89
	R. Lingual Gyrus	22, 86, -12	53
	L. Caudate	-8, 16, 4	49