CRUMBLER: A tool for the Prediction of Ancestry in Cattle

Tamar E. Crum[1], Robert D. Schnabel[1,2], Jared E. Decker[1,2], Luciana CA Regitano[3], and

Jeremy F. Taylor[1]

[1]Division of Animal Sciences, University of Missouri, Columbia, MO, USA 65211

[2]Informatics Institute, University of Missouri, Columbia, MO, USA 65211

[3]Embrapa Pecuária Sudeste, São Carlos, SP 13560-970, Brazil

Corresponding author: Jeremy F. Taylor. E-mail: taylorjerr@missouri.edu.

Short Title: Predicting the Ancestry of Cattle

E-mail addresses:

TEC: tamar.crum@mail.missouri.edu

RDS: schnabelr@missouri.edu

JED: deckerje@missouri.edu

LCAR: luciana.regitano@embrapa.br

JFT: taylorjerr@missouri.edu

Abstract

Background

In many beef and some dairy production systems, crossbreeding is used to take advantage of breed complementarity and heterosis. Admixed animals are frequently identified by their coat color and body conformation phenotypes, however, without pedigree information it is not possible to identify the expected breed composition of an admixed animal and in the presence of selection, the actual composition may differ from expectation. As the roles of DNA and genotype data become more pervasive in animal agriculture, a systematic method for estimating the breed composition (the proportions of an animal's genome originating from ancestral pure breeds) has utility for a variety of downstream analyses including the estimation of genomic breeding values for crossbred animals, the estimation of quantitative trait locus effects, and heterosis and heterosis retention in advanced generation composite animals. Currently, there is no automated or semi-automated ancestry estimation platform for cattle and the objective of this study was to evaluate the utility of extant public software for ancestry estimation and determine the effects of reference population size and composition and number of utilized single nucleotide polymorphism loci on ancestry estimation. We also sought to develop an analysis pipeline that would simplify this process for members of the livestock genomics research community.

Results

We developed and tested a tool, "CRUMBLER", to estimate the global ancestry of cattle using ADMIXTURE and SNPweights based on a defined reference panel. CRUMBLER, was developed and evaluated in cattle, but is a species agnostic pipeline

that facilitates the streamlined estimation of breed composition for individuals with

potentially complex ancestries using publicly available global ancestry software and a

specified reference population SNP dataset. We developed the reference panel from a

large cattle genotype data set and breed association pedigree information using

iterative analyses to identify purebred individuals that were representative of each

breed. We also evaluated the numbers of markers necessary for breed composition

estimation and simulated genotypes for advanced generation composite animals to

evaluate the precision of the developed tool.

Conclusion

The developed CRUMBLER pipeline extracts a specified subset of genotypes that is

common to all current commercially available genotyping platforms, processes these

into the file formats required for the analysis software, and predicts admixture

proportions using the specified reference population allele frequencies.

Background

Estimation of the breed composition of individuals with complex ancestries has utility for

estimating breed direct and heterosis effects as well as for the estimation of the additive

genetic merit of these individuals. It also has value for identifying the breed composition

of training populations used for genomic selection and hence the identification of target

breeds in which the developed prediction equations may have some relevance. Visual

classification of cattle based on breed characteristics suffers from similar problems as

the self-identification of ethnicity in humans [1], as most breed characteristics are

determined by alleles at relatively few loci. For example, recent extensive crossing with

Angus cattle in the U.S. produces a black hided animal which masks all other solid coat

colours found in other breeds and requires only a single dominant allele at the *MC1R*

locus. As a consequence, black-hided cattle have a "cryptic" population structure [1,2]

and the visual classification of black-hided animals for branded beef programs can

result in the marketing of animals with vastly different Angus genome content.

In the U.S. and many other countries, the breed of an animal is associated with its being

registered with a breed association which requires that both parents of the animal be

identified and also registered with the association. For the previous 50 years,

parentage has been validated by each breed association using blood or, more recently,

DNA typing. Many breed associations have closed herdbooks which means, in theory,

that the pedigrees of all animals can be traced back to the animals that founded the

breed's herdbook. Other breed associations have open herdbooks, which means that

crossbred animals can be registered with the breed if they have been graded up by

crossbreeding to have the expectation of a certain percentage of their genome (e.g.,

15/16ths) originating from the respective breed based upon pedigree records and

parentage validation. Pedigree errors that occurred prior to, or that were not identified

following the implementation of blood typing and DNA testing, lead to admixed animals

being incorrectly classified as fullblood and incorrectly identified admixture proportions

in purebred animals. The effects of recombination, random assortment of chromosomes

into gametes and selection can also lead to considerable variation in the extent of

identity by descent between relatives separated by more than a single meiosis and can

also lead to admixture proportions that differ substantially from expectation based on

pedigree.

Crossbreeding is extensively used in commercial beef production and in other livestock

species production systems to capitalize on the effects of breed complementarity and

heterosis resulting in herds of females that may have very complex ancestries that

frequently use fullblood or purebred bulls sourced from registered breeders. Changes in

the decision as to which breed of bull to use can result in large changes in admixture

proportions of replacement cows and marketed steers between years and large

differences can occur between herds for the same reason. When commercially sourced

animals are used to generate resource populations to study the genomics of

economically important traits such as feed conversion efficiency [3,4] or bovine

respiratory disease [5], the presence of extensive admixture in the phenotyped and

genotyped animals may impact the GWAA [3,4] and leads to the training of genomic

prediction models in populations for which the breed composition is not understood. As

a consequence, the utility of these models in other industry populations, including the

registered breeds in which the majority of genetic improvement is generated is also not

understood.

As the number of genotyped beef animals has increased, the need to classify the breed

composition of these animals has necessitated the development of a precise and

accurate method for estimating breed composition in cattle based on single nucleotide

polymorphism (SNP) data. Iterative ancestry estimation analyses performed using

different software input parameters may identify those that cause output sensitivity and

can lead to an interpretation of population structure that is close to the truth [6]. We

developed the CRUMBLER analysis pipeline to streamline the genomic estimation of

breed composition of crossbred cattle using high-density SNP genotype data, publicly

available software, and a reference panel containing genotypes for members of cattle

breeds that are numerically important in North America. The CRUMBLER pipeline is

species agnostic and could be adapted for breed composition estimation in other

species. CRUMBLER and the reference panel data are available on GitHub

(https://github.com/tamarcrum/CRUMBLER). This pipeline tool is released under the

GNU General Public License.

Materials and Methods

Genotype data

From among the numerically most important cattle breeds in North America, in terms of

their annual numbers of animal registrations, a list was compiled to define the target

breeds for reference panel development. Composite breeds, such as Brangus and

Braford, were not included in this list due to lack of available genotype data, but the

progenitor Angus, Hereford and Brahman breeds were included. Breeds such as

N'Dama, representing African taurine, and Nelore and Brahman, representing *Bos*

taurus indicus cattle, were included. We also initially included breeds that were likely to

be involved in early crossbreeding of cattle in the U.S. (Texas Longhorn).

From the 170,544 cattle with high-density SNP genotypes stored within the University of

Missouri Animal Genomics genotype database, we extracted genotypes for 48,776

animals identified as being registered with one of the numerically important U.S. Breed

Associations or belonging to other world breeds. Pedigree data were also obtained for

these animals from each of the Breed Associations, where available (Table 1). These

individuals had been genotyped using at least one of 9 different genotyping platforms

currently used internationally to genotype cattle including the GeneSeek (Lincoln, NE)

GGP-90KT, GGP-F250, GGP-HDV3, GGP-LDV1, GGP-LDV3, and GGP-LDV4 assays,

the Illumina (San Diego, CA) BovineHD and BovineSNP50 assays, and the Zoetis

(Kalamazoo, MI) i50K assay. The numbers of variants queried by each assay and the

number of individuals genotyped using each platform are shown in Table 2.

Marker set determination

To maximize the utility of the developed breed assignment tool, we identified the

intersection set of SNP markers located on the bovine assays for which we had

available genotype data (Table 2). However, during the process of identifying the

animals that would define the breed reference panel, only 16 individuals had been

genotyped using the GGP-LDV4 (n=2) and GGP-LDV3 (n=14) assays and no animals

had been genotyped using the GGP-LDV1 assay. To retain as many SNP markers as

possible for subsequent analysis, we identified the intersection of markers present on

the GGP-90KT, GGP-F250, GGP-HDV3, GGP-LDV3, GGP-LDV4, BovineHD,

BovineSNP50 and i50K assays. This intersection set included 6,799 SNP markers

(BC7K). The intersection of the markers representing 5 assays (GGP-90KT, GGP-

F250, GGP-HDV3, BovineHD, and BovineSNP50) was 13,291 markers (BC13K). By

removing only the 16 individuals from the breed reference panel that had been

genotyped on the GGP-LDV3 and GGP-LDV4 assays, we were able to compare

ancestry predictions using two marker set densities (BC13K and BC7K).

Pipeline

The developed CRUMBLER pipeline integrates the tools and the computational

efficiency of publicly available software, PLINK [7,8], EIGENSOFT [9,10] and

SNPweights [11] to generate ancestry estimates (Fig. 1). The pipeline integrates the

often cumbersome processes of data reformatting and sequentially processing the data

using analytical tools to generate ancestry proportions for targeted individuals based on

a curated breed reference panel.

PLINK

PLINK PED formatted genotypes are required as input to the pipeline. PLINK

(v1.90b3.31) was used for data filtering and formatting. Genotypes can arise from any

of the common bovine genotyping platforms (Table 2), provided that a PLINK

compatible MAP file is provided for each assay and data produced using only a single

genotyping assay is included in each PED file. The pipeline utilizes the PLINK marker

filtering tool (--extract) to extract the user-specified marker subset for ancestry analysis.

For analyses of animals genotyped on different genotyping platforms, the marker list

representing the intersection of the platforms can be provided to extract the markers

that are common to all assays. The pipeline allows multiple input genotype files and

uses the PLINK merge genotype files tool (--merge) to combine genotypes into a single

file for downstream analysis.

EIGENSOFT

The EIGENSOFT convertf package is used to convert all genotypes from PLINK PED

format into EIGENSTRAT format which is required by the SNPweights software. To

process the reference panel data, principal component analysis using EIGENSOFT

smartpca is used to generate the eigenvalues and eigenvectors that are required to

calculate SNP weights using SNPweights. However, the smartpca package included in

EIGENSOFT versions beyond 5.0.2 are not compatible with SNPweights. SNPweights

requires an input variable, "*trace*", to be located in the log file output from the smartpca

analysis. For versions of EIGENSOFT beyond 5.0.2, the source code can be edited to

ensure that the log file output is compatible with the SNPweights software (See

Supplementary Information).

SNPweights

SNPweights implements an ancestry inference model based on genome-wide SNP

weights computed using genotype data for an external panel of reference individuals.

To obtain SNP weights, the matrix (g_{ij}) of reference panel genotypes for SNP **i=1, …, M**

and individual **j=1, …, N** is normalized by subtracting the mean $\mu_i = N^{-1}\sum_j g_{ij}$ and

dividing by the standard deviation $[p_i(1-p_i)]^{0.5}$ for each SNP, where $p_i = \mu_i/2$, to improve

the results of the subsequent PCA analysis from which a kinship matrix is generated

[15]. A principal component decomposition is then used to generate the eigenvalues

and corresponding eigenvectors of the kinship matrix [11]. The SNP weights file only

needs to be recalculated if the reference panel is changed. EIGENSTRAT formatted

target animal genotypes are input into SNPweights, along with the precomputed

reference panel SNP weights. The SNP weights are then applied to the target

individuals to estimate their ancestry proportions [11].

Reference panel development

The definition of a set of reference individuals that define the genotype frequencies at

each SNP variant for each reference breed is technically demanding, but vitally

important to the process of defining ancestry. This process assumes that selection has

not operated to change gene frequencies between target and reference population

animals, and that each population is sufficiently large that drift has not impacted allele

frequencies. It also assumes that migration between different countries does not

influence population allele frequencies when registered animals are imported or

exported. FastSTRUCTURE [12] analysis and iterations of animal filtering using

SNPweights was performed using the genotypes of candidate reference panel

individuals to remove individuals with significant evidence of admixture from the

reference breed panel. An overview of the processes and iterations of filtering

conducted in the development of this reference panel set is shown in Fig. S1 and Table

1.

FastSTRUCTURE analysis to identify candidate reference panel individuals

Genotype data for 48,776 individuals produced by one of 8 different genotyping assays

were available for fastSTRUCTURE analysis (Table 1) [12]. We initially performed

focused fastSTRUCTURE analyses using small numbers of reference breeds including

Angus and Simmental, Angus and Gelbvieh, Angus and Limousin, Angus and Red

Angus, Red Angus, Hereford, Shorthorn and Salers, Red Angus, Hereford and

Shorthorn, and N'Dama, Nelore and Brahman (Figs. S2-S8). Individuals possessing an

ancestry assignment of at least 97% to their designated breed were retained for

subsequent analysis (see Supplementary Methods and Table 1). Following filtering

based on fastSTRUCTURE breed assignment, 17,852 individuals representing 19 of the

original breeds remained for further analysis (Supplementary Methods and Figs. S2-

S8). All of the Salers animals were removed in this filtering analysis which is consistent

with previous work that found that Salers and Limousin were very similar [4]. Variation in

reference population sample sizes has been shown to substantially influence the

estimation of the number of ancestral populations (K) in ancestry analyses [6,13,14]. To

minimize this effect and produce similar sample sizes for each of the reference breeds,

we randomly sampled 200 individuals from each reference breed for which at least 200

individuals remained after filtering on an ancestry assignment of at least 97%, otherwise

all remaining individuals were included for the breed (Table 1). Following

fastSTRUCTURE analysis using K=19 after removal of Salers and using the BC7K

marker set, Texas Longhorn was also removed from the reference panel breed list due

to the inability to distinguish Texas Longhorn as a distinct population (Figure 2).

Further, due to the known common ancestry [15] and similarity between Nelore and

Brahman (Figure 2), the breeds were combined to represent *Bos taurus indicus*.

SNPweights analyses to refine and validate reference panel members

Random sampling of reference breed individuals was performed to create sample sets

containing ≤n individuals per breed, for n = 50, 100, 150 and 200 individuals (Figs. 3 a-b

and Figs. S9-S10). Sampling was performed such that if a reference breed had ≥n

candidates then n individuals were randomly sampled, otherwise, all available

individuals were sampled. An analysis was performed using the BC7K marker set,

SNPweights was used to assign reference breed ancestries to the same sample of

individuals that was used to produce the SNP weights for each of the four samples of

individuals (Figs. 3 a-b and Figs. S9-S10). In the self-assignment analyses conducted

using the reference breed sample sets of ≤100 individuals per breed and ≤50 individuals

per breed, 7 individuals were removed due to their estimated breed ancestry being

≤60% to their registry breed (Holstein n=3, Jersey n=1, Japanese Black n=3) (Figs. 3 a-

b).

Breeds with open herdbooks

For the Gelbvieh, Limousin, Shorthorn, Simmental, and Braunvieh breeds that have

open U.S. herdbook registries, fullblood or 100% ancestry individuals were identified

based on pedigree data obtained from the respective breed associations (Table 1). The

term "fullblood" is used to identify cattle for which every ancestor is registered in the

herdbook and can be traced back to the breed founders. The term "purebred" refers to

animals that have been graded up via crossbreeding to purebred status. Charolais also

has an open herdbook registry in the U.S., however, access to Full French imported

Charolais breed members was limited. As a result, all Charolais individuals identified as

purebred in the association registry were retained for downstream analysis, however,

these individuals could contain up to 1/32 introgression from another breed. A random

sample of 200 individuals was taken for each breed with more than 200 identified

fullblood individuals, otherwise all animals were sampled. Individuals previously

included in the candidate reference panel following preliminary fastSTRUCTURE

filtering for the open herd book breeds were removed and replaced with the fullblood

individuals.

Additional reference panel filtering using SNPweights

After filtering animals identified to not be fullblood based on their pedigree information,

we randomly sampled ≤50 individuals per reference breed and utilized SNPweights to

estimate weights for each sample and also to estimate breed ancestries for members of

the same sample that was used to generate the SNP weights. Based on these

analyses, we created 5 overlapping reference breed sets, each containing individuals

with ≥90%, ≥85%, ≥80%, ≥75%, or ≥70% ancestry assignment to their registry breeds

(Table 3).

Simulated Genotypes

Using the phased BC7K genotypes for the final reference population of 803 individuals

(3 Nelore genotyped with the BovineHD assay were removed because they were

determined to cause the phasing software problems), we simulated genotypes for 803

individuals each generation (N = 1, 3, 5 and 10) by randomly sampling two individuals

as parents from generation N-1 and using a Poisson distribution to sample at random a

single recombinant chromosome from each parent. The number of recombination

events for each sampled chromosome was sampled from a Poisson distribution with

mean equal to chromosome length in Mb/100 (i.e. 1.58 Morgans for chromosome 1).

Simulated genotypes were produced for individuals 1 generation removed from the

fullblood/purebred reference population animals (i.e., 50% breed A and 50% breed B),

3, 5, and 10, generations, respectively, to evaluate the ability of CRUMBLER to detect

large through to small admixture proportions in animals with increasing numbers of

breeds represented in their ancestry. Breed composition estimates for these animals

were obtained by tracing the breed of origin of every allele present in each generation N

animal. For each marker, we attributed the genomic fragment from the centre points of

the intervals on each side of each marker to the breed of origin of the two alleles at

each marker and summed these across all loci. Finally, we normalized these sums by

dividing by the autosomal genome size using UMD3.1 coordinates.

Results and Discussion

The concept of breed and breed membership is man-made and does not inherently

exist in nature. Moreover, the formation of breeds of cattle is very recent, as cattle

domestication began about 10,000 years ago but the formation of herdbooks has

occurred only during the last 200-250 years [16]. Nevertheless, the effects of drift and

human selection over the last 200 years have caused sufficient divergence among

breeds that breed differences are identifiable at the molecular level. Such signals are

essential for breed ancestry analyses to be effective in modern admixed animals.

Previous work on assigning breed composition in admixed cattle utilized 50K genotype

data and a reference panel of 16 breeds, with the basis for reference panel inclusion

being breed association registration [17]. However, the continual evolution of

genotyping assays has led to content changes resulting in only a relatively small

proportion of markers in common among assays. Consequently, there is a need to

evaluate whether these markers are sufficient for breed content estimation, leading to

their conservation in the design of future assays. Furthermore the development of an

analytical pipeline based on these markers would simplify analysis for end-users and

the use of a single reference panel would allow the direct comparison of results

between applications.

Reference panel development

Previously developed cattle reference panels have relied on pedigree accuracy and

breed association registration for their definition [17]. Conversely, we used an iterative

approach for reference population curation that was able to validate the accuracy of the

pedigree information used to identify candidates. FastSTRUCTURE analyses

performed using the candidate individuals for each of the initial 19 reference breeds

suggested population subdivision in both the Hereford and Simmental (Figure 2).

Pedigree analysis for the Herefords within each subpopulation indicated that the

subpopulations comprised animals from the highly inbred USDA Miles City Line 1

Hereford population (L1) and other individuals representing broader U.S. Hereford

pedigrees. The Miles City L1 Hereford cattle were derived from two bulls, both sired by

Advance Domino 13 (AHA registration number 1668403) and 50 Hereford foundation

cows. Since the founding of the L1 Herefords, the migration of germplasm has been

unidirectional from L1 into the broader U.S. industry, as the L1 population has been

closed since its founding [18]. However, the L1 Herefords have profoundly influenced

the U.S. Hereford population. L1 Herefords do not segregate for recessive dwarfism,

which has been a threat to Hereford breeders since the 1950s, and this has led to L1

cattle becoming popular in the process of purging herds of the defect [19]. In 1980, the

average proportion of U.S. registered Herefords influenced by L1 genetics was 23%.

By 2008, this proportion had increased to 81% [18].

The detected subpopulation division within the Simmental breed (Figure 2) represents

the differentiation between purebred and fullblood animals. For example, progeny of a

popular Simmental sire are present in both subpopulations, however, in one

subpopulation the family members are all fullblood and in the other they are all purebred

or percentage Simmental animals. This result supports the need to identify fullblood

animals as reference panel breed representatives for breeds with open herdbooks.

Reference population sample size

By randomly sampling individuals from the candidate reference breed set and using

SNPweights to assign these individuals to reference populations, we found that

reference panel breed sample sizes of ≤50 or ≤100 individuals appeared to capture the

diversity within each breed and appropriately determined the ancestry of the tested

individuals (Fig. 3 a-b). For each breed, the percent ancestry predicted for the tested

reference samples was, on average, 3.86% higher when the SNP weights were

estimated using ≤50 individuals per breed than when ≤100 individuals per breed were

used (Table 4). This reflects the increased homogeneity of individuals within each

breed and a greater genetic distance between individuals from different breeds as

smaller samples of individuals from each breed are used to define the reference panel.

Further, due to limitations in the number of genotyped individuals for some breeds

(Table 1), as the sample size was increased globally, imbalances were created between

the reference panel breed sample sizes which impacted breed composition estimation

(Fig. S9-S10). It has previously been shown that the power to detect population

structure improves as the reference population sample sizes become more similar

[6,14].

<u>Marker density</u>

After the replacement of reference breed individuals with those identified to be fullblood

based on pedigree analysis for the open herdbook Gelbvieh, Simmental, Limousin,

Braunvieh, Shorthorn, and Charolais breeds, additional self-assignment analyses were

conducted to evaluate the effects of marker set size on ancestry prediction. Breed

reference panels were again constructed by randomly sampling ≤50 individuals per

breed and SNP weights were calculated using both the BC13K markers and BC7K

markers. The estimated SNP weights were then used to self-assign ancestry to

members of the reference panel animals representing the reference breed set. The

ancestry predictions for the reference breed individuals using either the BC7K (Fig. 4a;

Fig. S11) or BC13K (Fig. 4b; Fig. S12) marker sets indicate that use of the BC13K

marker set did not significantly impact the ancestry predictions. Consequently, the use

of the 6,799 markers common to the 8 commercially available genotyping platforms

appears to be sufficient to assign breed ancestry for the majority of animals produced in

the U.S. The CRUMBLER pipeline can accommodate samples genotyped using

alternative assays ,however, the produced breed composition estimates will be based

on the intersection of markers on the assay and the BC7K marker set.

Assignment thresholds

We next examined the effects of reference breed homogeneity on ancestry assignment

by identifying reference panel members that had been assigned to their breed of

registry using SNPweights with probabilities of ancestry of ≥90%, ≥85%, ≥80%, ≥75%,

and ≥70%, respectively (Table 3). From these individuals, reference breed panels were

obtained by randomly sampling ≤50 individuals per breed, until each individual was

represented in at least one sample set. SNP weights were then estimated using the

BC7K marker set and ancestry was assigned for these individuals using SNPweights

(Figs. 5-6 and Figs. S13-S15). Limiting the reference breed panel members to those

individuals with ≥90% ancestry assigned to their breed of registry produced a reference

panel that did not represent the extent of diversity within each of the breeds (Fig. 5). On

the other hand, using an ancestry assignment of ≥85% clearly captured greater diversity

within each breed (Fig. 6) and maximized the self-assignment of ancestry to the breed

of registration (Table 5).

Reference panel definition

To examine whether the specific individuals represented in the reference panel sample

influenced the self-assignment of ancestry to the sampled individuals, a second sample

of ≤50 distinct individuals per breed was obtained from the individuals with ≥85%

assignment to their breed of registration and analyzed with SNPweights (Fig. 7). Fig. 7

indicates that the ability to predict ancestry was not influenced by the specific individuals

sampled from the set of animals with ≥85% ancestry to their breed of registration.

Additionally, Figs. 6 and 7 suggest that the use of a reference breed panel constructed

by the random sampling of ≤50 individuals per breed from individuals with ≥85% self-

assigned ancestry to their breed of registration maintained sufficient within-breed

diversity to accurately estimate the ancestry of target individuals. However, these

figures also reveal small amounts of apparent introgression from other reference panel

breeds within each of the breeds. This does not appear to be an issue of marker

resolution since the analyses performed with the BC7K and BC13K marker sets

generated similar results (Fig. 4). We conclude that these apparent introgressions are

either due to a lack of power to discriminate among breeds using the common markers

designed onto commercial genotyping platforms, or represent the presence of common

ancestry among the breeds prior to the formation of breed herdbooks ~200 years ago.

Molecular evidence for this shared ancestry exists, for example, Hereford and Angus

cattle share the *Celtic* polled allele [20] and the segmental duplication responsible for

the white anterior, ventral and dorsal coat color pattern occurs only in Hereford and

Simmental cattle and their crosses [21]. These data clearly indicate that crossbreeding

was widespread prior to the formal conceptualization of breeds.

Reference panel validation

To evaluate the ability of the selected reference breed panel to identify breed

composition, an analysis was conducted for all 170,544 samples in the database which

required 60 processor minutes (Fig 6-7). We extracted animals with pedigree

information including fullblood and purebred animals registered with open herdbook

breed associations and 2,243 crossbred animals with varying degrees of admixture.

Considering the amount of available data, the number of pedigreed admixed animals

was very limited and the purebred animals all had similar expected admixture

proportions. Consequently, we next simulated genotypes for animals by assuming the

random mating of members of the reference breed panel for 1, 3, 5 and 10 generations

assuming non-overlapping generations to generate generations of animals with different

numbers of breeds and breed proportions represented in their genomes.

<u>Registered fullblood animals</u>

For the Gelbvieh, Limousin, Shorthorn, Simmental, and Braunvieh breeds that have

open herdbook registries, fullblood or 100% ancestry individuals were identified based

on pedigree data obtained from the respective breed associations (Table 1).

CRUMBLER estimates were obtained for these fullblood individuals and the distribution

of estimates by breed are in Fig. 8. For all breeds except Charolais, >50% of the

individuals had CRUMBLER estimated percentages of ≥80% to their respective breeds.

Average percentage estimates for fullblood Gelbvieh, Limousin, Shorthorn, Simmental,

and Braunvieh individuals were 76%, 78%, 83%, 79%, and 85%, respectively (Fig. 8b).

However, the number of genotyped imported Full French Charolais animals was limited

and so we also analyzed all purebred Charolais individuals which could contain up to

1/32nd of their genome introgressed from another breed. The average Charolais breed

assignment was 72% and the distribution of estimates was more variable than for the

fullblood animals from the other breeds (Fig. 8b).

Pedigreed crossbred animals

Based on pedigree, 2,005 individuals were identified as being primarily Hereford but

with varying degrees of Red Angus, Salers, Angus or unknown other breed influence.

The analysis results agreed with the pedigree data (Fig. 9a) To investigate the

correlations between pedigree and CRUMBLER estimated breed proportions, we

removed proportions for breeds that were less than 3% and normalized the remaining

values. CRUMBLER estimates were then correlated with the pedigree predicted

estimates of the proportion of Hereford in these individuals (Fig. 9c). CRUMBLER

tended to underestimate the Hereford proportion as the pedigree estimated Hereford

proportion tended to 100%.

The remaining 238 crossbred individuals were commercial, advanced generation

animals with an expected 50% Angus and 50% Simmental ancestry based on pedigree

data. Results of the CRUMBLER analysis support the pedigree data (Fig. 10).   The

presence of Red Angus ancestry in these animals reveals the inability of the analysis to

fully differentiate between Angus and Red Angus, which only diverged in the U.S. in

1954, and also the influence of Red Angus in the U.S. Simmental breed (Fig. S16).

Simulated genotypes

Genomes were simulated using the phased genotypes for 803 individuals from the

reference breed panel to contain varying breed numbers and admixture proportions

after 1, 3, 5, and 10 generations of random mating with nonoverlapping generations. In

generation 1, the admixed individuals were F_1 individuals with a 50:50 autosomal

genome composition unless both parents were randomly sampled from the same breed.

CRUMBLER estimates of breed composition using the simulated genotypes were

strongly correlated with the simulated compositions, especially for generations 1 and 3

(Fig 11). As the number of generations increases, the number of breeds represented in

the simulated genomes tends to increase and the proportion of the genome originating

from any one breed tends to decrease and the correlation between the simulated

proportions and CRUMBLER estimates decreased. Nevertheless, by generation 10

44% of animals had their genome proportions estimated with a correlation of at least

70%. In the U.S. commercial crossbreeding does not usually involve the use of more

than 3-4 breeds of cattle and while the number of generations of crossbreeding may

very well be 10 or perhaps more, many generations will involve the mating of animals

with similar genome ancestries and the proportions for each breed will be much greater

than present in the generation 10 animals in Fig 11. Consequently, the achieved

accuracies are likely to be closer to the generation 3 or 5 results where 99% and 68% of

animals, respectively, had their genome proportions estimated with a correlation of

greater than 80%.

Violation of model assumptions

The ancestry model assumes that neither drift or selection has acted to alter the allele

frequencies from those created by the initial admixture proportions. We examined

CRUMBLER estimates of breed composition for advanced generation members of the

Brangus (n=13,842), Beefmaster (n=4,010) and Santa Gertrudis (n=2,056) composite

breeds where selection has had the opportunity to change breed composition from

expectations at breed formation and Brangus individuals are expected to be ⅝ Angus

and ⅜ Brahman, Beefmaster individuals ¼ Hereford, ¼ Shorthorn and ½ Brahman, and

Santa Gertrudis ⅝ Shorthorn and ⅜ Brahman, respectively. However, CRUMBLER

assigned relatively large proportions of Holstein ancestry to these advanced generation

composite animals (Fig. 12). These American breeds do not contain any Holstein

introgression and they do not contain ancestry from a "Ghost Population", a population

not present in the reference set which would lead to a breed assignment to a reference

breed that it most closely resembles [6]. We speculate that this effect is caused by

selection creating a deviation from the founder breed proportions which is best

explained by an introgression from a distantly related breed, in this case, Holstein.

Stratifying these genotyped animals according to the number of generations from

foundation fullblood animals and examining the extent of estimated Holstein

introgression, which would be expected to increase with generation number, would

enable this to be tested, however, we did not have access to the necessary data.

However, this hypothesis is supported by the fact that the Santa Gertrudis had the least

Holstein introgression and the breed has published estimates of additive genetic merit

for many fewer years than the Beefmaster or Brangus.

Admixture

We also tested the ADMIXTURE software [22] for ancestry estimation and integration

into the CRUMBLER pipeline using the same reference breed panel that was developed

for use with SNPweights. ADMIXTURE uses maximum likelihood estimation to fit the

same statistical model as STRUCTURE, however, STRUCTURE does not allow the

specification of individuals of known descent to be used as a reference panel [22].

ADMIXTURE allows a supervised analysis, in which the user can specify a reference

set of individuals, by specifying the "--supervised" flag and requires an additional file

with a ".pop" suffix to specify the genotypes of the reference population individuals [22].

Unlike SNPweights, the reference population individuals' genotypes must be provided in

a genotype file for each analysis.

We first conducted an ADMIXTURE analysis in which we self-assigned ancestry for the

animals in the reference breed set formed with ≤50 individuals per breed from the

individuals that had ≥85% assignment to the breed of registration (Fig. 13). The results

shown in Fig. 13 are similar to those in Fig. 6 for the same reference panel, albeit with

perhaps less evidence of background introgression. We next conducted an analysis

using the reference panel used in Fig. 13 merged with data for the 2,005 high

percentage crossbred Herefords animals. The results shown in Fig. 14, reveal a

significant change in the ancestry proportions estimated for the reference panel

Guernsey, Gelbvieh and Romagnola individuals between the two analyses which used

exactly the same reference panel, but differed only in the number of individuals for

which ancestry was to be estimated. This suggests that ADMIXTURE may use the

target individuals to update information provided by the reference panel individuals

specified in the ".pop" file. Consequently, the ADMIXTURE estimated ancestry

proportions appear to be context dependent and may vary based on the other

individuals included in the analysis.

Moreover, the order in which the target individuals appear in the genotype input file also

appears to affect ADMIXTURE estimates of ancestry proportions for the target

individuals. Fig. 15 shows the results of an ADMIXTURE analysis in which the target

individuals were identical to those shown in Fig. 14, but for which the order of the

reference individuals and the 2,005 Hereford crossbred individuals was reversed in the

input files. In Fig. 14, the reference individuals appear before the 2,005 Hereford

crossbred individuals in the input file, whereas in Fig. 15, the 2,005 Hereford crossbred

individuals appeared before the reference individuals in the input file. The results reveal

a significant change in ancestry proportions for Guernsey and Gelbvieh, but the

Romagnola now appear to be non-admixed. Finally, we performed an ADMIXTURE

analysis for these animals in which the order of animals in the input genotype file was

completely randomized (Fig. 16). Following analysis, the individuals were sorted to

generate Fig. 16. Again, the ancestry proportions for the Guernsey, Gelbvieh and

Romagnola individuals suggest these breeds to be admixed.

STRUCTURE and ADMIXTURE are widely used for characterising admixed populations

[6], however, we have not found any reports in the literature that indicate that the

software is sensitive to the input order of individuals. However, we suspect that the

majority of users would have no need or motivation to run the software with permuted

data input files. Nevertheless, because of these inconsistencies between results, we

chose to not use ADMIXTURE for ancestry estimation within the CRUMBLER pipeline.

Broader application using additional commercially available assays

To broaden the spectrum of data from different commercially available assays that can

be evaluated, an additional intersection of markers was obtained using 11 commercially

available bovine assays including the GGP-90KT, GGP-F250, GGP-HDV3, GGP-LDV3,

GGP-LDV4, BovineHD, BovineSNP50, i50K, Irish Cattle Breeding Federation (Cork,

Ireland) IDBv3, and GeneSeek (Lincoln, NE) BOVG50v1 assays. The intersection SNP

set included 6,363 SNPs (BC6K). A SNPweights self-assignment analysis using the

reference set of individuals with ≥85% assignment to their breed of registration was

conducted to assess the effects of the reduction in number of markers used for ancestry

assignment. The ancestry proportions assigned based on the BC6K marker set (Fig.

17) did not differ appreciably from those obtained using the BC7K marker set (Fig. 6).

This result indicates the utility of CRUMBLER and the reference panel breed set across

the spectrum of commercially available genotyping platforms.

Conclusions

The determination of a set of reference population breeds and individuals that define

allele and genotype frequencies at each variant for each of the breeds is arguably the

most important, yet technically difficult step in the process of ancestry estimation. We

employed several iterations of filtering to remove recently admixed individuals and

identify a relatively homogeneous set of individuals that nevertheless represented the

variation that might be expected among individuals within a breed. Once determined,

the reference panel genotype data need only be processed once to obtain SNP weights

removing the need to share genotype data for reference individuals in subsequent

studies [11]. The upfront development of an external reference breed panel capitalizes

on the rich ancestry information available in large available datasets, and relatedness,

variation in sample sizes and diversity among the target individuals does not affect the

inference of ancestry [11].

In cattle, the visual evaluation of breed characteristics is a poor method for evaluating

the ancestry of individuals. Breed association pedigrees can be used to estimate

expected breed compositions, however, the random assortment of chromosomes into

gametes and selection can lead to ancestry proportions that differ from those expected

based upon pedigree. Moreover, the vast majority of commercial beef cattle in the U.S.

have no or very limited pedigree information and since these animals are frequently

used for genomic research [3–5], there is a need for a tool that can routinely provide

ancestry estimates for downstream use in GWAA or other genetic studies.

We tested ADMIXTURE and SNPweights and found that results from ADMIXTURE

appear to depend on the ancestry and order of appearance of individuals within the

genotype input file. We therefore developed an analysis pipeline, CRUMBLER, based

upon PLINK, EIGENSOFT and SNPweights to automate the process of ancestry

estimation. The developed bovine pipeline utilizes the 6,799 SNPs present on all 8

commercially utilized bovine SNP genotyping assays and results using these SNPs are

consistent with results obtained when 13,291 SNPs were used. From an available

48,776 genotyped individuals, we also developed a reference panel of 806 individuals

sampled from 17 breeds to have ≤50 individuals per breed that had ≥85% assignment

to their breed of registration. This panel appears to allow the robust estimation of the

ancestry of advanced generation admixed animals, however, all breeds share some

common ancestry which predates the recent development of breed association

herdbooks [16,23].

CRUMBLER is not limited to application in cattle and with the provision of suitable

reference breed allele frequencies can be applied to other species for ancestry

estimation. CRUMBLER pipeline scripts and reference panel breed SNP weights are

available on GitHub (https://github.com/tamarcrum/CRUMBLER).

Additional files

Supplementary Information (PDF). This file contains the source code changes in

SMARTPCA within versions of EIGENSOFT beyond 5.0.2 to enable compatibility with

SNPweights.

Supplementary Methods (PDF). This file describes the preliminary fastSTRUCTURE

analyses conducted on subsamples of breeds in the development of the reference

breed panel.

Supplementary Figures (PDF). This file contains Supplementary Figures S1-S16.

Fig. S1 An overview of the processes and iterations of filtering conducted in the

development of the reference panel.

Fig. S2 Preliminary FastSTRUCTURE analysis of candidate Angus and Simmental

reference population animals.

Fig. S3 Preliminary fastSTRUCTURE analysis of candidate Angus and Gelbvieh

reference population animals.

Fig. S4 Preliminary fastSTRUCTURE analysis of candidate Angus and Limousin

reference population animals.

Fig. S5 Preliminary fastSTRUCTURE analysis of candidate Angus and Red Angus

reference population animals.

Fig. S6 Preliminary fastSTRUCTURE analysis of candidate Red Angus, Hereford,

Shorthorn and Salers reference population animals.

Fig. S7 Preliminary fastSTRUCTURE analysis of candidate Red Angus, Hereford

and Shorthorn reference population animals.

Fig. S8 Preliminary fastSTRUCTURE analysis of candidate N'Dama, Nelore and

Brahman reference population animals.

Fig. S9 SNPweights self-assignment analysis for the reference sample set

containing ≤200 individuals per breed analyzed using the BC7K marker set.

Fig. S10 SNPweights self-assignment analysis for the reference sample set

containing ≤150 individuals per breed analyzed using the BC7K marker set.

Fig. S11 SNPweights self-assignment analysis for the reference sample set

containing ≤50 individuals per breed analyzed using the BC7K marker set.

Fig. S12 SNPweights self-assignment analysis for the reference sample sets

containing ≤50 individuals per breed analyzed using the BC13K marker set.

Fig. S13 SNPweights self-assignment analysis for the reference sample set with ≥80% ancestry to breed of registry and ≤50 individuals per breed using the BC7K marker set.

Fig. S14 SNPweights self-assignment analysis for reference sample set with ≥75% ancestry to breed of registry and ≤50 individuals per breed using the BC7K marker set.

Fig. S15 SNPweights self-assignment analysis for the reference sample set with ≥70% ancestry to breed of registry and ≤50 individuals per breed using the BC7K marker set.

Fig. S16 SNPweights self-assignment analyses using a reference panel with ≤50 individuals per breed and sampling from the individuals with ≥85% assignment to their breed of registry but with (a) Red Angus or (b) Angus excluded from the reference panel.

Authors' contributions

TC conceived the study and managed the project. TC, RS, JD, and JT contributed to defining the research questions and analytical approaches and interpretation of the results. TC programmed the CRUMBLER pipeline and carried out the data analyses. TC and JT drafted the manuscript. LR provided the Nelore samples but did not have

involvement in the scientific direction. All authors read and approved the final

manuscript.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

Project Name: CRUMBLER

Project Home Page: https://github.com/tamarcrum/CRUMBLER

Programming Language: Python

Other Requirements: PLINK, EIGENSOFT, and SNPweights

License: GNU GPL

Ethics approval and consent to participate

Not applicable.

Funding

JT and RS appreciate the support of NIH-USDA Dual Purpose with Dual Benefit

Program grant number NIH 1R01HD084353. JT and RS are supported by USDA-NIFA

grants 2013-68004-20364, 2015-67015-23183, 2016-67015-24923 and 2017-67015-

26760.

Consent for Publication

Not applicable.

Acknowledgements

Not applicable.

References

References

1. Burnett MS, Strain KJ, Lesnick TG, de Andrade M, Rocca WA, Maraganore DM.

Reliability of self-reported ancestry among siblings: implications for genetic

association studies. Am J Epidemiol. 2006 Mar 1;163(5):486–92.

2. Pritchard JK, Stephens M, Donnelly P. Inference of population structure using

multilocus genotype data. Genetics. 2000 Jun;155(2):945–59.

3. Saatchi M, Beever JE, Decker JE, Faulkner DB, Freetly HC, Hansen SL, et al.

QTLs associated with dry matter intake, metabolic mid-test weight, growth and feed

efficiency have little overlap across 4 beef cattle studies. BMC Genomics. 2014

Nov 20;15:1004.

4. Seabury CM, Oldeschulte DL, Saatchi M, Beever JE, Decker JE, Halley YA, et al.

Genome-wide association study for feed efficiency and growth traits in U.S. beef

cattle. BMC Genomics. 2017 May 18;18(1):386.

5. Neibergs HL, Seabury CM, Wojtowicz AJ, Wang Z, Scraggs E, Kiser JN, et al.

Susceptibility loci revealed for bovine respiratory disease complex in pre-weaned

holstein calves. BMC Genomics. 2014 Dec 22;15:1164.

6. Lawson DJ, van Dorp L, Falush D. A tutorial on how not to over-interpret

STRUCTURE and ADMIXTURE bar plots. Nat Commun. 2018 Aug 14;9(1):3258.

7. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al.

PLINK: a tool set for whole-genome association and population-based linkage

analyses. Am J Hum Genet. 2007 Sep;81(3):559–75.

8. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-

generation PLINK: rising to the challenge of larger and richer datasets.

Gigascience. 2015 Feb 25;4:7.

9. Patterson N, Price AL, Reich D. Population structure and eigenanalysis. PLoS

Genet. 2006 Dec;2(12):e190.

10. Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D. Principal

components analysis corrects for stratification in genome-wide association studies.

Nat Genet. 2006 Aug;38(8):904–9.

11. Chen C-Y, Pollack S, Hunter DJ, Hirschhorn JN, Kraft P, Price AL. Improved

ancestry inference using weights from external reference panels. Bioinformatics.

2013 Jun 1;29(11):1399–406.

12. Raj A, Stephens M, Pritchard JK. fastSTRUCTURE: Variational Inference of

Population Structure in Large SNP Data Sets. Genetics. 2014 Jun 1;197(2):573–

89.

13. Wang J. The computer program structure for assigning individuals to populations:

easy to use but easier to misuse. Mol Ecol Resour. 2017 Sep;17(5):981–90.

14. Puechmaille SJ. The program structure does not reliably recover the correct

population structure when sampling is uneven: subsampling and new estimators

alleviate the problem. Mol Ecol Resour. 2016 May;16(3):608–27.

15. Sanders JO. History and Development of Zebu Cattle in the United States. J Anim

Sci. 1980 Jun 1;50(6):1188–200.

16. Decker JE, McKay SD, Rolf MM, Kim J, Molina Alcalá A, Sonstegard TS, et al.

Worldwide patterns of ancestry, divergence, and admixture in domesticated cattle.

PLoS Genet. 2014 Mar;10(3):e1004254.

17. Kuehn LA, Keele JW, Bennett GL, McDaneld TG, Smith TPL, Snelling WM, et al.

Predicting breed composition using breed frequencies of 50,000 markers from the

US Meat Animal Research Center 2,000 Bull Project. J Anim Sci. 2011

Jun;89(6):1742–50.

18. Leesburg VLR, MacNeil MD, Neser FWC. Influence of Miles City Line 1 on the

United States Hereford population. J Anim Sci. 2014 Jun;92(6):2387–94.

19. McCann LP. battle of bull runts. 1974; Available from: http://agris.fao.org/agris-

search/search.do?recordID=US201300539215

20. Wiedemar N, Tetens J, Jagannathan V, Menoud A, Neuenschwander S,

Bruggmann R, et al. Independent polled mutations leading to complex gene

expression differences in cattle. PLoS One. 2014 Mar 26;9(3):e93435.

21. Whitacre L. Structural variation at the KIT locus is responsible for the piebald

phenotype in Hereford and Simmental cattle. 2014; Available from:

http://search.proquest.com/openview/45eba5fa3c5757a2c4c2ab18af1a8a98/1?pq-

origsite=gscholar&cbl=18750&diss=y

22. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in

unrelated individuals. Genome Res. 2009 Sep;19(9):1655–64.

23. Decker JE, Pires JC, Conant GC, McKay SD, Heaton MP, Chen K, et al. Resolving

the evolution of extant and extinct ruminants with high-throughput phylogenomics.

Proc Natl Acad Sci U S A. 2009 Nov 3;106(44):18644–9.

Table 1. Genotype data for 48,776 registered individuals from 20 breeds were
used to establish the reference population.

Breed	No. Registered Individuals	No. FullBlood Individuals[a]	No. Individuals Assigned to Breed[b]	Sampled Individuals[c]	No. Individuals After Pedigree and SNPweights[d]
Angus	5552	5552	485	200	200
Hereford	969	969	348	200	200
Limousin	2734	321	367	200	200
Charolais	1542	1489	1542	200	200
Simmental	15858	337	1583	200	196
Japanese Black	97	97	97	97	94
Braunvieh	148	69	148	148	69
Gelbvieh	12835	51	6000	200	51
Romagnola	37	37	37	37	37
Salers	68	68	0	0	0
Texas Longhorn	45	45	45	0	0
Shorthorn	291	178	166	166	178
Red Angus	1377	1377	124	124	124
Holstein	5816	5816	5816	200	197
Jersey	119	119	119	119	118
Brown Swiss	92	92	92	92	90
Guernsey	30	30	30	30	30
N'Dama	98	98	59	59	59
Brahman	127	127	86	86	50
Nelore	941	941	708	200	50
Total	48776	17813	17852	2558	2143

[a]Number of registered animals determined by pedigree analysis to be fullblood for breed
associations with open herdbooks.
[b]Number of registered animals assigned to their identified breed with $P \geq 0.97$ by
fastSTRUCTURE in preliminary analyses and retained for subsequent analyses.
[c]A random sample of 200 individuals was obtained for breeds with >200 individuals after
fastSTRUCTURE analysis and all individuals were sampled for breeds with ≤ 200 per breed and
the data were again analyzed by fastSTRUCTURE with K=19 after removal of the Salers.
[d]Animals that were determined to not be fullblood by pedigree analysis and animals
assigned with $P \leq 0.60$ by SNPweights to their breed of registry were removed.
809

810 **Table 2. The number of variants queried by each assay and the number of**
811 **individuals from the 20 reference breeds genotyped using each assay.**
812

Assay	No. of Variants	No. of Registered Individuals
BovineSNP50	58336	20485
BovineHD	777962	2303
GGP-F250	227234	3068
GGP-90KT	76999	4407
GGP-LDV3	26504	6065
GGP-HDV3	139977	3630
GGP-LDV4	30105	8653
GGP-LDV1	8762	165
Zoetis i50K	59825	0
ICBF IDBv3	53450	0
BOVGv1	47843	0
Total		48776

814 **Table 3. Number of individuals for each reference breed assigned to their breed of**
815 **registration by minimum ancestry threshold.**
816

Breed	Breed Assignment Probability				
	≥90%	≥85%	≥80%	≥75%	≥70%
Angus	51	136	184	199	200
Hereford	58	136	184	200	200
Limousin	93	127	144	162	173
Charolais	52	92	119	132	147
Simmental	21	43	81	103	121
Japanese Black	52	73	78	83	86
Braunvieh	37	57	63	65	68
Gelbvieh	23	31	39	43	43
Romagnola	10	25	32	36	37
Shorthorn	34	98	159	170	177
Red Angus	48	88	110	120	123
Holstein	39	119	172	193	196
Jersey	52	77	91	108	116
Brown Swiss	38	64	73	82	86
Guernsey	12	22	29	30	30
N'Dama	27	45	59	59	59
Brahman	15	40	50	50	50
Nelore	32	50	50	50	50
Total	694	1323	1717	1885	1962

818 **Table 4. Ancestry proportion statistics for the self-assignment of reference panel**
819 **members from samples of ≤50 or ≤100 individuals from the candidate reference**
820 **breed individuals.**
821

Breed	Min % (≤50)	Avg % (≤50)	Max % (≤50)	Min % (≤100)	Avg % (≤100)	Max % (≤100)
Angus	86.22	90.40	95.54	78.49	87.05	94.13
Hereford	79.75	90.08	95.05	73.41	87.39	96.81
Limousin	69.52	88.53	98.16	18.36	86.40	98.81
Charolais	78.14	90.19	99.82	48.93	77.46	93.96
Simmental	81.06	90.37	97.66	61.36	73.05	88.11
Japanese Black	81.44	90.00	97.07	24.51	86.50	98.95
Braunvieh	71.59	89.46	98.61	65.46	88.36	98.70
Gelbvieh	73.03	76.27	81.63	60.92	74.59	80.33
Romagnola	75.05	87.18	96.66	74.79	85.99	95.12
Shorthorn	84.42	88.69	94.54	70.71	85.27	96.35
Red Angus	79.00	89.60	96.33	68.07	86.83	97.38
Holstein	85.82	90.30	97.51	62.95	86.97	97.81
Jersey	78.55	89.28	95.93	61.23	86.54	97.18
Brown Swiss	80.10	89.22	96.40	61.68	86.02	98.42
Guernsey	79.53	89.19	95.85	77.40	88.31	94.36
N'Dama	80.67	89.25	96.90	78.91	87.78	95.67
Bos taurus indicus	87.83	91.91	97.75	81.43	89.79	97.60

Table 5. Average predicted ancestry and variance in predicted ancestry for
 candidate reference breed individuals when filtered on minimum predicted
 ancestry.

Breed	Avg % (70%)	Var (70%)	Avg % (75%)	Var (75%)	Avg % (80%)	Var (80%)	Avg % (85%)	Var (85%)	Avg % (90%)	Var (90%)
Angus	86.50	0.21	87.95	0.19	87.33	0.22	88.86	0.13	72.34	0.97
Hereford	86.99	0.22	87.09	0.23	87.48	0.19	88.25	0.13	84.62	0.43
Limousin	86.77	0.55	89.03	0.44	87.92	0.38	88.48	0.43	80.62	1.19
Charolais	80.18	2.16	85.03	1.77	86.28	0.99	88.56	0.52	81.54	0.76
Simmental	72.73	0.89	78.45	0.58	83.81	0.36	89.65	0.15	87.82	0.50
Japanese Black	87.85	0.52	88.04	0.39	88.46	0.27	88.74	0.21	80.06	0.61
Braunvieh	87.01	0.37	87.84	0.36	87.33	0.38	88.71	0.21	80.47	1.24
Gelbvieh	86.68	0.41	87.10	0.43	87.52	0.34	88.43	0.34	83.31	1.25
Romagnola	86.16	0.33	86.37	0.32	87.16	0.32	86.22	0.29	86.38	1.16
Shorthorn	85.97	0.26	87.03	0.22	86.80	0.14	87.38	0.07	83.00	0.70
Red Angus	86.41	0.53	87.08	0.48	87.40	0.35	87.46	0.23	23.37	0.66
Holstein	86.44	0.27	87.82	0.21	87.54	0.13	88.77	0.12	79.71	0.61
Jersey	87.01	0.46	86.93	0.44	87.86	0.24	87.98	0.27	80.52	0.71
Brown Swiss	86.22	0.47	86.73	0.51	88.24	0.26	88.11	0.20	82.23	0.70
Guernsey	86.46	0.23	87.64	0.19	87.50	0.25	88.02	0.51	80.43	2.36
N'Dama	87.76	0.19	87.91	0.21	87.89	0.15	89.25	0.17	86.40	0.52
Bos taurus indicus	87.68	0.07	88.24	0.09	87.55	0.11	88.53	0.09	84.89	0.38
Average	85.58	0.48	86.84	0.41	87.30	0.30	88.32	0.24	78.69	0.87

Fig. 1 Flow diagram of the breed composition pipeline.

Fig. 2 FastSTRUCTURE results for a random sample of ≤200 individuals per breed from the pool of 17,852 potential reference individuals at K=19. Breed identification is shown below each colored block and each animal is represented as a vertical line within the block.

Fig. 3 SNPweights self-assignment analysis results for reference panel sample sets consisting of: (a) ≤100 individuals per breed, or (b) ≤50 individuals per breed. Seven individuals were filtered for ≤60% ancestry to their breed of registry (Holstein n=3, Jersey n=1, Japanese Black n=3).

a.

b.

Fig. 4 SNPweights self-assignment of ancestry for candidate reference breed individuals following evaluation of open herdbook breeds using: (a) the BC7K, or (b) the BC13K marker panels. Reference breed panels were constructed by random sampling ≤50 individuals per breed and SNP weights were estimated using the BC7K and BC13K marker sets.

Fig. 5 Reference breed panel constructed by the random sampling of ≤50 individuals per breed from individuals with ≥90% ancestry was self-assigned to reference breed ancestry using the BC7K marker set.

Fig. 6 Reference breed panel constructed by the random sampling of ≤50 individuals per breed from individuals with ≥85% ancestry was self-assigned to reference breed ancestry using the BC7K marker set.

Fig. 7 Reference breed panel constructed by the independent random sampling of a second sample of ≤50 individuals per breed from individuals with ≥85% ancestry after eliminating individuals represented in the first sample was self-assigned to reference breed ancestry using the BC7K marker set.

Fig. 8 (a) Distribution by breed of SNPweights ancestry assignment results for 2,408 registered fullblood animals from open herd book breeds. (b) Pictorial representation of CRUMBLER estimates for 2,408 registered fullblood animals from open herd book breeds.

Fig. 9 (a) SNPweights ancestry results for 2,005 crossbred Hereford individuals with *a-priori* breed composition estimates determined by pedigree. (b) Breed assignment reference breed key. (c) Hereford SNPweights estimated proportions using CRUMBLER are plotted against the pedigree estimates. Data point color indicates the breed for which SNPweights assigned the highest proportion for each individual.

Fig. 10 (a) SNPweights ancestry results for 238 crossbred individuals with *a-priori* breed composition estimates of 50% Angus and 50% Simmental based on a reference panel with ≤50 individuals per breed sampled from individuals with ≥85% assignment to their breed of registry. (b) Breed assignment for the crossbred individuals can be determined using this reference breed key.

Fig. 11 Genotypes were simulated for the indicated number of generations of random mating, with generation 1 (G1) animals being 50:50 proportion except when two parents from the same breed were mated. SNPweights results were obtained using CRUMBLER pipeline parameters correlations between these estimates and the known simulated breed compositions were produced and the proportion of individuals within each correlation class is indicated.

Fig. 12 (a) SNPweights ancestry results using CRUMBLER pipeline for 4,010 Beefmaster, 13,842 Brangus, and 2,056 Santa Gertrudis individuals. (b) Breed assignment for these advanced generation composite animals can be determined using this reference breed key.

Fig. 13 Self-assignment of ancestry for the animals in the reference breed set formed with ≤50 individuals per breed from the individuals that had ≥85% assignment to their breed of registration using ADMIXTURE.

Fig. 14 ADMIXTURE analysis conducted using the same data as shown in Figure 11 (first four rows), merged with an additional 2,005 high percentage crossbred Hereford target individuals (last row).

Fig. 15 ADMIXTURE analysis conducted using the same data as shown in Fig. 12 where the reference individuals appear before the 2,005 Hereford crossbred individuals in the input file. Here, the 2,005 Hereford crossbred individuals appear before the reference individuals in the input file. The first row represents the 2005 Hereford crossbred samples. Rows 2 to 5 show the reference panel individuals.

Fig. 16 ADMIXTURE analysis conducted using the same data as shown in Figs. 12 and 13, but with the order of the individuals in the input genotype file randomized. The first four rows represent the reference panel individuals, the fifth row shows the 2,005 Hereford crossbred animals.

Fig. 17 Reference breed panel constructed by the random sampling of ≤50 individuals per breed from individuals with ≥85% ancestry was self-assigned to reference breed ancestry using the BC6K marker set.