PARylation modulates stress granule assembly, phase separation, and neurotoxicity of ALS-related RNA-binding proteins

Yongjia Duan[1,2,*], Aiying Du[1,*], Jinge Gu[1,2,*], Gang Duan[1], Chen Wang[1,2], Le Sun[1,2], Beituo Qian[1,2], Xue Deng[1,2], Kai Zhang[1,2], Kuili Tian[1], Hong Jiang[1,2], Yaoyang Zhang[1,2], Cong Liu[1,2,#], and Yanshan Fang[1,2,3,#]

[1]Interdisciplinary Research Center on Biology and Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai, 201210, China

[2]University of Chinese Academy of Sciences, Beijing, 100049, China

[3]Lead Contact

*Y.D., A.D. and J.G. contributed equally to this work.

#**Co-corresponding authors**:

Cong Liu: liulab@sioc.ac.cn

Yanshan Fang: fangys@sioc.ac.cn

       Tel: +86-21-6858.2510

       Fax: +86-21-6858.2502

Running title: PARylation regulates SG and LLPS of ALS-related RNPs

Keywords: ALS, RNP granule, TDP-43, hnRNP A1, phase separation, PARP1, PARG

SUMMARY

Mutations in RNA-binding proteins such as hnRNP A1 and TDP-43 cause aberrant protein aggregations, which are pathological hallmarks in neurodegenerative diseases including amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD). PARylation is an important protein posttranslational modification that is involved in many cellular functions such as DNA repair, transcriptional regulation and cell death. In this study, we investigate the function of PARylation in RNP-related neurodegeneration. We reveal that PARylation levels are a major regulator of the assembly-disassembly dynamics of the disease-related RNP granules. We show that hnRNP A1 and TDP-43 can both be PARylated and bind to PARylated proteins, and we further identify the PARylation site and the PAR-binding motif of hnRNP A1. PARylation not only promotes the LLPS of hnRNP A1 *in vitro*, but also regulates its translocation to SGs and the interaction with TDP-43 *in vivo*. Finally, we establish that both genetic and pharmacological inhibition of PARP mitigates hnRNP A1 and TDP-43-mediated neurotoxicity in cell and *Drosophila* models of ALS. Together, our findings propose PARP as a potential therapeutic target for treating ALS and related diseases.

INTRODUCTION

Eukaryotic cells encode a large number of RNA-binding proteins that associate with RNAs to form ribonucleoprotein (RNP) complexes. RNPs contain conserved RNA binding domain(s) and protein-protein interaction domain(s). They are presented in both nucleus and cytoplasm, where they play a major role in RNA metabolism including RNA processing, transport and turnover (Anko and Neugebauer, 2012; Lunde et al., 2007; Muller-McNicoll and Neugebauer, 2013). RNPs can form granules by liquid-liquid phase separation (LLPS), and aberrant RNP granules can develop into irreversible fibers and aggregations that contribute to the pathogenesis of human neurodegeneration diseases including amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD) (Lin et al., 2015). Posttranslational modifications (PTMs) such as phosphorylation, ubiquitination and acetylation are known to regulate the assembly and function of RNP granules (Brady et al., 2011；Cohen et al., 2015；Dammer et al., 2012; Li et al., 2017), and methylation is recently reported to modulate phase transition of hnRNPA2 and FUS (Hofweber et al., 2018; Qamar et al., 2018; Ryan et al., 2018). However, the function of other important PTMs in regulating LLPS and RNP granules remains to be explored.

Poly(ADP-ribosyl)ation (PARylation) is a reversible PTM process by which poly(ADP-ribose) (PAR) polymerases (PARPs) add ADP-ribose (ADPr) units to the Glu, Asp, Lys, Arg or Ser residue of a protein (Martello et al., 2016; Zhang et al., 2013) whereas enzymes such as PAR glycohydrolase (PARG) remove them (Niere et al., 2012; Slade et al., 2011). The opposing effects of PARPs and PARG in regulating protein PARylation play an important role in a variety of cellular functions, including chromatin remodeling, DNA repair, transcription regulation and cell death (Ahel et al., 2008; Andrabi et al., 2006; Frizzell et al., 2009; Singh et al., 2017). PARylation is therefore involved in various disease conditions such as cancer, neurodegeneration, oxidative stress, neural injury, and regeneration (Deng, 2009; Hanai et al., 2004; Martire et al., 2015; Brochier et al., 2015). Interestingly, PAR as well as some PARPs and PARG are found in cytoplasmic stress granules (SG) and may regulate microRNA-mediated translational repression and mRNA cleavage (Gagné et al., 2008; Kotova et al., 2009; Leung et al., 2011).

Mutations in RNA-binding proteins such as heterogeneous nuclear ribonucleoprotein A1 (hnRNP A1, encoded by *HNRNPA1*) and TAR DNA binding protein 43 kDa (TDP-43, encoded by *TARDBP*) are known to promote protein aggregation and cause familial ALS (Kim et al., 2013; Neumann et al., 2006). In this study, we show that PARylation plays an important role in regulating SG dynamics, phase separation, protein-protein interaction, and the neurotoxicity of hnRNP A1 and TDP-43. Together, the findings of this study have strong implications of therapeutic values of developing PARP inhibitors for use to treat ALS and other RNP-related diseases.

RESULTS

The levels of PARylation regulate the dynamics of ALS-related RNP granules during stress

Although PARylation occurs primarily on PARP proteins, association of PAR with SGs and ALS-related RNPs was observed (Gagné et al., 2003; Gagné et al., 2008; Leung et al., 2011). To determine whether PARylation levels regulated disease-related RNP granules, we tested how Olaparib, an Food and Drug Administration (FDA)-approved PARP inhibitor targeted for cancer treatment (Fong et al, 2009), affected the response of TDP-43 and hnRNP A1 when cells were stressed (arsenite, 100 μM). SGs were examined by immunocytochemistry using the SG marker T-cell-restricted intracellular antigen 1 (TIA-1)-related protein (TIAR) (Gottschald et al., 2010). TDP-43 and hnRNP A1 were recruited to SGs when cells were stressed, and inhibition of PARP by Olaparib significantly delayed the assembly of SGs as well as the translocation of TDP-43 (Fig. 1A-1C) and hnRNP A1 (Fig. S1A-S1C) to SGs. Next, we downregulated the PAR hydrolysis enzyme PARG with small interfering RNA (siRNA) to increase PARylation levels in the cells. SGs were induced by arsenite treatment for 30 min, followed by the washout experiment to determine the kinetics of SG disassembly as stress was released (Fig. 1E). Increase of PARylation levels by si-PARG drastically delayed SG disassembly as well as the retrieval of TDP-43 (Fig. S1A-S1C) and hnRNP A1 (Fig. 1D-1F) from SGs. Thus, PARylation levels are a major factor regulating the assembly-disassembly dynamics of SGs associated with ALS-related RNPs.

PARylation and PAR-binding of TDP-43 and hnRNP A1

Since PARylation levels affected the dynamics of disease-related RNP granules containing TDP-43 and hnRNP A1, we examined whether these proteins were PARylated or associated with PAR in cells. HA-tagged TDP-43 or Flag-tagged hnRNP A1 was expressed and immunoprecipitated from Hela cells with anti-HA or anti-Flag, and then examined by Western blotting with a pan-PAR antibody (anti-PAR) that recognizes both mono-ADPr and poly-ADPr. Although there were proteins co-immunoprecipitated with TDP-43 that were immunoblotted positively with anti-PAR, we did not detect obvious PARylation of TDP-43 *per ser* (Fig. 2A). To

boost the PARP1 activity in cells, we treated the cells with H_2O_2 (500 µM, 10 min) to induce PARP1 activation (Martello et al., 2016), which markedly increased the PARylation levels of hnRNP A1 (Fig. 2B). However, no PARylation of TDP-43 was detected even under this condition. Nevertheless, this result did not exclude the possibility that PARylation of TDP-43 in cells occurred at a very low level that was below our detection sensitivity. In contrast, hnRNP A1 immunoprecipitated from Hela cells showed robust stead-state and induced levels of PARylation. In addition, there were multiple PAR$^+$ bands of proteins of higher molecular weights that were co-immunoprecipitated with hnRNP A1, especially when treated with H_2O_2 (Fig. 2B). These were likely PARylated proteins that were associated with hnRNP A1 in the cells.

Next, we assessed whether these proteins could be PARylated *in vitro*. Of note, full-length TDP-43 (TDP-43-FL) protein expressed in bacteria was extremely insoluble. We tried several different expression vectors with different purification tags and induction temperatures, but failed to produce sufficient amount of soluble TDP-43-FL protein for the *in vitro* assays (Fig. S3A-S3E). We thus took an alternative approach and divided TDP-43 into two truncations, TDP-43$^{1-274}$ and TDP-43$^{274-414}$ (Fig. S3A and S3F-S3G). The purified His-tagged TDP-43 truncations as well as full-length hnRNP A1 were then subject to the *in vitro* PARylation reaction. Single strand DNA (ssDNA) was added to activate PARP1 in the *in vitro* system (see Methods). The PARylation levels were examined by SDS-PAGE and immunoblotting with anti-PAR. Activation of PARP1 by ssDNA was evident by the dramatic increase of PARylation of PARP1 itself (Fig. 2C). With PARP1 activation, TDP-43$^{1-274}$ and hnRNP A1 showed remarkable PARylation bands and up-shifting smears due to heterogeneity in the length of ADPr polymer attached. The purified TDP-43$^{274-414}$ protein showed a basal level of PARylation (possibly occurred during expression in bacteria), which was slightly increased in the *in vitro* PARlation assay. The PARylation smear of TDP-43$^{274-414}$ was much less intense than TDP-43$^{1-274}$, while hnRNP A1 exhibited the most robust PARylation smear in the *in vitro* assay (Fig. 2C).

The human hnRNP A1 protein contains two closely-related RNA recognition motif (RRMs) in the N-terminal region and a low complexity (LC), glycine-rich domain (GRD) in the C-terminal

region that includes an RGG box RNA binding domain and a M9 nuclear targeting sequence (Fig. 2D; He and Smith, 2009). In addition, previous mass spectrometry-based studies suggested that hnRNP A1 might contain a PARylation site at K298 and a putative PAR-binding motif (PBM) between the two RRM domains at amino acid (aa) 92–113 (Gagné et al., 2008; Martello et al., 2016). To confirm, we generated constructs to express Flag-tag hnRNP A1 of a PARylation site mutant (K298A) or a PBM mutant (R92A-K105/106V, referred to as PBM^{mut} thereafter) in Hela cells. Interestingly, we found that hnRNP A1-PBM^{mut} protein showed decreased solubility (Fig. 2E and Fig. 2F, input). Nonetheless, cells transfected with Flag-tagged wild-type (WT), K298A or PBM^{mut} hnRNP A1 were treated with H_2O_2, and the cell lysates were examined by immunoprecipitation (IP) with anti-Flag and Western blotting with anti-PAR (Fig. 2E). Compared to WT hnRNP A1, PARylation of K298A was dramatically reduced, whereas the co-immunoprecipitation (co-IP) of other PARylated proteins was not affected in the K298A mutant. In contrast, PBM^{mut} showed drastically decreased association of other PARylated proteins, whereas its own PARylation was not reduced but unexpectedly increased (Fig. 2F). This might be due to increased accessibility of ADPr to the K298 site in PBM^{mut} when association of other PARylated protein was removed. Of note, hnRNP A1 showed an up-shifting PARylation smear to a similar extent in the *in vitro* PARylation assay (Fig. 2C), suggesting that the hnRNP A1 protein was capable of being massively PARylated when at optimal conditions. Thus, binding to PAR and/or PARylated proteins via the PBM might prevent hyper-PARylation of hnRNP A1 at K298.

PARylation and PAR-binding are required for hnRNP A1 translocation to SGs

We showed that the cellular PARylation levels affected the recruitment and recovery of hnRNP A1 to and from SGs (Fig. S1 and Fig. 1D-1F). However, it was unclear if PARylation directly regulated the translocation of hnRNP A1 to SGs or it was an indirect effect resulted from altered dynamics of the SGs. To address this question, we expressed the PARylation mutant K298A and PAR-binding deficient PBM^{mut} of hnRNP A1 in Hela cells and examined their cellular localization before and after stress by immunostaining. In the absence of stress, WT and K298A of hnRNP A1 were

predominantly nuclear. PBM^{mut} was mainly localized to the nucleus but also showed cytoplasmic foci that did not co-localize with the SG marker TIAR (Fig. 3A). This data was in line with the reduced solubility of PBM^{mut} in the biochemistry analysis (Fig. 2E). Of note, the PBM is not located within the RGG or M9 domain, the known nuclear localization sequences of hnRNP A1 (Fig. 2D; Siomi and Dreyfuss, 1995; Nichols et al., 2000). Thus, it is unlikely that the PBM^{mut} cytoplasmic foci are formed due to a defect in nuclear importing of hnRNP A1.

Next, we stressed the cells with arsenite (100 µM, 30 min) and it significantly induced the formation of SGs in both WT and the two mutant hnRNP A1 K298A and PBM^{mut} (Fig. 3A). Thus, the overall ability of the cells to form SGs was not affected by expressing K298A or PBM^{mut} hnRNP A1 (Fig. 3B). In response to stress, WT hnRNPA1 showed a significant cytoplasmic translocation and co-localization with TIAR-labeled SGs (Fig. 3A and 3C). In contrast, no recruitment of K298A was observed and it remained predominantly in the nucleus (Fig. 3A and 3C). For PBM^{mut}, it could respond to the stress and co-localization of PBM^{mut} with TIAR-labeled SGs was detected (Fig. 3A), but the induction was to a less extent than WT hnRNP A1 (Fig. 3C). The percentage of cells with "PBM^{mut}+/TIAR-" cytoplasmic foci showed no difference before and after stress (Fig. 3A and 3D), indicating that stress did not affect the formation or turnover of the PBM^{mut} cytoplasmic foci. Together, these data suggest that PARylation of hnRNP A1 is required for its cytoplasmic translocation, whereas binding to PAR or PARylated proteins regulates the sorting and/or delivery of hnRNP A1 to SGs.

PAR promotes phase separation of hnRNP A1

RNPs such as hnRNP A1 can phase separate *in vitro*, which may regulate the assembly of RNP granules and the pathological progression to protein aggregations *in vivo* (Lin et al., 2015). The findings that hnRNP A1 was not only PARylated but also associated with PARylated proteins *in vitro* and *in vivo* (Fig. 2) prompted us to test whether PAR could regulate the LLPS of hnRNP A1. As previously reported (Lin et al., 2015), recombinant purified hnRNP A1 formed dynamic liquid droplets (LDs) *in vitro*, which fused and increased in size with decreasing concentration of the salt

(NaCl, 25–300 mM) and increasing concentration of hnRNP A1 (10–60 μM) (Fig. 4A).

Next, to determine if PAR affected the LLPS of hnRNP A1, we chose a condition at which spontaneous LLPS of hnRNP A1 barely occurred (20 μM hnRNPA1 and 25 mM, Fig. 4A). We added purified PAR polymers with increasing concentrations (1–7 μM) into the above *in vitro* demixing system, which induced the LLPS of hnRNP A1 in a dose-dependent manner (Fig. 4B). As a control, addition of heparin (7 μM) did not promote the LLPS of hnRNP A1 (Fig. 4C). Nor did PAR (7 μM) alone phase separate *in vitro* at the same condition (Fig. 4D). Thus, it is a specific effect of PAR to promote LLPS of hnRNP A1.

hnRNP A1 and TDP-43 co-phase separate *in vitro*

A recent study showed that the LC domain of another hnRNP family protein, hnRNP A2, could co-phase separate with TDP-43 and induced co-aggregation (Ryan et al., 2018). Thus, we determined whether hnRNP A1 and TDP-43 could co-phase separate *in vitro*. Since the PBM we identified located outside the LC domain of hnRNP A1, we tested the full-length hnRNP A1 instead of only the LC domain. As shown in Figure 5A, at the condition of 50 μM of hnRNP A1 and 100 mM of NaCl, hnRNP A1 did not phase separate spontaneously. As a control, BSA (50 μM) did not phase separate on its own or trigger LLPS of hnRNP A1 (Fig. 5B-5B'). TDP-43$^{1-274}$ (50 μM) at this condition showed mild LLPS, as only few small LDs were spotted (Fig. 5C'). When it was mixed with hnRNP A1, massive and much larger LDs were formed (Fig. 5C), suggesting that interaction with each other promoted the co-LLPS of hnRNP A1 and TDP-43$^{1-274}$. Interestingly, addition of TDP-43$^{274-414}$ with hnRNP A1 did not increase the number or size of the LDs (Fig. 5D-5D'). Since 50 μM of TDP-43$^{274-414}$ on its own already formed big LDs at this condition (Fig. 5D'), it might make the promotion of co-LLPS difficult to happen. To exclude this possibility, we lowered the concentration of TDP-43$^{274-414}$ to 20 μM, at which it showed a similar level of spontaneous LLPS to that of TDP-43$^{1-274}$ (Fig. 5C' and 5E'). However, still no co-LLPS of hnRNP A1 and TDP-43$^{274-414}$ was observed (Fig. 5F-5F'). Thus, hnRNP A1 could co-phase separate with TDP-43$^{1-274}$ *in vitro*.

PARylation modulates the interaction between hnRNP A1 and TDP-43

Next, to investigate whether hnRNP A1 indeed interacted with TDP-43 *in vivo* and whether PARylation could regulate this process, we conducted the co-IP experiments. We found that the endogenous TDP-43 protein could be co-immunoprecipitated with transiently expressed hnRNP A1-Flag, which was consistent with the observations that TDP-43 bound to a few hnRNP family proteins (Buratti et al., 2005; D'Ambrogio et al., 2009). Further, we showed that the interaction between hnRNP A1 and TDP-43 depended on the presence of RNA (Fig. S4). Next, we tested the effect of the PARP inhibitor Olaparib, which significantly reduced the amount of endogenous TDP-43 that were co-immunoprecipitated with hnRNP A1; on the other hand, activation of PARP1 by H_2O_2 significantly increased the co-IP of TDP-43 with hnRNP A1 (Fig. 6A-6B).

To understand how PARylation and PAR-binding regulated the interaction of hnRNP A1 with TDP-43, we examined the binding affinity of WT, K298A and PBM^mut with TDP-43 by co-IP. Compared to WT hnRNP A1, the PARylation site mutant K298A showed a similar capability of pull-down of TDP-43, however, the association of TDP-43 with PBM^mut was dramatically reduced (Fig. 6C-6D). Of note, the protein level of PBM^mut in the input was lower due to reduced solubility (Fig. 2E). To be able to compare the co-IP efficiency with a similar input level, cells transfected with 2 times of the PBM^mut expression plasmids (2xPBM^mut) were also examined. It increased the protein levels of PBM^mut in the input, however, co-IP of TDP-43 was still barely detected (Fig. 6C-6D). Together, these data indicate that the association of PAR or PARylated proteins via the PBM of hnRNP A1 is required for the interaction between hnRNP A1 and TDP-43.

Inhibition of PARylation reduces the cytotoxicity of hnRNP A1 and TDP-43 in motor neuron-like NSC-34 cells

We extended the study of the impact of PARylation on biochemical properties of the ALS-related RNPs to functional readouts such as the cytotoxicity of these proteins. We used the mouse motor neuron (MN)-like hybrid cell line, NSC-34 cells, to overexpress hnRNP A1 or TDP-43 by lentiviral

infection. The NSC-34 cells overexpressing hnRNP A1 or TDP-43 OE cells exhibited remarkable morphological changes and significant reduction of cell viability (Fig. 7A-7B and Fig. S5), which confirmed the cytotoxicity of overexpression (OE) of hnRNP A1 or TDP-43 in the MN-like cell model.

The K298A and PBM^{mut} mutants exhibited different levels of cytotoxicity compared to WT hnRNP A1 (Fig. 7C). The K298A mutant was more toxic than WT hnRNP A1 at both low and high concentrations of lentiviral infection. Considering that PARylation at K298A was required for cytoplasmic translocation of hnRNP A1 to SGs (Fig. 3), this result suggests that the inability of promptly responding to cellular stress enhances the cytotoxicity of disease-associated RNPs. PBM^{mut} showed a similar level of cytotoxicity to WT at a low concentration of lentiviral infection. At a higher infection concentration, the cytotoxicity of WT and K298A increased drastically, but that of PBM^{mut} did not change much and hence became less toxic than WT and K298A at this condition (Fig. 7C). It was possible that the decreased binding to PARylated proteins made PBM^{mut} and the SGs it was associated with more dynamic and less solid than those of WT hnRNP A1. This would be especially important when the proteins were expressed at high levels, as it may reduce the chance of the SGs to develop into gel-like or solid protein aggregations that resemble the disease pathology.

As K298A and PBM^{mut} affected the cytotoxicity of hnRNP A1 differently, we were keen to know the overall consequence of reducing PARylation levels in cells and whether it could mitigate the cytotoxicity of hnRNP A1 or TDP-43. Thus, we treated the NSC-34 cells with siRNA of PARP1 (si-PARP1). We found that downregulation of PARP1 significantly suppressed the decrease of cell viability induced by hnRNP A1 or TDP-43 OE (Fig. 7D-7F). Furthermore, as an attempt to testing small-molecule for treating ALS, we examined the PARP inhibitor Olaparib. PARP inhibitors are clinically used drugs to kill cancer cells. Thus, we tested different doses and found that 5 μM of Olaparib did not reduce the viability of NSC-34 cells but showed a remarkable suppression of TDP-43 OE-mediated cytotoxicity (Fig. 7G). Thus, both genetic and pharmacological inhibition of PARP significantly suppressed the cytotoxicity of ALS-associated RNPs in MN-like NSC-34 cells.

Downregulation of *Parp* suppresses TDP-43-mediated neurodegeneration in a *Drosophila* model of ALS

Finally, we validated these findings in an *in vivo* model of ALS using transgenic flies expressing human TDP-43 (hTDP-43). hTDP-43 OE in the fly photoreceptor cells (GMR driver) caused age-dependent eye degeneration, which was drastically suppressed by transgenic downregulation of fly *Parp* (RNAi-*Parp*) compared to RNAi-Ctrl (Fig. 8A-8B). We examined the KD efficiency of RNAi-*Parp* by quantitative real-time PCR (qPCR) assay and the mRNA level of *Parp* was reduced to below 40% of the RNAi-ctrl group (Fig. S6A). Also, we showed that the reduction of *Parp* did not affect the protein levels or solubility of hTDP-43 (Fig. S6B-S6C), confirming that the suppression by RNAi-*Parp* was not due to a reduction of transgenically expressed hTDP-43 protein in the system. Furthermore, we performed the climbing and lifespan assays to evaluate the behavioral consequences, which might represent more closely to the disease-relevant symptoms in ALS. We induced hTDP-43 OE in adult fly neurons using an *elav*GS driver and added RU486 to the fly food (80 µg/ml) starting from day one of the adulthood). It caused an age-dependent decline of the climbing capability and a significant shortening of the lifespan, both of which could be suppressed by RNAi-*Parp* in the fly neurons (Fig. 8C-8D). Together, these data indicate that PARP can modify TDP-43-mediated neurodegeneration *in vivo*.

DISCUSSION

PARylation regulates SG dynamics and phase separation of RNPs

In this study, we reveal that decrease of the cellular PARylation levels suppresses the formation of SGs and the recruitment of ALS-related RNPs such as TDP-43 and hnRNPA1 to the SG, while increase of PARylation levels delays the disassembly of SGs and the recovery of the RNPs. Interestingly, the function of PARylation appears to be specifically enriched in regulation of RNA/DNA-binding proteins and the associated complexes. For example, PARylation is involved in the regulation of the mitotic spindle (Chang et al., 2004), Cajal bodies (Kotova et al., 2009), DNA damage repair (Ahel et al., 2009), and microRNA-mediated translational repression (Leung et al., 2011). Because of their essential role in RNA processing and protein translation (Dreyfuss et al., 1993; Lee et al., 2012), proper regulation of these RNP granules is of particular importance and misregulation of them contributes to the pathogenesis of neurodegenerative diseases (Anderson and Kedersha, 2009; Buchan and Parker, 2009; Zhang et al., 2018). This is at least in part because many of the RNPs contain intrinsically disordered LC domains that can undergo spontaneous self-assembly via LLPS to generate higher-order structures such as gel-like LDs, fibrils and protein aggregations. PAR is a nuclear-acid like polymer and its accumulation in the cytosol may act as a molecular seed to accelerate the nucleation process of LLPS. Indeed, we find that adding PAR to the demixing system dramatically promote the LLPS of hnRNP A1. Consistently, when the manuscript of this study was being prepared, a recent paper published online reporting that the liquid demixing of TDP-43 could be affected by PAR (McGurk et al., 2018). Thus, the reversible modification of PARylation may be an important regulator of RNP granules, which in case of lasting stimuli or excessive PARylation, pathological fibrils or irreversible RNP aggregates may form.

PARylation regulates the interaction of hnRNP A1 with other RNPs and its recruitment to SGs

hnRNP A1 can form protein complexes with TDP-43 and other RNPs to regulate alternative

splicing as well as other aspects of RNAs (Buratti et al., 2005; Mohagheghi et al., 2016). Indeed, we find that hnRNP A1 and TDP-43 can co-phase separate *in vitro* and PARylation strongly regulates their interaction in cells. The hnRNP family proteins constitute an important component of the SGs and are highly regulated at molecular and cellular levels. For example, pathogenic mutations in the GRD of hnRNP A1 accelerate its recruitment to SGs and the fibrillization (Kim et al., 2013), while methylation of hnRNP A2 in the RGG domain suppresses the phase separation of hnRNP A2 (Ryan et al., 2018).

In this study, we show that PARylation regulates the LLPS of hnRNP A1 *in vitro* and the translocation of hnRNP A1 to SGs *in vivo*. We further reveal that hnRNP A1 contains a PARylation site at K298 that is in the GRD as well as a PBM that resides in between the two RRMs. K298 localizes at the C-terminus of the M9 domain, a non-classical nuclear localization sequence of hnRNP A1. Of note, K298 is not within the M9 core sequence (SNFGPMKGGNFGGRSSGPY) that is crucial for the nuclear import and binding to transportin (Iijima et al., FEBS Lett., 2006). Indeed, the PARylation deficient K298A mutant shows no defect in nuclear importing. Instead, it does not translocate to the cytoplasm in response to stress and exhibits the strongest toxicity among the three hnRNP A1 species tested in this study (see Fig. 7). These data implicate the PARylation at K298 as an important mechanism for hnRNP A1 to sense stress and/or serve as a nuclear exporting signal.

The PAR-binding deficient PBM[mut] of hnRNP A1 can respond to stress and translocate to SGs, likely because the signaling mechanims by PARylatino at K298 is intact. However, the SG recruitment of PBM[mut] is less effective than WT and a portion of PBM[mut] forms mis-localized cytoplasmic foci that do not co-localize to SGs. Of note, PBM[mut] shows hyper-PARylation in the *in vitro* assay (see Fig. 2C), possibly at K298 of hnRNP A1, which may result in abnormal activation of the nuclear exporting of PBM[mut]. In the meanwhile, since PBM[mut] shows a greatly reduced capacity to pull down PARylated proteins, we speculate that the association with PAR or PARylated proteins ensures the proper transport and anchoring of hnRNP A1 with SGs. Despite of forming cytoplasmic foci, PBM[mut] is not more toxic than WT hnRNP A1. Instead, PBM[mut] is less

toxic especially when expressed at high levels (see Fig. 7). It is possibly because the reduced association of PBM^{mut} with SGs makes the granules less prone to develop into pathogenic aggregates. And this effect may be particular important when the protein is at high abundance as LLPS of hnRNP A1 occurs with increasing concentrations and PAR promotes this process in a dose-dependent manner. It is worth noting that this is unlike the PBM mutant of TDP-43, which is thought to be more toxic because it could not be recruited to SGs and formed hyper-phosphorylated cytoplasmic foci (McGurk et al., 2018). Nevertheless, downregulation of PARylation levels reduces the cytotoxicity of hnRNP A1 and TDP-43 in both cases (see Fig. 7), which suggests that, under disease conditions, reduction of PAR-mediated association of RNPs and restore of the dynamics of RNP granules may be beneficial.

PARP inhibitors as potential therapeutics for treating ALS

Small-molecule inhibitors of PARP enzymes have received increasing interests since the initial discovery of them in killing *BRCA1/2*-mutant cancer cells. While the involvement of PARP in DNA damage repair has heralded the development of PARP inhibitors in cancer biology and therapy (Davar et al., Curr Med Chem, 2012), the functions of PARP are beyond DNA repair and oncology (Bai, 2015). In this study, we show that Olaparib, the PARP inhibitor approved by FDA for treating ovarian cancer and breast cancer, can significantly reduce the neurotoxicity of hnRNP A1 and TDP-43 in a motor neuron-like cell line, likely due to the pivotal role of PARP in regulating the protein-protein interaction and the dynamics of the disease-associated, aggregation-prone RNP granules. Therefore, we propose Olaparib as a promising candidate for developing ALS drugs, while further animal and clinical studies are needed to evaluate the potential value of Olaparib and other PARP inhibitors as therapeutic targets in ALS and other related neurodegenerative diseases.

In summary (Fig. 8E), in this study, we have demonstrated an important role of PARylation in regulating disease-related RNPs such as RNP A1 and TDP-43. hnRNP A1 is PARylated at K298

in the C-terminus and binds to PAR and/or PARylated proteins via the N-terminal PBM. PARylation and PAR-binding regulates the translocation of hnRNP A1 to SGs – K298A mutant does not respond to stress and remains predominantly nuclear; PBM^{mut} can be recruited to SGs, but a portion of it forms mis-localized cytoplasmic foci that do not co-localize with SGs. The mis-localization of PBM^{mut} is likely due to reduced interaction with other RNPs such as TDP-43. The association and assembly of RNP granules require RNAs, while the protein-protein interaction between the RNPs is regulated by PARylation. Thus, manipulation of PARylation levels modulates the assembly-disassembly dynamics of RNP granules as well as the toxicity of these aggregation-prone proteins. Finally, pharmacological and genetic inhibition of PARP mitigates the neurotoxicity of hnRNP A1 and TDP-43 in MN-like cells and the *Drosophila in vivo* model, which implicates the FDA-approved PARP inhibitor Olaparib as a potential drug candidate for developing therapeutics to treat ALS.

MATERIALS AND METHODS

Plasmids and constructs

To generate pCAG-TDP-43-HA and pCMV-TDP-43-3xFlag plasmids, human TDP-43 DNA was amplified from a TDP-43-Myc plasmid (Jiang et al., 2013) by PCR using the primers specified below. The desired PCR products were subcloned into a pCAG vector (Chen et al., 2014) or a pCMV-3Tag-3B vector (Agilent Technologies) using the ClonExpress™ II One Step Cloning Kit (Vazyme). To generate pCAG-hnRNP A1-Flag plasmid, human hnRNP A1 coding sequence was amplified from the cDNA of Hela cells using by PCR and sub-cloned into a pCAG vector by homologous recombination using the ClonExpress™ II One Step Cloning Kit (Vazyme). The pCAG-K298A-Flag and pCAG-PBM^{mut}-Flag of hnRNP A1 were generated by PCR using the pCAG-hnRNP A1-Flag as a template and hnRNP A1 by site-directed mutagenesis using the Fast Mutagenesis Kit II (Vazyme).

The pET-28a-TDP-43$^{1-274}$-6xhis and pET-28a-TDP-43$^{274-414}$-6xhis constructs for *Escherichia coli* (*E. coli*) expression, the two truncated TDP-43 fragments were amplified from the full-length TDP-43-Myc plasmid (Jiang et al., 2013) and inserted between the BamHI and Xho1 sites in pET-28a-6*his vector.

To generate lentiviral packaging constructs to express Flag, TDP-43-Flag or hnRNP A1-HA in NSC-34 cells, the coding sequence was amplified by PCR using the above expression plasmids as the templates and then sub-cloned into the pCDH-CMV-MCS-EF1-Puro vector using the restriction enzyme sites EcoRI, XbaI, XhoI, BstBI and BamHI.

Primers used for PCR to generate the expression plasmids are summarized below. All constructs were verified by sequencing to ensure the integrity of the cloned open reading frames.

Plasmid	Primer
pCMV-Myc-TDP-43	5'-ATGGCCATGGAGGCCGAATTCATGTCTGAATATATT-3'
	5'-CATGTCTGGATCCCCGCGGCCGCCTACATTCCCCAGCCAGA-3';
pCMV-Myc-TDP-43$^{1-274}$	5'-ATGGCCATGGAGGCCGAATTCATGTCTGAATATATT-3'
	5'-CATGTCTGGATCCCCGCGGCCGCCTAACTTCTTTCTAACTGTCTATT-3'
pCMV-TDP-43-Flag	5'-CGCTCTAGCCCGGGCGGATCCCACCATGTCTGAATATATTCGGGTAAC-3'
	5'-GGTATCGATAAGCTTGATATCTTCATTCCCCAGCCAGAAGA-3'
pCAG-TDP-43-HA	5'-CATCATTTTGGCAAAGAATTCCACCATGTCTGAATATATTCGGGTAAC-3'
	5'-GCTCCCCGGGGGTACCTCGAGTTAAGCGTAGTCTGGGACGTCGTATGGGT ACATTCCCCAGCCAGAAGACTT-3'
pCAG-TDP-43$^{1-274}$-HA	5'-ATGGCCATGGAGGCCGAATTCATGTCTGAATATATT-3'
	5'-CATGTCTGGATCCCCGCGGCCGCCTAACTTCTTTCTAACTGTCTATT-3'
pCAG-hnRNP A1-Flag	5'-CATCATTTTGGCAAAGAATTCCACCATGTCTAAGTCAGAGTCTCCTAAAGAG -3'
	5'-GCTCCCCGGGGGTACCTCGAGCTACTTGTCATCGTCGTCCTTGTAG-3'
pCAG-hnRNP A1 K298A-Flag	5'-CTTTGCAGCACCACGAAACCAAGGTGGCTATGGCGG-3'
	5'-TTCGTGGTGCTGCAAAGTATTGGCCTCCACCGCCATAGG-3'
pCAG-hnRNP A1 K298A-Flag	5'-AAGAGAGCTGTCTCCGCAGAAGATTCTCAAAGACCAGGTGCCCACTTA-3'
	5'- CTTTGAGAATCTTCTGCGGAGACAGCTCTCTTTGGTTCCACAACTCTT-3'
pCAG-hnRNPA1 PBM^{mut}-Flag	5'-AGGTGCCCACTTAACTGTGGTTGTTATATTTGTTGGTGGCATTAAAGAAGA CA CTGAAGAAC-3'
	5'-CCACCAACAAATATAACAACCACAGTTAAGTGGGCACCTGGTCTTTGAGAA-3'
pCDH-CMV-MCS-EF1-Puro-hTDP-43-Flag	5'-GCTCTAGAGCCACCATGTCTGAATATATTCGGGTAACC-3'
	5'-CCGCTCGAGCATTCCCCAGCCAGAAGACTTAG-3'
pET-28a-6*his-TDP-43$^{1-274}$	5'-ATGGCCATGGAGGCCGAATTCATGTCTGAATATATT-3'
	5'-CATGTCTGGATCCCCGCGGCCGCCTAACTTCTTTCTAACTGTCTATT-3'
pET-28a-6*his-TDP-43$^{274-414}$	5'-CAGCAAATGGGTCGCGCCACCGGATCCGGAAGATTTGGTGGT-3'
	5'-GTGGTGGTGGTGGTGCTCGAGCATTCCCCAGCCAGA-3'
Flag oligos	5'-AATTCCTCGAGTTCGAAGCCACCATGGATTACAAGGATGACGACGATAAG GACTATAAGGACGATGATGACAAGGACTACAAAGATGATGACGATAAATAGG-3'
	5'-GATCCCTATTTATCGTCATCATCTTTGTAGTCCTTGTCATCATCGTCCTTAT AGTCCTTATCGTCGTCATCCTTGTAATCCATGGTGGCTTCGAACTCGAGG-3'

Cell culture and transfection

Both Hela and HEK-293T cells were cultured in Dulbecco's modified Eagle's medium (sigma, D0819) supplemented with 10% fetal bovine serum (FBS, BioWest). NSC-34 cells were cultured in RPMI 1640 medium (Gibco, 11875-093) containing 10% (v/v) FBS. All cells were incubated at 37 °C in a humidified atmosphere of 95% air and 5% CO_2. PolyJet™ reagent (SignaGen, SL100688) was used for transfection of plasmids into HeLa and HEK-293 T cells. Cells were transfected for 24–72 h before harvest for the subsequent experiments. For knockdown experiments, siRNA (Genepharma,Shanghai) were transfected into cells using Lipofectamine™ RNAiMAX Transfection Reagent (Invitrogen, 13778150) according to the manufacture's instruction. siRNA was incubated for 48-72 h before cells were harvest. The siRNA oligos used in this study are summarized below.

Table 2. Oligos for siRNA knockdown used in this study.

Gene	siRNA
Si-Ctrl	5'-GCGGUGAAGUUAGAUUACATT-3'
Si-human PARG	5'-GCGGUGAAGUUAGAUUACATT-3'
Si-mouse PARP1	5'-CGACGCUUAUUACUGUACUTT-3'

Stress granule induction

HeLa cells grown on coverslips in a 12-well plate were treated with 100 µM of $NaAsO_2$ or PBS for 30 min before they were fixed with 4% paraformaldehyde. For recovery, the medium containing NaAsO2 was removed and the cells were incubated in fresh medium for indicated time before fixation. The formation of stress granules was evaluated by the subsequent immunocytochemistry assays.

Immunocytochemistry and confocal imaging

After transfection and drug treatments, cells grown on the cover slips in 24-well plate were fixed with 4% paraformaldehyde in PBS for 15 min at room temperature, permeabilized in 0.5% Triton X-100 in PBS for 10 min, and blocked with 3% goat serum in PBST (PBS + 0.1% Triton X-100) for

30 min at room temperature. The primary and secondary antibodies in the blocking buffer were then incubated at 4°C overnight or at room temperature for 1 h. After 3 washes with PBST, cells were mounted on glass slides using Vectashield Antifade Mounting Medium with DAPI (Vector Laboratories). Fluorescent images were taken with Leica TCS SP8 confocal microscopy system using a 63X oil objective (NA=1.4). Images were processed and assembled into figures using X (Leica) and Adobe Photoshop CS6.

Protein extraction and Western blotting

Total protein was extracted from cells in 2% SDS extraction buffer (50 mM Tris pH 6.8, 2% SDS, mercaptoethanol, 12.5% glycerol and 0.04% bromophenol blue) containing the protease inhibitor cocktail (Roche, 04693132001), 20 µM Olaparib (Selleck, S1060) and 8 µM ADP-HPD (Millopore, 118415). To separation of soluble and insoluble proteins, cells or fly heads were lysed on ice in RIPA buffer (50 mM Tris pH 8.0, 150 mM NaCl, 1% NP-40, 5 mM EDTA, 0.5% sodium deoxycholate, 0.1% SDS) supplemented with protease and phosphatase inhibitors. After sonication, the homogenates were centrifuged at 13,000 x g for 10-20 min at 4°C. The supernatant was used as the soluble fraction and the pellets containing insoluble fractions were dissolved in a 9 M urea buffer (9 M urea, 50 mM Tris buffer, pH 8.0) after wash.

Proteins were separated by 10% Bis-Tris SDS-PAGE (Invitrogen), immunoblotted with the primary and secondary antibodies. Detection was performed using the High-sig ECL Western Blotting Substrate (Tanon). Images were captured using an Amersham Imager 600 (GE Healthcare) and densitometry was measured using ImageQuant TL Software (GE Healthcare). The contrast and brightness were optimized equally using Adobe Photoshop CS6 (Adobe Systems Inc.). All experiments were normalized to GAPDH or tubulin as indicated in the figures.

Co-immunoprecipitation

HeLa cells were lysed in IP buffer (25 mM Tris•HCl pH 7.4, 150 mM NaCl, 1% NP-40, 1 mM EDTA, 5% glycerol) containing protease inhibitor cocktails, 20 µM Olaparib and 8 µM ADP-HPD.

To test if the interaction between hnRNP A1 and TDP-43 was RNA-dependent, cell lysates were treated with 100 mg/ml of RNase A for 30 min (Qiagen), and then incubated with mouse anti-Flag or rabbit anti-HA antibody on a rotary shaker at 4°C overnight. Mouse or rabbit IgG (Santa Cruz, sc-2025 or sc2027) was used as a control for the pull-down specificity. Anti-FLAG® M2 Affinity Gel (Sigma, A2220) or Dynabeads® Protein G beads (Novex) were then added and incubated at room temperature for 2 h. The gel or beads were then collected according to the manufacture's instruction and eluted in the 2xLDS sample buffer (Invitrogen) before the subsequent Western blotting assay.

Antibodies

The following antibodies were used for Western blotting, immunoprecipitation and immunocytochemistry assays: mouse anti-FLAG (Sigma, F3165), mouse anti-HA (Proteintech, 66006-1), anti-pan-ADP-ribose binding reagent (Millipore, MABE1016), rabbit anti-HA (CST, 3724T), rabbit anti-TDP-43 (Proteintech, 10782-2-AP), rabbit anti-TIAR (CST, 8509S), rabbit anti-hnRNPA1 (CST, 8443S), rabbit anti-PAR (ENZO, ALX-210-890A-0100), mouse anti-GAPDH (Proteintech, 60004-1) and mouse anti-Tubulin (MBL, PM054). HRP conjugated secondary antibodies, goat anti-mouse (Sigma, A4416) and goat anti-rabbit (Sigma, A9169). Fluorescent secondary antibodies: goat anti-mouse-Alexa Fluor 488 (Life Technologies, A11012); goat anti-rabbit-Alexa Fluor 568 (Life Technologies, A11012).

RNA extraction and real-time quantitative PCR (qPCR)

Total RNAs were extracted from cells or fly heads (40 heads per group) using TRIzol (Invitrogen) according to the manufacturer's instructions. After DNase (Promega) treatment to remove genomic DNA, the reverse transcription was performed using the High-Capacity cDNA Reverse Transcription Kit (Applied biosystems). The cDNA was then used in real-time PCR experiment using the SYBR Green qPCR Master Mix (Bimake) with the QuantStudio™ 6 Flex Real-Time PCR system (Life Technologies). The mRNA levels of *actin* were used as an internal control to

normalize the mRNA levels of genes of interest. The primers for qPCR are listed below.

Table 3. qPCR primers used in this study.

Gene	Primer
Fly Parp	5'-ATGAAGTACGGAGGCCAACC-3'
	5'-TCTTCACCTGACGCAAACCA-3'
Fly Actin	5'-GAGCGCGGTTACTCTTTCAC-3'
	5'-GCCATCTCCTGCTCAAAGTC-3'

TDP-43 and hnRNP A1 protein purification

Purification of TDP-43 aa1-274: BL21(DE3) competent E. coli (TransGenBiotech, CD601-03) transformed with pET-28a-6*his-TDP-43 aa1-274 was cultivated on LB plate containing Kanamycin (50 μg/ml) at 37°C overnight. Colonies were subsequently inoculated into 5ml of kanamycin-containing LB medium and cultivated at 37 °C for 12 h, then the cultures were diluted 10-fold using LB medium. When the absorbance value at 600nm measured in the diluted cultures reached 0.6, the culture temperature was shift to 16°C and protein expression was induced by 250 μM of IPTG. 18 h later, cells were harvested by centrifugation at 4000 rpm for 20 min at 4 °C and lysed with 50 mL of lysis buffer (50mM NaH2PO4, 500mM NaCl, pH 8.0, 10mM imidazole, 1mM DTT, 1mM PMSF) for 15 min. After cell lysates were filtered with a 0.2 mm filter, protein was purified by Ni column (GE Healthcare, USA), eluted with elution buffer (50mM NaH2PO4, 500mM NaCl, pH 8.0 and 50mM imidazole), transfer to storage buffer (50mM Tris-HCl at pH 7.5, 300mM NaCl), freshly frozen in liquid N2 and stored at -80 °C.

Purification of TDP-43 aa274-414: Different from the method for purification of TDP-43 aa1-274, TDP-43 aa274-414 expressing BL21(DE3) cells were lysed in a lysis buffer (50mM Tris-HCl, pH 8.0, and 6M guanidine hydrochloride) at room temperature and sonicated at 4 °C. Cell lysate was centrifuged at 14,000 rpm for 1h at 4°C. Proteins purified from the supernatant by using a Ni column were eluted with the buffer containing 50mM Tris-HCl at pH 8.0, 6M guanidine hydrochloride and 50mM imidazole. After further purified by HPLC (Agilent) with the elution buffer containing 35% (v/v) acetonitrile, the purified TDP-43 aa274-414 were freeze-dried by

FreeZone lyophilizers (Thermo Fisher) and stored at -20 °C.

Purification of hnRNPA1-WT: The pET9d-hnRNPA1-WT was transformed into Plyss strains. hnRNPA1-WT protein expression was induced by 0.5mM of IPTG at 25°C for 14 h. Cells were harvested and lysed in a lysis buffer (50mM Tris-HCl at pH 7.5, 2mM DTT, 1mM PMSF and 5% glycerin) at 4 °C. The cell lysate was centrifuged at 4°C, 14,000 rpm for 1h. The cleared supernatant was filtered using a 0.2 mm filter and the protein was purified over SP column (GE Healthcare, USA). The proteins in the column were eluted with the elution buffer (50mM Tris-HCl, pH 7.5, 1M NaCl, 2mM DTT, 5% glycerin and 50mM imidazole). Subsequently, the eluted proteins were further purified by superdex 75 (GE Healthcare, USA) using the buffer containing 50mM Tris-HCl, pH 7.5, 500mM NaCl, 2mM DTT and 5% glycerin. The purified proteins were freeze-dried by FreeZone lyophilizers (Thermo Fisher) and was stored at –80 °C.

All the purified proteins were confirmed by Coomassie brilliant blue staining and Western blotting.

In vitro PARylation assay

In vitro PARylation assay was performed according to the protocol adapted from (Slade et al., 2011). Briefly, substrate proteins were incubated with PARP1 (Sino Biological) in the presence of NAD in reaction buffer (50 mM Tris-Hcl, pH 7.4, 2 mM $MgCl_2$) with or without 2.5 μg of ssDNA (Sigma) at 37 °C for 30 min. The reactions were stopped by adding 20 μM Olaparib and the products were examined by SDS-PAGE and Western blotting.

In vitro phase separation

For the *in vitro* demixing experiments, purified hnRNP A1, PAR polymers (Trevigen, 4336-100-01) and NaCl at indicated concentrations were mixed into the phase separation buffer (50mM Tris-HCl, pH 7.5, 1% PEG (Sigma)) and incubated for 3 min at room temperature. For co-LLPS, hnRNP A1 protein were incubated with TDP-43 in the reaction buffer (50mM Tris-HCl, pH 7.5, 100mM NaCl) at the similar condition. 5 μL of each reaction samples were dropped onto a

coverslip and imaged using a Leica microscope with differential interference contrast (DIC) within 10 min.

Lentivirus production and infection

293T cells were co-transfected with psPAX2, pMD2.G and pCDH-CMV-MCS-EF1-Puro plasmids of Flag, TDP-43-Flag, WT and the mutant hnRNP A1-HA (K298A and PBM^{mut}), using the PolyJet™ reagent. Cell culture medium was collected and filtered through a 0.45 μm syringe filter (Millipore, SLHV033RB) at 48 h after transfection. The lentivirus was then concentrated using Lenti-X™ Concentrator (Clontech, PT4421-2). The 20-fold concentrated medium containing the desired lentivirus was used to infect NSC-34 cells in the subsequent experiments.

Cell viability assay

NSC-34 cells were seeded into 96-well plates (Corning) at the density of $1.28×10^4$ cells/well and cultured in 100 μL of cultre medium containing the lentivirus particles for the expression of the proteins of interest. 48 h later, the infection medium containing lentivirus was removed and replaced with fresh medium. 48-72 h post infection, cell viability was examined using the Cell Counting Kit-8 (CCK-8) (Dojindo) according to the manufacturer's instructions. Briefly, 10 μL of CCK-8 solution was added to each well. After incubation at 37℃ for 2.5 h, the absorbance at at 450 nm was measured with a Synergy2 microplate reader (BioTek Instruments).

TUNEL staining

TUNEL staining assay was performed using the TMR red *in situ* Cell Death Detection Kit (Sigma-Aldrich) according to the manufacturer's instruction. The transfected NSC-34 cells grown on coverslips in a 24-well plate were fixed with 4% paraformaldehyde for 15 min and permeabilized with PBST (1xPBS + 0.5% Triton X-100) for 10 min at room temperature, followed by incubation with 18 μl labeling solution plus 2 μl enzyme solution at 37 °C for 1 h. Cell were

mounted on glass slides using Vectashield Antifade Mounting Medium with DAPI (Vector Laboratories) and imaged using the Leica TCS SP8 confocal microscopy system.

Drosophila strains

The following strains were obtained from the Bloomington *Drosophila* Stock Center (BDSC): RNAi-*Parp* (#57265), *elav*GS (#43642), RNAi-*mCherry* (#35785, a control for *in vivo* RNAi knockdown), UAS-*lacZ* (#8529, a control for UAS transgene expression). UAS-TDP-43 and GRM-Gal4 flies were previously described (Kim et al., 2013). All the flies were raised on standard cornmeal media and maintained at 25 °C and 60% relative humidity.

Climbing ability and Lifespan assays

For the climbing assays, 20 flies per vial, 5-8 vials per group were tested. All flies were transferred into an empty polystyrene vial and gently tapped down to the bottom before 15 min adapted time for flies. The number of flies climbing over the height of 3 cm within 10 seconds was recorded. The test was repeated three times for each vial and 6~8 vials per group were tested.

For the lifespan experiments, 20 flies per vial, 7-9 vials per group were tested. Flies were transferred to fresh fly food every 3 days. The flies lost prior to natural death because of escape or accidental death were excluded from the final analysis. The median lifespan was calculated as the mean of the medians of each vial in a group, whereas the "50% survival" shown on the survival curves is derived from compilation of all vials of a group. For adult-onset, neuronal expression of the RNAi transgenes using the *elav*-GS driver (Osterwalder et al., 2001), flies were raised at 25 °C and 60% relative humidity on regular fly food supplemented with 80 µg/ml RU486 (TCI).

Statistical analysis

Unless otherwise noted, statistical significance in this study is determined by unpaired, two-tailed Student's t-test at $*p < 0.05$, $**p < 0.01$, and $***p < 0.001$. Error bars represent the standard error of the mean (SEM).

ACKNOWLEDGEMENTS

We thank the BDSC for providing fly strains, X. Gui for assistance in protein pruification, B. Yang for help in *in vitro* PARylation assay, S. Qiu and S. Zhang for technical supports, and J. Yuan, A. Li and members of the Liu lab and the Fang lab for helpful discussion and critical reading of the manuscript. This study was supported by grants from the National Key R&D Program of China (No. 2016YFA0501902) and the National Natural Science Foundation of China (NSFC, No. 31471017 and No. 81671254) to Y.F., the NSFC grant (No. 31470748) to C.L., and funding from the Shanghai Science and Technology Committee (No. 18ZR1448300) to A.D.

AUTHOR CONTRIBUTIONS

YD, CL and YF conceived the research; YD, AD, CL and YF designed the experiments; YD, AD, JG, LS, GD, CW, LS, BQ, and KT performed the experiments; YD, AD, GD, CW, XD and HJ contributed important new reagents; YD, AD, JG, GD, CW, LS, KZ, and YZ analyzed the data; YD, AD, JG, CL and YF interpreted and discussed the results; YD, AD, GD and YF prepared the figures; and YD, AD, GD, CL and YF wrote the paper. All authors read and approved the final manuscript.

DECLARATION OF INTERESTS

The authors declare that they have no competing interests.

SUPPORTING INFORMATION

Supporting information includes 6 figures.

REFERENCE

Ahel, I., Ahel, D., Matsusaka, T., Clark, A.J., Pines, J., Boulton, S.J., and West, S.C. (2008). Poly(ADP-ribose)-binding zinc finger motifs in DNA repair/checkpoint proteins. Nature 451, 81-85.

Ahel, D., Horejsi, Z., Wiechens, N., Polo, S.E., Garcia-Wilson, E., Ahel, I., Flynn, H., Skehel, M., West, S.C., Jackson, S.P., et al. (2009). Poly(ADP-ribose)-dependent regulation of DNA repair by the chromatin remodeling enzyme ALC1. Science 325, 1240-1243.

Anderson, P., and Kedersha, N. (2009). RNA granules: post-transcriptional and epigenetic modulators of gene expression. Nat Rev Mol Cell Biol 10, 430-436.

Andrabi, S.A., Kim, N.S., Yu, S.W., Wang, H., Koh, D.W., Sasaki, M., Klaus, J.A., Otsuka, T., Zhang, Z., Koehler, R.C., et al. (2006). Poly(ADP-ribose) (PAR) polymer is a death signal. Proc Natl Acad Sci U S A 103, 18308-18313.

Anko, M.L., and Neugebauer, K.M. (2012). RNA-protein interactions in vivo: global gets specific. Trends Biochem Sci 37, 255-262.

Bai, P. (2015). Biology of Poly(ADP-Ribose) Polymerases: The Factotums of Cell Maintenance. Mol Cell 58, 947-958.

Brady, O.A., Meng, P., Zheng, Y., Mao, Y., and Hu, F. (2011). Regulation of TDP-43 aggregation by phosphorylation and p62/SQSTM1. J Neurochem 116, 248-259.

Brochier, C., Jones, J.I., Willis, D.E., and Langley, B. (2015). Poly(ADP-ribose) polymerase 1 is a novel target to promote axonal regeneration. Proc Natl Acad Sci U S A 112, 15220-15225.

Buchan, J.R., and Parker, R. (2009). Eukaryotic stress granules: the ins and outs of translation. Mol Cell 36, 932-941.

Buratti, E., Brindisi, A., Giombi, M., Tisminetzky, S., Ayala, Y.M., and Baralle, F.E. (2005). TDP-43 binds heterogeneous nuclear ribonucleoprotein A/B through its C-terminal tail: an important region for the inhibition of cystic fibrosis transmembrane conductance regulator exon 9 splicing. J Biol Chem 280, 37572-37584.

Chang, P., Jacobson, M.K., and Mitchison, T.J. (2004). Poly(ADP-ribose) is required for spindle

assembly and structure. Nature 432, 645-649

Chen, Y., Wang, Y., Erturk, A., Kallop, D., Jiang, Z., Weimer, R.M., Kaminker, J., and Sheng, M. (2014). Activity-induced Nr4a1 regulates spine density and distribution pattern of excitatory synapses in pyramidal neurons. Neuron 83, 431-443.

Cohen, T.J., Hwang, A.W., Restrepo, C.R., Yuan, C.X., Trojanowski, J.Q., and Lee, V.M. (2015). An acetylation switch controls TDP-43 function and aggregation propensity. Nat Commun 6, 5845.

D'Ambrogio, A., Buratti, E., Stuani, C., Guarnaccia, C., Romano, M., Ayala, Y.M., and Baralle, F.E. (2009). Functional mapping of the interaction between TDP-43 and hnRNP A2 in vivo. Nucleic Acids Res 37, 4116-4126.

Dammer, E.B., Fallini, C., Gozal, Y.M., Duong, D.M., Rossoll, W., Xu, P., Lah, J.J., Levey, A.I., Peng, J., Bassell, G.J., et al. (2012). Coaggregation of RNA-binding proteins in a model of TDP-43 proteinopathy with selective RGG motif methylation and a role for RRM1 ubiquitination. PLoS One 7, e38658.

Davar, D., Beumer, J.H., Hamieh, L., and Tawbi, H. (2012). Role of PARP inhibitors in cancer biology and therapy. Current medicinal chemistry 19, 3907-3921

Deng, W. (2009). PARylation: strengthening the connection between cancer and pluripotency. Cell Stem Cell 5, 349-350.

Dreyfuss, G., Matunis, M.J., Pinol-Roma, S., and Burd, C.G. (1993). hnRNP proteins and the biogenesis of mRNA. Annu Rev Biochem 62, 289-321

Fong, P.C., Boss, D.S., Yap, T.A., Tutt, A., Wu, P., Mergui-Roelvink, M., Mortimer, P., Swaisland, H., Lau, A., O'Connor, M.J., et al. (2009). Inhibition of poly(ADP-ribose) polymerase in tumors from BRCA mutation carriers. The New England journal of medicine 361, 123-134.

Frizzell, K.M., Gamble, M.J., Berrocal, J.G., Zhang, T., Krishnakumar, R., Cen, Y., Sauve, A.A., and Kraus, W.L. (2009). Global analysis of transcriptional regulation by poly(ADP-ribose) polymerase-1 and poly(ADP-ribose) glycohydrolase in MCF-7 human breast cancer cells. J Biol Chem 284, 33926-33938.

Gagne, J.P., Hunter, J.M., Labrecque, B., Chabot, B., and Poirier, G.G. (2003). A proteomic approach to the identification of heterogeneous nuclear ribonucleoproteins as a new family of poly(ADP-ribose)-binding proteins. The Biochemical journal 371, 331-340.

Gagne, J.P., Isabelle, M., Lo, K.S., Bourassa, S., Hendzel, M.J., Dawson, V.L., Dawson, T.M., and Poirier, G.G. (2008). Proteome-wide identification of poly(ADP-ribose) binding proteins and poly(ADP-ribose)-associated protein complexes. Nucleic Acids Res 36, 6959-6976.

Gottschald, O.R., Malec, V., Krasteva, G., Hasan, D., Kamlah, F., Herold, S., Rose, F., Seeger, W., and Hanze, J. (2010). TIAR and TIA-1 mRNA-binding proteins co-aggregate under conditions of rapid oxygen decline and extreme hypoxia and suppress the HIF-1alpha pathway. J Mol Cell Biol 2, 345-356.

Hanai, S., Kanai, M., Ohashi, S., Okamoto, K., Yamada, M., Takahashi, H., and Miwa, M. (2004). Loss of poly(ADP-ribose) glycohydrolase causes progressive neurodegeneration in Drosophila melanogaster. Proc Natl Acad Sci U S A 101, 82-86.

He, Y., and Smith, R. (2009). Nuclear functions of heterogeneous nuclear ribonucleoproteins A/B. Cell Mol Life Sci 66, 1239-1256.

Hofweber, M., Hutten, S., Bourgeois, B., Spreitzer, E., Niedner-Boblenz, A., Schifferer, M., Ruepp, M.D., Simons, M., Niessing, D., Madl, T., et al. (2018). Phase Separation of FUS Is Suppressed by Its Nuclear Import Receptor and Arginine Methylation. Cell 173, 706-719 e713.

Ji, Y., and Tulin, A.V. (2009). Poly(ADP-ribosyl)ation of heterogeneous nuclear ribonucleoproteins modulates splicing. Nucleic Acids Res 37, 3501-3513

Jiang, L.L., Che, M.X., Zhao, J., Zhou, C.J., Xie, M.Y., Li, H.Y., He, J.H., and Hu, H.Y. (2013). Structural transformation of the amyloidogenic core region of TDP-43 protein initiates its aggregation and cytoplasmic inclusion. J Biol Chem 288, 19614-19624.

Kim, H.J., Kim, N.C., Wang, Y.D., Scarborough, E.A., Moore, J., Diaz, Z., MacLea, K.S., Freibaum, B., Li, S., Molliex, A., et al. (2013). Mutations in prion-like domains in hnRNPA2B1 and hnRNPA1 cause multisystem proteinopathy and ALS. Nature 495, 467-473.

Kim, H.J., Raphael, A.R., LaDow, E.S., McGurk, L., Weber, R.A., Trojanowski, J.Q., Lee, V.M., Finkbeiner, S., Gitler, A.D., and Bonini, N.M. (2014). Therapeutic modulation of eIF2alpha phosphorylation rescues TDP-43 toxicity in amyotrophic lateral sclerosis disease models. Nat Genet 46, 152-160.

Kotova, E., Jarnik, M., and Tulin, A.V. (2009). Poly (ADP-ribose) polymerase 1 is required for protein localization to Cajal body. PLoS Genet 5, e1000387.

Lee, E.B., Lee, V.M., and Trojanowski, J.Q. (2012). Gains or losses: molecular mechanisms of TDP43-mediated neurodegeneration. Nat Rev Neurosci 13, 38-50.

Leung, A.K., Vyas, S., Rood, J.E., Bhutkar, A., Sharp, P.A., and Chang, P. (2011). Poly(ADP-ribose) regulates stress responses and microRNA activity in the cytoplasm. Mol Cell 42, 489-499.

Li, W., Reeb, A.N., Lin, B., Subramanian, P., Fey, E.E., Knoverek, C.R., French, R.L., Bigio, E.H., and Ayala, Y.M. (2017). Heat Shock-induced Phosphorylation of TAR DNA-binding Protein 43 (TDP-43) by MAPK/ERK Kinase Regulates TDP-43 Function. J Biol Chem 292, 5089-5100.

Iijima, M., Suzuki, M., Tanabe, A., Nishimura, A., and Yamada, M. (2006). Two motifs essential for nuclear import of the hnRNP A1 nucleocytoplasmic shuttling sequence M9 core. FEBS Lett 580, 1365-1370.

Lin, Y., Protter, D.S., Rosen, M.K., and Parker, R. (2015). Formation and Maturation of Phase-Separated Liquid Droplets by RNA-Binding Proteins. Mol Cell 60, 208-219.

Lunde, B.M., Moore, C., and Varani, G. (2007). RNA-binding proteins: modular design for efficient function. Nat Rev Mol Cell Biol 8, 479-490.

Martello, R., Leutert, M., Jungmichel, S., Bilan, V., Larsen, S.C., Young, C., Hottiger, M.O., and Nielsen, M.L. (2016). Proteome-wide identification of the endogenous ADP-ribosylome of mammalian cells and tissue. Nature Communications 7, 12917.

Martire, S., Mosca, L., and d'Erme, M. (2015). PARP-1 involvement in neurodegeneration: A focus on Alzheimer's and Parkinson's diseases. Mech Ageing Dev 146-148, 53-64.

McGurk, L., Gomes, E., Guo, L., Mojsilovic-Petrovic, J., Tran, V., Kalb, R.G., Shorter, J., and

Bonini, N.M. (2018). Poly(ADP-Ribose) Prevents Pathological Phase Separation of TDP-43 by Promoting Liquid Demixing and Stress Granule Localization. Molecular Cell.

Mohagheghi, F., Prudencio, M., Stuani, C., Cook, C., Jansen-West, K., Dickson, D.W., Petrucelli, L., and Buratti, E. (2016). TDP-43 functions within a network of hnRNP proteins to inhibit the production of a truncated human SORT1 receptor. Hum Mol Genet 25, 534-545

Muller-McNicoll, M., and Neugebauer, K.M. (2013). How cells get the message: dynamic assembly and function of mRNA-protein complexes. Nat Rev Genet 14, 275-287.

Neumann, M., Sampathu, D.M., Kwong, L.K., Truax, A.C., Micsenyi, M.C., Chou, T.T., Bruce, J., Schuck, T., Grossman, M., Clark, C.M., et al. (2006). Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. Science 314, 130-133.

Nichols, R.C., Wang, X.W., Tang, J., Hamilton, B.J., High, F.A., Herschman, H.R., and Rigby, W.F. (2000). The RGG domain in hnRNP A2 affects subcellular localization. Exp Cell Res 256, 522-532.

Niere, M., Mashimo, M., Agledal, L., Dolle, C., Kasamatsu, A., Kato, J., Moss, J., and Ziegler, M. (2012). ADP-ribosylhydrolase 3 (ARH3), not poly(ADP-ribose) glycohydrolase (PARG) isoforms, is responsible for degradation of mitochondrial matrix-associated poly(ADP-ribose). J Biol Chem 287, 16088-16102.

Qamar, S., Wang, G., Randle, S.J., Ruggeri, F.S., Varela, J.A., Lin, J.Q., Phillips, E.C., Miyashita, A., Williams, D., Strohl, F., et al. (2018). FUS Phase Separation Is Modulated by a Molecular Chaperone and Methylation of Arginine Cation-pi Interactions. Cell 173, 720-734 e715.

Osterwalder T, Yoon KS, White BH, Keshishian H (2001) A conditional tissue-specific transgene expression system using inducible GAL4. Proc. Natl Acad. Sci. USA 98, 12596-12601.

Ryan, V.H., Dignon, G.L., Zerze, G.H., Chabata, C.V., Silva, R., Conicella, A.E., Amaya, J., Burke, K.A., Mittal, J., and Fawzi, N.L. (2018). Mechanistic View of hnRNPA2 Low-Complexity Domain Structure, Interactions, and Phase Separation Altered by Mutation and Arginine Methylation. Mol Cell 69, 465-479 e467.

Siomi, H. (1995). A nuclear localization domain in the hnRNP A1 protein. The Journal of Cell

Biology 129, 551-560.

Singh, H.R., Nardozza, A.P., Moller, I.R., Knobloch, G., Kistemaker, H.A.V., Hassler, M., Harrer, N., Blessing, C., Eustermann, S., Kotthoff, C., et al. (2017). A Poly-ADP-Ribose Trigger Releases the Auto-Inhibition of a Chromatin Remodeling Oncogene. Mol Cell 68, 860-871 e867.

Slade, D., Dunstan, M.S., Barkauskaite, E., Weston, R., Lafite, P., Dixon, N., Ahel, M., Leys, D., and Ahel, I. (2011). The structure and catalytic mechanism of a poly(ADP-ribose) glycohydrolase. Nature 477, 616-620.

Virag, L., Robaszkiewicz, A., Rodriguez-Vargas, J.M., and Oliver, F.J. (2013). Poly(ADP-ribose) signaling in cell death. Mol Aspects Med 34, 1153-1167.

Zhang, K., Daigle, J.G., Cunningham, K.M., Coyne, A.N., Ruan, K., Grima, J.C., Bowen, K.E., Wadhwa, H., Yang, P., Rigo, F., et al. (2018). Stress Granule Assembly Disrupts Nucleocytoplasmic Transport. Cell.

Zhang, Y., Wang, J., Ding, M., and Yu, Y. (2013). Site-specific characterization of the Asp- and Glu-ADP-ribosylated proteome. Nat Methods 10, 981-984.

Figure 1. PARylation levels influence the assembly-disassembly dynamics of SGs containing ALS-related RNPs.

(**A-C**) PARP inhibition suppresses SG assembly. (**A**) Representative confocal images of Hela cells treated with PBS (no stress) or 100 μM arsenite (stress) for indicated time in the absence (DMSO) or presence of the PARP inhibitor Olaparib (20 μM). All cells are transfected with TDP-43-HA (anti-HA) and stained for SGs by anti-TIAR (arrowheads), and the merge images with DAPI staining of DNA (blue) are shown. (**B**) The percentage of cells with SGs and TDP-43+ SGs from (A) is quantified. (**C**) Western blot analysis confirms the decrease of overall PARylation levels by Olaparib, with GAPDH as a loading control. (**D-F**) Hydrolysis of PAR by PARG is required for SG disassembly. (**D**) Cells transfected with siRNA (si-PARG) or scrambled siRNA (si-Ctrl) are stressed with arsenite (100 μM) for 30 min and then washout for indicated time. All cells are transfected with hnRNP A1-Flag (anti-Flag) and stained for SGs by anti-TIAR (arrowheads), and the merge images with DAPI staining of DNA (blue) are shown. (**E**) Percentage of cells with SGs or hnRNP A1+ SGs from (D). (**F**) Western blot analysis confirms PARG KD and increase of overall PARylation levels in the cells by si-PARG. Data are shown as mean ± SEM; n = over 100 cells each condition, pooled results of 3 independent repeats; *$p <$ 0.05, **$p <$ 0.01; ns, not significant; ud, undetected; Student's t-test. Scale bars: 10 μm.

Figure 2. PARylation and PAR binding of TDP-43 and hnRNP A1 *in vitro* and in cells

(**A-B**) HA-tagged TDP-43 (A) or Flag-tagged hnRNP A1 (B) is expressed and immunoprecipitated from Hela cells that are treated with or without H_2O_2, and examined by Western blotting with indicated antibodies. hnRNP A1 shows both steady state and induced levels of PARylation, whereas no PARylation of TDP-43 was detected even when the blot is overexposed to facilitate the detection. Asterisk, the rabbit anti-HA IgG heavy chain. (**C**) Recombinant purified His-tagged TDP-43$^{1-274}$, TDP-43$^{274-414}$ and full-length hnRNP A1 proteins (4 μg) were subjected to *in vitro* PARylation assay and then separated on SDS-PAGE gel. Coomassie Brilliant Blue staining confirms equal protein loading (upper panel, arrows indicate

the substrate proteins), followed by immunoblotting with anti-PAR (lower panel). ssDNA was added in the *in vitro* reaction system to activate PARP1. The PARylated bands and smears of recombinant TDP-43 and hnRNP A1 as well as PARP1 are indicated. (**D**) Functional domains of human hnRNP A1 protein. RRM: RNA recognition motif; RGG: RGG box RNA binding domain; M9: a nuclear targeting sequence termed as M9; GRD: glycine-rich domain; PBM: PAR-binding motif. The consensus sequence and the residues mutated in the PBM and putative PARylation site are indicated. h, hydrophobic amino acid residues; b, basic amino acid residues (Ji and Tulin, 2009). (**E**) Western blotting and quantification analysis of the solubility of WT, K298A and PBM[mut] hnRNP A1 proteins. S, soluble in RIPA buffer; I, insoluble in RIPA buffer, and the pellets are re-suspended with 9 M urea. Mean ± SEM; n = 3; *$p < 0.05$; two-way ANOVA. (**F**) Confirmation of PARylation or PAR-binding deficit of K298A and PBM[mut] compared to WT hnRNP A1. hnRNP A1-Flag was immunoprecipitated from Hela cells with anti-Flag and examined by Western blotting with anti-PAR. PARylated hnRNP A1 and other proteins (#) co-immunoprecipitated with hnRNP A1 are indicated. The blot is overexposed to facilitate the detection of less abundant PARylated proteins.

Figure 3. The PARylation and PAR-binding mutants of hnRNP A1 K298A and PBM[mut] exhibit deficits in translocation and targeting to SGs.

(**A**) Representative images of Hela cells transfected with WT, K298A or PBM[mut] of Flag-tagged hnRNP A1 and treated with PBS (no stress) or arsenite (stress) for 30 min. Arrows, hnRNP A1+/TIAR- cytoplasmic foci; arrowheads, hnRNP A1+/TIAR+ SGs; asterisks, hnRNP A1-/TIAR+ SGs. (**B-D**) The percentage of cells with SGs (B), hnRNP A1+/TIAR+ SGs (C), or hnRNP A1+/TIAR- cytoplasmic foci (D) is quantified. Mean ± SEM; n = over 300 cells each condition, pooled results of 3 independent repeats; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$; ns, not significant; Student's *t*-test. Scale bar: 10 μm.

Figure 4. PAR promotes LLPS of hnRNP A1 *in vitro*.

(**A**) hnRNP A1 forms dynamic LDs by LLPS *in vitro*. The concentrations of hnRNP A1 and the NaCl are shown. The higher hnRNP A1 and lower NaCl concentrations, the more and larger LDs are formed. (**B**) Addition of PAR promotes LLPS of hnRNP A1 in a dose-dependent manner. (**C**) The same concentration of heparin (7 μM) cannot induce LLPS of hnRNP A1 at the same condition. (**D**) PAR (7 μM) alone does not phase separate *in vitro*. Scale bars: 10μm.

Figure 5. Co-phase separation of hnRNP A1 and TDP-43 *in vitro* and the effect of PAR on this process.

(**A**) hnRNP A1 alone does not form spontaneous LDs by LLPS at the indicated condition *in vitro*. (**B-E**) co-LLPS experiments of hnRNP A1 with BSA (B), TDP-43$^{1-274}$ (C), or TDP-43$^{274-414}$ (D-E) with the indicated concentrations of each protein. (**B'-E'**) LLPS experiments of BSA (B'), TDP-43$^{1-274}$ (C'), or TDP-43$^{274-414}$ (D'-E') alone at the same condition as in (B-E). All tests in A-E' are carried out at the salt concentration of 100 mM (NaCl). Scale bars: 10 μm.

Figure 6. PARrlation modulates the interaction between hnRNP A1 and TDP-43.

(**A**) Flag-tagged hnRNP A1 is expressed in Hela cells treated with vehicle control (DMSO), Olaparib (20 μM) or H_2O_2 (500 μM), and immunoprecipitated with anti-Flag. They are then separated by SDS-PAGE and examined by Western blotting. (**B**) The level of co-IP of TDP-43 with hnRNP A1 is quantified as ratio of co-IP of TDP-43/input TDP-43 and shown as percentage to the DMSO control group. (**C**) Western blot analysis of cells transfected with WT, K298A or PBM^{mut} of Flag-tagged hnRNP A1, immunoprecipitated with anti-Flag, and blotted with indicated antibodies. 2xPBM^{mut}, cells transfected with 2 times of the PBM^{mut} expression plasmids. (**D**) The relative level of TDP-43 co-immunoprecipitated with WT or mutant hnRNP A1 is normalized to TDP-43 level in the input and shown as percentage to the WT hnRNP A1 group. Means ± SEM, n = 5 (B) or 3 (D); *p < 0.05, **p < 0.01, ***p < 0.001; One-way ANOVA.

Figure 7. PARP inhibition mitigates the cytotoxicity of hnRNP A1 and TDP-43 in MN-like

NSC-34 cells.

(**A**) NSC-34 cells infected with increased concentrations of lenti-hnRNP A1-HA show decreased cell viability in the Cell Counting Kit-8 (CCK-8) assay, indicating that hnRNP A1-induced cytotoxicity is dose-dependent. (**B**) Representative bright field images of NSC-34 cells infected with Lenti-Ctrl or Lenti-hnRNP A1-HA (1% virus concentration) are shown. Scale bar: 50 μm. (**C**) WT, K298A and PBM^{mut} of hnRNP A1 exhibit different levels of cytotoxicity in NSC-34 cells. (**D-E**) NSC-34 cells overexpressing hnRNP A1 (C) or TDP-43 (D) by lentivirus infection show decreased cell viability in the CCK-8 assay, which is significantly suppressed by KD of PARP1 by (si-PARP1) compared to the scrambled siRNA control (si-Ctrl). (**F**) Western blot analysis confirms PARP1 KD and decrease of overall PARylation levels by siRNA of PARP1 (si-PARP1). (**G**) PARP inhibitor Olaparib rescues TDP-43 OE-induced reduction of cell viability in the NSC-34 model. Mean ± SEM; n = 3; *$p < 0.05$, **$p < 0.01$; ns, not significant; Student's t-test.

Figure 8. Downregulation of *Parp* alleviates neurodegeneration in a *Drosophila* model of ALS.

(**A-B**) Representative images (A) and quantification (B) of degeneration scores of the fly eyes expressing hTDP-43 with RNAi-*Parp* or RNAi-*mCherry* (Ctrl). (**C-D**) RNAi KD of *Parp* in adult fly neurons using an *elav*GS driver alleviates TDP-43-mediated climbing decline (C) and shortened lifespan (D). The climbing capability is evaluated as the average percentage of flies that climb over 3 cm within 10 seconds. The median lifespan and the number of flies tested for each genotype are indicated. Mean ± SEM; **$p < 0.01$, ***$p < 0.001$; ns, not significant; Student's t-test. Scale bar, 100 μm. (**E**) A schematic model of the role of PARylation in regulating RNP granules. PARylation regulates the SG translocation, phase separation, and neurotoxicity of disease-related RNPs such as hnRNP A1 and TDP-43 (see Discussion). PAR, poly(ADP-ribose); PBM, PAR-binding motif; PPI, protein-protein interaction; WT, wild-type hnRNP A1; K298A, PARylation site mutant of hnRNP A1; PBM^{mut}, PAR-binding motif mutant of hnRNP A1; SG, stress granule; cyto. foci, cytoplasmic PBM^{mut} foci.

Figure 2_YDuan

Figure 4_YDuan

A

hnRNP A1

10 μM   20 μM   40 μM   60 μM

NaCl

25 mM

50 mM

150 mM

300 mM

Large droplets

LLPS

No droplet

B   20 μM hnRNP A1 + 25 mM NaCl

0 μM PAR   1 μM PAR   3 μM PAR   5 μM PAR   7 μM PAR

C   20 μM hnRNP A1 + 25 mM NaCl

0 μM Heparin   7 μM Heparin

D   25 mM NaCl

0 μM PAR   7 μM PAR

100 mM NaCl

50 µM hnRNP A1

50 µM BSA
+
50 µM hnRNP A1

50 µM TDP-43¹⁻²⁷⁴
+
50 µM hnRNP A1

50 µM TDP-43²⁷⁴⁻⁴¹⁴
+
50 µM hnRNP A1

20 µM TDP-43²⁷⁴⁻⁴¹⁴
+
50 µM hnRNP A1

A B C D E

50 µM BSA

50 µM TDP-43¹⁻²⁷⁴

50 µM TDP-43²⁷⁴⁻⁴¹⁴

20 µM TDP-43²⁷⁴⁻⁴¹⁴

B' C' D' E'

Figure 6_YDuan

Figure 7_YDuan

Figure 8_YDuan

A GMR>*hTDP-43* +

RNAi-Ctrl   RNAi-*Parp*

Day 5

Day 20

B Eye deg. score

GMR>*hTDP-43* +
■ RNAi-Ctrl (27)
■ RNAi-*Parp* (21)

ns   ***
Day 5   Day 20

C % of flies that climb over 3 cm in 10s

*elav*GS>*hTDP-43* +
■ RNAi-Ctrl (100)
■ RNAi-*Parp* (180)

***   **
Day 3   Day 20   Day 30

D % of surviving flies

*elav*GS>Ctrl (168)
(54.3 ± 1.0 d)

*elav*GS>*hTDP-43* + RNAi-Ctrl (137)
(45.0 ± 0.9 d)

*elav*GS>*hTDP-43* + RNAi-*Parp* (225)
(53.1 ± 1.0 d)

***
***

Day: 0   20   40   60

E

WT   PBM^{mut}

hnRNP A1   TDP-43   ?

SG

hnRNP A1

WT
K298A
PBM^{mut}

Nucleus

PBM   PAR   RNA   PPI

SG
More dynamic
Less toxic

si-PARG
si-PARP
Olaparib

SG
Less dynamic
More toxic

cyto.
foci