Article

Zebrafish Otolith Biomineralization Requires Polyketide Synthase

Kevin D. Thiessen [1], Steven J. Grzegorski [2], Lisa Higuchi [1], Jordan A. Shavit [2], and Kenneth L. Kramer [1],*

[1] Department of Biomedical Sciences, Creighton University School of Medicine, Omaha, NE, United States
[2] Department of Pediatrics, University of Michigan, Ann Arbor, MI, United States
* Correspondence: kenkramer@creighton.edu; Tel.: 402-280-2763

Abstract: Deflecting biomineralized crystals attached to vestibular hair cells are necessary for maintaining balance. Zebrafish (*Danio rerio*) are useful organisms to study these biomineralized crystals called otoliths, as many required genes are homologous to human otoconial development. We sought to identify and characterize the causative gene in a trio of mutants, *no content* (*nco*) and *corkscrew* (*csr*), and *vanished* (*vns*), which fail to develop otoliths during early ear development. We show that *nco*, *csr*, and *vns* have potentially deleterious mutations in polyketide synthase (*pks1*), a multi-modular protein that has been previously implicated in biomineralization events in chordates and echinoderms. We found that Otoconin-90 (Oc90) expression within the otocyst is normal in *nco* and *csr*; therefore, it is not sufficient for otolith biomineralization in zebrafish. Similarly, normal localization of Otogelin, a protein required for otolith tethering in the otolithic membrane, is not sufficient for Oc90 attachment. Furthermore, eNOS signaling and Endothelin-1 signaling were the most up- and down-regulated pathways during otolith agenesis in *nco*, respectively. Our results demonstrate distinct processes for otolith nucleation and biomineralization in vertebrates and will be a starting point for models that are independent of Oc90-mediated seeding. This study will serve as a basis for investigating the role of eNOS signaling and Endothelin-1 signaling during otolith formation.

Keywords: inner ear, otolith, biomineralization, calcium carbonate, polyketide synthase, zebrafish, endothelin-1, eNOS

1. Introduction

Otoconia and otoliths act as a mass load that increase the sensitivity of mechanosensory hair cells to the effects of gravity and linear acceleration in mammals and fish, respectively. While the morphology of otoconia ("ear particles") and otoliths ("ear stones") differ, the initial formation of bio-crystals rely on many homologous proteins [1]. Zebrafish otoliths are primarily composed of calcium carbonate in the form of aragonite, which accounts for ~99% of the total otolithic mass [2, 3]. The center of the otolith contains a proteinaceous core that acts as a site for otolith nucleation and biomineralization. This matrix lays the foundation for further otolith growth, which is mediated by daily deposition of additional otoconins and calcium carbonate molecules [2].

In zebrafish, otolith nucleation occurs when the otolith precursor particles or OPPs bind to the tips of the immotile kinocilia of tether cells within the otic vesicle [4, 5]. Subsequent studies have demonstrated that the critical period of otolith seeding and nucleation starts at 18-18.5 hpf and ceases by 24 hpf. [1, 4, 6-8]. In mammalian inner ear development, Otoconin-90 (Oc90; the major protein component of otoconia) is necessary for otoconial seeding and nucleation [9-11]. Oc90 can bind Otolin-1 to establish a protein-rich matrix that serves as a scaffold for subsequent deposition of

calcium carbonate [12, 13]. While it is not the major protein component in zebrafish otoliths, Oc90
plays an important role in otolith seeding and early development as *oc90*-morphants do not develop
otoliths [1, 14, 15]. While additional gene mutations have been identified that lead to otolith agenesis
[16-20], the genes responsible for several zebrafish otolith mutants have been undetermined.

In this study, we sought to identify and characterize the causative gene in a trio of zebrafish
mutants, *no content* (*nco*) and *corkscrew* (*csr*), and *vanished* (*vns*), which fail to develop otoliths during
early inner ear development. We provide genetic evidence that the causative gene is polyketide
synthase (*pks1*; currently *wu:fc01d11*), a candidate gene that was recently identified as a key factor of
otolith biomineralization in Japanese medaka (*Oryzias latipes*) [21].

2. Materials and Methods

All zebrafish were maintained in a temperature-controlled (28.5ºC) and light-controlled (14h
on/10h off) room per standardized conditions. *nco* strain (jj149) was generated by an ENU screen and
obtained from ZIRC (Eugene, OR, USA). *csr* was a spontaneous mutant in a *bre*-KO2/*ntl*-GFP line. *vns*
was a spontaneous mutant in a AB/TL background. All protocols were approved by Creighton
University and the University of Michigan Animal Care and Use Committees.

Mutant *nco* embryos and wild-type (WT) clutchmates were collected during the critical period
of otolith nucleation and seeding (24 hours post fertilization, hpf) and submitted for RNA sequencing.
Analysis was completed using MMAPPR (Mutation Mapping Analysis Pipeline for Pooled RNA-seq)
[22]. Whole genome sequencing of *csr* was performed and analyzed using MegaMapper [23]. All
sequencing was conducted at the University of Nebraska Medical Center Genomics Core Facility.
Accession numbers for *nco* RNA-seq and *csr* genome sequencing will be provided during review.

WT mRNA and *pks1*^{L905P} were synthesized using mMessage Machine from a clone provided by
Dr. Hiroyuki Takeda (University of Tokyo), cleaned on an RNeasy column, and subsequently
injected into single-cell *csr* and *nco* embryos. Naked plasmid was injected into *vns* embryos and
overall penetrance of otolith formation was determined. Site-directed mutagenesis was used to
generate the mutant clone containing the causative mutation in *csr* (*pks1*^{L905P} in Japanese medaka;
pks1^{A911P} in zebrafish). Primers used for site-directed mutagenesis were:

pks1_L905P_Forward: 5'-GATATGGCGTGATGTCCGGTGACAGGTTGAAGATC-3'
pks1_L905P_Reverse: 5'-ATCTTCAACCTGTCACCGGACATCACGCCATATC-3'

Pathway analysis of *nco* was performed using Ingenuity Pathway Analysis (IPA,
http://www.ingenuity.com) [24]. The Ensembl Gene IDs for differentially expressed genes were
uploaded to IPA. Cut-off for gene expression analysis was set at 0.75 RPKM. The calculated z-score
indicates a pathway with genes exhibiting increased mRNA levels (positive) or decreased mRNA
levels (negative). No change in mRNA levels results in a z-score of zero.

csr, *nco*, and *vns* samples were PCR-amplified and submitted for Sanger sequencing using
the following primers:

nco_Forward: 5'-GGGAGGATGCTTGTTGTTGG-3'
nco_Reverse: 5'-GTGGCCCAGAATAGGATCCA-3'
csr_Forward: 5'-AAGACGGGGACATGACTCAG-3'
csr_Reverse: 5'-TTCAACAAACAGTGCTCCGG-3'
vns_Forward: 5'-GCCATCATTGGAATTGGATG-3'
vns_Reverse: 5-GGTGTTCCAGTCCCATGAGC-3'

csr and *nco* embryos were collected during key stages in ear development, fixed with hydrogel
and washed in CHAPS-based CLARITY-clearing solution [25]. Embryos were decalcified with EDTA
before blocking, incubating in primary and secondary antibodies diluted in PBS-Triton (0.1%), and
imaging by confocal microscopy. Affinity-purified rabbit polyclonal antibodies were generated to
Otogelin (CGNRVDGPSASKG; 1:1000) or Oc90 (CNTQSDTVDRKPTQSKPQ; 1:1000) by
conventional methods and directly labelled before immunofluorescence. Other antibodies used were
Keratan Sulfate (MZ15; 1:2000; DSHB), Hair Cell Specific-1 (HCS-1; 1:500; DSHB), and acetylated-
tubulin (1:500; Sigma T6793). Phalloidin (ThermoFisher A12379) was used at a concentration of 1:500.
Mitotracker Red (ThermoFisher #M22425) was resuspended in DMSO (0.25 mM) and diluted to
200 nM in E3 embryo medium. Embryos were then incubated for 20 minutes before removing
Mitotracker solution and replacing with fresh E3 embryo medium. Samples were allowed to stabilize
for 30 minutes before imaging at 21 hpf. Embryos were phenotyped at 27 hpf.
To test the effects of exogenous ions on otolith formation, embryos were kept in E3 Medium
until early gastrulation. Embryos were washed, dechorionated, and transferred to 1X Basic Solution
(58 mM NaCl, 0.4 mM $MgSO_4$ and 5 mM HEPES) supplemented with 0.7 mM potassium chloride, 0.6
mM calcium nitrate or 0.6 mM calcium chloride. Embryos were then transferred to fresh 1X Basic
Solution with respective supplement for the remaining development. Embryos were scored by the
presence or absence of otoliths at 27 hpf.
Statistical significance was calculated using Fisher's Exact Test, G-test for Independence, and
Chi-Squared Distribution.

3. Results

3.1 csr and nco are genetically-linked

The most apparent phenotype of *csr* and *nco* mutants is that they fail to form otoliths or any
observable complex calcium deposits within the inner ear (Fig. 1A-C). Furthermore, the mutant
larvae are homozygous lethal as the swim bladder fails to inflate (Fig. 1A'-C') and they are unable to
feed. While it is still unknown why the swim bladder fails to inflate when otoliths are absent, it is a
common phenotype in other mutants with otolith agenesis [14, 16-18, 22]. Due to this commonality
within *csr* and *nco content*, we sought to determine if these phenotypes would complement each other.
The results of the complementation test showed that some offspring failed to develop otoliths (n =
31/106), supporting that *nco* and *csr* likely are allelic.

3.2 Exogenous ions influence otolith nucleation in csr embryos; not nco or vns embryos

As an aquatic species, the environment of zebrafish can be easily controlled and adapted to
assess its impact on embryonic development. Previously, small molecules have been used to block
otolith development by inhibiting otolith nucleation [6]. We hypothesized that there was an error in
ion homeostasis that could be affected by exogenous solutions. In water treatments supplemented
with potassium chloride (n=59), we found a significant increase in *csr* embryos lacking otoliths
(35.60%) compared to treatments supplemented with calcium chloride (10.94%; n=64) or calcium
nitrate (20.17%; n=119)(G-test; p=0.004). Additionally, we observed no significant change in *nco*
mutant embryos for water treatments supplemented with potassium chloride (17.76%; n=107),
calcium chloride (16.67%; n=120) or calcium nitrate (16.9%; n=112)(G-test; p=0.975). Similarly, the
penetrance of otolith formation in *vns* was not affected by exogenous salts (data not shown).

Furthermore, Mitotracker was used to mark mitochondria-rich cells in *csr* and *nco* embryos. While

nco embryos appear normal, we observed that *csr* embryos show a lack of Mitotracker localization at

21 hpf (Fig. S1). Altogether, this suggests the nature of each mutation, while likely allelic, are

inherently different.

3.3 Potentially deleterious mutations identified in polyketide synthase for csr, nco, and vns

To positionally clone the gene responsible for *nco* and *csr*, we used complementary approaches

for each strain. MMAPPR analysis of *nco*-derived RNA sequencing (Fig. 2A) [22] and MegaMapper

analysis of *csr*-derived whole genome sequencing (Fig. 2B) [23] both identified a genomic region with

high homology surrounding the *pks1* locus. While several other genes were in that region, a previous

study on otolith biomineralization in Japanese medaka made *pks1* the likely gene candidate [21].

Potentially deleterious mutations were identified in *pks1* for *csr* (A911P) and *nco* (L681*), which were

both located within a conserved acyl transferase domain (Fig. 2C). In regards to *csr* mutation,

homozygosity at the locus is compatible with proper development in the AB/TL background and

with otolith agenesis in the AB background (data not shown). Furthermore, a deleterious mutation

in *vns* (G239R) was serendipitously found to be linked to a neighboring gene during a separate study.

The deleterious point mutation was identified by Sanger sequencing of the *pks1* locus and confirmed

by relatively high penetrance of otolith agenesis (95%).

3.4 Japanese medaka pks1 mRNA rescues otolith biomineralization in csr, nco, and vns

While the last common ancestor of Japanese medaka and zebrafish was estimated to be 150

million years ago [26], we sought to assess if the function of *pks1* within the inner ear is conserved.

We injected Japanese medaka *pks1* mRNA or DNA into single-cell embryos of *csr*, *nco*, and *vns*

heterozygous incrosses. Microinjection of Japanese medaka *pks1* mRNA (300 ng/μL) rescued otolith

biomineralization in both *csr* (p<0.0001; χ^2<0.0001; n=93) and *nco* (p=0.0032; χ^2=0.0022; n=84) mutants

(Fig. 3B). Additionally, microinjection of the Japanese medaka *pks1* plasmid (20 ng/uL) provided by

Dr. Takeda rescued otolith biomineralization in *vns* (p<0.0001; χ^2=0.0004; n=39). Using site-directed

mutagenesis, we introduced the non-synonymous mutation (A911P) in *csr* to the Japanese medaka

mRNA construct (L905P). We repeated injections into single-cell embryos and failed to rescue otolith

biomineralization in *csr* and *nco*. WT medaka *pks1*, but not *pks1*^{L905P}, rescued otolith biomineralization

in *csr* and *nco* embryos (Fig. 3C).

3.5 Ingenuity pathway analysis of nco embryos

While *pks1* is thought to produce an otolith nucleation factor [21], its broader role during inner

ear development is unknown. Ingenuity Pathway Analysis of *nco* at 24 hpf identified eNOS and

Endothelin-1 signaling as the top up- and down-regulated pathways, respectively (Fig. 4A). Among

the down regulated genes was *rdh12l*, a gene adjacent to *pks1*, suggesting that there is local control of

transcription at that locus. *mir-92a*, the top down-regulated gene, has a predicted binding site in the

3′UTR of *rdh12l* (Fig. S2) [27]. In addition, several genes listed in the top ten up- or down-regulated

lists are also enriched in adult mechanosensory hair cells such as *il11b*, *fosab*, *fosb*, *fosl1a*, *socs3a*, *scg5*,

and *dnaaf3* (Figs. 4B-C) [28]. Of these genes, *il11b* is up-regulated during neuromast hair cell

regeneration [29]. Notably, *dnaaf3* causes primary ciliary dyskinesia and morpholino knockdown of

dnaaf3 causes abnormal otolith growth [30]. While its role in inner ear development is unknown, *scg5*

is expressed within the anterior and posterior poles of the otic placode during the critical period of
otolith nucleation [31].

3.6   Aberrant expression of proteins involved in otolith development in csr and nco

In mammalian inner ear development, Oc90 is necessary for otoconial seeding and nucleation
[9, 10]. Similarly, the role of Oc90 is evolutionarily conserved in zebrafish and has been previously
thought to be sufficient for otolith nucleation [14]. Using immunofluorescence (IF), we saw diffuse
expression of Oc90 in *csr* and *nco* otocysts (Figs. 5B-D), which demonstrated that Oc90 expression
within the otocyst is not sufficient for otolith biomineralization in zebrafish. Similarly, normal
localization of Otogelin (Otog), a protein required for otolith tethering in the otolithic membrane is
not sufficient for Oc90 attachment. Additionally, other otoconins that are important for calcium
deposition and growth were detected with diffuse expression within the otocyst such as Starmaker
and Keratan Sulfate (data not shown) [32, 33].

3.7 Polyketide synthase expression is enriched in adult mechanosensory hair cells

Otolith nucleation is thought to be mediated by a tether-cell specific otolith precursor binding
factor (OPBF), which lays the foundation for the successive biomineralization of the otolith [5, 7, 34].
The presence of an OPBF was proposed almost two decades ago and its identification proves to be
elusive [34]. Recent studies suggest that one or more OPBFs are expressed by tether-cells and help to
mediate otolith nucleation by binding other OPPs [5, 7, 35].
We sought to assess if *pks1* or its enzymatic product is a tether-cell OPBF. First, we demonstrated
that the total number of hair cells remain unchanged during early development in *nco*, suggesting
there are no differences in tether cell maturation and maintenance (Figs. 5E-G). Then, using publicly
available RNA-seq data, *pks1* mRNA can be detected during the critical period of otolith nucleation
[36]. Previous data has shown its localization in the otic vesicle at 19 hpf [21], supporting its role as
an OPBF. *pks1* is enriched (7.46-fold increase) in mechanosensory hair cells compared to support cells
within the adult zebrafish inner ear (Table S1). Additionally, RNA-seq data suggests *pks1* appears to
be differentially expressed in support cells. Support cells predominantly express a 300bp region of
the 5′UTR of the *pks1* transcript [28]. A search for transcriptional regulatory motifs in the 5′UTR of
pks1 found a predicted binding site for TCF-3 [37], a transcription factor highly expressed in adult
mechanosensory hair cells [28]. While the role of TCF-3 in the inner ear is unknown, it is expressed
within the otic vesicle during the critical period of otolith nucleation [31].

4. Discussion

The mutants *csr*, *nco*, and *vns* were chosen for this study because each lack the necessary factors
such as an OPBF for otolith seeding and biomineralization. To determine the genes responsible for
otolith agenesis in these mutants, we used two complementary approaches. The first approach was
Whole Genome Sequencing of the *csr* mutant genome to identify regions of high homology. This
indeed was difficult as the *csr* background strain was heavily inbred, resulting in multiple peaks of
high homology. Since we demonstrated *csr* and *nco* are genetically-linked, we sought to further clarify
the responsible locus using a second method (i.e. RNA-seq of the *nco* transcriptome) for comparison.
This result pinpointed a region of high homology near the end of the 24th chromosome. While
deciphering potentially deleterious mutations within that region, we focused on *pks1* following

evidence that it is responsible for otolith nucleation in Japanese medaka [21]. While these species are
evolutionarily divergent, the shared phenotype between medaka and our mutants suggested that the
role of *pks1* is conserved. As a result, we chose to use medaka *pks1* nucleic acid to rescue otolith
formation in *csr, nco* and *vns* mutants. Similarities can also be drawn with other zebrafish mutants
such as *keinstein,* which has diffuse expression of Starmaker within the otocyst and exhibits similar
circling swimming behaviors [38, 39]. Furthermore, *keinstein* may be another *pks1* allele due to its
predicted chromosomal location [40].

While WT medaka *pks1* rescues otolith biomineralization in *csr* and *nco*, differences in penetrance
of exogenous ions on otolith formation suggested the nature of each mutation is fundamentally
different. This was confirmed by Sanger sequencing that *nco* has a premature stop codon while *csr*
likely makes a defective protein that may be stabilized by exogenous ions. This defective protein may
be the explanation for the differences in Mitotracker localization in *csr*. Due to its surface stain
expression, we hypothesize that Mitotracker was localized to mitochondria-rich ionocytes [41].
Ionocytes have previously been implicated in otolith formation as mutations in *gcm2*, which is
responsible for ionocyte maturation, leads to otolith agenesis [42]. We hypothesize that the
endolymph in *csr* and *nco* mutants has the necessary components for otolith nucleation [2] but lack a
trigger produced by *pks1*. Additionally, the absence of *pks1* does not visibly appear to affect hair cell
development that are required for otolith nucleation [5]. IF of *csr* and *nco* embryos demonstrated that
expression of a critical otoconial seeding protein, Oc90, within the otocyst is not sufficient for otolith
biomineralization in the presence of the otolithic membrane.

One caveat is that the penetrance of otolith formation is influenced by the genetic background
of zebrafish. When treated with the small molecule 31N3, WT embryos in the AB/EKW background
fail to develop otoliths [6]. However, 31N3 fails to inhibit otolith formation in the TL and TU strains,
suggesting that there are potential genetic modifiers that influence otolith nucleation in these
backgrounds. In the case of *csr*, homozygosity at the locus is compatible with otolith agenesis in the
AB background and, with proper development, in the TL background. This suggests *csr* may be a
hypomorphic allele and the AB background can overcome the loss of Pks1 function with enhanced
ion flux. Ironically, the mutant phenotype was lost when *csr* was outcrossed to the WIK background.
It was only until *csr* was backcrossed to the AB background that the mutants were recovered.
Altogether, we suggest that the AB background heavily influences the penetrance of otolith
formation.

While *pks1* likely acts as an enzyme whose expression is enriched in adult mechanosensory hair
cells [28], its product acts as an OPBF and is required for otolith nucleation in zebrafish. However,
the molecular function of *pks1* remains unknown. Using *nco* RNA-seq data, we performed an
Ingenuity Pathway Analysis, which identified eNOS and Endothelin-1 signaling as the most up- and
down-regulated pathways, respectively. eNOS signaling could be impacted by *pks1* metabolites such
as iromycin, which has been shown to inhibit this pathway [43]. Both eNOS and Endothelin-1 have
been implicated in inner ear development and function. Notably, it has been demonstrated that these
pathways are inversely related in sensorineural hearing loss [44]. An example of this is Waardenburg
syndrome, caused by mutations in endothelins, which cause abnormal pigmentation and
sensorineural hearing loss [45]. During early development, Endothelin-1 mRNA turns on during the
critical period of otolith nucleation [31, 36] and is detected in the otic vesicle at 24 hpf [46]. Endothelin-
1 and its receptor (*ednraa*) are both enriched in adult zebrafish inner ear support cells [28].

Additionally, Endothelin-1 has been implicated with the FOS-family of genes (*fosab*, *fosb*, and *fosl1a*) and *socs3a*, which are all differentially expressed in *nco* at 24 hpf. They are all part of a regulatory network during hypergravity-mediated bone formation [47], which might suggest a common mechanism between bone mineralization and otolith biomineralization. Future studies will attempt to clarify the roles of Endothelin-1 and eNOS signaling pathways during otolith biomineralization.

Author Contributions: Conceptualization, K.T. and K.K.; Methodology, K.T, S.G. and K.K..; Validation, K.T., S.G., and L.H.; Formal Analysis, K.T. and S.G.; Investigation, K.T., S.G., and L.H.; Resources, J.S. and K.K; Data Curation, J.S. and K.K.; Writing-Original Draft Preparation, K.T.; Writing-Review & Editing, K.T., S.G., J.S., and K.K; Visualization, K.T. and K.K.; Supervision, J.S. and K.K.; Project Administration, J.S. and K.K.; Funding Acquisition, J.S. and K.K.

Funding: The Kramer lab are grateful for funding through grants from the State of Nebraska (LB-692), the National Center for Research Resources (5P20RR018788-09), and the National Institute of General Medical Sciences (8 P20 GM103471-09). The Shavit lab acknowledges support from National Heart, Lung, and Blood Institute grants (R01HL124232 and HL125774).

Acknowledgments: We recognize the University of Nebraska Medical Center Genomics Core Facility for assistance with sequencing and bioinformatics. We thank Dr. Hiroyuki Takeda from the University of Tokyo for supplying the Japanese medaka *pks1* mRNA construct. We acknowledge Creighton University Integrated Biomedical Imaging Facility for assistance with confocal microscopy. Finally, we express gratitude to the members of the Kramer Lab at Creighton University for their support with zebrafish husbandry.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Lundberg, Y.W., et al., *Mechanisms of otoconia and otolith development.* Dev Dyn, 2015. **244**(3): p. 239-53.

2. Payan, P., et al., *Endolymph chemistry and otolith growth in fish.* Comptes Rendus Palevol, 2004. **3**(6-7): p. 535-547.

3. Borelli, G., et al., *Biochemical relationships between endolymph and otolith matrix in the trout (Oncorhynchus mykiss) and turbot (Psetta maxima).* Calcif Tissue Int, 2001. **69**(6): p. 356-64.

4. Riley, B.B., et al., *A critical period of ear development controlled by distinct populations of ciliated cells in the zebrafish.* Dev Biol, 1997. **191**(2): p. 191-201.

5. Stooke-Vaughan, G.A., et al., *The role of hair cells, cilia and ciliary motility in otolith formation in the zebrafish otic vesicle.* Development, 2012. **139**(10): p. 1777-87.

6. Peterson, R.T., et al., *Small molecule developmental screens reveal the logic and timing of vertebrate development.* Proc Natl Acad Sci U S A, 2000. **97**(24): p. 12965-9.

7. Stooke-Vaughan, G.A., et al., *Otolith tethering in the zebrafish otic vesicle requires Otogelin and alpha-Tectorin.* Development, 2015. **142**(6): p. 1137-45.

8. Riley, B.B., *Genes Controlling the Development of the Zebrafish Inner Ear and Hair Cells*, in *Current Topics in Developmental Biology*. 2003, Academic Press. p. 357-388.

9. Zhao, X., et al., *Otoconin-90 deletion leads to imbalance but normal hearing: a comparison with other otoconia mutants.* Neuroscience, 2008. **153**(1): p. 289-99.

10. Zhao, X., et al., *Gene targeting reveals the role of Oc90 as the essential organizer of the otoconial organic matrix.* Dev Biol, 2007. **304**(2): p. 508-24.

11. Wang, Y., et al., *Otoconin-90, the mammalian otoconial matrix protein, contains two domains of*
homology to secretory phospholipase A2. Proc Natl Acad Sci U S A, 1998. **95**(26): p. 15345-50.

12. Deans, M.R., J.M. Peterson, and G.W. Wong, *Mammalian Otolin: a multimeric glycoprotein specific*
to the inner ear that interacts with otoconial matrix protein Otoconin-90 and Cerebellin-1. PLoS
One, 2010. **5**(9): p. e12765.

13. Moreland, K.T., et al., *In vitro calcite crystal morphology is modulated by otoconial proteins otolin-*
1 and otoconin-90. PLoS One, 2014. **9**(4): p. e95333.

14. Petko, J.A., et al., *Otoc1: a novel otoconin-90 ortholog required for otolith mineralization in*
zebrafish. Dev Neurobiol, 2008. **68**(2): p. 209-22.

15. Murayama, E., et al., *Otolith matrix proteins OMP-1 and Otolin-1 are necessary for normal otolith*
growth and their correct anchoring onto the sensory maculae. Mech Dev, 2005. **122**(6): p. 791-803.

16. Hughes, I., et al., *Otopetrin 1 is required for otolith formation in the zebrafish Danio rerio.* Dev Biol,
2004. **276**(2): p. 391-402.

17. Stawicki, T.M., et al., *The zebrafish merovingian mutant reveals a role for pH regulation in hair cell*
toxicity and function. Dis Model Mech, 2014. **7**(7): p. 847-56.

18. Sumanas, S., J.D. Larson, and M. Miller Bever, *Zebrafish chaperone protein GP96 is required for*
otolith formation during ear development. Dev Biol, 2003. **261**(2): p. 443-55.

19. Kiss, P.J., et al., *Inactivation of NADPH oxidase organizer 1 results in severe imbalance.* Curr Biol,
2006. **16**(2): p. 208-13.

20. Colantonio, J.R., et al., *The dynein regulatory complex is required for ciliary motility and otolith*
biogenesis in the inner ear. Nature, 2008. **457**: p. 205.

21. Hojo, M., et al., *Unexpected link between polyketide synthase and calcium carbonate*
biomineralization. Zoological Lett, 2015. **1**(1): p. 3.

22. Hill, J.T., et al., *MMAPPR: Mutation Mapping Analysis Pipeline for Pooled RNA-seq.* Genome
Research, 2013. **23**(4): p. 687-697.

23. Obholzer, N., et al., *Rapid positional cloning of zebrafish mutations by linkage and homozygosity*
mapping using whole-genome sequencing. Development, 2012. **139**(22): p. 4280-90.

24. Kramer, A., et al., *Causal analysis approaches in Ingenuity Pathway Analysis.* Bioinformatics, 2014.
30(4): p. 523-30.

25. Chung, K., et al., *Structural and molecular interrogation of intact biological systems.* Nature, 2013.
497(7449): p. 332-7.

26. Kirchmaier, S., et al., *The Genomic and Genetic Toolbox of the Teleost Medaka (Oryzias latipes).*
Genetics, 2015. **199**(4): p. 905-918.

27. Ulitsky, I., et al., *Extensive alternative polyadenylation during zebrafish development.* Genome Res,
2012. **22**(10): p. 2054-66.

28. Barta, C.L., et al., *RNA-seq transcriptomic analysis of adult zebrafish inner ear hair cells.* Sci Data,
2018. **5**: p. 180005.

29. Jiang, L., et al., *Gene-expression analysis of hair cell regeneration in the zebrafish lateral line.*
Proceedings of the National Academy of Sciences, 2014. **111**(14): p. E1383-E1392.

30. Mitchison, H.M., et al., *Mutations in axonemal dynein assembly factor DNAAF3 cause primary*
ciliary dyskinesia. Nature Genetics, 2012. **44**: p. 381.

31. Thisse, B., et al., *Expression of the zebrafish genome during embryogenesis.* 2001: ZFIN Direct Data
Submission.

32. Yang, H., et al., *Matrix recruitment and calcium sequestration for spatial specific otoconia*
development. PLoS One, 2011. **6**(5): p. e20498.

33. Sollner, C., et al., *Control of crystal size and lattice formation by starmaker in otolith*
biomineralization. Science, 2003. **302**(5643): p. 282-6.

34. Riley, B.B. and D.J. Grunwald, *A mutation in zebrafish affecting a localized cellular function*
required for normal ear development. Dev Biol, 1996. **179**(2): p. 427-35.

35. Yu, X., et al., *Cilia-driven fluid flow as an epigenetic cue for otolith biomineralization on sensory*
hair cells of the inner ear. Development, 2011. **138**(3): p. 487-94.

36. White, R.J., et al., *A high-resolution mRNA expression time course of embryonic development in*
zebrafish. Elife, 2017. **6**.

37. Chang, T.H., et al., *An enhanced computational platform for investigating the roles of regulatory*
RNA and for identifying functional RNA motifs. BMC Bioinformatics, 2013. **14 Suppl 2**: p. S4.

38. Sollner, C., et al., *Mutated otopetrin 1 affects the genesis of otoliths and the localization of*
Starmaker in zebrafish. Dev Genes Evol, 2004. **214**(12): p. 582-90.

39. Whitfield, T.T., et al., *Mutations affecting development of the zebrafish inner ear and lateral line.*
Development, 1996. **123**: p. 241-54.

40. Geisler, R., et al., *Large-scale mapping of mutations affecting zebrafish development.* BMC
Genomics, 2007. **8**: p. 11.

41. Esaki, M., et al., *Mechanism of development of ionocytes rich in vacuolar-type H(+)-ATPase in the*
skin of zebrafish larvae. Developmental biology, 2009. **329**(1): p. 116-129.

42. Kumai, Y., R.W.M. Kwong, and S.F. Perry, *A role for transcription factor glial cell missing 2 in*
Ca2+ homeostasis in zebrafish, Danio rerio. Pflügers Archiv - European Journal of Physiology,
2015. **467**(4): p. 753-765.

43. Surup, F., et al., *The iromycins, a new family of pyridone metabolites from Streptomyces sp. I.*
Structure, NOS inhibitory activity, and biosynthesis. J Org Chem, 2007. **72**(14): p. 5085-90.

44. Liu, Q., et al., *[The study on plasma ET and NO of patients with sudden hearing loss].* Lin Chuang
Er Bi Yan Hou Ke Za Zhi, 2003. **17**(11): p. 668-9.

45. Pingault, V., et al., *Review and update of mutations causing Waardenburg syndrome.* Hum Mutat,
2010. **31**(4): p. 391-406.

46. Miller, C.T., et al., *sucker encodes a zebrafish Endothelin-1 required for ventral pharyngeal arch*
development. Development, 2000. **127**(17): p. 3815-3828.

47. Aceto, J., et al., *Zebrafish Bone and General Physiology Are Differently Affected by Hormones or*
Changes in Gravity. PLoS ONE, 2015. **10**(6): p. e0126928.

372   **Figure 1:** (**A-D**) The *csr*, *nco,* and *vns* mutant phenotypes fail to form otoliths within the inner ear. However,

373   semicircular canal formation appears to be normal. (**A'-D'**) All mutants fail to inflate their swim bladders, which

374   is lethal. Imaged at 5 days post fertilization (dpf). Magnification 6.3X. (*) indicates swim bladder.

Figure 2: Complementary approaches for causative gene discovery. MMAPPR analysis of RNA sequencing data for *nco* (**A**) and whole genome homology mapping for *csr* (**B**) identified regions of high homology on the 24th chormosome near the *pks1* locus (~33 Gb). (**C**) Deleterious mutations were identified in *pks1* for *nco* and *csr* within the acyl transferase (AT) domain and *vns* within the polyketide synthase (PKS) domain. Sanger sequencing confirmed SNPs in *csr*, *nco*, and *vns* mutants. Other domains include Ketoacyl Synthetase (KS), Medium Chain Reductase (MDR), NAD(P)-dependent dehydrogenase (NDD), and Phosphopanthetheine-Binding (PP).

384 **3:** WT *pks1* nucleic acid rescues otolith formation in *csr*, *nco*, and *vns*. (**A**) Normal frequencies of mutant phenotypes in each uninjected strain. All four pairings follow homozygous

385 recessive mode of inheritance. (**B**) Results of injected embryos show that Japanese medaka *pks1* mRNA (300 pg) rescues both *csr* and *nco* mutants and *pks1* DNA (20 pg) rescues *vns*

386 mutants. (*, $p < 0.0001$, paired *t*-test)(**, $p < 0.0032$, paired *t*-test)(***, $p = 0.0001$, paired *t*-test),. Site-directed mutagenesis was used to introduce a conserved mutation in *csr* (A911P)

387 into the Japanese medaka construct (L905P) (**C**) Injection of pks1[L905P] (300 pg) fails to rescue *csr* or *nco* mutant phenotypes.

388
389 **Figure 4:** Gene expression and pathway analysis of *nco* embryos. (**A**) Ingenuity Pathway Analysis shows the top up-regulated and down-regulated pathways, which are eNOS
390 Signaling and Endothelin-1 Signaling, respectively. Positive z-score indicated increased mRNA levels. Negative z-score indicates decreased mRNA levels. No change in mRNA levels
391 results in a z-score of zero. (**B**) Differential gene expression in the top up-regulated genes. (**C**) Differential gene expression in the top down-regulated genes. (**, expressed in adult
392 zebrafish mechanosensory hair cells) [28].
393

394
395 **Figure 5:** Aberrant expression of proteins invovled in otolith development in *csr* and *nco*. (**A**) Schematic of otic vesicle at 27 hpf. Anterior to right. (**B**) In WT, Oc90 is expressed
396 within the mineralized otolith, which is situated atop the otolithic membrane (Otog), at 27 hpf. Scale bar = 5 μm. (**C-D**) Oc90 has diffuse expression within the otocyst of *csr* and *nco*.
397 In *csr* and *nco*, Otog is localized near the apical surface of hair cells. (**E-F**) Expression showing hair cells in WT and *nco* larvae at 5dpf. Scale bar = 25 μm. (**G**) Quantification of hair
398 cell numbers in the posterior and anterior macula of WT and *nco* (n = 4).

Figure S1: Spatial differences in mitochondrial membrane potentials. (**A**) While Mitotracker marks active mitochondria in WT, (**B**) *csr* embryos show a lack of Mitotracker expression during early development. Arrow indicates otic vesicle.

405 **Figure S2**: *miR-92a* binding site in the 3' UTR of *rdh12l*. TargetScanFish 6.2 of *rhd12l* in zebrafish shows potential microRNA binding sites including *miR-92a,* which is the most

406 down-regulated gene in *nco* embryos at 24 hpf.

Hair cells (SRA)	Total Reads	ORF - Read Counts	ORF RPKM	5' UTR Read Counts
SRX3022431	14413064	40	0.389182443	0
SRX3022432	100567605	390	0.543821111	0
SRX3022433	50912071	151	0.415916114	0
Support cells (SRA)	Total Reads	ORF - Read Counts	ORF RPKM	5' UTR Read Counts
SRX3022434	54844980	3	0.007670681	14
SRX3022435	59741039	0	0	38
SRX3022436	45498619	0	0	14
			LOG2	
Hair cells RPKM average - ORF		0.44963989	-1.1531	
Hair cells SD		0.082651371		
Support cells RPKM average - ORF		0.002556894	-8.611	
Support cells SD		0.00442867		
Fold Change		7.4579		

408   **Table S1.** Differential expression of pks1 in hair and support cells.