A spike-timing-dependent plasticity rule for single, clustered and distributed dendritic spines

Authors

Sabrina Tazerart[1#], Diana E. Mitchell[1#], Soledad Miranda-Rottmann[1], and Roberto Araya[1*]

Affiliations

[1]Department of Neurosciences, Faculty of Medicine, University of Montreal

[#]These authors contributed equally to this work

*Corresponding author: Roberto Araya Ph.D. Assistant Professor, Department of Neurosciences, Faculty of Medicine, University of Montreal, C.P. 6128 Succursale Centre-Ville, Montreal, Quebec H3C 3J7, T: 514-343-7815, e-mail: roberto.araya@umontreal.ca

SUMMARY

Spike-timing-dependent plasticity (STDP) has been extensively studied in cortical pyramidal neurons, however, the precise structural organization of excitatory inputs that supports STDP, as well as the structural, molecular and functional properties of dendritic spines during STDP remain unknown. Here we performed a spine STDP protocol using two-photon glutamate uncaging to mimic presynaptic glutamate release (pre) paired with somatically generated postsynaptic spikes (post). We found that the induction of STDP in single spines follows a classical Hebbian STDP rule, where pre-post pairings at timings that trigger LTP (t-LTP) produce shrinkage of the activated spine neck and a concomitant increase in its synaptic strength; and post-pre pairings that trigger LTD (t-LTD) decrease synaptic strength without affecting the activated spine shape. Furthermore, we tested whether the single spine-Hebbian STDP rule could be affected by the activation of neighboring (clustered) or distant (distributed) spines. Our results show that the induction of t-LTP in two clustered spines (<5 μm apart) enhances LTP through a mechanism dependent on local spine calcium accumulation and actin polymerization-dependent

neck shrinkage, whereas t-LTD was disrupted by the activation of two clustered spines but

recovered when spines were separated by >40 μm. These results indicate that synaptic

cooperativity, induced by the co-activation of only two clustered spines, provides local dendritic

depolarization and local calcium signals sufficient to disrupt t-LTD and extend the temporal

window for the induction of t-LTP, leading to STDP only encompassing LTP.

INTRODUCTION
39
Dendritic spines, the main recipient of excitatory information in the brain (Araya, 2014), are tiny

protrusions with a small head (~1 μm in diameter and <1 fL volume) separated from the dendrite

by a slender neck. Spines can undergo structural remodeling that is tightly coupled with synaptic

function (Araya, 2014; Araya et al., 2014; Matsuzaki et al., 2001; Matsuzaki et al., 2004), and

are the preferential site for the induction of long-term potentiation (LTP) (Harvey and Svoboda,

2007; Harvey et al., 2008; Matsuzaki et al., 2004; Nishiyama and Yasuda, 2015) and long-term

depression (LTD) (Oh et al., 2013), thought to be the underlying mechanisms for learning and

memory in the brain (Hayashi-Takagi et al., 2015). A variation of LTP and LTD has been

described in pyramidal neurons that involves the pairing of pre- and postsynaptic action

potentials, known as spike-timing dependent plasticity (STDP) (Bi and Poo, 1998; Feldman,

2012). In this process, the timing between pre- and postsynaptic action potentials modulates

synaptic strength, triggering LTP or LTD (Bi and Poo, 1998). The sign and magnitude of the

change in synaptic strength depends on the relative timing between spikes of two connected

neurons (the pre- and postsynaptic neuron (Zhang et al., 1998)). The STDP learning rules and

their dependency on postsynaptic dendritic depolarization (Sjostrom and Hausser, 2006;

Sjostrom et al., 2001), firing rate (Sjostrom et al., 2001), and somatic distance of excitatory

inputs (Froemke et al., 2010; Froemke et al., 2005; Sjostrom and Hausser, 2006) have been

extracted from studies using connected neuronal pairs or by using extracellular stimulating

electrodes, but the precise location and structural organization of excitatory inputs capable of

supporting STDP at its minimal functional unit – the dendritic spine – are unknown.

Activity-dependent spine morphological changes (spine head (Matsuzaki et al., 2004), neck

(Araya et al., 2014), or both (Tonnesen et al., 2014)) have been correlated with changes in

synaptic strength in cortical pyramidal neurons by mechanisms involving biochemical and

electrical spine changes (Araya, 2014; Nishiyama and Yasuda, 2015). Thus, here we asked what

patterns of activity and structural organization of excitatory synaptic inputs support the

generation of t-LTP and t-LTD, and which morphological, biophysical and molecular changes

observed in dendritic spines can account for the induction of t-LTP and t-LTD?

To induce synapse-specific STDP we performed a protocol whereby two-photon (2P) uncaging

of a caged glutamate (MNI-glutamate (Matsuzaki et al., 2001)) at a single spine – to mimic

synaptic release – is preceded or followed in time (STDP timing window (Bi and Poo, 1998)) by

a backpropagating action potential (bAP) to trigger t-LTP (Figure 1A, pre-post) or t-LTD (Figure

2A, post-pre), respectively. Postsynaptic spikes were triggered by current injection via a whole-

cell recording pipette. Two-photon uncaging of a caged glutamate at a single spine triggered

excitatory postsynaptic potentials (uncaging(u)EPSP) that were recorded in the soma of layer 5

(L5) pyramidal neurons before and after the induction of STDP, while the morphology of the

activated spine neck and head was monitored (Figure 1A and 2A). To induce synapse-specific

STDP and monitor calcium levels in the activated spines and parent dendrites we performed a

protocol during which we perform nearly simultaneous 2P uncaging of glutamate and 2P calcium

imaging of the activated spines and nearby dendrites.

Here, we provide evidence showing that the induction of STDP in single or distributed spines

follows a bidirectional Hebbian STDP rule. Furthermore, we show that synaptic cooperativity,

induced by the co-activation of only two clustered spines, disrupt t-LTD (< 40 μm distance

between spines) and extend the temporal window for the induction of t-LTP (< 5 μm distance

between spines) via the generation of differential local calcium signals leading to an STDP rule

for clustered inputs only encompassing LTP.

RESULTS

Induction of t-LTP in single dendritic spines

To induce t-LTP, we used a repetitive spike-timing protocol (40 times, 0.5 Hz) in which 2P uncaging of glutamate at a single spine was closely followed in time (+7 or +13 ms later, see Methods section) by a bAP (Figure 1A). We evaluated spine morphology and uEPSP amplitude for 40 min following STDP induction to establish the time course of STDP at individual synapses (Figure 1C.1 and D.1). In addition, the maximum uEPSP change and concomitant changes in spine morphology observed in each experiment are shown (Figure 1C.2 and D.2).

A repetitive pre-post pairing protocol of +13 ms reliably induced t-LTP (significant increase in the uEPSP amplitude over time, $P < 0.001$, $n = 7$ experiments, from 6 neurons, from 6 mice, Figure 1B.1 and C.1), and shortening of the activated spine neck within a few minutes ($P < 0.001$), with no significant change in spine head volume ($n = 7$, Figure 1B.2 and C.1). These results were also consistent when we considered the maximum uEPSP change in amplitude from each experiment and concomitant changes in spine morphology (uEPSP = 134.21 ± 3.29%, $P < 0.001$, $n = 7$; neck length = 71.88 ± 10.66%, $P < 0.05$, $n = 7$; spine head volume = 98.11 ± 7.34%, $P = 0.81$, $n = 7$) (Figure 1C.2). We obtained similar results when we instead considered the average of all the values obtained following t-LTP induction for uEPSP amplitude, neck length and head volume (Figure S1). This effect was specific to the activated spine (Figure 1B.2), with neighbouring spines having no appreciable changes in their neck length or head volume (neck length = 98.09 ± 5.06%, $P = 0.71$, $n = 13$; head volume = 103.01 ± 3.61%, $P = 0.42$, $n = 14$). Control experiments showed that there was no significant change in uEPSP amplitude or spine morphology following the STDP protocol when either action potentials or

synaptic stimulation were applied in isolation, as well as when we monitored the long-term

stability of these parameters without any STDP protocol (Figure S2).

A pre-post pairing of + 7 ms showed a non-significant tendency for the induction of LTP

following t-LTP induction, and a non-significant tendency for the shrinkage of the activated

spine neck (n = 8 experiments, from 8 neurons, from 8 mice, Figure 1D.1). When the maximum

change in uEPSP amplitude from each experiment and the concomitant changes in spine

morphology were analyzed, we saw no significant changes in uEPSP amplitude and spine

morphology (Figure 1D.2, uEPSP = 112.27 ± 14.19%, P = 0.42, n = 8; neck length = 88.90 ±

5.89%, P = 0.10, n = 8; head volume = 97.81 ± 4.54%, P = 0.64, n = 8) (Figure 2E). Similar

results were observed when we instead considered the average of all the values obtained

following t-LTP induction (+7 ms) for uEPSP amplitude, neck length and head volume (Figure

S1). Because voltage has been shown to be an important factor in the induction of t-LTP and t-

LTD, we verified that the initial uEPSP amplitudes were not significantly different for pre-post

pairing protocols of +13 ms versus +7 ms (uEPSP: 0.62 ± 0.14 versus 0.53 ± 0.16 mV, P = 0.68;

Figure S3).

These results indicate that for the pairing times tested, there is a preferred pre-post t-LTP pairing

time-window (+ 13 ms) at which activated spines in basal dendrites from L5 pyramidal neurons

undergo a significant increase in synaptic strength, and a concomitant neck shrinkage (Figure

2E).

Induction of t-LTD in single dendritic spines

We then studied t-LTD in single spines by using a repetitive spike-timing protocol (40 times, 0.5

Hz) in which 2P uncaging of glutamate at a single spine was preceded in time (-15 or -23 ms) by

a bAP (post-pre protocol, Figure 2A). When postsynaptic spikes preceded presynaptic firing by 15 ms (i.e., -15 ms), a significant reduction of the uEPSP amplitude occurred within a few minutes following induction of t-LTD (n= 6 experiments, from 5 neurons and 5 mice, Figure 2B.1 and C.1, P < 0.001), with no significant changes in spine neck length or head dimension (Figure 2B.2 and C.1). Furthermore, when the maximum change in uEPSP amplitude after the induction of t-LTD in single spines at pairings of -15 ms was analyzed (Figure 2C.2) we also observed a significant depression of uEPSP amplitude (uEPSP = 71.52 ± 7.07%, P < 0.01, n = 6), with no significant changes in spine morphology (Figure 2B.2 and C.2, neck length = 105.54 ± 9.85%, P = 0.62, n = 6; head volume = 103.25 ± 3.02%, P = 0.33, n = 6). Interestingly, after the induction of t-LTD in single spines when postsynaptic spikes preceded presynaptic firing by 23 ms (i.e., -23ms) there were no significant changes in the amplitude of the uEPSPs or in the spine neck length and head dimensions for the duration of the recordings (n= 7 experiments, from 7 neurons and 7 mice, Figure 2D.1). These results were also consistent with analyses of the maximal uEPSP change in each experiment and concomitant spine morphology (Figure 2D.2: uEPSP = 82.09 ± 9.89%, P = 0.12, n = 7; neck length = 84.15 ± 7.73%, P = 0.09, n = 7; head volume = 98.81 ± 5.57%, P = 0.84, n = 7). There was no significant difference between the initial uEPSP amplitude for post-pre pairing protocols of -15 ms versus -23 ms (EPSP: 0.59 ± 0.07 versus 0.49 ± 0.08 mV, P = 0.42; Figure S3). We obtained similar results when we instead considered the average of all the values obtained following t-LTD induction in single spines for uEPSP amplitude, neck length and head volume (Figure S1). This indicates that for the pairing times tested, only a post-pre t-LTD pairing time-window of -15 ms can effectively induce LTD in single dendritic spines in the basal dendrites from L5 pyramidal neurons.

Taken together these results show that the induction of t-LTP and t-LTD in single spines follows

a Hebbian-STDP learning rule that is bidirectional, and favors presynaptic inputs that precede

postsynaptic spikes and depresses presynaptic inputs that are uncorrelated with postsynaptic

spikes at a very precise and narrow temporal window (+13 ms for the generation of t-LTP and -

15 ms for t-LTD, Figure 2E-F). The single spine STDP rule we observed has a narrower post-pre

LTD induction pairing time window than previously observed in connected pairs of L2/3

(Feldman, 2000; Froemke et al., 2005) and L5 pyramidal neurons (Sjostrom et al., 2001) – where

the presynaptic control of t-LTD via an mGluR and/or cannabinoid type 1 receptor-dependent

mechanism (Bender et al., 2006; Hashimotodani et al., 2005; Nakamura et al., 1999; Sjostrom et

al., 2003) could plausibly account for these differences.

Induction of t-LTP in clustered dendritic spines

It has been suggested that STDP not only depends on spike timing and firing rate but also on

synaptic cooperativity and the amount of voltage generated at the postsynaptic site (Sjostrom and

Hausser, 2006; Sjostrom et al., 2001). Furthermore, it has been shown in mouse hippocampal

CA1 pyramidal cells that the induction of t-LTP in a single spine in low extracellular Mg^{2+}

conditions can reduce the threshold for the induction of plasticity of a neighboring spine

activated 90 seconds later (Harvey and Svoboda, 2007). However, a direct demonstration of

synaptic cooperativity for synchronous, or nearly synchronous synaptic inputs, in physiological

extracellular Mg^{2+} conditions, and at the level of single spines for the induction of STDP in the

dendrites of pyramidal neurons remains unknown. Hence, we next directly tested if synaptic

cooperativity, marked by the co-induction of t-LTP in clustered dendritic spines from basal

dendrites of L5 pyramidal neurons, and the local spatio-temporal summation of inputs, can

generate a local dendritic depolarization and local calcium signals, that are high enough to disrupt the single spine STDP learning rule described in Figure 2E. To test this, a two spine STDP protocol (forty 2P uncaging pulses, pulse duration 2ms, 0.5Hz, see methods section) was performed, whereby nearly simultaneous 2P uncaging of caged glutamate in clustered (distance between spines < 5 μm) spines was followed in time by a bAP to trigger t-LTP (Figure 3A). With this protocol, we investigated whether activating clustered spines extended the pre-post timing window capable of generating LTP by increasing the degree of depolarization immediately before the postsynaptic spike at timings where plasticity was not reliably generated. Specifically, we induced t-LTP in two clustered spines at pre-post timings of +7 ms, and surprisingly found that this protocol was in fact capable of effectively and significantly generating increases in uEPSP amplitude (Figure 3B.1) and the concomitant shrinkage of the activated spine neck, with no apparent changes in its spine head size (Figure 3B.2). Pooled data showed that significant increases in uEPSP amplitude and shrinkage of the spine neck of the activated spines occurs only a few minutes post t-LTP induction and lasted for the duration of the recordings (Figure 3C, $P < 0.01$, n = 10 spines from 8 experiments, 6 neurons and 6 mice), with no significant changes in the spine head size (Figure 3C, n = 16 spines from 8 experiments, 6 neurons, 6 mice). Similar results were observed when we analyzed the maximal change in uEPSP amplitude in each experiment and the concomitant spine neck length and head size of the two clustered spines after induction of t-LTP at pairings of + 7 ms (Figure 3H; uEPSP = 130.86 ± 8.18%, $P < 0.01$, n = 8; neck length = 73.22 ± 5.84%, $P < 0.01$, n = 10; head volume =102.00 ±2.58%, $P = 0.45$, n = 16). We obtained similar results when we instead considered the average of all the values obtained following t-LTP induction in two clustered spines for uEPSP amplitude, neck length and head volume (Figure S1). In control experiments, no significant

change in uEPSP amplitude or spine morphology was observed when we monitored the long-

term stability of these parameters without any STDP protocol (Figure S2). These results indicate

that synaptic cooperativity of nearly simultaneous synaptic inputs – shown by the induction of t-

LTP in only two clustered spines (< 5 μm apart) – is sufficient to significantly trigger synaptic

potentiation and shrinkage of the activated spine necks at a pre-post timing that is otherwise

ineffective at generating significant morphological changes and synaptic potentiation when only

one spine is being activated (for comparison between one versus two clustered spines see Figure

S4). Hence, the synaptic cooperativity of only two neighbouring synaptic inputs onto spines (< 5

μm apart) in the basal dendrites of L5 pyramidal neurons extends the pre-post timing window

that can trigger potentiation (Figure 3H, and compare Figure 3C with Figure 1D.1).

Induction of t-LTP in distributed dendritic spines

To precisely study the effect that inter-spine distance of the nearly simultaneously activated

spines has on synaptic cooperativity and the induction of t-LTP at pairings of +7 ms, we also

performed experiments where the activated spines were further away (> 5 μm apart) from each

other. When we correlated the inter-spine distance and the maximal uEPSP change following t-

LTP induction, we found that as two activated spines were further away from each other, the less

t-LTP was induced (Figure 3E). The induction of t-LTP decayed exponentially as a function of

the inter-spine distance with a length constant (λ) of 5.7 μm. Therefore we used this value as the

boundary between clustered (< 5 μm) and distributed (> 5 μm) spines for the induction of t-LTP

at pairings of + 7ms. Clustered spines were located in the same basal dendrite (n = 8/8 pairs),

while distributed spines were either in sister branches emanating from the same bifurcation point

(n = 7/13 pairs) or in the same basal dendrite (n = 6/13 pairs). By separating the data in this

manner, the t-LTP protocol (+ 7ms) in distributed spines (> 5 μm, Figure 3D) failed to induce t-

LTP (Figure 3F.1) or changes in spine head size and neck length at all the time tested post t-LTP

induction (Figure 3G, n = 13 experiments, n = 26 spines, from 11 neurons and 7 mice). For

comparisons between the activation of two clustered versus two distributed spines with a pre-

post timing of + 7ms see Figure S4.   When we analyzed the maximum uEPSP change in

amplitude in each experiment and the concomitant changes in spine head volume and neck

length in the activated spines after t-LTP induction in two distributed spines, we also observed a

complete loss of t-LTP induction, with no changes in spine morphology (Figure 3H; uEPSP =

92.52 ± 6.570%, P = 0.28, n = 13 experiments; neck length = 94.76 ± 4.51%, P = 0.26, n = 20

spines; head volume = 98.65 ± 2.41%, P = 0.58, n = 26 spines, from 13 experiments performed

in 11 neurons from 7 mice). Similar results were obtained when we instead considered the

average of all the values obtained following t-LTP induction (+7ms) in distributed spines for

uEPSP amplitude, neck length and head volume (Figure S1). In summary, this data shows that

the induction of t-LTP in nearly synchronously activated spines at pre-post timings of + 7 ms is

disrupted when the two activated spines were more than 5 μm apart in the basal dendrites of L5

pyramidal neurons.   These results uncover a two-spine activation crosstalk-spatial barrier of 5

μm for the induction of t-LTP to occur at pre-post timings of + 7 ms.

Molecular mechanisms responsible for the generation of t-LTP in dendritic spines

The results led us to consider the possible mechanisms underlying the generation of t-LTP at the

level of single spines. Specifically, we asked why the induction of t-LTP in single and clustered

spines is associated with the shrinkage of the activated spine neck. We and others have reported

that the induction of LTP can trigger activity dependent changes in neck length (Araya et al.,

2014; Tonnesen et al., 2014) and spine head size (Matsuzaki et al., 2004; Nishiyama and Yasuda,

2015; Tonnesen et al., 2014), and that the amplitude of uEPSP recorded at the cell soma is

inversely proportional to the length of the activated spine neck (Araya et al., 2006b; Araya et al.,

2014; Richardson et al., 2009). However the mechanisms by which the t-LTP-induced neck

shrinkage is associated with synaptic plasticity remains unknown. Numerical simulations show

that the EPSP amplitude/neck length correlation can be explained by variations in synaptic

conductance, electrical attenuation through the neck, or a combination of the two (Araya et al.,

2014). Nevertheless, solutions that rely exclusively on the passive electrical attenuation of

synaptic inputs through the spine neck assume very high (> 2 GOhm) neck resistance (Araya et

al., 2014), which are at odds with recent spine neck resistance estimations (Harnett et al., 2012;

Jayant et al., 2017). These results suggest that the control of AMPA receptor content in spines

could contribute significantly to the observed t-LTP-dependent changes in synaptic strength. To

experimentally study the contribution of AMPA receptors to these phenomena, we performed t-

LTP experiments in two clustered spines from L5 pyramidal neurons recorded via patch pipettes

loaded with intracellular solution containing 200 μM PEP1-TGL – a peptide that inhibits AMPA

receptor incorporation to the postsynaptic density (PSD) by blocking GluR1 C-terminus

interaction with PDZ domains at the PSD (Hayashi et al., 2000) (Figure 4A). PEP1-TGL

incubation by itself did not trigger a run-down of uEPSP amplitude or changes in spine

morphology over time (Figure S5A). Pooled data from experiments where a repetitive pre-post

pairing protocol of + 7 ms was used to activate clustered spines in the presence of PEP1-TGL

show that the peptide completely inhibited t-LTP for the duration of the experiment (Figure 4B.1

and C.1), but had no effect on the t-LTP-induced shrinkage of the activated spine necks (Figure

4B.2 and C.1, P < 0.001, n = 6 spines, from 5 experiments, 5 neurons and 5 mice) or in

modifying spine head size (n = 10 spines, from 5 neurons and 5 mice, Figure 4C.1). Furthermore,

when we analyzed the maximum change in uEPSP amplitude and the concomitant spine

morphology after the induction of t-LTP in the presence of PEP1-TGL, we also found an

inhibition of t-LTP, but no effect on the t-LTP-induced shrinkage of the spine neck (Figure 4B.2-

C.2, uEPSP = 94.82 ± 14.82%, P = 0.74, n = 5 experiments, from 5 neurons and 5 mice; neck

length = 83.92 ± 5.35%, P < 0.05, n = 6; head volume = 100.08 ± 3.23%, P = 0.98, n = 10). We

obtained similar results when we instead considered the average of all the values obtained

following t-LTP induction in two clustered spines in the presence of PEP1-TGL for uEPSP

amplitude, neck length and head volume (Figure S1). No significant difference was observed

between the initial uEPSP amplitude for pre-post pairing protocols of +7 ms with versus without

PEP1-TGL (uEPSP: 1.06 ± 0.2 versus 1.16 ± 0.28 mV, P = 0.81; Figure S3). These results

indicate that GluR1 receptor incorporation into the PSD - via its interaction with PDZ domains -

is required for the induction of t-LTP in spines. However, the role of the spine neck shrinkage in

AMPA receptor incorporation into the PSD and ultimately on the induction of t-LTP remains

open.

Experimental and theoretical studies have indicated that lateral diffusion of AMPA receptors into

and out of the spine head can be restricted by the spine neck geometry (Ashby et al., 2006;

Holcman and Schuss, 2011; Holcman and Triller, 2006; Kusters et al., 2013). In particular,

lateral diffusion of AMPA receptors into and out of mushroom spines (long-necked spines) has

been shown to be significantly slower than that observed in stubby spines (small-necked spines)

(Ashby et al., 2006) – which is supported by studies that show reduced diffusion of membrane

proteins located in spine necks (Shim et al., 2012). In addition, quantitative models using

realistic spine morphologies indicate that decreasing the radius and increasing the spine neck

length increases the retention of AMPA receptors at the synapse (Kusters et al., 2013), even

when their interaction with scaffolding cytoskeletal proteins is neglected (Holcman and Schuss,

2011). Actin is highly enriched in the spine neck and head (Frost et al., 2010), and plays an

important role in anchoring AMPA receptors in the spine (Allison et al., 1998) and AMPA

receptor trafficking (Hanley, 2014), being instrumental for synaptic transmission and plasticity

(Honkura et al., 2008; Kim and Lisman, 1999; Ramachandran and Frey, 2009). Hence, to address

the role that t-LTP-induced neck shrinkage has on AMPA receptor lateral trafficking to the PSD,

and the generation of t-LTP in the activated spines we focused on actin dynamics. We used the

actin polymerization inhibitor latrunculin A (Lat-A) (Allison et al., 1998; Kim and Lisman,

1999; Ramachandran and Frey, 2009) (Figure 4D) − which did not trigger any run-down of

uEPSP amplitude or changes in spine morphology over time in the absence of STDP induction

(Figure S5B) − to test the potential role of actin dynamics on the spine induction of t-LTP, and

on the neck shrinkage and AMPA receptor incorporation into the PSD in the activated spines

(Figure 4D). The induction of t-LTP at pre-post pairings of +7 ms in two clustered spines in the

presence of 100 nM Lat-A completely blocked the shrinkage of the activated spine necks and the

induction of t-LTP (Figure 4E and F.1, n = 8 spines, from 3 neurons and 2 mice), inducing

instead a significant reduction in uEPSP amplitude over time (Figure 4F.1, $P < 0.001$, n = 4

experiments, from 3 neurons and 2 mice). These observations were also consistent with analyses

of the maximal change in uEPSP amplitude in each experiment and concomitant spine

morphology post t-LTP induction in the presence of Lat-A (Figure 4F.2, uEPSP= 55.45 ± 7.13%,

$P < 0.01$, n = 4 experiments; neck length = 87.20 ± 9.15%, $P = 0.21$, n = 8 spines; head volume =

101.25 ± 10.99%, $P = 0.91$, n = 8 spines).

We obtained similar results when we instead considered the average of all the values obtained

following t-LTP induction in two clustered spines in the presence of 100 nM Lat-A for uEPSP

amplitude, neck length and head volume (Figure S1).

No significant difference was observed between initial uEPSP amplitudes for pre-post pairing

protocols of +7 ms with versus without Lat-A (uEPSP: 0.94 ± 0.20 versus 1.16 ± 0.28 mV, P =

0.61; Figure S3). The lack of run-down of uEPSP amplitude over time in neurons treated with

Lat-A in the absence of STDP induction (Figure S5B), but the significant depression in uEPSP

after the induction of t-LTP suggests that the induction of plasticity, and the rearrangement of

actin filaments de-stabilized AMPA receptors, leading to removal from the PSD.

In summary, these results show that actin polymerization is required for the t-LTP-dependent

neck shrinkage and the induction of plasticity. Our findings further suggest that the induction of

t-LTP occurs via a mechanism that involves a neck-shrinkage-dependent facilitated diffusion of

GluR1 subunits from the spine neck to the head, and subsequent incorporation into the PSD. We

hypothesize that a shorter and wider neck facilitates the transport of AMPA receptors into the

spine head (Figure 4D), a mechanism that is required for the induction of t-LTP.

Induction of t-LTD in clustered and distributed dendritic spines

We then studied whether the induction t-LTD in single spines observed at pairings of -15 ms

could be affected by synaptic cooperativity. Our reasoning was based on two previous

observations which suggest that 1) t-LTP induction in the distal dendrites of L5 pyramidal

neurons (layer 3- L5 pyramidal neuron pairs) triggers LTD instead of LTP, and 2) that LTD can

be converted into LTP by increasing the local voltage (Sjostrom and Hausser, 2006). We

hypothesized that the induction of t-LTD in single spines depends on the degree of local

depolarization and hence, LTD can be disrupted by the activation of neighboring spines. To test

this, we performed repetitive spike-timing protocol (40 times, 0.5 Hz) in which 2P uncaging of

glutamate at two spines (separated by up to 100 μm) was preceded in time (-15 ms) by a bAP

(Figure 5D and Figure S6A). Surprisingly, we found that this t-LTD protocol failed to induce

any change in uEPSP amplitude or spine head volume with only a slight but significant reduction

in spine neck length (Figure S6A). When we analyzed the maximum uEPSP change in amplitude

in each experiment and the concomitant morphological alterations in the activated spines after t-

LTD induction, we observed a complete inhibition in the induction of t-LTD, and no change in

spine morphology (Figure S6B; uEPSP = 93.22 ± 6.29%, P = 0.30, n = 17 experiments from 14

neurons and 14 mice; neck length = 88.56 ± 5.69%, P = 0.06, n = 23 spines; head volume =

102.41 ± 6.10%, P = 0.69, n = 34 spines). To more precisely characterize the effect of activating

two spines on the induction of t-LTD, we correlated the inter-spine distance and the uEPSP

change following STDP induction (see Methods). We found that as the two activated spines were

further away from each other, the more t-LTD was recovered (Figure 5E). Specifically, the

uEPSP change decayed exponentially as a function of inter-spine distance with a length constant

(λ) of 43.5 μm. Therefore, we used this value as a boundary between clustered (< 40 μm) and

distributed (> 40 μm) spines. Using this classification, clustered spines were located in the same

dendrite (n = 11/12 pairs) or in sister branches emanating from the same bifurcation point (n =

1/12 pairs), while distributed spines were always located on separate dendrites (n = 5/5 pairs).

When we separated our data in this manner, the t-LTD protocol in two clustered spines (Figure

5A) failed to induce LTD (Figure 5B.1) or changes in spine head size at all the times tested post

t-LTD induction (Figure 5B.2 and C, n = 12 experiments, n = 24 spines, from 11 neurons and 11

mice), with only a slight but significant induction of shrinkage of the spine neck at some time

points (Figure 5C, P < 0.05, n = 19 spines, from 11 neurons and 11 mice). For comparison

between the activation of one versus two clustered spines with a post-pre timing of -15 ms see

Figure S7. When we analyzed the maximum uEPSP change in amplitude in each experiment and

the concomitant morphological alterations in the activated spines after t-LTD induction in two

clustered spines, we also observed a complete inhibition in the induction of LTD, a slight but

significant reduction in spine neck length, and no changes in spine head size (Figure 5H; uEPSP

= 101.70 ± 7.02%, P = 0.81, n = 12 experiments; neck length = 85.28 ± 5.96%, P < 0.05, n = 19

spines; head volume = 102.98 ± 8.86%, P = 0.73, n = 24 spines, from 12 experiments performed

in 11 neurons from 11 mice). We obtained similar results when we instead considered the

average of all the values obtained following t-LTD induction in clustered spines for uEPSP

amplitude, neck length and head volume (Figure S1). These results were surprising since not

only did we not observe t-LTD in clustered spines, but we also observed significant neck

shrinkage with no LTP (see Figure 1 and 3). To account for this observation, we explored if there

was a correlation between the induction of plasticity in these experiments and both the shrinkage

of the spine neck and the distance between the activated clustered spines – since the local

voltage, and hence the induction of plasticity, could be affected by the distance between the

activated clustered spines. Indeed, we found that the distance between the activated spines under

these experimental conditions (t-LTD induction protocol in clustered spines) is correlated with

the induction of plasticity and the shrinkage of the activated spine necks (Equation 1 in Methods;

P < 0.01; Figure S6D). This analysis suggests that during t-LTD induction the structural

arrangement of clustered spines (< 40 μm) determines the sign and magnitude of the change in

synaptic strength and concomitant neck shrinkage.

We next investigated the mechanisms underlying the disruption of t-LTD by activating spines

separated by increasingly larger distances (Figure 5D). Interestingly, the induction of t-LTD in

spines separated by more than 40 μm (distributed spines) was capable of recovering the

generation of LTD (Figure 5E-F). Pooled data from all experiments demonstrate that the

activation of distributed spines reliably induces t-LTD (Figure 5G, significant reduction in

uEPSP amplitude, $P < 0.01$, n = 5 experiments from 5 neurons and 5 mice), without triggering

changes in neck length or spine head size (Figure 5F and G). When we analyzed the maximal

change in uEPSP amplitude in each experiment and concomitant spine morphological changes,

we saw a significant induction of t-LTD and no change in spine morphology (Figure 5H, uEPSP

$= 72.86 \pm 8.08\%$, $P < 0.05$, n = 5 experiments, neck length $= 97.85 \pm 15.47\%$, $P = 0.89$, n = 6

spines; head volume $= 101.06 \pm 4.59\%$, $P = 0.82$, n = 10 spines) as what was found in

experiments where t-LTD was generated at pairing times of -15 ms in single dendritic spines

(Figure 2B-C). We obtained similar results when we instead considered the average of all the

values obtained following t-LTD induction in distributed spines for uEPSP amplitude, neck

length and head volume (Figure S1). No significant difference was observed between the initial

uEPSP amplitude for clustered versus distributed spines activated with post-pre pairings of -15

ms (EPSP: 1.06 ± 0.13 versus 1.31 ± 0.19 mV, $P = 0.25$; Figure S3). For comparison between the

activation of clustered versus distributed spines after post-pre pairings of -15 ms, see Figure S7.

In summary, this data shows that the induction of t-LTD at pairing times of -15 ms was disrupted

when only two clustered spines (< 40 μm apart) were nearly simultaneously activated in the

basal dendrites of L5 pyramidal neurons, but could be recovered if the activated spines are

distributed (> 40 μm) in the dendritic tree.

Spine calcium transients during the induction of t-LTP and t-LTD in single and clustered dendritic spines

Calcium is critical for the induction of synaptic plasticity (Artola and Singer, 1993; Cummings et al., 1996; Fino et al., 2010; Lynch et al., 1983; Malenka et al., 1988), and high or low local concentration difference in dendrites and spines are thought to be associated with gating LTP or LTD, respectively (Ismailov et al., 2004; Lisman, 1989; Nevian and Sakmann, 2006). Therefore, to investigate the different mechanisms – with respect to local calcium accumulations – underlying the induction of t-LTP and t-LTD in single versus two clustered spines, we performed 2P calcium imaging in a region of interest (ROI) of the activated spines and their parent dendrites during STDP induction protocols throughout each of the 40 pre-post or post-pre repetitions (see Methods). The "before" images correspond to the calcium signals observed in the ROI right before the pairing in each repetition – uncovering the lack or presence of local calcium accumulation during the 40 pairing repetitions. The "after" images correspond to the calcium signals observed in the ROI right after the pairing in each repetition – uncovering a proxy for the amplitude and local calcium accumulation during the 40 pairing repetitions.

To dissect potential differences in local calcium signals and accumulation that can account for the presence or absence of t-LTP and t-LTD induction in clustered versus distributed spines, we imaged 2P calcium activity during five different STDP induction protocols: (1) pre-post pairing of +7 ms in one spine; (2) pre-post pairing of +7 ms in two clustered spines; (3) pre-post pairing of +13ms in one spine; (4) post-pre pairing of -15 ms in one spine; (5) post-pre pairing of -15 ms in two clustered spines.

During the pre-post (+7 ms) pairing protocol in single spines we found that, across the 40 repetitions, there was little to no calcium accumulation in the spine or dendrite (Figure 6A and

left panels in Figure 6B, n = 7 experiments, from 6 neurons, and 4 mice). As expected, there was,

however, a significant increase in calcium immediately following the stimulation (left panels in

Figure 6D) that due to the lack of accumulation throughout the 40 repetitions, did not build up a

local calcium signal in the activated dendrites and spines. In contrast, when we applied the exact

same pairing protocol (pre-post + 7ms) in two clustered spines, there was a striking calcium

accumulation in both the activated spines and dendrite that was evident when we analyzed the

images taken before (Figure 6C and middle panels in Figure 6B) and after stimulation (middle

panels in Figure 6D, n = 6 experiments, from 4 neurons, and 4 mice). Thus, activating just one

additional spine using the same pairing protocol alters the calcium dynamics (compare black and

green traces in right panels of Figure 6B and D), possibly through a mechanism that is incapable

of extruding calcium increases in spines in between pre-post repetitions, leading to its build up in

spines and parent dendrites, which ultimately guide the induction of plasticity.

Since the activation of a single spine with a repetitive pre-post pairing protocol of +13 ms

reliably induced t-LTP (Figure 1B-C), we performed experiments to measure the calcium signals

in the activated spine and parent dendrite during the pre-post (+13 ms) pairing protocol. We

found that, across the 40 repetitions, there was a significant calcium accumulation in both the

activated spine and dendrite that was evident when we analyzed the images taken before (Figure

7A and B) and after stimulation (Figure 7A and C).

In summary, the induction of t-LTP in a single spine at a pre-post pairing protocol of +13 ms

(Figure 7) and in clustered spines at a pairing time of +7ms (which is otherwise ineffective in

triggering t-LTP if only one spine is being activated) (Figure 6) are correlated with an

accumulation of calcium in spines and parent dendrites, likely being responsible for the induction

of t-LTP.

Next, we performed the same experiments with a post-pre (-15 ms) pairing protocol in both

single and clustered spines. In single spines, we observed moderate calcium increases (Figure

8A, n = 5 experiments, from 4 neurons, and 4 mice) that were observed when we analyzed

images taken before (left panels in Figure 8B) and after the post-pre stimulation (left panels in

Figure 8D). Surprisingly, we found similar results to those observed with single spine t-LTD

induction protocols, when we applied the same pairing protocol in two clustered spines (Figure

8C, n = 6 experiments, from 4 neurons, and 3 mice) before (middle panels in Figure 8B) or after

the induction protocol (middle panels in Figure 8D), even though no plasticity is induced in this

condition. As suggested by previous studies (Graupner and Brunel, 2010), we hypothesize that

the range of spine calcium levels required for the induction of t-LTD is relatively narrow, and

that the resolution with our current experimental set-up is not sufficient to tease apart

significantly different calcium dynamics in one versus two clustered spines during a post-pre

pairing protocol of -15 ms. Moreover, models of STDP provide evidence that, in addition to

overall calcium levels, the detailed time course of calcium levels in the postsynaptic cell during a

pairing protocol also guide the induction of plasticity (Rubin et al., 2005). Nonetheless, these

results suggest that the induction of t-LTD does not require significant calcium accumulations

during the 40 repetitions, and most likely depends on the amplitude of calcium signals right after

the stimulation.

DISCUSSION

We uncovered the STDP rules for single, clustered and distributed dendritic spines in the basal dendrites of L5 pyramidal neurons. Our results show that the induction of STDP in single spines follows a classical Hebbian STDP learning rule that is bidirectional, in which presynaptic input leading postsynaptic spikes generates t-LTP and postsynaptic spikes preceding presynaptic activation of single dendritic spines results in t-LTD. Furthermore, we found that the induction of t-LTP triggers the shrinkage of the activated spine neck, without any significant changes in the spine head size, building upon our previous findings of the activity-dependent shrinkage of the spine neck (Araya et al., 2014). Our results indicate that the induction of t-LTP requires 1) the incorporation of new GluR-1 receptors with PDZ-domain containing proteins in the PSD and, 2) an actin polymerization-dependent neck shrinkage of the activated spine neck (Figure 4). We showed that the induction of t-LTP triggers actin-dependent neck shrinkage, which is likely required for the lateral diffusion of GluR-1 receptors from the spine neck to the spine head, and its incorporation to the PSD – generating plasticity. In support of this spine mechanism of LTP induction is a recent report showing that AMPA receptor surface diffusion is fundamental for the induction of hippocampal LTP and contextual learning (Penn et al., 2017). In addition, we found that the induction of t-LTD was not accompanied with spine neck or head changes, which is at odds with previous findings suggesting structural changes in spine head volume during the induction of LTP or LTD (Matsuzaki et al., 2004; Oh et al., 2013; Tanaka et al., 2008). The discrepancy between our results and those observed previously after the induction of t-LTP (head enlargement (Harvey and Svoboda, 2007; Tanaka et al., 2008)), LTP (Matsuzaki et al., 2004), or LTD (head shrinkage, (Oh et al., 2013)) using glutamate uncaging are likely explained by methodological differences. While our data was obtained using ACSF with physiological

concentrations of magnesium and calcium, those from other reports were done in low or a magnesium-free ACSF (Matsuzaki et al., 2004; Oh et al., 2013), low calcium extracellular solution for the induction of LTD (Oh et al., 2013), or in a magnesium-free ACSF and an intracellular solution containing 5 µM actin that was required for the t-LTP-mediated spine head enlargements (Tanaka et al., 2008).

Nonetheless, it has been shown in vivo that a spike-timing protocol triggers receptive field plasticity in layer 2/3 pyramidal neurons is correlated with spine head volume changes (enlargement and shrinkage) observed after 1.5-2 hours (El-Boustani et al., 2018). Taken together, these data, and previous observations (Araya et al., 2014), suggest that there is a new form of structural spine plasticity during t-LTP that involves rapid neck shrinkage without head volume enlargements. In addition, we show that the induction of t-LTD does not require structural spine changes. Although spines have the machinery and do undergo structural head changes, we propose that our results represent a stage during memory formation that occurs before structural head volume changes, a process likely linked with memory consolidation. Importantly, our data suggest that during STDP, the use of spine volume changes as the sole proxy for LTP or LTD (El-Boustani et al., 2018) is not a complete representation of plasticity in spines from dendrites in cortical pyramidal neurons.

We then explored the functional consequences of synaptic cooperativity of nearly simultaneous excitatory inputs on STDP. Our results show that the induction of t-LTP in two clustered spines - separated by less than 5 µm - is sufficient to induce LTP and shrinkage of the activated spine necks at a pre-post timing (+ 7ms) that is otherwise ineffective at triggering significant morphological changes and synaptic potentiation when only one spine is being activated. To uncover the distance between the nearly simultaneously activated spines capable of supporting

synaptic cooperativity and the induction of t-LTP, we varied the distance of the activated spines. Our results show that the induction of t-LTP is suppressed when spines are separated by more than 5 μm apart (Figure 3), with an effective length constant (λ), that represents the length at which the electrotonic potential decays to a value of 37% of that at the point of origin, of 5.7 μm These results show that the nearly simultaneous activation of clustered spines, less than < 5 μm apart, can extends the pre-post timing window that can trigger potentiation. This data, together with results showing that the induction of t-LTP in one spine can change the threshold for the induction of plasticity in a spine located at less than 10 μm away and activated 90 seconds later (Harvey and Svoboda, 2007), indicate that there is a spatially restricted dendritic compartment that is required for spine crosstalk and the broadening of the STDP timing required for triggering t-LTP (Figure 9).

On the other hand, the induction of t-LTD in two clustered spines disrupts the generation of LTD leading to a STDP learning rule that is incapable of supporting LTD, but only encompasses LTP (Figure 9). We next investigated the dendritic mechanisms responsible for the disruption of t-LTD, and found that the induction of t-LTD is recovered when the activated spines are separated by more than 40 μm (Figure 5, Figure 9). Interestingly, the effective length constant (λ) in the basal dendrites of L5 pyramidal neurons has been reported to be 50 μm (Nevian et al., 2007). This value of λ supports the idea that significant voltage attenuations – capable of recovering LTD – can be expected when the t-LTD induction protocol is triggered in spines that are separated by more than 40 μm in the basal dendrites of L5 pyramidal neurons (Figure 5 D-H). However, we cannot discard that other mechanisms, such as the diffusion of active molecules (Harvey et al., 2008), could contribute to the switch from LTD to no-LTD induction observed in distributed/single spines and clustered spines, respectively. These results are in discrepancy with

observations showing that in connected pairs of L5-L5 pyramidal cells, t-LTD is reliably

generated after post-pre pairing protocols (Sjostrom et al., 2001). A likely explanation for this

apparent controversy is that the synaptic inputs from one L5 pyramidal neuron to another are

distributed (Markram et al., 1997). Importantly, clustered and distributed excitatory inputs have

been described in the dendrites of pyramidal neurons both *in vitro* and *in vivo* (Araya, 2014;

Bloss et al., 2018; Kasthuri et al., 2015; Schmidt et al., 2017). Our results clearly show the

functional importance that the structural and temporal organization of excitatory synaptic inputs

have on the induction of t-LTP and t-LTD, and how just two clustered excitatory synaptic inputs

are capable of altering the STDP learning rule in the basal dendrites of L5 pyramidal neurons

(Figure 9).

How the synaptic activation of just one extra clustered spine is capable of (1) inducing t-LTP at a

pre-post timing that is otherwise ineffective in inducing potentiation and (2) disrupting the

induction of t-LTD? To explore the mechanisms that may be responsible for these observations

we imaged local calcium signals in the activated spines and parent dendrites before and after

each of the 40 pairings performed during t-LTP and t-LTD induction protocols. Our reasoning

for performing these experiments was based on findings that different levels of depolarization

gate local calcium signals, which depending on its magnitude and kinetics, can generate LTP

(high calcium) or LTD induction (sustained but moderate calcium signals) (Feldman, 2012;

Lisman, 1989; Yang et al., 1999). In addition, calcium-based modeling studies of STDP have

shown that different calcium dynamics mediate the induction of t-LTP versus t-LTD (Graupner

and Brunel, 2010; Rubin et al., 2005). Specifically, the calcium control hypothesis indicates that

large levels of calcium (above a plasticity threshold, Θ_p) are thought to lead to t-LTP whereas

more moderate, prolonged levels (between the depression threshold, Θ_{dSTART}, and Θ_{dSTOP})) give

rise to t-LTD and a mid-level range in which t-LTD does not occur (below Θ_{dSTART}) (Karmarkar

and Buonomano, 2002; Rubin et al., 2005; Shouval et al., 2002). A major assumption of these

models is infinite time constants for synaptic variables at resting calcium levels so that the

synaptic changes do not to decay after the presentation of the stimulus (Graupner and Brunel,

2010) - a significant constraint for the stabilization of synaptic changes. A potential solution to

this problem is the degree of local calcium accumulation observed in the activated spines

throughout the t-LTP or t-LTD induction protocol. In fact, these models are consistent,

fundamentally, with our results which show that a pre-post pairing of +7 ms protocol in two

clustered spines, and a pre-post pairing of +13 ms protocol in a single spine gives rise to t-LTP

accompanied by a substantial increase in the intracellular calcium levels following each pairing

repetition, and a significant accumulation of local calcium levels throughout the induction

protocol – likely mediated by the inability to efficiently extrude the local calcium signals in

between each pre-post pairing at this induction frequency (Figure 6 and 7). We propose that the

local spine calcium accumulation we observe provides a new and key variable for the induction

of plasticity, which reduces the constraints imposed by calcium-base models for the stabilization

of synaptic changes (Graupner and Brunel, 2010; Karmarkar and Buonomano, 2002; Rubin et

al., 2005; Shouval et al., 2002).

These changes in local spine and dendritic calcium signals observed when clustered spines are

activated at pairings of + 7ms (Figure 6) or in a single spine activated at pairings of + 13ms

(Figure 7) suggest that perhaps Θ_p can be reached only with ~ 10-20 pre-post pairings (~20-40

seconds). In contrast a pairing protocol of + 7ms in one spine induces no plasticity, and calcium

levels that are effectively extruded in between pairings leading to no calcium accumulation

during the induction protocol (reaching levels below Θ_p and Θ_{dSTART}). These results suggest that

it is not only the amplitude of the local calcium signals after each pairing, but also the local

calcium accumulation during the induction protocol (40 pairings, ~80 seconds) in spines and

dendrites that are required to reach Θ_p for the induction of t-LTP in clustered spines. As

mentioned before, recently it has been demonstrated in vivo that spike timing-induced receptive

field plasticity, with millisecond time delays between visual stimulus (pre) and optogenetic

stimulation in layer 2/3 pyramidal neurons (post), is correlated with increases in synaptic

strength (El-Boustani et al., 2018). These results together with evidence from other in vivo

studies showing that layer 5 pyramidal neurons can spike up to frequencies of 20 Hz during

movement (Schiemann et al., 2015), suggest that our pairing protocol, and findings, are likely

present under certain in vivo conditions and are relevant for plasticity of networks and ultimately

behaviour.

Our results further show that a post-pre protocol of -15 ms in a single spine induces t-LTD and

moderate intracellular calcium signals in spines and parent dendrites after each pairing, without

an evident increase in local calcium accumulation. These results possibly reflect that the calcium

signal generated during the induction protocol passed Θ_{dSTART} and remained for several seconds

in this permissive calcium concentration window – between Θ_{dSTART} and Θ_{dSTOP} – generating

LTD. Activating two clustered spines with the same protocol, however, does not induce

plasticity and gives rise to an apparent smaller initial calcium accumulation than that observed

with the activation of a single spine but with a slow build-up of calcium. These results possibly

reflect that the spine calcium levels crossed Θ_{dSTART} only after > 20 repetitions and then crossed

Θ_{dSTOP} and Θ_{pSTART} after a few (<10) repetitions reaching slightly higher local calcium levels.

Further experiments to prove this calcium control hypothesis of t-LTD induction, with better

temporal detection tools are needed.

These findings presented here are quite remarkable since stimulating just one additional spine

during a STDP protocol can alter the calcium dynamics and the induction of t-LTP and t-LTD.

To our knowledge, this is the first demonstration of the functional relevance that the structural

organization and nearly simultaneous subthreshold activation of only a few clustered inputs in

the dendrites of pyramidal neurons have on plasticity. We propose the term *micro clusters* to

describe this structural and functional modality of synaptic connectivity. In fact, the relevance of

synaptic *micro clusters* on the input/output properties of pyramidal neurons is also supported by

three dimensional electron microscopy and neuronal reconstruction studies that have shown the

presence of postsynaptic innervation of the same axon spaced at less than 10 μm in the basal

dendrites of L2/3 pyramidal neurons from the medial entorhinal cortex (Schmidt et al., 2017), L5

pyramidal neurons from somatosensory cortex (Kasthuri et al., 2015) and in the distal apical tuft

dendrites in stratum lacunosum-moleculare of hippocampal CA1 pyramidal neurons (Bloss et al.,

2018). In addition to having spines innervated by the same axon, it is likely that functional

synaptic *micro clusters* can be gated by the convergence of different axons, which could increase

the computational power of cortical circuits through a multi-neuronal control of synaptic

cooperativity and ultimately the implemented STDP learning rule. Furthermore, it has been

shown that orientation selectivity in visual cortex is correlated with the degree of spatial synaptic

clustering of co-tuned synaptic inputs within the dendritic field (Wilson et al., 2016), and that

functional clusters of dendritic spines separated by less than 10 μm share similar spatial

receptive field properties, spontaneous and sensory-driven activity (Scholl, 2017). Interestingly,

is has been recently shown in the mouse visual cortex that a single axon can contact clustered

synapses in the postsynaptic neuron, but that local clustering is not favoured over widespread

spacing of synaptic inputs (Lee et al., 2016b). In addition, recently it has been demonstrated in

L2/3 pyramidal neurons from mouse primary visual cortex that synaptic *micro clusters* reflects

the interaction of functionally similar inputs - with similar orientation preference - from different

sources – callosal and non-callosal inputs (Lee et al., 2019), providing functional and anatomical

data for the presence of multi-neuronal control of synaptic cooperativity in a short dendritic

segment. Moreover, it has been shown that coactive neighboring synapses drive the maturation

of cluster synapses in CA1 pyramidal neurons (Lee et al., 2016a), suggesting that *micro clusters*

also play an important role in the development and shaping of network connectivity. Taken

together these reported findings and our data suggest that the functional specificity and structural

arrangement of synaptic inputs, distributed or forming *micro clusters* in the dendrites of

pyramidal neurons, are fundamental for guiding the rules for sensory perception, affecting the

STDP learning rule, learning and memory, and ultimately cognition.

METHODS

Brain slice preparation and electrophysiology. Brains from postnatal day 14-21 C57B/6 mice - anesthetized with isoflurane - were removed and immersed in cold (4°C) oxygenated sucrose cutting solution containing (in mM) 27 $NaHCO_3$, 1.5 NaH_2PO_4, 222 Sucrose, 2.6 KCl, 1 $CaCl_2$, and 3 $MgSO_4$. Coronal brain slices (300-μm-thick) of visual cortex were prepared as described (Araya et al., 2006b). Brain slices were incubated for 1/2 hour at 32°C in artificial cerebrospinal fluid (ACSF, in mM: 126 NaCl, 26 $NaHCO_3$, 10 Dextrose, 1.15 NaH_2PO_4, 3 KCl, 2 $CaCl_2$, 2$MgSO_4$) and then transferred to a recording chamber. Electrophysiological recordings were performed in whole-cell current-clamp configuration with MultiClamp 700B amplifiers (Molecular Devices) in L5 pyramidal neurons with a patch electrode (4-7 MΩ) filled with internal solution containing (in mM) 0.1 Alexa Fluo 568, 130 D-Gluconic Acid, 2 $MgCl_2$, 5 KCl, 10 HEPES, 2 MgATP, 0.3 NaGTP, pH 7.4, and 0.4% Biocytin. All experiments were conducted at room temperature (~20-22°C). We did not extend our experiments to include voltage-clamp recordings since recent evidence indicates that the high spine neck resistance can prevent the voltage-clamp control of excitatory synapses and that these measurements can be significantly distorted in spiny neurons (Beaulieu-Laroche and Harnett, 2018).

Two-photon imaging and two-photon uncaging of glutamate. Two-photon imaging was performed with a custom-built two-photon laser scanning microscope, consisting of 1) a Prairie scan head (Bruker) mounted on an Olympus BX51WI microscope with a 60X, 0.9 N.A. water immersion objective; 2) a tunable Ti-Sapphire laser (Chameleon Ultra-II, Coherent, >3 W, 140-fs pulses, 80 MHz repetition rate), 3) two photomultiplier tubes (PMT) for fluorescence detection. Fluorescence images were detected with Prairie software (Bruker).

Fifteen minutes after break-in, two-photon scanning images of basal dendrites were obtained

with 720 nm and low power (~5 mW on sample (i.e., after the objective)) excitation light and

collected with a PMT. Two-photon uncaging of 4-methoxy-7-nitroindolinyl (MNI)-caged L-

glutamate (2.5 mM; Tocris) was performed as previously described (Araya et al., 2006a). This

concentration of MNI-glutamate completely blocked IPSCs (Fino et al., 2009), thus, our results

represent excitatory inputs only. Uncaging was performed at 720 nm (~25-30 mW on sample).

Note that the laser power used for imaging is not sufficient to result in any partial uncaging of

glutamate (Figure S8). Activated spines were mostly on the second and third branch of the basal

dendrites and were on average 40.22 ± 1.62 μm away from the soma (range of distances: ~10 to

80 μm from soma, n = 104 spines, Figure S9). Only spines with a clear head contour and that

were separated by >1 μm from neighboring spines were selected. The location of the uncaging

spot was positioned at ~ 0.3 μm away from the upper edge of the selected spine head (red dot in

figures), which has a spatial resolution of 0.71 and 0.88 μm for one and two spines respectively

(Figure S10). Care was taken to maintain the position of the uncaging spot. After each uncaging

sequence, the spot was repositioned to keep the same distance of ~ 0.3 µm from the edge of the

soma and to avoid artificial potentiation or depression. The uncaging-induced excitatory

postsynaptic potentials (uEPSP) were recorded at the soma of L5 pyramidal neurons.

The kinetics of uEPSPs from the present study are not significantly different (10/90 rate of rise:

0.07 ± 0.014 mV/ms, p=0.92; duration: 115.5 ± 15.3 ms, p=0.65) from those triggered at 37°C

(Araya et al., 2006b).

Spike-timing-dependent plasticity (STDP) induction protocol: To induce t-LTP in single spines,

we used two-photon uncaging of MNI-glutamate (40 times every 2 seconds, with each uncaging

pulse lasting 2 ms), which, after 7 or 13 ms, was followed by a backpropagating action potential

(bAP) (triggered by a brief (10 ms) current injection (0.4 -0.6 nA) in the soma). To induce t-LTD

in single spines, two-photon uncaging of MNI-glutamate (40 times every 2 seconds, with each

uncaging pulse lasting 2 ms) was preceded for -15 or -23 ms by bAP. When we evaluated t-LTP

and t-LTD in two spines, we used similar protocols to those described above, but the spines were

activated with two-photon uncaging of MNI-glutamate sequentially with an onset delay of ~2.1

ms for the second spine. No significant difference was observed in the in 10/90 rise time of the

uEPSPs triggered when one versus two spines were activated (9.05 ± 1.19 ms versus 9.49 ± 0.54

ms, respectively; $p = 0.71$).

To evaluate the morphological and synaptic strength of the activated spines before and after the

STDP induction protocol, we performed 2P imaging, and low frequency 2P uncaging (0.5 Hz) in

single or multiple spines. To establish the time course of the changes in uEPSP amplitude, neck

length and head volume following STDP induction, for each experiment, we interpolated the

data taken at different time points using the *interp1* function in MATLAB (MathWorks) with the

pchip option, which performs a shape-preserving piecewise cubic interpolation. Note that we

constrained this fit so that it terminated with a slope of zero following the last data point. Then,

for each condition, we averaged the uEPSP amplitude, neck length and head volume temporal

traces. The length of the x-axis was set as the time at which the last data point was obtained for

those sets of experiments. Shaded area represents ±SE. To determine at which time the uEPSP

amplitude, neck length and head volume temporal traces are significantly different from baseline,

we binned the temporal traces every 5min and tested whether it was significantly different from

baseline (100%).

The time at which the maximal change in uEPSP was observed after t-LTP and t-LTD induction

was used to calculate the percent change from control, and the percent changes in neck length

and head volume. These analyses are displayed in Figures 1 to 5.

Experimental checkpoints and data analysis: Electrophysiological data were analyzed with

Wavemetrics software (Igor Pro) and MATLAB. The resting membrane potential of the recorded

L5 pyramidal neurons was -58.27 ± 2.08 mV (n = 58 neurons). After taking this measurement,

pyramidal neurons were maintained at − 65 mV in current clamp configuration throughout the

recording session. Only neurons for which the injected current to hold the cell at − 65 mV was <

100 pA were included in this study. For the generation of bAP, only action potentials with

amplitude of > 45 mV were considered for analysis. In most experiments, two control tests (each

consisting of 10 uncaging pulses at 0.5 Hz), spaced by 5 min were performed to assess the

reliability of the uEPSP amplitude. Only experiments for which uEPSP amplitudes were not

significantly different before and after 5 minutes in control conditions were considered for

analysis (less than 10% variation).

Synaptic plasticity was assessed by two parameters: the uEPSP amplitude and the spine

morphology (neck length and head volume). The peak uEPSP amplitude was measured from

each individual uEPSP by taking the peak value and averaging 2 ms before and after using

Wavemetrics (Igor Pro). Only uEPSPs that were > 0.1 mV when one spine was activated in the

control condition (i.e., before the induction of plasticity) were included in the analysis.

Synaptic plasticity was determined by the relative change of uEPSP amplitude (average of 10

uEPSP) measured before and after the STDP protocol. For each experiment, we evaluated

whether the STDP protocol generated potentiation or depression by determining how many

uEPSP data points fell above or below baseline values over the course of the experiment.

Potentiation was defined as the majority of uEPSP amplitude data points measured over time

increasing relative to baseline, and the maximum uEPSP increase was used for statistical test.

Depression was defined as the majority of uEPSP amplitude data points measured over time

decreasing relative to baseline, and the maximum uEPSP decrease was used for statistical test.

The spike timings +7 ms and +13 ms (pre leads post) or -23 ms and -15 ms (post leads pre)

correspond to the delta time offset between the beginning of the uncaging pulse (pre) and the

beginning of the bAP pulse (post) repeated 40 times.

The analysis of spine morphology was performed from z-projections of the whole spine using

ImageJ (Rasband and Bright, 1995) (neck length) and MATLAB (MathWorks) (head volume).

The neck length was measured from the bottom edge of the spine head to the edge of its parent

dendritic shaft using the segmented line tool in ImageJ. We selected mostly spines with a spine

neck longer than 0.2 μm. For those with a shorter neck, we did not report their length for analysis

and statistics due to the diffraction limited resolution of our images. For spines whose necks

shrunk after the STDP protocol below the diffraction limited resolution of our microscope, we

set their length as the minimal measurement of spine neck length reported by Tonnesen et al.,

using STED microscopy (0.157 μm) (Tonnesen et al., 2014). We estimated the relative spine

head volume using the ratio of the maximum spine fluorescence and the maximum fluorescence

observed in the dendrite measured from z-projections of the whole spine (Holtmaat et al., 2005;

Nimchinsky et al., 2004). To obtain the spine volume, we then multiplied this ratio by the PSF of

our microscope (0.11 fL) (Ruttinger et al., 2008). Linear optimization techniques were used to

determine the correlation between EPSP change, neck length change and distance between 2

activated spines. Specifically, the change in uEPSP amplitude was modeled using the following

equation:

$$uEPSP = c_1 \times NL + c_2 \times D \qquad\qquad \textit{Equation 1}$$

Where *uEPSP* and *NL* are the percent change in uEPSP and neck length, respectively, following

the STDP protocol, *D* is the distance between the 2 spines, and c_1, c_2 and c_3 are constant

coefficients. These parameters were estimated using a least squares technique to obtain an

optimal fit of the data that minimized the sum of the residuals squared. The relationship between

inter-spine distance and the percent change in uEPSP was fit with the following exponential

equation:

$$y = \alpha e^{\frac{-x}{\lambda}} + \beta \qquad\qquad \textit{Equation 2}$$

where α, β and λ are constants, y represents the change in uEPSP, and x is the inter-spine

distance.

Calcium imaging: During calcium imaging experiments, we performed whole-cell current-clamp

recordings of L5 pyramidal neurons with a patch electrode containing calcium indicator Fluo-4

(300 µM; Thermo Fisher) and Alexa-594 (100 µM) diluted in an internal solution containing (in

mM) 130 D-Gluconic Acid, 2 $MgCl_2$, 5 KCl, 10 HEPES, 2 MgATP, 0.3 NaGTP, pH 7.4, and

0.4% Biocytin. To perform sequential 2P calcium imaging and 2P uncaging of caged glutamate

in selected spines at one wavelength (810 nm), we used ruthenium-bipyridine-

trimethylphosphine caged glutamate (RuBi-glu, Tocris) (Fino et al., 2009), diluted into the bath

solution for a final concentration of 800 µM. Uncaging of Rubi-glu was performed at 810 nm

(~25-30 mW on sample). The location of the uncaging spot was positioned at ~ 0.3 µm away

from the upper edge of the selected spine head (red dot in Figures 6-8). Changes in calcium were

monitored by imaging 2P calcium signals and detecting the fluorescence with 2 PMTs placed

after wavelength filters (525/70 for green, 595/50 for red). We performed 2P calcium imaging

during five different STDP induction protocols triggered at 0.5Hz: (1) pre-post pairing of +7 ms

in one spine; (2) pre-post pairing of +7 ms in two clustered spines; (3) pre-post pairing of +13 ms

in one spine; (4) post-pre pairing of -15 ms in one spine; (5) post-pre pairing of -15 ms in two

clustered spines. We restricted the image acquisition to a small area (~150 x 150 pixels) which

contained the spine(s) that we uncaged and the shaft. Images were acquired at ~ 30 Hz, averaged

8 times, with 8 μs dwell time. Calcium signals were imaged 500 ms before STDP induction

protocol and right after (4ms) the stimulation for more than 600 ms. We focused our analysis on

the images obtained before and immediately after the stimulation in each pairing repetition. ROI

drawing was performed using custom algorithms (MATLAB; MathWorks). For spine heads, the

ROI was a circle whereas for dendrites it was a polygon. Fluorescence was computed as the

mean of all pixels within the ROI. We quantified the relative change in calcium concentration

($\frac{\Delta F}{F}$) using the following formula:

$$\frac{\Delta F}{F} = \frac{G - G_{baseline}}{R} \qquad\qquad \textit{Equation 3}$$

where G is the fluorescence from the Fluo-4 dye and R is the fluorescence from the Alexa-594

dye. $G_{baseline}$ is the mean of all pixels of Fluo-4 signal within the ROI taken from the first image

at the first stimulation.

Statistical analysis: Statistics were performed with GraphPad Prism 5. Statistical significance

was determined using two-tailed Student's paired *t*-test when we analyzed the maximum change

in uEPSP amplitude after the induction of t-LTP or t-LTD in each experiment and the

concomitant changes in the activated spine morphology. Statistical significance was determined

using one-way repeated measures ANOVA when we analyzed the time course of the uEPSP

amplitude and spine morphological changes after induction of t-LTP or t-LTD with post-hoc

pairwise comparisons using Dunnett's test. *P<0.05; **P<0.01; ***P<0.001.

Pharmacology: Latrunculin A (Lat-A, Tocris Bioscience) was dissolved in DMSO at 1/1000 and

added to the recording chamber containing the brain slice at 100 nM for 15 min before starting

the STDP protocol. PEP1-TGL (Tocris Bioscience) was added in the pipette at 200 μM; after 15

min in whole cell condition, electrophysiological recording and synaptic plasticity experiments

were started. MNI-glutamate (Tocris Bioscience) was diluted in ACSF from stock solution and

bath applied at 2.5 mM. Fresh vials of MNI-glutamate were used for each experiment.

Ethics
Animal experimentation: these studies were performed in compliance with experimental

protocols (13-185, 15-002, 16-011 and 17-012) approved by the *Comité de déontologie de*

l'expérimentation sur les animaux (CDEA) of the University of Montreal.

ACKNOWLEDGEMENTS

We thank P.J Sjöström and A. Kolta for critical discussion and reading of the manuscript, and are grateful to all other members of Roberto Araya's laboratory for kind support. We also thank members of the *Groupe de Recherche sur le système nerveux central* (GRSNC) for support and equipment shearing. This work was funded by the Canadian Institutes of Health Research (CIHR) grant MOP-133711 to R.A., a Canada Foundation for Innovation (CFI) equipment grant *Fonds des leaders* 29970 to R.A., and a Natural Sciences and Engineering Research Council of Canada (NSERC Discovery Grant) grant application No. 418113-2012 (NSERC PIN 392027) to R.A. S.T. was supported in part by a salary support from the GRSNC at Université of Montréal. D.E.M. was supported in part by a postdoctoral fellowship from the Fonds de recherche du Québec – Santé (FRQS).

AUTHOR CONTRIBUTIONS

R.A. conceived the project. S.T. and D.E.M. performed the experiments. S.T. and D.E.M. performed data analyses. S.M-R. performed control experiments. R.A., S.T., and D.E.M. designed experiments. R.A. and D.E.M. wrote the manuscript. R.A. supervised the project. All authors read and approved the contents of the manuscript.

DECLARATION OF INTEREST

The authors declare no competing interests.

REFERENCES

Allison, D.W., Gelfand, V.I., Spector, I., and Craig, A.M. (1998). Role of actin in anchoring postsynaptic receptors in cultured hippocampal neurons: Differential attachment of NMDA versus AMPA receptors. Journal of Neuroscience 18, 2423-2436.

Araya, R. (2014). Input transformation by dendritic spines of pyramidal neurons. Frontiers in neuroanatomy 8.

Araya, R., Eisenthal, K.B., and Yuste, R. (2006a). Dendritic spines linearize the summation of excitatory potentials. Proceedings of the National Academy of Sciences of the United States of America 103, 18799-18804.

Araya, R., Jiang, J., Eisenthal, K.B., and Yuste, R. (2006b). The spine neck filters membrane potentials. Proceedings of the National Academy of Sciences of the United States of America 103, 17961-17966.

Araya, R., Vogels, T.P., and Yuste, R. (2014). Activity-dependent dendritic spine neck changes are correlated with synaptic strength. Proceedings of the National Academy of Sciences of the United States of America.

Artola, A., and Singer, W. (1993). Long-term depression of excitatory synaptic transmission and its relationship to long-term potentiation. Trends Neurosci 16, 480-487.

Ashby, M.C., Maier, S.R., Nishimune, A., and Henley, J.M. (2006). Lateral diffusion drives constitutive exchange of AMPA receptors at dendritic spines and is regulated by spine morphology. The Journal of neuroscience : the official journal of the Society for Neuroscience 26, 7046-7055.

Beaulieu-Laroche, L., and Harnett, M.T. (2018). Dendritic Spines Prevent Synaptic Voltage Clamp. Neuron 97, 75-82 e73.

Bender, V.A., Bender, K.J., Brasier, D.J., and Feldman, D.E. (2006). Two coincidence detectors for spike timing-dependent plasticity in somatosensory cortex. The Journal of neuroscience : the official journal of the Society for Neuroscience 26, 4166-4177.

Bi, G.Q., and Poo, M.M. (1998). Synaptic modifications in cultured hippocampal neurons: dependence on spike timing, synaptic strength, and postsynaptic cell type. The Journal of neuroscience : the official journal of the Society for Neuroscience 18, 10464-10472.

Bloss, E.B., Cembrowski, M.S., Karsh, B., Colonell, J., Fetter, R.D., and Spruston, N. (2018). Single excitatory axons form clustered synapses onto CA1 pyramidal cell dendrites. Nature neuroscience 21, 353-363.

Cummings, J.A., Mulkey, R.M., Nicoll, R.A., and Malenka, R.C. (1996). Ca2+ signaling requirements for long-term depression in the hippocampus. Neuron 16, 825-833.

El-Boustani, S., Ip, J.P.K., Breton-Provencher, V., Knott, G.W., Okuno, H., Bito, H., and Sur, M. (2018). Locally coordinated synaptic plasticity of visual cortex neurons in vivo. Science 360, 1349-1354.

Feldman, D.E. (2000). Timing-based LTP and LTD at vertical inputs to layer II/III pyramidal cells in rat barrel cortex. Neuron 27, 45-56.

Feldman, D.E. (2012). The Spike-Timing Dependence of Plasticity. Neuron 75, 556-571.

Fino, E., Araya, R., Peterka, D.S., Salierno, M., Etchenique, R., and Yuste, R. (2009). RuBi-Glutamate: Two-Photon and Visible-Light Photoactivation of Neurons and Dendritic spines. Frontiers in neural circuits 3, 2.

Fino, E., Paille, V., Cui, Y., Morera-Herreras, T., Deniau, J.M., and Venance, L. (2010). Distinct
coincidence detectors govern the corticostriatal spike timing-dependent plasticity. The Journal of
physiology 588, 3045-3062.
Froemke, R.C., Letzkus, J.J., Kampa, B.M., Hang, G.B., and Stuart, G.J. (2010). Dendritic
synapse location and neocortical spike-timing-dependent plasticity. Frontiers in synaptic
neuroscience 2, 29.
Froemke, R.C., Poo, M.M., and Dan, Y. (2005). Spike-timing-dependent synaptic plasticity
depends on dendritic location. Nature 434, 221-225.
Frost, N.A., Shroff, H., Kong, H.H., Betzig, E., and Blanpied, T.A. (2010). Single-Molecule
Discrimination of Discrete Perisynaptic and Distributed Sites of Actin Filament Assembly within
Dendritic Spines. Neuron 67, 86-99.
Graupner, M., and Brunel, N. (2010). Mechanisms of induction and maintenance of spike-timing
dependent plasticity in biophysical synapse models. Frontiers in computational neuroscience 4.
Hanley, J.G. (2014). Actin-dependent mechanisms in AMPA receptor trafficking. Front Cell
Neurosci 8.
Harnett, M.T., Makara, J.K., Spruston, N., Kath, W.L., and Magee, J.C. (2012). Synaptic
amplification by dendritic spines enhances input cooperativity. Nature 491, 599-602.
Harvey, C.D., and Svoboda, K. (2007). Locally dynamic synaptic learning rules in pyramidal
neuron dendrites. Nature 450, 1195-1200.
Harvey, C.D., Yasuda, R., Zhong, H.N., and Svoboda, K. (2008). The spread of Ras activity
triggered by activation of a single dendritic spine. Science 321, 136-140.
Hashimotodani, Y., Ohno-Shosaku, T., Tsubokawa, H., Ogata, H., Emoto, K., Maejima, T.,
Araishi, K., Shin, H.S., and Kano, M. (2005). Phospholipase Cbeta serves as a coincidence
detector through its Ca2+ dependency for triggering retrograde endocannabinoid signal. Neuron
45, 257-268.
Hayashi-Takagi, A., Yagishita, S., Nakamura, M., Shirai, F., Wu, Y.I., Loshbaugh, A.L.,
Kuhlman, B., Hahn, K.M., and Kasai, H. (2015). Labelling and optical erasure of synaptic
memory traces in the motor cortex. Nature 525, 333-+.
Hayashi, Y., Shi, S.H., Esteban, J.A., Piccini, A., Poncer, J.C., and Malinow, R. (2000). Driving
AMPA receptors into synapses by LTP and CaMKII: requirement for GluR1 and PDZ domain
interaction. Science 287, 2262-2267.
Holcman, D., and Schuss, Z. (2011). Diffusion laws in dendritic spines. Journal of mathematical
neuroscience 1, 10.
Holcman, D., and Triller, A. (2006). Modeling synaptic dynamics driven by receptor lateral
diffusion. Biophysical journal 91, 2405-2415.
Holtmaat, A.J., Trachtenberg, J.T., Wilbrecht, L., Shepherd, G.M., Zhang, X., Knott, G.W., and
Svoboda, K. (2005). Transient and persistent dendritic spines in the neocortex in vivo. Neuron
45, 279-291.
Honkura, N., Matsuzaki, M., Noguchi, J., Ellis-Davies, G.C.R., and Kasai, H. (2008). The
subspine organization of actin fibers regulates the structure and plasticity of dendritic spines.
Neuron 57, 719-729.
Ismailov, I., Kalikulov, D., Inoue, T., and Friedlander, M.J. (2004). The kinetic profile of
intracellular calcium predicts long-term potentiation and long-term depression. The Journal of
neuroscience : the official journal of the Society for Neuroscience 24, 9847-9861.
Jayant, K., Hirtz, J.J., Plante, I.J., Tsai, D.M., De Boer, W.D., Semonche, A., Peterka, D.S.,
Owen, J.S., Sahin, O., Shepard, K.L., and Yuste, R. (2017). Targeted intracellular voltage

recordings from dendritic spines using quantum-dot-coated nanopipettes. Nature nanotechnology
12, 335-342.

Karmarkar, U.R., and Buonomano, D.V. (2002). A model of spike-timing dependent plasticity:
one or two coincidence detectors? J Neurophysiol 88, 507-513.

Kasthuri, N., Hayworth, K.J., Berger, D.R., Schalek, R.L., Conchello, J.A., Knowles-Barley, S.,
Lee, D., Vazquez-Reina, A., Kaynig, V., Jones, T.R., *et al*. (2015). Saturated Reconstruction of a
Volume of Neocortex. Cell 162, 648-661.

Kim, C.H., and Lisman, J.E. (1999). A role of actin filament in synaptic transmission and long-
term potentiation. Journal of Neuroscience 19, 4314-4324.

Kusters, R., Kapitein, L.C., Hoogenraad, C.C., and Storm, C. (2013). Shape-induced asymmetric
diffusion in dendritic spines allows efficient synaptic AMPA receptor trapping. Biophysical
journal 105, 2743-2750.

Lee, K.F., Soares, C., Thivierge, J.P., and Beique, J.C. (2016a). Correlated Synaptic Inputs Drive
Dendritic Calcium Amplification and Cooperative Plasticity during Clustered Synapse
Development. Neuron 89, 784-799.

Lee, K.S., Vandemark, K., Mezey, D., Shultz, N., and Fitzpatrick, D. (2019). Functional
Synaptic Architecture of Callosal Inputs in Mouse Primary Visual Cortex. Neuron.

Lee, W.C., Bonin, V., Reed, M., Graham, B.J., Hood, G., Glattfelder, K., and Reid, R.C.
(2016b). Anatomy and function of an excitatory network in the visual cortex. Nature 532, 370-
374.

Lisman, J. (1989). A mechanism for the Hebb and the anti-Hebb processes underlying learning
and memory. Proceedings of the National Academy of Sciences of the United States of America
86, 9574-9578.

Lynch, G., Larson, J., Kelso, S., Barrionuevo, G., and Schottler, F. (1983). Intracellular
injections of EGTA block induction of hippocampal long-term potentiation. Nature 305, 719-
721.

Malenka, R.C., Kauer, J.A., Zucker, R.S., and Nicoll, R.A. (1988). Postsynaptic calcium is
sufficient for potentiation of hippocampal synaptic transmission. Science 242, 81-84.

Markram, H., Lubke, J., Frotscher, M., Roth, A., and Sakmann, B. (1997). Physiology and
anatomy of synaptic connections between thick tufted pyramidal neurones in the developing rat
neocortex. J Physiol-London 500, 409-440.

Matsuzaki, M., Ellis-Davies, G.C.R., Nemoto, T., Miyashita, Y., Iino, M., and Kasai, H. (2001).
Dendritic spine geometry is critical for AMPA receptor expression in hippocampal CA1
pyramidal neurons. Nature neuroscience 4, 1086-1092.

Matsuzaki, M., Honkura, N., Ellis-Davies, G.C.R., and Kasai, H. (2004). Structural basis of
long-term potentiation in single dendritic spines. Nature 429, 761-766.

Nakamura, T., Barbara, J.G., Nakamura, K., and Ross, W.N. (1999). Synergistic release of Ca2+
from IP3-sensitive stores evoked by synaptic activation of mGluRs paired with backpropagating
action potentials. Neuron 24, 727-737.

Nevian, T., Larkum, M.E., Polsky, A., and Schiller, J. (2007). Properties of basal dendrites of
layer 5 pyramidal neurons: a direct patch-clamp recording study. Nature neuroscience 10, 206-
214.

Nevian, T., and Sakmann, B. (2006). Spine Ca2+ signaling in spike-timing-dependent plasticity.
Journal of Neuroscience 26, 11001-11013.

Nimchinsky, E.A., Yasuda, R., Oertner, T.G., and Svoboda, K. (2004). The number of glutamate
receptors opened by synaptic stimulation in single hippocampal spines. The Journal of
neuroscience : the official journal of the Society for Neuroscience 24, 2054-2064.
Nishiyama, J., and Yasuda, R. (2015). Biochemical Computation for Spine Structural Plasticity.
Neuron 87, 63-75.
Oh, W.C., Hill, T.C., and Zito, K. (2013). Synapse-specific and size-dependent mechanisms of
spine structural plasticity accompanying synaptic weakening. Proceedings of the National
Academy of Sciences of the United States of America 110, E305-312.
Penn, A.C., Zhang, C.L., Georges, F., Royer, L., Breillat, C., Hosy, E., Petersen, J.D., Humeau,
Y., and Choquet, D. (2017). Hippocampal LTP and contextual learning require surface diffusion
of AMPA receptors. Nature 549, 384-388.
Ramachandran, B., and Frey, J.U. (2009). Interfering with the actin network and its effect on
long-term potentiation and synaptic tagging in hippocampal CA1 neurons in slices in vitro. The
Journal of neuroscience : the official journal of the Society for Neuroscience 29, 12167-12173.
Rasband, W.S., and Bright, D.S. (1995). Nih Image - a Public Domain Image-Processing
Program for the Macintosh. Microbeam Anal 4, 137-149.
Richardson, R.J., Blundon, J.A., Bayazitov, I.T., and Zakharenko, S.S. (2009). Connectivity
patterns revealed by mapping of active inputs on dendrites of thalamorecipient neurons in the
auditory cortex. The Journal of neuroscience : the official journal of the Society for
Neuroscience 29, 6406-6417.
Rubin, J.E., Gerkin, R.C., Bi, G.Q., and Chow, C.C. (2005). Calcium time course as a signal for
spike-timing-dependent plasticity. J Neurophysiol 93, 2600-2613.
Ruttinger, S., Buschmann, V., Kramer, B., Erdmann, R., Macdonald, R., and Koberling, F.
(2008). Comparison and accuracy of methods to determine the confocal volume for quantitative
fluorescence correlation spectroscopy. J Microsc 232, 343-352.
Schiemann, J., Puggioni, P., Dacre, J., Pelko, M., Domanski, A., van Rossum, M.C., and Duguid,
I. (2015). Cellular mechanisms underlying behavioral state-dependent bidirectional modulation
of motor cortex output. Cell Rep 11, 1319-1330.
Schmidt, H., Gour, A., Straehle, J., Boergens, K.M., Brecht, M., and Helmstaedter, M. (2017).
Axonal synapse sorting in medial entorhinal cortex. Nature 549, 469-475.
Scholl, B., Wilson, D.E., Fitzpatrick, D. (2017). Local Order within Global Disorder: Synaptic
Architecture of Visual Space. In Neuron, pp. 1127-1138.
Shim, S.H., Xia, C.L., Zhong, G.S., Babcock, H.P., Vaughan, J.C., Huang, B., Wang, X., Xu, C.,
Bi, G.Q., and Zhuang, X.W. (2012). Super-resolution fluorescence imaging of organelles in live
cells with photoswitchable membrane probes. Proceedings of the National Academy of Sciences
of the United States of America 109, 13978-13983.
Shouval, H.Z., Bear, M.F., and Cooper, L.N. (2002). A unified model of NMDA receptor-
dependent bidirectional synaptic plasticity. Proceedings of the National Academy of Sciences of
the United States of America 99, 10831-10836.
Sjostrom, P.J., and Hausser, M. (2006). A cooperative switch determines the sign of synaptic
plasticity in distal dendrites of neocortical pyramidal neurons. Neuron 51, 227-238.
Sjostrom, P.J., Turrigiano, G.G., and Nelson, S.B. (2001). Rate, timing, and cooperativity jointly
determine cortical synaptic plasticity. Neuron 32, 1149-1164.
Sjostrom, P.J., Turrigiano, G.G., and Nelson, S.B. (2003). Neocortical LTD via coincident
activation of presynaptic NMDA and cannabinoid receptors. Neuron 39, 641-654.

Tanaka, J.I., Horiike, Y., Matsuzaki, M., Miyazaki, T., Ellis-Davies, G.C.R., and Kasai, H.
(2008). Protein synthesis and neurotrophin-dependent structural plasticity of single dendritic
spines. Science 319, 1683-1687.
Tonnesen, J., Katona, G., Rozsa, B., and Nagerl, U.V. (2014). Spine neck plasticity regulates
compartmentalization of synapses. Nature neuroscience 17, 678-685.
Wilson, D.E., Whitney, D.E., Scholl, B., and Fitzpatrick, D. (2016). Orientation selectivity and
the functional clustering of synaptic inputs in primary visual cortex. Nature neuroscience 19,
1003-1009.
Yang, S.N., Tang, Y.G., and Zucker, R.S. (1999). Selective induction of LTP and LTD by
postsynaptic [Ca2+]i elevation. J Neurophysiol 81, 781-787.
Zhang, L.I., Tao, H.W., Holt, C.E., Harris, W.A., and Poo, M.M. (1998). A critical window for
cooperation and competition among developing retinotectal synapses. Nature 395, 37-44.
1034

Figure 1: **Induction of t-LTP in single dendritic spines**. (A) Experimental protocol for the induction of t-LTP with pre-post pairings (two-photon (2P) uncaging of glutamate followed by a bAP) of +7 and +13 ms in single dendritic spines (sp). (B) Example of a t-LTP protocol with pre-post pairing protocol of + 13 ms. (B.1) Average uEPSP response recorded via a whole-cell pipette at the soma of L5 pyramidal neurons before (Control) and after the induction of t-LTP in a selected dendritic spine located in basal dendrites (B.2). Traces in B.1 correspond to an average of 10 depolarizations generated by 2P uncaging over the indicated spine (B.2 red dot) before (black trace) and after the induction of t-LTP (red trace). (C.1) Time course of uEPSP amplitude (black line), neck length (red line) and spine head volume (blue line) changes over the course of 40 min following STDP induction in individual spines at a pre-post timing of + 13 ms. *ns*, not significant; *P < 0.05; **P < 0.01; ***P < 0.001, one-way repeated measures ANOVA followed by post hoc Dunnet's test. (C.2) Maximum changes in uEPSP amplitude (black bar and dots) and concomitant changes in neck length (red bar and dots) and head volume (blue bar and dots) of the activated spine after the induction of t-LTP at a pre-post timing of +13 ms. *P < 0.05, ***P < 0.001, paired t-test. (D.1) Time course of uEPSP amplitude (black line), neck length (red line) and spine head volume (blue line) changes over the course of ~40 min following STDP induction in individual spines at a pre-post timing of + 7 ms. *ns*, not significant; *P < 0.05, one-way repeated measures ANOVA followed by post hoc Dunnet's test. (D.2) Maximum changes in uEPSP amplitude (black bar and dots) and corresponding changes in neck length (red bar and dots) and head volume (blue bar and dots) of the activated spine after the induction of t-LTP at a pre-post timing of +7 ms. Shaded light area in C.1 and D.1, and error bars in C.2 and D.2 represent ±SEM. Time 0 in C.1 and D.1 is the time when the 40 pre-post repetitions of the induction protocol were completed.

Figure 2: **Induction of t-LTD in single dendritic spines**. (A) Experimental protocol for the

induction of t-LTD with post-pre pairings protocols of -15 and -23 ms in single dendritic spines.

(B) Example of a t-LTD protocol with post-pre pairings protocol of -15 ms. (B.1) Average

uEPSP response recorded via a whole-cell pipette at the soma of L5 pyramidal neurons before

(Control) and after the induction of t-LTD in selected dendritic spines located in basal dendrites

(B.2). Traces in B.1 correspond to an average of 10 uEPSP generated by the 2P uncaging of

glutamate at the indicated spine (B.2, red dot) before (black trace) and after the induction of t-

LTD (red trace). (C.1) Time course of uEPSP amplitude (black line), neck length (red line) and

spine head volume (blue line) changes over the course of ~35 min following STDP induction in

individual spines at a post-pre timing of -15 ms. *ns*, not significant; ***P < 0.001, one-way

repeated measures ANOVA followed by post hoc Dunnet's test. (C.2) Maximum changes in

uEPSP amplitude (black bar and dots) and concomitant changes in neck length (red bar and dots)

and head size of the activated spine (blue bar and dots) after the induction of t-LTD at a post-pre

timing of -15 ms. **P < 0.01, paired t test. (D.1) Time course of uEPSP amplitude (black line),

neck length (red line) and spine head volume (blue line) changes over the course of ~ 40 min

following STDP induction in individual spines at a post-pre timing of -23 ms. *ns*, not significant,

one-way repeated measures ANOVA followed by post hoc Dunnet's test. (D.2) Maximum

changes in uEPSP amplitude (black bar and dots) and corresponding changes in neck length (red

bar and dots) and head volume (blue bar and dots) of the activated spine after the induction of t-

LTD at a post-pre timing of -23 ms. Shaded light area in C.1 and D.1, and error bars in C.2 and

D.2 represent ±SEM. Time 0 in C.1 and D.1 is the time when the 40 pre-post repetitions of the

induction protocol were completed. (E) STDP learning rule for single dendritic spines: Plot

illustrating the maximum changes in uEPSP amplitude (black data points), and concomitant

changes in neck length (red data points) and head volume (blue data points) after the induction of

t-LTP (temporal offset +13 ms and +7 ms) and t-LTD (temporal offset -15 ms and -23 ms). *P <

0.05, **P < 0.01, paired t test. (F) Diagram indicating the uEPSP amplitude and spine neck

morphological changes observed after the induction of t-LTP and t-LTD.

Figure 3: **Induction of t-LTP in clustered and distributed dendritic spines** (A) Experimental protocol for the induction of t-LTP (pre-post timing of + 7 ms) in two clustered dendritic spines (< 5 μm distance between spines). (B) Example of one experiment where two neighbouring dendritic spines were activated with a pre-post t-LTP protocol. (B.1) Average uEPSP response recorded via a whole-cell pipette at the soma of L5 pyramidal neurons before (Control) and after the induction of t-LTP in two selected dendritic spines located in a basal dendrite (red dots, B.2). Traces in B.1 correspond to an average of 10 uEPSP generated by the 2P uncaging at the indicated spines (B.2) before (black trace) and after the induction of t-LTP (red trace). (C) Time course of uEPSP amplitude (black line), neck length (red line) and spine head volume (blue line) changes over the course of ~25 min following STDP induction in clustered spines (< 5μm distance between spines) at a pre-post timing of + 7 ms. *ns*, not significant; **P < 0.01, ***P < 0.001, one-way repeated measures ANOVA followed by post hoc Dunnet's test. (D) Experimental protocol for the induction of t-LTP (pre-post timing of + 7 ms) in two distributed spines (> 5μm distance between spines). (E) Induction of t-LTP (increase in uEPSP amplitude, more than 100%) in two nearly simultaneously activated spines at a pre-post timing of + 7 ms is dependent on inter-spine distance. Data points represent mean. We use the extracted value of λ as a boundary between clustered and distributed spines. (F) Example of one experiment where two distributed dendritic spines (> 5μm distance between spines) were activated with a pre-post timing of + 7ms. (F.1) Average uEPSP response recorded via a whole-cell pipette at the soma of L5 pyramidal neurons before (Control) and after the induction of t-LTP in two selected dendritic spines located in a basal dendrite (red dots, F.2). Traces in F.1 correspond to an average of 10 uEPSP generated by the 2P uncaging at the indicated spines (F.2) before (black trace) and after the induction of t-LTP (red trace). (G) Time course of uEPSP amplitude (black line), neck length

(red line) and spine head volume (blue line) changes over the course of ~30 min following STDP

induction in distributed spines (> 5μm distance between spines) at a pre-post timing of + 7 ms.

ns, not significant; **P < 0.01, ***P < 0.001, one-way repeated measures ANOVA followed by

post hoc Dunnet's test. (H) Pooled data of the maximum changes in uEPSP amplitude (black bar

and dots) and concomitant changes in neck length (red bar and dots) and head volume (blue bar

and dots) of individual (1sp), clustered (< 5μm) and distributed (> 5μm) spines (2sp) after the

induction of t-LTP at a pre-post timing of +7ms. **P < 0.01, paired t test. Shaded light area in

B.1, F.1C and G and error bars in H represent ±SEM. Time 0 in C and G is the time when the 40

pre-post repetitions of the induction protocol are completed. NL = neck length, HV = head

volume.

Figure 4: **Molecular mechanisms responsible for the induction of t-LTP**. (A) Experimental

protocol for the induction of t-LTP (pre-post timing of + 7 ms) in two clustered dendritic spines

(< 5 µm) in the presence of PEP1-TGL (200 µM). (B) An example of one experiment where two

neighbouring dendritic spines were activated with a pre-post t-LTP protocol in the presence of

PEP1-TGL. (B.1) Average uEPSP recorded via a whole-cell pipette at the soma of L5 pyramidal

neurons before (Control) and after the induction of t-LTP in two selected dendritic spines located

in basal dendrites (B.2). Traces in B.1 correspond to an average of 10 uEPSP generated by the 2P

uncaging of glutamate at the two indicated spines (red dots, B.2) before (black trace) and after

the induction of t-LTP in the presence of PEP1-TGL (red traces). (C.1) Time course of uEPSP

amplitude (black line), neck length (red line) and spine head volume (blue line) changes over the

course of ~28 min following STDP induction in clustered spines at a pre-post timing of + 7 ms in

the presence of PEP1-TGL. *ns*, not significant; **P < 0.01, ***P < 0.001, one-way repeated

measures ANOVA followed by post hoc Dunnet's test. (C.2) Pooled data of the maximum

changes in uEPSP amplitude (black bar and dots) and concomitant changes in neck length (red

bar and dots) and head volume (blue bar and dots) of the activated clustered spines after the

induction of t-LTP at a pre-post timing of +7ms in control conditions (Cont) and in the presence

of PEP1-TGL (PEP1-TGL). *P < 0.05, **P < 0.01, paired t test. Time 0 is the time when the 40

pre-post repetitions of the induction protocol were completed. (D) Experimental protocol for the

induction of t-LTP (pre-post timing of + 7 ms) in two clustered dendritic spines (< 30µm) in the

presence Latrunculin-A (Lat-A, 100nM). (E) Example of one experiment where two

neighbouring dendritic spines were activated with a pre-post t-LTP protocol in the presence of

Lat-A. (E.1) Average uEPSP recorded via a whole-cell pipette at the soma of L5 pyramidal

neurons before (Control) and after the induction of t-LTP in two selected dendritic spines (red

dots, E.2) located in basal dendrites (E.2). Traces in E.1 correspond to an average of 10 uEPSP

generated by the 2P uncaging over the two indicated spines (red dots, E.2) before (black trace)

and after the induction of t-LTP in the presence of Lat-A (red trace). (F.1) Time course of uEPSP

amplitude (black line), neck length (red line) and spine head volume (blue line) changes over the

course of ~12 min following STDP induction in clustered spines at a pre-post timing of + 7 ms in

the presence of Lat-A. *ns*, not significant; **P < 0.01, ***P < 0.001, one-way repeated measures

ANOVA followed by post hoc Dunnet's test. (F.2) Pooled data of the maximum changes in

uEPSP amplitude (black bar and dots) and concomitant changes in neck length (red bar and dots)

and head volume (blue bar and dots) of the activated clustered spines after the induction of t-LTP

at a pre-post timing of +7 ms in control conditions (Cont) and in the presence of Lat-A (Lat-A).

**P < 0.01, ***P < 0.001, paired t test. NL = neck length, HV = head volume. Shaded light area

in C.1 and F.1 and error bars in C.2 and F.2 represent ±SEM. Time 0 is the time when 40 pre-

post repetitions of the induction protocol were completed.

Figure 5. Induction of t-LTD in clustered and distributed dendritic spines: (A) Experimental

protocol for the induction of t-LTD protocol (pairings of - 15 ms) in two clustered dendritic

spines (< 40 μm). (B) An example of one experiment where two neighbouring dendritic spines

were activated with a post-pre t-LTP protocol of -15 ms. (B.1) Average uEPSP recorded via a

whole-cell pipette at the soma of L5 pyramidal neurons before (Control) and after the induction

of t-LTD in two selected dendritic spines (red dots, B.2) located in basal dendrites. Traces in B.1

correspond to an average of 10 uEPSP generated by 2P uncaging of glutamate at the indicated

spines before (black trace) and after the induction of t-LTD (red trace). (C) Time course of

uEPSP amplitude (black line), the neck length (red line) and spine head volume (blue line) of the

activated clustered spines after the induction of t-LTD at pairings of -15 ms. *ns*, not significant;

*$P < 0.05$, one-way repeated measures ANOVA followed by post hoc Dunnet's test. (D)

Experimental protocol for the induction of t-LTD (post-pre timing of -15 ms) in two distributed

spines (> 40 μm distance between spines). (E) t-LTD recovery (decrease in uEPSP amplitude,

less than 100%) is dependent on inter-spine distance. Data points represent mean. We use the

extracted value of λ as a boundary between clustered and distributed spines. (F) Example of one

experiment where two distributed dendritic spines (> 40 μm distance between spines) were

activated with a post-pre timing -15 ms. (F.1) Average uEPSP response recorded via a whole-

cell pipette at the soma of L5 pyramidal neurons before (Control, underneath image, F.2) and 3

min after the induction of t-LTD (F.2, over imposed image) in two selected dendritic spines

located in basal dendrites. (F.2) Inset shows a low magnification image of the recorded neuron

with the marked location of the selected spines. Traces in F.1 correspond to an average of 10

uEPSP generated by 2P uncaging at the indicated spines (red dots, F.2) before (black trace) and

after the induction of t-LTD (red trace). (G) Time course of uEPSP amplitude (black line), the

neck length (red line) and spine head volume (blue line) of the activated distributed spines (> 40

μm) after the induction of t-LTD at pairings of -15 ms. *ns*, not significant; $*P < 0.05$, $**P < 0.01$,

one-way repeated measures ANOVA followed by post hoc Dunnet's test. (H) Pooled data of the

maximum changes in uEPSP amplitude (black bar and dots) and concomitant changes in neck

length (red bar and dots) and head volume (blue bar and dots) of individual (1sp), clustered (2sp

at < 40 μm) and distributed spines (2sp at > 40 μm) after the induction of t-LTD at a post-pre

timing of -15 ms. $*P < 0.05$, $**P < 0.01$, paired t test. Time 0 in C and G is the time when the 40

pre-post repetitions of the induction protocol were completed. NL = neck length, HV = head

volume.

Figure 6. Calcium dynamics in single and two clustered spines during a pre-post pairing protocol of + 7ms. (A.1) Single 2P images of a spine and dendrite from a L5 pyramidal neuron loaded with Alexa 594 (100uM) and Fluo4 (300uM). Red ellipses and blue polygons indicate the ROIs selected for the calcium signal analysis. (A.2) Two photon calcium signal images before (left panels) and after (right panels) a pre-post pairing protocol (+7 ms). The 1^{st}, 2^{nd}, and 40^{th} repetitions of the pairing protocol are shown here. The change in calcium fluorescence from baseline (ΔF) is color coded. (B) Population averages of the calcium signals ($\Delta F/F$) measured in spines (B.1) and dendrites (B.2) before the pairing protocol performed in 1 spine (left panels) and 2 spines (middle panels). The right panels shows the superimposed $\Delta F/F$ population averages in 1 spine (black lines) and 2 spines (green lines). (C.1) Single 2P images of two clustered spines and dendrite from a L5 pyramidal neuron loaded with Alexa 594 (100μM) and Fluo4 (300μM). Red ellipses and blue polygons indicate the ROIs selected for the calcium signal analysis. (C.2) Two photon calcium signal images before (left panel) and after (right panels) a pre-post pairing protocol (+ 7ms). The 1^{st}, 2^{nd}, and 40^{th} repetitions of the pairing protocol are shown here. The change in calcium fluorescence from baseline (ΔF) is color coded. (D) Population averages of the calcium signals ($\Delta F/F$) measured in spines (D.1) and dendrites (D.2) after the pairing protocol performed in one spine (left panels) and two spines (middle panels). The right panels shows the superimposed $\Delta F/F$ population averages in one spine (black lines) and two spines (green lines). *ns*, not significant; *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$, one-way repeated measures ANOVA followed by post hoc Dunnet's test.

Figure 7. Calcium dynamics in single spines during a pre-post pairing protocol of +13ms.
(A.1) Single 2P images of a spine and dendrite from a L5 pyramidal neuron loaded with Alexa 594 (100uM) and Fluo4 (300uM). Red ellipses and blue polygons indicate the ROIs selected for the calcium signal analysis. (A.2) Two photon calcium signal images before (left panels) and after (right panels) a pre-post pairing protocol of +13 ms. The 1^{st}, 2^{nd}, and 40^{th} repetitions of the pairing protocol are shown here. The change in calcium fluorescence from baseline (ΔF) is color coded. (B) Population averages of the calcium signals ($\Delta F/F$) measured in spines (B.1 and C.1) and dendrites (B.2 and C.2) before (B) and after (C) the pairing protocol. *ns*, not significant; $*P < 0.05$; $**P < 0.01$; $***P < 0.001$, one-way repeated measures ANOVA followed by post hoc Dunnet's test.

Figure 8. Calcium dynamics in single and two clustered spines during post-pre pairing

protocol. (A1) Single 2P images of a spine and dendrite from a L5 pyramidal neuron loaded

with Alexa 594 (100uM) and Fluo4 (300uM). Red ellipses and blue polygons indicate the ROIs

selected for the calcium signal analysis. (A.2) Two photon calcium signal images before (left

panels) and after (right panels) a post-pre pairing protocol (-15 ms). The 1^{st}, 2^{nd}, and 40^{th}

repetitions of the pairing protocol are shown here. The change in calcium fluorescence from

baseline (ΔF) is color coded. (B) Population averages of the calcium signals ($\Delta F/F$) measured in

spines (B.1) and dendrites (B.2) before the pairing protocol performed in one spine (left panels)

and two spines (middle panels). The right panels shows the superimposed $\Delta F/F$ population

averages in one spine (black lines) and two clustered spines (green lines). (C.1) Single 2P images

of two clustered spines and dendrite from a L5 pyramidal neuron loaded with Alexa 594

(100μM) and Fluo4 (300μM). Red ellipses and blue polygons indicate the ROIs selected for the

calcium signal analysis. (C.2) Two photon calcium signal images before (left panel) and after

(right panels) a pre-post pairing protocol (-15 ms). The 1^{st}, 2^{nd}, and 40^{th} repetitions of the pairing

protocol are shown here. The change in calcium fluorescence from baseline (ΔF) is color coded.

(D) Population averages of the calcium signals ($\Delta F/F$) measured in spines (D.1) and dendrites

(D.2) after the pairing protocol performed in one spine (left panels) and two spines (middle

panels). The right panels shows the superimposed $\Delta F/F$ population averages in 1 spine (black

lines) and 2 spines (green lines). *ns*, not significant; *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$, one-

way repeated measures ANOVA followed by post hoc Dunnet's test.

1251

Figure 9. STDP learning rule for single distributed and clustered dendritic spines. STDP learning rule in the basal dendrites of L5 pyramidal neurons as a function of the structural organization of excitatory inputs in basal dendrites of L5 pyramidal neurons. Note how STDP in single, or distributed spines (separated by > 40 μm), follow a bidirectional Hebbian STDP rule. Importantly, our model indicate that the induction of t-LTD can be disrupted by the co-activation of two clustered spines separated by < 40 μm, and t-LTP enhanced by the co-activation of two clustered spines separated by < 40 μm by synaptic cooperativity. We propose that synaptic cooperativity generates local dendritic depolarization high enough to disrupt bidirectional STDP, leading to STDP that only encompasses LTP.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A.1

A.2   Pairing protocol pre-post (+7 ms)

C.1

C.2   Pairing protocol pre-post (+7 ms)

Rep #1

Rep #2

Rep #40

1μm

ΔF 0 — 919

ΔF 0 — 1811

B.1   Spine before stim
1 spine   2 spines

B.2   Dendrite before stim
1 spine   2 spines

D.1   Spine after stim
1 spine   2 spines

D.2   Dendrite after stim
1 spine   2 spines

ΔF/F

Repetition

Figure 7

A.1

Rep #1

Rep #2

Rep #40

2 µm

A.2

Pairing protocol pre-post (+13 ms)

25   ΔF   623

B.1 Spine before stim

C.1 Spine after stim

B.2 Dendrite before stim

C.2 Dendrite after stim

ΔF/F

Repetition

Figure 8

Figure 9