A spike timing-dependent plasticity rule for single, clustered and distributed dendritic spines

Sabrina Tazerart[1#], Diana E. Mitchell[1#], Soledad Miranda-Rottmann[1], and Roberto Araya[1*]

[1]Department of Neurosciences, Faculty of Medicine, University of Montreal

[#]These authors contributed equally to this work

[*]Corresponding author

Correspondence: Roberto Araya Ph.D. Assistant Professor, Department of Neurosciences, Faculty of Medicine, University of Montreal, C.P. 6128 Succursale Centre-Ville, Montreal, Quebec H3C 3J7, T: 514 343 7815, e-mail: roberto.araya@umontreal.ca

ABSTRACT

Dendritic spines can undergo structural remodeling, and are the preferential site for the induction of long-term potentiation (LTP) and long-term depression (LTD). In a variant of LTP and LTD, known as spike-timing dependent plasticity (STDP), the sign and magnitude of the change in synaptic strength depends on the timing between the spikes of two connected neurons. Although STDP has been extensively studied in cortical pyramidal neurons, the precise structural organization of excitatory inputs that supports STDP, as well as the structural, molecular and functional properties of dendritic spines during STDP remain unknown. Here we developed a spine STDP protocol, in which two-photon glutamate uncaging over single or multiple spines from the basal dendrites of layer 5 pyramidal

neurons, which mimics presynaptic release of glutamate (pre), was paired with somatically

generated postsynaptic spikes (post). We found that the induction of STDP in single spines

follows a classical Hebbian STDP rule, where pre-post pairings at timings that trigger LTP

(t-LTP) produce shrinkage of the activated spine neck and a concomitant increase in its

synaptic strength; and post-pre pairings that trigger LTD (t-LTD) decrease synaptic

strength without affecting the activated spine shape. Furthermore, we tested whether the

single spine-Hebbian STDP rule could be affected by the activation of neighboring

(clustered) or distant (distributed) spines. Our results show that the induction of t-LTP in

two clustered spines (< 5 μm apart) enhances LTP via a mechanism that is accompanied by

local spine calcium increases that accumulates during the induction protocol, and that

requires actin polymerization-dependent neck shrinkage, which permits AMPA receptor

transport to the spine head and insertion into the postsynaptic density (PSD). Moreover, the

induction of t-LTD is disrupted when two clustered spines are activated, with no calcium

accumulation in spines or dendrites, but can be recovered if the activated spines are

separated by > 40 μm. These results indicate that the induction of STDP in single, or

distributed spines, follow a Hebbian STDP rule. Interestingly, synaptic cooperativity,

induced by the co-activation of only two clustered spines and the local spatio-temporal

summation of clustered synaptic inputs, provides local dendritic depolarization and local

calcium signals that are sufficient to disrupt t-LTD and extend the temporal window for the

induction of t-LTP, leading to STDP only encompassing LTP.

Dendritic spines, the main recipient of excitatory information in the brain [1], are tiny protrusions

with a small head (~1 μm in diameter and <1 fL volume) separated from the dendrite by a slender

neck. Spines can undergo structural remodeling that is tightly coupled with synaptic function [1-4],

and are the preferential site for the induction of long-term potentiation (LTP) [4-6] and long-term

depression (LTD) [7], thought to be the underlying mechanisms for learning and memory in the brain

[8]. A variation of LTP and LTD has been described in pyramidal neurons that involves the pairing

of pre- and postsynaptic action potentials, known as spike-timing dependent plasticity (STDP) [9,

10]. In this process, the timing between pre- and postsynaptic action potentials modulates synaptic

strength, triggering LTP or LTD [10]. The sign and magnitude of the change in synaptic strength

depends on the relative timing between spikes of two connected neurons (the pre- and postsynaptic

neuron [11]). The STDP learning rules and their dependency on postsynaptic dendritic depolarization

[12, 13], firing rate [12], and somatic distance of excitatory inputs [13-15] have been extracted from studies

using connected neuronal pairs or by using extracellular stimulating electrodes, but the precise

location and structural organization of excitatory inputs capable of supporting STDP at its minimal

functional unit – the dendritic spine – are unknown.

Activity-dependent spine morphological changes (spine head [4], neck [2], or both [16]) have been

correlated with changes in synaptic strength in cortical pyramidal neurons by mechanisms

involving biochemical and electrical spine changes [1, 6]. Thus, here we asked what patterns of

activity and structural organization of excitatory synaptic inputs support the generation of t-LTP

and t-LTD, and which morphological, biophysical and molecular changes observed in dendritic

spines can account for the induction of t-LTP and t-LTD?

To induce synapse-specific STDP we developed a protocol whereby two-photon (2P) uncaging of

a caged glutamate (MNI-glutamate [3]) at a single spine – to mimic synaptic release – is preceded

or followed in time (STDP timing window [10]) by a backpropagating action potential (bAP) to

trigger t-LTP (Figure 1A, pre-post) or t-LTD (Figure 2A, post-pre), respectively. Postsynaptic

spikes were triggered by current injection via a whole-cell recording pipette.   Two-photon

uncaging of a caged glutamate at a single spine triggered excitatory postsynaptic potentials

(uncaging(u)EPSP) that were recorded in the soma of layer 5 (L5) pyramidal neurons before and

after the induction of STDP, while the morphology of the activated spine neck and head was

monitored (Figure 1A and 2A). To induce synapse-specific STDP and monitor calcium levels in

the activated spines and parent dendrites we developed a protocol during which we perform nearly

simultaneous 2P uncaging of glutamate and 2P calcium imaging of the activated spines and nearby

dendrites.

Here, we provide evidence showing that the induction of STDP in single or distributed spines

follows a bidirectional Hebbian STDP rule. Furthermore, we show that synaptic cooperativity,

induced by the co-activation of only two clustered spines, disrupt t-LTD (< 40 µm distance

between spines) and extend the temporal window for the induction of t-LTP (< 5 µm distance

between spines) via the generation of differential local calcium signals leading to an STDP rule

for clustered inputs only embracing LTP.

92   **RESULTS**

Induction of t-LTP in single dendritic spines

To induce t-LTP, we used a repetitive spike-timing protocol (40 times, 0.5 Hz) in which 2P

uncaging of glutamate at a single spine was closely followed in time (+7 or +13 ms later, see

Methods section) by a bAP (Figure 1A). We evaluated spine morphology and uEPSP amplitude

for 40 min following STDP induction to establish the time course of STDP at individual synapses

(Figure 1C.1 and D.1). In addition, the maximum uEPSP change and concomitant changes in spine

morphology observed in each experiment are shown (Figure 1C.2 and D.2).

A repetitive pre-post pairing protocol of +13 ms reliably induced t-LTP (significant increase in the

uEPSP amplitude over time, $P < 0.001$, n = 7 experiments, from 6 neurons, from 6 mice, Figure

1B.1 and C.1), and shortening of the activated spine neck within a few minutes ($P < 0.001$), with

no significant change in spine head volume (n = 7, Figure 1B.2 and C.1). These results were also

consistent when we considered the maximum uEPSP change in amplitude from each experiment

and concomitant changes in spine morphology (uEPSP = $134.21 \pm 3.29\%$, $P < 0.001$, n = 7; neck

length = $71.88 \pm 10.66\%$, $P < 0.05$, n = 7; spine head volume = $98.11 \pm 7.34\%$, $P = 0.81$, n = 7)

(Figure 1C.2). We obtained similar results when we instead considered the average of all the values

obtained following t-LTP induction for uEPSP amplitude, neck length and head volume

(Supplementary Figure 1). This effect was specific to the activated spine (Figure 1B.2), with

neighbouring spines having no appreciable changes in their neck length or head volume (neck

length = $98.09 \pm 5.06\%$, $P = 0.71$, n = 13; head volume = $103.01 \pm 3.61\%$, $P = 0.42$, n = 14).

Control experiments showed that there was no significant change in uEPSP amplitude or spine

morphology following the STDP protocol when either action potentials or synaptic stimulation

were applied in isolation, as well as when we monitored the long-term stability of these parameters

without any STDP protocol (Supplementary Figure 2).

A pre-post pairing of + 7 ms showed a non-significant tendency for the induction of LTP following

t-LTP induction, and a non-significant tendency for the shrinkage of the activated spine neck (n =

8 experiments, from 8 neurons, from 8 mice, Figure 1D.1). When the maximum change in uEPSP

amplitude from each experiment and the concomitant changes in spine morphology were analyzed,

we saw no significant changes in uEPSP amplitude and spine morphology (Figure 1D.2, uEPSP =

$112.27 \pm 14.19\%$, P = 0.42, n = 8; neck length = $88.90 \pm 5.89\%$, P = 0.10, n = 8; head volume =

$97.81 \pm 4.54\%$, P = 0.64, n = 8) (Figure 2E). Similar results were observed when we instead

considered the average of all the values obtained following t-LTP (+7 ms) induction for uEPSP

amplitude, neck length and head volume (Supplementary Figure 1). Because voltage has been

shown to be an important factor in the induction of t-LTP and t-LTD, we verified that the initial

uEPSP amplitudes were not significantly different for pre-post pairing protocols of +13 ms versus

+7 ms (uEPSP: 0.62 ± 0.14 versus 0.53 ± 0.16 mV, P = 0.68; Supplementary Figure 3).

These results indicate that there is a preferred pre-post t-LTP pairing time-window (+ 13 ms) at

which activated spines in basal dendrites from L5 pyramidal neurons undergo a significant increase

in synaptic strength, and a concomitant neck shrinkage (Figure 2E).

Induction of t-LTD in single dendritic spines

We then studied t-LTD in single spines by using a repetitive spike-timing protocol (40 times, 0.5

Hz) in which 2P uncaging of glutamate at a single spine was preceded in time (-15 or -23 ms) by

a bAP (post-pre protocol, Figure 2A). When postsynaptic spikes preceded presynaptic firing by 15

ms (i.e., -15 ms), a significant reduction of the uEPSP amplitude occurred within a few minutes

following induction of t-LTD (n= 6 experiments, from 5 neurons and 5 mice, Figure 2B.1 and C.1,

$P < 0.001$), with no significant changes in spine neck length or head dimension (Figure 2B.2 and

C.1). Furthermore, when the maximum change in uEPSP amplitude after the induction of t-LTD

in single spines at pairings of -15 ms was analyzed (Figure 2C.2) we also observed a significant

depression of uEPSP amplitude (uEPSP = 71.52 ± 7.07%, $P < 0.01$, n = 6), with no significant

changes in spine morphology (Figure 2B.2 and C.2, neck length = 105.54 ± 9.85%, P = 0.62, n =

6; head volume = 103.25 ± 3.02%, P = 0.33, n = 6). Interestingly, after the induction of t-LTD in

single spines when postsynaptic spikes preceded presynaptic firing by 23 ms (i.e., -23ms) there

were no significant changes in the amplitude of the uEPSPs or in the spine neck length and head

dimensions for the duration of the recordings (n= 7 experiments, from 7 neurons and 7 mice, Figure

2D.1). These results were also consistent with analyses of the maximal uEPSP change in each

experiment and concomitant spine morphology (Figure 2D.2: uEPSP = 82.09 ± 9.89%, P = 0.12,

n = 7; neck length = 84.15 ± 7.73%, P = 0.09, n = 7; head volume = 98.81 ± 5.57%, P = 0.84, n =

7). There was no significant difference between the initial EPSP amplitude for post-pre pairing

protocols of -15 ms versus -23 ms (EPSP: 0.59 ± 0.07 versus 0.49 ± 0.08 mV, P = 0.42;

Supplementary Figure 3). We obtained similar results when we instead considered the average of

all the values obtained following t-LTD induction in single spines for uEPSP amplitude, neck

length and head volume (Supplementary Figure 1). This indicates that only a post-pre t-LTD

pairing time-window of -15 ms can effectively induce LTD in single dendritic spines in the basal

dendrites from L5 pyramidal neurons.

Taken together these results show that the induction of t-LTP and t-LTD in single spines follows

a Hebbian-STDP learning rule that is bidirectional, and favors presynaptic inputs that precede

postsynaptic spikes and depresses presynaptic inputs that are uncorrelated with postsynaptic spikes

at a very precise and narrow temporal window (+13 ms for the generation of t-LTP and -15 ms for

t-LTD, Figure 2E-F). The single spine STDP rule we observed has a narrower post-pre LTD

induction pairing time window than previously observed in connected pairs of L2/3 [14, 17] and L5

pyramidal neurons [12] – where the presynaptic control of t-LTD via an mGluR and/or cannabinoid

type 1 receptor-dependent mechanism [18-21] could plausibly account for these differences.

Induction of t-LTP in clustered dendritic spines

It has been suggested that STDP not only depends on spike timing and firing rate but also on

synaptic cooperativity and the amount of voltage generated at the postsynaptic site [12, 13]. However,

a direct demonstration of synaptic cooperativity at the level of single spines in the dendrites of

pyramidal neurons remains unknown. Hence, an experiment was designed to directly test if

synaptic cooperativity, marked by the co-induction of t-LTP in clustered dendritic spines from

basal dendrites of L5 pyramidal neurons, and the local spatio-temporal summation of inputs, can

generate a local dendritic depolarization and local calcium signals, that are high enough to disrupt

the single spine STDP learning rule described in Figure 2E. To test this, a two spine STDP protocol

(forty 2P uncaging pulses, pulse duration 2ms, 0.5Hz, see methods section) was performed,

whereby 2P uncaging of caged glutamate in clustered (distance between spines < 5 µm) spines

was followed in time by a bAP to trigger t-LTP (Figure 3A). With this protocol, we investigated

whether activating clustered spines extended the pre-post timing window capable of generating

LTP by increasing the degree of depolarization immediately before the postsynaptic spike at

timings where plasticity was not reliably generated. Specifically, we induced t-LTP in two

clustered spines at pre-post timings of +7 ms, and surprisingly found that this protocol was in fact

capable of effectively and significantly generating increases in uEPSP amplitude (Figure 3B.1)

and the concomitant shrinkage of the activated spine neck, with no apparent changes in its spine

head size (Figure 3B.2). Pooled data showed that significant increases in uEPSP amplitude and

shrinkage of the spine neck of the activated spines occurs only a few minutes post t-LTP induction

and lasted for the duration of the recordings (Figure 3C.1, $P < 0.01$, n = 10 spines from 8

experiments, 6 neurons and 6 mice), with no significant changes in the spine head size (Figure

3C.1, n = 16 spines from 8 experiments, 6 neurons, 6 mice). Similar results were observed when

we analyzed the maximal change in uEPSP amplitude in each experiment and the concomitant

spine neck length and head size of the two clustered spines after induction of t-LTP at pairings of

+ 7 ms (Figure 3C.2; uEPSP = 130.86 ± 8.18%, $P < 0.01$, n = 8; neck length = 73.22 ± 5.84%, $P <$

0.01, n = 10; head volume =102.00 ±2.58%, $P = 0.45$, n = 16). We obtained similar results when

we instead considered the average of all the values obtained following t-LTP induction in two

clustered spines for uEPSP amplitude, neck length and head volume (Supplementary Figure 1). In

control experiments, no significant change in uEPSP amplitude or spine morphology was observed

when we monitored the long-term stability of these parameters without any STDP protocol

(Supplementary Figure 2). These results indicate that synaptic cooperativity – shown by the

induction of t-LTP in only two clustered spines (< 5 µm apart) – is sufficient to significantly trigger

synaptic potentiation and shrinkage of the activated spine necks at a pre-post timing that is

otherwise ineffective at generating significant morphological changes and synaptic potentiation

when only one spine is being activated (for comparison between one versus two cluster spines see

Supplementary Figure 4). Hence, the synaptic cooperativity of only two neighbouring synaptic

inputs onto spines (< 5 µm apart) in the basal dendrites of L5 pyramidal neurons extends the pre-

post timing window that can trigger potentiation (Figure 3C.2, and compare Figure 3C.1 with

Figure 1D.1).

Molecular mechanisms responsible for the generation of t-LTP in dendritic spines

The results led us to consider the possible mechanisms underlying the generation of t-LTP at the

level of single spines. Specifically, we asked why the induction of t-LTP in single and clustered

spines is associated with the shrinkage of the activated spine neck. We and others have reported

that the induction of LTP can trigger activity dependent changes in neck length [2, 16] and spine head

size [4, 6, 16], and that the amplitude of uEPSP recorded at the cell soma is inversely proportional to

the length of the activated spine neck [2, 22, 23]. However the mechanisms by which the t-LTP-induced

neck shrinkage is associated with synaptic plasticity remains unknown. Numerical simulations

show that the EPSP amplitude/neck length correlation can be explained by variations in synaptic

conductance, electrical attenuation through the neck, or a combination of the two [2]. Nevertheless,

solutions that rely exclusively on the passive electrical attenuation of synaptic inputs through the

spine neck assume very high (> 2 GOhm) neck resistance [2], which are at odds with recent spine

neck resistance estimations [24, 25]. These results suggest that the control of AMPA receptor content

in spines could contribute significantly to the observed t-LTP-dependent changes in synaptic

strength. To experimentally study the contribution of AMPA receptors to these phenomena, we

performed t-LTP experiments in two clustered spines from L5 pyramidal neurons recorded via

patch pipettes loaded with intracellular solution containing 200 µM PEP1-TGL – a peptide that

inhibits AMPA receptor incorporation to the postsynaptic density (PSD) by blocking GluR1 C-

terminus interaction with PDZ domains at the PSD [26] (Figure 4A). PEP1-TGL incubation by itself

did not trigger a run-down of uEPSP amplitude or changes in spine morphology over time

(Supplementary Figure 5A). Pooled data from experiments where a repetitive pre-post pairing

protocol of + 7 ms was used to activate clustered spines in the presence of PEP1-TGL show that

the peptide completely inhibited t-LTP for the duration of the experiment (Figure 4B.1 and C.1),

but had no effect on the t-LTP-induced shrinkage of the activated spine necks (Figure 4B.2 and

C.1, $P < 0.001$, n = 6 spines, from 5 experiments, 5 neurons and 5 mice) or in modifying spine

head size (n = 10 spines, from 5 neurons and 5 mice, Figure 4C.1). Furthermore, when we analyzed

the maximum change in uEPSP amplitude and the concomitant spine morphology after the

induction of t-LTP in the presence of PEP1-TGL, we also found an inhibition of t-LTP, but no

effect on the t-LTP-induced shrinkage of the spine neck (Figure 4B.2-C.2, uEPSP = 94.82 ±

14.82%, P = 0.74, n = 5 experiments, from 5 neurons and 5 mice; neck length = 83.92 ± 5.35%, P

< 0.05, n = 6; head volume = 100.08 ± 3.23%, P = 0.98, n = 10). We obtained similar results when

we instead considered the average of all the values obtained following t-LTP induction in two

clustered spines in the presence of PEP1-TGL for uEPSP amplitude, neck length and head volume

(Supplementary Figure 1). No significant difference was observed between the initial uEPSP

amplitude for pre-post pairing protocols of +7 ms with versus without PEP1-TGL (uEPSP: 1.06 ±

0.2 versus 1.16 ± 0.28 mV, P = 0.81; Supplementary Figure 3). These results indicate that GluR1

receptor incorporation into the PSD - via its interaction with PDZ domains - is required for the

induction of t-LTP in spines. However, the role of the spine neck shrinkage in AMPA receptor

incorporation into the PSD and ultimately on the induction of t-LTP remains open.

Experimental and theoretical studies have indicated that lateral diffusion of AMPA receptors into

and out of the spine head can be restricted by the spine neck geometry [27-30]. In particular, lateral

diffusion of AMPA receptors into and out of mushroom spines (long-necked spines) has been

shown to be significantly slower than that observed in stubby spines (small-necked spines) [27] –

which is supported by studies that show reduced diffusion of membrane proteins located in spine

necks [31]. In addition, quantitative models using realistic spine morphologies indicate that

decreasing the radius and increasing the spine neck length increases the retention of AMPA

receptors at the synapse [29], even when their interaction with scaffolding cytoskeletal proteins is

neglected [30]. Actin is highly enriched in the spine neck and head [32], and plays an important role in

anchoring AMPA receptors in the spine [33] and AMPA receptor trafficking [34], being instrumental

for synaptic transmission and plasticity [35-37]. Hence, to address the role that t-LTP-induced neck

shrinkage has on AMPA receptor lateral trafficking to the PSD, and the generation of t-LTP in the

activated spines we focused on actin dynamics. We used the actin polymerization inhibitor

latrunculin A (Lat-A) [33, 35, 37] (Figure 4D) − which did not trigger any run-down of uEPSP

amplitude or changes in spine morphology over time in the absence of STDP induction

(Supplementary Figure 5B) − to test the potential role of actin dynamics on the spine induction of

t-LTP, and on the neck shrinkage and AMPA receptor incorporation into the PSD in the activated

spines (Figure 4D). The induction of t-LTP at pre-post pairings of +7 ms in two clustered spines

in the presence of 100 nM Lat-A completely blocked the shrinkage of the activated spine necks

and the induction of t-LTP (Figure 4E and F.1, n = 8 spines, from 3 neurons and 2 mice), inducing

instead a significant reduction in uEPSP amplitude over time (Figure 4F.1, $P < 0.001$, n = 4

experiments, from 3 neurons and 2 mice). These observations were also consistent with analyses

of the maximal change in uEPSP amplitude in each experiment and concomitant spine morphology

post t-LTP induction in the presence of Lat-A (Figure 4F.2, uEPSP= $55.45 \pm 7.13\%$, $P < 0.01$, n =

4 experiments; neck length = $87.20 \pm 9.15\%$, $P = 0.21$, n = 8 spines; head volume = $101.25 \pm$

10.99%, $P = 0.91$, n = 8 spines).

We obtained similar results when we instead considered the average of all the values obtained

following t-LTP induction in two clustered spines in the presence of 100 nM Lat-A for uEPSP

amplitude, neck length and head volume (Supplementary Figure 1).

No significant difference was observed between initial uEPSP amplitudes for pre-post pairing

protocols of +7 ms with versus without Lat-A (uEPSP: 0.94 ± 0.20 versus 1.16 ± 0.28 mV, P =

0.61; Supplementary Figure 3). The lack of run-down of uEPSP amplitude over time in neurons

treated with Lat-A in the absence of STDP induction (Supplementary Figure 5B), but the

significant depression in uEPSP after the induction of t-LTP suggests that the induction of

plasticity, and the rearrangement of actin filaments de-stabilized AMPA receptors, leading to

removal from the PSD.

In summary, these results show that actin polymerization is required for the t-LTP-dependent neck

shrinkage and the induction of plasticity. Our findings further suggest that the induction of t-LTP

occurs via a mechanism that involves a neck-shrinkage-dependent facilitated diffusion of GluR1

subunits from the spine neck to the head, and subsequent incorporation into the PSD. We

hypothesize that a shorter and wider neck facilitates the transport of AMPA receptors into the spine

head (Figure 4D), a mechanism that is required for the induction of t-LTP.

Induction of t-LTD in clustered and distributed dendritic spines

We then studied whether the induction t-LTD in single spines observed at pairings of -15 ms could

be affected by synaptic cooperativity. Our reasoning was based on two previous observations

which suggest that 1) t-LTP induction in the distal dendrites of L5 pyramidal neurons (layer 3- L5

pyramidal neuron pairs) triggers LTD instead of LTP, and 2) that LTD can be converted into LTP

by increasing the local voltage [13]. We hypothesised that the induction of t-LTD in single spines

depends on the degree of local depolarization and hence, LTD can be disrupted by the activation

of neighboring spines. To test this, we performed repetitive spike-timing protocol (40 times, 0.5

Hz) in which 2P uncaging of glutamate at two spines (separated by up to 100 µm) was preceded

in time (-15 ms) by a bAP (Figure 5D and Supplementary Figure 6A). Surprisingly, we found that

this t-LTD protocol failed to induce any change in uEPSP amplitude or spine head volume with

only a slight but significant reduction in spine neck length (Supplementary Figure 6B). When we

analyzed the maximum uEPSP change in amplitude in each experiment and the concomitant

morphological alterations in the activated spines after t-LTD induction, we observed a complete

inhibition in the induction of t-LTD, and no change in spine morphology (Supplementary Figure

6C; uEPSP = 93.22 ± 6.29%, P = 0.30, n = 17 experiments from 14 neurons and 14 mice; neck

length = 88.56 ± 5.69%, P = 0.06, n = 23 spines; head volume = 102.41 ± 6.10%, P = 0.69, n = 34

spines). To more precisely characterize the effect of activating two spines on the induction of t-

LTD, we correlated the inter-spine distance and the uEPSP change following STDP induction (see

Methods). We found that as the two activated spines were further away from each other, the more

t-LTD was recovered (Supplementary Figure 6D). Specifically, the uEPSP change decayed

exponentially as a function of inter-spine distance with a length constant (λ) of 43.5 µm. Therefore,

we used this value as a boundary between clustered (< 40 µm) and distributed (> 40 µm) spines.

Using this classification, clustered spines were located in the same dendrite (n = 11/12 pairs) or in

sister branches emanating from the same bifurcation point (n = 1/12 pairs), while distributed spines

were always located on separate dendrites (n = 5/5 pairs). When we separated our data in this

manner, the t-LTD protocol in two clustered spines (Figure 5A) failed to induce LTD (Figure 5B.1)

or changes in spine head size at all the times tested post t-LTD induction (Figure 5B.2 and C.1, n

= 12 experiments, n = 24 spines, from 11 neurons and 11 mice), with only a slight but significant

induction of shrinkage of the spine neck at some time points (Figure 5C.1, P < 0.05, n = 19 spines,

from 11 neurons and 11 mice). For comparison between the activation of one versus two clustered

spines with a post-pre timing of -15 ms see Supplementary Figure 7. When we analyzed the

maximum uEPSP change in amplitude in each experiment and the concomitant morphological

alterations in the activated spines after t-LTD induction in two clustered spines, we also observed

a complete inhibition in the induction of LTD, a slight but significant reduction in spine neck

length, and no changes in spine head size (Figure 5C.2; uEPSP = 101.70 ± 7.02%, P = 0.81, n =

12 experiments; neck length = 85.28 ± 5.96%, P < 0.05, n = 19 spines; head volume = 102.98 ±

8.86%, P = 0.73, n = 24 spines, from 12 experiments performed in 11 neurons from 11 mice). We

obtained similar results when we instead considered the average of all the values obtained

following t-LTD induction in clustered spines for uEPSP amplitude, neck length and head volume

(Supplementary Figure 1). These results were surprising since not only did we not observe t-LTD

in clustered spines, but we also observed significant neck shrinkage with no LTP (see Figure 1 and

3). To account for this observation, we explored if there was a correlation between the induction

of plasticity in these experiments and both the shrinkage of the spine neck and the distance between

the activated clustered spines – since the local voltage, and hence the induction of plasticity, could

be affected by the distance between the activated clustered spines. Indeed, we found that the

distance between the activated spines under these experimental conditions (t-LTD induction

protocol in clustered spines) is correlated with the induction of plasticity and the shrinkage of the

activated spine necks (Equation 1 in Methods; P < 0.01; Supplementary Figure 6E). This analysis

suggests that during t-LTD induction the structural arrangement of clustered spines (< 40 µm)

determines the sign and magnitude of the change in synaptic strength and concomitant neck

shrinkage.

We next investigated the mechanisms underlying the disruption of t-LTD by activating spines

separated by increasingly larger distances (Figure 5D). Interestingly, the induction of t-LTD in

spines separated by more than 40 µm (distributed spines) was capable of recovering the generation

of LTD (Figure 5E.1 and Supplementary Figure 6D). Pooled data from all experiments

demonstrate that the activation of distributed spines reliably induces t-LTD (Figure 5F.1,

significant reduction in uEPSP amplitude, $P < 0.01$, n = 5 experiments from 5 neurons and 5 mice),

without triggering changes in neck length or spine head size (Figure 5E.2 and F.1). When we

analyzed the maximal change in uEPSP amplitude in each experiment and concomitant spine

morphological changes, we saw a significant induction of t-LTD and no change in spine

morphology (Figure 5F.2, uEPSP = 72.86 ± 8.08%, $P < 0.05$, n = 5 experiments, neck length =

97.85 ± 15.47%, $P = 0.89$, n = 6 spines; head volume = 101.06 ± 4.59%, $P = 0.82$, n = 10 spines)

as what was found in experiments where t-LTD was generated at pairing times of -15 ms in single

dendritic spines (Figure 2B-C). We obtained similar results when we instead considered the

average of all the values obtained following t-LTD induction in distributed spines for uEPSP

amplitude, neck length and head volume (Supplementary Figure 1). No significant difference was

observed between the initial uEPSP amplitude for clustered versus distributed spines activated

with post-pre pairings of -15 ms (EPSP: 1.06 ± 0.13 versus 1.31 ± 0.19 mV, $P = 0.25$;

Supplementary Figure 3). For comparison between the activation of clustered versus distributed

spines after post-pre pairings of -15 ms, see Supplementary Figure 8.

In summary, this data shows that the induction of t-LTD at pairing times of -15 ms was completely

disrupted when only two clustered spines (< 40 μm apart) were activated in the basal dendrites of

L5 pyramidal neurons, but could be recovered if the activated spines are distributed (> 40 μm) in

the dendritic tree.

Spine calcium transients during the induction of t-LTP and t-LTD in single and clustered

dendritic spines

Calcium is critical for the induction of synaptic plasticity [38-42], and high or low local concentration

difference in dendrites and spines are thought to be associated with gating LTP or LTD,

respectively [43-45]. Therefore, to investigate the different mechanisms – with respect to local calcium

accumulations – underlying the induction of t-LTP and t-LTD in single versus two clustered

spines, we performed 2P calcium imaging in a region of interest (ROI) of the activated spines and

their parent dendrites during STDP induction protocols throughout each of the 40 pre-post or post-

pre repetitions (see Methods). The "before" images correspond to the calcium signals observed in

the ROI right before the pairing in each repetition – uncovering the lack or presence of local

calcium accumulation during the 40 pairing repetitions. The "after" images correspond to the

calcium signals observed in the ROI right after the pairing in each repetition – uncovering a proxy

for the amplitude and local calcium accumulation during the 40 pairing repetitions.

To dissect potential differences in local calcium signals and accumulation that can account for the

presence or absence of t-LTP and t-LTD induction in clustered versus distributed spines, we

imaged 2P calcium activity during four different STDP induction protocols: (1) pre-post pairing

of +7 ms in one spine; (2) pre-post pairing of +7 ms in two clustered spines; (3) post-pre pairing

of -15 ms in one spine; (4) post-pre pairing of -15 ms in two clustered spines.

During the pre-post (+7 ms) pairing protocol in single spines we found that, across the 40

repetitions, there was little to no calcium accumulation in the spine or dendrite (Figure 6A-B and

left panels in Figure 6C, n = 7 experiments, from 6 neurons, and 4 mice). As expected, there was,

however, a significant increase in calcium immediately following the stimulation (left panels in

Figure 6F) that due to the lack of accumulation throughout the 40 repetitions, did not build up a

local calcium signal in the activated dendrites and spines. In contrast, when we applied the exact

same pairing protocol (pre-post + 7ms) in two clustered spines, there was a striking calcium

accumulation in both the activated spines and dendrite that was evident when we analyzed the

images taken before (Figure 6D-E and middle panels in Figure 6C) and after stimulation (middle

panels in Figure 6F, n = 6 experiments, from 4 neurons, and 4 mice). Thus, activating just one

additional spine using the same pairing protocol alters the calcium dynamics (compare black and

green traces in right panels of Figure 6C and F), possibly through a mechanism that is incapable

of extruding calcium increases in spines in between pre-post repetitions, leading to its build up in

spines and parent dendrites, which ultimately guide the induction of plasticity.

We performed the same experiments with a post-pre (-15 ms) pairing protocol in both single and

clustered spines. In single spines, we observed moderate calcium increases (Figure 7A-B, left

panels in Figure 6C and F, n = 5 experiments, from 4 neurons, and 4 mice) that were observed

when we analyzed images taken before and after the post-pre stimulation. Surprisingly, we found

similar results to those observed with single spine t-LTD induction protocols, when we applied the

same pairing protocol in two clustered spines (Figure 7D-E, middle panels in Figure 7A and F, n

= 6 experiments, from 4 neurons, and 3 mice) even though no plasticity is induced in this condition.

As suggested by previous studies [46], we hypothesize that the range of spine calcium levels required

for the induction of t-LTD is relatively narrow, and that the resolution with our current

experimental set-up is not sufficient to tease apart significantly different calcium dynamics in one

versus two clustered spines during a post-pre pairing protocol of -15 ms. Moreover, modeling

STDP provide evidence that, in addition to overall calcium levels, the detailed time course of

calcium levels in the postsynaptic cell during a pairing protocol also guide the induction of

plasticity [47]. Nonetheless, these results suggest that the induction of t-LTD does not require

significant calcium accumulations during the 40 repetitions, and most likely depends on the

amplitude of calcium signals right after the stimulation. An interesting observation is that when

we fit the calcium signals with a linear regression, the fits from all the different induction protocols

have different slopes, which goes in line with our hypothesis (Supplementary Figure 9). More

specifically, in single spines, a pre-post pairing protocol of +7 ms induced a relatively low calcium

signal with a shallow slope (black lines in Supplementary Figure 9) whereas in clustered spines

this same protocol caused a robust increase in calcium with a steep slope (blue lines in

Supplementary Figure 9). A post-pre pairing protocol of -15 ms, caused a modest increase in

calcium in both single and clustered spines before the stimulation (red and green lined in left panels

of Supplementary Figure 9), whereas after the stimulation the calcium increase is more prominent

in clustered spines (red and green lined in right panels of Supplementary Figure 9). These results

provide evidence that the calcium levels needed to induce of t-LTD are restricted to a narrow range

and that surpassing this range biases towards the induction of weak levels of LTP.

DISCUSSION

We uncovered the STDP rules for single, clustered and distributed dendritic spines in the basal

dendrites of L5 pyramidal neurons. Our results show that the induction of STDP in single spines

follows a classical Hebbian STDP learning rule that is bidirectional, in which presynaptic input

leading postsynaptic spikes generates t-LTP and postsynaptic spikes preceding presynaptic

activation of single dendritic spines results in t-LTD. Furthermore, we found that the induction of

t-LTP triggers the shrinkage of the activated spine neck, without any significant changes in the

spine head size. Our results indicate that the induction of t-LTP requires 1) the incorporation of

new GluR-1 receptors with PDZ-domain containing proteins in the PSD and, 2) an actin

polymerization-dependent neck shrinkage of the activated spine neck (Figure 4). We showed that

the induction of t-LTP triggers actin-dependent neck shrinkage, which is likely required for the

lateral diffusion of GluR-1 receptors from the spine neck to the spine head, and its incorporation

to the PSD – generating plasticity. In support of this spine mechanism of LTP induction is a recent

report showing that AMPA receptor surface diffusion is fundamental for the induction of

hippocampal LTP and contextual learning [48]. In addition, we found that the induction of t-LTD

was not accompanied with spine neck or head changes, which is at odds with previous findings

suggesting structural changes in spine head volume during the induction of LTP or LTD [4, 7, 49]. The

discrepancy between our results and those observed previously after the induction of t-LTP (head

enlargement [49]), LTP [4], or LTD (head shrinkage, [7]) using glutamate uncaging are likely explained

by methodological differences. While our data was obtained using ACSF with physiological

concentrations of magnesium and calcium, those from other reports were done in a magnesium-

free ACSF [4, 7], low calcium extracellular solution for the induction of LTD [7], or in a magnesium-

free ACSF and an intracellular solution containing 5 µM actin that was required for the t-LTP-

mediated spine head enlargements [49].

Nonetheless, it has been shown in vivo that a spike-timing protocol triggers receptive field

plasticity in layer 2/3 pyramidal neurons is correlated with spine head volume changes

(enlargement and shrinkage) observed after 1.5-2 hours [50]. Taken together, our data suggest that

there is a new form of structural spine plasticity during t-LTP that involves rapid neck shrinkage

without head volume enlargements. In addition, we show that the induction of t-LTD does not

require structural spine changes. Although spines have the machinery and do undergo structural

head changes, we propose that our results represent a stage during memory formation that occurs

before structural head volume changes, a process likely linked with memory consolidation.

Importantly, our data suggest that during STDP, the use of spine volume changes as the sole proxy

for LTP or LTD [50] is not a complete representation of plasticity in spines from dendrites in cortical

pyramidal neurons.

We then explored the functional consequences of synaptic cooperativity on STDP. Our results

show that the induction of t-LTP in two clustered spines - separated by less than 5 µm - is sufficient

to induce LTP and shrinkage of the activated spine necks at a pre-post timing that is otherwise

ineffective at triggering significant morphological changes and synaptic potentiation when only

one spine is being activated. These results show that the activation of clustered spines extends the

pre-post timing window that can trigger potentiation. On the other hand, the induction of t-LTD in

two clustered spines disrupts the generation of LTD leading to a STDP learning rule that is

incapable of supporting LTD, but only encompasses LTP (Figure 8A). We next investigated the

dendritic mechanisms responsible for the disruption of t-LTD, and found that the induction of t-

LTD is fully recovered when the activated spines are separated by more than 40 µm (Figure 5,

Figure 8A and Supplementary Figure 6). Interestingly, the effective length constant (λ), that

represents the length at which the electrotonic potential decays to a value of 37% of that at the

point of origin, in the basal dendrites of L5 pyramidal neurons has been reported to be 50 µm [51].

This value of λ supports the idea that significant voltage attenuations – capable of recovering LTD

– can be expected when the t-LTD induction protocol is triggered in spines that are separated by

more than 40 µm in the basal dendrites of L5 pyramidal neurons (Figure 5 D-F). However, we

cannot discard that other mechanisms, such as the diffusion of active molecules [5], could contribute

to the switch from LTD to no-LTD induction observed in distributed/single spines and clustered

spines, respectively. These results are in discrepancy with observations showing that in connected

pairs of L5-L5 pyramidal cells, t-LTD is reliably generated after post-pre pairing protocols [12]. A

likely explanation for this apparent controversy is that the synaptic inputs from one L5 pyramidal

neuron to another are distributed [52]. Importantly, clustered and distributed excitatory inputs have

been described in the dendrites of pyramidal neurons both *in vitro* and *in vivo* [1, 53-55]. Our results

clearly show the functional importance that the structural and temporal organization of excitatory

synaptic inputs have on the induction of t-LTP and t-LTD, and how just two clustered excitatory

synaptic inputs are capable of altering the STDP learning rule in the basal dendrites of L5

pyramidal neurons (Figure 8A).

How the synaptic activation of just one extra clustered spine is capable of (1) inducing t-LTP at a

pre-post timing that is otherwise ineffective in inducing potentiation and (2) disrupting the

induction of t-LTD? To explore the mechanisms that may be responsible for these observations

we imaged local calcium signals in the activated spines and parent dendrites before and after each

of the 40 pairings performed during t-LTP and t-LTD induction protocols. Our reasoning for

performing these experiments was based on findings that different levels of depolarization gate

local calcium signals, which depending on its magnitude and kinetics, can generate LTP (high

calcium) or LTD induction (sustained but moderate calcium signals) [9, 43, 56]. In addition, calcium-

based modeling studies of STDP have shown that different calcium dynamics mediate the

induction of t-LTP versus t-LTD [46, 47]. Specifically, the calcium control hypothesis indicates that

large levels of calcium (above a plasticity threshold, Θ_p) are thought to lead to t-LTP whereas more

moderate, prolonged levels (between the depression threshold, Θ_{dSTART}, and Θd_{STOP})) give rise to

t-LTD and a mid-level range in which t-LTD does not occur (below Θ_{dSTART}) (Figure 8B) [47, 57, 58].

A major assumption of these models is infinite time constants for synaptic variables at resting

calcium levels so that the synaptic changes do not to decay after the presentation of the stimulus

[46] - a significant constraint for the stabilization of synaptic changes. A potential solution to this

problem is the degree of local calcium accumulation observed in the activated spines throughout

the t-LTP or t-LTD induction protocol. In fact, these models are consistent, fundamentally, with

our results which show that a pre-post pairing (+7 ms) protocol in two clustered spines gives rise

to t-LTP accompanied by a substantial increase in the intracellular calcium levels following each

pairing repetition, and a significant accumulation of local calcium levels throughout the induction

protocol – likely mediated by the inability of the two clustered activated spines to efficiently

extrude the local calcium signals in between each pre-post pairing (Figure 6). We propose that the

local spine calcium accumulation we observe provides a new and key variable for the induction of

plasticity, which reduces the constraints imposed by calcium-base models for the stabilization of

synaptic changes [46, 47, 57, 58].

These changes in local spine and dendritic calcium signals (Figure 6) suggest that perhaps Θ_p can

be reached only with ~ 10 pre-post pairings (~20-30 seconds). In contrast this same protocol in

one spine induces no plasticity, producing calcium signals right after the pairing stimuli that are

effectively extruded in between pairings leading to no calcium accumulation during the induction

protocol (reaching levels below Θ_p and Θ_{dSTART}; Figure 8B). These results suggest that it is not

only the amplitude of the local calcium signals after each pairing, but also the local calcium

accumulation during the induction protocol (40 pairings, ~80 seconds) in spines and dendrites that

are required to reach Θ_p for the induction of t-LTP in clustered spines. As mentioned before,

recently it has been demonstrated in vivo that spike timing-induced receptive field plasticity, with

millisecond time delays between visual stimulus (pre) and optogenetic stimulation in layer 2/3

pyramidal neurons (post), is correlated with increases in synaptic strength [50]. These results together

with evidence from other in vivo studies showing that layer 5 pyramidal neurons can spike up to

frequencies of 20 Hz during movement [59], suggest that our pairing protocol, and findings, are likely

present under in vivo conditions and are relevant for plasticity of networks and ultimately

behaviour.

Our results further show that a post-pre protocol of -15 ms in a single spine induces t-LTD and

moderate intracellular calcium signals in spines and parent dendrites after each pairing, without an

evident increase in local calcium accumulation. These results possibly reflect that the calcium

signal generated during the induction protocol passed Θ_{dSTART} and remain for several seconds in

this permissive calcium concentration window – between Θ_{dSTART} and Θ_{dSTOP} – generating LTD

(Figure 8B). Activating two clustered spines with the same protocol, however, does not induce

plasticity and gives rise to an apparent smaller initial calcium accumulation than that observed

with the activation of a single spine but with a slow build-up of calcium. These results possibly

reflect that the spine calcium levels crossed Θ_{dSTART} only after > 20 repetitions and then crossed

Θ_{dSTOP} and Θ_{pSTART} after a few (<10) repetitions reaching slightly higher local calcium levels. This

calcium control hypothesis of t-LTD induction is based on the average linear fits of each

experiment, and a tendency, that although clear, is not statistically significant with our

measurements (Figure 8B and Supplementary Figure 9).

These findings presented here are quite remarkable since stimulating just one additional spine

during a STDP protocol can completely alter the calcium dynamics and the induction of t-LTP and

t-LTD. To our knowledge, this is the first demonstration of the functional relevance that the

structural organization and simultaneous subthreshold activation of only a few clustered inputs in

the dendrites of pyramidal neurons have on plasticity. We propose the term *micro clusters* to

describe this structural and functional modality of synaptic connectivity. In fact, the relevance of

synaptic *micro clusters* on the input/output properties of pyramidal neurons is also supported by

three dimensional electron microscopy and neuronal reconstruction studies that have shown the

presence of postsynaptic innervation of the same axon spaced at less than 10 µm in the basal

dendrites of L2/3 pyramidal neurons from the medial entorhinal cortex [53], L5 pyramidal neurons

from somatosensory cortex [54] and in the distal apical tuft dendrites in stratum lacunosum-

moleculare of hippocampal CA1 pyramidal neurons [55]. In addition to having spines innervated by

the same axon, it is likely that functional synaptic *micro clusters* can be gated by the convergence

of different axons, which could increase the computational power of cortical circuits through a

multi-neuronal control of synaptic cooperativity and ultimately the implemented STDP learning

rule. Furthermore, recently it has been shown that orientation selectivity in visual cortex is

correlated with the degree of spatial synaptic clustering of co-tuned synaptic inputs within the

dendritic field [60], and that functional clusters of dendritic spines separated by less than 10 µm share

similar spatial receptive field properties, spontaneous and sensory-driven activity [61]. Taken

together these reported findings and our data suggest that the functional specificity and structural

arrangement of synaptic inputs, distributed or forming *micro clusters* in the dendrites of pyramidal

neurons, are fundamental for guiding the rules for sensory perception, affecting the STDP learning

rule, learning and memory, and ultimately cognition.

METHODS

Brain slice preparation and electrophysiology. Brains from postnatal day 14-21 C57B/6 mice -

anesthetized with isoflurane - were removed and immersed in cold (4°C) oxygenated sucrose

cutting solution containing (in mM) 27 $NaHCO_3$, 1.5 NaH_2PO_4, 222 Sucrose, 2.6 KCl, 1 $CaCl_2$,

and 3 $MgSO_4$. Coronal brain slices (300-µm-thick) of visual cortex were prepared as described [22].

Brain slices were incubated for 1/2 hour at 32°C in artificial cerebrospinal fluid (ACSF, in mM:

126 NaCl, 26 $NaHCO_3$, 10 Dextrose, 1.15 NaH_2PO_4, 3 KCl, 2 $CaCl_2$, $2MgSO_4$) and then

transferred to a recording chamber. Electrophysiological recordings were performed in whole-cell

current-clamp configuration with MultiClamp 700B amplifiers (Molecular Devices) in L5

pyramidal neurons with a patch electrode (4-7 MΩ) filled with internal solution containing (in

mM) 0.1 Alexa Fluo 568, 130 D-Gluconic Acid, 2 $MgCl_2$, 5 KCl, 10 HEPES, 2 MgATP, 0.3

NaGTP, pH 7.4, and 0.4% Biocytin. All experiments were conducted at room temperature (~20-

22°C). We did not extend our experiments to include voltage-clamp recordings since recent

evidence indicates that the high spine neck resistance can prevent the voltage-clamp control of

excitatory synapses and that these measurements can be significantly distorted in spiny neurons[62].

Two-photon imaging and two-photon uncaging of glutamate. Two-photon imaging was performed

with a custom-built two-photon laser scanning microscope, consisting of 1) a Prairie scan head

(Bruker) mounted on an Olympus BX51WI microscope with a 60X, 0.9 N.A. water immersion

objective; 2) a tunable Ti-Sapphire laser (Chameleon Ultra-II, Coherent, >3 W, 140-fs pulses, 80

MHz repetition rate), 3) two photomultiplier tubes (PMT) for fluorescence detection. Fluorescence

images were detected with Prairie software (Bruker).

Fifteen minutes after break-in, two-photon scanning images of basal dendrites were obtained with

720 nm and low power (~5 mW on sample (i.e., after the objective)) excitation light and collected

with a PMT. Two-photon uncaging of 4-methoxy-7-nitroindolinyl (MNI)-caged L-glutamate (2.5

mM; Tocris) was performed as previously described [63]. This concentration of MNI-glutamate

completely blocked IPSCs [64], thus, our results represent excitatory inputs only. Uncaging was

performed at 720 nm (~25-30 mW on sample). Note that the laser power used for imaging is not

sufficient to result in any partial uncaging of glutamate (Supplementary Figure 10). Activated

spines were mostly on the second and third branch of the basal dendrites and were on average ~

40 µm away from the soma (Supplementary Figure 11). Only spines with a clear head contour and

that were separated by >1 µm from neighboring spines were selected. The location of the uncaging

spot was positioned at ~ 0.3 µm away from the upper edge of the selected spine head (red dot in

figures), which has a spatial resolution of 0.71 and 0.88 µm for one and two spines respectively

(Supplementary Figure 12). Care was taken maintain the position of the uncaging spot. After each

uncaging sequence, the spot was repositioned to keep the same distance of ~ 0.3 µm from the edge

of the soma and to avoid artificial potentiation or depression. The uncaging-induced excitatory

postsynaptic potentials (uEPSP) were recorded at the soma of L5 pyramidal neurons. The kinetics

of uEPSPs from the present study are not significantly different (10/90 rate of rise: 0.07 ± 0.014

mV/ms, p=0.92; duration: 115.5 ± 15.3 ms, p=0.65) from those triggered at 37°C [22].

Spike timing-dependent plasticity (STDP) induction protocol: To induce t-LTP in single spines,

we used two-photon uncaging of MNI-glutamate (40 times every 2 seconds, with each uncaging

pulse lasting 2 ms), which, after 7 or 13 ms, was followed by a backpropagating action potential

(bAP) (triggered by a brief (10 ms) current injection (0.4 -0.6 nA) in the soma). To induce t-LTD

in single spines, two-photon uncaging of MNI-glutamate (40 times every 2 seconds, with each

uncaging pulse lasting 2 ms) was preceded for -15 or -23 ms by bAP. When we evaluated t-LTP

and t-LTD in two spines, we used similar protocols to those described above, but the spines were

activated with two-photon uncaging of MNI-glutamate sequentially with an onset delay of ~2.1

ms for the second spine. No significant difference was observed in the in 10/90 rise time of the

uEPSPs triggered when one versus two spines were activated (9.05 ± 1.19 ms versus 9.49 ± 0.54

ms, respectively; p=0.71).

To evaluate the morphological and synaptic strength of the activated spines before and after the

STDP induction protocol, we performed 2P imaging, and low frequency 2P uncaging (0.5 Hz) in

single or multiple spines. To establish the time course of the changes in uEPSP amplitude, neck

length and head volume following STDP induction, for each experiment, we interpolated the data

taken at different time points using the *interp1* function in MATLAB (MathWorks) with the *pchip*

option, which performs a shape-preserving piecewise cubic interpolation. Note that we constrained

this fit so that it terminated with a slope of zero following the last data point. Then, for each

condition, we averaged the uEPSP amplitude, neck length and head volume temporal traces. The

length of the x-axis was set as the time at which the last data point was obtained for those sets of

experiments. Shaded area represents ±SE. To determine at which time the EPSP amplitude, neck

length and head volume temporal traces are significantly different from baseline, we binned the

temporal traces every 5min and tested whether it was significantly different from baseline (100%).

The time at which the maximal change in uEPSP was observed after t-LTP and t-LTD induction

was used to calculate the percent change from control, and the percent changes in neck length and

head volume. These analyses are displayed in Figures 1 to 5.

Experimental checkpoints and data analysis: Electrophysiological data were analyzed with

Wavemetrics software (Igor Pro) and MATLAB. The resting membrane potential of the recorded

L5 pyramidal neurons was -58.27 ± 2.08 mV (n = 58 neurons). After taking this measurement,

pyramidal neurons were maintained at − 65 mV in current clamp configuration throughout the

recording session. Only neurons for which the injected current to hold the cell at − 65 mV was <

100 pA were included in this study. For the generation of bAP, only action potentials with

amplitude of > 45 mV were considered for analysis. In most experiments, two control tests (each

consisting of 10 uncaging pulses at 0.5 Hz), spaced by 5 min were performed to assess the

reliability of the uEPSP amplitude. Only experiments for which uEPSP amplitudes were not

significantly different before and after 5 minutes in control conditions were considered for analysis

(less than 10% variation).

Synaptic plasticity was assessed by two parameters: the uEPSP amplitude and the spine

morphology (neck length and head volume). The peak uEPSP amplitude was measured from each

individual uEPSP by taking the peak value and averaging 2 ms before and after using Wavemetrics

(Igor Pro). Only uEPSPs that were >0.1 mV in the control condition (i.e., before the induction of

plasticity) were included in the analysis.

Synaptic plasticity was determined by the relative change of uEPSP amplitude (average of 10

uEPSP) measured before and after the STDP protocol. For each experiment, we evaluated whether

the STDP protocol generated potentiation or depression by determining how many uEPSP data

points fell above or below baseline values over the course of the experiment. Potentiation was

defined as the majority of uEPSP amplitude data points measured over time increasing relative to

baseline, and the maximum uEPSP increase was used for statistical test. Depression was defined

as the majority of uEPSP amplitude data points measured over time decreasing relative to baseline,

and the maximum uEPSP decrease was used for statistical test. The spike timings +7 ms and +13

ms (pre leads post) or -23 ms and -15 ms (post leads pre) correspond to the delta time offset

between the beginning of the uncaging pulse (pre) and the beginning of the bAP pulse (post)

repeated 40 times.

The analysis of spines morphology was performed from z-projections of the whole spine using

ImageJ [65] (neck length) and MATLAB (MathWorks) (head volume). The neck length was

measured from the bottom edge of the spine head to the edge of its parent dendritic shaft using the

segmented line tool in ImageJ. We selected mostly spines with a spine neck longer than 0.2 µm.

For those with a shorter neck, we did not report their length for analysis and statistics due to the

diffraction limited resolution of our images. For spines whose necks shrunk after the STDP

protocol below the diffraction limited resolution of our microscope, we set their length as the

minimal measurement of spine neck length reported by Tonnesen et al., using STED microscopy

(0.157 µm) [16]. We estimated the relative spine head volume using the ratio of the maximum spine

fluorescence and the maximum fluorescence observed in the dendrite measured from z-projections

of the whole spine [66, 67]. To obtain the spine volume, we then multiplied this ratio by the PSF of

our microscope (0.11 fL) [68]. Linear optimization techniques were used to determine the correlation

between EPSP change, neck length change and distance between 2 activated spines. Specifically,

the change in EPSP amplitude was modeled using the following equation:

$$uEPSP = c_1 \times NL + c_2 \times D \qquad\qquad\qquad Equation\ 1$$

Where *uEPSP* and *NL* are the percent change in uEPSP and neck length, respectively, following

the STDP protocol, *D* is the distance between the 2 spines, and c_1, c_2 and c_3 are constant

coefficients. These parameters were estimated using a least squares technique to obtain an optimal

fit of the data that minimized the sum of the residuals squared. The relationship between inter-

spine distance and the percent change in uEPSP was fit with the following exponential equation:

$$y = \alpha e^{\frac{-x}{\lambda}} + \beta \qquad\qquad\qquad Equation\ 2$$

where α, β and λ are constants, y represents the change in uEPSP, and x is the inter-spine distance.

Calcium imaging: During calcium imaging experiments, we performed whole-cell current-clamp

recordings of L5 pyramidal neurons with a patch electrode containing calcium indicator Fluo-4

(300 μM; Thermo Fisher) and Alexa-594 (100 μM) diluted in an internal solution containing (in

mM) 130 D-Gluconic Acid, 2 $MgCl_2$, 5 KCl, 10 HEPES, 2 MgATP, 0.3 NaGTP, pH 7.4, and 0.4%

Biocytin. To perform sequential 2P calcium imaging and 2P uncaging of caged glutamate in

selected spines at one wavelength (810 nm), we used ruthenium-bipyridine-trimethylphosphine

caged glutamate (RuBi-glu, Tocris) [64], diluted into the bath solution for a final concentration of

600 μM. Uncaging of Rubi-glu was performed at 810 nm (~25-30 mW on sample). The location

of the uncaging spot was positioned at ~ 0.3 μm away from the upper edge of the selected spine

head (red dot in Figures 6-7). Changes in calcium were monitored by imaging 2P calcium signals

and detecting the fluorescence with 2 PMTs placed after wavelength filters (525/70 for green,

595/50 for red). We performed 2P calcium imaging during 4 different STDP induction protocols

triggered at 0.5Hz: (1) pre-post pairing of +7 ms in one spine; (2) pre-post pairing of +7 ms in two

clustered spines; (3) post-pre pairing of -15 ms in one spine; (4) post-pre pairing of -15 ms in two

clustered spines. We restricted the image acquisition to a small area (~150 x 150 pixels) which

contained the spine(s) that we uncaged and the shaft. Images were acquired at ~ 30 Hz, averaged

8 times, with 8 μs dwell time. Calcium signals were imaged 500 ms before STDP induction

protocol and right after (4ms) the stimulation for more than 600 ms. We focused our analysis on

the images obtained before and immediately after the stimulation in each pairing repetition. ROI

drawing was performed using custom algorithms (MATLAB; MathWorks). For spine heads, the

ROI was a circle whereas for dendrites it was a polygon. Fluorescence was computed as the mean

of all pixels within the ROI. We quantified the relative change in calcium concentration ($\frac{\Delta F}{F}$) using

the following formula:

$$\frac{\Delta F}{F} = \frac{G - G_{baseline}}{R}$$ *Equation 3*

where G is the fluorescence from the Fluo-4 dye and R is the fluorescence from the Alexa-594

dye. $G_{baseline}$ is the mean of all pixels of Fluo-4 signal within the ROI taken from the first image

at the first stimulation. We estimated the calcium signal during each condition and using the

following equation:

$$\frac{\widehat{\Delta F}}{F} = a \times x + b$$ *Equation 4*

where $\frac{\widehat{\Delta F}}{F}$ is the estimated change in calcium signal, x is the repetition (binned every 5 repetitions),

a the slope and b a constant coefficient.

Statistics: Statistics were performed with GraphPad Prism 5. Statistical significance was

determined using two-tailed Student's paired t-test when we analyzed the maximum change in

uEPSP amplitude after the induction of t-LTP or t-LTD in each experiment and the concomitant

changes in the activated spine morphology. Statistical significance was determined using one-way

repeated measures ANOVA when we analyzed the time course of the uEPSP amplitude and spine

morphological changes after induction of t-LTP or t-LTD with post-hoc pairwise comparisons

using Dunnett's test. *$P<0.05$; **$P<0.01$; ***$P<0.001$.

Pharmacology: Latrunculin A (Lat-A, Tocris Bioscience) was dissolved in DMSO at 1/1000 and

added to the recording chamber containing the brain slice at 100 nM for 15 min before starting the

STDP protocol. PEP1-TGL (Tocris Bioscience) was added in the pipette at 200 µM; after 15 min

in whole cell condition, electrophysiological recording and synaptic plasticity experiments were

started. MNI-glutamate (Tocris Bioscience) was diluted in ACSF from stock solution and bath

applied at 2.5 mM. Fresh vials of MNI-glutamate were used for each experiment.

726   **ETHICS**

Animal experimentation: these studies were performed in compliance with experimental protocols

(13-185, 15-002, 16-011 and 17-012) approved by the *Comité de déontologie de l'expérimentation*

sur les animaux (CDEA) of the University of Montreal.

REFERENCES

1.   Araya, R. Input transformation by dendritic spines of pyramidal neurons. *Frontiers in*
neuroanatomy **8** (2014).
2.   Araya, R., Vogels, T.P. & Yuste, R. Activity-dependent dendritic spine neck changes are correlated
with synaptic strength. *Proceedings of the National Academy of Sciences of the United States of America*
(2014).
3.   Matsuzaki, M*., et al.* Dendritic spine geometry is critical for AMPA receptor expression in
hippocampal CA1 pyramidal neurons. *Nature neuroscience* **4**, 1086-1092 (2001).
4.   Matsuzaki, M., Honkura, N., Ellis-Davies, G.C.R. & Kasai, H. Structural basis of long-term
potentiation in single dendritic spines. *Nature* **429**, 761-766 (2004).
5.   Harvey, C.D., Yasuda, R., Zhong, H.N. & Svoboda, K. The spread of Ras activity triggered by
activation of a single dendritic spine. *Science* **321**, 136-140 (2008).
6.   Nishiyama, J. & Yasuda, R. Biochemical Computation for Spine Structural Plasticity. *Neuron* **87**, 63-
75 (2015).
7.   Oh, W.C., Hill, T.C. & Zito, K. Synapse-specific and size-dependent mechanisms of spine structural
plasticity accompanying synaptic weakening. *Proceedings of the National Academy of Sciences of the*
United States of America **110**, E305-312 (2013).
8.   Hayashi-Takagi, A*., et al.* Labelling and optical erasure of synaptic memory traces in the motor
cortex. *Nature* **525**, 333-+ (2015).
9.   Feldman, D.E. The Spike-Timing Dependence of Plasticity. *Neuron* **75**, 556-571 (2012).
10.   Bi, G.Q. & Poo, M.M. Synaptic modifications in cultured hippocampal neurons: dependence on
spike timing, synaptic strength, and postsynaptic cell type. *The Journal of neuroscience : the official journal*
of the Society for Neuroscience **18**, 10464-10472 (1998).
11.   Zhang, L.I., Tao, H.W., Holt, C.E., Harris, W.A. & Poo, M.M. A critical window for cooperation and
competition among developing retinotectal synapses. *Nature* **395**, 37-44 (1998).
12.   Sjostrom, P.J., Turrigiano, G.G. & Nelson, S.B. Rate, timing, and cooperativity jointly determine
cortical synaptic plasticity. *Neuron* **32**, 1149-1164 (2001).
13.   Sjostrom, P.J. & Hausser, M. A cooperative switch determines the sign of synaptic plasticity in
distal dendrites of neocortical pyramidal neurons. *Neuron* **51**, 227-238 (2006).
14.   Froemke, R.C., Poo, M.M. & Dan, Y. Spike-timing-dependent synaptic plasticity depends on
dendritic location. *Nature* **434**, 221-225 (2005).
15.   Froemke, R.C., Letzkus, J.J., Kampa, B.M., Hang, G.B. & Stuart, G.J. Dendritic synapse location and
neocortical spike-timing-dependent plasticity. *Frontiers in synaptic neuroscience* **2**, 29 (2010).
16.   Tonnesen, J., Katona, G., Rozsa, B. & Nagerl, U.V. Spine neck plasticity regulates
compartmentalization of synapses. *Nature neuroscience* **17**, 678-685 (2014).
17.   Feldman, D.E. Timing-based LTP and LTD at vertical inputs to layer II/III pyramidal cells in rat barrel
cortex. *Neuron* **27**, 45-56 (2000).
18.   Nakamura, T., Barbara, J.G., Nakamura, K. & Ross, W.N. Synergistic release of Ca2+ from IP3-
sensitive stores evoked by synaptic activation of mGluRs paired with backpropagating action potentials.
Neuron **24**, 727-737 (1999).
19.   Hashimotodani, Y*., et al.* Phospholipase Cbeta serves as a coincidence detector through its Ca2+
dependency for triggering retrograde endocannabinoid signal. *Neuron* **45**, 257-268 (2005).
20.   Bender, V.A., Bender, K.J., Brasier, D.J. & Feldman, D.E. Two coincidence detectors for spike
timing-dependent plasticity in somatosensory cortex. *The Journal of neuroscience : the official journal of*
the Society for Neuroscience **26**, 4166-4177 (2006).
21.   Sjostrom, P.J., Turrigiano, G.G. & Nelson, S.B. Neocortical LTD via coincident activation of
presynaptic NMDA and cannabinoid receptors. *Neuron* **39**, 641-654 (2003).

22.   Araya, R., Jiang, J., Eisenthal, K.B. & Yuste, R. The spine neck filters membrane potentials.
Proceedings of the National Academy of Sciences of the United States of America **103**, 17961-17966
(2006).
23.   Richardson, R.J., Blundon, J.A., Bayazitov, I.T. & Zakharenko, S.S. Connectivity patterns revealed
by mapping of active inputs on dendrites of thalamorecipient neurons in the auditory cortex. *The Journal*
of neuroscience : the official journal of the Society for Neuroscience **29**, 6406-6417 (2009).
24.   Harnett, M.T., Makara, J.K., Spruston, N., Kath, W.L. & Magee, J.C. Synaptic amplification by
dendritic spines enhances input cooperativity. *Nature* **491**, 599-602 (2012).
25.   Jayant, K.*, et al.* Targeted intracellular voltage recordings from dendritic spines using quantum-
dot-coated nanopipettes. *Nature nanotechnology* **12**, 335-342 (2017).
26.   Hayashi, Y.*, et al.* Driving AMPA receptors into synapses by LTP and CaMKII: requirement for GluR1
and PDZ domain interaction. *Science* **287**, 2262-2267 (2000).
27.   Ashby, M.C., Maier, S.R., Nishimune, A. & Henley, J.M. Lateral diffusion drives constitutive
exchange of AMPA receptors at dendritic spines and is regulated by spine morphology. *The Journal of*
neuroscience : the official journal of the Society for Neuroscience **26**, 7046-7055 (2006).
28.   Holcman, D. & Triller, A. Modeling synaptic dynamics driven by receptor lateral diffusion.
Biophysical journal **91**, 2405-2415 (2006).
29.   Kusters, R., Kapitein, L.C., Hoogenraad, C.C. & Storm, C. Shape-induced asymmetric diffusion in
dendritic spines allows efficient synaptic AMPA receptor trapping. *Biophysical journal* **105**, 2743-2750
(2013).
30.   Holcman, D. & Schuss, Z. Diffusion laws in dendritic spines. *Journal of mathematical neuroscience*
1, 10 (2011).
31.   Shim, S.H.*, et al.* Super-resolution fluorescence imaging of organelles in live cells with
photoswitchable membrane probes. *Proceedings of the National Academy of Sciences of the United States*
of America **109**, 13978-13983 (2012).
32.   Frost, N.A., Shroff, H., Kong, H.H., Betzig, E. & Blanpied, T.A. Single-Molecule Discrimination of
Discrete Perisynaptic and Distributed Sites of Actin Filament Assembly within Dendritic Spines. *Neuron* **67**,
86-99 (2010).
33.   Allison, D.W., Gelfand, V.I., Spector, I. & Craig, A.M. Role of actin in anchoring postsynaptic
receptors in cultured hippocampal neurons: Differential attachment of NMDA versus AMPA receptors.
Journal of Neuroscience **18**, 2423-2436 (1998).
34.   Hanley, J.G. Actin-dependent mechanisms in AMPA receptor trafficking. *Front Cell Neurosci* **8**
(2014).
35.   Kim, C.H. & Lisman, J.E. A role of actin filament in synaptic transmission and long-term
potentiation. *Journal of Neuroscience* **19**, 4314-4324 (1999).
36.   Honkura, N., Matsuzaki, M., Noguchi, J., Ellis-Davies, G.C.R. & Kasai, H. The subspine organization
of actin fibers regulates the structure and plasticity of dendritic spines. *Neuron* **57**, 719-729 (2008).
37.   Ramachandran, B. & Frey, J.U. Interfering with the actin network and its effect on long-term
potentiation and synaptic tagging in hippocampal CA1 neurons in slices in vitro. *The Journal of*
neuroscience : the official journal of the Society for Neuroscience **29**, 12167-12173 (2009).
38.   Lynch, G., Larson, J., Kelso, S., Barrionuevo, G. & Schottler, F. Intracellular injections of EGTA block
induction of hippocampal long-term potentiation. *Nature* **305**, 719-721 (1983).
39.   Malenka, R.C., Kauer, J.A., Zucker, R.S. & Nicoll, R.A. Postsynaptic calcium is sufficient for
potentiation of hippocampal synaptic transmission. *Science* **242**, 81-84 (1988).
40.   Artola, A. & Singer, W. Long-term depression of excitatory synaptic transmission and its
relationship to long-term potentiation. *Trends Neurosci* **16**, 480-487 (1993).
41.   Cummings, J.A., Mulkey, R.M., Nicoll, R.A. & Malenka, R.C. Ca2+ signaling requirements for long-
term depression in the hippocampus. *Neuron* **16**, 825-833 (1996).

42.   Fino, E.*, et al.* Distinct coincidence detectors govern the corticostriatal spike timing-dependent
plasticity. *The Journal of physiology* **588**, 3045-3062 (2010).
43.   Lisman, J. A mechanism for the Hebb and the anti-Hebb processes underlying learning and
memory. *Proceedings of the National Academy of Sciences of the United States of America* **86**, 9574-9578
(1989).
44.   Ismailov, I., Kalikulov, D., Inoue, T. & Friedlander, M.J. The kinetic profile of intracellular calcium
predicts long-term potentiation and long-term depression. *The Journal of neuroscience : the official
journal of the Society for Neuroscience* **24**, 9847-9861 (2004).
45.   Nevian, T. & Sakmann, B. Spine Ca2+ signaling in spike-timing-dependent plasticity. *Journal of
Neuroscience* **26**, 11001-11013 (2006).
46.   Graupner, M. & Brunel, N. Mechanisms of induction and maintenance of spike-timing dependent
plasticity in biophysical synapse models. *Frontiers in computational neuroscience* **4** (2010).
47.   Rubin, J.E., Gerkin, R.C., Bi, G.Q. & Chow, C.C. Calcium time course as a signal for spike-timing-
dependent plasticity. *J Neurophysiol* **93**, 2600-2613 (2005).
48.   Penn, A.C.*, et al.* Hippocampal LTP and contextual learning require surface diffusion of AMPA
receptors. *Nature* **549**, 384-388 (2017).
49.   Tanaka, J.I.*, et al.* Protein synthesis and neurotrophin-dependent structural plasticity of single
dendritic spines. *Science* **319**, 1683-1687 (2008).
50.   El-Boustani, S.*, et al.* Locally coordinated synaptic plasticity of visual cortex neurons in vivo.
Science **360**, 1349-1354 (2018).
51.   Nevian, T., Larkum, M.E., Polsky, A. & Schiller, J. Properties of basal dendrites of layer 5 pyramidal
neurons: a direct patch-clamp recording study. *Nature neuroscience* **10**, 206-214 (2007).
52.   Markram, H., Lubke, J., Frotscher, M., Roth, A. & Sakmann, B. Physiology and anatomy of synaptic
connections between thick tufted pyramidal neurones in the developing rat neocortex. *J Physiol-London*
500, 409-440 (1997).
53.   Schmidt, H.*, et al.* Axonal synapse sorting in medial entorhinal cortex. *Nature* **549**, 469-475 (2017).
54.   Kasthuri, N.*, et al.* Saturated Reconstruction of a Volume of Neocortex. *Cell* **162**, 648-661 (2015).
55.   Bloss, E.B.*, et al.* Single excitatory axons form clustered synapses onto CA1 pyramidal cell
dendrites. *Nature neuroscience* **21**, 353-363 (2018).
56.   Yang, S.N., Tang, Y.G. & Zucker, R.S. Selective induction of LTP and LTD by postsynaptic [Ca2+]i
elevation. *J Neurophysiol* **81**, 781-787 (1999).
57.   Karmarkar, U.R. & Buonomano, D.V. A model of spike-timing dependent plasticity: one or two
coincidence detectors? *J Neurophysiol* **88**, 507-513 (2002).
58.   Shouval, H.Z., Bear, M.F. & Cooper, L.N. A unified model of NMDA receptor-dependent
bidirectional synaptic plasticity. *Proceedings of the National Academy of Sciences of the United States of
America* **99**, 10831-10836 (2002).
59.   Schiemann, J.*, et al.* Cellular mechanisms underlying behavioral state-dependent bidirectional
modulation of motor cortex output. *Cell Rep* **11**, 1319-1330 (2015).
60.   Wilson, D.E., Whitney, D.E., Scholl, B. & Fitzpatrick, D. Orientation selectivity and the functional
clustering of synaptic inputs in primary visual cortex. *Nature neuroscience* **19**, 1003-1009 (2016).
61.   Scholl, B., Wilson, D.E., Fitzpatrick, D. Local Order within Global Disorder: Synaptic Architecture of
Visual Space. in *Neuron* 1127-1138 (2017).
62.   Beaulieu-Laroche, L. & Harnett, M.T. Dendritic Spines Prevent Synaptic Voltage Clamp. *Neuron* **97**,
75-82 e73 (2018).
63.   Araya, R., Eisenthal, K.B. & Yuste, R. Dendritic spines linearize the summation of excitatory
potentials. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 18799-
18804 (2006).

64.   Fino, E*., et al.* RuBi-Glutamate: Two-Photon and Visible-Light Photoactivation of Neurons and
Dendritic spines. *Frontiers in neural circuits* **3**, 2 (2009).
65.   Rasband, W.S. & Bright, D.S. Nih Image - a Public Domain Image-Processing Program for the
Macintosh. *Microbeam Anal* **4**, 137-149 (1995).
66.   Nimchinsky, E.A., Yasuda, R., Oertner, T.G. & Svoboda, K. The number of glutamate receptors
opened by synaptic stimulation in single hippocampal spines*. The Journal of neuroscience : the official
journal of the Society for Neuroscience* **24**, 2054-2064 (2004).
67.   Holtmaat, A.J*., et al.* Transient and persistent dendritic spines in the neocortex in vivo. *Neuron* **45**,
279-291 (2005).
68.   Ruttinger, S*., et al.* Comparison and accuracy of methods to determine the confocal volume for
quantitative fluorescence correlation spectroscopy. *J Microsc* **232**, 343-352 (2008).

Tazerart, Mitchell et al.

Acknowledgements. We thank A. Kolta and P.J Sjöström for critical discussion and reading of the manuscript, and are grateful to all other members of Roberto Araya's laboratory for kind support. We also thank members of the *Groupe de Recherche sur le système nerveux central* (GRSNC) for support and equipment shearing. This work was funded by the Canadian Institutes of Health Research (CIHR) grant MOP-133711 to R.A., a Canada Foundation for Innovation (CFI) equipment grant *Fonds des leaders* 29970 to R.A., and a Natural Sciences and Engineering Research Council of Canada (NSERC Discovery Grant) grant application No. 418113-2012 (NSERC PIN 392027) to R.A. S.T. was supported in part by a salary support from the GRSNC at Université of Montréal. D.E.M. was supported in part by a postdoctoral fellowship from the Fonds de recherche du Québec – Santé (FRQS).

Author contributions. R.A. conceived the project. S.T. and D.E.M. performed the experiments. S.T. and D.E.M. performed data analyses. S.M-R. performed control experiments. R.A., S.T., and D.E.M. designed experiments. R.A. and D.E.M. wrote the manuscript. R.A. supervised the project. All authors read and approved the contents of the manuscript.

Competing financial interests. The authors declare no competing financial interest.

Materials & Correspondence. Correspondence and material requests should be addressed to R.A.

Figure 1: **Induction of t-LTP in single dendritic spines**. (A) Experimental protocol for the

induction of t-LTP with pre-post pairings (two-photon (2P) uncaging of glutamate followed by a

bAP) of +7 and +13 ms in single dendritic spines (sp). (B) Example of a t-LTP protocol with pre-

post pairing protocol of + 13 ms. (B.1) Average uEPSP response recorded via a whole-cell pipette

at the soma of L5 pyramidal neurons before (Control) and after the induction of t-LTP in a selected

dendritic spine located in basal dendrites (B.2). Traces in B.1 correspond to an average of 10

depolarizations generated by 2P uncaging over the indicated spine (B.2 red dot) before (black

trace) and after the induction of t-LTP (red trace). (C.1) Time course of uEPSP amplitude (black

line), neck length (red line) and spine head volume (blue line) changes over the course of 40 min

following STDP induction in individual spines at a pre-post timing of + 13 ms. *ns*, not significant;

*P < 0.05; **P < 0.01; ***P < 0.001, one-way repeated measures ANOVA followed by post hoc

Dunnet's test. (C.2) Maximum changes in uEPSP amplitude (black bar and dots) and concomitant

changes in neck length (red bar and dots) and head volume (blue bar and dots) of the activated

spine after the induction of t-LTP at a pre-post timing of +13 ms. *P < 0.05, ***P < 0.001, paired

t-test. (D.1) Time course of uEPSP amplitude (black line), neck length (red line) and spine head

volume (blue line) changes over the course of ~40 min following STDP induction in individual

spines at a pre-post timing of + 7 ms. *ns*, not significant; *P < 0.05, one-way repeated measures

ANOVA followed by post hoc Dunnet's test. (D.2) Maximum changes in uEPSP amplitude (black

bar and dots) and corresponding changes in neck length (red bar and dots) and head volume (blue

bar and dots) of the activated spine after the induction of t-LTP at a pre-post timing of +7 ms.

Shaded light area in C.1 and D.1, and error bars in C.2 and D.2 represent ±SEM. Time 0 in C.1

and D.1 is the time when the 40 pre-post repetitions of the induction protocol were completed.

Figure 2: **Induction of t-LTD in single dendritic spines**. (A) Experimental protocol for the

induction of t-LTD with post-pre pairings protocols of -15 and -23 ms in single dendritic spines.

(B) Example of a t-LTD protocol with post-pre pairings protocol of -15 ms. (B.1) Average uEPSP

response recorded via a whole-cell pipette at the soma of L5 pyramidal neurons before (Control)

and after the induction of t-LTD in selected dendritic spines located in basal dendrites (B.2). Traces

in B.1 correspond to an average of 10 uEPSP generated by the 2P uncaging of glutamate at the

indicated spine (B.2, red dot) before (black trace) and after the induction of t-LTD (red trace).

(C.1) Time course of uEPSP amplitude (black line), neck length (red line) and spine head volume

(blue line) changes over the course of ~35 min following STDP induction in individual spines at

a post-pre timing of -15 ms. *ns*, not significant; ***$P < 0.001$, one-way repeated measures ANOVA

followed by post hoc Dunnet's test. (C.2) Maximum changes in uEPSP amplitude (black bar and

dots) and concomitant changes in neck length (red bar and dots) and head size of the activated

spine (blue bar and dots) after the induction of t-LTD at a post-pre timing of -15 ms. **$P < 0.01$,

paired t test. (D.1) Time course of uEPSP amplitude (black line), neck length (red line) and spine

head volume (blue line) changes over the course of ~ 40 min following STDP induction in

individual spines at a post-pre timing of -23 ms. *ns*, not significant, one-way repeated measures

ANOVA followed by post hoc Dunnet's test. (D.2) Maximum changes in uEPSP amplitude (black

bar and dots) and corresponding changes in neck length (red bar and dots) and head volume (blue

bar and dots) of the activated spine after the induction of t-LTD at a post-pre timing of -23 ms.

Shaded light area in C.1 and D.1, and error bars in C.2 and D.2 represent ±SEM. Time 0 in C.1

and D.1 is the time when the 40 pre-post repetitions of the induction protocol were completed. (E)

STDP learning rule for single dendritic spines: Plot illustrating the maximum changes in uEPSP

amplitude (black data points), and concomitant changes in neck length (red data points) and head

volume (blue data points) after the induction of t-LTP (temporal offset +13 ms and +7 ms) and t-

LTD (temporal offset -15 ms and -23 ms). *P < 0.05, **P < 0.01, paired t test. (F) Diagram

indicating the uEPSP amplitude and spine neck morphological changes observed after the

induction of t-LTP and t-LTD.

Figure 3: **Induction of t-LTP in clustered dendritic spines** (A) Experimental protocol for the

induction of t-LTP (pre-post timing of + 7 ms) in two clustered dendritic spines (< 5 μm distance

between spines). (B) Example of one experiment where two neighbouring dendritic spines were

activated with a pre-post t-LTP protocol. (B.1) Average uEPSP response recorded via a whole-cell

pipette at the soma of L5 pyramidal neurons before (Control) and after the induction of t-LTP in

two selected dendritic spines located in a basal dendrite (red dots, B.2). Traces in B.1 correspond

to an average of 10 uEPSP generated by the 2P uncaging at the indicated spines (B.2) before (black

trace) and after the induction of t-LTP (red trace). (C.1) Time course of uEPSP amplitude (black

line), neck length (red line) and spine head volume (blue line) changes over the course of ~25 min

following STDP induction in clustered spines at a pre-post timing of + 7 ms. *ns*, not significant;

**P < 0.01, ***P < 0.001, one-way repeated measures ANOVA followed by post hoc Dunnet's

test. (C.2) Pooled data of the maximum changes in uEPSP amplitude (black bar and dots) and

concomitant changes in neck length (red bar and dots) and head volume (blue bar and dots) of

individual (1sp) or clustered spines (2sp) after the induction of t-LTP at a pre-post timing of +7ms.

**P < 0.01, paired t test. Shaded light area in C.1 and error bars in C.2 represent ±SEM. Time 0 in

C.1 is the time when the 40 pre-post repetitions of the induction protocol are completed. NL =

neck length, HV = head volume.

Figure 4: **Molecular mechanisms responsible for the induction of t-LTP**. (A) Experimental

protocol for the induction of t-LTP (pre-post timing of + 7 ms) in two clustered dendritic spines

(< 5 µm) in the presence of PEP1-TGL (200 µM). (B) An example of one experiment where two

neighbouring dendritic spines were activated with a pre-post t-LTP protocol in the presence of

PEP1-TGL. (B.1) Average uEPSP recorded via a whole-cell pipette at the soma of L5 pyramidal

neurons before (Control) and after the induction of t-LTP in two selected dendritic spines located

in basal dendrites (B.2). Traces in B.1 correspond to an average of 10 uEPSP generated by the 2P

uncaging of glutamate at the two indicated spines (red dots, B.2) before (black trace) and after the

induction of t-LTP in the presence of PEP1-TGL (red traces). (C.1) Time course of uEPSP

amplitude (black line), neck length (red line) and spine head volume (blue line) changes over the

course of ~28 min following STDP induction in clustered spines at a pre-post timing of + 7 ms in

the presence of PEP1-TGL. *ns*, not significant; **P < 0.01, ***P < 0.001, one-way repeated

measures ANOVA followed by post hoc Dunnet's test. (C.2) Pooled data of the maximum changes

in uEPSP amplitude (black bar and dots) and concomitant changes in neck length (red bar and

dots) and head volume (blue bar and dots) of the activated clustered spines after the induction of

t-LTP at a pre-post timing of +7ms in control conditions (Cont) and in the presence of PEP1-TGL

(PEP1-TGL). *P < 0.05, **P < 0.01, paired t test. Time 0 is the time when the 40 pre-post

repetitions of the induction protocol were completed. (D) Experimental protocol for the induction

of t-LTP (pre-post timing of + 7 ms) in two clustered dendritic spines (< 30µm) in the presence

Latrunculin-A (Lat-A, 100nM). (E) Example of one experiment where two neighbouring dendritic

spines were activated with a pre-post t-LTP protocol in the presence of Lat-A. (E.1) Average

uEPSP recorded via a whole-cell pipette at the soma of L5 pyramidal neurons before (Control) and

after the induction of t-LTP in two selected dendritic spines (red dots, E.2) located in basal

dendrites (E.2). Traces in E.1 correspond to an average of 10 uEPSP generated by the 2P uncaging

over the two indicated spines (red dots, E.2) before (black trace) and after the induction of t-LTP

in the presence of Lat-A (red trace). (F.1) Time course of uEPSP amplitude (black line), neck

length (red line) and spine head volume (blue line) changes over the course of ~12 min following

STDP induction in clustered spines at a pre-post timing of + 7 ms in the presence of Lat-A. *ns*, not

significant; **P < 0.01, ***P < 0.001, one-way repeated measures ANOVA followed by post hoc

Dunnet's test. (F.2) Pooled data of the maximum changes in uEPSP amplitude (black bar and dots)

and concomitant changes in neck length (red bar and dots) and head volume (blue bar and dots) of

the activated clustered spines after the induction of t-LTP at a pre-post timing of +7 ms in control

conditions (Cont) and in the presence of Lat-A (Lat-A). **P < 0.01, ***P < 0.001, paired t test.

NL = neck length, HV = head volume. Shaded light area in C.1 and F.1 and error bars in C.2 and

F.2 represent ±SEM. Time 0 is the time when 40 pre-post repetitions of the induction protocol

were completed.

Figure 5. Induction of t-LTD in clustered and distributed dendritic spines: (A) Experimental

protocol for the induction of t-LTD protocol (pairings of - 15 ms) in two clustered dendritic spines

(< 40 µm). (B) An example of one experiment where two neighbouring dendritic spines were

activated with a post-pre t-LTP protocol of -15 ms. (B.1) Average uEPSP recorded via a whole-

cell pipette at the soma of L5 pyramidal neurons before (Control) and after the induction of t-LTD

in two selected dendritic spines (red dots, B.2) located in basal dendrites. Traces in B.1 correspond

to an average of 10 uEPSP generated by 2P uncaging of glutamate at the indicated spines before

(black trace) and after the induction of t-LTD (red trace). (C.1) Time course of uEPSP amplitude

(black line), the neck length (red line) and spine head volume (blue line) of the activated clustered

spines after the induction of t-LTD at pairings of -15 ms. *ns*, not significant; *P < 0.05, one-way

repeated measures ANOVA followed by post hoc Dunnet's test. (C.2) Pooled data of the maximum

changes in uEPSP amplitude (black bar and dots) and concomitant changes in neck length (red bar

and dots) and head volume (blue bar and dots) of individual (1sp) or clustered spines (2sp) after

the induction of t-LTD at a post-pre timing of -15 ms. *P < 0.05, **P < 0.01, paired t test. Time

0 in C.1 is the time when the 40 pre-post repetitions of the induction protocol were completed. NL

= neck length, HV = head volume. (D) Experimental protocol for the t-LTD induction protocol

(pre-post timing of – 15 ms) in two distributed dendritic spines (> 40 µm). (E) An example of one

experiment where two distributed dendritic spines were activated with a post-pre t-LTP induction

protocol of -15 ms. (E.1) Average uEPSP recorded via a whole-cell pipette at the soma of L5

pyramidal neurons before (Control, underneath image, E.2) and 3 min after the induction of t-LTD

(E.2, over imposed image) in two selected dendritic spines located in basal dendrites. (E.2) Inset

shows a low magnification image of the recorded neuron with the marked location of the selected

spines. Traces in E.1 correspond to an average of 10 uEPSP generated by 2P uncaging at the

indicated spines (red dots, E.2) before (black trace) and after the induction of t-LTD (red trace).

(F.1) Time course of uEPSP amplitude (black line), the neck length (red line) and spine head

volume (blue line) of the activated distributed spines (> 40 µm) after the induction of t-LTD at

pairings of -15 ms. *ns*, not significant; *P < 0.05, **P < 0.01, one-way repeated measures ANOVA

followed by post hoc Dunnet's test. (F.2) Pooled data of the maximum changes in uEPSP

amplitude (black bar and dots) and corresponding changes in neck length (red bar and dots) and

head volume (blue bar and dots) of clustered (2sp at < 40 µm) or distributed spines (2sp at > 40

µm) after the induction of t-LTD at a post-pre timing of -15ms. *P < 0.05, paired t test. Shaded

light area in C.1 and F.1 and error bars in C.2 and F.2 represent ±SEM. Time 0 in C.1 and F.1 is

the time when the 40 pre-post repetitions of the induction protocol were completed. NL = neck

length, HV = head volume.

1040

Figure 6. Calcium dynamics in single and two clustered spines during pre-post pairing

protocol. (A.1) Single 2P images of a spine and dendrite from a L5 pyramidal neuron loaded with

Alexa 594 (100uM) and Fluo4 (300uM). Red ellipses and blue polygons indicate the ROIs selected

for the calcium signal analysis. (A.2) Two photon calcium signal images before (left panels) and

after (right panels) a pre-post pairing protocol (+7 ms). The 1^{st}, 2^{nd}, and 40^{th} repetitions of the

pairing protocol are shown here. The change in calcium fluorescence from baseline (ΔF) is color

coded. (B) Calcium signals ($\Delta F/F$) in the spine (B.1) and dendrite (B.2) from the experiment

depicted in A before (dotted lines) and after (solid lines) the pairing protocol. (C) Population

averages of the calcium signals ($\Delta F/F$) measured in spines (C.1) and dendrites (C.2) before the

pairing protocol performed in 1 spine (left panels) and 2 spines (middle panels). The right panels

shows the superimposed $\Delta F/F$ population averages in 1 spine (black lines) and 2 spines (green

lines). (D.1) Single 2P images of two clustered spines and dendrite from a L5 pyramidal neuron

loaded with Alexa 594 (100uM) and Fluo4 (300uM). Red ellipses and blue polygons indicate the

ROIs selected for the calcium signal analysis. (D.2) Two photon calcium signal images before (left

panel) and after (right panels) a pre-post pairing protocol (+ 7ms). The 1^{st}, 2^{nd}, and 40^{th} repetitions

of the pairing protocol are shown here. The change in calcium fluorescence from baseline (ΔF) is

color coded. (E) Calcium signals ($\Delta F/F$) in the spines (E.1) and dendrite (E.2) before (dotted lines)

and after (solid lines) the pairing protocol. (F) Population averages of the calcium signals ($\Delta F/F$)

measured in spines (F1) and dendrites (F.2) after the pairing protocol performed in one spine (left

panels) and two spines (middle panels). The right panels shows the superimposed $\Delta F/F$ population

averages in one spine (black lines) and two spines (green lines). *ns*, not significant; *$P < 0.05$; **P

< 0.01; ***$P < 0.001$, one-way repeated measures ANOVA followed by post hoc Dunnet's test.

Figure 7. Calcium dynamics in single and two clustered spines during post-pre pairing

protocol. (A1) Single 2P images of a spine and dendrite from a L5 pyramidal neuron loaded with

Alexa 594 (100uM) and Fluo4 (300uM). Red ellipses and blue polygons indicate the ROIs selected

for the calcium signal analysis. (A.2) Two photon calcium signal images before (left panels) and

after (right panels) a post-pre pairing protocol (-15 ms). The 1^{st}, 2^{nd}, and 40^{th} repetitions of the

pairing protocol are shown here. The change in calcium fluorescence from baseline (ΔF) is color

coded. (B) Calcium signals ($\Delta F/F$) in the spine (B.1) and dendrite (B.2) from the experiment

depicted in A before (dotted lines) and after (solid lines) the pairing protocol. (C) Population

averages of the calcium signals ($\Delta F/F$) measured in spines (C.1) and dendrites (C.2) before the

pairing protocol performed in one spine (left panels) and two spines (middle panels). The right

panels shows the superimposed $\Delta F/F$ population averages in one spine (black lines) and two spines

(green lines). (D.1) Single 2P images of two clustered spines and dendrite from a L5 pyramidal

neuron loaded with Alexa 594 (100uM) and Fluo4 (300uM). Red ellipses and blue polygons

indicate the ROIs selected for the calcium signal analysis. (D.2) Two photon calcium signal images

before (left panel) and after (right panels) a pre-post pairing protocol (-15 ms). The 1^{st}, 2^{nd}, and

40^{th} repetitions of the pairing protocol are shown here. The change in calcium fluorescence from

baseline (ΔF) is color coded. (E) Calcium signals ($\Delta F/F$) in the spines (E.1) and dendrite (E.2)

before (dotted lines) and after (solid lines) the pairing protocol. (F) Population averages of the

calcium signals ($\Delta F/F$) measured in spines (F1) and dendrites (F2) after the pairing protocol

performed in one spine (left panels) and two spines (middle panels). The right panels shows the

superimposed $\Delta F/F$ population averages in 1 spine (black lines) and 2 spines (green lines). *ns*, not

significant; *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$, one-way repeated measures ANOVA followed

by post hoc Dunnet's test.

1086

Figure 8. STDP learning rule for single distributed and clustered dendritic spines. (A) STDP

learning rule in the basal dendrites of L5 pyramidal neurons as a function of the structural

organization of excitatory inputs in basal dendrites of L5 pyramidal neurons. Note how STDP in

single, or distributed spines (separated by > 40 μm), follow a bidirectional Hebbian STDP rule that

can be disrupted by synaptic cooperativity [co-activation of two clustered spines]. We propose that

synaptic cooperativity generates local dendritic depolarization high enough to disrupt bidirectional

STDP, leading to STDP that only encompasses LTP. (B) Model showing a hypothetical

relationship between postsynaptic calcium levels and the induction of plasticity. Large levels of

calcium (above a potentiation threshold, Θ_{pSTART}) are thought to lead to t-LTP whereas more

moderate, prolonged levels (between the depression threshold, Θ_{dSTART}, and Θd_{STOP}) give rise to

t-LTD and a mid-level range in which neither t-LTP nor t-LTD occur (below Θ_{dSTART}). The average

linear fits of the calcium signal measured in all experiments throughout the different STDP

induction protocols are superimposed.

Figure 1

Figure 2

Figure 3

A

Pairing Protocol pre-post (+7ms)

Δt = 7 ms   +Δt   40 x

2 s

2P Glu Uncaging 2 sp

bAP

Before t-LTP   ⇌   After t-LTP

STDP-LTP

B.1

9 min after

Control

0.5 mV

20 ms

B.2

Control   9 min

2 µm

C.1

Clustered spines:
Positive spike timing (+7ms)

uEPSP, Neck Length & Head Volume changes (%)

ns   ns   ns   ns   ns
**   ***   ***   ***   ***
ns   ***   ***   **   ***

Time (min)

C.2

uEPSP, Neck Length & Head Volume changes (%)

**   **

uEPSP 1sp uEPSP 2sp NL 1sp NL 2sp HV 1sp HV 2sp

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8