The geometry of dependence: solitary bee larvae prioritize carbohydrate over protein in parentally provided pollen

Alexander J. Austin[1,2†] and James D. J. Gilbert[1]

[1]Department of Biological and Marine Sciences, University of Hull, UK

[2]Environment & Sustainability, Ku-ring-gai Municipal Council, NSW, Australia

[†]To whom correspondence should be addressed. Email: alexander.j.austin17@gmail.com

Acknowledgements

The authors thank Victor Swetez, Toby Bagnall, Fiona Tainsh and Shannon Woodmansey for logistical assistance; Stephen Simpson, Audrey Dussutour, Mathieu Lihoreau, Michal Filipiak, Lori Lawson Handley and Domino Joyce for valuable discussions and comments on the manuscript. The study was funded by the University of Hull.

Author Contributions

JDJG and AA conceived and designed the study. AA gathered and analysed all data. Both authors wrote and edited the manuscript.

Abstract

1. Most organisms must regulate their nutritional intake in an environment full of complex food choices. While this process is well understood for self-sufficient organisms, dependent offspring, such as bee larvae, in practice have limited food choices because food is provided by parents. Nutrient balancing may therefore be achieved by offspring, by parents on offspring's behalf, or cooperatively by both.

2. We used the Geometric Framework to investigate the capacity of dependent larval mason bees (*Osmia bicornis*) to regulate their intake of protein and carbohydrate. Herbivorous insect larvae are typically expected to balance protein and carbohydrate to maximise growth and reproduction.

3. Contrary to prediction, carbohydrate and not protein mediated both growth and survival to pupation. Accordingly, larvae prioritised maintaining a constant intake of carbohydrate while tolerating excesses and deficiencies of protein, and self-selected a relatively carbohydrate biased diet compared to other hymenopterans. Reasons for prioritising carbohydrate may include (1) the relative abundance of protein in their normal pollen diet, (2) the relative paucity of nectar in parental provisions making carbohydrate a scarce resource, or (3) the requirement for diapause for all *O. bicornis* larvae. Larvae were intolerant of moderate dietary dilution, reflecting an evolutionary history of nutrient-dense food.

4. Our results demonstrate that dependent *O. bicornis* offspring remain capable of balancing nutrition and thus play an active role in their own nutrient balancing, but they are nevertheless tolerant of wide swings in protein, an important macronutrient for development. Research should now aim to establish whether protein is regulated by parents, or by larvae via post-ingestive processing.

Keywords: bees, parental care, offspring provisioning, nutritional ecology, nutritional geometry, limiting nutrient, diapause, environmental change

Introduction

Most animals manage their nutrient intake by combining nutritionally different foods (Simpson and Raubenheimer 2012). However, the importance of this ability depends upon the nutritional variability of the animals' typical food (Despland and Noseworthy 2006; Raubenheimer et al. 2009); extreme specialists, for example, can lose the capacity to regulate nutrition (Warbrick-Smith et al. 2009; Poissonnier et al. 2018 Aug 19). One way in which organisms can experience limited nutritional choice is if they are dependent upon others for nutrition, or "alloregulation" (Lihoreau et al. 2014), such as dependent offspring of altricial birds, human toddlers, and many larval insects. Offspring tend to have different requirements from parents (Harper and Turner 2000; Michaelsen et al. 2003) - particularly for protein, given their elevated rates of somatic growth and development. Accordingly, parents often make different nutritional choices for their offspring versus when foraging for themselves (Royama 1970; Dussutour and Simpson 2009; Burt and Amin 2014). For example, granivorous birds usually provision young with insects, rather than the seed diets of adults, to fulfil protein requirements (Wiens and Johnston 2012). Under these circumstances, by what rules offspring regulate their own consumption should depend upon provisioning rules of parents. If parents alloregulate offspring nutrition tightly, then offspring should have no need for self-regulation, like extreme specialists (Poissonnier et al. 2018).

The Geometric Framework for Nutrition (GF) allows us to investigate foraging decisions made by animals in multi-dimensional "nutrient space" (Simpson and Raubenheimer 1993). The GF can be used to determine animals' nutritional choices relative to their "intake target" - the optimal amount and balance of multiple macronutrients - as well as their "rule of compromise" that governs their choices when restricted to suboptimal food (Raubenheimer and Simpson 1999a). The GF has provided insights into the nutritional ecology of a broad range of taxa (reviewed in Simpson and Raubenheimer 2012). Its application to dependent offspring, though, has typically been as part of studies of social insect systems (e.g. Helm et al. 2017) and often inferring offspring requirements from patterns of alloparental feeding in studies more broadly focused on adult foraging (see Dussutour and Simpson 2009; Cook et al. 2010; Vaudo et al. 2016). In such systems, multiple adults normally contact offspring, progressively feeding and adjusting nutrition in response to feedback (Field 2005; Schmickl and Karsai 2017), making the responses of individual larvae difficult both to follow and interpret.

In solitary bees, by contrast, typically females provision offspring individually with a pollen ball before sealing the cell and leaving. This behaviour makes solitary bees an ideal, manipulable model for directly studying the nutrition of dependent larvae (Strohm et al. 2002). Larvae of bees, like most aculeate hymenopterans, rely on parents or alloparents for nutrition (Field 2005). Nutritional requirements for bee adults and offspring differ, often radically (Weeks et al. 2004; Filipiak 2019); adults primarily feed on carbohydrate-rich nectar (although see Cane 2016) and larvae feed mostly on protein-rich pollen (Filipiak 2019). Solitary bees, along with most other hymenopterans and many other parental insects, typically have a simple one-to-one parent-offspring relationship whereby parents "mass provision" their young, providing a finite, fixed-mass food provision, and have no contact with their young during development (Costa 2006). Such systems are almost unstudied in a rigorous nutritional context (but see e.g. Roulston and Cane 2002). In these species, there is no opportunity for parents to adjust nutrition according to offspring feedback, and the larva must therefore make the best of what it is given. It may be that offspring regulate their own nutrition to compensate for variation, as in more independent insect larvae (Lee et al. 2002). Alternatively, they may have lost this capacity, like extreme specialists (Warbrick-Smith et al. 2009; Poissonnier et al. 2018). We know very little about how bee larvae deal with variable nutrition (but see Helm et al. 2017) - a knowledge gap with potentially important consequences, considering the proposed link between nutritional stress and bee declines (Roulston and Goodell 2011; Goulson et al. 2015).

In this study, we used a commercially important solitary bee species, *Osmia bicornis*, to investigate how dependent larvae cope with varying nutrition, and whether they can regulate their own intake. *O. bicornis* are pollen generalists (Falk 2015) and the pollen balls that parents provide to offspring are variable in species composition (Haider et al. 2014). Natural variation in pollen ball nutrient content is largely unquantified (although see Budde and Lunau 2007), so there is no prior expectation about the capacity of larvae to regulate their consumption. We used a classic GF design (e.g. Lee et al. 2008), focusing on protein and carbohydrate, with two experimental phases. In the first "no-choice" phase we raised *O. bicornis* larvae on fixed diets of differing protein to carbohydrate ratios to determine their rules of compromise and the diet composition that maximised fitness. In a second "choice" phase, we then provided larvae with targeted choices between sets of two imbalanced diets that differed in their protein:carbohydrate ratios to determine whether larvae defend an intake target. Given the central role of protein in

growth of insect larvae (Scriber and Slansky 1981; Behmer 2009), we predicted that (1) protein

would be a key driver of fitness in larval *O. bicornis*, (2) larvae would accordingly aim for a

relatively protein-biased intake target, and (3) larvae would prioritize regulating intake of protein

over carbohydrate.

Methods

Study organism

Osmia bicornis is a common, cavity-nesting solitary bee (Falk 2015), and a commercially

important pollinator (Jauker et al. 2012). *O. bicornis* larvae were obtained as diapausing adults in

cocoons (Mauerbienen®). These were released at the nesting site at the University of Hull in

April 2017, and emerging adults allowed to breed. Early trials revealed that fresh eggs and

newly emerged larvae were too fragile for manipulation. Therefore, newly emerged larvae were

left alone for two days before we transferred them to a single-occupancy nest and assigned each

to an experimental treatment. Details of nesting apparatus and monitoring protocols are available

in the supplementary methods.

Diet Formulation

Existing artificial diet protocols for solitary bees have met with limited success in terms

of larval survival (Nelson et al. 1972; Fichter et al. 1981). We used six diets, consisting of three

different protein:carbohydrate (P:C) ratios (Diet A = 1:1.2, Diet B = 1:2.3 & Diet C = 1:3.4) and

two total macronutrient concentrations (concentration 1 = 90% nutrients, 10% diluent, or

concentration 2 = 70% nutrients, 30% diluent; see table S1). Diet ratios were chosen based on a

combination of the nutrient ratios in honeybee-collected pollen loads and published data for

protein content of *O. bicornis* pollen balls (Budde and Lunau 2007). Diets were diluted with

sporopollenin, the primary constituent of the exine of pollen (Mackenzie et al. 2015) an

extraordinarily stable natural polymer. Sporopollenin is a novel dietary diluent for bees; we have

demonstrated its suitability in a separate study (manuscript submitted). For a more detailed

description of sporopollenin and its preparation, see supplementary methods.

Experiment 1: No-choice phase

Two-day-old larvae, randomized by parentage, were allocated to one of 6 treatments

corresponding to our 6 artificial diets (n = 20/treatment). Provisions were made to resemble the

size of natural provisions (mean initial artificial provision weight = 0.323g +/- 0.034g). Once

provisioned, larvae were placed in an incubation chamber (Gallenkamp, IH-270) at 23°C and

80% RH. Provisions were replaced weekly, to avoid desiccation and mould formation, or when

fully consumed by larvae, ensuring the diet was always available in excess. Weight of provision

consumed was recorded upon provision replacement. Should they finish this provision, it was

replenished with a fresh one. A "water control" group, containing artificial diets but no larvae,

was used to track water loss from the diets, going through the same weighing regime as above

with weight loss recorded at each swap. Nests were checked daily for mortality. Cocoon weight

was recorded at the completion of pupation.

Experiment 2: Choice phase

In the choice experiment, 32 two-day-old larvae of mixed parentage were randomly

divided among four treatments. Treatments consisted of strategic pairwise combinations (see Fig

1; Table 1) of four possible diets: A1 (1P:1.2C, 90%), A2 (1P:1.2C, 70%), C1 (1P:3.4C, 90%),

C2 (1P:3.4C, 70%). Because *O. bicornis* larvae are sedentary and receive a single provision, it is

not biologically appropriate to present choices between two diets simultaneously. Therefore,

choices were offered temporally by swapping the provision every other day, presenting one diet

at a time. This required the larvae to differentially feed over time to compensate for temporal

imbalance, in order to converge on an intake target (see e.g. Raubenheimer and Jones 2006). All

larvae were kept on the same treatment from two days post-hatching up to pupation, whereupon

diet replenishment ceased. The diet that the larvae would be fed first was randomly assigned

prior to the experiment.

Statistical Analysis

All analyses were conducted in R version 3.4.2 (R Core Team, 2017). For the no-choice

experiment, we calculated total nutrients consumed (protein and carbohydrate) from raw diet

consumption data for each swap, adjusted for water loss and dilution. Values were then summed

for each larva.

To investigate consumption rules, including rules of compromise, we first asked whether

diet ratio and concentration affected consumption of (a) the total provision, (b) protein, or (c)

carbohydrate, using models of each respective variable with "ratio" and "concentration" as

predictors. Rules of compromise can include nonlinear effects (Simpson and Raubenheimer

1993); we investigated linear effects first using ordinary linear models, assessing goodness of fit

before progressing to evaluating nonlinear effects.

To assess fitness consequences of macronutrient consumption, we assessed cocoon

weight at pupation and survival to pupation. We analysed cocoon weight using a linear model

with "cocoon weight" as a predictor and "protein consumed", "carbohydrate consumed" and

their interaction as predictors. To analyse survival, we used parametric survival analysis in the

survival package in R. In both cases, we first assessed the fit of linear models before attempting

to fit more complex nonlinear effects. To visualise these fitness effects, we calculated response

surfaces for cocoon weight and survival, and visualised them using non-parametric thin-plate

splines.

In the choice experiment, the mean final consumption of each nutrient was investigated

using linear models with diet combination, dilution and their interaction as predictors, and using

Tukey's post hoc tests to compare individual treatments against each other. Under a null

expectation we would expect larvae to eat randomly from each diet (Fig 1). Thus, for each larva

we calculated the deviation from this null expectation. We then tested whether these values

systematically departed from zero for protein and carbohydrate, and whether these departures

from random consumption differed by treatment group. We used a linear model with "deviation

from random consumption" as the response variable and "treatment group" as a predictor.

Larvae that died pre-pupation were not used in the calculation of the mean protein and

carbohydrate consumption for diets in either experiment, or for cocoon weight, but were used in

analyses involving survival.

Results

No-choice phase

Dietary P:C ratio had a significant effect on the overall amount of provision consumed,

with larvae consuming more provision on high P:C ratio diets (Fig. 2a). Total consumption was

also affected by diet concentration; larvae on less concentrated diets consumed more provision,

indicating compensatory feeding (Table 2a, Fig. 2a).

Dietary P:C ratio had a strong effect on the total amount of P eaten; more protein was

eaten by larvae raised on the higher P:C diets (Fig. 3). Diet concentration had no effect on the

amount of P consumed; neither was there a ratio:concentration interaction (Table 2b).

In contrast, larvae consumed similar amounts of C across all diets, with neither

concentration nor dietary P:C ratio having an influence on the amount of C consumed (Table 2c).

A median of 0.22 ± 0.06 g of C was consumed by (surviving) larvae across all diet treatments

(Fig. 3).

Cocoon weight varied differently with macronutrient intake depending on the overall

concentration of the diet (Table 3a). At 90% nutrient density, cocoon weight was correlated

positively with the amount of carbohydrate consumed, and negatively with protein (Fig. 4a). For

our range of diets, the greatest weights were obtained by larvae that ate above approx. 0.3g C and

below 0.15g P. In contrast, at 70% nutrient density, cocoons were lower in weight than on the

90% diets, and were fairly uniform in weight irrespective of macronutrient intake (Fig 4b). Note

that the linear model fitted well and no non-linear effects were observed for the range of diets

used - that is, we did not identify an optimal amount of P or C that maximised cocoon weight.

The relationship between survival and nutrition similarly depended upon dietary

concentration. Survival of larvae fed diets at 90% concentration depended primarily upon

carbohydrate consumption (Fig 5a). Those larvae that consumed high amounts of carbohydrate

saw the lowest mortality irrespective of how much protein was consumed. Interestingly,

however, at lower levels of carbohydrate, mortality increased with decreasing amounts of protein

(Table 3b). Survival of larvae raised on the more dilute diets was greater, and was not

substantially affected by intake of P or C (Fig 5). Again, all linear models showed a good fit, so

we did not look for non-linear patterns in larval survival.

Choice phase

We found no evidence of larvae defending a common intake target at the choice phase.

Nevertheless, they converged upon a P:C ratio of approx. 1:1.8 (i.e. a common line, or

"nutritional rail"; Fig. 6). The amount of protein consumed by larvae was significantly affected

by diet combination: more protein was consumed by individuals offered diet combinations that

were overall more concentrated (Table 4a, Fig 6). Similarly, carbohydrate consumption was

significantly affected by diet combination (Table 4b). However, unlike with protein, this pattern

appeared to be driven by the diets at the extreme, with significantly more carbohydrate eaten

only by those in the most concentrated diet pair compared to the least concentrated pair (Fig 6);

other pairwise comparisons were not significant.

Despite the lack of a common intake target, larvae were not consuming diets at random.

For both carbohydrate and protein we saw differences in consumption from what would have

been expected for each larva based on random consumption, and this effect was dependent on the

specific set of diet choices (Fig 7a, Table 4c, d). When visualised as the amounts of protein and

carbohydrate consumed during each 48h treatment period, it is clear that larvae were achieving a

degree of homeostasis in carbohydrate consumption compared to what would be expected under

random consumption of each diet choice, whereas their consumption of protein aligned closely

with what would be expected under random consumption (Fig. 7b-d).

Discussion

Although we did not identify an optimal (fitness-maximising) intake for either

carbohydrate or protein, carbohydrate was positively associated with both body size and survival

in *Osmia bicornis* larvae (Figs. 4a, 5a). Accordingly, given a choice, larvae converged on a

relatively carbohydrate-biased protein:carbohydrate ratio of 1:1.8 (Fig. 6). Moreover, larvae

prioritised carbohydrate over protein intake, showing tighter control over carbohydrate

consumption than over protein consumption (Fig 7), and pupating after eating about 0.22 g

carbohydrate irrespective of protein and of dietary dilution (Fig 3). This carbohydrate target fell

short of the amount of carbohydrate that maximised cocoon weight or survival to pupation. Yet

achieving even this carbohydrate target carried costs for larvae on more dilute diets, which

suffered greater mortality and lower cocoon weights (Figs 4b, 5b). Taken together, these results

show that (1) larval *O. bicornis* are at least partially responsible for their own nutritional

regulation, and (2) their performance and consumption rules suggest adaptation to a pattern of

carbohydrate-limited growth and survival. In what follows, we suggest how and why these

patterns in *O. bicornis* depart from expected results based on studies of related organisms, and

more generally what these findings suggest about nutritional cooperation between parents and

offspring in provisioning species.

Larvae grew and survived best on our highest carbohydrate (i.e. lowest

protein:carbohydrate) diets. Accordingly, across our range of diets, larvae maintained a constant

carbohydrate intake while tolerating excesses or deficiencies of protein (a "no-interaction" rule

of compromise; Raubenheimer and Simpson 1999a) - although it is conceivable that this

represented a narrow window on a more curved intake array that would be visible over a broader

diet range. Both these patterns are unusual because insect herbivores are generally considered to

be limited by protein (e.g. Bernays and Chapman 2007; although see Le Gall and Behmer 2014).

In the few existing studies involving larval bees, e.g. honeybees (Helm et al. 2017), bumblebees

(Kraus et al. 2019) and sweat bees (Roulston and Cane 2002), protein and not carbohydrate

mediated larval growth and/or survival. More broadly, insect larvae often grow and survive best

on balanced or moderately high protein:carbohydrate ratios (Roeder and Behmer 2014;

Rodrigues et al. 2015), although low protein:carbohydrate ratios are associated with longevity in

adults (e.g. Lee et al. 2008). Moreover, animals generally prioritise regulation of the nutrient that

is typically limiting in their normal diet, and tolerate variation in nutrients that are abundant

(Raubenheimer and Simpson 1999b). Tolerance of wide variation in protein is thus usually seen

in predators (e.g. Raubenheimer et al. 2007; Kohl et al. 2015). In contrast, herbivores often

regulate protein more tightly than carbohydrate (Lee et al. 2002; Le Gall and Behmer 2014;

VanOverbeke et al. 2017).

During the choice phase, when allowed to self-select diets, *O. bicornis* larvae converged

on a protein:carbohydrate ratio of 1:1.8. This ratio is considerably more carbohydrate-biased than

that preferred by bumblebees foraging on behalf of microcolonies (1:0.25, Vaudo et al. 2018;

1:0.08, Kraus et al. 2019), and (to a lesser extent) than ants foraging for colonies with offspring

(1:1.5, Dussutour and Simpson 2009). It is also more carbohydrate-biased than that selected by

reproductive, solitary phytophages such as grasshopper adults and lepidopteran larvae (1:0.25 -

1:1.4, reviewed in Behmer 2009) and is closer to diets selected by *Drosophila* larvae (1:2,

Rodrigues et al. 2015). Notably, though, 1:1.8 was more protein-biased than the ratio that we

found maximised both cocoon weight and survival (1:3.4), suggesting *O. bicornis* larvae may

choose diets that favour other fitness-related quantities (such as reproduction and/or

developmental time) over body size/survival, as in *Drosophila* (Lee et al. 2008; Rodrigues et al.

2015).

Two main features of *O. bicornis'* ecology may help to explain their prioritization of

carbohydrate, and their relative preference for this macronutrient compared to what we know of

related taxa. First, the relative paucity of carbohydrate in *O. bicornis'* larval diet may help to

explain these findings. Despite being herbivorous, *Osmia* larvae are unlikely to be protein-

limited, because pollen is among the most protein-rich of plant tissues (Mattson 1980).

Moreover, in *Osmia* specifically, nectar constitutes only a tiny fraction of the pollen ball, less

than 4% (Maddocks and Paulus 1987; see Radmacher and Strohm 2010), in contrast to many

other bees where nectar is a principal source of carbohydrate for larvae (e.g. Kraus et al. 2019).

O. bicornis larvae may therefore be limited more by the amount of digestible carbohydrate

within pollen than by dietary protein (see Roulston and Cane 2000). Second, *O. bicornis* is (to

our knowledge) the first truly solitary hymenopteran studied under the GF; other studies have

concerned individuals likely to become workers of social species. Unlike social hymenopterans,

O. bicornis offspring are all reproductive and undergo diapause (Fliszkiewicz et al. 2012) - both

activities dependent on the fat body, where carbohydrate-derived fat is stored (Kawooya and

Law 1988; Ziegler and Van Antwerpen 2006; Hahn and Denlinger 2007; Wasielewski et al.

2013). Thus, *O. bicornis* larvae may have additional requirements for carbohydrate over and

above those of developing nonreproductive, nondiapausing hymenopteran workers. These

contrasting findings reinforce the idea that bees' nutritional needs may be just as diverse as their

ecologies.

Although larvae retained the ability to regulate carbohydrate by over- or under-eating

protein, they nevertheless coped very poorly with dietary dilution (Fig 4b, 5b), despite displaying

compensatory feeding behaviour (Fig 2) that suggests they both detected and responded to such

dilution. The dilution was not excessive (70% nutrient density) compared to similar studies

offering very highly dilute diets (14%, Raubenheimer and Simpson 1993; 16.8%, Lee et al.

2004). The locusts and caterpillars in those studies, though, are adapted for diets that vary greatly

in nutrient density, beginning dilute and becoming even more dilute over the season (Scriber and

Slansky 1981). By contrast, pollen is among the most consistently nutrient-rich parts of a plant

(Roulston and Cane 2000) and does not broadly vary in composition over a season (DeGrandi-

Hoffman et al. 2018). With a normal diet of unadulterated pollen and very little nectar, *Osmia*

larvae may have had no need to evolve mechanisms to cope with dilution. In comparison,

caterpillars reared on an invariant diet for generations lost the ability both to regulate intake and

to cope with dilution (Warbrick-Smith et al. 2009). *Osmia* larvae appear to have retained the

former capacity, but lost the latter, suggesting a normal diet that is dense in nutrients, but

variable in composition.

In systems where parents gather food for offspring from the environment, both parents

and offspring can be active participants in nutritional regulation. The lack of protein regulation

shown by *O. bicornis* larvae highlights the importance of understanding (a) whether mother *O.*

bicornis adjust protein content of provisions in response to imbalances in the landscape, and (b)

whether larvae have physiological adaptations (e.g. post-ingestive processing) for tolerating

protein imbalance. Budde & Lunau (2007) found that *O. bicornis* provisions contained about

19% protein regardless of pollen species used, suggesting a degree of homeostasis by parents.

Yet human activity is reducing floral diversity (Papanikolaou et al. 2017). Evidence is mixed

concerning whether, in practice, parent bees assess pollen nutrients at the flower (reviewed by

Nicholls and Hempel de Ibarra 2016). Both bumblebees and ants balance nutrition on behalf of

colonies (Dussutour and Simpson 2009; Vaudo et al. 2018), regulating more tightly when

foraging for offspring - protein in both taxa (Dussutour and Simpson 2009; Kraus et al. 2019)

and carbohydrate in ants (Dussutour and Simpson 2008; Cook et al. 2010). On the other hand,

protein gathered by honeybees varied passively with landscape usage (while maintaining

carbohydrate and lipid; Donkersley et al. 2014). Which regulatory strategy *Osmia* parents and

larvae collectively pursue may have important implications for their vulnerability to human-

induced landscape change, and so should now be a focus for research.

More broadly, understanding the relative roles of offspring (intake regulation and post-

ingestive processing) versus parents in nutrient balancing will be key to understanding the

nutritional ecology of species with parental provisioning. Such species include not just bees, but

other important ecosystem service providers such as dung beetles (Frank et al. 2017) and burying

beetles (Hopwood et al. 2013), as well as altricial birds (Wiens and Johnston 2012) and even

humans (Burt and Amin 2014). Alloregulation by parents is not a given; the relative roles of

parent and offspring in these groups are likely to reflect species' ecologies. Recent studies have

found nutritional mismatches between oviposition sites selected by parents and the nutritional

requirements of the offspring that will develop in those sites (Rodrigues et al. 2015; Lihoreau et

al. 2016). Parent sweat bees (*Lasioglossum zephyrum*) appear not to regulate protein in larval

provisions, despite protein mediating offspring performance (Roulston and Cane 2002). In *O.*

bicornis, we have shown that offspring retain the ability to regulate their nutritional intake

despite all food selection being done by parents. Larvae appeared to pay closest attention to

regulating dietary carbohydrate, consistent with this nutrient mediating both growth and survival.

Yet protein remains a key requirement for development; key now is to establish specifically how

protein balance is achieved, and whether parents or larvae carry that responsibility.

References

Behmer ST. 2009. Insect herbivore nutrient regulation. Annu Rev Entomol. 54:165–187.

Bernays EA, Chapman RF. 2007. Host-Plant Selection by Phytophagous Insects. Springer
Science & Business Media.

Budde J, Lunau K. 2007. Rezepte für ein Pollenbrot--heute: Osmia rufa. Entomologie heute.
19:173–179.

Burt NM, Amin M. 2014. A mini me?: exploring early childhood diet with stable isotope
ratio analysis using primary teeth dentin. Arch Oral Biol. 59(11):1226–1232.

Cane JH. 2016. Adult Pollen Diet Essential for Egg Maturation by a Solitary Osmia Bee. J
Insect Physiol. doi:10.1016/j.jinsphys.2016.09.011.
http://dx.doi.org/10.1016/j.jinsphys.2016.09.011.

Cook SC, Eubanks MD, Gold RE, Behmer ST. 2010. Colony-level macronutrient regulation in
ants: mechanisms, hoarding and associated costs. Anim Behav. 79(2):429–437.

Costa JT. 2006. The Other Insect Societies. Harvard University Press.

DeGrandi-Hoffman G, Gage SL, Corby-Harris V, Carroll M, Chambers M, Graham H, Watkins
deJong E, Hidalgo G, Calle S, Azzouz-Olden F, et al. 2018. Connecting the nutrient
composition of seasonal pollens with changing nutritional needs of honey bee (Apis
mellifera L.) colonies. J Insect Physiol. 109:114–124.

Despland E, Noseworthy M. 2006. How well do specialist feeders regulate nutrient intake?
Evidence from a gregarious tree-feeding caterpillar. J Exp Biol. 209(Pt 7):1301–1309.

Donkersley P, Rhodes G, Pickup RW, Jones KC, Wilson K. 2014. Honeybee nutrition is linked
to landscape composition. Ecol Evol. 4(21):4195–4206.

Dussutour A, Simpson SJ. 2008. Carbohydrate regulation in relation to colony growth in
ants. J Exp Biol. 211(Pt 14):2224–2232.

Dussutour A, Simpson SJ. 2009. Communal nutrition in ants. Curr Biol. 19(9):740–744.

Falk SJ. 2015. Field guide to the bees of Great Britain and Ireland. British Wildlife
Publishing.

Fichter BL, Stephen WP, Vandenberg JD. 1981. An Aseptic Technique for Rearing Larvae of
the Leafcutting Bee Megachile Rotundata (Hymenoptera, Megachilidae). J Apic Res.
20(3):184–188.

Field J. 2005. The evolution of progressive provisioning. Behav Ecol.
http://beheco.oxfordjournals.org/content/16/4/770.short.

Filipiak M. 2019 Apr 15. Key pollen host plants provide balanced diets for wild bee larvae:
A lesson for planting flower strips and hedgerows. Rader R, editor. J Appl Ecol.
doi:10.1111/1365-2664.13383. https://onlinelibrary.wiley.com/doi/abs/10.1111/1365-
2664.13383.

Fliszkiewicz M, Giejdasz K, Wasielewski O, Krishnan N. 2012. Influence of winter
temperature and simulated climate change on body mass and fat body depletion during
diapause in adults of the solitary bee, Osmia rufa (Hymenoptera: Megachilidae). Environ
Entomol. 41(6):1621–1630.

Frank K, Brückner A, Hilpert A, Heethoff M, Blüthgen N. 2017. Nutrient quality of vertebrate
dung as a diet for dung beetles. Sci Rep. 7(1):12141.

Goulson D, Nicholls E, Botías C, Rotheray EL. 2015. Bee declines driven by combined stress
from parasites, pesticides, and lack of flowers. Science. 347(6229):1255957.

Hahn DA, Denlinger DL. 2007. Meeting the energetic demands of insect diapause: nutrient
storage and utilization. J Insect Physiol. 53(8):760–773.

Haider M, Dorn S, Sedivy C, Muller A. 2014. Phylogeny and floral hosts of a predominantly
pollen generalist group of mason bees (Megachilidae: Osmiini). Biol J Linn Soc Lond.
111(1):78–91.

Harper EJ, Turner CL. 2000. Nutrition and energetics of the canary (Serinus canarius).
Comp Biochem Physiol B Biochem Mol Biol. 126(3):271–281.

Helm BR, Slater GP, Rajamohan A, Yocum GD, Greenlee KJ, Bowsher JH. 2017. The
geometric framework for nutrition reveals interactions between protein and carbohydrate
during larval growth in honey bees. Biol Open. 6(6):872–880.

Hopwood PE, Moore AJ, Royle NJ. 2013. Nutrition during sexual maturation affects
competitive ability but not reproductive productivity in burying beetles. Blanckenhorn W,
editor. Funct Ecol. 27(6):1350–1357.

Jauker F, Bondarenko B, Becker HC, Steffan-Dewenter I. 2012. Pollination efficiency of wild
bees and hoverflies provided to oilseed rape. Agric For Entomol. 14(1):81–87.

Kawooya JK, Law JH. 1988. Role of lipophorin in lipid transport to the insect egg. J Biol
Chem. 263(18):8748–8753.

Kohl KD, Coogan SCP, Raubenheimer D. 2015. Do wild carnivores forage for prey or for
nutrients? Bioessays. 37(6):701–709.

Kraus S, Gómez-Moracho T, Pasquaretta C, Latil G, Dussutour A, Lihoreau M. 2019.
Bumblebees adjust protein and lipid collection rules to the presence of brood. Curr Zool.
65(4):437–446.

Lee KP, Behmer ST, Simpson SJ, Raubenheimer D. 2002. A geometric analysis of nutrient
regulation in the generalist caterpillar Spodoptera littoralis (Boisduval). J Insect Physiol.
48(6):655–665.

Lee KP, Raubenheimer D, Simpson SJ. 2004. The effects of nutritional imbalance on
compensatory feeding for cellulose-mediated dietary dilution in a generalist caterpillar.
Physiol Entomol. 29(2):108–117.

Lee KP, Simpson SJ, Clissold FJ, Brooks R, Ballard JWO, Taylor PW, Soran N, Raubenheimer
D. 2008. Lifespan and reproduction in Drosophila: New insights from nutritional geometry.
Proc Natl Acad Sci U S A. 105(7):2498–2503.

Le Gall M, Behmer ST. 2014. Effects of protein and carbohydrate on an insect herbivore: the
vista from a fitness landscape. Integr Comp Biol. 54(5):942–954.

Lihoreau M, Buhl J, Charleston MA, Sword GA, Raubenheimer D, Simpson SJ. 2014.
Modelling nutrition across organizational levels: from individuals to superorganisms. J
Insect Physiol. 69:2–11.

Lihoreau M, Poissonnier L-A, Isabel G, Dussutour A. 2016. Drosophila females trade off
good nutrition with high-quality oviposition sites when choosing foods. J Exp Biol. 219(Pt
16):2514–2524.

Mackenzie G, Boa AN, Diego-Taboada A, Atkin SL, Sathyapalan T. 2015. Sporopollenin, The
Least Known Yet Toughest Natural Biopolymer. Front Mater Sci. 2:129.

Maddocks R, Paulus HF. 1987. Quantitative Aspekte der Brut-biologie von Osmia rufa L.
und Osmia cornuta Latr.(Hymenoptera, Megachilidae): Eine vergleichende Untersuchung
zu Mechanismen der Konkurrenzminderunt zweier nahverwandter Bienenarten.
Zoologische Jahrbücher Abteilung für Systematik, Ökologie und Geographie der Tiere.
114(1):15–44.

Mattson WJ. 1980. Herbivory in Relation to Plant Nitrogen Content. Annu Rev Ecol Syst.
11(1):119–161.

Michaelsen KF, Weaver L, Branca F, Robertson A. 2003. Feeding and nutrition of infants and
young children. WHO Regional Publications, European Series Report No.: 87.

http://www.euro.who.int/__data/assets/pdf_file/0004/98302/WS_115_2000FE.pdf?ua=1.

Nelson EV, Roberts RB, Stephen WP. 1972. Rearing Larvae of the Leaf-Cutter Bee Megachile
Rotundata on Artificial Diets. J Apic Res. 11(3):153–156.

Nicholls E, Hempel de Ibarra N. 2016. Assessment of pollen rewards by foraging bees. Funct
Ecol. doi:10.1111/1365-2435.12778. http://dx.doi.org/10.1111/1365-2435.12778.

Papanikolaou AD, Kühn I, Frenzel M, Kuhlmann M, Poschlod P, Potts SG, Roberts SPM,
Schweiger O. 2017. Wild bee and floral diversity co-vary in response to the direct and
indirect impacts of land use. Ecosphere. 8(11). doi:10.1002/ecs2.2008.
http://dx.doi.org/10.1002/ecs2.2008.

Poissonnier L-A, Arganda S, Simpson SJ, Dussutour A, Buhl J. 2018. Nutrition in extreme
food specialists: An illustration using termites. Overgaard J, editor. Funct Ecol.
32(11):2531–2541.

Poissonnier L-A, Arganda S, Simpson SJ, Dussutour A, Buhl J. 2018 Aug 19. Nutrition in
extreme food specialists: an illustration using termites. Funct Ecol. doi:10.1111/1365-
2435.13200. http://doi.wiley.com/10.1111/1365-2435.13200.

Radmacher S, Strohm E. 2010. Factors affecting offspring body size in the solitary bee
Osmia bicornis (Hymenoptera, Megachilidae). Apidologie . 41(2):169–177.

Raubenheimer D, Jones SA. 2006. Nutritional imbalance in an extreme generalist omnivore:
tolerance and recovery through complementary food selection. Anim Behav. 71(6):1253–
1262.

Raubenheimer, D., Mayntz, D., Simpson, S.J., Tøft, S., 2007. Nutrient-specific compensation
following diapause in a predator: implications for intraguild predation. Ecology 88, 2598–
2608.

Raubenheimer D, Simpson SJ. 1993. The geometry of compensatory feeding in the locust.
Anim Behav. 45(5):953–964.

Raubenheimer D, Simpson SJ. 1999a. Integrating nutrition: a geometrical approach.
Entomol Exp Appl. 91(1):67–82.

Raubenheimer D, Simpson SJ. 1999b. Integrating nutrition: a geometrical approach. In:
Proceedings of the 10th International Symposium on Insect-Plant Relationships. Springer
Netherlands. (Series Entomologica). p. 67–82.

Raubenheimer D, Simpson SJ, Mayntz D. 2009. Nutrition, ecology and nutritional ecology:
toward an integrated framework. Funct Ecol. 23(1):4–16.

Rodrigues MA, Martins NE, Balancé LF, Broom LN, Dias AJS, Fernandes ASD, Rodrigues F,
Sucena É, Mirth CK. 2015. Drosophila melanogaster larvae make nutritional choices that

minimize developmental time. J Insect Physiol. 81:69–80.

Roeder KA, Behmer ST. 2014. Lifetime consequences of food protein-carbohydrate content
for an insect herbivore. Davidowitz G, editor. Funct Ecol. 28(5):1135–1143.

Roulston TH, Cane JH. 2000. Pollen nutritional content and digestibility for animals. Plant
Syst Evol. 222(1-4):187–209.

Roulston TH, Cane JH. 2002. The effect of pollen protein concentration on body size in the
sweat bee Lasioglossum zephyrum (Hymenoptera: Apiformes). Evol Ecol. 16(1):49–65.

Roulston TH, Goodell K. 2011. The role of resources and risks in regulating wild bee
populations. Annu Rev Entomol. 56:293–312.

Royama T. 1970. Factors governing the hunting behaviour and selection of food by the
great tit (Parus major. J Anim Ecol. 39(3):619–668.

Schmickl T, Karsai I. 2017. Resilience of honeybee colonies via common stomach: A model
of self-regulation of foraging. PLoS One. 12(11):e0188004.

Scriber JM, Slansky F Jr. 1981. The Nutritional Ecology of Immature Insects. Annu Rev
Entomol. 26(1):183–211.

Simpson SJ, Raubenheimer D. 1993. A Multi-Level Analysis of Feeding Behaviour: The
Geometry of Nutritional Decisions. Philos Trans R Soc Lond B Biol Sci. 342(1302):381–402.

Simpson SJ, Raubenheimer D. 2012. The nature of nutrition: a unifying framework. Aust J
Zool. 59(6):350–368.

Strohm E, Daniels H, Warmers C, Stoll C. 2002. Nest provisioning and a possible cost of
reproduction in the megachilid bee Osmia rufa studied by a new observation method. Ethol
Ecol Evol. 14(3):255–268.

VanOverbeke DR, Thompson SN, Redak RA. 2017. Dietary self-selection and rules of
compromise by fifth-instar Vanessa cardui. Entomol Exp Appl. 163(2):209–219.

Vaudo AD, Farrell LM, Patch HM, Grozinger CM, Tooker JF. 2018. Consistent pollen
nutritional intake drives bumble bee (Bombus impatiens) colony growth and
reproduction across different habitats. Ecol Evol. 7:40568.

Vaudo AD, Patch HM, Mortensen DA, Tooker JF, Grozinger CM. 2016. Macronutrient ratios
in pollen shape bumble bee (Bombus impatiens) foraging strategies and floral preferences.
Proc Natl Acad Sci U S A. 113(28):E4035–42.

Warbrick-Smith J, Raubenheimer D, Simpson SJ, Behmer ST. 2009. Three hundred and fifty
generations of extreme food specialisation: testing predictions of nutritional ecology.
Entomol Exp Appl. 132(1):65–75.

Wasielewski O, Wojciechowicz T, Giejdasz K, Krishnan N. 2013. Overwintering strategies in
the red mason solitary bee—physiological correlates of midgut metabolic activity and
turnover of nutrient reserves in females of Osmia bicornis. Apidologie . 44(6):642–656.

Weeks RD, Wilson LT, Vinson SB, James WD. 2004. Flow of Carbohydrates, Lipids, and
Protein Among Colonies of Polygyne Red Imported Fire Ants, Solenopsis invicta
(Hymenoptera: Formicidae). Ann Entomol Soc Am. 97(1):105–110.

Wiens J, Johnston R. 2012. Adaptive correlates of granivory in birds. Granivorous Birds in
Ecosystems: Their Evolution, Populations, Energetics, Adaptations, Impact and Control,
(Eds J Pinowski and SC Kendeigh) pp.:301–340.

Ziegler R, Van Antwerpen R. 2006. Lipid uptake by insect oocytes. Insect Biochem Mol Biol.
36(4):264–272.

Tables & Figures

Table 1. Sample sizes for each diet combination used for choice phase (allocated by random

coin toss). "Order" refers to diet order - e.g. for A1C1, Order 1 would receive A1 first

whereas Order 2 would receive C1 first, determined by coin toss. Surviving larvae are in

parentheses.

	Order 1	Order 2
A1C1	1 (1)	6 (5)
A1C2	5 (5)	4 (3)
A2C1	3 (2)	5 (5)
A2C2	5 (2)	3 (1)

 Table 2. Model tables for models of consumption in no-choice phase. Key: ratio = dietary
protein:carbohydrate ratio (A = 1P:1.2C, B=1P:2.3C, C=1P:3.4C). Conc=overall nutrient
concentration.

(a) Total amount consumed (linear model)

Model term	Coefficients of minimal model		Effect of dropping term from full model			
	Estimate	Std. Error	df1	df2	F	p
Intercept (ratio A, conc 90%)	0.67	0.03				
ratio	**-**	**-**	**2**	**80**	**10.86**	**<0.0001**
ratio B	-0.13	0.05				
ratio C	-0.20	0.04				
conc	**-**	**-**	**1**	**79**	**14.04**	**<0.001**
conc 70%	0.15	0.04				
ratio × conc	-	-	2	78	0.4	0.67

(b) Protein consumed (linear model)

Model term	Estimate	Std. Error	df1	df2	F	p
Intercept (ratio A)	0.18	0.01				
ratio	**-**	**-**	**2**	**80**	**75.87**	**<0.0001**
ratio B	-0.07	0.01				
ratio C	-0.11	0.01				

conc	-	-	1	79	1.85	0.1
ratio × conc	-	-	2	78	0.0032	0.99

(c) Carbohydrate consumed (linear model)

Model term	Estimate	Std. Error	df1	df2	F	p
Intercept	0.23	0.01				
ratio	-	-	2	80	1.52	0.22
conc	-	-	1	79	2.97	0.09
ratio × conc	-	-	2	78	0.20	0.82

531
532
533

Table 3. Model tables for models of fitness correlates in no-choice phase

(a) Cocoon weight (linear model)

Model term	Coefficients of minimal model		Effect of dropping term from full model			
	Estimate	Std. Error	df1	df2	F	p
Intercept (conc 90%)	0.032	0.015	-	-	-	-
Protein	-0.317	0.065	-	-	-	-
Carbohydrate	0.408	0.066	-	-	-	-
Conc 70%	-0.011	0.027	-	-	-	-
Protein:Carbohydrate	-	-	1	72	0.145	0.705
Protein:Conc 70%	**0.392**	**0.126**	**1**	**72**	**8.942**	**0.004**
Carbohydrate:Conc 70%	**-0.337**	**0.132**	**1**	**72**	**6.310**	**0.014**

(b) Survival to pupation (parametric survival regression)

Model term	Coefficients of minimal model		Effect of dropping term from full model		
	Estimate	Std. Error	df1	χ^2	p
(Intercept)	2.114	0.556	-	-	-
Protein	6.009	2.257	-	-	-

Carbohydrate	9.236	3.188	-	-	-	-
Conc 70%	0.467	0.543	-	-	-	-
Protein:Carbohydrate	**-18.505**	**8.426**	**1**		**-3.977**	**0.046**
Protein:Conc 70%	**-**	**-**	1		-0.072	0.789
Carbohydrate:Conc 70%	**-6.279**	**3.138**	**1**		**-4.875**	**0.027**

535
536
537
538

Table 4. Model tables for models of consumption in choice phase. For details of diet combinations, see text.

(a) Protein consumed (amount)

| Model term | Coefficients of minimal model | | Effect of dropping term from full model | | | |
	Estimate	Std. Error	df1	df2	F	p
Diet combination	-	-	**3**	**23**	**7.43**	**0.002**
A1C2	0.19	0.01				
A2C1	0.15	0.01				
A1C1	0.21	0.01				
A2C2	0.12	0.02				

(b) Carbohydrate consumed (amount)

| Model term | Coefficients of minimal model | | Effect of dropping term from full model | | | |
	Estimate	Std. Error	df1	df2	F	p
Diet combination	-	-	**3**	**23**	**4.58**	**0.013**
A1C2	0.32	0.02				
A2C1	0.28	0.02				
A1C1	0.37	0.02				

A2C2	0.23	0.03

(c) Protein consumed (departure from random)

Model term	Coefficients of minimal model		Effect of dropping term from full model			
	Estimate	Std. Error	df1	df2	F	p
Diet combination	**-**	**-**	**4**	**20**	**19.67**	**<0.0001**
A1C2	0.007	0.002				
A2C1	0.006	0.002				
C1A1	0.013	0.002				
C2A2	-0.001	0.003				

(d) Carbohydrate consumed (departure from random)

Model term	Coefficients of minimal model		Effect of dropping term from full model			
	Estimate	Std. Error	df1	df2	F	p
Diet combination	**-**	**-**	**4**	**20**	**51.647**	**<0.0001**
A1C2	-0.001	0.003				
A2C1	-0.035	0.003				
C1A1	-0.020	0.003				

	C2A2	0.002	0.004

Figure captions

Figure 1. Expected protein and carbohydrate consumption if larvae ate indiscriminately between

two diets. Diet choices are pairwise combinations of diets A, B, C and D, which each contain

protein and carbohydrate at different ratios and concentrations. Solid lines represent P:C ratios;

black points represent actual nutrient content of each diet, which depends upon dilution as well

as P:C ratio. Red points represent expected consumption if larvae eat randomly (i.e. equally)

from each of a choice of two diets (choices denoted by the red point labels).

Figure 2. Amount of provision in grams consumed by larvae raised on the 3 different P:C ratio

artificial diets at the 2 different macronutrient concentrations (90% and 70% macronutrient

content).

Figure 3. Mean total (+/- 1 SD) amount of P and C consumed in grams by larvae on each

diet before pupation. Solid lines and letters represent three P:C ratios (A = 1:1.2, B = 1:2.3,

C = 1:3.4). Numbers following letters denote diet concentration (1 = 90%, 2 = 70%).

Figure 4. Effects of P and C consumption upon cocoon weight (g) in larvae fed diets at (a)

90% and (b) 70% nutrient density. Transition from blue to red indicates heavier cocoons.

For context, mean total consumption of P and C for each diet is plotted (data as in Fig. 2).

Solid lines and letters represent three P:C ratios (A = 1:1.2, B = 1:2.3, C = 1:3.4).

Figure 5. Effects of P and C consumption upon survival probability (black = died, white =

pupated) and duration (colour; transition from blue to red indicates longer survival

duration) in larvae fed diets at (a) 90% and (b) 70% nutrient density. For context, mean

total consumption of P and C for each diet is plotted (data as in Fig. 2). Solid lines and

letters represent three P:C ratios (A = 1:1.2, B = 1:2.3, C = 1:3.4).

Figure 6. The mean (+/- SE) amount of protein (P) and carbohydrate (C) eaten by larvae in

the choice experiment. Each point label denotes a choice of two diets, one A and one C;

black labels show observed intake, red labels show expected intake under random

consumption. Letters represent diet P:C ratio (A = 1:1.2, C = 1:3.4); numbers represent diet

concentration (1 = 90%, 2 = 70%). Solid lines represent dietary P:C ratios (Top line = Diet

A, Bottom line = Diet C). Dashed red line shows expected average P:C ratio based on

random consumption. Dashed black line shows average P:C ratio of observed intake across

larvae. Black bars show comparisons *among* treatment groups' observed intake (Tukey's

post-hoc comparisons; for details of linear model see text) in P and C.

Figure 7. (a) Deviation from random intake of protein and carbohydrate for larvae in

different treatment groups during the choice phase. Bars with similar letters displayed

above or below are not statistically significantly different (Tukey's post-hoc comparisons).

(b, c) Mean (+/- 1SD) observed intake of protein and carbohydrate over successive diet

swaps, irrespective of the concentrations of the diet choices, for larvae starting on diet A

(b) or diet C (c), versus expected nutrient intake under random consumption of diets (d,

showing larvae starting on diet C, chosen arbitrarily). For calculations of expected

consumption, see text.

589
590   Fig. 1.

592   Fig. 2.

594
595   Fig. 3.

597   Fig. 4.

600
601   Fig. 5.
602

603
604 Fig. 6.
605

609   Fig. 7.