Widespread methane formation by *Cyanobacteria* in aquatic and terrestrial ecosystems

Bižić-Ionescu M.[1*#], Klintzsch T.[2*], Ionescu D.[1*], Hindiyeh M. Y.[3], Günthel M.[4], Muro-Pastor A.M.[5], Eckert W.[6], Keppler F.[2,7], Grossart H-P[1,8#]

1. Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), Alte Fischerhuette 2, D-16775 Stechlin, Germany
2. Institute of Earth Sciences, Biogeochemistry Group, Heidelberg University, Heidelberg, Germany
3. Department of Water and Environmental Engineering, German Jordanian University, Amman, Jordan
4. Department of Biosciences, Swansea University, SA2 8PP Swansea, United Kingdom.
5. Instituto de Bioquímica Vegetal y Fotosíntesis, Consejo Superior de Investigaciones Científicas and Universidad de Sevilla, Sevilla, Spain
6. Israel Oceanographic and Limnological Research, The Yigal Alon Kinneret Limnological Laboratory, Migdal 14650, Israel
7. Heidelberg Center for the Environment (HCE), Heidelberg University, D-69120 Heidelberg, Germany
8. Institute of Biochemistry and Biology, Potsdam University, Maulbeerallee 2, 14469 Potsdam, Germany

* These authors contributed equally to this work

\# Correspondence should be addressed to:

Mina Bižić-Ionescu, Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), Department of Experimental Limnology, Alte Fischerhuette 2, Neuglobsow, OT Stechlin, 16775, Germany
Email: mbizic@igb-berlin.de; Tel +49-33082-69969

Hans-Peter Grossart, Leibniz Institute of freshwater ecology and inland fisheries (IGB), Department of Experimental Limnology, Alte Fischerhuette 2, Neuglobsow, OT Stechlin, 16775, Germany
Email: hgrossart@igb-berlin.de; Tel +49-33082-69991

The authors declare no conflict of interest

Abstract

Evidence is accumulating to challenge the paradigm that biogenic methanogenesis, traditionally considered a strictly anerobic process, is exclusive to *Archaea*. Here we demonstrate that Cyanobacteria living in marine, freshwater and terrestrial environments produce methane at substantial rates under light and dark oxic and anoxic conditions, forming a link between light driven primary productivity and methane production in globally relevant group of phototrophs. Biogenic methane production was enhanced during oxygenic photosynthesis and directly attributed to the cyanobacteria by applying stable isotope labelling techniques. We suggest that formation of methane by *Cyanobacteria* may contribute to methane accumulation in oxygen-saturated surface waters of marine and freshwater ecosystems. Moreover, in these environments, cyanobacterial blooms already do, and might further occur more frequently during future global warming and thus have a direct feedback on climate change. We further highlight that cyanobacterial methane production not only affects recent and future global methane budgets, but also has implications for inferences on Earth's methane budget for the last 3.5 billion years, when this phylum is thought to have first evolved.

Introduction

Methane (CH_4) is the second most important anthropogenic greenhouse gas after CO_2 and is estimated to have 28-34 times higher warming effect than the latter over a 100-year period (Intergovernmental Panel on Climate Change, 2014). The mixing ratio of CH_4 in the troposphere has increased from 715 ppbv in the preindustrial era to currently 1,860 ppbv (Nov. 2017 NOAA). Estimated global CH_4 emissions to the atmosphere average at ca. 560 Tg per year (1Tg = 10^{12} g) exceeding the current estimated sinks by ca. 13 Tg per year (Tian et al., 2016). Thus, to mitigate

the constant increase in atmospheric CH_4 a comprehensive understanding of global CH_4 sources

and the environmental parameters that affect them is necessary.

Traditionally, biogenic methanogenesis is the formation of methane under strictly anoxic

conditions by microbes from the domain *Archaea* (phylogenetically distinct from both eukaryotes

and *Bacteria*). However, in the past decade there has been growing evidence that also eukaryotes

such as algae (Lenhart et al., 2016), plants (Keppler, Hamilton, Braß, & Röckmann, 2006), animals

(Tuboly et al., 2013), fungi (Lenhart et al., 2012) and probably humans (Keppler et al., 2016)

produce methane under oxic conditions albeit at considerably lower rates. These recent findings

suggest that the phenomenon may not be solely limited to methanogenic *Archaea* and could

include new metabolic pathways. For example, the conversion of methylated substrates such as

methylphosphonates to CH_4 by *Bacteria* has been extensively addressed in recent years with

regards to the "methane paradox" (Repeta et al., 2016; Wang, Dore, & McDermott, 2017).

Recently, Zheng *et al.* (2018) have shown CH_4 formation by *Rhodopseudomonas palustris* during

N_2 fixation. Methane emission was also detected from cryptogamic covers, i.e. phototrophic

assemblages on plant, rock and soil surfaces (Lenhart et al., 2015).

Accumulation of CH_4 in oxygenated freshwater environments has been repeatedly associated with

the presence of *Cyanobacteria* (see also Fig. S1). Methane production by *Cyanobacteria* has been

attributed to either demethylation of methylphosphonates (Beversdorf, White, Björkman, Letelier,

& Karl, 2010; Gomez-Garcia, Davison, Blain-Hartnung, Grossman, & Bhaya, 2011; Yao, Henny,

& Maresca, 2016) or to the association of filamentous *Cyanobacteria* with methanogenic *Archaea*,

providing the latter with the necessary hydrogen for CH_4 production (Berg, Lindblad, & Svensson,

2014). *Cyanobacteria* are ubiquitous, found literally in any illuminated environment as well as

unexpectedly in some dark subsurface ones as well (Hubalek et al., 2016; Puente-Sánchez et al.,

2018). Furthermore, this phylum predominated Earth whilst the environment was still reductive and ca. 1.3 billion years prior to the great oxidation event which occurred 2.4 billion years ago (Gumsley et al., 2017). Therefore, we tested whether this phylum contributes to the global CH_4 budget independent of naturally occurring, extra-cellular precursor substances or the presence of methanogenic *Archaea*. If so, their ubiquitous nature, their expected future increase in abundance (Huisman et al., 2018; Visser et al., 2016) and their proximity to the interface with the atmosphere makes them potential key players in the global CH_4 cycle.

Here we demonstrate that unicellular as well as filamentous, freshwater, marine and terrestrial members of the prominent and ubiquitous phylum *Cyanobacteria*, a member of the domain *Bacteria*, produce CH_4 at substantial rates under both light and dark, oxic and anoxic conditions.

Material and Methods:

Cyanobacterial cultures

Seventeen different cyanobacterial cultures were obtained from various sources and grown using the media described in Table 1.

Strain name	Source	Morphology	Growth medium	Incubation Temp. °C
Anabaena sp. PCC7120	IBVF	Filamentous	BG11	30
Anabaena cylindrica ATCC29414	IBVF	Filamentous	BG11	30
Anabaena borealis	CCALA	Filamentous	BG11	30
Scytonema hofmanni PCC7110	IBVF	Filamentous	BG11	30
Leptolyngbya sp. (desert crust)	HUJI	Filamentous	BG11	30
Phormidium persicinum	IBVF	Filamentous	f/2	26
Trichodesmium erythraeum	MPI-MM	Filamentous	YBCII	26
Nodularia spumigena	IOW	Filamentous	f/2 (8 psu)	20
Chroococcidiopsis sp. PCC9317	IBVF	Unicellular	BG11	30
Microcystis aeruginosa PCC7806	IGB	Unicellular	BG11	30
Prochlorococcus sp. MIT9313	Uni Freiburg	Unicellular	AMP1	22
Prochlorococcus sp. MIT9312	Haifa Uni	Unicellular	ASW-Pro99	22

Prochlorococcus sp. MIT0604	Haifa Uni	Unicellular	FSW-Pro99	22
Prochlorococcus sp. NATL2A	Haifa Uni	Unicellular	ASW-Pro99	22
Prochlorococcus sp. MED4	Haifa Uni	Unicellular	ASW-Pro99	22
Synechococcus sp. WH7803	Haifa Uni	Unicellular	ASW-Pro99	22
Synechococcus sp. WH8102	Haifa Uni	Unicellular	ASW-Pro99	22

Table 1. Cyanobacterial cultures used in this study and their growth conditions. Shaded cultures are fully axenic while others are mono-cyanobacterial. Sources abbreviations: IBVF: Culture collection of the Institute for Plant Biochemistry and Photosynthesis, Sevilla Spain; CCALA: Culture collection of autotrophic organisms; HUJI: Laboratory of Aaron Kaplan, Hebrew University of Jerusalem, Jerusalem Israel; IOW: Laboratory of Falk Pollehne, Leibniz Institute for Baltic Sea research, Warnemünde, Germany; MPI-MM: Max Planck Institute for Marine Microbiology, Bremen, Germany; IGB: Leibniz Institute of Freshwater Ecology and Inland Fisheries, Neuglobsow, Germany; Uni. Freiburg, Laboratory of Claudia Steglich, Freiburg University, Freiburg, Germany. Haifa University, Laboratory of Daniel Sher. Media source: BG11 (Stanier, Deruelles, Rippka, Herdman, & Waterbury, 1979); f/2 (Guillard & Ryhter, 1962); YBCII (Chen, Zehr, & Mellon, 1996); AMP1 (Moore et al., 2007); Filtered sea water (FSW) / Artificial sea water Pro99 (Moore et al., 2007).

Stable isotope labeling experiments

Culturing and treatments

To investigate the production of *Cyanobacteria*-derived CH_4, 60 ml vials with 40 ml liquid and 20 ml head space volume (ambient air) were used and sealed with septa suitable for gas sampling. For the ^{13}C labelling experiments $NaH^{13}CO_3$ (99 % purity, Sigma-Aldrich, Germany) was added amounting to 10 % of the initial dissolved inorganic carbon (DIC) in BG11 (Stanier et al., 1979) (DIC = 0.4 mM, enriched by added $NaHCO_3$; pH ≈ 7.0) and 4.5 % of the DIC in f/2 medium (Guillard & Ryhter, 1962) (DIC = 2015 μmol L^{-1}; pH ≈ 8.2) and 1 % of the DIC in the Pro99 (Moore et al., 2007) based medium used for axenic *Synechococcus* and *Prochlorococcus* cultures. Four different examination groups were used: (1) Sterile medium; (2) Sterile medium with $NaH^{13}CO_3$; (3) Sterile medium with culture; (4) Sterile medium with culture and $NaH^{13}CO_3$; Four replicates of each cyanobacteria culture (n = 4).

The cultures were grown under a light–dark cycle of 16 and 8 hours at 22.5 °C at a light intensity

of $\approx$ 30 μmol quanta m^{-2} s^{-1} for a total period of 3 days.

Continuous-flow isotope ratio mass spectrometry (CF-IRMS)

CF-IRMS was employed for measurement of the δ^{13}C-CH_4 values in the head space gas above the

cultures. Head space gas from exetainers was transferred to an evacuated sample loop (40 mL) and

interfering compounds were then separated by GC and CH_4 trapped on Hayesep D. The sample

was then transferred to the IRMS system (ThermoFinniganDeltaplus XL, Thermo Finnigan,

Bremen, Germany) via an open split. The working reference gas was carbon dioxide of high purity

(carbon dioxide 4.5, Messer Griesheim, Frankfurt, Germany) with a known δ^{13}C value of -23.64

‰ relative to Vienna Pee Dee Belemnite (V-PDB). All δ^{13}C-CH_4 values were corrected using three

CH_4 working standards (isometric instruments, Victoria, Canada) calibrated against IAEA and

NIST reference substances. The calibrated δ^{13}C-CH_4 values of the three working standards were -

23.9 $\pm$ 0.2 ‰, -38.3 $\pm$ 0.2 ‰, and -54.5 $\pm$ 0.2 ‰. The average standard deviations (n = 3) of the

CF-IRMS measurements were in the range of 0.1 to 0.3 ‰. All ^{13}C / ^{12}C-isotope ratios are

expressed in the conventional δ notation in per mille (‰) vs. V-PDB, using the equation 1.

$$\delta = \left[\frac{(^{13}C/^{12}C)_{sample}}{(^{13}C/^{12}C)_{standard}} \right] - 1 \qquad\qquad (1)$$

Membrane inlet mass spectrometer experiments

Experiments were conducted using a Bay Instruments (MD, USA) Membrane Inlet Mass

Spectrometer (MIMS) consisting of a Pfeiffer Vacuum HiCube 80 Eco turbo pumping station

connected to a QMG 220 M1, PrismaPlus®, C-SEM, 1-100 amu, Crossbeam ion source mass

spectrometer (Pfeiffer Vacuum, Germany). Culture samples were pumped (Minipuls3, peristaltic

pump, Gilson) through a capillary stainless tubing connected to Viton® pump tubing as described in Kana *et al*. (Kana, Cornwell, & Zhong, 2006). The coiled stainless-steel tubing was immersed in a water bath to stabilize the sample temperature. Temperatures were set according to the growth conditions of the different cultures. Inside the vacuum inlet the sample passed through an 8 mm long semipermeable microbore silicone membrane (Silastic®, DuPont) before exiting the vacuum and returning to the culture chamber forming a closed system with respect to liquids. This required a 3.5 ml experimental chamber which consisted of an inner chamber where cultures were placed, and an isolated outer chamber connected to a water bath to maintain the culture at a constant temperature. The experimental chamber was placed on a magnetic stirrer and the cultures were continuously stirred for the duration of the experiments to prevent the formation of concentration gradients.

Cultures were transferred to fresh medium before the onset of each experiment after which 3.5 ml of the culture were transferred to the experimental chamber and an equal volume was used for determination of cell counts or dry weight. The latter was determined by filtering the samples on pre-weighed combusted GFF filters (Millipore) and drying at 105 °C for 48 h. In the case of non-homogenous cultures, the biomass from the experimental chamber was used at the end of the experiment for determination of dry weight. Marine picophytoplankton cultures were normalized by cell counting using a FACSAria II flow cytometer (BD Bioscienses, Heidelberg, Germany) at a flow rate of 23.5 µl / min for 2.5 min. Autofluorescence was used to separate cells from salt precipitates in the medium. Cells for counting were collected from the same batch used in the experimental chamber.

The light regime for the experiments was as follows: dark from 19:30 to 09:00 then light intensity was programmed to increase to 60, 120, 180, 400 µmol quanta m^{-2} s^{-1} with a hold time of 1.5 h at

each intensity. After maximum light the intensity was programmed to decrease in reverse order with the same hold times until complete darkness again at 19:30. Experiments lasted a minimum of 48 h with at least one replicate longer than 72 h. A minimum of 3 replicate experiments were conducted for each culture.

As negative controls ultrapure water as well as autoclave cyanobacterial biomass were measured to test for non-biogenic methane prosecution by the experimental system (Fig. S2).

Methane calculations

Methane (and oxygen) concentrations were calculated using the ratio relative to the inert gas Ar (m/z 40). Methane concentration was deduced from mass 15 which does not overlap with other gases in the sample (Schlüter & Gentz, 2008). The CH_4, O_2 and Ar concentration in the different media were calculated based on known solubility constants (Powell, 1972) and were calibrated to the measured signals in MQ water and growth media at different temperatures. To further calibrate the $CH_{4(mz15)}$/Ar ratio, measurements were conducted on air saturated water at different salinities (Fig. S3).

Methane production rates were calculated as the 1[st] derivative of the Savizky-Golay (Savitzky & Golay, 1964) smoothed data using the sgolay function in the R package signal (http://r-forge.r-project.org/projects/signal/). To account for the continuous degassing from the CH_4 supersaturation experimental chamber, the degassing rate was determined experimentally using rapid heating of air-saturated water from 18 to 30 °C leading to an instant (super)saturation of 127 % and 130 % for CH_4 and Ar, respectively. This procedure was repeated under two mixing conditions: I) mixing was applied via magnetic steering as conducted for most cultures; II) mixing occurred only via the cyclic pumping of sample through the MIMS membrane as applied to

Synechococcus and *Prochlorococcus* cultures. The change in concentration of CH_4 was monitored

and a linear ($R^2 = 0.95$) saturation degree dependent rate was determined. The determined rate

given in Equations 2 and 3 for type I and type II mixing, respectively, was similar to that

determined by comparing the most negative slopes of the culture experiments, when

cyanobacterial production rates are expected to be minimal or zero, and the supersaturation state

of the culture. Final rates were calculated by adding the absolute values of the local CH_4 slope (1^{st}

derivative) and the local degassing rate (equations. 2,3).

$$R_{degassing} = -2.2365 \times 10^{-12} \times Sat_{(CH_4)} + 2.12656 \times 10^{-12} \qquad (2)$$

$$R_{degassing} = -8.8628 \times 10^{-14} \times Sat_{(CH_4)} + 3.5819 \times 10^{-14} \qquad (3)$$

Where: $R_{degassing}$ is the degassing rate in mol CH_4 sec^{-1} and $Sat_{(CH4)}$ is the fraction CH_4

saturation state >1 (and <1.3) determined by measured concentration vs. calculated

solubility.

DNA extraction and sequencing

To evaluate the presence of methanogenic *Archaea* in non-axenic cultures DNA was extracted as

described in Nercessian *et al.* (Nercessian, Noyes, Kalyuzhnaya, Lidstrom, & Chistoserdova,

2005). The resulting DNA was sent for Illumina sequencing at MrDNA (Shallowater, TX, USA)

on a Miseq platform 2x300 bp using the Arch2A519F (CAGCMGCCGCGGTAA) and

Arch1017R (GGCCATGCACCWCCTCTC) primers (Fischer, Güllert, Neulinger, Streit, &

Schmitz, 2016). Archaeal community composition was analyzed using the SILVA-NGS pipeline

(Ionescu et al., 2012) (Fig. S4). After a standard PCR for the *mcrA* gene resulted in no visible

products from any of the cultures a qPCR assay was conducted as well resulting in low copy

numbers of the gene (Fig. S4). The sequences were submitted to the European Nucleotide Archive

under project number: PRJEB25851.

Results and Discussion

To test the hypothesis that *Cyanobacteria* directly produce CH_4 independent of methylated

precursors (e.g. methylphosphonates) in ambient water, thirteen different filamentous and

unicellular cyanobacterial cultures (for details of chosen cultures see Table 1) that are known to

grow in marine, freshwater and terrestrial environments were incubated under sterile conditions

with ^{13}C labelled sodium hydrogen carbonate ($NaH^{13}CO_3$) as carbon source. All investigated

cyanobacterial cultures showed CH_4 production with increasing stable isotope values ($\delta^{13}C$-CH_4

values) clearly indicating that ^{13}C carbon was implemented into CH_4, whereas no ^{13}C enrichment

occurred in the control experiments (Fig. 1). These results unambiguously show that

Cyanobacteria produce CH_4 *per se* and that the process is most likely linked to general cell

metabolism such as photoautotrophic carbon fixation. The different enrichment of ^{13}C indicated

by $\delta^{13}C$-CH_4 values ranging from 1.71 to 1337 ‰ observed in the different cultures is a result of

different production rates as well as differences in biomass. The involvement of methanogenic

Archaea in this process can be ruled out. First, five of the cultures were axenic. Second, the oxygen

concentrations during CH_4 production were in most cases above saturation level (Fig. 2 and Fig.

S2) and while methanogenic *Archaea* were recently reported from oxic environments (Angle et

al., 2017), their activity is attributed to anoxic microniches. Third, sequencing analysis of non-

axenic cultures and quantitative real-time PCR of the *mcrA* gene showed methanogenic *Archaea*

are either absent or present in negligible numbers (Fig. S4).

Furthermore, demethylation of methylphosphonates from the spent growth medium is unlikely to

be the mechanism involved in this instance even though some *Cyanobacteria* do possess the

necessary enzymatic machinery (Beversdorf et al., 2010; Gomez-Garcia et al., 2011) for the
following reasons. First, thus far, demethylation of methylphosphonates has been shown to occur
only under phosphorus starvation, which was highly unlikely in this study since the culture
medium contained ca. 200 µmol P L^{-1}. Indeed, publicly available transcriptomic data for *Anabaena*
sp. PCC 7120 (Flaherty, Van Nieuwerburgh, Head, & Golden, 2011; Mitschke, Vioque, Haas,
Hess, & Muro-Pastor, 2011) and *Trichodesmium erythraeum* (Pfreundt, Kopf, Belkin, Berman-
Frank, & Hess, 2014) show no evidence for activity of the phosphonate C-P lyase genes under
standard (P-rich) culture conditions. Secondly, some of the *Cyanobacteria* used in this study (i.e.
Microcystis aeruginosa PCC 7806, *Synechococcus* WH7803 and WH8102, as well as all
sequenced species of *Chroococcidiopsis* sp., *Leptolyngbya* sp. and *Phormidium* sp. and
Prochlorococcus) do not possess the known C-P lyase genes necessary for conversion of
methylphosphonates to CH$_4$. The lack of the *phn* genes (gene operon for phosphonate metabolism)
was demonstrated to be a common feature of the genus *Prochlorococcus* (Luo & Konstantinidis,
2011). *T. erythraeum* was shown to internally produce phosphonates as P storage later to be freed
by demethylation (Dyhrman, Benitez-Nelson, Orchard, Haley, & Pellechia, 2009), a process that
is likely to release CH$_4$. Nevertheless, the same study shows, though not focusing on cyanobacteria
alone, that marine unicellular organisms such as *Synechococcus* and *Crocosphaera*, do not contain
a detectable phosphonate storage.

Despite the recent finding of CH$_4$ production during N$_2$ fixation by *Rhodopseudomonas palustris*
(Zheng et al., 2018) we suggest that this is not the pathway leading to CH$_4$ production in our
experiments. First, most *Cyanobacteria* used in this study are unable (i.e marine *Synechococcus*,
Prochlorococcus, *Microcystis aeruginosa*) or unknown (*Leptolyngbya* sp., *Phormidium*
persicinum) to fix N$_2$. Second, all experiments were conducted in NO$_3^-$ or NH$_4^+$ rich, fresh, media,

and therefore N_2 fixation in capable cyanobacteria is likely to be inhibited to a certain degree

(Knapp, 2012). Thus, given the rapid and tight response of CH_4 production with the onset of light,

we consider that the mechanism by which *Cyanobacteria* readily convert fixed CO_2 to CH_4 under

light conditions must revolve around their central metabolism. Inhibitors of photosynthesis such

as Atrazine and DBMIB (2,5-Dibromo-6-isopropyl-3-methyl-1,4-benzoquinone) inhibit the

methane production under light conditions, however, the exact biochemical pathway(s) involved

in cyanobacteria-derived CH_4 formation remain so far unknown and thus require further

investigation.

Patterns and rates of CH_4 production were investigated in seventeen cultures over several days of

continuous measurement of CH_4 concentration using a membrane inlet mass spectrometry system

(MIMS). Our measurements, lasting 2-5 days, showed that CH_4 production occurs both under light

and dark conditions (Fig. 2 and Fig. S2). This is evident by a positive production rate at almost all

times in all experiments. Replicate experiments revealed that, while *Cyanobacteria* repeatedly

produced CH_4, rates and patterns were not consistent, particularly so for production during the

periods of darkness. Often, a period with lower rates of CH_4 production was observed between

light and dark phases (Fig. 2 and Fig. S2). The latter is evidenced as a decrease in CH_4

concentration resulting from degassing of our incubation system. This suggests that different

mechanisms may be involved in CH_4 production under light and dark conditions, presumably

dependent on freshly generated photosynthetic products during light and on storage compounds

during dark periods. Fermentation of storage compounds by *Cyanobacteria* has been previously

described and known to produce among other compounds acetate and hydrogen which are known

precursors of acetoclastic CH_4 formation (Stal & Moezelaar, 1997). Interestingly, most of the

genes required for methanogenesis are present in non-methanogenic organisms, including

Cyanobacteria. Nevertheless, in this instance since the methyl-coenzyme reductase (*mcr*) gene is absent this would suggest that if *Cyanobacteria* produce CH₄ via conventional pathways, an ortholog of the *mcr* gene exists.

Figure 1. δ^{13}C-CH₄ values measured during incubation experiments of fifteen different filamentous and unicellular freshwater, soil and marine cyanobacterial cultures with and without NaH^{13}CO₃ supplementation. All cyanobacterial cultures produced CH₄. Using NaH^{13}CO₃ as carbon source (CL) resulted in increasing stable δ^{13}C-CH₄ values as compared to the starting condition. This establishes the direct link between carbon fixation and CH₄ production. The ^{13}C enrichment is not quantitative and thus not comparable between cultures. Error bars represent standard deviation (n=4).

Methane production rates (Fig. 3) were calculated using the slope of CH₄ profiles and were normalized on a cyanobacterial biomass dry weight basis for larger cyanobacteria or cell counts for small-celled marine picophytoplankton. The latter to obtain high accuracy for the small-cell-sized picophytoplankton, *Synechococcus* and *Prochlorococcus*. Hourly CH₄ production rates

across cultures of larger cyanobacteria were in the range of 0.1 to 3.4 µmol g^{-1} h^{-1} in individual

experiments and a mean of 0.51 ± 0.26 µmol g^{-1} h^{-1}. Among the marine picophytoplankton

Synechococcus sp. exhibited low rates ranging between 10^{-4} and 10^{-2} µmol CH_4 per 10^6 cells, while

Prochlorococcus cultures produced methane at rates ranging from 0.01 to 5 µmol CH_4 per 10^6

cells. When compared to production rates of typical methanogenic *Archaea*, CH_4 production rates

of freshwater, soil and large marine cyanobacteria are three to four orders of magnitude lower than

the CH_4 production rates noted for typical methanogenic *Archaea* in culture under optimal

conditions (oxygen free) but one to three orders of magnitude higher than rates observed in

eukaryotes (Fig. S5). Due to their small size, conversion of CH_4 production rates of

picophytoplankton to µmol g^{-1} h^{-1} results in values exceeding those of methanogenic *Archaea*.

Nevertheless, to obtain 1 g of *Prochlorococcus* cells one would need to integrate 0.1-10 m^2 over

a depth of 200 m (Lange et al., 2018) as compared to ca. 20 Kg of soil for methanogenic *Archaea*

(assuming 10^9 cells per g sediment of which 50 % are methanogens). In our experiments,

Prochlorococcus and *Synechococcus* cultures produced CH_4 only at light intensities above 20

µmol quanta m^{-2} s^{-1} and therefore, it is likely that only surface *Procholorococcus* and

Syenchococcus communities contribute to the oceanic CH_4 flux to the atmosphere.

Figure 2. Continuous measurement of CH_4 and oxygen under light/dark periods using a membrane inlet mass spectrometer (MIMS). Examples are shown for two cultures. Data for other cultures can be found in Fig. S2. A decrease in CH_4 concentration is a result of either reduced, or no, production coupled with degassing from the supersaturated, continuously-mixing, semi-open incubation chamber towards equilibrium with atmospheric CH_4 (2.5 nM and 2.1 nM for freshwater and seawater, respectively). Calculated CH_4 production rates account for the continuous emission of CH_4 from the incubation chamber for as long as the CH_4 concentrations are supersaturated. The light regime for the experiments was as follows: dark (black bar) from 19:30 to 09:00 then light intensity (yellow bar) was programmed to increase to 60, 120, 180, 400 µmol quanta m^{-2} s^{-1} with a hold time of 1.5 h at each intensity. After the maximum light period the intensity was programmed to decrease in reverse order with the same hold times until complete darkness again at 19:30.

Methane production in oxic soils has been previously discussed and attributed mainly to abiotic

factors (Jugold et al., 2012) or methanogenic *Archaea* (Hao, Scharffe, Crutzen, & Sanhueza,

1988), although the latter was thought unlikely (Jugold et al., 2012; Kammann, Hepp, Lenhart, &

Müller, 2009). Here we show that a typical desert crust *Cyanobacteria* (identified in this study as

Leptolyngbya sp.), as well as the most common endolithic cyanobacterium *Chroococcidiopsis*

(Garcia-Pichel, Belnap, Neuer, & Schanz, 2003) produce CH_4 both under light and dark conditions

 (Fig. 1, Fig. S2), thus inferring a new but as yet unknown and unaccounted for source of CH_4 from

 oxic soils.

Figure 3. Average CH_4 production rates (μmol g DW^{-1} h^{-1}) obtained from multiple long-term measurements (2-5 days) using a membrane inlet mass spectrometer. The rates are designated by colour according to the environment from which the *Cyanobacteria* were originally isolated; dark blue, light blue and green for marine, freshwater and soil environments, respectively. Black and red lines represent median and mean values, respectively.

Cyanobacteria are ubiquitous in nature and their presence in aquatic systems is expected to

increase with eutrophication and rising global temperatures (Visser et al., 2016). The "methane

paradox" describing the production of CH_4 in oxic water layers has been known for four decades

(Scranton & Farrington, 1977). Though values may vary between water bodies, a recent study

suggests that up to 90 % of CH_4 emitted from freshwater lakes can be produced in the oxic layer

(Donis et al., 2017) with *Cyanobacteria* often being associated with elevated CH_4 concentration

in oxygen supersaturated freshwater systems (Grossart, Frindte, Dziallas, Eckert, & Tang, 2011).

In open oceanic environments, distant from any coastal, the contribution of lateral transport from

anoxic environments is expected to nonexistent. Nevertheless, based on the emission rates of our

laboratory investigations it is difficult to extrapolate the contribution of cyanobacteria to marine,

freshwater and terrestrial environments and finally to the global scale. First, only one attempt has

been done to estimate the global cyanobacterial biomass (Garcia-Pichel et al., 2003). This study

does not account for the increase in blooms of toxic and non-toxic cyanobacteria in freshwater

systems (Bowling, Blais, & Sinotte, 2015; Glibert, Maranger, Sobota, & Bouwman, 2014;

Huisman et al., 2018; Paerl & Huisman, 2008; Visser et al., 2016), nor for less monitored

cyanobacterial environments such under the ice-cover of frozen lakes (Bižić-Ionescu, Amann, &

Grossart, 2014). Recent evaluations of *Prochlorococcus* (Lange et al., 2018) suggest a global

biomass larger by 33 % than estimated in 2003 by Garcia-Pichel *et al*. Second, while our

experiments demonstrate inarguably the ability of *Cyanobacteria* to produce CH_4 independent of

external substrates, as well as to transfer fixed CO_2 to CH_4 under laboratory conditions, we cannot

account for the effect of nutrients concentrations and light quality in the natural environment.

Nevertheless, to get a first idea of what the laboratory rates might sum up to when applied to the

natural environment we performed a simple mathematical exercise for the oceanic

Prochlorococcus community. As suggested before, since experiments with low light (< 20 µmol

quanta m^{-2} s^{-1}) showed no detectable CH_4 production only surface *Prochlorococcus* communities

were used for this calculation. As such, only rates from High-Light *Prochlorococcus* strains were

used i.e. MIT9312, MIT0604 and MED4 averaging at 0.4 µmol CH_4 h^{-1} 10^{-6} cells. Based on recent

estimates of *Prochlorococcus* abundances (Lange et al., 2018) the standing stock global surface

communities were estimated to consist of 2.48×10^{19} cells (out of a total of 3.4×10^{27}). When taken

together, these numbers result in a potential production by global surface *Prochlorococcus*

communities of 1.39 Tg CH_4 y^{-1}. This number is not to be confused with the oceanic CH_4 flux,

estimated at 1.2 Tg CH_4 y^{-1} (Rhee, Kettle, & Andreae, 2009), which is the result of production by

multiple processes, transport and consumption by methanotrophs.

In this study, we show that *Cyanobacteria* can readily convert fixed inorganic carbon directly to

CH_4 and emit the potent greenhouse gas under both light and dark conditions. This is in addition

to the already established ability of *Cyanobacteria* to produce CH_4 by the demethylation of

methylphosphonates (Beversdorf et al., 2010; Gomez-Garcia et al., 2011). *Cyanobacteria* as a

phylum are the most ubiquitous group of organisms on Earth, thriving in most, naturally and

artificially, illuminated environments almost regardless of temperatures, salinity and nutrient

concentrations. Accordingly, their ability to produce CH_4 via different pathways, likely related to

their surroundings, makes them important to the present and future global CH_4 cycle and budget.

Even more so, as blooms of cyanobacteria are increasing with eutrophication and rising global

temperatures (Huisman et al., 2018; Visser et al., 2016). Furthermore, as phototrophic prokaryotes

such as *Cyanobacteria* have been inhabiting Earth for more than 3.5 billion years (Falcón,

Magallón, & Castillo, 2010; Frei et al., 2016) they may have had a major contribution to Earth's

CH_4 cycle such as during the great oxygenation event or even earlier when the conditions on Earth

were more reductive favoring CH_4 production.

Further research, however, is needed to elucidate the biochemical pathways of CH_4 formation in

Cyanobacteria and fully assess its possible relevance for ecology and the global CH_4 budget

throughout Earth history and how it might change in the future.

Acknowledgements

In-situ probe data was obtained from the LakeLab (www.lakelab.de) as part of the routine

monitoring of Lake Stechlin run by the Leibniz Institute of Freshwater Ecology and Inland

Fisheries. We thank the technical assistants of department III for making this data available. We

thank the members of the MIBI group and particularly Jason Woodhouse for assistance in

sampling, analysis and discussion of the data. We thank Tim Urich from Greifswald University

for the qPCR analysis. We thank Claudia Steglich and Wolfgang Hess from Freiburg University

for the *Prochlorococcus* culture. We thank Falk Pollehne from the Institute of Baltic Sea Research

for the *Nodularia* culture. We thank Meri Eichner from the Max Planck Institute for Marine

Microbiology for the *Trichodesmium* culture. We thank Daniel Sher and Dalit Roth from Haifa

University for the cultures of *Synechococcus* WH7803 and WH8102 and *Prochlorococcus*

MIT9312, NATL2A, MIT0604, MED4. Funding was provided to M.B.I., D.I. and H.P.G. via the

DFG-Aquameth project (GR1540-21-1), the BMBF-BIBS project (01LC1501G) and the Human

Frontiers Science project (HFSP 2039371). F.K. and T.K. were supported by the German Research

Foundation (DFG; KE 884/8-2, KE 884/11-1 and KE 884/16-2).

Author Contributions

M.B.I., T.K., D.I., F.K., H.P.G., conceived the study and designed the experiments; M.B.I., D.I.,

M.Y.H., M.G., A.M.M.P., W.E., performed MIMS experiments and *in-situ* measurements and

analyzed the data; T.K. performed stable isotope measurements and together with F.K. analyzed

the data. M.B.I., D.I., performed microbial community data analysis; A.M.M.P. analyzed

transcriptomics data. M.B.I., T.K., D.I., M.Y.H., M.G., A.M.M.P., W.E., F.K., H.P.G. discussed

the results and wrote the paper.

References

Angle, J. C., Morin, T. H., Solden, L. M., Narrowe, A. B., Smith, G. J., Borton, M. A., … Wrighton, K. C. (2017). Methanogenesis in oxygenated soils is a substantial fraction of wetland methane emissions. *Nature Communications*, *8*(1), 1567. https://doi.org/10.1038/s41467-017-01753-4

Berg, A., Lindblad, P., & Svensson, B. H. (2014). Cyanobacteria as a source of hydrogen for methane formation. *World Journal of Microbiology and Biotechnology*, *30*(2), 539–545. https://doi.org/10.1007/s11274-013-1463-5

Beversdorf, L. J., White, A. E., Björkman, K. M., Letelier, R. M., & Karl, D. M. (2010). Phosphonate metabolism by *Trichodesmium* IMS101 and the production of greenhouse gases. *Limnology and Oceanography*, *55*(4), 1768–1778. https://doi.org/10.4319/lo.2010.55.4.1768

Bižić-Ionescu, M., Amann, R., & Grossart, H.-P. (2014). Massive regime shifts and high activity of heterotrophic bacteria in an ice-covered lake. *PLoS ONE*, *9*(11), e113611. https://doi.org/10.1371/journal.pone.0113611

Bowling, L. C., Blais, S., & Sinotte, M. (2015). Heterogeneous spatial and temporal cyanobacterial distributions in Missisquoi Bay, Lake Champlain: An analysis of a 9 year data set. *Journal of Great Lakes Research*, *41*(1), 164–179. https://doi.org/10.1016/J.JGLR.2014.12.012

Chen, Y.-B., Zehr, J. P., & Mellon, M. (1996). Growth and nitrogen fixation of the diazotrophic filamentous nonheterocystous cyanobacterium trichodesmium sp. IMS 101 in defined media: evidence for a circadian rhythm. *Journal of Phycology*, *32*(6), 916–923. https://doi.org/10.1111/j.0022-3646.1996.00916.x

Donis, D., Flury, S., Stöckli, A., Spangenberg, J. E., Vachon, D., & McGinnis, D. F. (2017). Full-scale evaluation of methane production under oxic conditions in a mesotrophic lake. *Nature Communications*, *8*(1), 1661. https://doi.org/10.1038/s41467-017-01648-4

Dyhrman, S. T., Benitez-Nelson, C. R., Orchard, E. D., Haley, S. T., & Pellechia, P. J. (2009). A microbial source of phosphonates in oligotrophic marine systems. *Nature Geoscience*, *2*(10), 696–699. https://doi.org/10.1038/ngeo639

Falcón, L. I., Magallón, S., & Castillo, A. (2010). Dating the cyanobacterial ancestor of the chloroplast. *The ISME Journal*, *4*(6), 777–783. https://doi.org/10.1038/ismej.2010.2

Fischer, M. A., Güllert, S., Neulinger, S. C., Streit, W. R., & Schmitz, R. A. (2016). Evaluation of 16S rRNA Gene Primer Pairs for Monitoring Microbial Community Structures Showed High Reproducibility within and Low Comparability between Datasets Generated with Multiple Archaeal and Bacterial Primer Pairs. *Frontiers in Microbiology*, *7*. https://doi.org/10.3389/fmicb.2016.01297

Flaherty, B. L., Van Nieuwerburgh, F., Head, S. R., & Golden, J. W. (2011). Directional RNA deep sequencing sheds new light on the transcriptional response of Anabaena sp. strain PCC 7120 to combined-nitrogen deprivation. *BMC Genomics*, *12*, 332.

https://doi.org/10.1186/1471-2164-12-332

Frei, R., Crowe, S. A., Bau, M., Polat, A., Fowle, D. A., & Døssing, L. N. (2016). Oxidative elemental cycling under the low O2 Eoarchean atmosphere. *Scientific Reports*, *6*(1), 21058. https://doi.org/10.1038/srep21058

Garcia-Pichel, F., Belnap, J., Neuer, S., & Schanz, F. (2003). Estimates of global cyanobacterial biomass and its distribution. *Algological Studies*, *109*(1), 213–227. https://doi.org/10.1127/1864-1318/2003/0109-0213

Glibert, P. M., Maranger, R., Sobota, D. J., & Bouwman, L. (2014). The Haber Bosch–harmful algal bloom (HB–HAB) link. *Environmental Research Letters*, *9*(10), 105001. https://doi.org/10.1088/1748-9326/9/10/105001

Gomez-Garcia, M. R., Davison, M., Blain-Hartnung, M., Grossman, A. R., & Bhaya, D. (2011). Alternative pathways for phosphonate metabolism in thermophilic cyanobacteria from microbial mats. *The ISME Journal*, *5*(1), 141–149. https://doi.org/10.1038/ismej.2010.96

Grossart, H.-P., Frindte, K., Dziallas, C., Eckert, W., & Tang, K. W. (2011). Microbial methane production in oxygenated water column of an oligotrophic lake. *Proceedings of the National Academy of Sciences*, *108*(49), 19657–19661. https://doi.org/10.1073/pnas.1110716108

Guillard, R. R. L., & Ryhter, J. H. (1962). Studies of marine planktonic diatoms. *Can. J. Microbiol.*, *8*, 229–239.

Gumsley, A. P., Chamberlain, K. R., Bleeker, W., Söderlund, U., de Kock, M. O., Larsson, E. R., & Bekker, A. (2017). Timing and tempo of the Great Oxidation Event. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(8), 1811–1816. https://doi.org/10.1073/pnas.1608824114

Hao, W. M., Scharffe, D., Crutzen, P. J., & Sanhueza, E. (1988). Production of N2O, CH4, and CO2 from soils in the tropical savanna during the dry season. *Journal of Atmospheric Chemistry*, *7*(1), 93–105. https://doi.org/10.1007/BF00048256

Hubalek, V., Wu, X., Eiler, A., Buck, M., Heim, C., Dopson, M., … Ionescu, D. (2016). Connectivity to the surface determines diversity patterns in subsurface aquifers of the Fennoscandian shield. *The ISME Journal*, *10*(10), 2447–2458. https://doi.org/10.1038/ismej.2016.36

Huisman, J., Codd, G. A., Paerl, H. W., Ibelings, B. W., Verspagen, J. M. H., & Visser, P. M. (2018). Cyanobacterial blooms. *Nature Reviews Microbiology*, *16*(8), 471–483. https://doi.org/10.1038/s41579-018-0040-1

Intergovernmental Panel on Climate Change (Ed.). (2014). *Climate Change 2013 - The Physical Science Basis*. Cambridge: Cambridge University Press. https://doi.org/10.1017/CBO9781107415324

Ionescu, D., Siebert, C., Polerecky, L., Munwes, Y. Y., Lott, C., Häusler, S., … de Beer, D. (2012). Microbial and chemical characterization of underwater fresh water springs in the Dead Sea. *PloS One*, *7*(6), e38319. https://doi.org/10.1371/journal.pone.0038319

Jugold, A., Althoff, F., Hurkuck, M., Greule, M., Lenhart, K., Lelieveld, J., & Keppler, F. (2012). Non-microbial methane formation in oxic soils. *Biogeosciences*, *9*(12), 5291–5301. https://doi.org/10.5194/bg-9-5291-2012

Kammann, C., Hepp, S., Lenhart, K., & Müller, C. (2009). Stimulation of methane consumption by endogenous CH4 production in aerobic grassland soil. *Soil Biology and Biochemistry*, *41*(3), 622–629. https://doi.org/10.1016/j.soilbio.2008.12.025

Kana, T. M., Cornwell, J. C., & Zhong, L. (2006). Determination of denitrification in the Chesapeake Bay from measurements of N2 accumulation in bottom water. *Estuaries and Coasts*, *29*(2), 222–231. https://doi.org/10.1007/BF02781991

Keppler, F., Hamilton, J. T. G., Braß, M., & Röckmann, T. (2006). Methane emissions from terrestrial plants under aerobic conditions. *Nature*, *439*(7073), 187–191. https://doi.org/10.1038/nature04420

Keppler, F., Schiller, A., Ehehalt, R., Greule, M., Hartmann, J., & Polag, D. (2016). Stable isotope and high precision concentration measurements confirm that all humans produce and exhale methane. *Journal of Breath Research*, *10*(1), 016003. https://doi.org/10.1088/1752-7155/10/1/016003

Knapp, A. N. (2012). The sensitivity of marine N2 fixation to dissolved inorganic nitrogen. *Frontiers in Microbiology*, *3*(OCT), 374. https://doi.org/10.3389/fmicb.2012.00374

Lange, P. K., Brewin, R. J. W., Dall'Olmo, G., Tarran, G. A., Sathyendranath, S., Zubkov, M., & Bouman, H. A. (2018). Scratching beneath the surface: A model to predict the vertical distribution of Prochlorococcus using remote sensing. *Remote Sensing*, *10*(6), 847. https://doi.org/10.3390/rs10060847

Lenhart, K., Bunge, M., Ratering, S., Neu, T. R., Schüttmann, I., Greule, M., … Keppler, F. (2012). Evidence for methane production by saprotrophic fungi. *Nature Communications*, *3*, 1046. https://doi.org/10.1038/ncomms2049

Lenhart, K., Klintzsch, T., Langer, G., Nehrke, G., Bunge, M., Schnell, S., & Keppler, F. (2016). Evidence for methane production by the marine algae Emiliania huxleyi. *Biogeosciences*, *13*(10), 3163–3174. https://doi.org/10.5194/bg-13-3163-2016

Lenhart, K., Weber, B., Elbert, W., Steinkamp, J., Clough, T., Crutzen, P., … Keppler, F. (2015). Nitrous oxide and methane emissions from cryptogamic covers. *Global Change Biology*, *21*(10), 3889–3900. https://doi.org/10.1111/gcb.12995

Luo, C., & Konstantinidis, K. T. (2011). Phosphorus-related gene content is similar in Prochlorococcus populations from the North Pacific and North Atlantic Oceans. *Proceedings of the National Academy of Sciences of the United States of America*, *108*(16), E62-3; author reply E64-6. https://doi.org/10.1073/pnas.1018662108

Mitschke, J., Vioque, A., Haas, F., Hess, W. R., & Muro-Pastor, A. M. (2011). Dynamics of transcriptional start site selection during nitrogen stress-induced cell differentiation in Anabaena sp. PCC7120. *Proceedings of the National Academy of Sciences of the United States of America*, *108*(50), 20130–20135. https://doi.org/10.1073/pnas.1112724108

Moore, L. R., Coe, A., Zinser, E. R., Saito, M. A., Sullivan, M. B., Lindell, D., … Chisholm, S.

W. (2007). Culturing the marine cyanobacterium Prochlorococcus. *Limnology and Oceanography: Methods*, *5*(10), 353–362. https://doi.org/10.4319/lom.2007.5.353

Nercessian, O., Noyes, E., Kalyuzhnaya, M. G., Lidstrom, M. E., & Chistoserdova, L. (2005). Bacterial populations active in metabolism of C1 compounds in the sediment of Lake Washington, a freshwater lake. *Applied and Environmental Microbiology*, *71*(11), 6885–6899. https://doi.org/10.1128/AEM.71.11.6885-6899.2005

Paerl, H. W., & Huisman, J. (2008). CLIMATE: Blooms Like It Hot. *Science*, *320*(5872), 57–58. https://doi.org/10.1126/science.1155398

Pfreundt, U., Kopf, M., Belkin, N., Berman-Frank, I., & Hess, W. R. (2014). The primary transcriptome of the marine diazotroph Trichodesmium erythraeum IMS101. *Scientific Reports*, *4*, 6187. https://doi.org/10.1038/srep06187

Powell, R. J. (1972). Solubility of 16 gases in heptacosafluorotributylamine and carbon disulfide. *Journal of Chemical & Engineering Data*, *17*(3), 302–304. https://doi.org/10.1021/je60054a024

Puente-Sánchez, F., Arce-Rodríguez, A., Oggerin, M., García-Villadangos, M., Moreno-Paz, M., Blanco, Y., … Parro, V. (2018). Viable cyanobacteria in the deep continental subsurface. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(42), 10702–10707. https://doi.org/10.1073/pnas.1808176115

Repeta, D. J., Ferrón, S., Sosa, O. A., Johnson, C. G., Repeta, L. D., Acker, M., … Karl, D. M. (2016). Marine methane paradox explained by bacterial degradation of dissolved organic matter. *Nature Geoscience*, *9*(12), 884–887. https://doi.org/10.1038/ngeo2837

Rhee, T. S., Kettle, A. J., & Andreae, M. O. (2009). Methane and nitrous oxide emissions from the ocean: A reassessment using basin-wide observations in the Atlantic. *Journal of Geophysical Research*, *114*(D12), D12304. https://doi.org/10.1029/2008JD011662

Savitzky, A., & Golay, M. J. E. (1964). Smoothing and differentiation of data by simplified least squares procedures. *Analytical Chemistry*, *36*(8), 1627–1639. https://doi.org/10.1021/ac60214a047

Schlüter, M., & Gentz, T. (2008). Application of membrane inlet mass spectrometry for online and in situ analysis of methane in aquatic environments. *Journal of the American Society for Mass Spectrometry*, *19*(10), 1395–1402. https://doi.org/10.1016/j.jasms.2008.07.021

Scranton, M. I., & Farrington, J. W. (1977). Methane Production in the Waters off Walvis Bay. *Journal of Geophysical Research*, *82*(31), 4947–4953. https://doi.org/10.1029/JC082i031p04947

Stal, L. J., & Moezelaar, R. (1997). Fermentation in Cyanobacteria. *FEMS Microbiology Reviews*, *21*(2), 179–211. https://doi.org/10.1111/j.1574-6976.1997.tb00350.x

Stanier, R. Y., Deruelles, J., Rippka, R., Herdman, M., & Waterbury, J. B. (1979). Generic assignments, strain histories and properties of pure cultures of Cyanobacteria. *Microbiology*, *111*(1), 1–61. https://doi.org/10.1099/00221287-111-1-1

Tian, H., Langenfelds, R., Thornton, B. F., Kleinen, T., Locatelli, R., Etiope, G., … Covey, K.

(2016). The global methane budget 2000–2012. *Earth System Science Data*, *8*(2), 697–751. https://doi.org/10.5194/essd-8-697-2016

Tuboly, E., Szabó, A., Garab, D., Bartha, G., Janovszky, Á., Ero″s, G., … Boros, M. (2013). Methane biogenesis during sodium azide-induced chemical hypoxia in rats. *American Journal of Physiology-Cell Physiology*, *304*(2), C207–C214. https://doi.org/10.1152/ajpcell.00300.2012

Visser, P. M., Verspagen, J. M. H., Sandrini, G., Stal, L. J., Matthijs, H. C. P., Davis, T. W., … Huisman, J. (2016). How rising CO2 and global warming may stimulate harmful cyanobacterial blooms. *Harmful Algae*, *54*, 145–159. https://doi.org/10.1016/j.hal.2015.12.006

Wang, Q., Dore, J. E., & McDermott, T. R. (2017). Methylphosphonate metabolism by Pseudomonas sp. populations contributes to the methane oversaturation paradox in an oxic freshwater lake. *Environmental Microbiology*, *19*(6), 2366–2378. https://doi.org/10.1111/1462-2920.13747

Yao, M., Henny, C., & Maresca, J. A. (2016). Freshwater bacteria release methane as a by-product of phosphorus acquisition. *Applied and Environmental Microbiology*, *82*(23), 6994–7003. https://doi.org/10.1128/AEM.02399-16

Zheng, Y., Harris, D. F., Yu, Z., Fu, Y., Poudel, S., Ledbetter, R. N., … Harwood, C. S. (2018). A pathway for biological methane production using bacterial iron-only nitrogenase. *Nature Microbiology*, *3*(3), 281–286. https://doi.org/10.1038/s41564-017-0091-5

Fig. S1. Temporal profiles of CH_4, O_2 and cyanobacterial derived Chl a between July 2014 and

July 2016. The CH_4 data was measured every 1 - 4 weeks depending on season using a GC-FID

as described in Grossart *et al.* (Grossart *et al.*, 2011). O_2 and Chl a were measured hourly using a

YSI and a BBE probe (see www.lake-lab.de), respectively.

Fig. S2. Continuous measurements of CH_4 and oxygen under light/dark cycles using a membrane

inlet mass spectrometer (MIMS) in 17 different cyanobacterial cultures. A decrease in CH_4

concentration is a result of reduced (or no) production coupled with degassing from the

supersaturated, continuously-mixing, semi-open incubation chamber towards equilibrium with

atmospheric CH_4 (2.5 nM and 2.1 nM for freshwater and seawater, respectively). Calculated CH_4

production rates account for the continuous emission of CH_4 from the incubation chamber for as

long as the CH_4 concentrations are supersaturated. The light regime for the experiments was as

follows: dark (black bar) from 19:30 to 09:00 then light intensity (yellow bar) was programmed to

increase to 60, 120, 180, 400 µmol quanta m^{-2} s^{-1} with a hold time of 1.5 h at each intensity. After

maximum light period the intensity was programmed to decrease in reverse order with the same

hold times until complete darkness again at 19:30.

Fig. S3. Raw signal obtained from mass 15 (CH_4) and mass 40 (Argon) plotted against the

calculated solubility at different salinities at 30 °C. The signal in both cases is linearly correlated

to the concentration of the dissolved gas. The ratio between the two masses was extrapolated

between 0 and 50 ppt and was used for calculating the CH_4 concentration.

Fig. S4. Community composition of the cyanobacterial cultures as obtained when sequenced using

Archaea specific primers. Methanogenic *Archaea* (red) are in most cases lower that 0.1% of the

obtained sequences. In the absence of *Archaea* DNA template, the primers amplify DNA of

Bacteria. The background presence or complete absence of methanogens was confirmed by qPCR

of the *mcrA* gene.

Fig. S5. Average CH4 production rates (µmol gDW-1 h-1) obtained from multiple long-term measurements (2-5 days) with a membrane inlet mass spectrometer. The rates are shown by colour according to the environment from which the *Cyanobacteria* were originally isolated; dark blue, light blue and green for marine, freshwater and soil environments, respectively. The rates are presented in comparison to three known methanogens. Rates for the methanogens were obtained from references: Mountfort and Asher (Mountfort and Asher, 1979), Kröninger *et al*. (Kröninger *et al.*, 2017) and Gerhard *et al*.(Gerhard *et al.*, 1993). Rates for eukaryotes including marine algae and terrestrial plants were taken from Lenhart *et al*. (Lenhart *et al.*, 2016), Keppler *et al*. (Keppler *et al.*, 2006), Brüggemann *et al*. (Brüggemann *et al.*, 2009), Wishkermann *et al*. (Wishkerman *et al.*, 2011) and Qaderi *et al*. (Qaderi and Reid, 2009) No emission rates (on a dry weight basis) are available for fungi and animals.