Gene flow improves fitness at a range edge under climate change

Megan Bontrager[1,3] and Amy L. Angert[2]

[1]Department of Botany, University of British Columbia, Vancouver, British Columbia V6T
1Z4, Canada

[2]Departments of Botany and Zoology, University of British Columbia, Vancouver, British
Columbia V6T 1Z4, Canada

[3]Author for correspondence (mgbontrager@gmail.com)

Article type: Letter

Running title: Gene flow at a range edge under climate change

Keywords: *Clarkia pulchella*, climate change, common garden, gene flow, geographic range limits,
local adaptation, swamping gene flow

Abstract word count: 291

Total word count: 5215

1 Abstract

Populations at the margins of a species' geographic range are often thought to be poorly adapted

to their environment. According to theoretical predictions, gene flow can inhibit these range edge

populations if it disrupts adaptation to local conditions. Alternatively, if range edge populations are

small or isolated, gene flow can provide beneficial genetic variation, and may facilitate adaptation

to environmental change. We tested these competing predictions in the annual wildflower *Clarkia*

pulchella using greenhouse crosses to simulate gene flow from sources across the geographic range

into two populations at the northern range margin. We planted these between-population hybrids in

common gardens at the range edge, and evaluated how genetic differentiation and climatic differences

between edge populations and gene flow sources affected lifetime fitness. During an anomalously

warm study year, gene flow from populations occupying historically warm sites improved fitness at

the range edge, and plants with one or both parents from warm populations outperformed local

plants. The effects of the temperature provenance of gene flow sources were most apparent at early

life history stages, but precipitation provenance also affected reproduction. We also found benefits

of gene flow that were independent of climate: after climate was controlled for, plants with parents

from different populations performed better at later lifestages than those with parents from the

same population, indicating that gene flow may improve fitness via relieving homozygosity. Further

supporting this result, we found that increasing genetic differentiation of parental populations had

positive effects on fitness of hybrid seeds. Gene flow from warmer populations, when it occurs,

is likely to contribute adaptive genetic variation to populations at the northern range edge as the

climate warms. On heterogeneous landscapes, climate of origin may be a better predictor of gene

flow effects than geographic origin.

Impact summary

What limits species geographic ranges on the landscape? One process of interest when trying to answer this question is gene flow, which is the movement of genetic material between populations, as might occur in plants when seeds or pollen move across the landscape. One hypothesis that has been proposed is that gene flow from populations in other environments prevents populations at range edges from adapting to their local habitats. Alternatively, it has been suggested that these populations might benefit from gene flow, as it would provide more genetic material for natural selection to act upon.

We tested these predictions in an annual wildflower, *Clarkia pulchella*. We simulated gene flow by pollinating plants from the range edge with pollen from other populations. Then we planted the resulting seeds into common gardens in the home sites of the range edge populations and recorded their germination, survival, and reproduction. The weather during our experiment was much warmer than historic averages in our garden sites, and perhaps because of this, we found that gene flow from warm locations improved the performance of range edge populations. This result highlights the potential role of gene flow and dispersal in aiding adaptation to warming climates. We also found some positive effects of gene flow that were independent of climate. Even after we statistically controlled for adaptation to temperature and precipitation, plants that were the result of gene flow pollinations produced more seeds and fruits than plants with both parents from the same population. Rather than preventing adaptation, in our experiment, gene flow generally had positive effects on fitness.

Introduction

Species are limited in their geographic extents on the landscape. In many cases, the limits of species' geographic distributions are the result of niche limitation, rather than simply an inability to disperse to suitable areas beyond their current distribution (Lee-Yaw et al., 2016). This raises the question of what prevents populations on the range periphery from adapting to sites beyond the range edge (Antonovics, 1976; Bridle and Vines, 2007), particularly when boundaries are not co-incident with an abrupt shift in the abiotic environment. The putative causes of limits to adaptation at the range edge hinge upon demographic and genetic features of metapopulations (Sexton et al., 2009).

If range limits represent limits to adaptation, this could be the result of insufficient genetic variation in range edge populations. Limited genetic variation at range edges is predicted because range edge populations are often characterized as small, with more frequent or severe changes in population size and elevated rates of turnover relative to central populations (Vucetich and Waite, 2003). Populations that are on the leading edge of range expansions may also exhibit patterns of low genetic variation as a result of successive founder events (Pujol and Pannell, 2008). Significant declines in neutral genetic variation near range edges is a common (though not ubiquitous) pattern (Eckert et al., 2008; Pironon et al., 2017), indicating that some of these processes are likely to affect some range edges in some species. If the observed declines in neutral variation also reflect reduced adaptive genetic variation, this might result in marginal populations being less locally adapted when compared to central populations, as they have less capacity to respond to local selection pressures. Maladaptation is expected to lead to poor demographic performance, reducing colonization opportunities in sites beyond the range, and potentially creating (or reinforcing) a range edge at equilibrium along an environmental gradient (Kirkpatrick and Barton, 1997).

Swamping gene flow is another often-invoked hypothesis for how equilibrial range limits might form and persist (Lenormand, 2002; Sexton et al., 2009). Under swamping gene flow, peripheral populations are unable to adapt to their local conditions because they experience maladaptive gene flow from central populations (Kirkpatrick and Barton, 1997). This process is predicted to occur when populations are arranged along an environmental gradient where individuals are well-adapted and abundant in the center of that gradient. Because of this asymmetry in abundance, net gene

flow is asymmetric and brings alleles that are adaptive in central environments to edge popula-

tions, disrupting local adaptation to edge environments. This causes edge populations to become

demographic sinks, where death rates exceed birth rates, and prevents further range expansion.

According to this model, the fitness of edge populations will depend upon the rate of gene flow

from center to edge as well as the steepness of the environmental gradient (i.e., the magnitude of

environmental differences between the sources of the gene flow and the recipient populations).

Comprehensive empirical tests of the swamping gene flow hypothesis are difficult to conduct

because they require demonstrating both the negative effects of gene flow on edge populations

as well as the occurrence of asymmetric gene flow on the landscape. Evidence to-date indicates

that swamping gene flow might limit adaptation along geographic gradients in some systems (Paul

et al., 2011) and sometimes limit the geographic range (Fedorka et al., 2012; Holliday et al., 2012).

However, in other systems there are no detectable fitness costs of gene flow across environmental

gradients (Emery, 2009; Moore and Hendry, 2009; Samis et al., 2016) and strong local adaptation

persists despite gene flow (Yeaman and Jarvis, 2006; Gould et al., 2014). Outbreeding depression

may generate patterns similar to those resulting from the disruption of local adaptation, but the

effects of genetic incompatibilities can be discerned from those of swamping by experimental designs

that allow for decoupling of environmental and genetic differentiation.

Most theory about swamping gene flow at range edges has been developed with the assumption

of smooth environmental gradients underlying the range, however, this assumption is unrealistic

for most species. Topography, continentality, and other landscape features make transects from

range centers to edges heterogenous with regards to climate. Other habitat variables, such as soil

type or the biotic community (which may mediate responses to climate in addition to imposing

selection on their own), are also likely to be spatially heterogeneous. This complicates predictions

of the swamping gene flow hypothesis: range edge populations may experience gene flow from

environmentally divergent neighboring populations, or environmentally similar central populations,

as well as combinations falling anywhere in between. In this case, geography cannot be used as

a proxy for predictions about the effects of gene flow, rather, these predictions must be informed

by the environmental differences between populations. Gene flow between populations in similar

environments may be beneficial, even when populations are geographically disparate, because gene

flow can allow for the spread of environment-specific beneficial alleles that arise in a single population

(Sexton et al., 2011). Abundant-center distribution patterns and asymmetric gene flow have been

documented in some species, but are not ubiquitous (Sagarin and Gaines, 2002), perhaps at least

in part as a result of complex environmental gradients.

In addition to contributing alleles that are adaptive or maladaptive in a given environment, gene

flow may provide relief from homozygosity caused by drift or inbreeding. Gene flow is expected to

increase heterozygosity and reintroduce variation that can allow for masking or purging of fixed

deleterious alleles. As a result, gene flow can improve fitness in peripheral populations (Sexton

et al., 2011). The extent to which gene flow causes heterosis depends upon the genetic divergence of

populations (Ingvarsson and Whitlock, 2000), but not explicitly on the magnitude of the environ-

mental differences between the source and recipient of gene flow, though environmental differences

are correlated with genetic differentiation in some species (Sexton et al., 2014).

Gene flow may also be beneficial when maladaptation arises due to disequilibrium between a

populations' optimal conditions and the environment. This could occur when a species is undergoing

a range expansion, or when the environmental landscape is moving out from under individuals, as is

occurring under climate change (Aitken and Whitlock, 2013). If a population is locally adapted to

historic conditions in a site, and the environment changes rapidly, then gene flow from populations

with historic conditions that are more similar to these new local conditions is expected to improve

population performance.

To investigate how gene flow affects peripheral populations, we simulated gene flow among

populations spanning the northern half of the range of an annual wildflower, *Clarkia pulchella*,

and measured lifetime fitness of individuals in two common gardens at the species' northern range

edge. We asked 1) Are range edge populations of *C. pulchella* locally adapted? 2) What climatic

factors predict fitness at the northern range edge? 3) Does gene flow positively or negatively affect

edge populations? and 4) How does the effect of gene flow from other populations depend upon

the genetic differentiation and climatic distances of these populations? Under conditions where

the range edge is not at equilibrium with climate, we expect that gene flow from sites that are

historically similar to the experimental conditions will improve performance. Under conditions in which this species' range is at equilibrium with climate, and if this edge is limited by adaptation, we expect that gene flow from populations in similar climates will have a positive effect on fitness via heterosis or the contribution of adaptive alleles, but that gene flow from strongly contrasting climates will be detrimental. If populations have genetic incompatibilities (which need not be the result of divergent selection, but could simply be the result of drift under prolonged separation) then we expect the offspring of crosses between populations that are more genetically divergent to perform worse, regardless of the conditions of the test environment. However, if heterozygosity is positively related to fitness we expect a greater benefit from gene flow between more genetically divergent populations.

Methods

Study system, seed collection, and site selection

Clarkia pulchella Pursh (Onagraceae) is a winter annual that grows on sparsely vegetated, south-facing slopes with low canopy cover throughout eastern Washington and Oregon, Idaho, and western Montana (United States) and southeastern British Columbia (Canada). This species germinates in fall and overwinters as a seedling before flowering in late May, June, and early July. It has no observed seed dormancy, but seeds will not germinate immediately upon dehiscing and require an after-ripening period of several weeks. It has showy pink flowers and is visited by a diverse array of pollinators though it has some capacity to self-pollinate in the absence of pollinators or mates (MacSwain et al., 1973; Bontrager et al., 2018). Individual plants typically produce fewer than 10 flowers, though some larger individuals may produce up to *c.* 100 on occasion.

Seeds of *C. pulchella* were collected from 15 populations in July 2014 (Figure 1A; Table S1). We used the two northwestern-most populations of the continuous distribution of *C. pulchella* as common garden sites (hereafter referred to as focal populations). Other populations (hereafter, donor populations) were selected with the goal of sampling representative variation in major climatic axes (temperature, precipitation, and seasonality of these variables; Figure 1B) across the northern

153   half of the species' range. In each of these populations, seeds were collected haphazardly from plants
154   spaced >0.5 m apart.

Greenhouse generation and crossing design

We grew field-collected seeds in the greenhouse and implemented a controlled crossing design to
generate seeds for the field transplant. We planted maternal families from each population into
22 randomized blocks, with each block containing one family from each population. Two types
of crosses were performed: "within-population" crosses and "between-population" crosses. For
within-population crosses, dams were pollinated using pollen from the plant of the same population
in the subsequent block (Figure S1A). Each plant from each population was therefore used as both
a sire and a dam with other plants from the same population. For between-population crosses,
flowers on plants from the two focal populations within each block were pollinated using each of
the donor plants in that block (Figure S1B). These crosses simulate an early stage of gene flow:
the progeny of a mating event that is the result of long distance pollen dispersal (or the progeny of
a cross between a native individual and a recent immigrant). We collected the seeds from crosses
as fruits ripened. Further details about our greenhouse conditions and crossing design are in the
Supplementary Methods.

Common garden design and installation

For the transplant, we used seeds from 15 of our greenhouse blocks, and substituted seeds from the
same type of cross from other greenhouse blocks when they were unavailable from our primary 15.
Seeds were glued to toothpicks to expedite planting and monitoring in the field. Two seeds were
glued to each toothpick with a tiny dab of water-soluble glue (when seeds were limited, just one
seed was glued to each toothpick). At each of the two sites, toothpicks were planted into 10 fully
randomized plots. Each plot contained two toothpicks from each cross type from each of the 15
replicates. We only planted between-population crosses with local dams at each of the two focal sites
(i.e., Blue Lake plots only contained between-population crosses performed on Blue Lake plants,
and Rock Creek plots only contained between-population crosses performed on Rock Creek plants).

Within-population crosses from all populations were planted out at both sites. Therefore, each plot

contained two replicates of each of 15 crosses from 29 cross types (14 between-population groups

and 15 within-population groups). For some cross types, less than 15 families had sufficient seeds

for the full design, therefore each plot contained 832 toothpicks at Rock Creek and 836 toothpicks

at Blue Lake. In total, our design included 16,680 toothpicks and 32,755 seeds. Seeds were planted

in September 2015. Details of transplant installation are in the Supplementary Methods.

Monitoring and measuring

Germination was censused 16-20 November 2015. We documented the emergence of either 0, 1, or 2

germinants at each toothpick. If two germinants were present, these were randomly thinned so that

just one remained. Overwinter survival was assessed 17-21 March 2016. During both germination

and overwinter survival censuses, plant size was measured as the distance in mm from the tip of

one leaf to the tip of the other (most plants had just one pair of leaves). In early June 2016

we censused the spring survival of all plants and began measuring reproduction every 2-3 days.

As a proxy for seed production, we measured the ovary length of each flower produced on each

plant. Pollinators were excluded from our plots to prevent the escape of non-native genotypes,

so we calibrated a conversion from ovary length to seed production using hand pollinations (see

Supplementary Methods). We continued reproductive censuses until all plants had senseced, but

reduced census frequency when flowering slowed in July.

Climatic and genetic differentiation

We compiled monthly temperature and precipitation data from 1951-1980 for all seed sources, as

well as the gardens during the months of the experiment (September 2015-July 2016) from PRISM

(PRISM Climate Group, 2017). We calculated historic (pre-warming, 1951-1980) climate averages

for each site, which we compared to conditions experienced by plants during the transplant in our

analyses. Inter-annual climate variability was very similar in each of the seed collection sites (data

not shown) so we do not further consider variability in climate, and focus on averages only. Pairwise

population genetic differentiation (F_{ST}) was calculated from 2982 SNPs that were genotyped in up

to 12 parental individuals from each population. F_{ST} was calculated using the implementation of
Weir and Cockerham (1984) in the R package hierfstat (Goudet and Jombart, 2015). These methods
are described in full in Bontrager and Angert (2018).

Statistical analyses

Did local populations outperform foreign populations?

We tested whether local populations were, on average, superior to foreign populations using within-
population crosses only. We compared lifetime fitness of local vs. foreign individuals in a generalized
linear mixed model (GLMM) with a zero-inflated negative binomial distribution using the package
glmmTMB (Brooks et al., 2017). These zero-inflated models allow specification of fixed effects
for both the zero-inflation part of the model (the probability of a non-zero value) as well as the
conditional part of the model (the effect on the response after accounting for zero-inflation). Gen-
erally, we consider the zero-inflation part of the model to reflect early lifestages, as the majority of
plants that produced zero seeds did so as a result of failing to germinate or survive winter. The
conditional part of the model may reflect both differences during reproduction as well as differences
among individuals accumulated across all lifestages. In addition to testing for local adaptation
represented by lifetime fitness, we tested whether local populations performed better than foreign
populations at any component lifestage: germination, size after germination, overwinter survival,
size after winter, fruit count, and estimated seed production. Plant size was modeled with a Gaus-
sian response distribution, germination and survival with binomial response distributions and logit
link functions, and fruit counts and seed production with zero-inflated negative binomial response
distributions and log link functions. For component lifestage analyses, we included only individuals
that had survived to the preceding census, and always included plant size at the previous census
to account for differences that had accumulated at earlier lifestages. For all of these models we
initially included a random effect structure of block within site, dam within dam population, and
sire within sire population. However, models of later lifestages and lifetime fitness frequently failed
to converge with this parameterization. When convergence failed, we reduced random effects to
only block within site and sire population. Predictions and confidence intervals were visualized

with ggeffects (Lüdecke, 2018).

Does climate of origin explain performance in common gardens?

We built GLMMs using the methods described above to evaluate the effects of mismatch between
the experiment conditions and population's historic climates on lifetime fitness and all fitness com-
ponents, again using only within-population crosses. We calculated the absolute difference between
the garden conditions and the historic temperature and precipitation of each source population. We
use absolute differences because we expect that mismatch in either direction along a climate axis
will negatively impact fitness. Very few source populations were from sites drier or hotter than con-
ditions during the experiment, so absolute differences mostly result from source populations being
historically cooler or wetter than the experiment. Our lifetime fitness model included absolute dif-
ferences in temperature (for the experiment duration, September-July) in both the conditional and
zero-inflation parts of the model, as well as absolute differences in spring and summer precipitation
(April-July) in the conditional part of the model only. We isolated the specific lifestages affected by
each of these climatic predictors using the methods described above. In lifestage-specific analyses,
we calculated climate differences using only the months in each census window. In all analyses, all
continuous predictors were scaled and centered.

Does gene flow help or hurt edge populations?

Based on the results of the analyses above, we expected that gene flow from some populations was
likely to confer benefits by contributing adaptive genetic variation to focal populations experiencing
an anomalous climate. To evaluate whether there were benefits of gene flow that were independent
of these climate effects, we calculated the midparent historic temperature average for all individuals
(that is, the average temperature of dam and sire sites) and then calculated the absolute difference
between this temperature and the experimental temperature (Figure S2). We calculated a metric
of absolute midparent precipitation difference (averaged over April-July) in the same manner. We
used GLMMs as described above to test for an effect of gene flow in addition to an anticipated effect
of midparent climate differences on lifetime fitness and each component lifestage. In these models,

gene flow was included as a categorical fixed effect (within-population cross vs. between-population

cross) along with midparent temperature and precipitation differences. We included gene flow and

temperature differences in both the conditional and zero-inflation parts of the lifetime fitness model,

and precipitation differences in only the conditional part. We had difficulty disentangling effects

of precipitation differences and gene flow (see Results), so we ran these models with and without

precipitation differences.

264 Do the effects of gene flow depend upon the genetic differentiation between focal and

265 donor populations?

We examined whether the genetic differentiation (F_{ST}) between the two parental populations of the

between-population crosses positively or negatively affected offspring fitness. We could only esti-

mate genetic differentiation between parental populations for individuals with parents from different

populations, so in these analyses we use between-population crosses only. We built zero-inflated

GLMMs as described above using lifetime fitness and included predictors of absolute midparent tem-

perature and precipitation differences as well as F_{ST}. We also tested the effects of these parameters

on each component lifestage. Our ability to detect significant effects of climate in full models was

limited, likely due to the narrow range of midparent climatic variability across between-population

crosses, so we also built separate models of each of our three predictors on lifetime fitness and each

lifestage. All statistical analyses were implemented in R version 3.4.3 (R Core Team, 2017).

276 Results

277 Climate of origin explains performance in common gardens

Local populations were not superior to the average foreign population in their cumulative fitness

across all lifestages, or in any component lifestage (Figure S3, Table S2). Populations that were best

matched to experimental temperatures performed best in our gardens; lifetime fitness declined with

increasing absolute temperature differences between the source and the experimental conditions

(Figure 2A). This occurred via effects on both the probability of producing any seeds (the zero-

inflation part of the model; $\beta = -0.341$, SE $= 0.033$, $P < 0.001$; Figure 2B), and the number of

seeds produced (the conditional part of the model; $\beta = -0.114$, SE $= 0.050$, $P = 0.022$; Figure 2C).
Local populations, which are historically intermediate in temperature (Figure 1B), were mismatched
from the experiment conditions and performed worse than populations from warmer sites that were
more climatically similar to the garden conditions.

Analyses of component lifestages support these inferences (Figure 2D, Table S3): being poorly
matched to experimental temperatures had negative effects on germination proportion, overwinter
survival, and the size of plants after winter. While precipitation differences were not significant in
the model of lifetime fitness ($\beta = -0.067$, SE $= 0.049$, $P = 0.177$, Figure 2C), they did have a
negative effect on seed production among plants surviving the winter (Figure 2D, Table S3).

293 Gene flow may confer some benefits to edge populations

As in the analyses of within-population plants only, both midparent temperature differences and
midparent precipitation differences had negative effects on lifetime fitness in our common gar-
dens (Table S4A; Figure 3AB). Gene flow (i.e., being a between-population rather than a within-
population cross) did not have a significant effect in the lifetime fitness model that also included
climate differences.

It is difficult to disentangle the effects of precipitation differences from the effects of gene flow in
these analyses. This is because our focal populations are already among the driest provenances in
our experiment. Therefore, the average between-population plant is better matched to the experi-
mental conditions than the average within-population plant, because the midparent precipitation of
between-population plants is always calculated with at least one very dry focal parent (Figure S2B).
This was not an issue with temperature differences, because our focal populations are intermedi-
ate to other provenances in terms of temperature (Figure S2A). When lifetime fitness was analyzed
without precipitation differences in the model, we found that gene flow (being a between-population
cross, rather than a within-population cross), had a positive effect on lifetime fitness in addition to
effects of temperature (Table S4B).

A small positive effect of gene flow, independent of climatic differences, is supported by analyses
of some lifestage components (Figure 3C; Table S4C). Negative effects of precipitation and tem-

perature differences were similar to those found in the analyses of climatic drivers of performance, while gene flow (i.e., being from a between-population vs. a within-population cross) had a positive effect on fruit production and a marginal positive effect on seed production.

Genetic differentiation between parental populations positively affects fitness

Both midparent temperature difference from the garden conditions and genetic differentiation between parental populations had significant effects on fitness (Figure 4, Table S5AB, Table S6A). Genetic differentiation between parental populations had a positive effect on lifetime fitness via increasing the probability of producing seeds (the zero-inflation part of the model). The effects of genetic differentiation on lifetime fitness were mirrored in the analyses of single lifestages: F_{ST} had a positive effect on germination, size after winter, fruit count, and seed production (Figure 4C, Table S5A, Table S6A).

The negative effect of midparent temperature differences in between-population crosses is generally consistent with our analyses of climatic drivers of performance in within-population crosses. Between-population plants with donor parents that were well-matched to temperatures during the experiment were more likely to produce seeds, as indicated by the significant negative effect of midparent temperature differences in the zero-inflation part of that model (Table S5B, Table S6A). Midparent temperature differences did not significantly affect any single lifestage, but had marginal negative effects on size after germination, size after winter, and fruit number (Figure 4C, Table S5B). Midparent precipitation differences did not significantly affect lifetime fitness or component lifestages (Table S5C).

It is important to note that genetic differentiation is not especially correlated with temperature differences (r = -0.25), though genetic differentiation and precipitation differences are correlated (r = 0.64), with plants whose parents are more genetically differentiated also having larger differences between their historic midparent precipitation and conditions during the experiment. We think it is unlikely that the apparent positive effects of genetic differentiation are actually driven by precipitation differences, because we would expect high precipitation differences to negatively affect fitness. The overall weak or absent effects of temperature and precipitation differences in these

338 models may be due to a narrower range of variation in midparent climate for the between-population
339 crosses relative to the within-population crosses (Figure S2).

Discussion

We conducted a common garden experiment at the northern range margin of *Clarkia pulchella*
to examine how the effects of gene flow on peripheral populations vary with climatic and genetic
differentiation between focal and source populations. We examined predictors of fitness of within-
population crosses, in which both parents originated from the same source population, as well as
between-population crosses, in which one parent was local to the common gardens and the other was
from another population from across the northern half of the range of *C. pulchella*. In our experi-
ment, provenances of *C. pulchella* from climates that most closely matched the conditions during
the experiment performed best, even better than local populations. Populations of *C. pulchella* at
the northern range margin benefited from gene flow from warm source locations during the warm
year of our experiment. Gene flow also conferred some benefits independent of climate, as evidenced
by the positive effect of gene flow on later lifestages when controlling for temperature differences
and the positive effect of increasing genetic differentiation between the parental populations.

Climate of origin predicts performance

Populations of *Clarkia pulchella* are adapted to their historic climate regimes, a pattern consistent
with findings in many other species (Anderson et al., 2015; Wilczek et al., 2014). When grown in
common sites, the performance of individuals was determined by the degree to which conditions
during the experiment deviated from historic temperature and precipitation averages of each prove-
nance (Figure 2). Because of this local adaptation to climate, gene flow from sites that deviate
from local conditions (in our experiment, sites that are cooler than the focal populations) had the
potential to disrupt local adaptation, as indicated by the somewhat negative effects of midparent
temperature differences on between-population plants (Figure 4, Table S5). Although simulated
gene flow from divergent environments had the potential to reduce fitness in this experiment, it is
unlikely that gene flow occurs at a high enough rate among natural populations to swamp local

adaptation. However, our results highlight that gene flow and dispersal need not be from populations that are geographically distant (or from the center of the range) to be climatically divergent from historic or current conditions. Rather, two of the most temperature-mismatched populations used in the experiment are from sites nearest to our common gardens (populations D1 and D3; Figure 1).

Under climate change, local adaptation to historic climate regimes may generate local maladaptation in field trials. We see this in our results, where populations from warmer locations performed best in our gardens (Figure 2, Table S3), and gene flow from warmer locations had positive effects on some lifestages (Figure 4, Table S5). This lagging adaptation to climate has been documented in other recent common garden studies. In a reciprocal transplant experiment of a long-lived sedge, McGraw et al. (2015) found that populations were displaced 140 km south of their optimal climate conditions. Wilczek et al. (2014) found that local genotypes of *Arabidopsis thaliana* from across Europe were consistently outperformed in common gardens by accessions from historically warmer locations. These results indicate that dispersal and gene flow are important processes promoting range stasis as climate warms, as they allow alleles that are beneficial in warm environments to spread from historically warm populations to recently warming sites. However, climate is multivariate, and as the climate changes it may generate combinations of conditions that no population has historically experienced (Williams and Jackson, 2007; Mahony et al., 2017). The particular combination of hot and dry conditions in our common gardens was unlike any of our populations' historic temperature and precipitation combinations (Figure 1), though they are similar to normals from some populations not included in our experiment, primarily from the southern half of the species range (data not shown). While precipitation conditions were similar to those historically experienced by the focal populations, temperature conditions favored another set of populations. Whether the optimal traits for different climatic axes are antagonistic and whether segregation and recombination will allow adaptation to novel climates are important considerations in predicting climate change responses.

Gene flow confers benefits independent of climate

We saw some additional positive effects of gene flow once the effects of climate are controlled for (Figure 3, Table S4). These positive effects may be the result of increased heterozygosity when parental plants come from two different populations; this inference is supported by the positive effect of genetic differentiation between parental populations on performance (Figure 4, Table S5). These results are also generally consistent with previous work in which experimental populations of *Clarkia pulchella* with higher genetic effective population sizes had lower extinction probabilities (Newman and Pilson, 1997). An interesting direction for future models of swamping gene flow along environmental gradients might be to explore whether incorporating heterosis-dependent increases in the effective migration rate (Ingvarsson and Whitlock, 2000) alters predictions (this could be done with various dispersal distances, under scenarios of various magnitudes of isolation-by-distance). However, an important question is whether the benefits of reduced homozygosity (or increased heterozygosity) are transient effects among F1s, how long they would persist in future generations if our between-population plants backcrossed into the focal populations, and whether these benefits may be counteracted by outbreeding depression as recombination disrupts co-adapted gene complexes. The answers to these questions are likely to depend on many factors, including the genetic architecture of local adaptation and population size (Willi et al., 2007), but fitness declines in subsequent generations are not uncommon after between-population crosses (Fenster and Galloway, 2000; Johansen-Morris and Latta, 2006). Novel environments may alter the costs and benefits of outbreeding: increases in variation among individuals might help populations adapt, despite temporary decreases in mean fitness due to outbreeding depression.

During this study, the effects of being well-matched to the experimental conditions seemed to dominate over potential benefits of being from a local population (for example, the benefits of being adapted to local soil conditions or herbivores). This inference is supported by the fact that lifetime fitness of local populations did not differ from that of foreign populations even once climate differences were controlled for (results not shown) though our experiment was not especially well-suited to test this because we have only two local populations.

Limited inference about population persistence

Our ability to make inferences from our results about the longer-term effects of gene flow on the persistence and adaptive potential of range edge populations is limited. While it seems clear that gene flow from warm sites is likely to accelerate adaptation to warming conditions, we do not know whether these populations were historically limited by adaptation, and whether the additional genetic variation introduced by gene flow would permit better adaptation to local conditions and range expansion on an evolutionary time scale. These types of questions are difficult to test in field systems (but see Etterson and Shaw 2001), but inferences can be made by examining genetic variance of wild populations in the lab (Kellermann et al., 2006; Hoffmann et al., 2003). The development of experimental evolution systems to test equilibrial range dynamics is an exciting avenue for future work—this would be a natural extension of recent studies of range expansion dynamics using experimental evolution in the lab (Ochocki and Miller, 2017; Williams et al., 2016).

It is also important to note that all populations in our experiment had reproductive rates that were well above replacement (one seed produced per seed planted, see y-axis on Figure 2A, Figure 4A), so we have no evidence that gene flow has the potential to drive populations extinct, or to rescue them from extinction. The high lifetime fitness we observed during our experiment could be due to several factors. First, perhaps warm conditions over the entire season are favorable for all sources, but are more favorable for warm-adapted populations. Alternatively, we could have increased fitness by limiting antagonistic biotic interactions. Finally, and perhaps most plausibly, our plot placement may have upwardly biased our germination and reproductive estimates. We placed plots in patches that appeared favorable to *C. pulchella*, but naturally dispersing seeds are likely to land in a mix of favorable and unfavorable patches. We do not know whether any of these factors might interact with provenance, in which case they might change the relative performances of populations in our experiment.

Conclusions and future directions

On heterogeneous landscapes, the effects of gene flow are likely to depend strongly on climate of origin, which may be decoupled from geography. Although gene flow from divergent climates

may result in reduced fitness, gene flow can also confer benefits to range edge populations via both preadapted alleles and reduced homozygosity. As a result, moderate rates of gene flow likely represent a net benefit to edge populations, particularly as the environment is changing. This study highlights the challenges of testing hypotheses about equilibrial range limits in the field, where climate change is a persistent reality. Even if populations were once locally adapted, they are likely no longer at equilibrium with climate. The signal of climate anomalies disrupting local adaptation can be detected in published literature to date (Bontrager et al., in prep.). In light of this, future studies of local adaptation at range edges should be designed in such a way that the results will be informative even in non-equilibrial conditions.

Author contributions

MB and ALA conceived of the project. MB performed greenhouse crosses, garden installation, and data collection. MB analysed the data in consultation with ALA. MB wrote the manuscript with advice and edits from ALA.

Data deposition

All data and code required to recreate these analyses is publicly available at github.com/megan bontrager/clarkia-pulchella-transplant. Upon publication, all data and code will be archived in an appropriate repository (e.g., Dryad).

Acknowledgements

We would like to thank M. Osmond, L. Bontrager, C. Leven, D. Gamble, A. Porter, A. Wilkinson, J. Chan, T. Mitchell, and P. Chen for their help in the field. A. Wilkinson and M. Zink Yi provided plant care in the greenhouse. B. Gass and D. Holtz also helped with plant care. E. Fitz and D. Gamble assisted with greenhouse pollinations and fruit collection. A. Porter, J. Ono, and J. Chan assisted with preparing the seeds for planting. British Columbia Provincial Parks permitted the garden installation and monitoring. M. Whitlock, J. Whitton, S. Aitken and T. Usui provided

468 helpful comments on an earlier draft of this manuscript. This work was supported by grants from
469 the Botanical Society of America and the Washington Native Plant Society. MB was supported by
470 a University of British Columbia Four-year Fellowship. The authors declare no conflicts of interest.

References

Aitken, S. N. and M. C. Whitlock (2013). Assisted gene flow to facilitate local adaptation to climate
change. *Annual Review of Ecology, Evolution, and Systematics 44.* 6

Anderson, J. T., N. Perera, B. Chowdhury, and T. Mitchell-Olds (2015). Microgeographic pat-
terns of genetic divergence and adaptation across environmental gradients in *Boechera stricta*
(Brassicaceae). *The American Naturalist 186*(S1), S60–S73. 15

Antonovics, J. (1976). The nature of limits to natural selection. *Annals of the Missouri Botanical*
Garden, 224–247. 4

Bontrager, M. and A. L. Angert (2018). Genetic differentiation is determined by geographic distance
in clarkia pulchella. *bioRxiv*, 374454. 10

Bontrager, M., C. D. Muir, and A. Angert (2018). Geographic and climatic drivers of reproductive
assurance in clarkia pulchella. *bioRxiv*, 372375. 7

Bridle, J. R. and T. H. Vines (2007). Limits to evolution at range margins: when and why does
adaptation fail? *Trends in Ecology & Evolution 22*(3), 140–147. 4

Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielsen, H. J.
Skaug, M. Machler, and B. M. Bolker (2017). glmmTMB balances speed and flexibility among
packages for zero-inflated generalized linear mixed modeling. *The R Journal 9*(2), 378–400. 10

Eckert, C., K. Samis, and S. Lougheed (2008). Genetic variation across species' geographical ranges:
the central–marginal hypothesis and beyond. *Molecular Ecology 17*(5), 1170–1188. 4

Emery, N. C. (2009). Ecological limits and fitness consequences of cross-gradient pollen movement
in *Lasthenia fremontii*. *The American Naturalist 174*(2), 221–235. 5

Etterson, J. R. and R. G. Shaw (2001). Constraint to adaptive evolution in response to global warming. *Science 294* (5540), 151–154. 18

Fedorka, K., W. Winterhalter, K. Shaw, W. Brogan, and T. Mousseau (2012). The role of gene flow asymmetry along an environmental gradient in constraining local adaptation and range expansion. *Journal of Evolutionary Biology 25* (8), 1676–1685. 5

Fenster, C. B. and L. F. Galloway (2000). Population differentiation in an annual legume: genetic architecture. *Evolution 54* (4), 1157–1172. 17

Goudet, J. and T. Jombart (2015). hierfstat: Estimation and tests of hierarchical F-statistics. R package version 0.04-22. 10

Gould, B., D. A. Moeller, V. M. Eckhart, P. Tiffin, E. Fabio, and M. A. Geber (2014). Local adaptation and range boundary formation in response to complex environmental gradients across the geographical range of *Clarkia xantiana* ssp. *xantiana*. *Journal of Ecology 102* (1), 95–107. 5

Hoffmann, A., R. Hallas, J. Dean, and M. Schiffer (2003). Low potential for climatic stress adaptation in a rainforest *Drosophila* species. *Science 301* (5629), 100–102. 18

Holliday, J. A., H. Suren, and S. N. Aitken (2012). Divergent selection and heterogeneous migration rates across the range of Sitka spruce (*Picea sitchensis*). *Proceedings of the Royal Society of London B: Biological Sciences 279* (1734), 1675–1683. 5

Ingvarsson, P. K. and M. C. Whitlock (2000). Heterosis increases the effective migration rate. *Proceedings of the Royal Society of London B: Biological Sciences 267* (1450), 1321–1326. 6, 17

Johansen-Morris, A. and R. G. Latta (2006). Fitness consequences of hybridization between ecotypes of *Avena barbata*: hybrid breakdown, hybrid vigor, and transgressive segregation. *Evolution 60* (8), 1585–1595. 17

Kellermann, V. M., B. van Heerwaarden, A. A. Hoffmann, and C. M. Sgrò (2006). Very low additive genetic variance and evolutionary potential in multiple populations of two rainforest *Drosophila* species. *Evolution 60* (5), 1104–1108. 18

Kirkpatrick, M. and N. H. Barton (1997). Evolution of a species' range. *The American Naturalist 150*(1), 1–23. 4

Lee-Yaw, J. A., H. M. Kharouba, M. Bontrager, C. Mahony, A. M. Csergő, A. M. Noreen, Q. Li, R. Schuster, and A. L. Angert (2016). A synthesis of transplant experiments and ecological niche models suggests that range limits are often niche limits. *Ecology Letters 19*(6), 710–722. 4

Lenormand, T. (2002). Gene flow and the limits to natural selection. *Trends in Ecology & Evolution 17*(4), 183–189. 4

Lüdecke, D. (2018). *ggeffects: Create tidy data frames of marginal effects for 'ggplot' from model outputs*. R package version 0.3.3. 11

MacSwain, J. W., P. H. Raven, and R. W. Thorp (1973). Comparative behavior of bees and Onagraceae: 4. *Clarkia* bees of the Western United States. *University of California Publications in Entomology 70*. 7

Mahony, C. R., A. J. Cannon, T. Wang, and S. N. Aitken (2017). A closer look at novel climates: new methods and insights at continental to landscape scales. *Global Change Biology 23*(9), 3934–3955. 16

McGraw, J. B., J. B. Turner, S. Souther, C. C. Bennington, M. C. Vavrek, G. R. Shaver, and N. Fetcher (2015). Northward displacement of optimal climate conditions for ecotypes of *Eriophorum vaginatum* L. across a latitudinal gradient in Alaska. *Global Change Biology 21*(10), 3827–3835. 16

Moore, J.-S. and A. P. Hendry (2009). Can gene flow have negative demographic consequences? Mixed evidence from stream threespine stickleback. *Philosophical Transactions of the Royal Society B: Biological Sciences 364*(1523), 1533–1542. 5

Newman, D. and D. Pilson (1997). Increased probability of extinction due to decreased genetic effective population size: experimental populations of *Clarkia pulchella*. *Evolution 51*(2), 354–362. 17

Ochocki, B. M. and T. E. Miller (2017). Rapid evolution of dispersal ability makes biological invasions faster and more variable. *Nature Communications 8*, 14315. 18

Paul, J. R., S. N. Sheth, and A. L. Angert (2011). Quantifying the impact of gene flow on phenotype-environment mismatch: a demonstration with the scarlet monkeyflower *Mimulus cardinalis*. *The American Naturalist 178*(S1), S62–S79. 5

Pironon, S., G. Papuga, J. Villellas, A. L. Angert, M. B. García, and J. D. Thompson (2017). Geographic variation in genetic and demographic performance: new insights from an old biogeographical paradigm. *Biological Reviews 92*(4), 1877–1909. 4

PRISM Climate Group (2017). Oregon state university. Accessed: 2017-09-30. 9

Pujol, B. and J. R. Pannell (2008). Reduced responses to selection after species range expansion. *Science 321*(5885), 96–96. 4

R Core Team (2017). *R: a language and environment for statistical computing, version 3.4.4.* Vienna, Austria: R Foundation for Statistical Computing. 12

Sagarin, R. D. and S. D. Gaines (2002). The 'abundant centre' distribution: to what extent is it a biogeographical rule? *Ecology Letters 5*(1), 137–147. 6

Samis, K. E., A. López-Villalobos, and C. G. Eckert (2016). Strong genetic differentiation but not local adaptation toward the range limit of a coastal dune plant. *Evolution 70*(11), 2520–2536. 5

Sexton, J. P., S. B. Hangartner, and A. A. Hoffmann (2014). Genetic isolation by environment or distance: which pattern of gene flow is most common? *Evolution 68*(1), 1–15. 6

Sexton, J. P., P. J. McIntyre, A. L. Angert, and K. J. Rice (2009). Evolution and ecology of species range limits. *Annual Review of Ecology, Evolution, and Systematics 40*. 4

Sexton, J. P., S. Y. Strauss, and K. J. Rice (2011). Gene flow increases fitness at the warm edge of a species range. *Proceedings of the National Academy of Sciences 108*(28), 11704–11709. 6

Vucetich, J. A. and T. A. Waite (2003). Spatial patterns of demography and genetic processes
across the species' range: null hypotheses for landscape conservation genetics. *Conservation*
Genetics 4(5), 639–645. 4

Weir, B. S. and C. C. Cockerham (1984). Estimating F-statistics for the analysis of population
structure. *Evolution 38*(6), 1358–1370. 10

Wilczek, A. M., M. D. Cooper, T. M. Korves, and J. Schmitt (2014). Lagging adaptation to warming
climate in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences 111*(22), 7906–
7913. 15, 16

Willi, Y., M. Van Kleunen, S. Dietrich, and M. Fischer (2007). Genetic rescue persists beyond
first-generation outbreeding in small populations of a rare plant. *Proceedings of the Royal Society*
of London B: Biological Sciences 274(1623), 2357–2364. 17

Williams, J. L., B. E. Kendall, and J. M. Levine (2016). Rapid evolution accelerates plant population
spread in fragmented experimental landscapes. *Science 353*(6298), 482–485. 18

Williams, J. W. and S. T. Jackson (2007). Novel climates, no-analog communities, and ecological
surprises. *Frontiers in Ecology and the Environment 5*(9), 475–482. 16

Yeaman, S. and A. Jarvis (2006). Regional heterogeneity and gene flow maintain variance in a
quantitative trait within populations of lodgepole pine. *Proceedings of the Royal Society of London*
B: Biological Sciences 273(1594), 1587–1593. 5

Figure 1 Geographic locations and climate averages of populations used in this experiment. **(A)** Populations span the northern half of the geographic range of *Clarkia pulchella* (indicated by the dashed line). Focal population sites (where common gardens were installed) are indicated by "F" and donor population sites are indicated by "D". Identifying codes for each population correspond to the Map ID column in Table S1. **(B)** Temperature and precipitation conditions of common gardens and source provenances. Colored circles represent the 1951-1980 average temperature and precipitation in the home site of each population. Open circles represent weather conditions during the experiment. Focal populations are historically intermediate relative to donor populations in average historic temperature (x-axis), but are the from the driest sites of any population used in the experiment (y-axis). Conditions in common gardens during the experiment were hot relative to normal conditions at those sites, and hot and dry relative to average conditions of all populations in the experiment.

Figure 2 Effects of absolute temperature difference (September-July; T_{diff}) and absolute precipitation difference (April-July; P_{diff}) on performance of *Clarkia pulchella* in common gardens. These analyses include within-population plants only (no gene flow). **(A)** Lifetime fitness declines with increasing differences in temperature between the historic average of the source population and the experimental conditions in the transplant gardens. The shaded area represents the 95% confidence interval of the model estimate conditioned on fixed effects only. Though these temperature differences are expressed as absolute, almost all populations were from sources that are historically cooler than the transplant sites were during the experiment. Points are raw averages for each source population in each garden, colored by the distance between the source population and the transplant garden. Triangles are focal populations in their home sites. **(B)** Regression estimates and standard errors from the zero-inflation part of a model of lifetime fitness. **(C)** Regression estimates and standard errors from the conditional part of a model of lifetime fitness. **(D)** Schematic of effects of absolute temperature differences (T_{diff}) and absolute precipitation difference (P_{diff}) on component lifestages of *Clarkia pulchella*. Directionality of effects is illustrated with "-"; in these analyses all significant effects were negative. Predictors in boxes are significant ($P < 0.05$). Size in the previous lifestage is not shown here, but has a significant positive effect on overwinter and reproductive lifestages. This summarizes the significant results of separate models for each lifestage; full statistical results of these tests are in Table S3.

Figure 3 Effects of gene flow (differences between between-population and within-population crosses) on performance of *Clarkia pulchella*, accounting for midparent temperature and precipitation. **(A)** Regression estimates and standard errors from the zero-inflation part of a model of lifetime fitness. **(B)** Regression estimates and standard errors from the conditional part of a model of lifetime fitness. **(C)** Effects of gene flow (GF), absolute midparent temperature differences (T_{diff}), and absolute midparent precipitation differences (P_{diff}) on component lifestages of *Clarkia pulchella*. Directionality of effects is illustrated with "+" and "-". Marginally significant parameters ($0.05 < P < 0.10$) are shown in boxes with dashed margins, predictors in solid boxes are significant ($P < 0.05$). Size in the previous lifestage is not shown here, but has a significant positive effect on overwinter and reproductive lifestages. This summarizes the significant results of separate models for each lifestage; full statistical results of these tests are in Table S4.

Figure 4 Effects of genetic differentiation between parental populations, as well as midparent temperature and precipitation on performance of *Clarkia pulchella*. **(A)** Among between-population crosses, increased genetic divergence between parental populations had a positive effect on lifetime fitness. Points are raw averages for each gene flow source in each garden, colored by the distance between the source population and the garden (the home of the recipient population). **(B)** Regression estimates and standard errors of genetic differentiation (F_{ST}) and absolute midparent temperature differences (T_{diff}) on the probability of producing seeds. **(C)** Effects of genetic differentiation, temperature differences, and absolute precipitation differences (P_{diff}) on conditional seed production. **(D)** Effects of T_{diff} and F_{ST} on component lifestages of *Clarkia pulchella*. Precipitation differences were not significant when tested for component lifestages. Directionality of effects is illustrated with "+" and "-". Marginally significant parameters ($0.05 < P < 0.10$) are shown in boxes with dashed margins, predictors in solid boxes are significant ($P < 0.05$). Size in the previous lifestage is not shown here, but has a significant positive effect on overwinter and reproductive lifestages. * indicates predictors that are only significant in separate models, not in full models with all predictors. Complete statistical results of these tests are in Table S5 and Table S6.

 # Supplementary methods

 ## Greenhouse conditions and crossing design

Seeds were planted in the greenhouse 9-11 December, 2014 in conetainers (Stuewe and Sons, Tan-
gent, Oregon, USA) filled with Sunshine Mix No. 4 (Sun Gro Horticulture, Agawam, MA, USA).
For each of 22 maternal families per population, 3-5 seeds were planted on the soil surface in each
pot. For families from each of the two focal populations, three replicate pots were planted per
family because larger quantities of flowers would be needed from these families; other populations
were represented by one replicate per family. Pots were arranged into randomized blocks, with each
block containing one family from each population (one pot from each donor population and three
replicate pots from each of the two focal populations). The soil was kept moist until germination,
then plants were hand watered every 1-3 days as needed to prevent wilting. After germination,
plants were thinned randomly to one per cone and pumice was added to the soil surface to prevent
fungal growth. Plants began to flower in March 2015. Plants were bagged to prevent unintentional
pollination, and flowers were emasculated upon opening to prevent self-pollination. For the crosses,
20 of the 22 blocks were used, the other two were maintained in the same growing conditions to
provide alternate plants in case of mortality or sterility. We performed as many crosses as possible
using a single focal plant, but if flower production was too low on that plant we also used one of
the replicate focal plants from the same family. Most crosses had to be performed 2-3 times to
obtain adequate numbers of seeds for the experiment. Some crosses could not be performed due
to mortality, sterility, or limited flower production. As ripening progressed, the ends of fruits were
taped shut to prevent seed loss. Upon ripening, fruits were collected and stored in coin envelopes
in the lab. Crosses were performed March-May 2015 and and we collected fruits March-June 2015.

Transplant installation

Seeds attached to toothpicks were planted in the ground 18-21 September 2015. Plots were prepared
by removing litter, large rocks, and dried remains of herbaceous perennial plants. The ground surface
was minimally leveled to allow for placement of planting grids that aided in consistently spacing
the plants. Each toothpick was inserted into the ground gently so that seeds were not dislodged or

damaged until seeds were ~3 mm below the soil surface. Toothpicks were inserted at 5 cm spacing
into ~1 m by 2 m blocks. Block shape was varied to accommodate rocks and shrubs surrounding
the planting area. After planting, each block was protected with 20 cm high hardware cloth cages
supported by rebar. These cages were intended to prevent trampling by larger animals but did
not prevent entry of rodents and other small animals. The area surrounding the plots at each
site was sprayed with deer repellent several times during the course of the experiment. To ensure
germination, plots were watered at a rate of ~10 L per plot 27-29 October 2015, though at that
time most seeds that were checked already had radicles emerging. In May 2016 cattle fencing was
put around the plots at the Blue Lake site before cattle were released into the area for grazing; this
fencing succeeded in keeping the cows off the plots. No cattle were present at the Rock Creek site.

Details of monitoring and measuring

Our transplant gardens were installed in areas where *C. pulchella* occurs naturally, and because of
this we wanted to evaluate how frequently we might have mistaken naturally occurring plants for
experimental plants. During germination surveys we censused one more germinant than the number
of seeds that we planted at 23 out of 16,680 grid points (0.14%). This gives an estimate of the
minimum rate at which naturally occurring seeds were indistinguishable from our planted seedlings.
So, while it is probable that some naturally occurring plants were mistaken for experimental plants,
we consider the frequency of possible misidentification to be acceptably low.

During winter, some plots were affected by frost-heave and seedlings were uprooted from their
planting locations when their toothpick was forced out of the ground (1901/16680 grid points,
11.4%). In lightly affected areas, toothpicks and seedlings were gently settled back into the soil.
In more heavily affected areas, individual identity could no longer be determined confidently and
individuals were excluded from further measurements and analyses (95/16680 grid points, 0.57%).

Censuses of reproduction began on 2 June 2016. On 12-13 June 2016 we censused spring survival
of all plants. Once flowering began, we placed bridal-veil nets over the hardware cloth on each plot
to prevent pollen escape into local populations. In June we censused each plot every 2-3 days. We
recorded the date of first flowering of each plant at this temporal resolution. During each census,

the immature ovary length of each new flower was recorded to be used as a proxy for maximum seed

set. Flowers were marked as they were measured with a permanent marker and a running flower

count was kept for each plant to avoid double-counting as flowers senesced. We continued these

assessments as flowering slowed in July, but reduced the census interval to once a week. Damage

to plants, such as rodent activity or herbivory, was noted during monitoring. Any plants with

uncertain identities (due to frost damage as mentioned above, being far from their toothpick, or the

toothpick disappearing; n = 201/16680 toothpicks, 1.2%) were excluded from all analyses. Plants

that were killed by gophers, browsers, or galling insects were excluded from analyses that involved

lifestages downstream from these events (n = 525/16479 plants, 3.2%) because we do not think that

this mortality is related to population origin but rather to block-specific factors.

Ovary length as a fitness proxy

Pollinations were performed on a subset of plants to calibrate a conversion from immature ovary

lengths to seed production. On 596 flowers (mean = 29.8 per plot, range = 0–126) stigmas were

dusted with an ample pollen load using all four anthers from another plant in the plot. These

flowers were marked with strings around the pedicles and fruits were collected when ripe. Seeds in

each of these pollinated fruits were later counted in the lab. Total seed production per individual

was estimated by multiplying the total ovary length of each plant by the average number of seeds

per mm of immature ovary, as determined from the pollinated fruits. This resulted in an estimate

of 4.75 seeds per mm based on a linear regression of number of seeds predicted by ovary length

with the intercept set to 0 ($R_0^2 = 0.87$). We pollinated only a maximum of one flower per plant, so

these may be overestimates because they do not account for potential resource limitation of seed

set. However, we checked whether variation in seeds per mm of fruit was associated with individual

fitness (the overall fruit production per individual) or block quality (estimated based on the average

fruit production of a block), and we could not attribute variation in seeds per mm of fruit to either

of these factors. Therefore, while our conversion from fruit length to seeds may not be exact, we

do not expect it to be systematically biased.

 # Supplementary tables and figures

Table S1 Geographic information for the populations of *Clarkia pulchella* used in this experiment.

Population name	Map ID	Type	Latitude	Longitude	Elevation (m)
Blue Lake	F1	Focal	49.05	-119.56	842
Johnstone Creek	F2	Focal	49.04	-119.05	866
Border	D1	Donor	48.98	-118.99	1211
Day Creek Road	D2	Donor	48.94	-118.51	911
Bodie Mountain	D3	Donor	48.83	-118.83	1603
Boulder Creek Road	D4	Donor	48.76	-118.33	1115
Aeneas Valley	D5	Donor	48.54	-118.91	1126
Henry Creek Road	D6	Donor	47.45	-114.77	1103
Heyburn State Park	D7	Donor	47.34	-116.79	801
McCrosky State Park	D8	Donor	47.09	-116.98	1186
Graves Creek Road	D9	Donor	46.80	-114.41	1201
Bitterroot	D10	Donor	46.54	-113.89	1424
Abel's Ridge	D11	Donor	46.28	-117.60	1457
Tucannon	D12	Donor	46.24	-117.74	1022
Pendleton	D13	Donor	45.74	-118.25	649

Table S2 Results of generalized linear mixed effects models for the effect of local vs. foreign origin on performance of *Clarkia pulchella* in common gardens. There are no significant differences between populations of local vs. foreign origin in fitness components or lifetime fitness. Size during the previous census (November for overwinter survival and size, March for fruit counts and estimated seed production) is always a significant predictor of performance in subsequent lifestages.

Response	Foreign vs. local origin			Size during previous census		
	Estimate	SE	*P*-value	Estimate	SE	*P*-value
Germination	-0.034	0.088	0.702	-	-	-
Size after germination	0.027	0.073	0.708	-	-	-
Overwinter survival	0.176	0.126	0.161	**0.377**	**0.033**	**< 0.001**
Size after winter	-0.068	0.182	0.708	**0.750**	**0.042**	**< 0.001**
Fruit count	0.089	0.146	0.544	**0.544**	**0.041**	**< 0.001**
Seed production	-0.005	0.128	0.966	**0.388**	**0.032**	**< 0.001**
Lifetime fitness - zero inflation	-0.032	0.123	0.798	-	-	-
Lifetime fitness - conditional	-0.093	0.128	0.469	-	-	-

Table S3 Results of generalized linear mixed effects models of the effects of absolute precipitation and temperature differences on component lifestages of *Clarkia pulchella*. Temperature and precipitation differences refer to absolute differences between the historic conditions that a population experienced and the conditions in the common gardens during the experiment. These differences were calculated using climate data from only the months of that census period (i.e., September-November for germination and size after germination, December-March for overwinter survival and size after winter, and April-July for fruit counts and seed production). Analyses were conducted using only plants surviving the previous census window. Whenever applicable, size in the previous census was included as a covariate to account for differences accumulated during earlier lifestages. Significant parameters are indicated with bold text.

| Response | Absolute temperature difference | | | Absolute precipitation difference | | | Size in previous census | | |
	Estimate	SE	*P*-value	Estimate	SE	*P*-value	Estimate	SE	*P*-value
Germination	**-0.094**	**0.035**	**0.007**	-	-	-	-	-	-
Size after germination	-0.027	0.026	0.293	-	-	-	-	-	-
Overwinter survival	**-0.078**	**0.032**	**0.015**	-	-	-	**0.366**	**0.033**	**< 0.001**
Size after winter	**-0.518**	**0.095**	**< 0.001**	-	-	-	**0.748**	**0.042**	**< 0.001**
Fruit count	-0.114	0.071	0.108	-0.091	0.058	0.119	**0.543**	**0.041**	**< 0.001**
Seed production	-0.055	0.047	0.249	**-0.110**	**0.044**	**0.011**	**0.389**	**0.033**	**< 0.001**

Table S4 Results of generalized linear mixed effects models of the effects of being a within-population cross vs. a between-population cross, while accounting for effects of absolute precipitation and temperature differences. Temperature and precipitation differences refer to absolute differences between the average historic conditions of an individual's parental populations and the conditions in the common gardens during the experiment. Positive estimates of the effects of between-population vs. within-populations indicate that having parents from two different populations ("gene flow") is beneficial. **(A)** Effects on lifetime fitness of *Clarkia pulchella.* **(B)** Effects on lifetime fitness when midparent precipitation differences are not included in the model. **(C)** Effects on component lifestages. Climate differences were calculated using climate data from only the months of that census period (i.e., September-November for germination and size after germination, December-March for overwinter survival and size after winter, and April-July for fruit counts and seed production). Analyses were conducted using only plants surviving the previous census window. Whenever applicable, size in the previous census was included as a covariate to account for differences accumulated during earlier lifestages. Significant parameters are indicated with bold text.

Response	Between-populations vs. within-populations			Absolute midparent temperature difference			Absolute midparent precipitation difference			Size during previous census		
	Estimate	SE	*P*-value	Estimate	SE	*P*-value	Estimate	SE	*P*-value	Estimate	SE	*P*-value
A. Lifetime fitness												
Lifetime fitness - zero inflation	0.008	0.047	0.860	**-0.240**	**0.023**	**< 0.001**	-	-	-	-	-	-
Lifetime fitness - conditional	0.047	0.049	0.345	**-0.085**	**0.032**	**0.007**	**-0.072**	**0.036**	**0.045**	-	-	-
B. Lifetime fitness without precipitation in model												
Lifetime fitness - zero inflation	0.008	0.047	0.860	**-0.240**	**0.023**	**< 0.001**	-	-	-	-	-	-
Lifetime fitness - conditional	**-0.114**	**0.036**	**0.002**	**-0.059**	**0.029**	**0.041**	-	-	-	-	-	-
C. Component lifestages												
Germination	-0.035	0.078	0.653	**-0.088**	**0.029**	**0.003**	-	-	-	-	-	-
Size after germination	-0.033	0.041	0.421	-0.028	0.015	0.058	-	-	-	-	-	-
Overwinter survival	-0.012	0.047	0.794	**-0.058**	**0.022**	**0.009**	-	-	-	**0.336**	**0.023**	**< 0.001**
Size after winter	-0.052	0.127	0.678	**-0.274**	**0.068**	**< 0.001**	-	-	-	**0.721**	**0.031**	**< 0.001**
Fruit count	**-0.125**	**0.055**	**0.021**	**-0.081**	**0.032**	**0.011**	**-0.092**	**0.034**	**0.007**	**0.493**	**0.030**	**< 0.001**
Seed production	0.092	0.047	0.052	-0.030	0.028	0.285	**-0.092**	**0.030**	**0.002**	**0.357**	**0.024**	**< 0.001**

Table S5 Results of generalized linear mixed effects models separately testing the effects of **(A)** genetic differentiation, **(B)** absolute midparent temperature differences, and **(C)** absolute midparent precipitation differences on performance of *Clarkia pulchella* in common gardens. Absolute midparent temperature and precipitation differences refer to absolute differences between the conditions in the common gardens during the experiment and the average historic conditions of an individual's parental populations. These analyses were performed using between-population crosses only, that is, every plant has one parent from a focal population and one parent from a donor population. For analyses of lifetime fitness, temperature differences were calculated using the duration of the experiment and precipitation differences were calculated using April-July values. Precipitation differences are only included as an effect in the conditional part of the model of lifetime fitness because precipitation effects are expected to manifest at later lifestages. For analyses of component lifestages, climate differences were calculated using climate data from only the months of that census period (i.e., September-November for germination and size after germination, December-March for overwinter survival and size after winter, and April-July for fruit counts and seed production). Component lifestage analyses were conducted using only plants surviving the previous census window. Whenever applicable, size in the previous census was included as a covariate to account for differences accumulated during earlier lifestages. Significant parameters are indicated with bold text.

A. F_{ST}

Response	F_{ST} Estimate	SE	P-value	Size during previous census Estimate	SE	P-value
Lifetime fitness - zero inflation	**0.086**	**0.028**	**0.002**	-	-	-
Lifetime fitness - conditional	**0.080**	**0.031**	**0.010**	-	-	-
Germination	**0.124**	**0.059**	**0.034**	-	-	-
Size after germination	0.014	0.014	0.306	-	-	-
Overwinter survival	0.054	0.035	0.124	**0.336**	**0.037**	**< 0.001**
Size after winter	**0.104**	**0.050**	**0.035**	**0.743**	**0.044**	**< 0.001**
Fruit count	**0.067**	**0.031**	**0.028**	**0.451**	**0.044**	**< 0.001**
Seed production	**0.081**	**0.029**	**0.005**	**0.334**	**0.035**	**< 0.001**

B. Absolute midparent temperature differences

Response	Temperature difference Estimate	SE	P-value	Size during previous census Estimate	SE	P-value
Lifetime fitness - zero inflation	**-0.148**	**0.046**	**0.001**	-	-	-
Lifetime fitness - conditional	-0.090	0.056	0.105	-	-	-
Germination	-0.088	0.070	0.212	-	-	-
Size after germination	-0.029	0.017	0.096	-	-	-
Overwinter survival	-0.054	0.049	0.275	**0.335**	**0.037**	**< 0.001**
Size after winter	-0.132	0.070	0.060	**0.743**	**0.044**	**< 0.001**
Fruit count	-0.121	0.063	0.057	**0.457**	**0.044**	**< 0.001**
Seed production	-0.071	0.066	0.285	**0.336**	**0.035**	**< 0.001**

C. Absolute midparent precipitation differences

Response	Precipitation difference Estimate	SE	P-value	Size during previous census Estimate	SE	P-value
Lifetime fitness - conditional	0.123	0.078	0.113	-	-	-
Fruit count	0.047	0.083	0.574	**0.452**	**0.044**	**< 0.001**
Seed production	0.099	0.076	0.192	**0.334**	**0.035**	**< 0.001**

Table S6 Results of generalized linear mixed effects models of the effects of genetic differentiation between parental populations on performance of *Clarkia pulchella* in common gardens. Effects of absolute precipitation and temperature differences are also included in these models. Temperature and precipitation differences refer to the absolute midparent differences, i.e., the absolute differences between the conditions in the common gardens during the experiment and the average historic conditions of an individual's parental populations. These analyses were performed using between-population crosses only, that is, every plant has one parent from a focal population and one parent from a donor population. **(A)** Effects on lifetime fitness. **(B)** Effects on component lifestages. Climate differences were calculated using climate data from only the months of that census period (i.e., September-November for germination and size after germination, December-March for overwinter survival and size after winter, and April-July for fruit counts and seed production). Analyses were conducted using only plants surviving the previous census window. Whenever applicable, size in the previous census was included as a covariate to account for differences accumulated during earlier lifestages. Significant parameters are indicated with bold text.

Response	F_{ST} Estimate	SE	*P*-value	Temperature difference Estimate	SE	*P*-value	Precipitation difference Estimate	SE	*P*-value	Size during previous census Estimate	SE	*P*-value
A. Lifetime fitness												
Lifetime fitness - zero inflation	**0.064**	**0.030**	**0.030**	**-0.117**	**0.048**	**0.014**	-	-	-	-	-	-
Lifetime fitness - conditional	0.059	0.037	0.108	-0.031	0.055	0.574	0.053	0.082	0.520	-	-	-
B. Component lifestages												
Germination	0.111	0.062	0.073	-0.021	0.080	0.798	-	-	-	-	-	-
Size after germination	0.005	0.016	0.770	-0.026	0.020	0.179	-	-	-	-	-	-
Overwinter survival	0.047	0.044	0.286	-0.014	0.060	0.809	-	-	-	**0.335**	**0.037**	**< 0.001**
Size after winter	0.072	0.061	0.232	-0.073	0.082	0.376	-	-	-	**0.742**	**0.044**	**< 0.001**
Fruit count	0.058	0.034	0.089	-0.093	0.058	0.110	-0.010	0.084	0.905	**0.455**	**0.044**	**< 0.001**
Seed production	**0.071**	**0.033**	**0.031**	-0.035	0.055	0.522	0.028	0.077	0.719	**0.334**	**0.035**	**< 0.001**

Figure S1 Schematic of the greenhouse crossing design to generate seeds for our common gardens. Three blocks are diagrammed here, but crosses were performed in 22 blocks and seeds from 15 blocks were transplanted into the field. **(A)** Within-population crosses: for each seed family from each of 15 populations (three are shown), plants were crossed in a "daisy chain" design, in which each plant was hand-pollinated using pollen from another individual of the same population from the subsequent block. **(B)** Between-population crosses: we used pollen from each of 13 donor populations (two are shown) to pollinate flowers on plants from each of the two focal populations (one is shown). Each focal plant served as a dam for multiple between-population crosses, that is, each focal seed family had one flower pollinated by a plant from each of 13 donor populations and from the other focal population.

Figure S2 Distribution of climate differences of within- vs. between-population crosses of *Clarkia pulchella* relative to conditions during the experiment. Each dot represents a combination of maternal population, paternal population, and transplant site. Dark blue dots are within-population crosses of focal populations transplanted into their home sites. Gold dots are within-population crosses from donor populations planted into each of the two gardens, as well as the focal populations planted into each other's sites. Red dots are between-population crosses. Vertical blue bars are placed at zero, indicating where populations would be perfectly matched to the temperature or precipitation conditions during the experiment. **(A)** Distribution of the differences between the average temperature in the home sites of parental populations and conditions during the experiment. Focal populations are intermediate in temperature relative to other populations in the experiment; this results in similar average differences in temperature in between-population crosses and within-population crosses. **(B)** Distribution of differences between the average precipitation in the home sites of parental populations and conditions during the experiment. Focal populations are among the driest in the experiment; this results in smaller average differences in precipitation in between-population crosses compared to within-population crosses. Note that figures in the main text use absolute temperature differences: the absolute value of the midparent differences as they are plotted in this figure.

Figure S3 Lifetime fitness (seeds produced per seed planted) from populations of *Clarkia pulchella* with foreign vs. local parents. This analysis includes within-population plants only (no gene flow). Each point represents the average of a single family. Error bars are 95% confidence intervals of model estimated means, omitting variation from random effects.