Bayesian arrival model for Atlantic salmon smolt counts powered by environmental covariates and expert knowledge

Henni Pulkkinen, Panu Orell, Jaakko Erkinaro, Samu Mäntyniemi

H. Pulkkinen.* Natural Resources Institute Finland, Paavo Havaksen tie 3 P.O. Box 413, 90014 Oulu. Finland. henni.pulkkinen@luke.fi

P. Orell. Natural Resources Institute Finland, Paavo Havaksen tie 3 P.O. Box 413, 90014 Oulu. Finland. panu.orell@luke.fi

J. Erkinaro. Natural Resources Institute Finland, Paavo Havaksen tie 3 P.O. Box 413, 90014 Oulu. Finland. jaakko.erkinaro@luke.fi

S. Mäntyniemi. Natural Resources Institute Finland, Latokartanonkaari 9, 00790 Helsinki. Finland. samu.mantyniemi@luke.fi

Corresponding author: Henni Pulkkinen (phone: +358 40 485 0088)

Abstract

Annual run size and timing of Atlantic salmon smolt migration was estimated using Bayesian model framework and data from six years of a video monitoring survey. The model has a modular structure. It separates sub-processes of departing, traveling and observing, of which the first two together define the arrival distribution. The sub-processes utilize biological background and expert knowledge about the migratory behavior of smolts and about the probability to observe them from the video footage under varying environmental conditions. Daily mean temperature and discharge were used as environmental covariates. The model framework does not require assuming a simple distributional shape for the arrival dynamics and thus also allows for multimodal arrival distributions. Results indicate that 20 % - 43% of smolts passed the Utsjoki monitoring site unobserved during the years of study. Predictive studies were made to estimate daily run size in cases with missing counts either at the beginning or in the middle of the run, indicating good predictive performance.

Keywords: salmon, arrival model, passage count, Bayesian models, environmental covariates, expert knowledge, biological realism, missing data, prediction

Introduction

Migratory fish species are often monitored along their migration routes at different life stages to collect fishery-independent data for stock assessment. Monitoring methods include e.g. counting fences, video, sonar and snorkeling counts, and common to all is that they rarely provide perfect information about the number of individuals passing the system (Dempson et al. 1991; Romakkaniemi et al. 2000; Orell et al. 2007; 2011). Part of the fish run may be missed because of difficult environmental conditions or device failures, there may be double counting, or the data

achieved may be otherwise partial or biased (e.g. Holmes et al. 2006). Thus, the assumptions made

when interpreting the data can have a great influence on the estimated stock abundance.

Increasing the information beyond the actual monitoring data collected can improve the

trustworthiness of stock assessments (Kuparinen et al. 2012): such information can be based on the

resources available in other fields of biological research such as ecology or life history theory, but

also on the physical and technical details of the observation processes. Methods of Bayesian

inference enable both estimation of uncertainty and combination of various sources of information.

Statistical methods are often used, however, without considering the biological realism and just

focusing on statistical data analysis. Such a procedure easily leads to awkward model assumptions

that cannot be biologically interpreted and the potential effects of those modelling decisions may

pass undiscussed. We echo the idea of Kuparinen et al. (2012) and Mäntyniemi et al. (2015) in

asserting that biological plausibility has the priority over measures of statistical fit and information

theoretic parsimony when it comes to assessing the performance of stock assessment models.

During the course of time, various methods have been used to estimate run dynamics and passage

counts at point census stations. The simplest methods are based on "connect-the-dots" type of

linear interpolation (Gewin and WanHatten 2005, Johnson et al. 2007) but such approach requires

an observation before and after the missing time point and thus missing tails cannot be estimated. If

it is assumed that a proportion passing the site on a given date is constant over the years then

estimation of missing tails is possible using the methods proposed by e.g. Van Alen (2000) but

considering the variability, for example, in annual environmental conditions (e.g. Erkinaro et al.

1998; Orell et al. 2007; Otero et al. 2014), such assumption is problematic.

Hilborn et al. (1999) used a maximum likelihood method to estimate number of salmon with ground-

based stream survey, accounting also for estimate of uncertainty. Su et al. (2001) continued their

work with hierarchical Bayesian approach enabling learning from years with more data to those with

missing data. Furthermore, Sethi and Bradley (2016) introduced a Bayesian approach to estimate

missing passage at weirs with run curve model to account for arrival dynamics and process variation

model to describe the observed data. While these studies attempt to account for uncertainty, they

do not discuss their model assumptions against biological knowledge. This together with

consideration of difference between the observation and model prediction as "noise" instead of

proportion unobserved (as e.g. in Sethi and Bradley 2016), may cast some doubt on the biological

meaningfulness of the results.

In our study, a Bayesian model framework was introduced for estimating the annual number of

Atlantic salmon (*Salmo salar*) smolts passing the video monitoring site at a subarctic River Utsjoki, a

tributary of the large River Teno in northern Finland (cf. Erkinaro et al. 2019). The framework

consists of sub-processes of departing, travelling and observing. When combined, the processes of

departing and travelling form the arrival distribution that describes the timing of the smolt run.

Environmental covariates were used as a source of information in the sub-processes and expert

knowledge is utilized in providing biologically meaningful model structures and prior distributions.

Hierarchical structure was assumed over the study years making it possible to learn from the

processes, borrowing strength from data rich years of study to those with missing data. Predictive

studies illustrate the model performance in cases with missing data either at the beginning or in the

middle of the run. We show that a biologically meaningful passage model can be built without

assuming any specific shape for the arrival distribution. Furthermore, we claim that existing expert

knowledge about the observability of smolts is a key source of information in estimating the total

uncertainty about the annual total passage especially when environmental conditions have a great

influence on the probability to observe individuals passing the survey site. Eventually, the resulting

estimates describe the logical inference about the smolt passage that is formed by carefully

analyzing the structure of the problem and interpreting observed data in the light of expert's existing

knowledge about migratory behavior and accuracy of video counting.

Materials and Methods

Data

River Utsjoki (drainage area 1652 km^2) is a tributary of the large River Teno system at the northernmost border between Finland and Norway. Salmon production areas extend c. 60 km upstream in the Utsjoki mainstem as well as c. 30 km along two major tributaries (see Borgstrøm et al. 2010). Each spring, the monitoring system of 8 video cameras is set on the river bed at the main channel under a bridge close to the tributary outlet (Figure 1). The video footage of these cameras provides data on both smolts descending and adults ascending the river during the course of summer. Most of the individuals migrate via the main channel, but occasional video monitoring in the side channels has indicated that a few percentages of the smolts would pass through the sides. The installation and use of cameras is described in detail by Davidsen et al. (2005) and Orell et al. (2007). The camera technology and operation have been similar throughout the monitoring years (2002 - present), but some updates have been done to the data recording system, without affecting comparability of the data. In this study we analyze the smolt count data at the main channel from years 2005-2009 and 2014. These years were chosen to demonstrate the highly variable environmental conditions, especially the strongly varying discharge (Figure 2). The smolt data are aggregated in daily counts of individuals over 61 days of June-July.

The datasets on environmental covariates, air temperature and discharge, are aggregated into daily averages. Both covariates are measured near the video site, but conditions may differ between the video site and upper parts of the river where some of the smolts begin their migration from. These datasets are, however, considered as reasonable proxies for the environmental conditions affecting the migration behavior of smolts.

Data from the first 23 days of June in 2005 are missing because the high level of water delayed the

installation of the monitoring system. Thus, the missing counts are predicted for those dates.

Predictive studies were performed by treating data in 2007 and 2014 as partly missing: For 2007 the

first 17% of the run strength and for 2014 the peak +-2 days count was removed from the analysis.

These years were chosen to the predictive study as the environmental conditions in those years fell

in between the extreme conditions of the overall study years. To evaluate the predictive

performance of the model, the model predicted counts for the dates treated as missing were

compared with the observed data.

Expert elicitation as a source of informative priors

Model structure and informative priors were elicited from an expert, who is most familiar with the

behavior of salmon smolts and the video monitoring system at Utsjoki. In addition, the expert has

comprehensive personal experience on studying smolt migration in other tributaries of the River

Teno catchment, as well on other salmon rivers in northern Fennoscandia. The expert also has

extensive knowledge about various visual monitoring methods like video monitoring and snorkelling,

and has studied their possibilities and restrictions.

The expert was asked to base his views on his overall biological knowledge about the behavior of the

smolts: when do the smolts potentially make their decision to depart given the temperature, how

long their travelling may take from the nearest and furthest locations of origin, and how the

environmental conditions affect the proportion of smolts that may pass the video site unobserved.

Extra care was taken to ensure that the expert would express his views based on knowledge

independent of the data that would be analysed in the model since the avoidance of double usage of

the same data is vital for reliable estimation of total uncertainty.

Elicitation was carried out iteratively, first by asking preliminary questions on e.g. the biological

process in question, and then providing graphical illustration of the parameters as a function of

environmental conditions (as in Figures 7-9). Prior distributions were adjusted until the expert

agreed that the illustration is in line with his views. The questions were not defined – or the answers

translated – directly to inform about prior distributions of specific parameters as the marginal

distributions often lack good real-life interpretation. Instead, the focus was on the visual form of the

joint distributions of each sub-process as a function of an environmental covariate, as will later be

shown in figures 7-9. The marginal distributions of parameters related to each sub-process were

adjusted until the joint distribution corresponded roughly the view of the expert. The direct

questions mainly addressed the mean values of the sub-processes and the levels of environmental

covariate at which significant changes in the process were expected to take place. Questions elicited

from the expert are listed in table 1.

Probability model of the problem

The model is constructed by thinking of four steps, described in detail in the following sections:

1) Process of departing, in which the smolts make the decision to depart while daily
   temperature affects the probability to depart in a given day;
2) Process of travelling, in which the smolt's travel time (in days) from the place of departure to
   the video site is modeled with daily discharge as a covariate;
3) Arrival distribution, which links together the processes of departing and travelling into a
   timing of the smolt run;
4) Observation process, in which the discharge influences visibility and the probability that an
   individual smolt passing the site is observed from the video footage.

Process of departing

An individual smolt's probability to begin the migration in day i of year y given that the smolt has not

departed yet is considered to depend on the temperature on that day. We assume a logit-normally

linear relationship for the smolt's probability to depart $(p_{i,y})$ and temperature $t_{i,y}$:

(1) $logit(p_{i,y}) \sim N(\alpha^P + \beta^P\ t_{i,y}\ ,\ s^P)$

Informative prior distributions are given to parameters α^P, β^P and s^P according to expert view (see

tables 1 and 2).

The daily probabilities of departing (given those individuals have not departed yet; $p_{i,y}$) can be

transformed into expected proportions of the total smolt run that depart each day $(P_{i,y})$ as

(2) $P_{1,y} = p_{1,y}$

(3) $P_{i,y} = (1 - \sum_{k=1}^{i-1} p_{k,y})\ p_{i,y}$

(4) $P_{61,y} = 1 - \sum_{i=1}^{60} p_{i,y}$.

Process of travelling

According to the expert view, each smolt should arrive at the video site within 14 days after

departing. Thus we assume that the smolts that depart on day i pass the video site in one of the next

14 days according to discretized distribution function of lognormal distribution (Schwarz & Dempson

1994, Mäntyniemi & Romakkaniemi 2002):

(5) $Q^D_{i,j,y} = \phi\left(\frac{\log(j-i+0.5-M_{i,y})}{s}\right) - I_{i,j}\phi\left(\frac{\log(j-i+0.5-M_{i,y})}{s}\right)$.

Here ϕ is the cumulative density function of the standard normal distribution, $M_{i,y}$ is the mean of

log(travel time) to video site of smolts that depart on day i , and S is the standard deviation of

log(travel time). Thus, $Q^D_{i,j,y}$ is the conditional probability that a smolt that departed on day i will

arrive on day j, when j goes from i to $i + 13$.

To ensure that all smolts arrive within 14 days of departing, $Q^D_{i,j,y}$'s are standardized:

(6) $q^D_{i,j,y} = Q^D_{i,j,y} / \sum_{j=i}^{(i+13)} (Q^D_{i,j,y})$,

and thus $\sum_{j=i}^{(i+13)} (q^D_{i,j,y}) = 1$.

It is further considered that increasing discharge increases the speed at which the smolts travel,

shortening their travel time. The expected travel time on real scale $\mu^D_{j,y} = \exp(M_{i,y} + 0.5\,S^2)$ is

assumed to follow a lognormal distribution with expected value $\gamma_{i,y}$ and coefficient of variation cv^D

and it is assumed to depend on the discharge on the day of departure ($f_{i,y}$):

(7) $\gamma_{i,y} = \exp(\alpha^D + \beta^D \cdot f_{i,y})$.

Again, informative prior distributions are given for parameters α^D, β^D and $cv^{\mu D}$ according to expert

view (see tables 1 and 2).

Arrival distribution

By combining the processes of departing and travelling, we achieve joint probabilities that a smolt

departs on day i and passes the video site on day j:

(8) $q^J_{i,j,y} = q^D_{i,j,y}\, P_{i,y}$.

The expected proportions of smolts (out of the total run size) passing the video site each day can

then be found by marginalizing over the day of departure:

$\quad\quad$ (9) $\mu_{j,y}^N = \begin{cases} \sum q_{j,1:j,y}^J \ , \ j \le 14 \\ \left(\sum q_{j,(i-13):j,y}^J \ , j \ > 14. \right. \end{cases}$

Furthermore, we give a uniform prior distribution on log scale for the total annual size of the smolt

run passing the video site:

$\quad\quad$ (10) $\quad\quad \log(N_y^{tot}) \sim Unif(7,15).$

Parameterisation on the log-scale allows for a wide, minimally informative prior distribution for the

annual run size. One could also use a more biologically realistic option for estimating the annual

abundance based on adult escapement or on expert evaluated smolt production capacity of the river

(e.g. Uusitalo et al. 2005). Because the results of this analysis are going to be used in the future

research as building block of a sequential modelling framework for the total life cycle of Utsjoki

salmon (similarly as in Michielsens et al. 2008), a minimally informative prior distribution for annual

production is needed. This ensures that the information flowing from the full life cycle model will be

used only once.

The annual run size is assumed to be Dirichlet-multinomially distributed over the 61 days of June-

July. For computational simplicity, we follow Mäntyniemi et al. (2015) and approximate multinomial

distribution as

$\quad\quad$ (11) $\quad\quad N_{j,y} = \ q_{j,y}^N \ N_y^{tot},$

where $q_{j,y}^N$ is the proportion of smolts passing the video site in day j and $q_{j,y}^N$'s follow a Dirichlet

distribution parameterized with the expected proportions of smolts passing the video site each day

$\mu_{j,y}^N$ and overdispersion η^N:

(12) $q_{1:61,y}^{N} \sim Dirich(\mu_{1,y}^{N}\eta^{N}, \ldots, \mu_{61,y}^{N}\eta^{N}).$

To ease the computation we approximate the Dirichlet-distribution with a set of lognormal

distributions (see appendix A for details).

The annual proportional prior distributions for the the smolt run over 61 days of June-July are

illustrated in figure 3. These priors are based on the model structure and data on environmental

covariates (daily mean air temperature and discharge).

Observation process

In previous sections we have covered the processes assumed to affect the timing of the smolt

passage at the video site. To estimate the total uncertainty in the smolt run size, the process of

observing smolts from the video footage needs to be included. It seems natural to assume that some

of the individuals pass the site unobserved, and thus this proportion must be accounted for. We

introduce an observation process that builds on expert view on how high the probability to observe

a smolt may be in excellent vs. poor conditions on visibility and how the discharge is considered to

affect this probability. The number of smolts observed follows a Beta-binomial distribution:

(13) $N_{j,y}^{Obs} \sim Beta - bin(\mu_{j,y}^{B}\eta^{B}, \ (1 - \mu_{j,y}^{B})\eta^{B}, \ N_{j,y})$

Here $N_{j,y}$ is the total daily number of smolts passing the video site and the probability to observe a

smolt follows a $Beta(\mu_{j,y}^{B}\eta^{B}, \ (1 - \mu_{j,y}^{B})\eta^{B})$ distribution, where $\mu_{j,y}^{B}$ is the expected probability for

a smolt to be observed and η^{B} is the overdispersion parameter. Overdispersion in this context is

thought to arise from the schooling behavior of smolts. Groups of smolts can be either detected or

missed, which increases the variance of the observation process (Mäntyniemi & Romakkaniemi

2002, Linden & Mäntyniemi 2011, Mäntyniemi et al. 2015). The expected value of the process is

244 linked to discharge following the expert view that in very good visibility (low discharge) at maximum

245 90% of the smolts are observed. As the discharge increases, the visibility decreases, and the volume

246 of water to be covered by the video monitoring increases. Therefore, according to the expert,

247 observation probability decreases gradually to a minimum of 30%. The expected probability $\mu_{i,y}^B$ is

248 considered to follow logit-normally linear relationship on the interval [0.3,0.9]:

249  (14) $\mu_{j,y}^B = 0.6 \left(\frac{\exp(BB_{j,y})}{1+\exp(BB_{j,y})} \right) + 0.3$

250  (15) $BB_{j,y} \sim N\left(\alpha^B - \beta^B f_{j,y}, \ sd_{BB}^2 \right),$

251 where α^B, β^B and sd_{BB} have prior distributions according to the expert view (Tables 1 and 2).

252 *Bayesian inference and sensitivity analysis*

253 Bayesian analysis that is not sensitive to prior information simply cannot exist. The most influential

254 component of the inference just gradually shifts from prior information about the parameters

255 towards prior information about the data collection process as more data accumulates. From the

256 inferential point of view, both extremes and anything between are equally valid: the resulting

257 posterior shows what our logical state of mind should be after interpreting the data. Sometimes the

258 state of mind is that we learned nothing new about all or some of the parameters.

259 However, a commonly held belief is that large quantities of data "override" the prior knowledge and

260 thus decrease the sensitivity of the resulting posterior distribution to prior information. Here we

261 show why this is not the case. The key fact is that it is not the data itself that changes the prior, or

262 leaves it unchanged. Prior information becomes updated by interpreting data, and that

263 interpretation is based on prior information. This can be seen from the Bayes' rule

$$p(\theta|x) \propto p(x|\theta)p(\theta) = p(x,\theta)$$

which states that the posterior distribution of parameters (θ) is proportional to the joint probability

distribution of parameters and potentially observable data (x). This distribution summarizes what is

known about the entire problem before obtaining the observed data values.

All the joint probabilities $p(x, \theta)$ have a Bayesian interpretation of being measures of degree of

belief. In principle, the person whose belief is being measured could state these joint probabilities

directly. However, in practice most people find it easier to construct their joint beliefs by factorizing

their thinking between what they knew about the parameters $(p(\theta))$ and how they would use their

existing knowledge to predict data if they knew the parameter values $(p(x|\theta)$ (More elaborate

factorization can also be used, as demonstrated in previous sections).

The conditional probabilities $p(x|\theta)$ describe the degrees of belief about the link between

parameters and data, providing a predetermined interpretation of any potential data values. Once

the values are obtained, the posterior distribution (up to a constant) is found by keeping only those

joint probabilities that had been pre-assigned to values that became observed. In other words, the

observed data objectively selects the subjective joint prior degrees of belief that will be kept to form

the posterior distribution.

Why sensitivity analysis?

As shown above, the results of any Bayesian model are always guaranteed to be sensitive to some

aspects of prior knowledge. The role of a sensitivity analysis is to provide information about the

relative importance of these different pieces of prior information. Some idea of the relative roles of

the information sources can be obtained simply by comparing the priors and posteriors: if the

posterior distribution of a parameter is (almost) identical to its prior, then we can conclude that

compared to our existing knowledge, our interpretation of the data did not provide new

information. This is a case where a sensitivity analysis can reveal deeper insight on the structure of

the problem: would our interpretation of data provide at least some new information if we

hypothetically had less information about the parameters to begin with? Such a question can be

answered by repeating the analysis with priors that express more uncertainty than we actually have.

Sensitivity study on observability priors

Our preliminary results of the arrival model indicated that parameters α^B and β^B which describe the

relationship between the observability of smolts and the discharge, did not update from their priors.

In order to gain more in-depth understanding of this result, we performed a sensitivity analysis by

pretending that our prior knowledge would be weaker than it actually is: parameters α^B and β^B

were given more uncertain prior distributions compared to priors based on expert elicitation (Table

2). More details and results of this study can be found from Appendix B.

Details on model run

Model covering all the sub-processes was fitted using MCMC sampling with JAGS 4.2.0 software (Just

Another Gibbs Sampler; Plummer 2003) and R runjags package (Denwood 2016). Model was run

with 2 chains for 7 500 000 iterations which took around 20 days. Burn-in period of 500 000

iterations was removed from the beginning of the run. Convergence was diagnosed using Gelman-

Rubin statistics (Gelman and Rubin 1992), ensuring that the upper limit for the potential scale

reduction factor was below 1.1 for all parameters of interest. The complete JAGS code for the model

structure can be found from Appendix C.

Results

When compared with the observed smolt counts, annual mean posterior estimates of daily run size

indicate that 20 - 43 % of smolts passed the video site unobserved during the years of study (Figure

4). The highest proportion of unobserved was estimated for 2008, median being 43 % and 90 %

probability interval (PI) [38 %, 47 %]. This result reflects the high level of discharge in early/mid-

season in 2008 (Figure 2).

Scenarios to study predictive performance of the model ('first 17% of missing' -scenario for 2007 and

'peak day +- 2 days missing' –scenario for 2014) show that daily counts fall in the predicted 90% PI

indicating good predictive performance (Figure 5, Table 3). For 2005, when the high level of water

delayed the setup of the monitoring system until June 23rd, posterior mean indicates that 10% of

the run was missed during this period (90 % PI being [3 %, 23 %] of the run). Because the

temperature was below the average until mid-June in that year, it is not likely that the migration

would have started very early in the season (Figures 2 and 5).

When comparing marginal posterior distributions with corresponding priors, 7 key parameters out of

12 updated, indicating that the learning does not take place evenly about all parameters (Table 4).

The joint posterior distributions of the process of departing as a function of temperature (consisting

of parameters α^P, β^P and sd^P) updates quite heavily from the prior (Figure 6). Updating takes place

towards higher temperatures indicating, for example, that 50 % probability to depart is met with

50% certainty around 14 °C compared to a priori belief of this level being reached already at 11 °C.

The posterior cumulative distribution of the expected travel time to the video site (consisting of

parameters α^D, β^D, cv^D and $cv^{\mu D}$) supported somewhat longer travel times than expected a priori

both for low and high discharge (Figure 7). At low discharge (10m^3/s), 50 % of smolts were estimated

to arrive at the video site within 5 days (posterior median), whereas a priori the same proportion

was considered to be met around 3.5 days. At high discharge (100m^3/s), 50 % of smolts would arrive

by 3 days based on the posterior and by a bit less than 2 days based on the prior. Note that posterior

cumulative distributions at these specific levels of discharge were chosen for illustrative purposes

and that in practice the model allows any positive value for the discharge.

Joint priors and posteriors of observability parameters α^B, β^B and sd^{BB} are close the same (Figure

8) indicating that in this case the combination of priors and our interpretation of data did not result

in further learning about the process of observing. Sensitivity analysis (Appendix B) revealed that if

we happened to have more uncertainty about these parameters, we would have clearly learned

something new about the role of discharge (β^B) in the observability of smolts. The parameter

responsible for the baseline of the observability (α^B) was more sensitive to prior uncertainty about

it: no new information after seeing the data could be obtained within the range of parameter values

studied.

Discussion

In our study, an arrival model for Atlantic salmon smolts was designed by separating the processes

of departing, travelling and observing. The processes of departing and travelling together form the

arrival distribution, i.e. the timing of the smolt run. This modular structure enabled us to concentrate

on biologically meaningful mathematical assumptions related to each sub-process instead of

assuming that fish count data as such follows a specific distributional shape. The model framework

aimed at high biological realism by accounting for expert knowledge in the model formulation, firstly

by including choices for model structures and prior distributions, and secondly by linking

environmental covariates to provide information about the timing of the migration. The estimated

total uncertainty accounted for the uncertainty in the sub-processes, including the observation

process in which variability in discharge is known to have a large impact on the probability to

observe passing smolts.

One of the most important advantages of our approach is that the need to assume a specific

parametric shape for the arrival distribution is avoided. Past studies (Hilborn et al. 1999, Su et al.

2001, Sethi and Bradley 2016) have assumed unimodal shape (e.g. normal, skew-normal, Student's t)

for the arrival curve. However, when taking a look at the shape of (any) arrival time series it seems

clear that such assumption is a tolerable simplification at best (see e.g. figure 5 in this paper, figure 2

in Sethi and Bradley 2016).

Instead of basing the arrival curve on a specific distributional form without biological interpretation,

we have based our model structure largely on the expert knowledge. While one can state that the

expert may be wrong in his/her views, it can also be argued that replacing the available knowledge

with ad-hoc but conventional choices of distribution has even less of a chance to end up with a

"true" outcome. The expert elicitation requires ensuring that the experts are able to express their

views about the processes independent from the data that will be analysed. This can be problematic

at first as often the same experts have the best knowledge about the issues related to conditions at

the study area, behavior of the species, details on the data collection and the data itself. We

recommend that questions elicited should be designed (similarly as in our study) to target the sub-

processes and phenomenon behind the data, not the data itself.

Fitting the elicited information together with the model structure can also be challenging. For

example, when we combined the expert views on the observation process with a logistic functional

form, very low uncertainty was inevitably assumed at high (>60 m^3/s) discharge. However, such

discharges usually take place only at the beginning of the season when the smolt run has not really

started (Orell et al. 2007), and thus the impact of the assumption on the estimated total run size is

minor. Regardless of methods chosen on which the model is built upon it is important to keep in

mind that the results depend always on the combination of the objective data and subjective model

choice.

Earlier studies have paid little attention to the role of observation models in estimating the total

uncertainty. Hillborn et al. (1999) distinguished process and observation errors, assuming observer

efficiency either as known or unchanging inside the study period. Su et al. (2001) discussed the issue

of error in the observation process and mainly concluded that such error can cause convergence

problems in sampling. Sethi and Bradly (2016) incorporated uncertainty only on the estimates of

missing passage counts, treating the observed counts as true total number of passing without

uncertainty. As they assumed normal or negative Binomial models for the observed passage counts,

they in fact allowed model predicted passage counts to be either greater or smaller than the

observed counts. This in turn can lead to underestimation of the missing counts. We claim that it is

necessary to consider the observed count as a minimum estimate for the true number especially if

the back-and-forth movement of individuals is negligible. Along with this notion the need for a

realistic observation process becomes obvious. Depending on the approach, the estimate of total

annual passage can be very different. As Hilborn et al. (1999) put it, 'when we admit uncertainty in

observer efficiency we become much less certain about the actual escapement'.

The sensitivity analysis showed that in our case video counts could provide information about the

effect of discharge on the observability. This can be explained by the fact that the daily video counts

are thought to depend on the number of smolts migrating and their observability, both of which are

thought to depend on daily varying environmental covariates. The daily number of migrating smolts

is controlled by temperature and discharge -related parameters, but the model fit to observed

counts can be further improved by changing the observability according to discharge. Interestingly,

the sensitivity analysis showed that the interpretation of data favors the same parameter values

than the expert's prior. However, even though the signal from the data agrees with the expert's

existing knowledge, the signal is weak compared to expert's already existing certainty. Therefore the

certainty does not notably increase. The sensitivity analysis also revealed that in our case the general

level of the observability of smolts is highly dependent on the existing expert knowledge: very little

or no new information could be inferred from the video counts. This underlines the need for

rigorous expert elicitation.

If the purpose of the model would have been to replicate the observed data set with as few

parameters as possible, then the sensitivity of the results to the observability priors would be a sign

of overparameterisation: these parameters could be the first to be dropped from the model. The

concept of population size would probably be unnecessary too for such a purpose. However, our

goal is not to find the most parsimonious model for data, which would be the purpose of

information theory. Instead, we are dealing with an inverse problem: given the data and our existing

knowledge about migration and observation processes, what can we realistically say about the

population size? Within this approach the sensitivity is not of consequence when it comes to

simplifying the model. As shown earlier, any Bayesian model is completely made of prior knowledge

and always sensitive to it in one way or another. Any model simplification should be based on

biological plausibility, rather than on the ability of the model to replicate data (Kuparinen et al.

2012). Using information theoretic arguments to simplify a model meant for solving inverse

problems can lead to underestimation of actual uncertainty (Kuparinen et al. 2012) which causes

underestimation of the value of any new information that could be potentially gathered

(Mäntyniemi et al. 2009).

Sethi and Bradley (2016) fitted datasets in their study using several combinations of arrival and

process error models. They utilized information-theoretic model selection (DIC, Spiegelhalter et al.

2002) to find out the best fitting model for each dataset. However, no discussion took place whether

some of these models would be biologically more reasonable than another. We argue that the

model selection should primarily be based on biological realism and in case biological model

uncertainty is needed to be tackled, Bayesian model averaging (BMA, Hoeting et al. 1999) should be

preferred over DIC or other information criteria.

Su et al. (2001) used hierarchical structure in their study, assuming that years in which timing of the

run is well known can be used for passing information to years with missing data. We argue that

learning should rather take place over the (biological) process, assuming exchangeability over year

specific parameters of those processes. Our model structure can also be considered as hierarchical,

although we simply assume that the underlying processes are the same from year to year allowing

between- and within-years variation. If there were a need for a meta-analysis of similar studies over

several stocks of the same species, exchangeability in biological parameters could be considered

between the stocks. Such approach could also allow meta-analysis of data from different types of

monitoring systems if detailed observation processes were tailored for each system and

exchangeability was considered as reasonable assumption for the biological processes.

Our model framework could be fine-tuned and developed further. In an ideal situation,

environmental covariates would be available from upstream locations from where the smolts

originate. Data on air temperature would be replaced with water temperature to better explain the

physiological processes the smolts encounter. The total annual estimates could be improved by

including individuals that migrate via side channels. One year of video data from the side channels is

available but larger set of data would be needed to evaluate the effect of varying discharge on the

proportion of smolts passing via the main vs. side channels. To increase the realism even more, the

schooling behavior of smolts could be accounted for in more detail than done here. Salmon smolts

have a tendency to school during their migration (e.g. Bakshtanskiy et al. 1988, Davidsen et al. 2005)

and this creates more variation in the monitoring data compared to independent movement

(Mäntyniemi and Romakkaniemi 2002, Mäntyniemi et al. 2015). Schooling behavior could also be

linked with the process of travelling as the passing schools of fish have been suggested to stimulate

migratory behavior in the remaining smolts (e.g. Hansen and Jonsson, 1985). Schooling also affects

the observation process as schools are easier to spot than individual smolts, and on the other hand,

it can be difficult to have an exact count of individuals from a large school. For Utsjoki video survey,

daily average school sizes of smolts are recorded and incorporating those into the model framework

is an interesting topic for future research.

Despite the above possibilities for further development of the modeling framework, we claim that

our model already contains the elements of realism that are required to provide credible estimates

of the total annual smolt run. This approach can be applied to similar monitoring surveys where

point-checking the numbers of migrating fish (Trépanier et al. 1996), by e.g. sonar, trap, or weir is

used for estimating run sizes. As far as we know, environmental covariates have not been included

so far to estimate both the dynamics of passage and the observation process. Our model framework

also indicated a good predictive performance for missing dates both at the beginning and in the

middle of the run. Furthermore, the results from this study will act as a single step of sequential

model framework that covers the total life cycle of multiple populations in the River Teno complex

(Vähä et al. 2017; Erkinaro et al. 2019), similarly as the system designed for Baltic salmon stocks

(Michielsens et al. 2008).

Acknowledgements

465 *We thank the numerous field workers enabling the yearly monitoring programme and smolt data*

466 *collection at the River Utsjoki. We also acknowledge the crucial technical knowhow by Jorma Kuusela*

467 *in running the annual monitoring and the instrumental role of Dr. Martin Svenning and Anders*

468 *Lamberg in early phases of the River Utsjoki monitoring programme. Funding for this study was*

469 *provided by the Academy of Finland (project No. 286334).*

References

Bakshtanskiy, E., Nesterov, V., and Neklyodov, M. 1988. Development of schooling behaviour in

juvenile Atlantic salmon, Salmo salar, during seaward migration. J. Ichthyol. 28: 91–101.

Borgstrøm, R., Opdahl, J., Svenning, M.-A., Länsman, M., Orell, P., Niemelä, E., Erkinaro, J. &

Dempson, J. B. 2010. Temporal changes in ascendance and is-season exploitation of Atlantic salmon,

Salmo salar, inferred by a video camera array. Fish. Manage. Ecol. 17: 454-463.

Davidsen, J., Svenning, M. A., Orell, P., Yoccoz, N., Dempson, J. B., Niemelä, E., Klemetsen, A.,

Lamberg, A., and Erkinaro, J. 2005. Spatial and temporal migration of wild Atlantic salmon smolts

determined from a video camera array in the sub-Arctic River Tana. Fish. Res. 74: 210-222.

Dempson, J. B, and Stansbury, D. E. 1991. Using partial counting fences and atwo-sample stratified

design for mark–recapture estimation of an Atlantic salmonsmolt population. N. Am. J. Fish.

Manage. 11: 27–37.

Denwood, M.J. 2016. runjags: An R Package Providing Interface Utilities, Model Templates, Parallel

Computing Methods and Additional Distributions for MCMC Models in JAGS. J. Stat. Softw. 71(9): 1-

25. doi:10.18637/jss.v071.i09.

Erkinaro, J., Julkunen, M. and Niemelä, E. 1998. Migration of juvenile Atlantic salmon Salmo salar in

small tributaries of the subarctic River Teno, northern Finland. Aquaculture 168: 105-119.

Erkinaro, J., Czorlich, Y., Orell, P., Kuusela, J., Länsman, M., Falkegård, M., Pulkkinen, H., Primmer, C.

& Niemelä, E. 2019. Life history variation across four decades in a diverse population complex of

Atlantic salmon in a large subarctic river. Can. J. Fish. Aquat. Sci. 76: 42-55.

Gelman, A., and Rubin, D.B. 1992. Inference from iterative simulation using multiple sequences. Stat.

Sci. 7: 457–511.

Hilborn, R., Bue, B.G., and Sharr, S. 1999. Estimating spawning escapements from periodic counts: a

comparison of methods. Can. J. Fish. Aquat. Sci. 56(5): 888–896. doi:10.1139/f99-013.

Hansen, L.P., and Jonsson, B. 1985. Downstream migration of hatchery-reared smolts of Atlantic

salmon (Salmo salar L.) in the River Imsa. Aquaculture, 45: 237–248.

Holmes, J.A, Cronkite, G.M.W., Enzenhofer, H.J., and Mulligan, T.J. 2006. Accuracy and precision of

fish-count data from a "dual-frequency identification sonar" (DIDSON) imaging system. ICES J. Mar.

Sci. 63: 543–555. doi:10.1016/j.icesjms.2005.08.015

Kuparinen, A., Mäntyniemi, S., Hutchings, J.A., and Kuikka, S. 2012. Increasing biological realism of

fisheries stock assessment: towards hierarchical Bayesian methods. Environ. Rev. 20: 135–151.

doi:10.1139/A2012-006.

Michielsens, C.G.J., McAllister, M.K., Kuikka, S., Mäntyniemi, S., Romakkaniemi, A., Pakarinen, T.,

Karlsson, L., and Uusitalo, L. 2008. Combining multiple Bayesian data analyses in a sequential

framework for quantitative fisheries stock assessment. Can. J. Fish. Aquat. Sci. 65: 962-974.

doi:10.1139/F08-015

Mäntyniemi, S., and Romakkaniemi, A. 2002. Bayesian mark-recapture estimation with an

application to a salmonid smolt population. Can. J. Fish. Aquat. Sci. 59: 1748-1758. doi:10.1139/F02-

146.

Mäntyniemi, S.H.P., Whitlock, R.E., Perälä, T.A., Blomstedt, P.A., Vanhatalo, J.P., Rincon, M.M.,

Kuparinen, A.K., Pulkkinen, H.P., and Kuikka, O.S. 2015. General state-space population dynamics

model for Bayesian stock assessment. ICES J. Mar. Sci. 72: 2209-2222. doi:10.1093/icesjms/fsv117

Mäntyniemi, S., Kuikka, S., Rahikainen, M., Kell, L.T., Kaitala, V. 2009. The value of Information in

fisheries management: North Sea herring as an example. ICES J. Mar. Sci. nce 66: 2278-2283.

Orell, P., Erkinaro, J., Svenning, M., Davidsen, J., and Niemelä, E. 2007. Synchrony in the

downstream migration of smolts and upstream migration of adult Atlantic salmon in the sub-Arctic

River Utsjoki. J. Fish. Biol. 71: 1735–1750.

Orell, P., Erkinaro, J., and Karppinen, P. 2011. Accuracy of snorkelling counts in assessing spawning

stock of Atlantic salmon, Salmo salar, as verified by radio-tagged fish and underwater video

monitoring. Fish. Manag. Ecol. 18: 392–399.

Otero, J., L'Abée-Lund, J.H., Castro-Santos, T., Leonardsson, K., Storvik, G.O., Jonsson, B., Dempson,

B., Russell, I.C., Jensen, A.J., Baglinière, J.-L., Dionne, M., Armstrong, J.D., Romakkaniemi, A., Letcher,

B.H., Kocik, J.F., Erkinaro, J., Poole, R., Rogan, G., Lundqvist, H., MacLean, J.C., Jokikokko, E.,

Arnekleiv, J.V., Kennedy, R.J., Niemelä, E., Caballero, P., Music, P.A., Antonsson, T., Gudjonsson, S.,

Veselov, A.E. Lamberg, A., Groom, S., Taylor, B.H., Taberner, M., Dillane, M., Arnason, F., Horton, G.,

Hvidsten, N.A., Jonsson, I.R., Jonsson, N., McKelvey, S., Næsje, T., Skaala, Ø., Smith, G.W., Sægrov, H.,

Stenseth, N.C. and Vøllestad, L.A. 2014. Basin-scale phenology and effects of climate variability on

global timing of initial seaward migration of Atlantic salmon (Salmo salar). Glob. Change Biol. 20: 61-

75.

Plummer, M. 2003. JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling.

In proceedings of the 3[rd] International Workshop on Distributed Statistical Computing (DSC 2003),

20-22 March, Vienna, Austria. ISSN 1609-395X.

Romakkaniemi, A., Lilja, J., Nykänen, M., Marjomäki, T.J. and Jurvelius, J. 2000. Spawning run of

Atlantic Salmon (Salmo salar) in the River Tornionjoki monitored by horizontal split-beam

echosounding. Aquat. Living Resour. 13: 349-354.

Schwarz, C., and Dempson, J. 1994. Mark-recapture estimation of a salmon smolt population.

Biometrics. 50: 98-108.

Sethi, A.S., and Bradley, C. 2016. Statistical arrival models to estimate missed passage counts at

fish weirs. Can. J. Fish. Aquat. Sci. 73: 1251–1260. doi:10.1139/cjfas-2015-0318.

Spiegelhalter, D.J., Best, N.G., Carlin, B.P., and van der Linde, A. 2002. Bayesian measures of model

complexity and fit (with discussion). J. R. Stat. Soc. Ser. B. 64: 583–639.

Su, Z., Adkison, M.D., and Van Alen, B.W. 2001. A hierarchical Bayesian model for estimating

historical salmon escapement and escapement timing. Can. J. Fish. Aquat. Sci. 58: 1648–1662.

doi:10.1139/f01-099.

Trépanier, S., Rodríquez, M. A. and Magnan, P. 1996. Spawning migrations in landlocked Atlantic

salmon: time series modelling of river discharge and water temperature effects. J. Fish Biol. 48: 925–

936.

Laura Uusitalo, Sakari Kuikka, Atso Romakkaniemi; Estimation of Atlantic salmon smolt carrying

capacity of rivers using expert knowledge, ICES Journal of Marine Science, Volume 62, Issue 4, 1

January 2005, Pages 708–722, https://doi.org/10.1016/j.icesjms.2005.02.005

Vähä, J.P., Erkinaro, J., Falkegård, M., Orell, P. & Niemelä, E. 2017. Genetic stock identification of

Atlantic salmon and its evaluation in a large population complex. Can. J. Fish. Aquat. Sci. 74: 327–

338.

 Table 1. Questions elicited from the expert

Question	Expert view
Process of departing	
- At what range of temperature (air temperature at Utsjoki logger) first smolts are considered to begin their migration?	6-8 °C
- What level of temperature would be considered so high that the process of migration not be influenced by further increase in temperature (i.e. logit curve levels out)?	12 °C
Process of travelling	
- What is the the maximum travel time (in days) in which a smolt can be considered to arrive at the video site?	14 days
- What is the minimum travel time in which a smolt can be considered to arrive at the video site?	1-2 days
Process of observing	
- What is the upper (/lower) limit for the expected probability at which an average smolt is considered to be observed in best (/worst) possible environmental conditions, i.e. at minimum (/maximum) discharge?	0.9 (0.3)
- What is the level of discharge below which the visibility is considered optimal?	20 m^3/s
- At optimal visibility, what is the range of probability at which the fish are considered to be observed with around 90% of certainty?	0.75-0.9
- What is the level of discharge at which the visibility is considered to be poor?	50-60 m^3/s
- At poor visibility, what I the range of probability at which the fish are considered to be observed with around 90% of certainty?	0.3-0.6

587   Table 2: List of symbols, descriptions and prior distributions

Symbol	Description	Prior distribution
$logN_y^{tot}$	Total annual run size on log-scale	Uniform(7,15)
N_y^{tot}	Total annual run size on real scale	-
$N_{j,y}$	Daily number of smolts passing the video site	-
$q_{j,y}^N$	Daily proportion of smolts passing the video site	-
$\mu_{j,y}^N$	Expected proportion of smolts passing the video site	-
η^N	Overdispersion of the arrival process	Uniform(0.001,100000)
$p_{i,y}$	Probability to depart (if haven't departed yet)	-
$P_{i,y}$	Proportion of smolts departing each day	-
α^P	Intercept for probability to depart given temperature	N(-20,1)
β^P	Regression coefficient for probability to depart given temperature	logN(0.6,0.1)
s^P	Standard deviation of probability to depart	logN(0,1)
$Q_{i,j,y}^D$	Unstandardised proportion for travel time	-
$q_{i,j,y}^D$	Standardised proportion for travel time	-
$M_{i,y}$	Mean of log(travel time)	-
S	Standard deviation of log(travel time)	-
$\mu_{i,y}^D$	Expected travel time	-

$\gamma_{i,y}$	Expected value of expected travel time	-
cv^D	Coefficient of variation of log(travel time)	Uniform(0.001,2)
α^D	Intercept for the expected value of expected travel time given flow	logN(0.52,0.07)
β^D	Regression coefficient for the expected value of expected travel time given flow	logN(-4.6,0.04)
$cv^{\mu D}$	Coefficient of variation of expected travel time	Uniform(0.001,1)
$q^J_{i,j,y}$	Probability that a smolt departs on day i and passes the video site on day j	-
$\mu^B_{j,y}$	Expected probability to observe a smolt (M2)	-
η^B	Overdispersion of the observation process	Uniform(5,1000)
α^B	Intercept for probability to be observed	N(2.9,0.02)
β^B	Regression coefficient for probability to be observed	logN(-2.6,0.001)
sd_{BB}	Standard deviation of expected probability to be observed	logN(-0.23, 0.005)
Data		
$t_{i,y}$	Mean daily air temperature	-
$f_{j,y}$	Mean daily discharge	-
$N^{Obs}_{j,y}$	Daily smolt count	-

592 Table 3. Results from the predictive study in 2005 (real missing data), 2007 ('first 17% missed' -

593 scenario) and 2014 ('peak of the migration +-2 days missing' -scenario). For posterior distributions,

594 mean values and 90% probability intervals are illustrated.

Year	Sum of missing counts	% of missing counts out of total observed	Number of predicted missing	% of predicted missing out of total
2005	NA	NA	2122 (457,5159)	10% (3%,23%)
2007	2539	17%	3346 (723,8546)	15% (4%,33%)
2014	4085	21%	4677 (2580,7367)	18% (11%,26%)

598 Table 4. Prior and posterior means and 5 % and 95 % quantiles of key parameters

Variable	Prior				Posterior		
	mean	5%	95%		mean	5%	95%
α^P	-20.0	-21.6	-18.3		-19.9	-21.3	-18.5
β^P	1.92	1.09	3.09		1.37	1.25	1.49
s^P	1.65	0.20	5.15		2.89	2.36	3.54
α^D	1.74	1.09	2.61		2.63	2.21	3.09
β^D	0.010	0.007	0.014		0.010	0.007	0.013
cv^D	1.00	0.11	1.90		1.54	1.01	1.95
$cv^{\mu D}$	0.50	0.05	0.95		0.22	0.02	0.55
α^B	2.90	2.69	3.11		2.91	2.70	3.11
β^B	0.074	0.070	0.078		0.074	0.070	0.078
sd_{BB}	0.80	0.71	0.89		0.81	0.72	0.90
η^B	505	56	951		408	15	933
η^N	50092	5008	95069		289	208	403

606   Figure 1: Setup of the video monitoring system at the river bed in Utsjoki. Graph (A) illustrates the

607   profile of the bridge and river bed and graph (B) the river bed in the main channel. Arrows in graph

608   (B) illustrate the position and direction of each of the 8 cameras. Field of view (FoV) of the cameras

609   are: horizontal 78 degrees; vertical 58 degrees.

612
613   Figure 2. Data on daily mean air temperature and mean discharge at Utsjoki on study years 2005-

614   2009 and 2014.

618   Figure 3. Annual prior distributions for the timing of the smolt run given the environmental

619   covariates (daily mean air temperature and discharge).

Annual size of the smolt run

621   Figure 4. Annual number of smolts passing the video site. Posterior distributions are illustrated with

622   boxplots: middle line is median of the distribution, box covers 50% and whiskers 90% of the

623   probability mass. Count data are illustrated with dots.

Figure 5. Daily number of smolts that pass the video site in 2005-2009 and 2014. Posterior

distributions are illustrated with boxplots: middle line is median of the distribution, box covers 50%

and whiskers 90% of the probability mass. Count data are illustrated with dots. Data points included

in the study are marked with black and the ones excluded (counts treated as missing) marked with

grey.

Figure 6. Probability that a smolt begins the migration at given temperature. Priors and posteriors

are illustrated with grey and black boxplots, respectively. Middle line in the box illustrates median of

the distribution, whereas box covers 50% and whiskers 90% of the probability mass.

641
642   Figure 7. Cumulative distribution of expected travel time in days from the point of departure to the

643   video site. Above: Cumulative distribution at low discharge (10 m^3/s). Below: Cumulative distribution

644   at high discharge (100 m^3/s). Priors and posteriors are illustrated with grey and black boxplots,

645   respectively. Middle line in the box illustrates median of the distribution, whereas box covers 50%

646   and whiskers 90% of the probability mass.

653
Figure 8. The expected observation probability at given discharge. Priors and posteriors are

illustrated with grey and black boxplots, respectively. Middle line in the box illustrates median of the

distribution, whereas box covers 50% and whiskers 90% of the probability mass.

Appendix A: Lognormal approximation for Dirichlet-distribution

Let's consider a set of Dirichlet-distributed parameters:

(A.1) $q_{1:n} \sim Dirich(\mu_1\eta, \dots, \mu_n\eta),$

Where μ_i is the expected probability for slot i and η is the overdispersion parameter. Such Dirichlet-

distribution can be approximated with a set of lognormally distributed z_i by assuming

(A.2) $q_{1:n} = z_i / \sum_i z_i$

(A.3) $z_i \sim logN(M_i, T_i)$

(A.4) $M_i = \log(\mu_i) - \frac{0.5}{T_i}$ and

(A.5) $T_i = \log\left(\frac{1}{\mu_i \eta} + 1\right).$

Appendix B: Sensitivity study on priors of the observation process

The expert elicited priors for the parameters of the observation process (Figure 8) did not update

when the model was fitted to video counts. In order to study reasons behind this finding we made

another run with the model with more uncertain prior distributions for parameters α^B and β^B.

Parameter α^B was set to follow N(2.9,1) and parameter β^B logN(-2.6,1) distribution, respectively.

Other prior distributions and model assumptions remained the same as described in the methods

section. Model was run 5 100 000 iterations at which point all parameters of interest had converged

according to Gelman-Rubin shrink factor.

Unlike in the model run with expert elicited priors, the sensitivity model run with wider prior

distributions resulted with update for observability parameters (Figure B1). Parameter α^B updates

only slightly, and most of the learning takes place for parameter β^B. In comparison, in the model run

with expert elicited priors, posteriors of these parameters were identical to the prior distributions

(Figure 8 in Results section). The joint distributions of priors/posteriors of parameters α^B, β^B and

sd^{BB} as a function of discharge (Figure B2) show that learning takes place mainly at the lower ends

of the observation probability.

Joint posteriors from the model run with expert elicited priors and sensitivity model run are

compared in figure B3. The difference in the joint posteriors mainly takes place as increased

uncertainty on higher probabilities to observe smolts, whereas the lower limits of the distributions

are relatively close between the models.

Figure B4 illustrates the impact the prior choice on observability has on the posterior estimates on

the annual total number of smolts. The posteriors are close the same at the upper end of the

distributions, whereas at the lower ends of the distributions the estimates of the sensitivity model

support smaller annual abundances than the estimates from the run with expert elicited priors

(Figure B4). We learn that with the current model structure, the video survey provides mainly

information about the minimum level of observability (through realized discharges), but that expert

knowledge is vital in understanding the maximum limits of observability. Posterior distributions of

parameters related to processes of departing and travelling were identical with the estimates of the

model run with expert elicited priors on observation process (Figures 6 and 7 in the Results section).

Scatter plots of the posterior samples between the annual total number of smolts and parameters of

the observation process, α^B and β^B (Figures B5 and B6), give us more insight on the model behavior.

The interpretation of these parameters is that the α^B sets up the base level for the observation

probability and β^B defines on how quickly the probability decreases as the discharge increases. The

posteriors on total smolt abundance seem to be more sensitive on the choice of prior on α^B than for

choice of prior on β^B, this can be seen especially from the samples of the sensitivity model run

(Figures B5 and B6). On the other hand, observed data has more power to update parameter β^B

than α^B (Figure B1). Correlation between α^B and β^B balances the joint distribution in the sensitivity

model run, as with high values of α^B values of $-\beta^B$ decrease (Figure B7).

707
Figure B1. Marginal densities of parameters α^B and β^B in the sensitivity model run. Prior

distributions are illustrated with dotted and posterior distributions with solid lines.

Figure B2. The expected observation probability at given discharge in sensitivity model run. Priors

and posteriors are illustrated with grey and black boxplots, respectively.

Figure B3. The expected observation posterior probability at given discharge in model run with

expert elicited priors (grey box plots) and sensitivity model run (black boxplots).

Figure B4. Posterior estimates of annual number of smolts passing the video site based on model

run with expert elicited priors (grey boxplots) and priors from sensitivity analysis (white boxplots).

Middle line is the median of the distribution; box covers 50% and whiskers 90% of the probability

mass. Count data are illustrated with dots.

Figure B5. Scatter plots of sampled posterior values between annual total numer of smolts (in

thousands) and parameter α^B of the observation process. Dark grey dots illustrate the posterior

samples from the sensitivity analysis, whereas light grey dots illustrate the posterior samples from

the analysis with expert elicited priors.

Figure B6. Scatter plots of sampled posterior values between annual total numer of smolts (in

thousands) and parameter β^B of the observation process. Dark grey dots illustrate the posterior

samples from the sensitivity analysis, whereas light grey dots illustrate the posterior samples from

the analysis with expert elicited priors.

Figure B7. Scatter plots of sampled posterior of parameters α^B and β^B of the observation process.

Dark grey dots illustrate the posterior samples from the sensitivity analysis, whereas light grey dots

illustrate the posterior samples from the analysis with expert elicited priors.

Appendix C: JAGS code for the arrival model

model{

 # Process of departing

 # =========================

for(y in 1:nYears){

for(i in 1:nDays){

 # p: probability to start migration at day t, if haven't done so earlier

 # t: daily mean temperature

logit(p[i,y])<-p2[i,y]

p2[i,y]~dnorm(aP+bP*t[i,y],1/pow(sdP,2))

```
751        }

752      }

753      # P: expected proportion of smolts departing at day i

754      for(y in 1:nYears){

755        P[1,y]<-p[1,y]

756        for(i in 2:(nDays-1)){

757          P[i,y]<-(1-sum(P[1:(i-1),y]))*p[i,y]

758        }

759        P[nDays,y]<-1-sum(P[1:60,y])

760      }

761

762      aP~dnorm(-20,1)

763      bP~dlnorm(0.6,10)

764      sdP~dlnorm(0,1)

765

766      # Process of travelling

767      # =============

768      # qD: probability that a smolt that departs at day i passes the video site at day j

769      for(y in 1:nYears){

770        for(i in 1:nDays){

771          # i: day of departure

772          # j: day of passing the video site

773          # f: daily mean discharge

774          # j==i

775          qDx[i,i,y]<-phi((log(0.5)-MD[i,y])/SD)

776          # j>i

777          for(j in (i+1):(i+13)){

778            qDx[i,j,y]<-phi((log(j-i+0.5)-MD[i,y])/SD)-phi((log(j-i-0.5)-MD[i,y])/SD)

779          }
```

```
780        for(j in i:(i+13)){
781          qD[i,j,y]<-qDx[i,j,y]/(sum(qDx[i,i:(i+13),y])+0.0001)
782        }
783        MD[i,y]<-log(muD[i,y])-0.5/TD
784        muD[i,y]~dlnorm(log(exp(aD-bD*f[i,y]))-0.5/TmuD, TmuD)
785      }
786    }
787    SD<-1/sqrt(TD)
788    TmuD<-1/log(cvmuD*cvmuD+1)
789    TD<-1/log(cvD*cvD+1)
790
791    aD~dlnorm(0.52,14)
792    bD~dlnorm(-4.6,25)
793    cvmuD~dunif(0.001,1)
794    cvD~dunif(0.001,2)
795
796    # Arrival distribution
797    # ============================================
798    # Joint distribution of qD and P: prob that a smolt departing day i arrives day j
799    for(y in 1:nYears){
800      for(i in 1:nDays){ # day of departure
801        for(j in i:(i+13)){ # day of passing the video site
802          qJ[j,i,y]<-qD[i,j,y]*P[i,y]
803        }
804      }
805
806    # Expected proportion of smolts passing the video site each day
807    for(j in 1:14){
808      muqN[j,y]<-sum(qJ[j,1:j,y])+0.0001 # small constant added for numerical stability
```

```
809          }
810        for(j in 15:nDays){
811           muqN[j,y]<-sum(qJ[j,(j-13):j,y])+0.0001
812        }
813      }

815    for(y in 1:nYears){
816      Ntot[y]<-exp(LNtot[y]) # total run size in year y
817      LNtot[y]~dunif(7,15)

819      for(j in 1:(nDays-1)){
820        N[j,y]<-round(qN[j,y]*Ntot[y]) # daily number of fish passing
821      }
822      N[nDays,y]<-round(Ntot[y]*(1-sum(qN[1:(nDays-1),y])))

824      # qN: daily proportion of smolts
825      # dirichlet-distribution approximated with lognormal
826      qN[1:nDays,y]<-zN[1:nDays,y]/sum(zN[1:nDays,y])

828      for(i in 1:nDays){
829        zN[j,y]~dlnorm(MN[j,y], TauN[j,y])
830      }

832      alphaN[1:nDays,y]<-muqN[1:nDays,y]*eta_alphaN
833      MN[1:nDays,y]<-log(muqN[1:nDays,y])-0.5/TauN[1:nDays,y]
834      TauN[1:nDays,y]<-1/log((1/alphaN[1:nDays,y])+1)
835    }

837    eta_alphaN~dunif(0.001,100000)
```

```
838
839    # Observation process
840    # ====================
841    for(y in 1:nYears){
842      for(j in 1:nDays){
843        # Nobs: Observed number of smolts
844        Nobs[j,y]~dbetabin(100,10,N[j,y])  # M1
845        Nobs[j,y]~dbetabin(muB[j,y]*etaB,(1-muB[j,y])*etaB,N[j,y])  #M2
846
847        muB[j,y]<-0.6*(exp(BB[j,y])/(1+exp(BB[j,y])))+0.3
848        BB[j,y]~dnorm(aB-bB*f[j,y],1/pow(sdBB,2))
849
850      }
851    }
852    # priors for observation process M2
853    aB~dnorm(2.9,60)
854    bB~dlnorm(-2.6,984)
855    sdBB~dlnorm(-0.23,210)
856    etaB~dunif(5,1000)
857  }
858
```