Bayesian arrival model for Atlantic salmon smolt counts powered by environmental covariates and expert knowledge

Henni Pulkkinen, Panu Orell, Jaakko Erkinaro, Samu Mäntyniemi

H. Pulkkinen.* Natural Resources Institute Finland, Paavo Havaksen tie 3 P.O. Box 413, 90014-Oulu. Finland. henni.pulkkinen@luke.fi

P. Orell. Natural Resources Institute Finland, Paavo Havaksen tie 3 P.O. Box 413, 90014-Oulu. Finland. panu.orell@luke.fi

J. Erkinaro. Natural Resources Institute Finland, Paavo Havaksen tie 3 P.O. Box 413, 90014-Oulu. Finland. jaakko.erkinaro@luke.fi

S. Mäntyniemi. Natural Resources Institute Finland, Latokartanonkaari 9, 00790 Helsinki. Finland. samu.mantyniemi@luke.fi

Corresponding author: Henni Pulkkinen (phone: +358 40 485 0088)

Abstract

Annual run size and timing of Atlantic salmon smolt migration was estimated using Bayesian model framework and data from six years of a video monitoring survey. The model has a modular structure. It separates sub-processes of departing, traveling and observing, of which the first two together define the arrival distribution. The sub-processes utilize biological background and expert knowledge about the migratory behavior of smolts and about the probability to observe them from the video footage under varying environmental conditions. Daily mean temperature and discharge were used as environmental covariates. The model framework does not require assuming a simple distributional shape for the arrival dynamics and thus also allows for multimodal arrival distributions. Results indicate that 25 % - 75% of smolts pass the Utsjoki monitoring site unobserved. Predictive studies were made to estimate daily run size in cases with missing counts either at the beginning or in the middle of the run, indicating good predictive performance.

Keywords: salmon, arrival model, passage count, Bayesian models, environmental covariates, expert knowledge, biological realism, missing data, prediction

Introduction

Migratory fish species are often monitored along their migration routes at different life stages to collect fishery-independent data for stock assessment. Monitoring methods include e.g. counting fences, video, sonar and snorkeling counts, and common to all is that they rarely provide perfect information about the number of individuals passing the system (Dempson et al. 1991; Romakkaniemi et al. 2000; Orell et al. 2007; 2011). Part of the fish run may be missed because of difficult environmental conditions or device failures, there may be double counting, or the data

achieved may be otherwise partial or biased (e.g. Holmes et al. 2006). Thus, the assumptions made

when interpreting the data can have a great influence on the estimated stock abundance.

Increasing the information beyond the actual monitoring data collected can improve the

trustworthiness of stock assessments (Kuparinen et al. 2012): such information can be based on the

resources available in other fields of biological research such as ecology or life history theory, but

also on the physical and technical details of the observation processes. Methods of Bayesian

inference enable both estimation of uncertainty and combination of various sources of information.

Statistical methods are often used, however, without considering the biological realism and just

focusing on statistical data analysis. Such a procedure easily leads to awkward model assumptions

that cannot be biologically interpreted and the potential effects of those modelling decisions may

pass undiscussed.

During the course of time, various methods have been used to estimate run dynamics and passage

counts at point census stations. The simplest methods are based on "connect-the-dots" type of

linear interpolation (Gewin and WanHatten 2005, Johnson et al. 2007) but such approach requires

an observation before and after the missing time point and thus missing tails cannot be estimated. If

it is assumed that a proportion passing the site on a given date is constant over the years then

estimation of missing tails is possible using the methods proposed by, e.g, Van Alen (2000), but

considering the variability, for example, in annual environmental conditions (e.g. Erkinaro et al.

1998; Orell et al. 2007; Otero et al. 2014), such assumption is problematic.

Hilborn et al. (1999) used a maximum likelihood method to estimate number of salmon with ground-

based stream survey, accounting also for estimate of uncertainty. Su et al. (2001) continued their

work with hierarchical Bayesian approach enabling learning from years with more data to those with

missing data. Furthermore, Sethi and Bradley (2016) introduced a Bayesian approach to estimate

missing passage at weirs with run curve model to account for arrival dynamics and process variation

model to describe the observed data. While these studies attempt to account for uncertainty, they

do not discuss their model assumptions against biological knowledge. This together with

consideration of difference between the observation and model prediction as "noise" instead of

proportion unobserved (as e.g. in Sethi and Bradley 2016), may cast some doubt on the biological

meaningfulness of the results.

In our study, a Bayesian model framework was introduced for estimating the annual number of

Atlantic salmon (*Salmo salar*) smolts passing the video monitoring site at a subarctic River Utsjoki in

northern Finland. The framework consists of sub-processes of departing, travelling and observing.

When combined, the processes of departing and travelling form the arrival distribution that

describes the timing of the smolt run. Environmental covariates were used as a source of

information in the sub-processes and expert knowledge is utilized in providing biologically

meaningful model structures and prior distributions. Hierarchical structure was assumed over the

study years making it possible to learn from the processes, borrowing strength from data rich years

of study to those with missing data. Predictive studies illustrate the model performance in cases with

missing data either at the beginning or in the middle of the run. We show that a biologically

meaningful passage model can be built without assuming any specific shape for the arrival

distribution. Furthermore, we claim that expert knowledge is a key source of information in

estimating the total uncertainty in the annual total passage especially when environmental

conditions have a great influence on the probability to observe individuals passing the survey site.

Materials and Methods

Data

River Utsjoki is a tributary of the large River Teno system at the northernmost border between

Finland and Norway. Each spring, the monitoring system of 8 video cameras is set at the river bed

under a bridge close to the tributary outlet (see Orell et al. 2011; Fig 1). The video footage provides

data on both smolts descending and adults ascending the river during the course of summer. In this

study we analyze the smolt count data from years 2005-2009 and 2014. These years were chosen to

demonstrate the highly variable environmental conditions, especially the strongly varying discharge

(Figure 2). The smolt data are aggregated in daily counts of individuals over 61 days of June-July.

Data on environmental covariates contain air temperature and discharge measurements. These

datasets are aggregated into daily averages. Both covariates are measured near the video site, but

conditions may differ between the video site and upper parts of the river from where some of the

smolts begin their migration. These datasets are, however, considered as reasonable proxies for the

environmental conditions affecting the migration behavior of smolts.

Data from the first 23 days of June in 2005 are missing because the high level of water delayed the

setup of the monitoring system in that year. Thus, the missing counts are predicted for those dates.

Predictive studies were performed by treating data in 2007 and 2014 as partly missing: For 2007 the

first 17% of the run strength and for 2014 the peak +-2 days count was removed from the analysis.

To evaluate the predictive performance of the model, the model predicted counts for those dates

were compared with the observed data.

Expert elicitation as a source of informative priors

Model structure and informative priors were elicited from an expert that is most familiar with the

behavior of salmon smolts and the video monitoring system at Utsjoki. The expert was asked to base

his views on his background knowledge about the processes instead of considering the particular

data that will be included in the study. Elicitation was carried out iteratively, first by asking

preliminary questions on, e.g. biological process in question, and then providing graphical illustration

of the parameters. Prior distributions were adjusted until the expert agreed that the illustration is in

line with his views. Some examples of questions elicited from the expert are listed in table 1.

Probability model of the problem

The model is constructed by thinking of four steps, described in detail in the following sections:

1) Process of departing, in which the smolts make the decision to depart while daily

temperature affects the probability to depart in a given day;

2) Process of travelling, in which the smolt's travel time (in days) from the place of departure to

the video site is modeled with daily discharge as a covariate;

3) Arrival distribution, which links together the processes of departing and travelling into a

timing of the smolt run;

4) Observation process, in which the discharge influences visibility and the probability that an

individual smolt passing the site is observed from the video footage.

Process of departing

An individual smolt's probability to begin the migration in day i of year y given that the smolt has not

departed yet is considered to depend on the temperature on that day. We assume a logit-normally

linear relationship for the smolt's probability to depart $(p_{i,y})$ and temperature $t_{i,y}$:

(1) $logit(p_{i,y}) \sim N(\alpha^P + \beta^P \, t_{i,y} \, , \, s^P)$

Informative prior distributions are given to parameters α^P, β^P and s^P according to expert view (see

tables 1 and 2).

The daily probabilities of departing (given those individuals have not departed yet; $p_{i,y}$) can be

transformed into expected proportions of the total smolt run that depart each day $(P_{i,y})$ as

(2) $P_{1,y} = p_{1,y}$

(3) $P_{i,y} = (1 - \sum_{k=1}^{i-1} p_{k,y}) \, p_{i,y}$

(4) $P_{61,y} = 1 - \sum_{i=1}^{60} p_{i,y}$.

Process of travelling

According to the expert view, each smolt should arrive at the video site within 14 days after

departing. Thus we assume that the smolts that depart on day i pass the video site in one of the next

14 days according to discretized distribution function of lognormal distribution (Schwarz & Dempson

1994, Mäntyniemi & Romakkaniemi 2002):

$(5)\ Q_{i,j,y}^{D} = \phi\left(\frac{\log(j-i+0.5-M_{i,y})}{S}\right) - I_{i,j}\phi\left(\frac{\log(j-i+0.5-M_{i,y})}{S}\right).$

Here ϕ is the cumulative density function of the standard normal distribution, $M_{i,y}$ is the mean of

log(travel time) to video site of smolts that depart on day i , and S is the standard deviation of

log(travel time). Thus, $Q_{i,j,y}^{D}$ is the conditional probability that a smolt that departed on day i will

arrive on day j, when j goes from i to $i + 13$.

To ensure that all smolts arrive within 14 days of departing, $Q_{i,j,y}^{D}$'s are standardized:

$(6)\ q_{i,j,y}^{D} = Q_{i,j,y}^{D} / \sum_{j=i}^{(i+13)}(Q_{i,j,y}^{D}),$

and thus $\sum_{j=i}^{(i+13)}(q_{i,j,y}^{D}) = 1$.

It is further considered that increasing discharge increases the speed at which the smolts travel,

shortening their travel time. The expected travel time on real scale $\mu_{j,y}^{D} = \exp\left(M_{i,y} + 0.5\,S^2\right)$ is

assumed to follow a lognormal distribution with expected value $\gamma_{i,y}$ and coefficient of variation cv^{D}

and it is assumed to depend on the discharge $(f_{i,y})$:

$(7)\ \gamma_{i,y} = \exp\left(\alpha^{D} + \beta^{D} \cdot f_{i,y}\right).$

Again, informative prior distributions are given for parameters α^D, β^D and $cv^{\mu D}$ according to expert

view (see tables 1 and 2).

Arrival distribution

By combining the processes of departing and travelling, we achieve joint probabilities that a smolt

departs on day i and passes the video site on day j:

(8) $q^J_{i,j,y} = q^D_{i,j,y}\, P_{i,y}.$

The expected proportions of smolts (out of the total run size) passing the video site each day can

then be found by marginalizing over the day of departure:

(9) $\mu^N_{j,y} = \begin{cases} \sum q^J_{j,1:j,y} \ , \ j \leq 14 \\ \sum q^J_{j,(i-13):j,y} \ , j \ > 14. \end{cases}$

Furthermore, we give a uniform prior distribution on log scale for the total annual size of the smolt

run passing the video site:

(10) $\log (N^{tot}_y) \sim Unif(7,15).$

Parameterisation on the log-scale allows for a wide, minimally informative prior distribution for the

annual total abundance. This annual run size is assumed to be Dirichlet-multinomially distributed

over the 61 days of June-July. For computational simplicity, we follow Mäntyniemi et al. (2015) and

approximate multinomial distribution as

(11) $N_{j,y} = q^N_{j,y}\, N^{tot}_y,$

where $q_{j,y}^{N}$ is the proportion of smolts passing the video site in day j and $q_{j,y}^{N}$'s follow a Dirichlet

distribution parameterized with the expected proportions of smolts passing the video site each day

$\mu_{j,y}^{N}$ and overdispersion η^{N}:

(12) $q_{1:61,y}^{N} \sim Dirich(\mu_{1,y}^{N}\eta^{N}, \dots, \mu_{61,y}^{N}\eta^{N})$.

To ease the computation we approximate the Dirichlet-distribution with a set of lognormal

distributions (see appendix A for details).

The annual proportional prior distributions for the the smolt run over 61 days of June-July are

illustrated in figure 3. These priors are based on the model structure and data on environmental

covariates (daily mean air temperature and discharge).

Observation process

In previous sections we have covered the processes assumed to affect the timing of the smolt

passage at the video site. To estimate the total uncertainty in the smolt run size, the process of

observing smolts from the video footage needs to be included. It seems natural to assume that some

of the individuals pass the site unobserved, and thus this proportion must be accounted for. Next,

we introduce two versions for the observation process:

The first version (model M1) is simple yet not very realistic. It assumes that the number of smolts

observed at the video site on day i in year y ($N_{i,y}^{Obs}$) follows a Beta-binomial distribution:

(13) $N_{j,y}^{Obs} \sim Beta - bin(100,10, N_{i,y})$

Here $N_{j,y}$ is the total daily number of smolts passing the video site and numerical constants indicate

that the probability that an individual smolt is observed varies between days according to

$Beta(100,10)$ distribution with fixed mean 0.91 and standard deviation 0.03. Thus, 9% of smolts are

assumed to pass the video site unobserved on average, with 95% PI of [5%, 14%]. Albeit an

observation model with randomly varying probability for observing is not realistic, such a model can

be useful for illustrating how the processes of departing and travelling affect the model results when

considering assuming close to perfect process of observing.

The second version (model M2) for an observation process aims at higher realism, accounting for

expert views on how high the probability to observe a smolt may be in excellent vs. poor

circumstances and how the discharge would affect this probability. Again the number of smolts

observed follows a Beta-binomial distribution:

(14) $N_{j,y}^{Obs} \sim Beta-bin(\mu_{j,y}^B \eta^B,\ (1-\mu_{j,y}^B)\eta^B,\ N_{j,y})$

Now the probability to observe a smolt follows a $Beta(\mu_{j,y}^B \eta^B,\ (1-\mu_{j,y}^B)\eta^B)$ distribution, where

$\mu_{j,y}^B$ is the expected probability for a smolt to be observed and η^B is the overdispersion parameter.

Overdispersion is thought to arise from the schooling behavior of smolts. Groups of smolts can be

either detected or missed, which increases the variance of the observation process (Mäntyniemi &

Romakkaniemi 2002, Linden & Mäntyniemi 2011, Mäntyniemi et al. 2015). The expected value of the

process is linked to discharge following the expert view that in very good visibility (low discharge) at

maximum 90% of the smolts are observed. As the discharge increases, the visibility decreases, and

the volume of water to be covered by the video monitoring increases. Therefore, according to the

expert, observation probability decreases gradually to a minimum of 30%. The expected probability

$\mu_{i,y}^B$ is considered to follow logit-normally linear relationship on the interval [0.3,0.9]:

(15) $\mu_{j,y}^B = 0.6\left(\dfrac{\exp(BB_{j,y})}{1+\exp(BB_{j,y})}\right) + 0.3$

(16) $BB_{j,y} \sim N(\alpha^B - \beta^B f_{j,y},\ sd_{BB}^2),$

where α^B, β^B and sd^2_{BB} have prior distributions according to the expert view (Tables 1 and 2).

Models M1 and M2 were fitted using MCMC sampling with JAGS 4.2.0 software (Just Another Gibbs

Sampler; Plummer 2003) and R runjags package (Denwood 2016). Both models were run with 2

chains for 1 500 000 iterations this taking around 3 days per model. Convergence was diagnosed

using Gelman-Rubin statistics (Gelman and Rubin 1992). The complete JAGS code for the model

structure can be found from appendix B.

Results

When compared with the observed smolt counts, annual posterior estimates of daily run size (based

on M2) indicate that 25-75% of smolts pass the video site unobserved (Figure 4). Considering the

estimated uncertainty among the years without missing data, the proportion of unobserved is 90%

at most (in 2008 upper limit for 90% PI is 16000 smolts vs. observed data of 8400 smolts). Posterior

estimates of daily arrivals are illustrated in figures 5 (M1) and 6 (M2).

Table 3 shows the posteriors for the time periods with missing counts as well as the observed annual

counts for comparison when available. M2 predicts 24-33% higher number of smolts compared to

M1 as a result of more realistic observation model. Scenarios to study predictive performance of the

model contain 'first 17% of missing' -scenario for 2007 and 'peak day +- 2 days missing' –scenario for

2014. In both cases, all daily counts fall in the predicted 90% PI indicating good predictive

performance (Table 3). The percentage of predicted out of the estimated annual total is close to

identical between models in all 3 cases (Table 3).

Figures 7 and 8 illustrate the prior and posterior distributions from M2 for expected processes of

departing and travelling. Posterior distributions from M1 are nearly identical and thus not illustrated

here. The expected probability to begin the migration at given temperature updates quite heavily

from the prior distributions, supporting 4-5 degrees higher temperature at which the migration sets

off (Figure 7). The posterior cumulative distribution of the expected travel time to the video site

(Figure 8) supports somewhat longer travel times than expected a priori both for low ($10m^3/s$) and

high ($100m^3/s$) discharge. Note that posterior cumulative distributions at these specific levels of

discharge are chosen for illustrative purposes only and that in practice the models allow any positive

value for the discharge.

In M2, also the observation probability depends on the discharge. Figure 9 illustrates the prior and

posterior distributions of the observation process given the discharge. Priors and posteriors are close

the same, indicating that in this case the combination of priors and data did not result in further

learning about the process of observing.

Discussion

In our study, an arrival model for Atlantic salmon smolts was designed by separating the processes

of departing, travelling and observing. The processes of departing and travelling together form the

arrival distribution, i.e. the timing of the smolt run. This modular structure enabled us to concentrate

on biologically meaningful mathematical assumptions related to each sub-process instead of

assuming that fish count data as such follows a specific distributional shape. The model framework

aimed at high biological realism by accounting for expert knowledge in the model formulation, firstly

by including choices for model structures and prior distributions, and secondly by linking

environmental covariates to provide information about the timing of the migration. The estimated

total uncertainty accounted for the uncertainty in the sub-processes, including the observation

process in which variability in discharge is known to have a large impact on the probability to

observe passing smolts.

One of the most important advantages of our approach is that the need to assume a specific

parametric shape for the arrival distribution is avoided. Past studies (Hilborn et al. 1999, Su et al.

2001, Sethi and Bradley 2016) have assumed unimodal shape (e.g. normal, skew-normal, Student's t)

for the arrival curve. However, when taking a look at the shape of (any) arrival time series it seems

clear that such assumption is a tolerable simplification at best (see e.g. figure 5 in this paper, figure 2

in Sethi and Bradley 2016).

Earlier studies have paid little attention to the role of observation models in estimating the total

uncertainty. Hillborn et al. (1999) distinguished process and observation errors, assuming observer

efficiency either as known or unchanging inside the study period. Su et al. (2001) discussed the issue

of error in the observation process and mainly concluded that such error can cause convergence

problems in sampling. Sethi and Bradly (2016) incorporated uncertainty only on the estimates of

missing passage counts, treating the observed counts as true total number of passing without

uncertainty. As they assumed normal or negative Binomial models for the observed passage counts,

they in fact allowed model predicted passage counts to be either greater or smaller than the

observed counts. This in turn can lead to underestimation of the missing counts. We claim that it is

necessary to consider the observed count as a minimum estimate for the true number especially if

the back-and-forth movement of individuals is negligible. Along with this notion the need for a

realistic observation process becomes obvious. Depending on the approach, the estimate of total

annual passage can be very different. As Hilborn et al. (1999) put it, 'when we admit uncertainty in

observer efficiency we become much less certain about the actual escapement'.

Sethi and Bradley (2016) fitted datasets in their study using several combinations of arrival and

process error models. They utilized information-theoretic model selection (DIC, Spiegelhalter et al.

2002) to find out the best fitting model for each dataset. However, no discussion took place whether

some of these models would be biologically more reasonable than another. We argue that the

model selection should primarily be based on biological realism and in case biological model

uncertainty is needed to be tackled, Bayesian model averaging (BMA, Hoeting et al. 1999) should be

preferred over DIC or other information criteria. The purpose of an information criterion is to select

a statistical model that would be able to replicate the observed data with least amount of

parameters. Such a goal does not necessitate an ecological model: generic machine learning models

are probably going to perform much better in this regard.

Su et al. (2001) used hierarchical structure in their study, assuming that years in which timing of the

run is well known can be used for passing information to years with missing data. We argue that

learning should rather take place over the (biological) process, assuming exchangeability over year

specific parameters of those processes. Our model structure can also be considered as hierarchical,

although we simply assume that the underlying processes are the same from year to year allowing

between- and within-years variation. If there were a need for a meta-analysis of similar studies over

several stocks of the same species, exchangeability in biological parameters could be considered

between the stocks. Such approach could also allow meta-analysis of data from different types of

monitoring systems if detailed observation processes were tailored for each system and

exchangeability was considered as reasonable assumption for the biological processes.

Our model framework could be fine-tuned and developed further. In an ideal situation,

environmental covariates would be available from upstream locations from where the smolts

originate. Data on air temperature would be replaced with water temperature to better explain the

physiological processes the smolts encounter. To increase the realism even more, the schooling

behavior of smolts could be accounted for in more detail than done here. Salmon smolts have a

tendency to school during their migration (e.g. Bakshtanskiy et al. 1988, Davidsen et al. 2005) and

this creates more variation in the monitoring data compared to independent movement

(Mäntyniemi and Romakkaniemi 2002, Mäntyniemi et al. 2015). Schooling behaviour could also be

linked with the process of travelling as the passing schools of fish have been suggested to stimulate

migratory behavior in the remaining smolts (e.g. Hansen and Jonsson, 1985). Schooling also affects

the observation process as, on one hand, schools are easier to spot than individual smolts, and on

the other hand, it can be difficult to have an exact count of individuals from a large school. For

Utsjoki video survey, daily average school sizes of smolts are recorded and incorporating those into

the model framework is an interesting topic for future research.

Despite the above possibilities for further development of the modeling framework, we claim that

our model already contains the elements of realism that are required to provide credible estimates

of the total annual smolt run. This approach can be applied to similar monitoring surveys where

point-checking the numbers of migrating fish (Trépanier et al. 1996), by e.g. sonar, trap, or weir is

used for estimating run sizes. As far as we know, environmental covariates have not been included

so far to estimate both the dynamics of passage and the observation process. Our model framework

also indicated a good predictive performance for missing dates both at the beginning and in the

middle of the run.

Acknowledgements

329 *We thank the numerous field workers enabling the yearly monitoring programme and smolt data*

330 *collection at the River Utsjoki. We also acknowledge the crucial technical knowhow by Jorma Kuusela*

331 *in running the annual monitoring and the instrumental role of Dr. Martin Svenning and Anders*

332 *Lamberg in early phases of the River Utsjoki monitoring programme. Funding for this study was*

333 *provided by the Academy of Finland (project No. 286334).*

References

Bakshtanskiy, E., Nesterov, V., and Neklyodov, M. 1988. Development of schooling behaviour in

juvenile Atlantic salmon, Salmo salar, during seaward migration. J. Ichthyol. 28: 91–101.

Davidsen, J., Svenning, M. A., Orell, P., Yoccoz, N., Dempson, J. B., Niemelä, E., Klemetsen, A.,

Lamberg, A., and Erkinaro, J. 2005. Spatial and temporal migration of wild Atlantic salmon smolts

determined from a video camera array in the sub-Arctic River Tana. Fisheries Research 74: 210-222.

Dempson, J. B, and Stansbury, D. E. 1991. Using partial counting fences and atwo-sample stratified

design for mark–recapture estimation of an Atlantic salmonsmolt population. N. Am. J. Fish.

Manage. 11: 27–37.

Denwood, M.J. 2016. runjags: An R Package Providing Interface Utilities, Model Templates, Parallel

Computing Methods and Additional Distributions for MCMC Models in JAGS. J. Stat. Softw. 71(9): 1-

25. doi:10.18637/jss.v071.i09.

Erkinaro, J., Julkunen, M. and Niemelä, E. 1998. Migration of juvenile Atlantic salmon Salmo salar in

small tributaries of the subarctic River Teno, northern Finland. Aquaculture 168: 105-119.

Gelman, A., and Rubin, D.B. 1992. Inference from iterative simulation using multiple sequences. Stat.

Sci. 7: 457–511.

Hilborn, R., Bue, B.G., and Sharr, S. 1999. Estimating spawning escapements from periodic counts: a

comparison of methods. Can. J. Fish. Aquat. Sci. 56(5): 888–896. doi:10.1139/f99-013.

Hansen, L.P., and Jonsson, B. 1985. Downstream migration of hatchery-reared smolts of Atlantic

salmon (Salmo salar L.) in the River Imsa. Aquaculture, 45: 237–248.

Holmes, J.A, Cronkite, G.M.W., Enzenhofer, H.J., and Mulligan, T.J. 2006. Accuracy and precision of

fish-count data from a "dual-frequency identification sonar" (DIDSON) imaging system. ICES J. Mar.

Sci. 63: 543–555. doi:10.1016/j.icesjms.2005.08.015

Kuparinen, A., Mäntyniemi, S., Hutchings, J.A., and Kuikka, S. 2012. Increasing biological realism of

fisheries stock assessment: towards hierarchical Bayesian methods. Environ. Rev. 20: 135–151.

doi:10.1139/A2012-006.

Mäntyniemi, S., and Romakkaniemi, A. 2002. Bayesian mark-recapture estimation with an

application to a salmonid smolt population. Can. J. Fish. Aquat. Sci. 59: 1748-1758. doi:10.1139/F02-

146.

Mäntyniemi, S.H.P., Whitlock, R.E., Perälä, T.A., Blomstedt, P.A., Vanhatalo, J.P., Rincon, M.M.,

Kuparinen, A.K., Pulkkinen, H.P., and Kuikka, O.S. 2015. General state-space population dynamics

model for Bayesian stock assessment. ICES J. Mar. Sci. 72: 2209-2222. doi:10.1093/icesjms/fsv117

Orell, P., Erkinaro, J., Svenning, M., Davidsen, J., and Niemelä, E. 2007. Synchrony in the

downstream migration of smolts and upstream migration of adult Atlantic salmon in the sub-Arctic

River Utsjoki. J. Fish. Biol. 71: 1735–1750.

Orell, P., Erkinaro, J., and Karppinen, P. 2011. Accuracy of snorkelling counts in assessing spawning

stock of Atlantic salmon, Salmo salar, as verified by radio-tagged fish and underwater video

monitoring. Fisheries. Manag. Ecol. 18: 392–399.

Otero, J., L'Abée-Lund, J.H., Castro-Santos, T., Leonardsson, K., Storvik, G.O., Jonsson, B., Dempson,

B., Russell, I.C., Jensen, A.J., Baglinière, J.-L., Dionne, M., Armstrong, J.D., Romakkaniemi, A., Letcher,

B.H., Kocik, J.F., Erkinaro, J., Poole, R., Rogan, G., Lundqvist, H., MacLean, J.C., Jokikokko, E.,

Arnekleiv, J.V., Kennedy, R.J., Niemelä, E., Caballero, P., Music, P.A., Antonsson, T., Gudjonsson, S.,

Veselov, A.E. Lamberg, A., Groom, S., Taylor, B.H., Taberner, M., Dillane, M., Arnason, F., Horton, G.,

Hvidsten, N.A., Jonsson, I.R., Jonsson, N., McKelvey, S., Næsje, T., Skaala, Ø., Smith, G.W., Sægrov, H.,

Stenseth, N.C. and Vøllestad, L.A. 2014. Basin-scale phenology and effects of climate variability on

global timing of initial seaward migration of Atlantic salmon (Salmo salar). Glob. Change. Biol. 20: 61-

75.

Plummer, M. 2003. JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling.

In proceedings of the 3[rd] International Workshop on Distributed Statistical Computing (DSC 2003),

20-22 March, Vienna, Austria. ISSN 1609-395X.

Romakkaniemi, A., Lilja, J., Nykänen, M., Marjomäki, T.J. and Jurvelius, J. 2000. Spawning run of

Atlantic Salmon (Salmo salar) in the River Tornionjoki monitored by horizontal split-beam

echosounding. Aquat. Living. Resour. 13: 349-354.

Schwarz, C., and Dempson, J. 1994. Mark-recapture estimation of a salmon smolt population.

Biometrics. 50: 98-108.

Sethi, A.S., and Bradley, C. 2016. Statistical arrival models to estimate missed passage counts at

fish weirs. Can. J. Fish. Aquat. Sci. 73: 1251–1260. doi:10.1139/cjfas-2015-0318.

Spiegelhalter, D.J., Best, N.G., Carlin, B.P., and van der Linde, A. 2002. Bayesian measures of model

complexity and fit (with discussion). J. R. Stat. Soc. Ser. B. 64: 583–639.

Su, Z., Adkison, M.D., and Van Alen, B.W. 2001. A hierarchical Bayesian model for estimating

historical salmon escapement and escapement timing. Can. J. Fish. Aquat. Sci. 58: 1648–1662.

doi:10.1139/f01-099.

Trépanier, S., Rodríquez, M. A. and Magnan, P. 1996. Spawning migrations in landlocked Atlantic

salmon: time series modelling of river discharge and water temperature effects. J. Fish Biol. 48: 925–

936.

Table 1. Examples of questions elicited from the expert

Question	Expert view
Process of departing	
- At what range of temperature (air temperature at Utsjoki logger) first smolts are considered to begin their migration?	6-8 °C
- What level of temperature would be considered so high that the process of migration not be influenced by further increase in temperature (i.e. logit curve levels out)?	12 °C
Process of travelling	
- What is the the maximum travel time (in days) in which a smolt can be considered to arrive at the video site?	14 days
- What is the minimum travel time in which a smolt can be considered to arrive at the video site?	1-2 days
Process of observing	
- What is the upper (/lower) limit for the expected probability at which an average smolt is considered to be observed in best (/worst) possible environmental conditions, i.e. at minimum (/maximum) discharge?	90% (30%)
- What is the level of discharge at which the visibility can be considered optimal?	20 m^3/s
- What is the level of discharge at which the visibility is considered to be poorest?	50-60 m^3/s

 Table 2: List of symbols, descriptions and prior distributions

Symbol	Description	Prior distribution
$logN_y^{tot}$	Total annual run size on log-scale	Uniform(7,15)
N_y^{tot}	Total annual run size on real scale	-
$N_{j,y}$	Daily number of smolts passing the video site	-
$q_{j,y}^N$	Daily proportion of smolts passing the video site	-
$\mu_{j,y}^N$	Expected proportion of smolts passing the video site	-
η^N	Overdispersion of the arrival process	Uniform(0.001,100000)
$p_{i,y}$	Probability to depart (if haven't departed yet)	-
$P_{i,y}$	Proportion of smolts departing each day	-
α^P	Intercept for probability to depart given temperature	N(-20,1)
β^P	Regression coefficient for probability to depart given temperature	logN(0.6,0.1)
s^P	Standard deviation of probability to depart	logN(0,1)
$Q_{i,j,y}^D$	Unstandardised proportion for travel time	-
$q_{i,j,y}^D$	Standardised proportion for travel time	-
$M_{i,y}$	Mean of log(travel time)	-
S	Standard deviation of log(travel time)	
$\mu_{i,y}^D$	Expected travel time	-
$\gamma_{i,y}$	Expected value of expected travel time	-
cv^D	Coefficient of variation of log(travel time)	Uniform(0.001,2)
α^D	Intercept for the expected value of expected travel time given flow	logN(0.52,0.07)
β^D	Regression coefficient for the expected value of expected	logN(-4.6,0.04)

	travel time given flow	
$cv^{\mu D}$	Coefficient of variation of expected travel time	Uniform(0.001,1)
$q^{J}_{i,j,y}$	Probability that a smolt departs on day i and passes the video site on day j	-
$\mu^{B}_{j,y}$	Expected probability to observe a smolt (M2)	-
η^{B}	Overdispersion of the observation process	Uniform(5,1000)
α^{B}	Intercept for probability to be observed	N(2.9,0.02)
β^{B}	Regression coefficient for probability to be observed	logN(-2.6,0.001)
sd^{2}_{BB}	Standard deviation of expected probability to be observed	logN(-0.23, 0.005)
Data		
$t_{i,y}$	Mean daily air temperature	-
$f_{j,y}$	Mean daily discharge	-
$N^{Obs}_{j,y}$	Daily smolt count	-

435 Table 3. Results from the predictive study in 2005 (real missing data), 2007 ('first 17% missed' -

436 scenario) and 2014 ('peak of the migration +-2 days missing' -scenario). For posterior distributions,

437 median values and 90% probability intervals are illustrated.

Year	Sum of missing counts	% of missing counts out of total observed	model M1		model M2	
			Number of predicted	% of predicted out of total	Number of predicted	% of predicted out of total
2005	NA	NA	1620 (340,3930)	10% (2%,22%)	2020 (450,4850)	10% (3%,22%)
2007	2539	17%	2500 (510,6220)	14% (4%,31%)	3220 (700,7990)	15% (4%,32%)
2014	4085	21%	3790 (2030,6040)	18% (11%,26%)	4700 (2620,7390)	18% (11%,26%)

439
440 Figure 1: Setup of the video monitoring system at the river bed in Utsjoki. Arrows illustrate the

441 position and direction of each of the 8 cameras.

443
444   Figure 2. Data on daily mean air temperature and mean discharge at Utsjoki on study years 2005-

445   2009 and 2014.

448   Figure 3. Annual prior distributions for the timing of the smolt run given the environmental

449   covariates (daily mean air temperature and discharge).

450
451   Figure 4. Annual number of smolts passing the video site. Posterior distributions of M1 (grey

452   boxplots), M2 (black boxplots) and count data (black dots).

Figure 5. Daily number of smolts that pass the video site in 2005-2009 and 2014. Posterior

distributions of M1 (boxplots) and count data (dots). Data points included in the study are marked

with black and the ones excluded (counts treated as missing) marked with grey.

Figure 6. Daily number of smolts that pass the video site in 2005-2009 and 2014. Posterior

distributions of M2 (boxplots) and count data (dots). Data points included in the study are marked

with black and the ones excluded (counts treated as missing) marked with grey.

Figure 7. Probability that a smolt begins the migration at given temperature. Prior (grey boxplots)

and posterior distributions (black boxplots) from M2.

470
471   Figure 8. Cumulative distribution of expected travel time in days from the point of departure to the

472   video site. Above: Cumulative distribution at low discharge (10 m^3/s). Below: Cumulative distribution

473   at high discharge (100 m^3/s). Prior (grey boxplots) and posterior distributions (black boxplots) from

474   M2.

483 Figure 9. The expected observation probability at given discharge. Prior (grey boxplots) and

484 posterior distributions (black boxplots) from M2.

Appendix A: Lognormal approximation for Dirichlet-distribution

Let's consider a set of Dirichlet-distributed parameters:

(A.1) $\quad q_{1:n} \sim Dirich(\mu_1\eta, \dots, \mu_n\eta),$

Where μ_i is the expected probability for slot i and η is the overdispersion parameter. Such Dirichlet-

distribution can be approximated with a set of lognormally distributed z_i by assuming

(A.2) $q_{1:n} = z_i / \sum_i z_i$

(A.3) $z_i \sim logN(M_i, T_i)$

(A.4) $M_i = \log(\mu_i) - \frac{0.5}{T_i}$ and

(A.5) $T_i = \log\left(\frac{1}{\mu_i \eta} + 1\right).$

Appendix B: JAGS code for the arrival model

```
497  model{
498
499    # Process of departing
500    # ==========================
501    for(y in 1:nYears){
502     for(i in 1:nDays){
503       # p: probability to start migration at day t, if haven't done so earlier
504       # t: daily mean temperature
505       logit(p[i,y])<-p2[i,y]
506       p2[i,y]~dnorm(aP+bP*t[i,y],1/pow(sdP,2))
507      }
508     }
509    # P: expected proportion of smolts departing at day i
510    for(y in 1:nYears){
511     P[1,y]<-p[1,y]
512     for(i in 2:(nDays-1)){
513       P[i,y]<-(1-sum(P[1:(i-1),y]))*p[i,y]
```

```
514        }

515          P[nDays,y]<-1-sum(P[1:60,y])

516        }

517

518      aP~dnorm(-20,1)

519      bP~dlnorm(0.6,10)

520      sdP~dlnorm(0,1)

521

522      # Process of travelling

523      # =============

524      # qD: probability that a smolt that departs at day i passes the video site at day j

525      for(y in 1:nYears){

526        for(i in 1:nDays){

527          # i: day of departure

528          # j: day of passing the video site

529          # f: daily mean discharge

530          # j==i

531          qDx[i,i,y]<-phi((log(0.5)-MD[i,y])/SD)

532          # j>i

533          for(j in (i+1):(i+13)){

534            qDx[i,j,y]<-phi((log(j-i+0.5)-MD[i,y])/SD)-phi((log(j-i-0.5)-MD[i,y])/SD)

535          }

536          for(j in i:(i+13)){

537            qD[i,j,y]<-qDx[i,j,y]/(sum(qDx[i,i:(i+13),y])+0.0001)

538          }

539          MD[i,y]<-log(muD[i,y])-0.5/TD

540          muD[i,y]~dlnorm(log(exp(aD-bD*f[i,y]))-0.5/TmuD, TmuD)

541        }

542      }
```

```
543    SD<-1/sqrt(TD)

544    TmuD<-1/log(cvmuD*cvmuD+1)

545    TD<-1/log(cvD*cvD+1)

546

547    aD~dlnorm(0.52,14)

548    bD~dlnorm(-4.6,25)

549    cvmuD~dunif(0.001,1)

550    cvD~dunif(0.001,2)

551

552    # Arrival distribution

553    # ==============================================

554    # Joint distribution of qD and P: prob that a smolt departing day i arrives day j

555    for(y in 1:nYears){

556      for(i in 1:nDays){ # day of departure

557        for(j in i:(i+13)){ # day of passing the video site

558          qJ[j,i,y]<-qD[i,j,y]*P[i,y]

559        }

560      }

561

562      # Expected proportion of smolts passing the video site each day

563      for(j in 1:14){

564        muqN[j,y]<-sum(qJ[j,1:j,y])+0.0001 # small constant added for numerical stability

565      }

566      for(j in 15:nDays){

567        muqN[j,y]<-sum(qJ[j,(j-13):j,y])+0.0001

568      }

569    }

570

571    for(y in 1:nYears){
```

```
572    Ntot[y]<-exp(LNtot[y]) # total run size in year y

573    LNtot[y]~dunif(7,15)

574

575    for(j in 1:(nDays-1)){

576      N[j,y]<-round(qN[j,y]*Ntot[y]) # daily number of fish passing

577    }

578    N[nDays,y]<-round(Ntot[y]*(1-sum(qN[1:(nDays-1),y])))

579

580    # qN: daily proportion of smolts

581    # dirichlet-distribution approximated with lognormal

582    qN[1:nDays,y]<-zN[1:nDays,y]/sum(zN[1:nDays,y])

583

584    for(i in 1:nDays){

585      zN[j,y]~dlnorm(MN[j,y], TauN[j,y])

586    }

587

588    alphaN[1:nDays,y]<-muqN[1:nDays,y]*eta_alphaN

589    MN[1:nDays,y]<-log(muqN[1:nDays,y])-0.5/TauN[1:nDays,y]

590    TauN[1:nDays,y]<-1/log((1/alphaN[1:nDays,y])+1)

591    }

592

593    eta_alphaN~dunif(0.001,100000)

594

595    # Observation process

596    # ====================

597    for(y in 1:nYears){

598      for(j in 1:nDays){

599        # Nobs: Observed number of smolts

600        Nobs[j,y]~dbetabin(100,10,N[j,y])  # M1
```

601   *Nobs[j,y]~dbetabin(muB[j,y]*etaB,(1-muB[j,y])*etaB,N[j,y]) #M2*

603   *muB[j,y]<-0.6*(exp(BB[j,y])/(1+exp(BB[j,y])))+0.3*

604   *BB[j,y]~dnorm(aB-bB*f[j,y],1/pow(sdBB,2))*

606   *}*

607   *}*

608  *# priors for observation process M2*

609  *aB~dnorm(2.9,60)*

610  *bB~dlnorm(-2.6,984)*

611  *sdBB~dlnorm(-0.23,210)*

612  *etaB~dunif(5,1000)*

613  *}*