Potential for microbial anaerobic hydrocarbon degradation in naturally

occurring petroleum-associated deep-sea sediments

Xiyang Dong [1, *], Chris Greening [2], Jayne E. Rattray [1], Anirban Chakraborty [1],

Maria Chuvochina[2, 3], Daisuke Mayumi [1, 4], Carmen Li [1], James M. Brooks [5],

Bernie B. Bernard [5], Ryan A. Groves[1], Ian A. Lewis [1], Casey R.J. Hubert [1, *]

[1] Department of Biological Sciences, University of Calgary, Calgary, T2N 1N4, Canada

[2] School of Biological Sciences, Monash University, Clayton, VIC 3800, Australia

[3] Australian Centre for Ecogenomics, School of Chemistry and Molecular Biosciences, The

University of Queensland, QLD 4072, Australia

[4] Institute for Geo-Resources and Environment, Geological Survey of Japan, National Institute of

Advanced Industrial Science and Technology (AIST), 1-1-1 Higashi, Tsukuba, 305-8567, Japan

[5] TDI Brooks International, College Station, Texas, TX 77845, USA

* Corresponding author. E-mail: xiyang.dong@ucalgary.ca.

* Corresponding author. E-mail: chubert@ucalgary.ca. Telephone: +1 (403) 220-7794.

The authors declare no conflict of interest.

Abstract

The lack of cultured isolates and microbial genomes from the deep seabed means that very little

is known about the ecology of this vast habitat. Here, we investigated energy and carbon

acquisition strategies of microbial communities from three deep seabed petroleum seeps (3 km

water depth) in the Eastern Gulf of Mexico. Shotgun metagenomic analysis revealed that each

sediment harbored diverse communities of chemoheterotrophs and chemolithotrophs. We

recovered 82 metagenome-assembled genomes affiliated with 21 different archaeal and bacterial

phyla. Multiple genomes encoded enzymes for acetogenic fermentation of aliphatic and aromatic

compounds, specifically of candidate phyla *Aerophobetes*, *Aminicenantes*, TA06 and

Bathyarchaeota. Microbial interactions in these communities are predicted to be driven by

acetate and molecular hydrogen, as indicated by a high abundance of fermentation, acetogenesis,

and hydrogen utilization pathways. These findings are supported by sediment geochemistry,

metabolomics and thermodynamic modelling of hydrocarbon degradation. Overall, we infer that

deep-sea sediments experiencing thermogenic hydrocarbon inputs harbor phylogenetically and

functionally diverse communities potentially sustained through anaerobic hydrocarbon, acetate

and hydrogen metabolism.

Deep-sea sediments, generally understood to be those occurring in water depths greater than

~500 meters, represent one of the largest habitats on Earth. In recent years, culture-independent

16S rRNA gene surveys and metagenomic studies have revealed these sediments host a vast

abundance and diversity of bacteria and archaea [1-8]. Cell numbers decrease with sediment depth

and age, from between 10^6 and 10^{10} cm^{-3} in the upper cm at the sediment-water interface to

below 10^4 cm^{-3} several kilometers below the ocean floor [9, 10]. However, due to a lack of cultured

representatives and genomes recovered from deep-sea sediments, it remains largely unresolved

how microorganisms survive and function in these nutrient-limited ecosystems. Energy and

carbon sources are essential requirements that allow the buried microorganisms to persist. With

sunlight penetration not reaching the deep seabed, photosynthetic processes do not directly

support these communities [11]. It has therefore been proposed that deep sea benthic and

subsurface microbes are primarily sustained by complex detrital organic matter, including

carbohydrates, proteinaceous compounds, and humic substances, derived from the overlying

water column via sedimentation [11-13].

Another important potential carbon and energy source in deep-sea sediments are petroleum

geofluids that migrate from subseafloor reservoirs up to the seafloor [14]. Petroleum compounds

include smaller gaseous molecules, such as methane, propane and butane, and larger aliphatic

and aromatic liquids. Numerous studies have investigated the role of methane oxidation in

seabed sediments, which is mediated by anaerobic methanotrophic archaea (ANME), generally

in syntrophy with bacteria respiring sulfate or other electron acceptors [4, 6, 8, 15, 16]. In contrast,

little is known about the degradation of larger alkanes or aromatic compounds by deep seabed

microorganisms. Vigneron et al. [2] performed a comparative gene-centric study of hydrocarbon

and methane seeps of the Gulf of Mexico, and suggested that microorganisms in deep cold seeps

(water depth ~1 km) can potentially utilize a range of non-methane hydrocarbons. However, due

to the absence of metagenome binning in that study, relevant metabolic functions were not

assigned to specific pathways or taxa.

In addition to organic carbon compounds, microbial life in deep-sea sediments is also supported

by inorganic electron donors. Some microorganisms have been isolated from deep sediments that

are able to sustain themselves by oxidizing elemental sulfur, hydrogen sulfide, carbon monoxide,

ammonia and molecular hydrogen (H_2) [6, 8, 11]. Of these, H_2 is a particularly important energy

source given its production in large quantities by biological and geochemical processes. H_2 can

be generated as a metabolic byproduct of fermentation, together with volatile fatty acids such as

acetate, during organic matter degradation [9, 17]. H_2 can also be produced abiotically via

serpentinization, radiolysis of water, or thermal alteration of sedimentary organic matter [18]. For

example, the radiolysis of water by naturally occurring radionuclides (*e.g.* ^{40}K and ^{238}U) is

estimated to produce 10^{11} mol H_2 per year [8, 19]. Depending on the availability of electron

acceptors, H_2 oxidation can be coupled to sulfate, nitrate, and metal respiration, acetogenesis and

methanogenesis [8, 11].

To understand the role of hydrocarbon substrates and hydrogen in supporting microbial life in

deep-sea sediments, we performed metagenomic, geochemical and metabolomic analyses of

three deep seabed sediments (water depth ~3km). The three sites exhibited different levels of

migrated thermogenic hydrocarbons. Metagenomes generated from sediment samples obtained

from each site were assembled and binned to obtain metagenome-assembled genomes (MAGs)

and to reconstruct metabolic pathways for dominant members of the microbial communities.

Complementing this genome-resolved metagenomics, a gene-centric analysis was performed by

directly examining unassembled metagenomic data. Through the combination of metagenomics

with geochemistry and metabolomics, with supporting thermodynamic modeling, we provide

evidence that (1) deep-sea sediments harbor phylogenetically diverse heterotrophic and

lithotrophic microbial communities; (2) some members from the candidate phyla are engaged in

degradation of aliphatic and aromatic thermogenic hydrocarbons; and (3) microbial community

members are likely interconnected via acetate and hydrogen metabolism.

Results

Sediment geochemistry

This study tested three petroleum-associated near-surface sediments (referred to as Sites E26,

E29 and E44) sampled from the Eastern Gulf of Mexico [20]. Petroleum content and other

geochemical characteristics were analyzed for each of the three sites (Table 1). All sites had high

concentrations of aromatic compounds and liquid alkanes; aromatic compounds were most

abundant at Site E26, while liquid alkanes were at 2.5-fold higher concentration at Sites E26 and

E29 than Site E44. Alkane gases were only abundant at Site E29 and were almost exclusively

methane (CH_4). CH_4 sources can be inferred from stable isotopic compositions of CH_4 and molar

ratios of CH_4 to higher hydrocarbons [15]. Ratios of $C_1/(C_2+C_3)$ were greater than 1,000 and $\delta^{13}C$

values of methane were more negative than -60‰, indicating that the CH_4 in these sediments is

predominantly biogenic [15, 21]. GC-MS revealed an unresolved complex mixture (UCM) of

saturated hydrocarbons in the C_{15+} range in all three sites. Such UCM signals correspond to

degraded petroleum hydrocarbons and may indicate the occurrence of oil biodegradation at these

sites [22]. Signature metabolites for anaerobic biodegradation of alkanes and aromatic compounds

[23] were also detected, including benzoate, toluate and methyl- or trimethylsilyl esters (Table S1).

High concentrations of sulfate (>20 mM) were detected at each of the three sites (Table 1),

consistent with sulfate being present in high concentrations in seawater and diffusing into the

sediments. H_2 and acetate concentrations were both below limits of detection (0.005-0.02 ppm

and 2.5 µM, respectively) consistent with H_2 and acetate being present at extremely low steady-

state concentrations in deep-sea sediments due to tight coupling between producers and

consumers [3, 15].

Deep-sea sediments harbor phylogenetically diverse bacterial and archaeal communities

Illumina NextSeq sequencing of genomic DNA from deep-sea sediment communities produced

85,825,930, 148,908,270, and 138,795,692 quality-filtered reads for Sites E26, E29, and E44,

respectively (Table S2). The 16S rRNA gene amplicon sequencing results suggest the sediments

harbor diverse bacterial and archaeal communities, with Chao1 richness estimates of 359, 1375

and 360 using bacterial primers as well as 195, 180 and 247 using archaeal primers, for Sites E26,

E29 and E44, respectively (Table S3 and Figure S1). Taxonomic profiling of these metagenomes

using small subunit ribosomal RNA (SSU rRNA) marker genes reveals that the most abundant

phyla in the metagenomes were, in decreasing order, *Chloroflexi* (mostly classes

Dehalococcoidia and *Anaerolineae*), *Candidatus* Atribacteria, *Proteobacteria* (mostly class

Deltaproteobacteria), and *Candidatus* Bathyarchaeota (Figure 1a). While the three sites share a

broadly similar community composition, notable differences were *Ca.* Bathyarchaeota and

Proteobacteria being in higher relative abundance at the sites with more hydrocarbons (E29 and

E26; Table 1), whereas the inverse is true for *Actinobacteria*, the Patescibacteria group, and *Ca.*

Aerophobetes that are all present in higher relative abundance at Site E44 where hydrocarbon

levels are lower. Additional sampling is required to determine whether these differences are due

to the presence of hydrocarbons or other factors.

Assembly and binning for the three metagenomes resulted in a total of 82 MAGs with >50%

completeness and <10% contamination based on CheckM analysis [24]. Reconstructed MAGs

comprise taxonomically diverse members from a total of 6 archaeal and 15 bacterial phyla

(Figure 2 and Table S4). Within the domain Bacteria, members of the phylum *Chloroflexi* are

highly represented in each sample, especially from the classes *Dehalococcoidia* and

Anaerolineae. Within the domain Archaea, members of phylum *Bathyarchaeota* were recovered

from all three sites. Most other MAGs belong to poorly understood candidate phyla that lack

cultured representatives, including *Aminicenantes* (formerly OP8), *Aerophobetes* (formerly

CD12), *Cloacimonas* (formerly WWE1), *Stahlbacteria* (formerly WOR-3), *Atribacteria*

(formerly JS1 and OP9), TA06 and the Asgard superphylum including *Lokiarchaeota*,

Thorarchaeota, and *Heimdallarchaeota*.

Candidate phylum TA06, also known as GN04 or AC1, was originally discovered in a

hypersaline microbial mat [25]. First genomic representatives of this phylum were recovered from

estuarine sediments [26] with a small number of other MAGs recently reported to belong to this

lineage [27, 28]. Due to the paucity of available MAGs and misclassifications based on 16S rRNA

gene sequences, members of TA06 are often 'confused' with members of the phylum WOR-3

(*Stahlbacteria*) [28]. In addition to the phylogenetic inference here based on 43 concatenated

protein marker genes (Figure 2), the placement of two bins within the original TA06 phylum is

further supported by genome classification based on concatenation of 120 ubiquitous, single-

copy marker genes [29] as well as classification of 16S rRNA genes using the SILVA database [30]

(Tables S4 and S5).

In summary, while there are considerable community-level differences between the three sample

locations, the recovered MAGs share common taxonomic affiliations at the phylum and class

levels. Guided by sediment geochemistry (Table 1), we subsequently analyzed the metabolic

potential of these MAGs to understand how dominant bacterial and archaeal community

members generate energy and biomass in these natural petroleum-associated deep-sea

environments. Hidden Markov models (HMMs) and homology-based models were used to

search for the presence of different metabolic genes both in the recovered MAGs and unbinned

metagenomes. Where appropriate, findings were further validated through metabolomic analyses,

phylogenetic visualization, and analysis of gene context.

Capacity for hydrocarbon and detrital biomass degradation in sediment microbial

communities

In deep-sea marine sediments organic carbon is supplied either as detrital matter from the

overlying water column or as aliphatic and aromatic petroleum compounds that migrate upwards

from underlying petroleum-bearing sediments [11]. With respect to detrital matter, genes involved

in carbon acquisition and breakdown were prevalent across both archaeal and bacterial MAGs.

These include genes encoding intracellular and extracellular carbohydrate-active enzymes and

peptidases, as well as relevant transporters and glycolysis enzymes (Figure 3 and Table S6). The

importance of these carbon acquisition mechanisms is supported by the detection of

corresponding intermediate metabolites, such as glucose and amino acids, in all three sediments

(Table S1). The ability to break down fatty acids and other organic acids via the beta-oxidation

pathway was identified in 13 MAGs, including members of *Chloroflexi*, *Deltaproteobacteria*,

Aerophobetes and *Lokiarchaeota* (Figure 3 and Table S6). These results align with many other

studies suggesting that the majority of seabed microorganisms are involved in recycling of

residual organic matter, including complex carbohydrates, proteins and lipids [13, 31, 32].

Unlike in other studies, the presence of petroleum hydrocarbons is a defining feature of the

sediments investigated here; thus a key goal of this study was to identify the potential for

microbial degradation of hydrocarbons as a source of energy and carbon. To this end, we focused

on functional marker genes encoding enzymes that catalyze the activation of mechanistically

sophisticated C-H bonds, to initiate hydrocarbon biodegradation [33]. For anaerobic hydrocarbon

degradation, four oxygen-independent C-H activation reactions have been characterized: (1)

addition of fumarate by glycyl-radical enzymes, *e.g.* for activation of alkylbenzenes and straight

chain alkanes [34]; (2) hydroxylation with water by molybdenum cofactor-containing enzymes, *e.g.*

for activation of ethylbenzene [33]; (3) carboxylation catalyzed by UbiD-like carboxylases, *e.g.* for

activation of benzene and naphthalene [35]; and (4) reverse methanogenesis involving variants of

methyl-coenzyme M reductase, *e.g.* for activation of methane and butane [36]. Most of the

evidence for mechanisms 1-3 has come from studies of hydrocarbon contaminated aquifers,

whereas mechanism 4 has been studied extensively in marine sediments [23, 37].

Evidence for glycyl radical enzymes that catalyze fumarate addition was found in 15 out of the

82 MAGs based on identifying genes encoding alkylsuccinate synthase (AssA) (Figures 3 and

4a). The *assA* sequences identified, while phylogenetically distant from canonical fumarate-

adding enzymes and pyruvate formate lyases (Pfl), form a common clade with Pfl-like AssA

from *Archaeoglobus fulgidus* VC-16 and *Abyssivirga alkaniphila* L81 (Figure 4a). Both of these

organisms have been shown experimentally to be capable of anaerobic alkane degradation [38, 39].

The putative *assA* genes identified here are present in all three samples regardless of

hydrocarbon concentrations. They belong to MAGs affiliated with the bacterial phyla

Aerophobetes, *Aminicenantes* and *Chloroflexi* as well as the archaeal phyla *Bathyarchaeota*,

Lokiarchaeota and *Thorarchaeota.* The highest relative abundance of putative *assA* sequences

was found in Site E29 as idnciated by quality-filtered reads, consistent with this sediment

containing the highest concentration of aliphatic compounds (Tables 1 and S7). Additional

searching for other genes encoding fumarate-adding enzymes in the quality-filtered reads (e.g.

bssA and *nmsA*, and canonical *assA*) did not return significant counts, while some hits to

canonical *assA* were indicated, suggesting it may also be possible existence in these habitats

(Figure 4 and Table S7). Among the other three anaerobic hydrocarbon biodegradation

mechanisms mentioned above, a MAG classified as *Dehalococcoidia* (*Chloroflexi* E29_bin2)

contained genes encoding putative catalytic subunits of *p*-cymene dehydrogenase (Cmd) and

alkane C_2-methylene hydroxylase (Ahy) (Figures 3 and S2), known to support *p*-cymene and

alkane utilization [37]. Genes encoding enzymes catalyzing hydrocarbon carboxylation, reverse

methanogenesis and aerobic hydrocarbon degradation (e.g. *alkB*, *nahC* and *nahG*) were not

detected (Table S6). The latter result is expected due to the low concentrations of oxygen in the

top 20 cm of organic rich seabed sediments.

Considering the degradation of aromatic hydrocarbons, genes responsible for reduction of

benzoyl-CoA were detected in 12 MAGs (Figures 3 and 4b). Benzoyl-CoA is a universal

biomarker for anaerobic degradation of monoaromatic compounds as it is a common

intermediate to biochemical pathways catalyzing this process [40]. Benzoyl-CoA reduction to

cyclohex-1,5-diene-1-carboxyl-CoA is performed by Class I ATP-dependent benzoyl-CoA

reductase (BCR; BcrABCD) in facultative anaerobes (*e.g. Thauera aromatica*) or Class II ATP-

independent reductase (Bam; BamBCDEFGHI) in strict anaerobes like sulfate reducers [41]. The

bcr genes detected are all Class I, and were found in bacterial MAGs (*i.e.*, *Dehalococcoidia*,

Anaerolineae, *Deltaproteobacteria*, *Aminicenantes* and TA06) and archaeal MAGs (*i.e.*,

Thermoplasmata and *Bathyarchaeota*) (Figures 3 and 4b). Genes for further transformation of

dienoyl-CoA to 3-hydroxypimelyl-CoA were also identified (Figures 3 and 4b), *i.e.*, genes

encoding for 6-oxo-cyclohex-1-ene-carbonyl-CoA hydrolase (Oah), cyclohex-1,5-diencarbonyl-

CoA hydratase (Dch) and 6-hydroxycyclohex-1-ene-1-carbonyl-CoA dehydrogenases (Had) [42].

Together with detection of 23-162 nM benzoate in these sediments (Table 1) these results

strongly suggest that the organisms represented by these MAGs are engaged in the typical

downstream degradation of aromatic compounds employing the central benzoyl-CoA Bcr-Dch-

Had-Oah pathway. However, the possible origins of benzoate are unclear based on current data.

Widespread capacity for fermentative production and respiratory consumption of acetate

and hydrogen

Analysis of MAGs from these deep-sea hydrocarbon-associated sediments suggests that

fermentation, rather than respiration, is the primary mode of organic carbon turnover in this

environment. Most recovered MAGs with capacity for heterotrophic carbon degradation lacked

respiratory primary dehydrogenases and terminal reductases, with exceptions being several

Proteobacteria and one *Chloroflexi* (Table S6). In contrast, only 6 and 14 MAGs contained

genes indicating the capability for fermentative production of ethanol and lactate, respectively,

whereas 69 MAGs contained genes for fermentative acetate production (Figure 3 and Table S6).

These findings are consistent with other studies emphasizing the importance of fermentation,

including acetate production, in deep-sea sediments [12, 43].

Acetate can also be produced by acetogenic CO_2 reduction through the Wood-Ljungdahl

pathway using a range of substrates, including heterotrophic compounds [15]. Partial or complete

sets of genes for the Wood-Ljungdahl pathway were found in 50 MAGs (Figures 3 and S3),

including those affiliated with phyla previously inferred to mediate acetogenesis in deep-sea

sediments through either the tetrahydrofolate-dependent bacterial pathway (*e.g. Chloroflexi* and

Aerophobetes) [7, 44] or the tetrahydromethanopterin-dependent archaeal variant (*e.g.*

Bathyarchaeota and Asgard group) [45, 46]. In addition, the signature diagnostic gene for the Wood-

Ljungdahl pathway (*acsB*; acetyl-CoA synthase) is in high relative abundance in the quality-

filtered metagenome reads at all three sites (Table S7). These observations are in agreement with

mounting evidence that homoacetogens play a quantitatively important role in organic carbon

cycling in the marine deep biosphere [45, 47, 48].

Evidence for H_2 metabolism was also found in MAGs from all three sites. We screened putative

hydrogenase genes from various subgroups in MAGs as well as unbinned metagenomic

sequences (Figures 1, 3 and Tables S6, S7). Surprisingly few H_2 evolving-only hydrogenases

were detected, with only five Group A [FeFe]-hydrogenases and five Group 4 [NiFe]-

hydrogenases detected across the bacterial and archaeal MAGs. Instead, the most abundant

hydrogenases within MAGs as well as in quality-filtered unassembled reads were the Group 3b,

3c, and 3d [NiFe]-hydrogenases. Group 3b and 3d hydrogenases are physiologically reversible,

but generally support fermentation in anoxic environments by coupling NAD(P)H reoxidation to

fermentative H_2 evolution [49-51]. Group 3c hydrogenases mediate a central step in

hydrogenotrophic methanogenesis, bifurcating electrons from H_2 to heterodisulfides and

ferredoxin [52]. The functional role of Group 3c hydrogenases in Bacteria and non-methanogenic

Archaea remains unresolved [51] yet their corresponding genes co-occurred with heterodisulfide

reductases across multiple archaeal and bacterial MAGs (Figure 3). Various Group 1 [NiFe]-

hydrogenases were also detected, which supports hydrogenotrophic respiration in conjunction

with a wide range of terminal reductases. This is consistent with previous studies in the Gulf of

Mexico that experimentally measured the potential for hydrogen oxidation catalyzed by

hydrogenase enzymes [53].

Given the genomic evidence for hydrogen and acetate production in these sediments, we

investigated whether any of the MAGs encoded terminal reductases to respire using these

compounds as electron donors. In agreement with the high sulfate concentrations (Table 1), the

key genes for dissimilatory sulfate reduction (*dsrAB*) were widespread across the metagenome

reads, particularly at Site E29 (Table S7). These genes were recovered from MAGs affiliated

with *Deltaproteobacteria* and *Dehalococcoidia* (Table S6). We also identified 31 novel reductive

dehalogenase (*rdhA*) genes across 22 MAGs, mainly from *Aminicenantes* and *Bathyarchaeota*

(Figure 3 and Table S6), suggesting that organohalides – that can be produced through abiotic

and biotic processes in marine ecosystems [54] – may be electron acceptors in these deep-sea

sediments. All MAGs corresponding to putative sulfate reducers and dehalorespirers encoded the

capacity to completely oxidize acetate and other organic acids to CO_2 using either the reverse

Wood-Ljungdahl pathway or TCA cycle (Figure 3 and Table S6). Several of these MAGs also

harbored the capacity for hydrogenotrophic dehalorespiration *via* Group 1a and 1b [NiFe]-

hydrogenases (Figure 3). In addition to these dominant uptake pathways, one MAG belonging to

the epsilonproteobacterial genus *Sulfurovum* (E29_bin29) included genes for the enzymes

needed to oxidize either H_2 (group 1b [NiFe]-hydrogenase), elemental sulfur (SoxABXYZ), and

sulfide (Sqr), using nitrate as an electron acceptor (NapAGH); this MAG also has a complete set

of genes for autotrophic CO_2 fixation *via* the reductive TCA cycle (Figure 3 and Table S6). The

capacity for methanogenesis appears to be relatively low as revealed by the MAGs retrieved

from the three sediments. None of the MAGs contained *mcrA* genes, but these genes for

methanogenesis were detected in quality-filtered unassembled reads in all three sediments

(Figures 1d and S4) and were mainly affiliated with acetoclastic methanogens at Site E29, and

hydrogenotrophic methanogens at the other two sites (Figure 1d and S4). Overall, the

collectively weak *mcrA* signal in the metagenomes suggests that the high levels of biogenic

methane detected by geochemical analysis (Table 1) is due to methanogenesis in sediment layers

deeper than the top 20 cm.

Thermodynamic modelling of hydrocarbon degradation

Both the geochemistry data and biomarker gene survey suggest that hydrocarbon degradation is

occurring in the three deep-sea sediments sampled (Tables 1 and S1). Recreating environmental

conditions for cultivating the organisms represented by the retrieved MAGs is a challenging and

time consuming process, preventing further direct evidence of hydrocarbon degradation

capabilities (and other metabolisms) among the majority of the lineages represented by the

MAGs retrieved here [48]. Instead, we provide theoretical evidence that hydrocarbon degradation is

feasible in this environment by modelling whether these processes are thermodynamically

favorable in the conditions typical of deep sea sediments, namely high pressure and low

temperature.

As concluded from the genome analysis and supported by metabolomics (Table 1), it is likely

that most hydrocarbon oxidation occurs through fermentation rather than respiration. Taking

hydrogen production and the Wood-Ljungdahl pathway into consideration (Figures 3 and 4), we

compared the thermodynamic constraints on hydrocarbon biodegradation for two plausible

scenarios: (1) fermentation with production of hydrogen and acetate, and (2) fermentation with

production of acetate alone. Hexadecane and benzoate are used as representative aliphatic and

aromatic compounds, respectively, based on the geochemistry results (e.g. C_{2+} alkane detection)

and genomic analysis (e.g. *bcr* genes) [47, 55]. The calculated results show that the threshold

concentrations of acetate that result in favorable energetics ($\Delta G' < 0$) for fermentative co-

generation of acetate and hydrogen require acetate to be extremely low in a hexadecane

degradation scenario ($< 10^{-12}$ mM acetate), and acetate to be at moderate levels in a benzoate

degradation scenario (< 3.8 mM acetate) (Figure 5). By contrast, for fermentation leading to

production of only acetate, its concentration can be as high as 470 mM in a benzoate degradation

scenario and as high as 300 mM in a hexadecane degradation scenario (Figure 5). Fermentative

degradation of hexadecane to hydrogen and acetate in the deep seabed could therefore be less

favorable than acetate production alone via the Wood-Ljungdahl pathway. If microbial

communities are consuming hexadecane or more complex hydrocarbons as carbon and energy

sources, it might be likely that they employ the Wood-Ljungdahl pathway to produce acetate.

These lines of evidence emphasize the potential importance of acetogenic fermentation of

hydrocarbons in the deep seabed. However, other reactions such as fermentation to H_2 still

cannot be excluded, *e.g.*, for less complex hydrocarbons such as benzoate and related compounds.

Discussion

In this study, metagenomics revealed that the majority of the Bacteria and Archaea comprising deep-sea sediment microbial communities belong to candidate phyla that lack cultured representatives and sequenced genomes (Figures 1 and 2). As a consequence, it is challenging to link phylogenetic patterns with the microbial functional traits underpinning the biogeochemistry of deep seabed habitats. Here, we were able to address this by combining *de novo* assembly and binning of metagenomic data with geochemical and metabolomic analyses, and complementing our observations with thermodynamic modeling. Pathway reconstruction from 82 MAGs recovered from the three deep-sea surface sediments revealed that many community members were capable of anaerobic hydrocarbon degradation as well as acquiring and hydrolyzing residual organic matter (Figure 3), whether supplied as detritus from the overlying water column or as autochthonously produced necromass (Figure 6). Heterotrophic fermenters and acetogens were in considerably higher relative abundance than heterotrophic respirers, despite the abundance of sulfate in the sediments (Table 1). For example, while genomic coverage of putative sulfate reducers is relatively low (< 1% of the communities), the most abundant MAG at each site were all putative acetogenic heterotrophs, *i.e. Dehalococcoidia* E26_bin16, *Actinobacteria* E44_bin5, and *Aminicenantes* E29_bin47 for Sites E26, E44 and E29 respectively (~3.3-4.5% relative abundance, Table S4). Therefore, in contrast with coastal sediments [56], microbial communities in the deep seabed are likely influenced by the capacity to utilize available electron donors more so than by the availability of oxidants.

In this context, multiple lines of evidence indicate degradation of aliphatic or aromatic petroleum compounds as carbon and energy sources for anaerobic populations in these deep-sea

hydrocarbon seep environments (Table 1, Figures 3 - 5). Whereas capacity for detrital organic

matter degradation is a common feature in the genomes retrieved in this study, and from many

other environments [26], anaerobic hydrocarbon degradation is a more exclusive feature that was

detected in 23 out of 82 MAGs. Evidence of anaerobic alkane oxidation via fumarate addition

and hydroxylation pathways, as well as anaerobic aromatic compound degradation by the Class I

benzoyl-CoA reductase pathway, was found in all three sediments. The ability to utilize

hydrocarbons may explain the ecological dominance (high relative abundance) of certain

lineages of Bacteria and Archaea in these microbial communities (Figure 1a) as many of those

phyla have previously been found to be associated with hydrocarbons in various settings. For

example, *Aerophobetes* have been detected in other cold seep environments [7], *Aminicenantes* are

often found associated with fossil fuels [27], and *Chloroflexi* harboring genes for anaerobic

hydrocarbon degradation have been found in hydrothermal vent sediments [4]. While Archaea

have been reported to mediate oxidation of methane and other short-chain alkanes in sediments [5,

36], few have been reported to anaerobically degrade larger hydrocarbons [37]. The finding of

Bathyarchaeota and other archaeal phyla potentially capable of anaerobic hydrocarbon

degradation extends the potential hydrocarbon substrate spectrum for Archaea. More broadly,

these findings extend the breadth of lineages that include putative hydrocarbon-degrading

bacteria and archaea in deep-sea sediments. Current knowledge of anaerobic hydrocarbon

degradation remains limited, with the majority of studies focused on environments subject to

anthropogenic hydrocarbon contamination, most notably groundwater aquifers [37]. It is possible

that microorganisms inhabiting deep-sea sediments harbor novel mechanisms for anaerobic

hydrocarbon degradation that may be relevant for biotechnology and bioremediation in a variety

of other settings, *e.g.*, other cold habitats. Future studies of genome-enabled hydrocarbon

degradation using samples such as the sediments studied here may elucidate this further.

Genomic analyses of 12 MAGs harboring genes for central benzoyl-CoA pathway reveal that

they are likely a mixture of obligate fermenters and sulfate reducers. The finding that these

organisms use the ATP-consuming class I, not the reversible class II, benzoyl-CoA reductase is

surprising. It is generally thought that strict anaerobes must use class II BCRs because the

amount of energy available from benzoate oxidation during sulfate reduction or fermentation is

not sufficient to support the substantial energetic requirement of the ATP-dependent class I BCR

reaction [42]. However, acetogenic fermentation of hydrocarbons may explain how the Class I

reaction could be thermodynamically favorable, as shown in Figure 5. In agreement with this,

there are reported exceptions to the general Class I vs Class II observations, such as the

hyperthermophilic archaeon *Ferroglobus placidus* that couples benzoate degradation via the

Class I system with iron reduction [42], and fermentative deep-sea *Chloroflexi* strains DscP3 and

Dsc4 that contain genes for class I benzoyl-CoA reductases [44]. Indeed, acetogens can utilize

many different substrates and have relatively high ATP yields as well as thermodynamic

efficiencies toward heterotrophic substrates, which is consistent with the proposed importance of

acetogens in energy-limited seafloor ecosystems [45, 47].

Based on the evidence presented here we propose that acetate and hydrogen are the central

intermediates underpinning community interactions and biogeochemical cycling in these deep-

sea sediments (Figure 6). Maintaining low acetate and hydrogen concentrations in the

environment is important for promoting continuous fermentation of organic substrates, consistent

with thermodynamic constraints (Figure 5). Acetate and hydrogen in sediment porewater were

below detection limits, consistent with the high turnover rates of both compounds. This may

correspond with the genomic potential within these microbial communities for the coupling of

acetate consumption to sulfate reduction, organohalide respiration and acetoclastic

methanogenesis, as suggested in other studies [55, 57]. Some community members also appear to be

capable of H_2 consumption, including *via* putative heterodisulfide reductase-coupled

hydrogenases. In turn, hydrogen oxidation can support autotrophic carbon fixation and therefore

may provide a feedback loop for regeneration of organic carbon. Acetate- and hydrogen-

oxidizing community members are likely to promote upstream fermentative degradation of

necromass and hydrocarbons.

Overall, this metagenome dataset extended the knowledge of metabolic potential of microbial

communities inhabited in deep-sea sediments that receive an input of thermogenic hydrocarbon.

They are mostly likely sustained through fermentation, acetogenesis and hydrogen metabolisms.

More importantly, supported by geochemical and metabolomics analysis of hydrocarbons and

other metabolites as well as by thermodynamic modelling of different hydrocarbon degradation

scenarios, our findings expanded the diversity of microbial lineages with the potential for

anaerobic hydrocarbon degradation through *e.g.* the activity of glycyl radical enzymes. Together

with the recent discovery of anaerobic butane degradation in gas-rich hydrothermally-heated

sediments [36], it can be inferred that anaerobic degradation of hydrocarbons heavier than methane

might be more widespread than previously expected. They may play as an important chemical

energy in dark deep-sea sediments just like hydrogen, sulfide and methane [58].

Methods

Sampling and geochemical measurements

The three marine sediment samples used in this study were collected from the near-surface (top 20 cm) of the seafloor in the Eastern Gulf of Mexico as part of a piston coring survey, as described previously [20]. Samples for hydrocarbon characterization were sectioned on board the research vessel immediately following piston core retrieval, flushed with N_2 and sealed in hydrocarbon-free gas tight metal canisters then frozen until analysis. Interstitial gas analysis was later performed on the headspace in the canisters using GC with Flame Ionization Detector (GC-FID). Sediment samples for gas/liquid chromatography and stable isotope analysis were frozen, freeze-dried and homogenized then extracted using accelerated solvent extraction (ACE 200). Extracts were subsequently analyzed using GC/FID, a Perkin-Elmer Model LS 50B fluorometer, GC/MS and Finnigan MAT 252 isotope mass spectrometry as detailed elsewhere [59].

Sulfate and chloride concentrations were measured in a Dionex ICS-5000 reagent-free ion chromatography system (Thermo Scientific, CA, USA) equipped with an anion-exchange column (Dionex IonPac AS22; 4×250 mm; Thermo Scientific), an EGC-500 K_2CO_3 eluent generator cartridge and a conductivity detector. Organic acids were analysed in the 0.2 µm filtered sediment porewater using a Thermo RS3000 HPLC fitted with an Ultimate 3000 UV detector. Separation was achieved over an Aminex HPX-87H organic acid column (Biorad, USA) under isocratic conditions (0.05 mM H_2SO_4) at 60°C with a run time of 20 minutes. Organic acids were compared to the retention time of known standards and the limit of detection for acetate was determined to be 2.5 µM.

For the analysis of metabolites, sediment was spun down, the supernatant collected, diluted 1:1

in pure methanol, and filtered through 0.2 µm Teflon syringe filters. Extracts were separated

using Ultra High-Performance Liquid Chromatography (UHPLC) equipped with a hydrophilic

interaction liquid chromatography column (Syncronis HILIC, Thermo Fisher). A Thermo Fisher

Scientific Q-Exactive HF mass spectrometer in negative-mode electrospray ionization was used

to collect high-resolution full-scan MS data. In addition, benzoate ion (m/z 121.02943) was

subjected to fragmentation using collision induced dissociation with a collision energy of 10%

and activation time 20 ms (m/z 121.02943 > 77.03948). Metabolites were further identified using

accurate mass and retention times of standards using Thermo Xcalibur software and MAVEN

freeware. Larger compound lists were assigned identification using a combination of MAVEN

and the KEGG database.

DNA extraction and sequencing

For the three sediment samples, DNA was extracted from 10 g of sediment using the PowerMax

Soil DNA Isolation Kit (12988-10, QIAGEN) according to the manufacturer's protocol with

minor modifications for the step of homogenization and cell lysis *i.e.*, cells were lysed in

PowerMax Bead Solution tubes for 45 s at 5.5 m s^{-1} using a Bead Ruptor 24 (OMNI

International). DNA concentrations were assessed using a Qubit 2.0 fluorometer (Thermo Fisher

Scientific, Canada). Metagenomic library preparation and DNA sequencing was conducted at the

Center for Health Genomics and Informatics in the Cumming School of Medicine, University of

Calgary. DNA fragment libraries were prepared by shearing genomic DNA using a Covaris

sonicator and the NEBNext Ultra II DNA library preparation kit (New England BioLabs). DNA

was sequenced on a ~40 Gb (*i.e.* 130 M reads) mid-output NextSeq 500 System (Illumina Inc.)

300 cycle (2 × 150 bp) sequencing run.

To provide a high-resolution microbial community profile, the three samples were also subjected

to 16S rRNA gene amplicon sequencing on a MiSeq benchtop sequencer (Illumina Inc.). DNA

was extracted from separate aliquots of the same sediment samples using the DNeasy

PowerLyzer PowerSoil kit (MO BIO Laboratories, a Qiagen Company, Carlsbad, CA, USA) and

used as the template for different PCR reactions. The v3-4 region of the bacterial 16S rRNA

gene and the v4-8 region of the archaeal 16S rRNA gene were amplified using the primer pairs

SD-Bact-0341-bS17/SD-Bact-0785-aA21 and SD-Arch-0519-aS15/SD-Arch-0911-aA20,

respectively [60] as described previously [20] on a ~15 Gb 600-cyce (2 x 300 bp) sequencing run (for

results see Figure S1).

Metagenomic assembly and binning

Raw reads were quality-controlled by (1) clipping off primers and adapters and (2) filtering out

artifacts and low-quality reads as described previously [61]. Filtered reads were assembled using

metaSPAdes version 3.11.0 [62] and short contigs (<500 bp) were removed. Sequence coverage

was determined by mapping filtered reads onto assembled contigs using BBmap version 36

(https://sourceforge.net/projects/bbmap/). Binning of metagenome contigs was performed using

MetaBAT version 2.12.1 (--minContig 1500) [63]. Contaminated contigs in produced bins were

further removed based on genomic properties (GC, tetranucleotide signatures, and coverage) and

taxonomic assignments using RefineM version 0.0.22 [64]. Resulting bins were further examined

for contamination and completeness using CheckM version 1.0.8 with the lineage-specific

workflow [24].

Annotation

For MAGs, genes were called by Prodigal (-p meta) [65]. Metabolic pathways were predicted

against the KEGG GENES database using the GhostKOALA tool [66] and against the Pfam,

TIGRfam and custom HMM databases (https://github.com/banfieldlab/metabolic-hmms) using

MetaErg (https://sourceforge.net/projects/metaerg/). The dbCAN web server was used for

carbohydrate-active gene identification (cutoffs: coverage fraction: 0.40; e-value: 1e-18) [67].

Genes encoding proteases and peptidases were identified using BLASTP against the MEROPS

database release 12.0 (cutoffs: e-value, 1e-20; sequence identity, 30%) [68]. Genes involved in

anaerobic hydrocarbon degradation were identified using BLASTP against a custom database

(Table S8) (cutoffs: e-value, 1e-20; sequence identity, 30%). Hydrogenases were identified and

classified using a web-based search using the hydrogenase classifier HydDB with option of

Check sequences using CDD [69].

Full-length 16S rRNA genes were reconstructed from metagenomic reads using phyloFlash

version 3.1 (https://hrgv.github.io/phyloFlash/) together with the SILVA SSU 132 rRNA

database [30]. Diversity calculations were based on phyloFlash counts as well as the separate 16S

rRNA gene amplicon library results. Functional and taxonomic McrA gpkgs were used to assess

the diversity of methanogens against the metagenomic reads using GraftM with default

parameters [70]. Genes encoding the catalytic subunits of hydrogenases, *dsrA*, *acsB*, *assA*, *nmsA*

and *bssA* were retrieved from metagenomic reads through diamond BLASTx [71] queries against

comprehensive custom databases [46, 69, 72] (cutoffs: e-value, 1e-10; sequence identity, 70%).

Phylogenetic analyses

For taxonomic classification of each MAG, two methods were used to produce genome trees that were then used to validate each other. In the first method the tree was constructed using concatenated proteins of up to 16 syntenic ribosomal protein genes following procedures reported elsewhere [73]; the second tree was constructed using concatenated amino acid sequences of up to 43 conserved single-copy genes following procedures described previously [74]. Both trees were calculated using FastTree version 2.1.9 (-lg -gamma) [75] and resulting phylogenies were congruent. Reference genomes for relatives were accessed from NCBI GenBank including genomes selected from several recent studies representing the majority of candidate bacterial and archaeal phylogenetic groups [4, 64, 76-79]. The tree in Figure 2 was inferred based on concatenation of 43 conserved single-copy genes (Database S1). To be accurate, it was built using RAxML version 8 [80] implemented by the CIPRES Science Gateway [81]. It was called as follows: raxmlHPC-HYBRID -f a -n result -s input -c 25 -N 100 -p 12345 -m PROTCATLG -x 12345. The phylogeny resulting from RAxML is consistent with the taxonomic classification of MAGs that resulted from FastTree. Interactive tree of life (iTOL) version 3 [82] was used for tree visualization and modification.

For phylogenetic placements of functional genes, sequences were aligned using the MUSCLE algorithm [83] included in MEGA7 [84]. All positions with less than 95% site coverage were eliminated. Maximum likelihood phylogenetic trees were constructed in MEGA7 using a general time reversible substitution model and uniform rates among sites. These trees were bootstrapped with 100 replicates.

508   Taxonomic classification of MAGs inferred to belong to candidate phylum TA06 after

509   phylogenetic analyses were additionally confirmed by performing classify workflow using

510   GTDB-Tk version 0.0.6+ (https://github.com/Ecogenomics/GtdbTk).

Thermodynamic calculations

The values of Gibbs free energy of formation for substances were taken from Madigan et al. [85]

and Dolfing et al. [55]. The pH used in all calculations was 8.0 as reported in a previous

thermodynamic study of deep marine sediments [47], partial pressure was 300 atm based on water

depths at the three sites (http://docs.bluerobotics.com/calc/pressure-depth/), and temperature was

set as 4°C to represent deep sea conditions [86]. Calculations followed accepted protocols for

determining reaction kinetics and thermodynamics [87].

Acknowledgements

The work was supported by a Genome Canada Genomics Applied Partnership Program (GAPP)

grant to CRJH. We thank Xiaoli Dong and Marc Strous for bioinformatics support, the Centre

for Health Genomics and Informatics at University of Calgary for NextSeq sequencing,

Alexander Probst for providing the database of 16 syntenic ribosomal proteins, and Nina

Dombrowski and Brett Baker for providing a custom blast database for hydrocarbon degradation

genes.

Data availability

526   DNA sequences (amplicon sequences, genomes and raw sequence reads) have been deposited in

527   the NCBI BioProject database with accession number PRJNA415828 and PRJNASUB3936075

528   (https://www.ncbi.nlm.nih.gov/bioproject/). The authors declare that all other data supporting the

529   findings of this study are available within the article and its supplementary information files, or

530   from the corresponding authors upon request.

References

1. Petro, C., Starnawski, P., Schramm, A. & Kjeldsen, K.U. Microbial community assembly
in marine sediments. *Aquat Microb Ecol* **79**, 177-195 (2017).
2. Vigneron, A. et al. Comparative metagenomics of hydrocarbon and methane seeps of the
Gulf of Mexico. *Sci Rep* **7**, 16015 (2017).
3. Marshall, I.P.G., Karst, S.M., Nielsen, P.H. & Jørgensen, B.B. Metagenomes from deep
Baltic Sea sediments reveal how past and present environmental conditions determine
microbial community composition. *Mar Genomics* **37**, 58-68 (2018).
4. Dombrowski, N., Seitz, K.W., Teske, A.P. & Baker, B.J. Genomic insights into potential
interdependencies in microbial hydrocarbon and nutrient cycling in hydrothermal
sediments. *Microbiome* **5**, 106 (2017).
5. Lloyd, K.G. et al. Predominant archaea in marine sediments degrade detrital proteins.
Nature **496**, 215-218 (2013).
6. Orsi, W.D. Ecology and evolution of seafloor and subseafloor microbial communities.
Nat Rev Microbiol (2018).
7. Wang, Y. et al. Draft genome of an *Aerophobetes* bacterium reveals a facultative lifestyle
in deep-sea anaerobic sediments. *Sci Bull* **61**, 1176-1186 (2016).
8. Orcutt, B.N., Sylvan, J.B., Knab, N.J. & Edwards, K.J. Microbial ecology of the dark
ocean above, at, and below the seafloor. *Microbiol Mol Biol Rev* **75**, 361-422 (2011).
9. Inagaki, F. et al. Exploring deep microbial life in coal-bearing sediment down to ~2.5 km
below the ocean floor. *Science* **349**, 420-424 (2015).
10. Jørgensen, B.B. & Marshall, I.P. Slow Microbial Life in the Seabed. *Ann Rev Mar Sci* **8**,
311-332 (2016).
11. Jørgensen, B.B. & Boetius, A. Feast and famine--microbial life in the deep-sea bed. *Nat
Rev Microbiol* **5**, 770-781 (2007).
12. Arndt, S. et al. Quantifying the degradation of organic matter in marine sediments: A
review and synthesis. *Earth-Sci Rev* **123**, 53-86 (2013).
13. Orsi, W.D., Richards, T.A. & Francis, W.R. Predicted microbial secretomes and their
target substrates in marine sediment. *Nat Microbiol* **3**, 32-37 (2018).

14. Brooks, J.M. et al. Deep-sea hydrocarbon seep communities: evidence for energy and
nutritional carbon sources. *Science* **238**, 1138-1142 (1987).

15. Ijiri, A. et al. Deep-biosphere methane production stimulated by geofluids in the Nankai
accretionary complex. *Sci Adv* **4**, eaao4631 (2018).

16. Ruff, S.E. et al. Global dispersion and local diversification of the methane seep
microbiome. *Proc Natl Acad Sci U S A* **112**, 4015-4020 (2015).

17. Dong, X. et al. Fermentative Spirochaetes mediate necromass recycling in anoxic
hydrocarbon-contaminated habitats. *ISME J* **12**, 2039-2050 (2018).

18. Lin, Y.-S. et al. Towards constraining H_2 concentration in subseafloor sediment: A
proposal for combined analysis by two distinct approaches. *Geochim Cosmochim Acta* **77**,
186-201 (2012).

19. Sauvage, J., Flinders, A., Spivack, A. & D'Hondt, S. Global distribution of radiolytic H_2
production in marine sediment and implications for subsurface life. *AGU Fall Meeting*
Abstracts (2017).

20. Chakraborty, A. et al. Thermophilic endospores associated with migrated thermogenic
hydrocarbons in deep Gulf of Mexico marine sediments. *ISME J* **12**, 1895-1906 (2018).

21. Whiticar, M.J. Carbon and hydrogen isotope systematics of bacterial formation and
oxidation of methane. *Chem Geol* **161**, 291-314 (1999).

22. Killops, S.D. & Aljuboori, M.A.H.A. Characterization of the Unresolved Complex
Mixture (UCM) in the gas chromatograms of biodegraded petroleums. *Org Geochem* **15**,
147-160 (1990).

23. Gieg, L.M. & Toth, C.R.A. in Anaerobic Utilization of Hydrocarbons, Oils, and Lipids.
(ed. M. Boll) 1-30 (Springer International Publishing, Cham; 2017).

24. Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P. & Tyson, G.W. CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res* **25**, 1043-1055 (2015).

25. Ley, R.E. et al. Unexpected diversity and complexity of the Guerrero Negro hypersaline
microbial mat. *Appl Environ Microbiol* **72**, 3685-3695 (2006).

26. Baker, B.J., Lazar, C.S., Teske, A.P. & Dick, G.J. Genomic resolution of linkages in
carbon, nitrogen, and sulfur cycling among widespread estuary sediment bacteria.
Microbiome **3**, 14 (2015).

27.   Hu, P. et al. Genome-resolved metagenomic analysis reveals roles for candidate phyla
and other microbial community members in biogeochemical transformations in oil
reservoirs. *MBio* **7**, e01669-01615 (2016).

28.   Sieber, C.M.K. et al. Recovery of genomes from metagenomes via a dereplication,
aggregation and scoring strategy. *Nat Microbiol* **3**, 836-843 (2018).

29.   Parks, D.H. et al. A proposal for a standardized bacterial taxonomy based on genome
phylogeny. *bioRxiv* (2018).

30.   Quast, C. et al. The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic Acids Res* **41**, D590-596 (2013).

31.   Bradley, J.A., Amend, J.P. & LaRowe, D.E. Necromass as a limited source of energy for
microorganisms in marine sediments. *J Geophys Res-Biogeo* **123**, 577-590 (2018).

32.   Lipp, J.S., Morono, Y., Inagaki, F. & Hinrichs, K.U. Significant contribution of Archaea
to extant biomass in marine subsurface sediments. *Nature* **454**, 991-994 (2008).

33.   Boll, M. & Heider, J. in Handbook of Hydrocarbon and Lipid Microbiology. (ed. K.N.
Timmis) 1011-1024 (Springer Berlin Heidelberg, Berlin, Heidelberg; 2010).

34.   Tan, B. et al. Comparative analysis of metagenomes from three methanogenic
hydrocarbon-degrading enrichment cultures with 41 environmental samples. *ISME J* **9**,
2028-2045 (2015).

35.   Meckenstock, R.U. et al. Anaerobic degradation of benzene and polycyclic aromatic
hydrocarbons. *J Mol Microbiol Biotechnol* **26**, 92-118 (2016).

36.   Laso-Perez, R. et al. Thermophilic archaea activate butane via alkyl-coenzyme M
formation. *Nature* **539**, 396-401 (2016).

37.   Rabus, R. et al. Anaerobic Microbial Degradation of Hydrocarbons: From Enzymatic
Reactions to the Environment. *J Mol Microbiol Biotechnol* **26**, 5-28 (2016).

38.   Schouw, A. et al. *Abyssivirga alkaniphila* gen. nov., sp. nov., an alkane-degrading,
anaerobic bacterium from a deep-sea hydrothermal vent system, and emended
descriptions of *Natranaerovirga pectinivora* and *Natranaerovirga hydrolytica*. *Int J Syst
Evol Microbiol* **66**, 1724-1734 (2016).

39.   Khelifi, N. et al. Anaerobic oxidation of long-chain n-alkanes by the hyperthermophilic
sulfate-reducing archaeon, *Archaeoglobus fulgidus*. *ISME J* **8**, 2153-2166 (2014).

40. Porter, A.W. & Young, L.Y. in Advances in Applied Microbiology, Vol. 88. (eds. S.
Sariaslani & G.M. Gadd) 167-203 (Academic Press, 2014).

41. Dong, X. et al. Reconstructing metabolic pathways of a member of the genus
Pelotomaculum suggesting its potential to oxidize benzene to carbon dioxide with direct
reduction of sulfate. *FEMS Microbiol Ecol* **93**, fiw254-fiw254 (2017).

42. Holmes, D.E., Risso, C., Smith, J.A. & Lovley, D.R. Genome-scale analysis of anaerobic
benzoate and phenol metabolism in the hyperthermophilic archaeon *Ferroglobus*
placidus. *ISME J* **6**, 146-157 (2012).

43. Beulig, F., Røy, H., Glombitza, C. & Jørgensen, B.B. Control on rate and pathway of
anaerobic organic carbon degradation in the seabed. *Proc Natl Acad Sci U S A* **115**, 367-
372 (2018).

44. Sewell, H.L., Kaster, A.K. & Spormann, A.M. Homoacetogenesis in Deep-Sea
Chloroflexi, as inferred by single-cell genomics, provides a link to reductive
dehalogenation in terrestrial *Dehalococcoidetes*. *MBio* **8** (2017).

45. He, Y. et al. Genomic and enzymatic evidence for acetogenesis among multiple lineages
of the archaeal phylum *Bathyarchaeota* widespread in marine sediments. *Nat Microbiol* **1**,
16035 (2016).

46. Adam, P.S., Borrel, G. & Gribaldo, S. Evolutionary history of carbon monoxide
dehydrogenase/acetyl-CoA synthase, one of the oldest enzymatic complexes. *Proc Natl*
Acad Sci U S A **115**, E1166-E1173 (2018).

47. Lever, M.A. Acetogenesis in the energy-starved deep biosphere - a paradox? *Front*
Microbiol **2**, 284 (2011).

48. Yu, T. et al. Growth of sedimentary *Bathyarchaeota* on lignin as an energy source. *Proc*
Natl Acad Sci U S A **115**, 6022-6027 (2018).

49. Berney, M., Greening, C., Conrad, R., Jacobs, W.R., Jr. & Cook, G.M. An obligately
aerobic soil bacterium activates fermentative hydrogen production to survive reductive
stress during hypoxia. *Proc Natl Acad Sci U S A* **111**, 11479-11484 (2014).

50. Ma, K., Hao, Z. & Adams, M.W.W. Hydrogen production from pyruvate by enzymes
purified from the hyperthermophilic archaeon, *Pyrococcus furiosus*: A key role for
NADPH. *FEMS Microbiol Lett* **122**, 245-250 (1994).

51.   Greening, C. et al. Genomic and metagenomic surveys of hydrogenase distribution
652   indicate H_2 is a widely utilised energy source for microbial growth and survival. *ISME J*
653   **10**, 761 (2015).

654   52.   Wagner, T., Koch, J., Ermler, U. & Shima, S. Methanogenic heterodisulfide reductase
655   (HdrABC-MvhAGD) uses two noncubane [4Fe-4S] clusters for reduction. *Science* **357**,
656   699-703 (2017).

657   53.   Adhikari, R.R. et al. Hydrogen utilization potential in subsurface sediments. *Front*
658   *Microbiol* **7**, 8 (2016).

659   54.   Kawai, M. et al. High frequency of phylogenetically diverse reductive dehalogenase-
660   homologous genes in deep subseafloor sedimentary metagenomes. *Front Microbiol* **5**, 80
661   (2014).

662   55.   Dolfing, J., Larter, S.R. & Head, I.M. Thermodynamic constraints on methanogenic
663   crude oil biodegradation. *ISME J* **2**, 442-452 (2008).

664   56.   Canfield, D.E., Thamdrup, B. & Hansen, J.W. The anaerobic degradation of organic
665   matter in Danish coastal sediments: iron reduction, manganese reduction, and sulfate
666   reduction. *Geochim Cosmochim Acta* **57**, 3867-3883 (1993).

667   57.   Liu, Y.F. et al. Metabolic capability and in situ activity of microorganisms in an oil
668   reservoir. *Microbiome* **6**, 5 (2018).

669   58.   Colman, D.R., Poudel, S., Stamps, B.W., Boyd, E.S. & Spear, J.R. The deep, hot
670   biosphere: Twenty-five years of retrospection. *Proceedings of the National Academy of*
671   *Sciences* **114**, 6895-6903 (2017).

672   59.   Bernard, B.B. et al. Surface geochemical exploration and heat flow surveys in fifteen (15)
673   frontier Indonesian basins. *Thirty-Second Annual Convention & Exhibition May 2008*
674   (2008).

675   60.   Klindworth, A. et al. Evaluation of general 16S ribosomal RNA gene PCR primers for
676   classical and next-generation sequencing-based diversity studies. *Nucleic Acids Research*
677   **41**, e1-e1 (2013).

678   61.   Kleiner, M. et al. Assessing species biomass contributions in microbial communities via
679   metaproteomics. *Nat Commun* **8**, 1558 (2017).

680   62.   Nurk, S., Meleshko, D., Korobeynikov, A. & Pevzner, P.A. metaSPAdes: a new versatile
681   metagenomic assembler. *Genome Res* **27**, 824-834 (2017).

682   63.   Kang, D.D., Froula, J., Egan, R. & Wang, Z. MetaBAT, an efficient tool for accurately

683   reconstructing single genomes from complex microbial communities. *PeerJ* **3**, e1165

684   (2015).

685   64.   Parks, D.H. et al. Recovery of nearly 8,000 metagenome-assembled genomes

686   substantially expands the tree of life. *Nat Microbiol* **2**, 1533-1542 (2017).

687   65.   Hyatt, D. et al. Prodigal: prokaryotic gene recognition and translation initiation site

688   identification. *BMC Bioinformatics* **11**, 119 (2010).

689   66.   Lombard, V., Golaconda Ramulu, H., Drula, E., Coutinho, P.M. & Henrissat, B. The

690   carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Res* **42**, D490-495

691   (2014).

692   67.   Yin, Y. et al. dbCAN: a web resource for automated carbohydrate-active enzyme

693   annotation. *Nucleic Acids Res* **40**, W445-451 (2012).

694   68.   Rawlings, N.D. et al. The MEROPS database of proteolytic enzymes, their substrates and

695   inhibitors in 2017 and a comparison with peptidases in the PANTHER database. *Nucleic*

696   *Acids Res* **46**, D624-D632 (2018).

697   69.   Søndergaard, D., Pedersen, C.N. & Greening, C. HydDB: A web tool for hydrogenase

698   classification and analysis. *Sci Rep* **6**, 34212 (2016).

699   70.   Boyd, J.A., Woodcroft, B.J. & Tyson, G.W. GraftM: a tool for scalable, phylogenetically

700   informed classification of genes within metagenomes. *Nucleic Acids Res* **46**, e59 (2018).

701   71.   Buchfink, B., Xie, C. & Huson, D.H. Fast and sensitive protein alignment using

702   DIAMOND. *Nature Methods* **12**, 59 (2014).

703   72.   Anantharaman, K. et al. Expanded diversity of microbial groups that shape the

704   dissimilatory sulfur cycle. *ISME J* **12**, 1715-1728 (2018).

705   73.   Tully, B.J., Sachdeva, R., Graham, E.D. & Heidelberg, J.F. 290 metagenome-assembled

706   genomes from the Mediterranean Sea: a resource for marine microbiology. *PeerJ* **5**,

707   e3558 (2017).

708   74.   Kato, S. et al. Genome-enabled metabolic reconstruction of dominant chemosynthetic

709   colonizers in deep-sea massive sulfide deposits. *Environ Microbiol* **20**, 862-877 (2018).

710   75.   Price, M.N., Dehal, P.S. & Arkin, A.P. FastTree 2--approximately maximum-likelihood

711   trees for large alignments. *PLoS One* **5**, e9490 (2010).

712   76.   Anantharaman, K. et al. Thousands of microbial genomes shed light on interconnected
713   biogeochemical processes in an aquifer system. *Nat Commun* **7**, 13219 (2016).

714   77.   Zaremba-Niedzwiedzka, K. et al. Asgard archaea illuminate the origin of eukaryotic
715   cellular complexity. *Nature* **541**, 353-358 (2017).

716   78.   Probst, A.J. et al. Differential depth distribution of microbial function and putative
717   symbionts through sediment-hosted aquifers in the deep terrestrial subsurface. *Nat*
718   *Microbiol* **3**, 328-336 (2018).

719   79.   Hug, L.A. et al. A new view of the tree of life. *Nat Microbiol* **1**, 16048 (2016).

720   80.   Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
721   large phylogenies. *Bioinformatics* **30**, 1312-1313 (2014).

722   81.   Miller, M.A., Pfeiffer, W. & Schwartz, T. Creating the CIPRES Science Gateway for
723   inference of large phylogenetic trees. *2010 Gateway Computing Environments Workshop*
724   *(GCE)*, 1-8 (2010).

725   82.   Letunic, I. & Bork, P. Interactive tree of life (iTOL) v3: an online tool for the display and
726   annotation of phylogenetic and other trees. *Nucleic Acids Res* **44**, W242-245 (2016).

727   83.   Edgar, R.C. MUSCLE: multiple sequence alignment with high accuracy and high
728   throughput. *Nucleic Acids Res* **32**, 1792-1797 (2004).

729   84.   Kumar, S., Stecher, G. & Tamura, K. MEGA7: Molecular Evolutionary Genetics
730   Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* **33**, 1870-1874 (2016).

731   85.   Madigan, M.T., Martinko, J.M. & Parker, J. Brock biology of microorganisms, Vol. 13.
732   (Pearson, 2017).

733   86.   Xu, W.Y., Lowell, R.P. & Peltzer, E.T. Effect of seafloor temperature and pressure
734   variations on methane flux from a gas hydrate layer: Comparison between current and
735   late Paleocene climate conditions. *J Geophys Res-Sol Ea* **106**, 26413-26423 (2001).

736   87.   Dolfing, J. in Hydrocarbon and Lipid Microbiology Protocols. (eds. T.J. McGenity, K.N.
737   Timmis & B. Nogales Fernández) 155-163 (Springer Berlin Heidelberg, Berlin,
738   Heidelberg; 2015).

 Tables and Figures

740 **Table 1 Geochemical description of sediment samples from Sites E26, E29 and E44.** TSF

741 Max: total scanning fluorescence maximum intensity. UCM: uncharacterized complex mixture.

742 Σn-Alk: sum of C_{15}-C_{34} n-alkanes. ΣAlk Gas: total alkane gases. C_{2+} Alk: sum of alkane gases

743 larger than methane. T/D: thermogenic/diagenetic n-alkane ratio. BDL: below detection limit.

744 NA: not analyzed.

Core ID	Site E26	Site E29	Site E44
Latitude (N)	26.59	27.43	26.28
Longitude (W)	87.51	86.01	86.81
Water depth (km)	2.8	3.2	3.0
Sulfate (mM)	20.01	33.73	31.72
Benzoate (nM)	93.6	22.6	161.7
Succinate (nM)	11.7	5.0	16.6
Acetate (μM)	BDL	BDL	BDL
Chloride ($g\ L^{-1}$)	21.04	20.15	21.05
Total Scanning Fluorescence MAX	57326.7	26738.3	13502.3
Unresolved Complex Mixture ($\mu g\ g^{-1}$)	32	13	7.3
Σn-Alkanes ($ng\ g^{-1}$)	2845.3	2527	1045
Thermogenic/Diagenetic Ratio	1.0	2.6	0.8
ΣAlkane Gas (ppm)	9	36012	9.9
C_{2+} Alkanes (ppm)	0.3	17.5	0.5
$C_1/(C_2+C_3)$	NA	3974.2	NA
$\delta^{13}CH_4$ (‰, vs. PDB)	NA	-85.1	NA
H_2 (ppm)	BDL	BDL	BDL

Figure 1 Relative frequency of metagenomic sequence reads for different marker genes at

Sites E26, E29 and E44. (a) Community compositions based on reconstruction of full-length

16S rRNA genes from the metagenomes. Eukaryotes and unassigned reads are not shown. (b)

Relative occurrences of hydrogenases with different metal co-factors. (c) Relative occurrences of

different subtypes of NiFe hydrogenases. (d) Relative occurrences of *mcrA* genes indicative of

different types of methanogenesis.

Figure 2 Phylogenetic placement of 82 reconstructed metagenome-assembled genomes. A

maximum-likelihood phylogenomic tree was built based on concatenated amino acid sequences

of 43 conserved single copy genes using RAxML with the PROTGAMMALG model. Sequences

of *Altiarchaeales* ex4484_43 were used as an outgroup. The scale bar represents 1 amino acid

substitution per sequence position. Bootstrap values > 70% are indicated. Blue for Site E26

(E26_binX), red for Site E29 (E29_binY), and green for Site E44 (E44_binZ).

Figure 3 Identification of functional genes or pathways present in MAGs. The presence of

genes or pathways are indicated by orange shaded boxes. Aor, aldehyde:ferredoxin

oxidoreductase; Kor, 2-oxoglutarate/2-oxoacid ferredoxin oxidoreductase; Por,

pyruvate:ferredoxin oxidoreductase; Ior, indolepyruvate ferredoxin oxidoreductase. GHs,

glycoside hydrolases. Presence of the Embden-Meyerhof-Parnas pathway is indicated if at least

five genes were detected. Presence of the Beta-oxidation pathway is indicated if at least three

genes were detected. AssA, catalytic subunit of alkylsuccinate synthase. CmdA, catalytic subunit

of p-cymene dehydrogenase; AhyA, catalytic subunit of alkane C2-methylene hydroxylase.

Additional detail for the central benzoyl-CoA degradation pathway can be found in Figure 4.

Presence of the Wood-Ljungdahl pathway is indicated if at least four genes were detected.

767 Presence of the TCA cycle is indicated if at least six key genes were detected. Lactate and

768 ethanol fermentation are indicated if genes encoding respective dehydrogenases were detected.

769 H_2ase, hydrogenase. DsrAB, dissimilatory sulfite reductase. More details about these functional

770 genes and pathways can be found in the text and in Table S6.

771 **Figure 4 Evidence for anaerobic hydrocarbon degradation in MAGs.** (a) Phylogenetic

772 relationship of identified genes in MAGs with currently known alkyl-arylalkylsuccinate

773 synthases based on the respective catalytic alpha-subunits. Ass/Mas, n-alkanes (1-methylalkyl)

774 succinate synthase; Nms, 2-naphthylmethyl succinate synthase; Bss, benzyl succinate synthase;

775 Ibs, 4-isopropylbenzyl succinate synthase; Hbs, 4-hydroxybenzyl succinate synthase. Sequences

776 of pyruvate formate lyase (Pfl) from *E. coli* were used as an outgroup. The scale bar represents

777 0.1 amino acid substitutions per sequence position. Bootstrap values > 70% are indicated. The

778 full sequences can be found in Text S1. (b) Summary of identified enzymes involved in central

779 benzoyl-CoA processing in anaerobic aromatic hydrocarbon biodegradation. The MAGs were

780 shown only if it was at least partially complete (presence of at least three subunits within one

781 cluster for BcrABCD). Presence of genes or pathways are indicated by green boxes. Bcr,

782 benzoyl-CoA reductase; Oah, 6-oxo-cyclohex-1-ene-carbonyl-CoA hydrolase; Dch, cyclohex-

783 1,5-diencarbonyl-CoA hydratase; Had, 6-hydroxycyclohex-1-ene-1-carbonyl-CoA

784 dehydrogenases.

785 **Figure 5 Thermodynamic constraints on anaerobic benzoate and hexadecane degradation**

786 **in deep sea sediments.** Two possible scenarios are illustrated depending on the end products

787 based on metabolic predictions in Figure 3: (1) fermentation with production of hydrogen and

788 acetate, and (2) fermentation with production of acetate alone. Thermodynamics for each

reaction are indicated by a line in its corresponding color. If $\Delta G' < 0$, the reaction is energetically

favorable (yellow-shaded area), and if $\Delta G' > 0$ the reaction is assumed not to occur. The graph

shows that $\Delta G'$ for hexadecane fermentation to acetate alone (green reaction) will not reach

negative values unless the concentration for acetate is extremely low (far lower than the

detection limit for acetate of 2.5 µM in this study, see dash line) such that the other three

reactions are more realistic scenarios for anaerobic hydrocarbon degradation in the marine

sediments studied here.

Figure 6 Common organotrophic and hydrogenotrophic potentials of three hydrocarbon-

impacted microbial communities inferred from metagenomics and metabolomics.

Figure 1

Figure 2

Figure 3

<u>Figure 4</u>

Figure 5

Figure 6