The role of land use types and water chemical properties in structuring the microbiomes of a connected lake system

Sophi Marmen[1], Lior Blank[2], Ashraf Al-Ashhab[1,3], Assaf Malik[4], Lars Ganzert[5], Maya Lalzar[4], Hans-Peter Grossart[5,6] and Daniel Sher[1*]

[1] *Department of Marine Biology, Leon H. Charney School of Marine Sciences, University of Haifa, Israel*

[2] *Department of Plant Pathology and Weed Research, ARO, Volcani Center, Rishon Lezion, Israel*

[3] *Microbial Metagenomics Division, Dead Sea and Arava Science Center, Masada, Israel*

[4] *Bioinformatics Service Unit, University of Haifa, Israel*

[5] *Leibnitz Institute for Freshwater Ecology and Inland Fisheries, Department of Experimental Limnology, Stechlin, Germany*

[6] *Potsdam University, Institute of Bicochemistry and Biology, Maulbeerallee 2, 24469 Potsdam, Germany*

[*]Corresponding author: Dr. Daniel Sher, Department of Marine Biology, Leon H. Charney School of Marine Sciences, University of Haifa, Haifa 31905, Israel. email: dsher@univ.haifa.ac.il

Number of words- 7,188

Number of figures- 6

Running title: Land use and lake microbiome structur

Abstract

Lakes and other freshwater bodies are intimately connected to the surrounding land, yet to what extent land-use affects the quality of freshwater and the microbial communities living in various freshwater environments is largely unknown. We address this question through an analysis of the land use surrounding 46 inter-connected lakes located within 7 different drainage basins in northern Germany, and the microbiomes of these lakes during early summer. Lake microbiome structure was not determined by the specific drainage basin or by basin size, and bacterial distribution did not seem to be limited by distance. Instead, land use within the drainage basin could predict, to some extent, NO_2+NO_3 concentrations in the water, which (together with temperature, *chlorophyll a* and total phosphorus) affected water microbiome structure. Land use directly surrounding the water bodies, however, had little observable effects on water quality or the microbiome. Several microbial lineages, including environmentally important Cyanobacteria and Verrucomicrobia, were differentially partitioned between the lakes. As the amount of available data on land use (e.g. from remote sensing) increases, identifying relationships between land use, aquatic microbial communities

40 and their effect on water quality will be important to better manage freshwater resources
41 worldwide, e.g. by systemically identifying water bodies prone to ecological changes or the
42 presence of harmful organisms.
43
44 **Keywords:** microbiome; lakes; water parameters; land use; distribution; Cyanobacteria
45

1. Introduction

The world's population growth is maintained by a suite of human enterprises such as
agriculture, industry, fishing and international commerce (Vitousek*, et al.*, 1997). The use of
land for such enterprises has greatly altered the way ecosystems interact with the surrounding
land, atmosphere and aquatic systems (Vitousek*, et al.*, 1997). Freshwater ecosystems, in
particular, may be highly sensitive to human impact, mainly through eutrophication.
The microbial communities ("microbiomes") living in freshwater ecosystems are intimately
connected with water quality. Microbes take up, utilize and recycle elements such as carbon,
nitrogen, phosphorus and sulfur, and thus control and modulate local and global elemental
cycles (Newton*, et al.*, 2011). Some aquatic microorganisms are potential pathogens of
animals or humans, or can produce secondary metabolites or toxins (Abdelzaher*, et al.*, 2010,
Cabral, 2010, Valério*, et al.*, 2010). Thus, significant effort has been invested in elucidating
the environmental drivers that determine the composition and function of freshwater
microbiomes (reviewed by, e.g., Logue & Lindström, 2008, Williamson*, et al.*, 2009,
Newton*, et al.*, 2011, Zeglin, 2015). Environmental drivers of microbial community structure
may act at the local scale, where the physical-chemical characteristics of the water lead to a
"species sorting" process which enrich for bacteria adapted to occupy that specific niche
(Jones & McMahon, 2009, Langenheder & Székely, 2011, Van Rossum*, et al.*, 2015). Such
physical - chemical properties can include, for example, nutrient, organic carbon or metal
concentrations, pH, temperature, salinity and light intensity (Crump*, et al.*, 2004, Horner-
Devine*, et al.*, 2004, Yannarell & Triplett, 2005, Zeglin, 2015). The physical-chemical
properties of water depend, in turn, at least partly on the input of nutrients, organic matter and
other contaminants from the surrounding terrestrial ecosystem via the water shed (Chen*, et
al.*, 2018). Specifically, different types of land use surrounding the water bodies (e.g., forests,
agricultural lands, urban lands, etc.) can strongly affect the quantity and quality of terrestrial
input into water bodies (reviewed by (Solomon*, et al.*, 2015).
Local processes can be affected by processes occurring at larger spatial and temporal scales.
For example, physical connectivity between water bodies (e.g., within the same drainage
basin) may lead to bacterial transfer between environments, a process termed "mass effect",
(e.g., Quinn*, et al.*, 1997, Nelson*, et al.*, 2009, Crump*, et al.*, 2012, Lindström &
Langenheder, 2012, Logue*, et al.*, 2012). In very strongly connected habitats, dispersal rates
may be so high that they lead to ecosystem homogenization (Leibold & Norberg, 2004). As
the spatial scales increase, geographic distances and physical barriers may lead to limitations
on bacterial dispersal, resulting in "distance decay" patterns, whereby differences between
microbial communities increase with distance (Van der Gucht*, et al.*, 2007, Lear*, et al.*, 2013,
Niño-García*, et al.*, 2016, Marcelino*, et al.*, 2017, Logares*, et al.*, 2018). As the various
processes affecting freshwater microbial ecosystems are identified and characterized, many
questions remain: to what extent are local conditions within water bodies affected by those in
the surrounding terrestrial system? What is the relative importance of local conditions in
determining microbial community structure? What is the role of mass transport of microbes
from upstream sites, and how does this differ between freshwater systems? Are all microbial
taxa affected in the same manner by these processes? While many of these questions have

been addressed in ecosystems ranging from highly impacted agricultural lands (e.g. Griffin*,*
et al., 2017) to pristine Antarctic lakes (Logares*, et al.*, 2018), few studies have aimed to
integrate the effects of hydrology and water chemistry on lake biology in the context of land-
use within their respective catchment basins (e.g. Catherine*, et al.*, 2016). Answering some of
these questions is critical as we aim to understand how changes in the environment, including
land use, affect aquatic ecosystems.
In this study, we investigated the microbiomes of 46 German lakes, located in a network of 7
drainage basins connected through streams. The region encompassing these lakes and basins
comprises a tapestry of different natural and artificial land parcels (e.g., forests, pastures,
agricultural lands, water bodies, towns and villages, Figure 1). Our major goal was to
understand what shapes microbial population in such a system of connected natural water
bodies, located within relatively short distances and surrounded by a variety of different land-
use types. We hypothesize that: 1) land use directly affects water chemistry; 2) water
chemistry affects bacterial composition by habitat filtering of bacteria adapted to specific
water properties; 3) bacterial populations gradually change as the water moves downwards
within the connected lakes and streams; 4) there may be also a direct effect of land use types
on the water microbiome via direct input of live bacteria.
105
106
107

2. Materials and methods

2.1 Study region and delineation of drainage basins

The sampling area lies within the region of Brandenburg, in northern Germany (Figure 1,
Supplementary Figure 1, Supplementary text). The area is relatively flat (18-89 meters above
sea level) and contains hundreds of lakes, ponds and depressions that were formed by glacial
action during the Pleistocene. Three main basins drain this region, which are (from west to
east) the Rhin (which flows north), the Havel and the Ucker, the two latter flowing south in
the region of sampling. These three main basins can be subdivided into smaller basins. The
drainage basin of each lake was defined as the line connecting the highest points from which
the surface water flow enters each of the lakes, including all of the upstream lakes. We note
that groundwater flow may be significant in many lakes, and often does not flow as surface
water does in accordance with topographic basins (Winter*, et al.*, 2003). We also assume that
the water is fully mixed horizontally within each separate water body, and that our sampling
represents the epilimnion if the lakes are stratified. A detailed description of the sampling
region, methods of delineating the basins, identification of the canals connecting the basins
and GIS layer analysis, can be found in the supplementary information.
124

2.2. Sampling and analysis of environmental data

Each location was sampled once from the edge of the water body, as due to the range of
water bodies sampled and the inaccessibility to some by boat sampling from the deepest point
was not feasible. During sampling, dissolved oxygen, temperature pH and turbidity were
measured using a YSI multi probe (YSI 6920 V2-2; YSI Inc). At all locations, 5 liters of
surface water were collected using plastic jerry-cans that were washed with double distilled
water then with 70% ethanol and air dried prior to sampling. Care was taken not to raise any
sediment. Samples were brought to the lab and filtered within two hours. For *Chlorophyll a*
(chl a) measurement, water was filtrated on GF/C filters (Whatman glass fiber, 25 mm,
nominal pore size 1.2 μm) and placed into 1.5 mL sterile Eppendorf tubes. The water was

filtered until the filters were clogged, and the volume was recorded (50-800 mL). *Chlorophyll a* was extracted in absolute methanol for 12 h at room temperature in the dark, and the concentration was determined spectrophotometrically according to (Ritchie, 2008). For DNA extraction, water was filtered using a vacuum pump on GF/F filters (Whatman glass fiber, 25 mm, nominal pore size 0.7 μm). Preliminary experiments using a culture of the marine Pico-Cyanobacterium *Prochlorococcus* MED4 (~0.5 μm in diameter) showed that with the vacuum pumps we used approximately 99.9% of the cells of this size were retained on a GF/F membrane. The filters were placed into 1.5 mL sterile Eppendorf tubes with 1 mL of storage buffer (40 mM EDTA, 50 mM Tris-HCl, 0.75 M Sucrose) and stored at -80ºC until analyzed. The filtrate from the GF/F filters was collected for measurement of dissolved nutrients, and unfiltered water was used for total phosphorus measurements according to (Grasshoff & Kremling, 1983, Grasshoff, *et al.*, 2009) DNA extraction was performed using bead-beating followed by robotic extraction using a QiaCube robot, the gene encoding the 16S subunit of the ribosomal RNA was amplified using primers targeting the V3-V4 region, and the amplicons were sequenced using the 2×250 base pair format on a MiSeq flow cell (V3 chemistry). More details on the DNA extraction and 16S amplification procedures are found in the supplementary methods section.

2.3 Generalized Linear Model Analysis (GLM)

We used multivariable regression model in the framework of GLMs to relate the six land use types to the measured water chemical parameters. Multi-model inference based on the Akaike Information Criterion (AIC) was used to rank the importance of variables (Burnham & Anderson, 2003, Blank & Blaustein, 2014). The package ''glmulti'' was used to facilitate multi-model inference based on every possible first-order combination of the predictors in each scale (Calcagno & de Mazancourt, 2010). The coefficients associated with each variable and their relative importance were assessed using a multi-model average. Parameters were estimated by averaging across models, weighted by the probability that their originating models are the best in the set. Hence, the relative importance of a parameter is not based on *P*-values, but rather on its occurrence in the models that are best supported by the data. The importance value for a particular predictor is equal to the sum of the weights for the models in which the variable appeared. Thus, a variable that shows up in many of the models with large weights received a high importance value.

2.4 16S Sequence analysis and statistical tests

Sequences received from the sequencing facility were quality filtered to remove PhiX DNA, unmerged reads and low quality sequences, followed by classification into different Operational Taxonomic Units (OTUs) based on 97% sequence similarity threshold using the "pick_de_novo_otus.py" command in Qiime (Caporaso, *et al.*, 2010). OTUs with fewer than 5 reads across all samples were removed, resulting in 35,775 – 75,704 reads per sample. Alpha diversity (Shannon index) was calculated using the ''diversity'' function, and Bray–Curtis dissimilarity by ''vegdist'' function. Non-metric multidimensional scaling analysis (nMDS) was carried out in R with ''metaMDS'' function and a Bray-Curtis matrix. In order to relate environmental factors to specific OTUs we performed variation partitioning ("varpart" function in R) followed by canonical correspondence analaysis (CCA) using the "cca" and "anova.cca" functions in the "vegan" R package, on subsamples of 25,000 reads for each location which were then log10 transformed. For additional information on these analyses, see the supplementary methods.

2.5 Permutation test of phyla clustering significance

In CCA ordinations, the location of each OTU within the ordination is related to the effect of the variables constraining the CCA on that OTU. We wanted to address a different question - whether a group of OTUs (e.g. a phylogenetic lineage) behave coherently with respect to the environmental characteristics. In other words, we do not ask, for example, whether "a cyanobacterial OTU" is related to the CCA axes (as that is the direct output of the CCA), but whether the cyanobacteria as a group behave similarly with respect to these axes. We therefore designed a permutation test to determine: (a) the clustering significance of the OTUs within each clade; (b) the significance of association with the tested environmental factors. We obtained the coordinates of eigenvectors (e.g. pH, temperature, etc.), and of all OTUs belonging to the different clades, in the CCA data. For each clade p, we calculated two scores: d_p and $d_{p,v}$, where v is a given eigenvector. The score d_p equals the sum of Euclidean distances between all pairs of OTUs in p. The score $d_{p,v}$ is the sum of scalar projections of all OTUs in p, on a given eigenvector v. Accordingly: (a) for the clustering test, s_p OTUs were randomly chosen from all tested clades, to obtain the score $d_{p=random}$, where s_p is the count of OTUs in clade p. This calculation was repeated n times ($n=1000$), where the count of cases where $d_p \geq d_{p=random}$, divided by n, was defined as p-value; (b) for the association test, The score $d_{p,v}$ was compared to the score $d_{p=random,v}$ which was obtained from random coordinates normally distributed with the same x,y variance as the x,y variance as in clade p, and the same number of s_p datapoints. This calculation was repeated n times ($n=100$), where the count of cases with $d_{p,v} \geq d_{p=random,v}$ divided by n, was defined as p-value. Here, p-value < 0.025 or > 0.975 (after Benjamini-Hochberg correction for multipole hypothesis testing) indicates negative and positive association with the tested environmental factor.

3. Results

3.1 General characterization of the sampled water bodies
During the summer of 2015, we sampled 46 water bodies (lakes and one fish pond) within a relatively small area (~150 km by 100 km) of the Brandenburg/Mecklenburg lake district in northeastern Germany (Supplementary Excel file 1). The lakes ranged in trophic status from meso-oligotrophic (Lake Stechlin) to polytrophic (hyper-eutrophic), with total phosphorus concentrations between 0.009-0.589 mg/L and *chl* a concentrations ranging between 0.0005-0.320 mg/L. The water bodies belonged primarily to five major drainage basins (Figure 1, Supplementary Figure 1), and varied widely in their sizes and nutrient concentrations (Supplementary Data file 1). While the microbial diversity, expressed as the Shannon alpha-diversity index, between the basins was relatively similar (4.88-7, Figure 2A), clear differences were observed in community composition between the various lakes (Figure 2B). Comparing the microbiome composition between lakes (using the Bray- Curtis dissimilarity index) revealed that they can be broadly divided into two main groups: a cluster of sites with a high relative abundance of Cyanobacteria, and a second cluster with a high relative abundance of Actinobacteria, Proteobacteria and Bacteoidetes and low relative abundance of Cyanobacteria (Figure 2B).

Lakes are often broadly divided into trophic states, using parameters such as *chl* a or total phosphorus (TP) concentrations, and these trophic definitions are often used in water management. As shown in Supplementary Figure 2, the trophic status of the lakes based on *chl* a and total P concentrations was broadly in agreement with estimates based on multi-annual monitoring by the German federal states of Mecklenburg-Vorpommern and Brandenburg, although more of the lakes had an oligotrophic character at the time of

sampling. This is likely because we sampled during early summer where high production rates of zooplankton can reduce phytoplankton abundance, leading to a "clear water phase" and masking the real, higher trophic status of lakes high in nutrients. A slight tendency was seen for lakes with higher *chl* a and total P concentrations to have a higher relative abundance of Cyanobacteria (Supplementary Figure 2) manifested also as an observable trend in the NMDS ordination (Supplementary Figure 3A).

Interestingly, lakes from the same drainage basins did not cluster together, despite relatively large differences between the drainage basins in terms of their size and land-use in their water shed (Figure 2B, Supplementary Figure 3B, Supplementary Data File 1). Neither was a clear correlation observed between size of the drainage basin of each water body and chemistry of the water (Supplementary Figure 4A-F) or lake microbiome (Supplementary Figure 4G). One major factor, that might determine to what extent ecological communities differ, is the geographical distance between them. This phenomenon, identified by a "distance decay " relationship in community similarity (Nekola & White, 1999, Bell, 2010), has been demonstrated for both macro- and micro-organisms at various scales of geographic distances (Lear, *et al.*, 2013, Marcelino, *et al.*, 2017). No distance decay relationships were observed in our samples, neither within a single basin nor between samples belonging to different basins (Figure 2C). The lack of distance-decay patterns was still observed when the three major basins, Lychen, Rhin and Havel, were divided into sub-basins where all of the water bodies were directly connected (Supplementary Figure 5). Moreover, examination of the total species diversity in our landscape (i.e. γ-diversity) showed that 27% of the OTUs are core OTUs that were present among all sites, and that additional 22% OTUs were cosmopolitan, defined as being observed in >90% of the sites (Figure 2D). Rare and site unique OTUs combined represented only 3.4% of all OTUs. These rare and unique OTUs were also numerically scarce, representing at most 2.6% of the total sequence reads. The core microbiome was dominated by Actinobacteria and Proteobacteria (43% and 39% of the OTUs respectively, Supplementary Figure 6A), and contained very few Cyanobacteria OTUs (1%). In contrast, Cyanobacteria represented the dominant phylum of all cosmopolitan OTUs (47%). Thus, it seems that differences between microbial communities are primarily due to changes in the relative contribution of OTUs rather than to the presence of unique or endemic OTUs in different regions.

3.2 Land use in the drainage basins affects water chemistry, and through it, the microbiome

We next asked whether specific land use types may affect environmental conditions within the water body (e.g. nutrients, temperature, etc., referred to here as "aquatic environmental conditions", see supplementary information for the definition of the land-use types used). The effect of land use could be manifested at the level of the entire drainage basin (summing up the input of the entire draining into each water body), or, alternatively, at the level of local land use (adjacent to the water body). To test our hypotheses, we used General Linear Models (GLMs) to determine whether specific land use types could be associated with specific aquatic environmental conditions. On the basin scale, six land use types (agricultural area, pasture lands, arable lands, urban and forest areas and water bodies) explained a high proportion of the variability of nitrite and nitrate concentrations (NO_2+NO_3, $R_N^2= 0.74$, Figure 3, Supplementary Table 1), with the fraction of urban land being positively correlated to NO_2+NO_3 concentrations (Supplementary Table 1). When quantifying the independent contribution of each different land use separately, urban was the only significant variable explaining 74% of the variation in $NO_2+ NO_3$.

We next used VPA to explore whether and to what extent basin-level and local-level land
use types, as well as aquatic environmental conditions, explain the composition of the aquatic
microbiome. VPA results pointed to aquatic environmental parameters as key parameters, in
total explaining 15.5% of the variance in microbiome composition among all sites, 11%
independent of land-use parameters (Figure 4). In contrast, land-use types (at both basin and
local scale) had a very weak explanatory power for microbial community structure (Figure 4).
Following VPA results, we tested measured aquatic environmental parameters as explanatory
model for community composition using CCA followed by CCA permutation test
(Supplementary Table 4, Figure 5). The aquatic environmental parameter model explained
24.2% of the inertia in CCA and was significant ($P<0.05$). Similar to the NMDS
(unconstrained) ordination (Supplementary Figure 3A), the CCA did not show clear
separation of sampling sites by basin (Figure 5b). Four out of six aquatic environmental
parameters in our model were significant: total phosphorus (TP, $P < 0.01$) and *Chl a* ($P <$
0.05) concentration were positively associated with CCA1 (eigenvector for CCA1 and CCA2
were 0.89, -0.12 for *Chl a* and 0.45, -0.16 for TP), NO_2+NO_3 ($P < 0.05$) was positively
associated with CCA2 (eigenvector 0.30, 0.58) and water temperature ($P < 0.01$) was
negatively associated with CCA2 (eigenvector 0.08, -0.84) (Figure 5a, Supplementary Table
4). Notably, NO_2+NO_3 were identified by the GLMs as being significantly affected by land
use in the drainage basins.

3.3 Specific bacterial taxonomic groups are associated with sampling locations differing
in their aquatic environmental conditions

Having identified TP, *Chl* a, NO_2+NO_3 and temperature as variables affecting the
microbiome as a whole, we next asked whether we could identify specific microbial lineages
affected by these environmental conditions. To answer this question, we analyzed the
distribution patterns of OTUs belonging to different bacterial lineages along the CCA
ordination axes. Most phyla were widely distributed in the CCA biplot (Figure 5), suggesting
that, within each phylum, OTUs were affected differently by environmental factors driving
the CCA ordination. However, Actinobacteria OTUs formed a tight cluster whose center was
on the negative side of the x-axis (Figure 5C). This suggested that all OTUs in this group
were affected similarly by the measured environmental variables. To statistically test this, we
developed a permutation test to determine: a) whether OTUs in each lineage were clustered
together more than expected (i.e. the distances between them were smaller than those
between randomly selected OTUs. This suggests a coherent relationship with the factors
constraining the CCA at the lineage level); b) whether the OTUs were associated (positively
or negatively) with any of the eigenvectors of the specific aquatic parameters (i.e. the arrows
in Figure 5B). We then applied these two tests to lineages at different phylogenetic resolution
(phylum, class, order and family) to identify those that are coherently affected by the four
environmental parameters tested (Supplementary Data File 2).
Of the five most abundant phyla, only Actinobacteria were not randomly distributed in the
CCA plot, as determined by permutation analysis (Figure 5C-G, $P < 0.05$, Supplementary
Data File 2). As a phylum, they were negatively associated with *Chl* a, temperature and total
P. This suggests that, while the relative abundance of Actinobacteria was different between
different sites, within this phylum OTUs behaved similarly, i.e. they are found at similar
ratios at each location (Figure 2B). Proteobacteria showed a wider distribution range that was
close to significance in our test (adjusted p-value = 0.066, Figure 5D). These results are in
agreement with our observation that Actinobacteria and Proteobacteria OTUs dominate the
core microbiome of all sites (Supplementary Figure 6A) and that within each of these phyla, a

relatively large fraction of OTUs was core or cosmopolitan (Supplementary Figure 6B). In contrast, Verrucomicrobia, Cyanobacteria and Bacteroidetes did not cluster together in the CCA plot (although, as phyla, they were associated with some water parameters (Figure 5E-G, Supplementary Data File 2).

At a higher phylogenetic resolution, several lineages clustered in a non-random way and affected by the tested parameters (Supplementary Data File 2). In some cases, different lineages within the same phylum behaved similarly. For example, Acidimicrobiales and Actinomycetales orders, both belonging to the Actinobacteria, behaved similarly, being negatively associated with *Chl* a and total P concentrations, as well as with temperature. However, in other cases different lineages clearly clustered separately and were affected differently by the tested environmental factors, suggesting potential niche differentiation. Within the Cyanobacteria, the order Synechococcales was clearly separated in the CCA ordination from the Pseudoanabaenales and Nostocales (Figure 5H). The former was clustered non-randomly and was negatively associated with *Chl* a, NO_2+NO_3 and TP (Supplementary Data File 2), whereas the two latter orders were positively associated with these factors. Nostocales and Pseudanabaenales, in turn, differed in their correlation with temperature (positive and negative association with temperature, respectively). Notably, while Cyanobacteria appeared at all sampling sites (Figure 2B), few Cyanobacteria OTUs were part of the core microbiome (Supplementary Figure 6A), yet this phylum had the highest fraction of OTUs in the ''common'' category among all analyzed phyla (Supplementary Figure 6B). This further suggests that different Cyanobacteria of different water bodies, potentially correlated with water quality and trophic status. Differences in the ordination of groups at the class level were also observed within the phylum Verrucomicrobia, with Opitutae clustered non-randomly and associated with low NO_2+NO_3 and *Chl* a, clearly separating from other classes such as Methylacidiphilae and Spartobacteria, both of which were positively associated with *Chl* a (although not clustered, Figure 6B).

4. Discussion

Quality and chemical properties of freshwater ecosystems, aquatic microbial populations, and human health are tightly interlinked. In this study, we analyzed the microbiomes of a network of partly-interconnected lakes, which belong to several different drainage basins. We aimed to identify to what extent land-use in the drainage basins of the lakes affects both environmental conditions in the water and lake microbiomes. Several previous studies have identified correlations of land-use to various parameters of water chemistry, as well as to the abundance of specific members of the microbial community (Savage*, et al.*, 2010, Katsiapi*, et al.*, 2012, Beaver*, et al.*, 2014, Davis*, et al.*, 2015). However, few studies have attempted to determine whether land use controls the microbiome structure as a whole (Glöckner*, et al.*, 2000, Van Rossum*, et al.*, 2015). Importantly, in our analyses, we used tools that are relatively accessible to water quality managers: land-use maps are available for many regions of the world, and are often used for land management using GIS (Joerin*, et al.*, 2001, Zhang*, et al.*, 2015). The water quality measurements performed using hand-held instruments, and measurements of inorganic nitrogen compounds and total phosphorus were performed using commonly-applicable and standardized methods (Grasshoff & Kremling, 1983, Grasshoff & Ehrhardt, 1999, Louca*, et al.*, 2016). Our results suggest a conceptual model (schematically presented in Figure 6) whereby NO_2+NO_3 is related to a specific land use within the drainage basin of each water body (urban areas). In turn, NO_2+NO_3, together with total phosphorus *chl* a and temperature, are related to the relative abundance of specific microbial linages at different phylogenetic resolutions.

Previous studies have shown that specific environmental parameters affecting microbial community structure may vary between studies, freshwater systems and the time of sampling (e.g., Crump, *et al.*, 2007). Temperature is often identified as an explanatory variable (Van der Gucht, *et al.*, 2007), but this is not always the case (Savage, *et al.*, 2010). *Chlorophyll* a and total P concentrations are also sometimes identified as important variables (Yannarell & Triplett, 2005), but not always (Crump, *et al.*, 2007, Van der Gucht, *et al.*, 2007, Niño-García, *et al.*, 2016), and these two parameters may have a stronger effect on particle-associated rather than free-living bacteria (Savio, *et al.*, 2015). In our dataset, *chl* a and total P concentrations (which are often used as quantitative measures of lake trophic state) were somewhat related to the relative abundance of Cyanobacteria (Supplementary Figure 2), especially Nostocales and Pseudanabaenales (Figure 5). While pH is often identified as important (Yannarell & Triplett, 2005, Crump, *et al.*, 2007, Savio, *et al.*, 2015, Niño-García, *et al.*, 2016), it was not a statistically significant explanatory variable in our analysis. Finally, consistent with our results, NO_2+NO_3 have been shown to affect community structure in some cases (Kozak, *et al.*, 2019), whereas in other studies total N (which was not measured here) was shown to be important (e.g., Van der Gucht, *et al.*, 2007).

Below we first discuss specific links between land use types, water quality, and microbiome composition, as well as the association between specific lineages and water quality parameters. We then mention the caveats of our study and suggest ways to improve future studies.

4.1 What determines the microbiome - land use, mass transfer or biogeography?

Our initial hypothesis was that two types of factors would structure the microbial communities: i) environmental conditions within the water body, which are partly determined by land use within the drainage basin of each water body (often termed "species sorting", e.g., (Van der Gucht, *et al.*, 2007, Logue & Lindström, 2010), and ii) mass effects due to bacterial transport along the streams connecting the various sampled lakes. The relative importance of these two processes differs between different freshwater systems, with mass effects generally thought to be more prevalent in streams or smaller lakes with low water retention time and species sorting more prevalent in larger lakes with higher water retention times (Mašín, *et al.*, 2003, Lindström & Bergström, 2004, Crump, *et al.*, 2007, Logue & Lindström, 2010, Niño-García, *et al.*, 2016). However, not all lake systems are alike, example distance-decay relationships (which suggest mass effects) were observed in some lake systems (e.g. (Van der Gucht, *et al.*, 2007, Nelson, *et al.*, 2009), but not in others (e.g. Niño-García, *et al.*, 2016). Our results show no separation of bacterial communities by basin and no distance-decay patterns (Figure 2C, S2A). We note, however, that we did not measure the actual flux of water between the lakes, which could strongly affect the rate of emigration between the lakes (see "caveats", below). Given the currently available data, it seems likely that, at the geographic scale and time-points we studied, there is no clear limitation on the dispersal of bacteria between the lakes. Rather, environmental conditions within each water body strongly affected the microbiome structure, and these were partly determined by land-use in the drainage area of each water body (Figure 4).

The effect of water quality on the microbiome was quantitative rather than qualitative, resulting in differences in the relative abundances of different microbial groups or OTUs rather than driving some of them locally extinct (Supplementary Figure 6). Indeed ~50% of the OTUs were found in >90% of the sampling locations, suggesting a "core" microbiome that is found at all sites, albeit with different relative abundancesn (Savio, *et al.*, 2015). In organism-associated microbiomes, it has been suggested that microorganisms that are commonly found across locations or individuals (thus forming the core microbiome) may be

critical to the function of the microbial community, and indeed may define a "healthy"
community (Shade & Handelsman, 2011, Hernandez-Agreda, *et al.*, 2017). Such studies are
still rare for aquatic microbiomes, yet it is possible that better understanding of the dynamics
of the core microbiome may help understand how microbial communities are related to – and
affect – water quality.
Several studies have investigated the association between land use types and water properties
(Quinn, *et al.*, 1997, Larned, *et al.*, 2004, Mehaffey, *et al.*, 2005, Schoonover & Lockaby,
2006, Reimann, *et al.*, 2009). These studies generally suggest higher dissolved nitrogen and
phosphorus concentrations in waters surrounded by high coverage of urban areas and
agriculture lands (Wang & Yin, 1997, Ghai, *et al.*, 2014). In our study, urban land was an
important predictor for (and positively correlated with) concentrations of NO_2+NO_3 (Figure
3B-C, Supplementary Tables 1, 2). The correlation between urban area (which includes also
industry) and NO_2+NO_3 concentrations is in agreement with the previously mentioned
studies, and may be driven by discharge of waste and rain run-off from sealed surfaces, both
containing high nitrogen levels. Unlike previous studies, which observed a negative
correlation between forested area and total phytoplankton as well as Cyanobacteria biomass
(Catherine, *et al.*, 2008), we did not observe such an effect, despite having sampled in lakes
encompassing a wide array of trophic states. Neither did we observe an expected effect of
agricultural land, e.g. through the effect of nutrient runoff (Carpenter, *et al.*, 1998, Xu, *et al.*,
2010), potentially because this region is characterized by low P fertilizer application and low
livestock densities (Theobald, *et al.*, 2016). Thus, in the region we studied, no clear
correlation was observed between land use and total P or *chl* a concentrations (the low
number of oligotrophic-mesotrophic lakes precluded an analysis of the trophic levels
themselves). We note, however, that the interaction between land-use in lake catchments and
the physics, chemistry and biology of the lakes is complex, with potentially conflicting
drivers. For example, nutrients and dissolved organic matter exported from, e.g., forests into
lakes may, on the one hand, promote phytoplankton and bacterioplankton growth, while at
the same time increasing water color and subsequent light attenuation could result in changes
in lake stratification (reviewed by Solomon, 2015).
While we were able to explain, to some extent, water parameters via land-use types, and the
microbiome composition via water chemistry, we were unable to identify direct links
between land use and the specific microbiome composition. An interpretation of these results
might be that while land use in the drainage basin of each lake affects its microbiome
indirectly, e.g., by impacting water chemistry, there is a relatively low direct effect of land
use on the microbiome, e.g., by being a source for specific (presumably non-aquatic)
microbes. In support of this interpretation, relatively few OTUs belonged to Acidobacteria
(Supplementary Figure 6A) regarded as a representative phylum of soil bacteria (Janssen,
2006). Among these, only 0.04% of the total reads belonged to the *Solibacteres*, which have
been considered as a marker for soil bacteria (Battistuzzi & Hedges, 2009). In a study of
water bodies in the Ile-de-France region (Catherine, *et al.*, 2008), land use was able to partly
explain Cyanobacteria biomass, and in a subsequent study land use could predict to some
extent the phytoplankton structure through the effect of land-use on water quality (Catherine,
et al., 2016). Similarly, a land-use based model was able to partly predict the presence of
genes involved in the biosynthesis of the Cyanobacteria toxin microcystin (Marmen, *et al.*,
2016). In our dataset, even though the total fraction of Cyanobacteria is correlated with *Chl a*
($R^2 = 0.3$ Supplementary Figure 3C), and the main groups of Cyanobacteria partition clearly
into different niches based on the water chemistry (Figure 6A), we were unable to identify
any land-use based model that could explain the variation in the relative abundance of these
specific Cyanobacteria groups. Thus, while the potential of land-use to predict the abundance
of Cyanobacteria and their toxins has been demonstrated, further work is needed to develop

the appropriate methodologies and statistical tools that will identify linkages between land
use types and microbiome structure. Finally, it is possible that land use might affect the
function of the microbial population (at the genetic level) but that such an effect would not be
clearly seen in the population structure, as measured by 16S amplicon sequencing (e.g.,
Louca, 2016).

Additionally, increasing the number of sampled locations will help to better make use of
higher-resolution land-use maps, which may increase the ability to identify specific links
between land-use and microbiome structure. Given the good accessibility of land-use data,
which for much of the globe (including third world countries) can be derived from remote
sensing, developing these methods should be of high priority for future research (Ndlela, *et*
al., 2016).

4.2 Niche differentiation of freshwater bacterioplankton at the class and order levels
An important question in microbial ecology is whether there is a phylogenetic resolution at
which environmental conditions can be best related to microbiome structure and function.
Many studies focus on the phylum level (Lindström, *et al.*, 2005, Briée, *et al.*, 2007, Wolz, *et*
al., 2018), since, at a broad scale, bacterial phyla differ in some basic functional traits
including their ability to photosynthesize, or to utilize different carbon sources (Mitsui, *et al.*,
1986, Camp, *et al.*, 2009, Newton, *et al.*, 2011). Indeed, a clear differentiation had been
observed, in our dataset, between two groups of sites, those with a high relative abundance of
Cyanobacteria, and those with a high relative abundance of Actinobacteria and Proteobacteria
(Figure 2B). However, it has been suggested that this resolution is far too coarse to reliably
infer community function, due to the high genomic and functional variation within phyla and
even within individual bacterial lineages (Newton, *et al.*, 2011, Singer, *et al.*, 2016). Our
results suggest that some bacterial lineages clearly partition between different environments
already at low phylogenetic resolution, namely that of bacterial classes and orders. Several of
these patterns were in agreement with our previous expectations based on literature data. For
example, in freshwater environments, it is generally accepted that the relative contribution of
Pico-Cyanobacteria (e.g., Synechococcales) to total phytoplankton decreases with higher
trophic status. However, temporal patterns can be also important (Fukushima, *et al.*, 2017).
This may be attributed to the ability of Pico-Cyanobacteria to acquire major nutrients and
trace metals at sub-micromolar concentrations, a capability that enables them to occupy
relatively oligotrophic niches (Callieri, 2017). Larger phytoplankton (e.g., Nostocales) fills in
niches with higher nutrient concentrations, and hence is physiologically adapted to proliferate
at rich nutrient waters like fishponds and hypertrophic reservoirs (Cires & Ballot, 2016). The
presence and density of Nostocales are of great interest from a water quality perspective due
to their ability to produce a variety of secondary toxic metabolites (Sukenik, *et al.*, 2012). In
our samples, a clear separation could be observed in the CCA analyses between
Synechococcales, associated with conditions of low *Chl a* and TP, and Nostocales, associated
with the opposite conditions (Figure 5). Pseudanabaenales, which are filamentous
Cyanobacteria whose cells are smaller than those of the Nostocales, were associated also with
high NO_3+NO_3 concentrations. The physiology and ecology of Pseudanabaenales are
relatively poorly known (Acinas, *et al.*, 2008), yet these organisms are quite dominant in
many of the lakes we sampled (Supplementary Figure 6), providing an incentive for further
studies. The niche differentiation between Synechoccocales, Nostocales and
Pseudanabaenales is the process that underlies the relative scarcity of ubiquitous

Cyanobacteria OTUs, which instead dominate the "cosmopolitan" group (Supplementary
Figure 6), and further highlight the potential importance of understanding the dynamics of
"core" microbes in relation to water quality.
Potential niche differentiation was also observed within the phylum Verrocumicrobia, which
is ubiquitous but relatively less studied in freshwater environments (Figure. 5I, Newton, *et*
al., 2011). Some members of this phylum were found to be associated with high nutrient
concentrations or algal blooms (Eiler & Bertilsson, 2004, Kolmonen, *et al.*, 2004, Haukka, *et*
al., 2006). In our study Methylacidiphilae (and possibly also Spartobacteria) clearly separated
from the other three classes (Pedosphaerae, Opitutae and Verrucomicrobias), and may be
associated with high $NO_2 + NO_3$ concentrations and low temperatures. At least some
members of the Methylacidiphilae are methanotrophs (Camp, *et al.*, 2009), and one bacterium
from this class (*Methylacdiphilum fumariolicum*) is a diazotroph, able to actively fix
atmospheric N_2 (Khadem, *et al.*, 2010). While most methanotrophic Verrucomicrobia have
been isolated in extreme hot and acidic environments, the association of potential diazotrophs
(Methylacidiphilae as well as Pseudanabaenales) with high NO_2+NO_3 in lakes is intriguing,
and their potential contribution to high inorganic N concentrations is worth to perform
additional studies.
The separation between different orders in the CCA plots suggests that, in freshwater
bacteria, niche differentiation may be discernable at intermediate phylogenetic resolution -
higher than that of phyla but lower than that of specific species or strains. We also searched
for specific OTUs that were associated with aquatic environmental conditions or land use
categories, but found very few results, and when correlations were observed, they were
relatively weak (maximum Pearson R^2 was 0.53, Supplementary Table 3). Two OTUs, one
belonging to Phycisphaerales (phylum Planctomycetes) and one belonging to Saprospirales
(phylum Bacteroidetes), were positively associated with NO_2+NO_3 concentrations.
Phycisphaerales have been found in ANNAMOX processes (Hu, *et al.*, 2015), and
Saprospirales are associated with specific soil parameters (King, *et al.*, 2010, Hargreaves, *et*
al., 2015), yet, the role of these taxa in freshwater is mostly unclear. Thus, in our dataset, no
specific OTU can serve as an indicator species for specific aquatic environmental parameters
or land uses.
As the amount of genetic data (e.g. 16S rDNA surveys and metagenomes) from freshwater
locations rapidly increases, it will be important to explore the best phylogenetic resolution at
which to identify correlations between water quality and microbiome structure. Such
microbiome data can then be linked to the relative abundance of specific microbial groups
such as those involved in nutrient cycling, toxicity or pathogenesis, which in turn are
important traits for water quality management.
557
4.3 Caveats and Limitations

While we could explain a part of the variability in the microbiome (a maximum of 13.7% in
the VPA analysis, Figure. 4, similar to other studies, (Lear, *et al.*, 2013), a large part of the
variability remains unexplained. This variability may be due to other aquatic environmental
parameters we did not measure, for example the quantity and composition of the dissolved
organic carbon, which feeds heterotrophic bacteria and clearly has an effect on aquatic
microbiomes (Davis, *et al.*, 2015, Logue, *et al.*, 2015, Wei, *et al.*, 2016). Nor did we measure
hydrological properties of the systems such as the lake depth, turbulence, stratification or
retention times of the lakes, which can also strongly affect microbial population structure
(Kopylov, *et al.*, 2002, Nogueira, *et al.*, 2002, Humayoun, *et al.*, 2003, Carini, *et al.*, 2005,
Lindström, *et al.*, 2005, Percent, *et al.*, 2008). We also did not include in our analysis the
potential contribution of groundwater, even though this might be significant. Lake retention

times and potential groundwater inflow, for example, may affect the rate of population
change and the relative impact of neutral processes such as emigration compared to effects of
environmental drivers. While clearly of importance, often many of these hydrological
parameters remain unknown, including for the majority of the lakes we studied, some which
have been studied intensively for over 50 years (Allgaier & Grossart, 2006, Casper, 2012).
More broadly, we include in our analysis only "bottom-up" controls, whereas biotic
interactions such as grazing, phage infection and allelopathy can strongly impact population
structure (Louca, *et al.*, 2016).

Finally, our sampling approach, based on filtration on GF/F filters, assessed only part of the
community, likely under-estimating the relative abundance of ultramicrobacteria (Proctor, *et*
al., 2018). Many of these organisms belong to phyla that have yet to be cultured, and
understanding their contribution to lake microbial dynamics represents a major future
challenge. We also measured each location only once, relatively early during summer. Such a
snap-shot does not represent the dynamic nature of the aquatic microbiome. For example, in
several lakes that we studied, large Cyanobacteria blooms have been recorded, particularly
during August (Mantzouki, *et al.*, 2018). These blooms are often dominated by *Microcystis*
species, yet these were relatively rare in our samples. Clearly, extended time-series are
needed to gain a better understanding of the processes affecting aquatic microbiomes in
relation to land use.

Acknowledgements

We thank Elke Mach and Uta Mallok for chemical analyses, Liat Linde and Doron Fogel for
the DNA extraction and the MIBI group for valuable discussions

This manuscript has been released as a Pre-Print at (Marmen *et al.,* 2018).

Conflict of Interest Statement
The authors declare no conflict of interest
596

Funding source
This study was supported by a German-Israeli Cooperation in Water Technology Research
Young Scientists Exchange Program (to SM), by grant number 3-10342 from the Israeli
Ministry of Science and Technology (to DS) and by DFG project Microprime (GR1540/28-1,
to HPG). Also, we thank the Leibnitz Institute and the University of Haifa for additional
funding.

Author contributions:
SM, HPG and DS designed study, SM, LG, HPG and DS collected samples, LB, LG, HPG
and DS analyzed the geographic data, SM, LB, AA, AM, ML and DS analyzed the genetic
data, SM and DS wrote the manuscript with contributions from all co-authors.
608

All authors have approved the final manuscript for publication

References

Abdelzaher AM, Wright ME, Ortega C, *et al.* (2010) Presence of Pathogens and Indicator
Microbes at a Non-Point Source Subtropical Recreational Marine Beach. *Applied and*
Environmental Microbiology **76**: 724-732.
Acinas SG, Haverkamp THA, Huisman J & Stal LJ (2008) Phenotypic and genetic
diversification of Pseudanabaena spp. (cyanobacteria). *The Isme Journal* **3**: 31.
Allgaier M & Grossart H-P (2006) Diversity and seasonal dynamics of Actinobacteria
populations in four lakes in northeastern Germany. *Applied and Environmental Microbiology*
72: 3489-3497.
Battistuzzi FU & Hedges SB (2009) A Major Clade of Prokaryotes with Ancient Adaptations
to Life on Land. *Molecular Biology and Evolution* **26**: 335-343.
Beaver JR, Manis EE, Loftin KA, Graham JL, Pollard AI & Mitchell RM (2014) Land use
patterns, ecoregion, and microcystin relationships in U.S. lakes and reservoirs: A preliminary
evaluation. *Harmful Algae* **36**: 57-62.
Bell T (2010) Experimental tests of the bacterial distance–decay relationship. *The Isme*
Journal **4**: 1357.
Blank L & Blaustein L (2014) A multi-scale analysis of breeding site characteristics of the
endangered fire salamander (Salamandra infraimmaculata) at its extreme southern range
limit. *Hydrobiologia* **726**: 229-244.
Briée C, Moreira D & López-García P (200 (7Archaeal and bacterial community composition
of sediment and plankton from a suboxic freshwater pond. *Research in Microbiology* **158**:
213-227.
Burnham KP & Anderson DR (2003) *Model selection and multimodel inference: a practical*
information-theoretic approach. Springer Science & Business Media.
Cabral JPS (2010) Water Microbiology. Bacterial Pathogens and Water. *International*
Journal of Environmental Research and Public Health **7**: 3657.
Calcagno V & de Mazancourt C (2010) glmulti: an R package for easy automated model
selection with (generalized) linear models. *Journal of statistical software* **34**: 1-29.
Callieri C (2017) Synechococcus plasticity under environmental changes. *FEMS*
microbiology letters **364**: fnx229.
Camp HJMOd, Islam T, Stott MB, *et al* (2009) .Environmental, genomic and taxonomic
perspectives on methanotrophic Verrucomicrobia. *Environmental Microbiology Reports* **1**:
293-306.
Caporaso JG, Kuczynski J, Stombaugh J, *et al.* (2010) QIIME allows analysis of high-
throughput community sequencing data. *Nature Methods* **7**: 335.
Carini S, Bano N, LeCleir G & Joye SB (2005) Aerobic methane oxidation and methanotroph
community composition during seasonal stratification in Mono Lake, California (USA).
Environmental Microbiology **7**: 1127-1138.
Carpenter SR, Caraco NF, Correll DL, Howarth RW, Sharpley AN & Smith VH (1998)
NONPOINT POLLUTION OF SURFACE WATERS WITH PHOSPHORUS AND
NITROGEN. *Ecological Applications* **8**: 559-568.
Casper SJ (2012) *Lake Stechlin: A temperate oligotrophic lake*. Springer Science & Business
Media.
Catherine A, Troussellier M & Bernard C (2008) Design and application of a stratified
sampling strategy to study the regional distribution of cyanobacteria (Ile-de-France, France).
Water Research **42**: 4989-5001.
Catherine A, Selma M, Mouillot D, Troussellier M & Bernard C (2016) Patterns and multi-
scale drivers of phytoplankton species richness in temperate peri-urban lakes. *Science of the*
Total Environment **559**: 74-83.

Chen W, Wilkes G, KHAN IU-H, *et al.* (2018) Aquatic bacterial community associated with land use and environmental factors in agricultural landscapes using a metabarcoding approach. *Frontiers in Microbiology* **9**: 2301.

Cires S & Ballot A (2016) A review of the phylogeny, ecology and toxin production of bloom-forming Aphanizomenon spp. and related species within the Nostocales (cyanobacteria). *Harmful Algae* **54**: 21-43.

Crump BC, Amaral-Zettler LA & Kling GW (2012) Microbial diversity in arctic freshwaters is structured by inoculation of microbes from soils. *The Isme Journal* **6**: 1629.

Crump BC, Hopkinson CS, Sogin ML & Hobbie JE (2004) Microbial Biogeography along an Estuarine Salinity Gradient: Combined Influences of Bacterial Growth and Residence Time. *Applied and Environmental Microbiology* **70**: 1494-1505.

Crump BC, Adams HE, Hobbie JE & Kling GW (2007) BIOGEOGRAPHY OF BACTERIOPLANKTON IN LAKES AND STREAMS OF AN ARCTIC TUNDRA CATCHMENT. *Ecology* **88**: 1365-1378.

Davis TW, Bullerjahn GS, Tuttle T, McKay RM & Watson SB (2015) Effects of Increasing Nitrogen and Phosphorus Concentrations on Phytoplankton Community Growth and Toxicity During Planktothrix Blooms in Sandusky Bay, Lake Erie. *Environmental Science & Technology* **49**: 7197-7207.

Eiler A & Bertilsson S (2004) Composition of freshwater bacterial communities associated with cyanobacterial blooms in four Swedish lakes. *Environmental Microbiology* **6**: 1228-1243.

Fukushima M, Tomioka N, Jutagate T, *et al.* (2017) The dynamics of pico-sized and bloom-forming cyanobacteria in large water bodies in the Mekong River Basin. *PLOS ONE* **12**: e0189609.

Ghai R ,Mizuno CM, Picazo A, Camacho A & Rodriguez-Valera F (2014) Key roles for freshwater Actinobacteria revealed by deep metagenomic sequencing. *Molecular Ecology* **23**: 6073-6090.

Glöckner FO, Zaichikov E, Belkova N, Denissova L, Pernthaler J, Pernthaler A & Amann R (2000) Comparative 16S rRNA analysis of lake bacterioplankton reveals globally distributed phylogenetic clusters including an abundant group of actinobacteria. *Appl. Environ. Microbiol.* **66**: 5053-5065.

Grasshoff K & Ehrhardt M (1999) Methods of Seawater Analysis, Third, Completely Revised and Extenden Edition. *WILEY-VCH Verlag GmbH, D* **69469**: 203-223.

Grasshoff K, Kremling K & Ehrhardt M (2009) *Methods of seawater analysis*. John Wiley & Sons.

Grasshoff KE & Kremling M (1983) Methods of seawater analysis. Verlag Chemie. *Weinheim, Germany* **4**: 9.

Griffin JS, Lu N, Sangwan N, *et al.* (2017) Microbial diversity in an intensively managed landscape is structured by landscape connectivity. *FEMS microbiology ecology* **93**: fix120.

Hargreaves SK, Williams RJ & Hofmockel KS (2015) Environmental Filtering of Microbial Communities in Agricultural Soil Shifts with Crop Growth. *PLOS ONE* **10**: e0134345.

Haukka K, Kolmonen E, Hyder R, *et al.* (2006) Effect of Nutrient Loading on Bacterioplankton Community Composition in Lake Mesocosms. *Microbial Ecology* **51**: 137-146.

Hernandez-Agreda A, Gates RD & Ainsworth TD (2017) Defining the core microbiome in corals' microbial soup. *Trends in Microbiology* **25**: 125-140.

Horner-Devine MC, Lage M, Hughes JB & Bohannan BJM (2004) A taxa–area relationship for bacteria. *Nature* **432**: 750.

Hu S, Zeng RJ, Haroon MF, Keller J, Lant PA, Tyson GW & Yuan Z (2015) A laboratory investigation of interactions between denitrifying anaerobic methane oxidation (DAMO) and anammox processes in anoxic environments. *Scientific Reports* **5**: 8706.

Humayoun SB, Bano N & Hollibaugh JT (2003) Depth Distribution of Microbial Diversity in Mono Lake, a Meromictic Soda Lake in California. *Applied and Environmental Microbiology* **69**: 1030-1042.

Janssen PH (2006) Identifying the Dominant Soil Bacterial Taxa in Libraries of 16S rRNA and 16S rRNA Genes. *Applied and Environmental Microbiology* **72**: 1719-1728.

Joerin F, Thériault M & Musy A (2001) Using GIS and outranking multicriteria analysis for land-use suitability assessment. *International Journal of Geographical Information Science* **15**: 153-174.

Jones SE & McMahon KD (2009) Species-sorting may explain an apparent minimal effect of immigration on freshwater bacterial community dynamics. *Environmental Microbiology* **11**: 905-913.

Katsiapi M, Mazaris AD, Charalampous E & Moustaka-Gouni M (2012) Watershed land use types as drivers of freshwater phytoplankton structure. *Hydrobiologia* **698**: 121-131.

Khadem AF, Pol A, Jetten MSM & Op den Camp HJM (2010) Nitrogen fixation by the verrucomicrobial methanotroph 'Methylacidiphilum fumariolicum' SolV. *Microbiology* **156**: 1052-1059.

King AJ, Freeman KR, McCormick KF, Lynch RC, Lozupone C, Knight R & Schmidt SK (2010) Biogeography and habitat modelling of high-alpine bacteria. *Nature Communications* 1: 53.

Kolmonen E, Sivonen K, Rapala J & Haukka K (2004) Diversity of cyanobacteria and heterotrophic bacteria in cyanobacterial blooms in Lake Joutikas, Finland. *Aquatic Microbial Ecology* **36**: 201-211.

Kopylov AI, Kosolapov DB, Romanenko AV & Degermendzhy AG (2002) Structure of planktonic microbial food web in a brackish stratified Siberian lake. *Aquatic Ecology* **36**: 179-204.

Kozak A, Celewicz-Gołdyn S & Kuczyńska-Kippen N (2019) Cyanobacteria in small water bodies: The effect of habitat and catchment area conditions. *Science of the Total Environment* **646**: 1578-1587.

Langenheder S & Székely AJ (2011) Species sorting and neutral processes are both important during the initial assembly of bacterial communities. *The Isme Journal* **5**: 1086-1094.

Larned ST, Scarsbrook MR, Snelder TH, Norton NJ & Biggs BJF (2004) Water quality in low-elevation streams and rivers of New Zealand: Recent state and trends in contrasting land-cover classes. *New Zealand Journal of Marine and Freshwater Research* **38**: 347-366.

Lear G, Washington V, Neale M, Case B, Buckley H & Lewis G (2013) The biogeography of stream bacteria. *Global Ecology and Biogeography* **22**: 544-554.

Leibold MA & Norberg J (2004) Biodiversity in metacommunities: Plankton as complex adaptive systems? *Limnology and Oceanography* **49**: 1278-1289.

Lindström ES & Bergström A-K (2004) Influence of inlet bacteria on bacterioplankton assemblage composition in lakes of different hydraulic retention time. *Limnology and Oceanography* **49**: 125-136.

Lindström ES & Langenheder S (2012) Local and regional factors influencing bacterial community assembly. *Environmental Microbiology Reports* **4**: 1-9.

Lindström ES, Kamst-Van Agterveld MP & Zwart G (2005) Distribution of Typical Freshwater Bacterial Groups Is Associated with pH, Temperature, and Lake Water Retention Time. *Applied and Environmental Microbiology* **71**: 8201-8206.

Logares R, Tesson SV, Canbäck B, Pontarp M, Hedlund K & Rengefors K (2018) Contrasting prevalence of selection and drift in the community structuring of bacteria and microbial eukaryotes. *Environmental Microbiology* **20**: 2231-2240.

Logue JB & Lindström ES (2008) Biogeography of bacterioplankton in inland waters. *Freshwater Reviews* **1**: 99-115.

Logue JB & Lindström ES (2010) Species sorting affects bacterioplankton community composition as determined by 16S rDNA and 16S rRNA fingerprints. *The Isme Journal* **4**: 729.

Logue JB, Langenheder S, Andersson AF, Bertilsson S, Drakare S, Lanzén A & Lindström ES (2012) Freshwater bacterioplankton richness in oligotrophic lakes depends on nutrient availability rather than on species–area relationships. *The Isme Journal* **6**: 1127.

Logue JB, Stedmon CA, Kellerman AM*, et al.* (2015) Experimental insights into the importance of aquatic bacterial community composition to the degradation of dissolved organic matter. *The Isme Journal* **10**: 533.

Louca S, Parfrey LW & Doebeli M (2016) Decoupling function and taxonomy in the global ocean microbiome. *Science* **353**: 1272-1277.

Mantzouki E, Lürling M, Fastner J*, et al.* (2018) Temperature effects explain continental scale distribution of cyanobacterial toxins. *Toxins* **10**: 156.

Marcelino VR, van Oppen MJH & Verbruggen H (2017) Highly structured prokaryote communities exist within the skeleton of coral colonies. *The Isme Journal* **12**: 300.

Marmen S, Aharonovich D, Grossowicz M, Blank L, Yacobi YZ & Sher DJ (2016) Distribution and Habitat Specificity of Potentially-Toxic Microcystis across Climate, Land, and Water Use Gradients. *Frontiers in Microbiology* **7**.

Marmen S, Blank L, Al-Ashhab A, et al. (2018) The role of land use types and water chemical properties in structuring the microbiome of a connected lakes system. BioRxiv 401299

Mašín M, Jezbera J, Nedoma J, Straškrabová V, Hejzlar J & Šimek K (2003) Changes in bacterial community composition and microbial activities along the longitudinal axis of two canyon-shaped reservoirs with different inflow loading. *Hydrobiologia* **504**: 99-113.

Mehaffey MH, Nash MS, Wade TG, Ebert DW, Jones KB & Rager A (2005) Linking Land Cover and Water Quality in New York City'S Water Supply Watersheds. *Environmental Monitoring and Assessment* **107**: 29-44.

Mitsui A, Kumazawa S, Takahashi A, Ikemoto H, Cao S & Arai T (1986) Strategy by which nitrogen-fixing unicellular cyanobacteria grow photoautotrophically. *Nature* **323**: 720.

Ndlela LL, Oberholster PJ, Van Wyk JH & Cheng PH (2016) An overview of cyanobacterial bloom occurrences and research in Africa over the last decade. *Harmful Algae* **60**: 11-26.

Nekola JC & White PS (1999) The distance decay of similarity in biogeography and ecology. *Journal of Biogeography* **26**: 867-878.

Nelson CE, Sadro S & Melack JM (2009) Contrasting the influences of stream inputs and landscape position on bacterioplankton community structure and dissolved organic matter composition in high-elevation lake chains. *Limnology and Oceanography* **54**: 1292-1305.

Newton RJ, Jones SE, Eiler A, McMahon KD & Bertilsson S (2011) A Guide to the Natural History of Freshwater Lake Bacteria. *Microbiology and Molecular Biology Reviews* **75**: 14-49.

Niño-García JP, Ruiz-González C & del Giorgio PA (2016) Interactions between hydrology and water chemistry shape bacterioplankton biogeography across boreal freshwater networks. *The Isme Journal* **10**: 1755.

Nogueira R, Melo LsF, Purkhold U, Wuertz S & Wagner M (2002) Nitrifying and heterotrophic population dynamics in biofilm reactors: effects of hydraulic retention time and the presence of organic carbon. *Water Research* **36**: 469-481.

Percent SF, Frischer ME, Vescio PA, *et al* (2008) .Bacterial Community Structure of Acid-
Impacted Lakes: What Controls Diversity? *Applied and Environmental Microbiology* **74**:
1856-1868.
Proctor CR, Besmer MD, Langenegger T, *et al.* (2018) Phylogenetic clustering of small low
nucleic acid-content bacteria across diverse freshwater ecosystems. *The Isme Journal* **12**:
1344-1359.
Quinn JM, Cooper AB, Davies-Colley RJ, Rutherford JC & Williamson RB (1997) Land use
effects on habitat, water quality, periphyton, and benthic invertebrates in Waikato, New
Zealand ,hill-country streams. *New Zealand Journal of Marine and Freshwater Research* **31**:
579-597.
Reimann C, Finne TE, Nordgulen Ø, Sæther OM, Arnoldussen A & Banks D (2009) The
influence of geology and land-use on inorganic stream water quality in the Oslo region,
Norway. *Applied Geochemistry* **24**: 1862-1874.
Ritchie RJ (2008) Universal chlorophyll equations for estimating chlorophylls a, b, c, and d
and total chlorophylls in natural assemblages of photosynthetic organisms using acetone,
methanol, or ethanol solvents. *Photosynthetica* **46**: 115-126.
Savage C, Leavitt PR & Elmgren R (2010) Effects of land use, urbanization, and climate
variability on coastal eutrophication in the Baltic Sea. *Limnology and Oceanography* **55**:
1033-1046.
Savio D, Sinclair L, Ijaz UZ, *et al.* (2015) Bacterial diversity along a 2600 km river
continuum. *Environmental Microbiology* **17**: 4994-5007.
Schoonover JE & Lockaby BG (2006) Land cover impacts on stream nutrients and fecal
coliform in the lower Piedmont of West Georgia. *Journal of Hydrology* **331**: 371-382.
Shade A & Handelsman J (2011) Beyond the Venn diagram: the hunt for a core
microbiomeemi_2585. ed.^eds.), p.^pp.
Singer E, Bushnell B, Coleman-Derr D, *et al.* (2016) High-resolution phylogenetic microbial
community profiling. *The Isme Journal* **10**: 2020.
Solomon CT, Jones SE, Weidel BC, *et al.* (2015) Ecosystem consequences of changing
inputs of terrestrial dissolved organic matter to lakes: current knowledge and future
challenges. *Ecosystems* **18**: 376-389.
Sukenik A, Hadas O, Kaplan A & Quesada A (2012) Invasion of Nostocales (cyanobacteria)
to Subtropical and Temperate Freshwater Lakes – Physiological, Regional, and Global
Driving Forces. *Frontiers in Microbiology* **3**.
Theobald TF, Schipper M & Kern J (2016) Phosphorus flows in Berlin-Brandenburg, a
regional flow analysis. *Resources, Conservation and Recycling* **112**: 1-14.
Valério E, Chaves S & Tenreiro R (2010) Diversity and Impact of Prokaryotic Toxins on
Aquatic Environments: A Review. *Toxins* **2**: 2359.
Van der Gucht K, Cottenie K, Muylaert K, *et al.* (2007) The power of species sorting: local
factors drive bacterial community composition over a wide range of spatial scales.
Proceedings of the National Academy of Sciences **104**: 20404-20409.
Van Rossum T, Peabody MA, Uyaguari-Diaz MI, *et al* (2015) .Year-Long Metagenomic
Study of River Microbiomes Across Land Use and Water Quality. *Frontiers in Microbiology*
6.
Vitousek PM, Mooney HA, Lubchenco J & Melillo JM (1997) Human Domination of Earth's
Ecosystems. *Science* **277**: 494-499.
Wang X & Yin Z-Y (1997) Using GIS to assess the relationship between land use and water
quality at a watershed level. *Environment International* **23**: 103-114.
Wei Y, Wei Z, Cao Z, *et al.* (2016) A regulating method for the distribution of phosphorus
fractions based on environmental parameters related to the key phosphate-solubilizing
bacteria during composting. *Bioresource Technology* **211**: 610-617.

Williamson CE, Saros JE, Vincent WF & Smol JP (2009) Lakes and reservoirs as sentinels,
integrators, and regulators of climate change. *Limnology and Oceanography* **54**: 2273-2282.
Winter TC, Rosenberry DO & LaBaugh JW (2003) Where Does the Ground Water in Small
Watersheds Come From? *Groundwater* **41**: 989-1000.
Wolz CRM, Yarwood SA, Grant EHC, Fleischer RC, Lips KR & Hoye B (2018) Effects of
host species and environment on the skin microbiome of Plethodontid salamanders. *Journal*
of Animal Ecology **87**: 341-353.
Xu H, Paerl HW, Qin B, Zhu G & Gaoa G (2010) Nitrogen and phosphorus inputs control
phytoplankton growth in eutrophic Lake Taihu, China .*Limnology and Oceanography* **55**:
420-432.
Yannarell AC & Triplett EW (2005) Geographic and Environmental Sources of Variation in
Lake Bacterial Community Composition. *Applied and Environmental Microbiology* **71**: 227-
239.
Zeglin LH (2015) Stream microbial diversity in response to environmental changes: review
and synthesis of existing research. *Frontiers in Microbiology* **6**: 454.
Zhang J, Su Y, Wu J & Liang H (2015) GIS based land suitability assessment for tobacco
production using AHP and fuzzy set in Shandong province of China. *Computers and*
Electronics in Agriculture **114**: 202-211.

876
877 **Titles and legends to figures**

878
879

Figure 1. Land use and drainage basins in the sampling area. A total of 46 samples were
collected from lakes and a fish-pond in the Brandenburg region of northern Germany. Five
types of land use: urban areas, agriculture (further divided for GLM analysis to
heterogeneous agriculture, pasture lands and arable lands), forests, wetlands and water
bodies, were mapped using GIS, with black lines superimposed indicating drainage basins.
The arrows indicates the dominant flow direction within each basin. A map with the identities
of each sampling location is presented in Supplementary Figure 1, and the detailed
information on each sampling location can be found in Supplementary Tables 4-6.
890
891

Figure 2. An overview of the microbial populations in the sampled lakes. A) α-diversity of the microbial populations in the different lakes, measured by the Shannon index. The columns are colored by drainage basins. No clear patterns are observed. **B) Community composition and community clustering based on the Bray-Curtis similarity index**; Colors represent different phyla, lines within the same color represent OTUs at the 97% identity cutoff. The sampling sites can be divided into two groups: one dominated by Cyanobacteria, and the other by Actinobacteria, Proteobacteria and Bacteroidetes. **C) No distance decay patterns are observed**. Each point represents the Bray-Curtis dissimilarity between locations within the same basin (blue) or locations found in two separate basins (red). Spearman's r between geographic distance and Bray Curtis distance is 0.03, p=0.30. **D) Most of the OTUs are found in more than 50% of the sampling sites**. The diversity (γ diversity) of all sampling locations combined is presented, divided into five categories representing core OTUs (found in all sites), cosmopolitan OTUs (> 90% of the sites), common OTUs (> 50% of the sites), rare OTUs (< 10% of the sites), and unique OTUs (present at one site only).

Figure 3. Spesific land use typy affects aquatic environmental parameters: General Linear Models (GLMs) analysis were performed and resulted in **Urban areas are positively related to NO$_2$+NO$_3$ concentrations.** A strong correlation (R^2= 0.74) is observed between land uses and nitrite and nitrate concentrations, with the fraction of "urban areas" being the most important variable contributing to the correlation

919 **Figure 4. Water chemical properties represent the main explanatory variable for the**
920 **microbiome composition (ca. 11% among all sites).** Variation Partition Analysis performed
921 with 3 data sets (basin and local land use as well as water chemistry) to explain the variation
922 in the microbiome composition. The combination of land uses and water chemistry explain
923 13.7% of microbiome variability.

Figure 5. Differences between bacterial phyla in their association with different aquatic environmental conditions. Canonical Correspondence Analysis (CCA) plots are shown, with sampling locations and OTUs ordinated based on the four aquatic environmental parameters shown to significantly affect the microbiome composition. **A) An overview of the ordination, showing the 2845 most abundant OTUs.** The following panels each highlight a different aspect of this ordination. **B) The same ordination as in A, with the OTUs removed** to show the relative location of the eigenvectors for the aquatic environmental parameters and the ordination of the sampling sites. **C-G) The same plot as in A, with specific microbial phyla highlighted. H-I). Separation in the CCA ordination between selected lineages within the Cyanobacteria (h, orders) and Verrucomicrobia (I, classes).** Note the clear separation between the Synechococcoales and the Pseudanabaenales and Nostocales along the first CCA axis with some differentiation between the latter two orders primarily along the second CCA axis (h). Within the Verrucomicrobia, the Methylacidiphilae, which are limited to the upper-right hand quadrant, possibly in association with high NO_2+NO_3 levels, are clearly separated from the other orders, except of the Spartobacteria.

Figure 6. Conceptual model summarizing the observed effects of land use on water chemistry, and of water chemistry on selecetd microbial lineages. The first layer represents the effect of land use on aquatic environmental parameters (based on the GLMs), with the arrow showing the significant correlation between urban land and NO_2+NO_3. The seconds layer shows the effect of the four environmental parameters shown to sigificantly affect the microbiome structure (based on the CCA analysis) on selected microbial linages within the Cyanobacteria and Verrucomicrobia. Solid arrows show significant positive correlations (based on the permutation analysis), dotted blunt-ended lines show negative correlations. Lineages that are more clustered than randomly expected are in bold and italics.

Supplementary Material

1. Supplementary methods

1.1 Study region and delineation of drainage basins

The sampling area lies within the region of Brandenburg, in northern Germany (Figure 1,

Supplementary Figure 1). The area is relatively flat (18-89 meters above sea level) and

contains hundreds of lakes, ponds and depressions that were formed by glacial action during

the Pleistocene. Three main basins drain this region. From west to east, these are: 1) The

Rhin basin, which flows south and then south-east, joining the Havel basin near Oranienburg;

2) The Havel basin itself, which flows generally south from the area of Lake Müritz, joining

the Lychen sub-basin (which flows from the east) at Stolpsee (nearby Fürstenberg). It then

continues south through Oranienburg to Berlin. The Havel then turns west and drains into the

Elbe, itself draining into the North Sea, northwest of Hamburg; 3) the Ucker basin, flowing

north through Oberuckersee and Unteruckersee and finally draining into the Szczecin Lagoon

and the Baltic Sea. These three main basins can be subdivided into smaller basins.

The drainage basin of each lake was defined as the line connecting the highest points from

which the surface water flow enters each of the lakes. Since the lakes are inter-connected, the

drainage basin of each lake contains also those of the lakes upstream to it. Due to the flat

topography of the region automatic basin detection using GIS was not satisfactory, and thus

the basins depicted in Figure 1 and Supplementary Figure 1 were delineated manually using a

combination of topographic maps produced from the Digital Elevation Model layer with

a resolution of 0.0008333 decimal degree and high-resolution hiking and biking maps

(Kompass Fahrradkarte, 1:70,000). Human structures such as roads were not informative due

to the flat terrain. Literature surveys and site visits by the sampling team were used to clarify

connectivity and flow conditions at specific locations. Notably, due to the flat terrain and the

long history of water use for agriculture, aquaculture and transportation, the region includes

also many artificial canals, weirs and ponds, often connecting between different basins. In

order to simplify the analysis, the exchange between the basins was neglected. Additionally,

Lake Müritz primarily drains north-east into Lake Kölpinsee and from there east into the Elde

and ultimately into the Elbe. However, the lake also drains south at Rechlin into the Havel

basin. The border between the Havel and Elde basins was delineated, based on the

topography surrounding the lake, as shown in Figure 1.

Our 46 sampling sites varied between 7 different sub-basins as follow: 11 samples at Havel

basin, 10 at Lychen, 12 at Rhin, 5 at Stechlin, 4 at Ucker, 1 sample from Tollensesee, 1

sample from Berlin Spree and 2 samples from lakes that are not part from any drainage basin

(marked as "none" in Figure 1). Not all samples from the same basin are directly connected,

for instance, water from the lakes in the Rhin's northern part (12-15 in Supplementary Figure

1) meets water from the lakes in the Rhin's eastern part (18, 19, 21-23) only downstream,

near sites 16 and 17. Thus, in some cases sub-basins should be considered for the analyses

(see main text and Sup Supplementary Figure 5).

The land uses, obtained from the Corine Land Cover map

(CLC, http://www.eea.europa.eu/publications/COR0-landcover), were divided into five

categories: Artificial areas, Agriculture (further divided for GLM analysis to heterogeneous

agriculture, pasture lands and arable lands), Forests and natural areas, Wetlands and Water

bodies (Figure 1). Sample locations were selected in order to represent a wide diversity of

environmental parameters and land uses (Supplementary Data File 1, Supplementary Figure

S1).

1002

1003 **1.2 Estimation of the trophic status of the lakes**

The trophic classification of the lakes is based on measurements done or directed by the

environmental agencies of the federal states Mecklenburg-Vorpommern and Brandenburg,

following in general the rules of the water framework directive of the European Union. Such

assessments are available for 25 out of the 46 lakes in the following link

https://lfu.brandenburg.de/cms/detail.php/bb1.c.305410.de

Information for another 14 lakes was gathered from the following additional websites:

https://www-docs.b-tu.de/fg-gewaesserschutz/public/projekte/uba_2/02_meck_pom.pdf

https://www-docs.b-tu.de/fg-gewaesserschutz/public/projekte/uba_2/05_brandenburg.pdf

https://www.lung.mv-regierung.de/dateien/glrp_ms_06_2011.pdf

Finally, the trophic state of the rest of the lakes was assessed based on the local expertise of

the authors. Based on these assessments, there was one meso-oligotrophic lake, 8

mesotrophic lakes, 30 eutrophic lakes and 7 polytrophic lakes (including one fishpond,

Supplementary excel file).

1.3 DNA extraction, sequencing of 16S rRNA gene amplicons and sequence quality

control

DNA extraction was performed at the Biomedical Core Facility of the Technion, Israel, using

a combination of manual treatments and robotic extraction with the QiaCube robot. Briefly,

filters were thawed, centrifuged for 10 min at $15000 \times g$ and the preservation buffer was

removed. Lysis buffer from the DNeasy Blood & Tissue Kit (Qiagen) was added and the

samples were mechanically pounded with two 3mm sterile stainless steel beads at the speed

of 30 Hz for 1.5 min using a TissueLyser LT (Qiagen). 30 µl of lysozyme were added, the

tubes were incubated at 37°C for 30 min, followed by the addition of 25 mL proteinase K and

200□□l of the buffer AL with an additional incubation for 1 h at 56°C on a shaker. Finally,

the tubes were centrifuged for 10 min at 5000 x *g* and the upper liquid was transferred to a

new 2 mL Eppendorf tube for extraction by the QiaCube robot using the manufacturer's

instructions. The DNA quantity were assessed using Picogreen (Quant-iT PicoGreen dsDNA

reagent, Invitrogen), and the quality (apparent size and degradation products) was assessed

using a TapeStation (Agilent Technologies).

PCR reactions for amplifying the 16S rRNA gene fragments were performed in triplicate for

each DNA sample using the following primers targeting the V3-V4 region: forward

CS1_341F (5'-ACA CTG ACG ACA TGG TTC TAC ANN NNC CTA CGG GAG GCA

GCA G-3') and reverse CS2_806R (5'-TAC GGT AGC AGA GAC TTG GTC TGG ACT

ACH VGG GTW TCT AAT-3'). The PCR reactions were performed in a volume of 25 μl,

with 10 ng of DNA as PCR template. The PCR protocol was as follows: initial denaturation

stage at 95°C for 5 min, followed by 28 cycles at 95 °C for 30s, 50°C for 30s, and 72°C for

60 s, with a final extension step at 72°C for 5 min. The enzyme used was BIOLINE 2x

MyTaq Red Mix, and the PCR reaction was performed in a TProfessional Basic Gradient

thermocycler (Biometra). Following the first PCR, triplicate samples were combined, and the

products sent to the DNA Services Facility of the University of Illinois, Chicago, where a

second PCR was performed to incorporate barcodes and sequencing adapters (CS1 and CS2).

Sequencing was performed using a 2×250 base pair format on a MiSeq flow cell (V3

chemistry).

1047

1.4 16S analysis and statistical tests

Upon receiving the sequence data all sequences underwent quality control for Phix DNA

removal using bowtie2 (Langmead & Salzberg, 2012), unmerged reads using PEAR (Zhang,

et al., 2013) and incomplete, low quality scores and ambiguous bases sequences using

MOTHUR Software V.1.36.1 using a phred score of 30, removing homopolymers longer than

8 nucleotides long and allowing no ambiguous bases (Schloss, *et al.*, 2009). Following

quality filtering, sequences were classified into different Operational Taxonomic Units

(OTUs) based on 97% sequence similarity threshold using the "pick_de_novo_otus.py"

command in Qiime (Caporaso, *et al.*, 2010) and assigned taxonomy with the SILVA database

version 128. Upon classification, an OTU table was generated and OTUs with fewer than 5

reads across the entire table were removed. At the end of analysis, the range of sequence

number per sample was 35,775 – 75,704. This dataset, containing 45,240 OTUs, was used for

all diversity analyses. For all multivariate statistical analyses and for the graphs presented in

Figure 2B we used a smaller dataset which included only 3,611 OTUs with at least 50 reads

across all sites. OTU numbers were $\log_{10}$ transformed. All calculations were performed in R

software version 3.3.3 using the vegan package (Oksanen, *et al.*, 2007).

Alpha diversity (Shannon index) was calculated using the ''diversity'' function. The

similarity in bacterial community structure among samples was calculated using a Bray–

Curtis dissimilarity by ''vegdist'' function. Gamma diversity was calculated in Excel by

summing the number of sites where each OTU was present and dividing them into five

categories: 1) Core OTUs, which were present at all the locations; 2) Cosmopolitan OTUs,

which were present at more than 90% of the locations; 3) Common OTUS, present at more

than 50% of the sites; 4) Rare OTUs, present in less than 10% of the sites; and 5) Unique

OTUs, specific to only one site.

In order to explore whether samples from the same drainage basin have spatial similarity of

bacterial communities, a non-metric multidimensional scaling analysis (nMDS) was carried

out in R with ''metaMDS'' function and a Bray-Curtis matrix. Sites were colored by basins.

Before performing multivariate analyses, we removed environmental parameters which were

strongly correlated (Spearman $r > 0.8$, $P < 0.05$), with the exception of forests and arable

lands, as the removal of one of these dominant land-use types significantly reduced the total

fractional cover. Thus, the effect of these land-use types cannot be differentiated. The final

environmental parameters selected were pH, temperature, ammonium, $No_3 + NO_2$, total

phosphorus and *Chl a*, generally referred as ''aquatic environmental parameters''. Variation

partitioning analysis (VPA, Legendre*, et al.*, 2012) was used to describe the partitioning of

variation in water microbiome among three data sets: ''Local Land Use'' (the fractional land

use within 500 m of the water body) and ''Basin Land Use'' (the fractional land use within

the drainage basin of each water body) and aquatic environmental parameters, see

Supplementary Excel file 1, sheet "metadata". For the analysis, we included thirteen land use

types in the basin data-set and ten in the local (see Supplementary Data File 1). Land use

types which were measurable at less in four sites were removed from analysis (e.g. Basin

scale: Industrial areas, Construction areas, Artificial land cover, Crops, Vegetation, Open

1089   agriculture spaces and Wetlands; Local scale: Industrial areas, Artificial land cover,

1090   Herbaceous areas and Wetlands). We note that Forest and Arable lands have a strong

1091   negative correlation (Pearson $r = -0.86$), however, since in most basins these two land-use

1092   types contributes as much as 70-90% of the total area we did not remove one in order to

1093   avoid biases due to different total land-use values. The final table of land uses included six

1094   types in both scales: Urban fabric, Heterogenous agriculture, Pastures, Forests, Arable lands

1095   and Water bodies (see Supplementary Data file 1).

In order to validate the VPA results and further describe the effects of the different land use

types on OTU patters, we performed canonical correspondence analysis (CCA) and

permutation test (nperm=1000) using cca and anova.cca commands in R package 'vegan'.

We also considered whether the models have significant independent explanatory power by

testing conditioned CCA. In the case of testing for model "Basin Land Use", "Local Land

Use" parameters were used as conditioning variables and vice versa.

VPA and CCA were also used to evaluate partitioning of variation in the bacterial community

structure with three groups of explanatory variables: ''aquatic environmental parameters'',

''Local Land Use'', and ''Basin Land Use'' (Supplementary Data file 1). Since varpart

cannot be performed with missing data, we had to remove four sites (*i.e.* HAR, S.wen, Wen,

F.P, Supplementary Data file 1) from the data-set before performing the analysis. Part of the

removed samples are located around Berlin, for which we did not have detailed information

on the drainage basin, and part were missing aquatic environmental measurements due to

technical problems (I.e. Wan, Has and Hol).

1110

1111 **Supplementary Table 1:** Statistical values of each of the land uses in the GLMs correlations

1112   to water properties. The following abbreviations are used: U – Urban areas, Ar –

1113   Heterogenous agriculture, Pas – Pastures lands, Ag – Heterogeneous agriculture, For –

1114   Forests, W – Water bodies, *chl* a – *chlorophyll* a, TP – total phosphorus

Water parameter	Land uses	Level of land use	Model's R^2	Main contributing land uses for the correlation
NO_2+NO_3	U+Ar+Pas+Ag+For+W	Local	0.17	Arable (+)
	U+Ar+Pas+Ag+For+W	Basin	0.74	Urban (+)
pH	U+Ar+Pas+Ag+For+W	Local	0.13	–
	U+Ar+Pas+Ag+For+W	Basin	0.25	Forests (-)
Ammonium	U+Ar+Pas+Ag+For+W	Local	0.08	–
	U+Ar+Pas+Ag+For+W	Basin	0.06	–
P	U+Ar+Pas+Ag+For+W	Local	0.02	–
	U+Ar+Pas+Ag+For+W	Basin	0.14	–
Chl a	U+Ar+Pas+Ag+For+W	Local	0.06	–
	U+Ar+Pas+Ag+For+W	Basin	0.078	–

1115

1116

1117 **Supplementary Table 2** Importance values of each of the basin's land uses in the GLMs.

1118   Abbreviations: *Chl a – chlorophyll a*, TP – total phosphorus

	Chl a	NO$_2$+NO$_3$	pH	Ammonium	TP
Agriculture area	0.26	0.33	0.075	0.16	0.14
Pastures lands	0.23	0.17	0.45	0.13	0.13
Arable Lands	0.3	0.16	0.71	0.32	0.12
Urban area	0.24	1.00	0.14	0.14	0.12
Forests	0.36	0.19	0.19	0.3	0.12
Water bodies	0.31	0.15	0.44	0.13	0.31

1119

1120 **Supplementary Table 3: Spearman correlations between OTUs abundance and**
1121 **different environmental parameters.** The only significant correlations occur between two
1122 OTUs and the water concentrations of NO$_2$+NO$_3$

1123

Phylum	Order	Class	Family	Genus	Enviro-nmental parameter	r2	Adj p val
Planctomyce-tes	Phycisphaerae	Phycisphae-rales	NA	NA	NO$_2$+NO$_3$	0.53	0
Bacteroide-tes	Saprospir-ae	Saprospiral-es	Saprospira-ceae	Aquirestis	NO$_2$+NO$_3$	0.41	0

1124

1125 **Supplementary Table 4: VPA and CCA results to describe the partitioning of water**
1126 **microbiome by variation in aquatic environmental parameters and land uses.**
1127 Significance of the CCA models tested using Anova. * represents statistical significance.

Analysis		Water chemistry	Basin Land Use	Local Land use
Variation partitioning	Total VE (%)	11.3*	5.5	1.3
	Independent VE (%)	11*	4.5	0
CCA	Inertia explained (%)	24.2*	17.6	13.9
	Significant variables ($P<0.05$)	Temperature; NO$_2$+NO$_3$		

		Total P; *Chl* a
Partial CCA (conditioned)	Inertia explained (%)	21.4
	Significant variables (*P*<0.05)	Total P; NO₂+NO₃

1128

References

Caporaso JG, Kuczynski J, Stombaugh J, *et al.* (2010) QIIME allows analysis of high-
throughput community sequencing data. *Nature Methods* **7**: 335.
Langmead B & Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nature*
Methods **9**: 357.
Legendre P, Borcard D & Roberts DW (2012) Variation partitioning involving orthogonal
spatial eigenfunction submodels. *Ecology* **93**: 1234-1240.
Oksanen J, Kindt R, Legendre P, O'Hara B, Stevens MHH, Oksanen MJ & Suggests M
(2007) The vegan package. *Community ecology package* **10**: 631-637.
Schloss PD, Westcott SL, Ryabin T*, et al.* (2009) Introducing mothur: open-source, platform-
independent, community-supported software for describing and comparing microbial
communities. *Applied and Environmental Microbiology* **75**: 7537-7541.
Zhang J, Kobert K, Flouri T & Stamatakis A (2013) PEAR: a fast and accurate Illumina
Paired-End reAd mergeR. *Bioinformatics* **30**: 614-620.

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157 **Supplementery Figures**

1158

1159

1160

1161

Fig S1. Detailed map of the sampling location, with the different sampling sites noted.
Sampling sites were chosen to represent a high diversity in environmental gradients, e.g.
variety in land use and water chemistry. These differences between the sites can be visualized
- even before our measurement: **a-c)** Lakes of the Havel basin. **a)** Grosser Priepertsee is lake
located within a town called Priepert and is used for boating and recreation, while **b)**
Grienericksee is located next to the city of Rheinsberg with clear blue water. **c)** Thymensee is
located in the middle of a forest area far away from any town. Arrows with numbers indicate
sites as are presented in Supplementary Excel table. Red lines represent connectivity between
drainage basins, as mentioned in the text. Notice that two of the sites studied (Hölzerner See
and Tollensesee) are outside the scope of this map. Their exact coordinates can be found in
the Supplementary Excel table under the numbers 45 and 46.

1173

1174

1175

1176

Fig S2. Relationship between trophic state and microbial community composition. a, b) The population structure of the samples lakes, ordered by chlorophyll (a) and total P concentrations (b) at the time of sampling. The codes for the location sites are colored based on the trophic state shown in the supplementary excel file, which is based on the monitoring programs of the German federal states Mecklenburg-Vorpommern and Brandenburg, following in general the rules of the water framework directive of the European Union (see supplementary text for more information).

A slight tendency is seen towards increased relative abundance of cyanobacteria when *Chl* a (A) and total P (B) are high. C) **Weak correlation between chlorophyll concentration and relative abundance of cyanobacteria**. See also the NMDS of figure S3.

Fig S3. NMDS analysis of the bacterial population structure, colored by trophic level (a) and by drainage basin (b). The NMDS was calculated using Bray-Curtis dissimilarities. K=2, Stress= 0.16

1194

1195

Fig S4. No clear correlation between basin size, water chemistry or microbiome
composition. a-d) Drainage basin size vs ammonium (a), total phosphorus (b), temperature
(c) and pH (d); no clear correlation can be seen. e) Low correlation between basin size and
Chl *a* concentration, and f) No clear correlation between the fraction of cyanobacteria from
the total microbiome and the basin size. g) No correlation between basin size and microbiome
similarity. Each point represents the Bray-Curtis dissimilarity between two sampling points,
plotted against the difference in size between the drainage basins of the two locations.

1203

Fig S5. No distance decay patterns observed in sub-basins. The three large basins in the
analysis were each divided into two smaller sub-basins, where all of the water bodies are
directly connected. No distance-decay patterns were observed (Spearmans r ranged between
0.1 and 0.26, p values 0.3-0.7). The water bodies used for each sub-basin are as follows (see
Supplementary excel File for water body codes). **Lychen East**: Lyc, Zen, Kue, Hos, Has;
Lychen north: Lyc, Zen, HS, BL, Has; **Rhin North**: G.Rhe, Gri, Tie, Tor, Zer; **Rhin East**:
Tor, Gud, G.Dol, Huw, Wut, Zer; **Havel North-east**: Thy, S.L_1, S.L_2, Sto, Fur, Swt;
Havel North: G.Pri, Sto, EIF, Fur, .Swt

Fig S6. Relationships between different phyla and their relative abundance in the combined microbiome of all sampling sites. a) Relative abundance of bacterial phyla among core, cosmopolitan, common, rare and unique OTUs. The core OTUs, found in all sampling locations, are dominated by Proteobacteria and Actinobacteria. In contrast, few cyanobacterial OTUs are found in all sampling locations, whereas many are cosmopolitan (found in >90% of the locations)**. B) Distribution of core, cosmopolitan, common, rare and unique OTUs among the different phyla.** Note the difference in the relative number of core and cosmopolitan OTUs between Actinobacteria, suggested to have similar communities

1222 in all sampling sites, and Cyanobacteria, which reveal niche differences between different
1223 orders.

1224

1225

1226

1227 **Fig S7. Absences of separation in the CCA ordination between several bacterial**
1228 **lineages:** a) Proteobacteria orders, b) Bacteroidetes orders, c) Actinobacteria families. No
1229 clear differentiation is seen between the orders or families.

1230