Development and validation of the Evaluation Platform In COPD (EPIC): a population-based outcomes model of COPD for Canada

Mohsen Sadatsafavi MD, PhD[1]*, Shahzad Ghanbarian PhD[1]*, Amin Adibi MSc[1], Kate M. Johnson MSc[1], J Mark FitzGerald MD[2], William Flanagan MSc[3], Stirling Bryan PhD[4,5], Don Sin MD[2]; *for the Canadian Respiratory Research Network#*
*Co-first authors
Other collaborators: Shawn Aaron, Jean Bourbeau, Wan Tan, Teresa To, Penny Brasher, Deirdre Hennessy, Jacek Kopec

1. Faculty of Medicine and Faculty of Pharmaceutical Sciences, University of British Columbia, Vancouver, British Columbia, Canada
2. Faculty of Medicine, University of British Columbia, Vancouver, British Columbia, Canada
3. Statistics Canada, Ottawa, Ontario, Canada
4. Centre for Clinical Epidemiology and Evaluation, Vancouver Coastal Health Institute, Vancouver, British Columbia, Canada
5. School of Population and Public Health, The University of British Columbia, Vancouver, British Columbia, Canada

Corresponding author: Mohsen Sadatsafavi
Faculty of Pharmaceutical Sciences,
University of British Columbia
7th Floor, 828 West 10th Avenue
2405 Wesbrook Mall
Vancouver, BC Canada V6T 1Z3
Tel: 604.875.5178 | Fax: 604.875.5179
Email: msafavi@mail.ubc.ca

Contact for preprint request: Mohsen Sadatsafavi (msafavi@mail.ubc.ca)

Funding source: This study was supported by the Canadian Institutes of Health Research (Grant #142238), Genome Canada, and the Canadian Respiratory Research Network. Mohsen Sadatsafavi is supported by a New Investigator Salary Award from the Canadian Institutes of Health Research and a Scholar Award from Michael Smith Institute of Health Research. Don Sin is a Tier 1 Canada Research Chair holder in chronic obstructive pulmonary disease.

* This work has been presented at the Canadian Agency for Drugs and Technologies in Health 2018 Symposium

Keywords: Chronic obstructive pulmonary disease; Disease simulation; Whole disease modeling; Cost-effectiveness analysis; Epidemiological projection

Word count: 5,429 (body), 273 (abstract)

Abstract

Background: We report the development, validation, and implementation of a population-based outcomes model of Chronic Obstructive Pulmonary Disease (COPD) for Canada.

Methods: Evaluation Platform in COPD (EPIC) is a discrete event simulation model of Canadians 40 years of age or older. Three core features are its open-population design (incorporating projections of future population growth and aging, and smoking trends), its incorporation of heterogeneity in lung function decline and burden of exacerbations, and it modeling of the full natural history of COPD. Multiple original data analyses, undertaken as part of the EPIC research program, as well as values reported in the literature, were used to populate the model. Extensive face validity as well as internal and external validity evaluations were performed.

Results: The model was internally validated on demographic projections, mortality rates, lung function trajectories, COPD exacerbations, and stability of COPD prevalence over time within strata of risk factors. In external validation, simulated values for the rate of any exacerbations, severe exacerbations, and mortality were consistent with the reported values in two independent trials.

Conclusions: EPIC is the first whole-disease, outcome- and policy-focused model of COPD for Canada. A key feature of EPIC is modeling between-individual variability in important components of the natural history of COPD, which enables its use in modeling the outcomes of scenarios related to 'Personalized Medicine' interventions. By modeling the natural history of COPD from its inception, EPIC is capable of modeling the outcomes of decisions across the entire care pathway of COPD. By using a dynamic population, it can model the future population-level burden of COPD in the face of temporal changes in risk factors such as smoking rates.

Background

Chronic Obstructive Pulmonary Disease (COPD) is one of the most common chronic diseases globally(1). The prevalence of COPD is increasing in many jurisdictions. Worldwide, the prevalence of COPD has increased by 44.2% between 1990 and 2005(1). In Canada, there are 2.6 million individuals with COPD, comprising 17% of the adult population(2). COPD is characterized by the progressive and generally irreversible loss of lung function. A defining feature of COPD is the periods of intensified disease activity, referred to as acute exacerbations(3). Severe exacerbations that require inpatient care are the second leading cause of hospitalization in Canada, following only childbirth(4).

COPD is diagnosed using spirometry, and is diagnosed as such if the ratio of forced expiratory volume at one second (FEV_1) to forced vital capacity (FVC) is less than 70%, or alternatively less than the Lower Limit of Normal (LLN). The widely used Global Initiative for Chronic Obstructive Lung Disease (GOLD) severity grades are based on the ratio of FEV_1 to its predicted value, with 80%, 50%, and 30% thresholds separating severity grades I to IV(5). These classifications are common across several other guidelines(5–7), which has resulted in a large body of evidence relating COPD severity grades to disease outcomes such as exacerbation rates and costs.

A good understanding of the future burden of diseases and the long-term consequences of clinical and policy decisions can support the search for efficiency and equity in healthcare(8). In most instances, objective projection of disease burden and outcomes are based on computer simulations(9). A common approach to predicting the future burden of diseases and the outcomes of decisions is to develop a *de novo* computer model for the specific question at hand(10). For example, different groups have created cohort-based (e.g., Markov) models of COPD to predict the future costs and outcomes of COPD in specific countries(11), to evaluate the outcomes of various strategies (e.g., improving smoking cessation

rates, reducing the rate of exacerbations) for reducing the burden of COPD(12), and to evaluate the cost-effectiveness of bronchodilator therapies across different countries(13). However, this piecemeal approach has been criticized due to the resulting duplicate efforts, inconsistent assumptions, and risk of bias due to the influence of stakeholder interests when the model is developed with a particular research question in mind(14).

An alternative to this piecemeal approach is to create reference models that can be used for multiple purposes(8). Reference models are the result of dedicated scientific efforts, typically with more investment integrated into the overall design, better representation of disease mechanisms and outcomes, more comprehensive evidence synthesis, and more rigorous validation exercises than the *de novo* models(15). Once developed, these reference models can be used to evaluate the outcomes of many different scenarios within a consistent framework. A number of models have been developed for COPD, and a recent systematic review has identified the strengths and weaknesses of each of the different platforms(16). The majority of these models (with some exceptions(17)) simulate a cohort of patients who initially have the disease, and therefore they cannot be used to project the future incidence of the disease and the impact of preventive scenarios such as changing smoking rates and implementation of screening programs for earlier detection of COPD.

The Evaluation Platform In COPD (EPIC) was a nationally funded research project with the aim of creating an open-source, publicly available, population-based 'Whole Disease' COPD model for epidemiological projections and policy analyses in the Canadian context. The present paper discusses the concept and overall methodology, assumptions, data sources, and results of internal and external validation.

Methods

Conceptual framework

The conceptual framework underlying EPIC is that of Whole Disease Modeling(14). This framework emphasizes the transferability of the decision node across the care pathway of the disease, and the capacity for incorporating interactive decision nodes such that the effect of implementation of a policy (e.g., the provision of a nationwide screening program for COPD) will appropriately affect the outcomes of downstream interventions (e.g., pharmacotherapy for diagnosed patients)(18).

The investigator team included key stakeholder groups in the research program including clinical experts, methodologists, patients, and policy makers. We followed guideline recommendations on the proper conceptualization of a decision-analytic model(19). **Figure 1** provides a schematic illustration of the EPIC model. During the formative stages of this project, the team, during multiple rounds of discussions, agreed on the following four features to be the core components of the platform: 1) COPD should be modeled using the reference standard (spirometric definition) so that the model can make projections about the 'true' burden of COPD and the merits of screening/case detection strategies; 2) lung function trajectories and acute exacerbations are major factors impacting the natural history of COPD and should be simulated with high precision as they are ubiquitously the primary end points of clinical trials; 3) EPIC should realistically model heterogeneity in the natural history of COPD to enable modeling scenarios pertinent to Precision Medicine (e.g., the use of diagnostic and prognostic biomarkers for COPD diagnosis and treatment); 4) sex, age, and smoking, the three main risk factors for COPD incidence, should be modeled explicitly. A consequence of the latter feature is the assumption that any changes in the prevalence of COPD over time is due to changes in the distribution of these three risk factors in the population and/or changes in the life expectancy of the population with and without COPD. Additional features of the disease (e.g., symptoms, treatments), other risk factors (such

as environmental factors, history of asthma), and comorbid conditions will be added gradually through successive iterations.

<<Figure 1>>

EPIC is a Discrete Event Simulation (DES) of the general Canadian population. DES models the life history of individuals, one at a time, from entrance into the model to death (or end of the simulation time horizon)(20). The progression occurs during jumps in time (which is a continuous variable) from a set of pre-defined 'events' (e.g., COPD incidence, occurrence of an exacerbation, death – full list is available in **Appendix 1**). An individual is defined by a set of 'variables' (such as sex, current age, current FEV_1, individual-specific rate of FEV_1 decline and rate of exacerbations) whose values determine the rate of the occurrence of each event, and events in turn affect the value of variables defining the individual. Contrary to cohort-based models (such as the popular Markov models), individual-specific variables can create a virtually unlimited set of disease states, thus enabling robust modeling of heterogeneity and complex combination of disease states(20).

Model components and sources of evidence

EPIC can be seen as a series of inter-connected components (modules), each of which models a different aspect of the disease. Current modules include 1) demographic and risk factors, 2) COPD occurrence (prevalence and incidence), 3) lung function, 4) exacerbations, 5) costs and quality of life outcomes (payoffs), and 6) mortality. **Table 1** provides the structure of equations modeling various disease aspects for each module. The value of parameters (e.g., regression coefficients) are provided in separate tables in the appendices.

<<Table 1>>

1. Demographic and risk factor module

EPIC is a dynamic-population model of Canadian adults ≥40 years of age. As opposed to static models that simulate follow-up of a closed cohort of individuals for a given time (e.g., 10 years), dynamic models focus on the whole population within a specific date range (e.g., 2010 to 2030) and include births, deaths, immigration, and emigration. This is instrumental to the accurate modeling of temporal trends in risk factors such as population aging and changing smoking rates, and accommodating realistic aspects of policy implementations such as gradual market penetration of new medications or phased implementation of a population-based intervention such as a COPD screening program.

The demographic module of EPIC was based on Population Health Model (POHEM), a rigorously validated model of the risk factors and anthropomorphic and socio-demographic characteristics (e.g., height, weight, socioeconomic status) of Canadians, developed by Statistics Canada(21). POHEM has performed robustly in multiple internal and external validation challenges(21). POHEM models the Canadian population 20 years or older at year 2001 (prevalent population) based on the data from the 2001 Canadian Community Health Survey (CCHS). Individuals are followed over time and new individuals (incident population) enter the model in future years according to the projection of population growth and aging. This module accommodates the projected increase in life expectancy over time through the modification of future life tables. This module is also responsible for modeling smoking-related variables (current smoking status and pack-years of smoking). Changes in smoking status are modeled as transition rates estimated from various population-based Canadian surveys including the Canadian Health Study (1979), National Population Health Survey (1994), and CCHS (2008). The smoking pack-years variable is dynamically updated (annually or at the time smoking status changes).

2. COPD occurrence module

This module is responsible for simulating pre-existing and incident COPD. The former refers to COPD that already exists when an individual is first simulated (this is typically for the prevalent population in year 2001, and for immigrants who arrive to Canada after 2001 at older ages), while the latter occurs in simulated individuals as they age, who at the time of creation did not have COPD. COPD was defined as the ratio of FEV_1 to FVC being below its lower limit of normal (LLN)(22). The use of LLN-based definition, as opposed to the fixed (FEV_1/FVC< 0.70) criterion, was based on suggestions from the leading clinical experts in our team reflecting the growing support of the former definition due to its better properties (e.g., association with COPD risk factors) in the younger and older groups(22). EPIC does not model lung function in the general non-COPD population; instead, the occurrence of incident COPD is considered as an event whose risk is determined by the individual's characteristics. Lung function is then modeled in individuals in whom the COPD switch is turned on.

Pre-existing COPD: We used data from the Canadian Cohort of Obstructive Lung Disease (COLD) to assign a binary COPD status to individuals upon their creation. COLD was a nationally representative, population-based cross-sectional study of 3,042 Canadians 40 years of age or older(23). Using these data, we developed sex-specific regression equations for the probability of having COPD as a function of sex, age, current smoking status, and smoking pack-years (**Table 1** and **Appendix 2**).

Incident COPD: The sex-specific hazard of developing COPD was modeled as a log-linear function based on an individual's sex, age, smoking pack-years, and current smoking status (**Table 1** and **Appendix 2**). The incident equation was derived based on a 'steady state' assumption: that COPD incidence is such that COPD prevalence remains stable over time within the strata of risk factors (age, sex, and smoking pack-years) if the age-specific mortality in the population remains constant over time. These

assumptions establish a mathematical relationship between incidence and prevalence, enabling us to derive incidence equations from the prevalence equation(24). A stochastic optimization approach was adopted to solve for the coefficients in the incidence equation. Details of the calibration process are provided in *Appendix 2*.

3. Lung function module

Once the COPD designation is defined for an individual, an individual-specific initial FEV_1 value and an individual-specific annual rate of FEV_1 decline (representing heterogeneity in lung function decline) is assigned. The three components of this module are the initial FEV_1 value for pre-existing (prevalent) COPD cases, initial FEV_1 values for incident COPD cases, and the slope of decline in FEV_1 over time. For the first two components, we relied on patient-level data from the nationally representative COLD study(25). Sex-specific regression equations were used to estimate FEV_1 given risk factors (*Table 1* and *Appendix 3*).

FEV_1 decline was modeled based on individualized equations based on the Lung Health Study (LHS) data (n=5,887), as described in detail elsewhere(26). These equations performed robustly in internal validation as well as external validation in two independent cohorts(26). In brief, we fitted a random-intercept, random-slope model for FEV_1 decline based on 11 years of data from the LHS with covariates being age, sex, weight, height, and current smoking status (*Table 1*, given that the original equation had other risk variables such bronchial hyper-responsivity not modeled in EPIC, we refitted the original model with reduced variables – results are provided in *Appendix 3*). An additional quadratic term for time was included to capture the potential acceleration in lung function decline over time.

GOLD severity grades: Much of the evidence on COPD outcomes (e.g., costs, quality of life, exacerbation rates) is available according to the GOLD severity grades which are based on FEV_1 to its predicted ratio.

Calculating the predicted FEV_1 is based on individuals' age, sex, and height using the National Health and Nutrition Examination Survey equations(27).

4. Exacerbation module

The exacerbation module simulates the exact timing of acute exacerbations of COPD, as well as their duration and severity. Exacerbations were defined in four severity levels according to the event-based definition of exacerbations (which is based on healthcare services use)(28): mild exacerbations are those that result in the intensification of symptoms but do not require an encounter with the healthcare system; moderate exacerbations are those in which patient visits a physician or emergency department but is not hospitalized; severe exacerbations are those that result in hospital admissions; and very severe exacerbations are those that result in admission to the Intensive Care Unit.

We used separate sources for estimating the average rate of exacerbations and between-individual variability (heterogeneity) in rates (***Table 1***). The average rate was modeled as a function of the GOLD severity grades according to the pooled analysis by Hoogendoorn et. al.(29). The ratio of severe/very severe to total exacerbations was selected from the same study. However, this source did not provide the ratio of mild to moderate or severe to very severe exacerbations. These ratios were selected from a dedicated analysis of the Azithromycin for prevention of exacerbations of COPD study (the MACRO study), a large (n=1,142) one-year clinical trial of azithromycin therapy in preventing exacerbations(30). The instantaneous rate of exacerbation (hazard) was modeled as a log-linear function of variables; similarly, a proportional odds model was used to assign probabilities for various exacerbation severity levels (***Table 1***).

The rate and severity of exacerbations are substantially different among COPD patients(31). As such, projecting population-based outcomes of scenarios that require specific exacerbation history in patients (e.g., provision of preventive therapy in individuals with two or more exacerbations during the previous year) requires nuanced modeling of heterogeneity in the burden of exacerbations. In EPIC, simulated individuals have their specific rate of exacerbations, as well as their specific probability for an exacerbation, once it occurs, to be at a given level of severity. The extent of such heterogeneity was based on our previous work using data from the MACRO study(32). This heterogeneity was modeled through individual-specific (random-effect) intercepts for log-hazard of exacerbation rate and log-odds of exacerbation severity(32).

5. Mortality module

This module models death from two sources: death due to severe or very severe COPD exacerbations, and background mortality (due to other causes). Background mortality rates were taken from Statistics Canada's Canadian Life Tables and incorporate the projected declining mortality rates in the future(33). In addition, the probability of death from severe or very severe exacerbations was obtained from the literature (**Table 1**). Explicit modeling of COPD-related mortality in EPIC required calibration of background mortality rates so that the combined sources of mortality would produce mortality rates that resemble the sex-specific Canadian life tables. Finally, we assumed individuals will die of a severe exacerbation once they reach an FEV$_1$ of 0.5.

6. Payoff module

The payoff module assigns costs and health state utility values to events and collects the aggregate population values. These values were chosen from the literature after a dedicated literature review. EPIC currently does not model background medical costs and quality of life in the general non-COPD

population. Only COPD-specific direct medical costs are considered, and health state utility values are only assigned to patients with COPD. Two cost components are modeled. Annual 'maintenance' costs of COPD are those that accrue outside of episodes of exacerbations. These costs were modeled according to the GOLD grades from previously published studies(34). The second component is costs associated with exacerbations, which were modeled separately for each level of exacerbation severity based on Canadian studies(12,35). All costs are adjusted to 2015 Canadian dollars using annual inflation rates. Similarly, two sets of Quality Adjusted Life Years (QALY) weights (utilities) were modeled: utilities outside exacerbation episodes according to GOLD grades from the published literature, and a direct reduction in QALY associated with exacerbations given their level of severity(34).

Simulation platform

EPIC was programmed in C++. Two copies of the model were developed: one in Statistics Canada's Model Generator, a set of libraries and header files for DES programming implemented in Microsoft® Visual Studio, with pre-existing code representing POHEM(36). An identical open-source and open-access copy was developed with an interface in R (v3.4, with C++ code being called from the R functions using the Rcpp package)(37). The latter provides easy access to the model and online availability through a Web Application (http://resp.core.ubc.ca/ipress/EPIC). EPIC code is publicly available at https://github.com/aminadibi/epicR and can be downloaded and run as an R package.

Examining face validity and internal validity

Face validity refers to whether the model produces sensible results and if changing input parameters affects model output in the expected direction. Two important sets of face validity targets were the gradient of outcomes across categories of risk factors (e.g., higher COPD among ever-smokers than never-smokers), and the stability of outcomes over time within risk factor strata. Face validity results are not reported here but are available upon request. Internal validity was assessed by examining if the

model produced results that matched the patterns observed in input data sources or calibration targets. These included **1)** comparing the simulated population growth and aging with Statistics Canada's projections (validating demographic and mortality modules), **2)** comparing age-specific all-cause mortality rates with Statistic Canada's life table estimates (validating the mortality module), **3)** comparing COPD prevalence and GOLD grades by risk factor groups with estimates from the COLD study (validating the COPD occurrence and lung function modules), **4)** comparing average and 95% prediction interval of FEV_1 decline with prediction equations from LHS (validating the lung function module), and **5)** comparing the rate of total and severe exacerbations per GOLD severity grades with the original pooled estimates by Hoogendoorn et. al. (29) and results from the MACRO study (validating the exacerbation module).

Examining external validity

Ensuring external validity requires examining the model output against results from external sources of evidence that were not used in the model as input parameters or calibration targets. The external validity tests were based on the reported outcomes from the placebo arms of two independent clinical trials: TOwards a Revolution in COPD Health (TORCH)(38) and Understanding Potential Long-Term Impacts on Function with Tiotropium (UPLIFT)(39). These studies have been used by other COPD modeling groups as external validity targets and their use here also provides cross-validity between EPIC and existing COPD models(16,17,40–44). The external validity tests involved creating samples that mimicked the study sample of TORCH and UPLIFT to compare three quantities: rate of overall exacerbations, rate of severe exacerbations, and percentage of patients who died during follow-up. Rejection sampling was used to create EPIC samples that matched the baseline characteristics (age, sex, smoking status, FEV_1, and GOLD grades) of patients in TORCH and UPLIFT (details in ***Appendix 4***).

Results

Model calibration and internal validation

Demographic and mortality modules

After incorporating COPD-specific mortality and calibrating background mortality, the model generated age-specific all-cause mortality rates that were close to the Canadian life tables (*Figure 2)*.

<<Figure 2>>

The model performed robustly in predicting population growth and aging in the future. *Figure 3* presents the simulated (EPIC's) and projected (Statistics Canada's) population growth, as well as population age pyramid for three exemplary years (2015, 2025, and 2034).

<<Figure 3>>

COPD occurrence module

After the model was calibrated using the methodology presented in *Appendix 2*, the coefficient for calendar year in the logistic model of COPD prevalence (controlling for sex, age, and smoking) was 0.00008 (95%CI -0.00103 – 0.00119) for males and 0.00010 (95% CI -0.00008 – 0.00207) for females, indicating that the incidence and prevalence equations were coherent. The model performed robustly in replicating the COPD prevalence by sex and age groups as observed in the COLD study (*Figure 4*).

<<Figure 4>>

Lung function module

The equation for FEV_1 decline had shown robust internal validity (against LHS) as well as external validity against two independent cohorts. Details of the validation process for FEV_1 equations are provided in its original report[26]. Given these results, to validate this module we compared the simulated FEV_1s from EPIC against those predicted using the LHS-based equations. This enabled us to evaluate both the overall

similarity in FEV_1 values as well as the heterogeneity in FEV_1s, by determining the percentage of simulated FEV_1 values that fell within the 95% prediction interval provided by the LHS-based equation. *Figure 5* presents the results across four subgroups of subjects (based on permutations of sex and smoking status at baseline). Overall, the FEV_1 values simulated in EPIC were similar to the LHS-based predictions in terms of the mean and standard deviation across the range of possible values. The probabilities that the FEV_1 value simulated in EPIC fell within the LHS-based 95% prediction interval were between 94% and 96% for all four subgroups.

<<Figure 5>>

Exacerbation module

Figure 6 shows the EPIC-simulated annual rate of total and severe exacerbations stratified by GOLD grade compared with those reported in Hoogendoorn et. al. (29). EPIC mildly underestimated the rate of both total and severe exacerbations in GOLD grade I; nonetheless the simulated values were within the 95% confidence intervals of the original pooled estimates.

<<Figure 6>>

External validation

After rejection sampling, the baseline characteristics of the simulated sample in EPIC resembled those of the original samples of TORCH and UPLIFT (*Appendix 4*). *Figure 7* provides the results of external validation tests.

<<Figure 7>>

EPIC, along with two other COPD models (Hoogendoorn et. al.(17) and Samyshkin et. al. (44)) produced a higher total exacerbation rate than was observed in the UPLIFT trial (1.55/patient-year in EPIC versus

0.85/patient-year in UPLIFT, top panel of *Figure 7*). Similarly, EPIC and the same two models

overestimated exacerbation rate in TORCH (1.51/patient-year in EPIC versus 1.13/patient-year in

TORCH). EPIC-generated values were outside of the 95% confidence interval [95%CI] of both studies.

However, EPIC did replicate the higher exacerbation rate observed in TORCH compared with UPLIFT.

EPIC simulated a slightly higher rate of severe exacerbations than was observed in TORCH and UPLIFT

(0.21/patient-year in EPIC versus 0.17/patient-year in UPLIFT, and 0.22/patient-year in EPIC versus

0.19/patient-year in TORCH, middle panel of *Figure 7*). EPIC-generated estimate was outside of the

95%CI of UPLIFT. Three other models (Briggs(42), Hoogendoorn(17) and Samyshkin(44)) also generated

higher rates than was observed in the trials. EPIC simulated a lower mortality rate over the trial follow-

up (12.8%, bottom panel in *Figure 7*) than was observed in TORCH (15.2%); EPIC-generated value was

outside of the 95%CI. EPIC generated a similar mortality rate (17.3%) to what was observed in UPLIFT

(16.3%).

Discussion

We reported the conceptualization, development, implementation, as well as internal and external validation of a population-based microsimulation outcomes model of COPD in Canada. This platform was developed to provide the capacity to project the burden of COPD given changing risk factors and arrival of new technologies for primary, secondary, and tertiary care prevention. The major components of the model were simulated life trajectories of Canadians, equations describing the incidence and prevalence of spirometrically-defined COPD, the individual-specific progression of lung function decline in COPD, the individual-specific rate and severity of COPD exacerbations, COPD-related mortality rates, and estimates of costs and quality of life weights associated with different levels of disease severity and exacerbation status. The use of an open population is a key feature of the model as it provides the capacity to project future outcomes while considering temporal trends in risk factors such as aging and smoking. It also enables gradual market penetration of interventions and adoption of policies to be modeled, as well as the impact of sub-optimal adherence to treatments and guidelines. This was made possible by access to the population-based cohort in which COPD was defined spirometrically (as opposed to clinical cohorts that are typically based on convenience sampling of diagnosed patients in the community). Modeling heterogeneity in key components of the disease mechanisms (lung function decline and exacerbations) is another important feature of EPIC which should facilitate the evaluation of scenarios pertaining to complex decision rules for Precision Medicine (e.g., the provision of preventive exacerbation therapy for individuals with a given exacerbation history, or aggressive smoking cessation for patients above a threshold rate of lung function decline). In developing EPIC, we placed a major emphasis on using the most relevant sources of evidence for different components of the model. This included an extensively validated pre-existing model of socio-demographic characteristics of the Canadian population and trends in the distribution of risk factors (e.g., smoking), a secondary analysis of the landmark LHS study for modeling FEV_1 trajectories, a secondary analysis of a population-based

Canadian study for modeling COPD incidence and prevalence, and a secondary analysis of a recent clinical trial (whose primary outcome was exacerbations) to quantify heterogeneity in the burden of exacerbations.

The conceptual framework for EPIC was that of 'Whole Disease Modeling,' which recognizes that the outcome of a given decision across the care pathway of COPD is inevitably affected by the outcomes of other decisions, thus requiring the modeling of the entire disease pathway(14). As a case in point, critiques of population-based screening and opportunistic case finding strategies for undiagnosed COPD refer to the lack of effectiveness in available COPD treatments(45). However, with the arrival of more effective therapies, the benefit-risk and cost-effectiveness profiles of screening and case finding for COPD is likely to change. Implementation of biomarkers for earlier and more accurate detection of COPD exacerbations can in itself affect the outcomes of screening as well as therapies. A unified model that can project the impact of these scenarios can be used to periodically revisit the merits of each decision after upstream or downstream changes in the care pathway. Such a unified model can also provide opportunities for the redistribution of resources and potential disinvestments within the context of a given disease(14).

Overall the model performed robustly in face validity and internal validity tests. This should be reassuring in terms of the validity of the original statistical analyses, model calibration exercises, and model implementation. To some extent this also indicates the consistency of estimates from original data sources and input parameters taken from the literature. For instance, incidence and prevalence equations as well as equations for 'baseline' lung function were determined from the COLD study, whereas lung function decline equations were adapted from the LHS. These equations seem to be congruent resulting in a stationary distribution of COPD severity and distribution of COPD exacerbations over time. In general, the model performed satisfactorily in the external validity of the model against two independent studies that have also been used to validate other COPD models. It overestimated the

rate of overall exacerbation, but its simulated exacerbation rates were similar to two other COPD

models. For rate of severe exacerbations and proportion of patients who died during the follow-up

period of original studies, its predictions were better aligned with the reported values and were

generally around the mid-point of simulated values from five other COPD models. The model replicated

the direction of the differences in exacerbation and mortality rates between the two trials, which

indicates its responsiveness in propagating changes in risk factors to changes in outcomes.

The limitations of EPIC also need to be mentioned. This work represents the core structure of a COPD

'Whole Disease' model, but much work needs to be done to create a comprehensive COPD outcomes

model. While in its current form EPIC is capable of making projections about the future burden of COPD

for a broad range of policies, there are many other aspects of COPD which need to be incorporated.

Examples include treatment, symptoms, diagnosis, and comorbid conditions. Further, EPIC does not

currently incorporate uncertainty in the underlying evidence, which is a goal for the next iteration of

EPIC. Given that many parameters of EPIC are derived based on calibration techniques, this is not simply

a matter of replacing fixed input parameters with their probabilistic equivalent. More sophisticated

model calibration techniques might prove useful in this context(46). Further, EPIC is currently limited to

modeling the impact of sex, age, and smoking as risk factors for COPD, but does not consider other

factors such as occupational and environmental exposures or history of asthma. Because lung function

in the non-COPD population was not modeled, preventive strategies based on monitoring lung function

prior to persistent airflow obstruction cannot be assessed. Similarly, prodromal phase of exacerbations

and exacerbation phenotypes (e.g., infectious versus inflammatory) are not currently modeled, limiting

the applicability of EPIC in simulating scenarios pertaining to the early detection of exacerbations. As

well, the potential correlation between the propensity to exacerbate and the rate of lung function

decline were deemed too uncertain to be incorporated into the model. Overall, EPIC can currently be

used for modeling the outcomes of disease management based on lung function, exacerbation history,

and smoking history. However, it cannot provide this capacity for disease management based on symptoms or burden of comorbidity. The planned iterative upgrades to the platform should see these features gradually incorporated into the platform.

EPIC is not the first multi-purpose model of COPD. We have previously performed a systematic review of published COPD models that were used for epidemiological projections or economic evaluations(47). The majority of models were created for addressing a specific question (e.g., cost-effectiveness of a novel therapy). However, in some instances, subsequent models were built on previous ones in an iterative fashion. The presence of such an identifiable 'lineage' of models demonstrates the need for, and feasibility of, re-using an existing model for different objectives. Recently, a convention of model developers assessed the consistency of five different models in COPD and their external validity(16,48,49). Putting our developed model in the context of previous reference models is informative. Among the three outcomes of overall exacerbation, severe exacerbation, and mortality rates, EPIC-generated values are within the range generated by the other five models, with rate of exacerbations toward the higher end of the range. Among the previously reviewed models, only the model by Hoogendoorn et. al., which was developed for the Netherlands, is a dynamic population model(17). EPIC appears to be unique in modeling heterogeneity at the individual level in lung function and the rate and severity of exacerbations. Briggs et. al., developed a model of COPD disease progression based on the ECLIPSE(50) data (with resource-use estimates from UPLIFT)(42). The model was based on a rigorous practice of model conceptualization(51), and has performed satisfactorily in internal and external validation studies(16,42). A major advantage of this model is its incorporation of multiple aspects of COPD (such as symptoms and six-minute walking distance, a measure of functional capacity). However, reliance on mainly one study (ECLIPSE) makes this model less relevant to the Canadian context, and three-year follow-up of ECLIPSE makes long-term projections less certain.

Predicting the future using the best available evidence is a cornerstone of many disciplines. In healthcare, the use of modeling to reconcile evidence from multiple sources to project the outcomes of a disease and healthcare decision making is considered "an unavoidable fact of life"(9). Model-based projections can critically inform recommendations for best practices as well as policies. Reference models that encapsulate concerted efforts in evidence synthesis, validation, and implementation exists for many health conditions(52–54). For cardiovascular diseases, reference models have critically informed policies for more than 30 years(55) and continue to do so with high impact, paradigm shifting evaluations(56). Given the comparable and escalating burden of COPD, the need for such projections for this disease clearly exists. The recent efforts in developing reference COPD models are valuable responses to such a need, and EPIC adds to the portfolio of such efforts. While EPIC is developed for the Canadian context, many of the components of EPIC pertain to the natural history and biology of COPD and are independent of a particular setting. Other components can be updated (mainly by changing numerical values of input parameters rather than modifying the structure of equations) with the specifics of a healthcare setting. As such, this platform can have applicability for many other jurisdictions. The recent organized efforts in ensuring external validity and cross-comparability of such models can help the development of robust platforms that address the ongoing needs of both the research and policy community.

Table 1: Input parameters (βs) and their related equations in Evaluation Platform In COPD (EPIC) <<<u>Return to text</u>>>

Parameter(s)	Description	Value(s)	Source/Reference
Demographic and risk factor module			
Demographic variables and risk factors	Sex, age, smoking status, height and weight	Complex equations representing the baseline 2001 Canadian population (20 years or older) from 2001 CCHS	Based on Population Health Model(21)
COPD occurrence module			
COPD prevalence equation	Assigns COPD status to simulated individuals at the time of creation	$logit(p \mid sex) = \beta_{0,sex} + \beta_{1,sex} \times age + \beta_{2,sex} \times pack_{years} + \beta_{3,sex} \times current_{smoker} + e$	Regression on COLD data considering lower limit of normal criteria (***Appendix 2***)
COPD incidence equations	Estimated for patients without COPD	$log(hazard \mid sex) = \beta_{0,sex} + \beta_{1,sex} \times age + \beta_{2,sex} \times pack\_year + e$	Stochastic calibration based on COPD prevalence equations (***Appendix 2***)
Lung function module			
FEV$_1$ for prevalent cases	Baseline FEV$_1$ value for individuals who have COPD at the time of creation	$FEV1 \mid sex \sim Normal(\beta_{0,sex} + \beta_{1,sex} \times age + \beta_{2,sex} \times PackYears + \beta_{3,sex} \times height^2, e)$	Regression on COLD data (***Appendix 3***)
FEV$_1$ at incidence	Baseline FEV$_1$ value for individuals who develop COPD during the time in the model	$FEV_1 \mid sex \sim Normal\left(\beta_{0,sex} + \beta_{1,sex} \times age + \beta_{2,sex} \times PackYears + \beta_{3,sex} \times height^2 + \beta_{4,sex} \times \frac{\frac{FEV1}{FVC}}{LLN}, e\right)$	Regression on COLD data (***Appendix 3***)
FEV$_1$ decline over time	Models FEV$_1$ at time t (in years) in the future based on a random-intercept-random-slope model	$FEV_1(t) \sim Normal(Y_0 + Y_1 \times t + Y_2 \times t^2, e)$ $Y_0 = \beta_0 + \beta_1 \times age + \beta_2 \times sex + \beta_3 \times weight + \beta_4 \times height + \beta_5 \times height^2 + \beta_6 \times smokingstatus + \beta_7 \times age \times height^2$ $Y_1 = \beta'_0 + \beta'_1 \times age + \beta'_2 \times sex + \beta'_3 \times weight + \beta'_4 \times height + \beta'_5 \times height^2 +$	From the Lung Health Study(26) (***Appendix 3***)

		$\beta'_6 \times smokingstatus + \beta'_7 \times age * height^2$ $$Y_2 = \beta''_0$$	
Predicted FEV$_1$	Assign GOLD severity grades	$E(FEV_1) = \beta_0 + \beta_1 \times age + \beta_2 \times GOLD2 + \beta_3 \times GOLD3 + \beta_4 \times GOLD4$	From NHANES III(27)
		Exacerbation module	
Exacerbation rate	Log-hazard of exacerbation	$log(hazard) = \beta_0 + \beta_1 \times age + \beta_2 \times GOLD + e$	Pooled estimates from Hoogendoorn et. al.(29); variability estimated form MACRO(30) (***Appendix 4***)
Severity of exacerbation	Assigns one of four levels of severity to exacerbations	$logit(\pi(Yj < k))$ $= \beta_{0,k} + \beta_1 \times age + \beta_2 \times sex + \beta_3 \times FEV_1 + \beta_4 \times smokingStatus + \beta_5 \times packYear + \beta_6 \times BMI + e$	From Hoogendoorn et. al.(29) and MACRO study(30) (***Appendix 4***)
		Mortality module	
Background mortality rate	Death due to non-COPD reasons	Canadian life tables with calibration to remove COPD-specific mortality	Modification of the Canadian life tables(33) such that the combined sources of mortality match the life table values
Mortality due to COPD exacerbations	Short-term mortality due to severe exacerbations	$logit(p) = \beta_0 + \beta_1 \times age$	(55)(***Appendix 4***)
		Pay-off module	
Direct maintenance costs of maintenance COPD	Annual direct medical costs not associated with exacerbations (in 2015 $CAD)	$cost = \beta_1 \times GOLD_1 + \beta_2 \times GOLD_2 + \beta_3 \times GOLD_3 + \beta_4 \times GOLD_4$	Spencer et. al. (34) and Mitmann et. al. (56)(***Appendix 5***)
Direct costs associated with exacerbations	Direct medical costs pertaining to an episode of exacerbation according to its level of severity (in 2015	$cost = \beta_1 \times mild + \beta_2 \times moderale + \beta_3 \times severe + \beta_4 \times very.severe$	Spencer et. al. (34) and Mitmann et. al. (56)(***Appendix 5***)

	$CAD)		
Background utility values in COPD	Health state utility values in patients with COPD according to GOLD grades outsides of an exacerbation episode	$$utility = \beta_1 \times GOLD_1 + \beta_2 \times GOLD_2 + \beta_3 \times GOLD_{3.4}$$	Spencer et. al. (34) and Najafzadeh et. al. (12) (**Appendix 5**)
Reduction in QALY due to exacerbations	Two sets of regression coefficients (s=0 & s=1) separately modeled reduction in QALY values for mild/moderate and severe/very severe exacerbations which were dependent on the underlying GOLD grade	$$\Delta_Q = \beta_{1,s} \times GOLD_1 + \beta_{2,s} \times GOLD_2 + \beta_{3,s} \times GOLD_{3.4}$$	Spencer et. al. (34), Najafzadeh et. al. (12), and Goossens et. al.(57) (**Appendix 5**)
EPIC: Evaluation Platform In COPD, **GOLDx:** Global initiative for Obstructive Lung Disease severity level x (x ranges from 1 to 4 representing mild, moderate, severe, and very severe grade)			

Figure 1: Schematic illustration of Evaluation Platform In COPD (EPIC) <<Return to text>>

Background mortality risk is applied to each simulated individual once a year, no matter where the individual is in the model pathway. Payoff module collects outcomes (number of deaths, hospitalizations, costs, quality-adjusted life years, etc) as they occur/accrue during the entire process.

Figure 2: Comparison of the results of EPIC for mortality within age groups by sex with Statistics Canada's life tables <<Return to text>>

Uncertainty in the estimates were negligible and are not presented on this graph. All values belong to year 2009

EPIC: *Evaluation Platform In COPD,* **STATCAN:** *Statistics Canada*

V 2018.06.08

Figure 3: Statistics Canada's projected versus EPIC's simulated growth and aging patterns of the Canadian population 40 years or older. Top-left: population size, top-right: age pyramid in year 2015, bottom-left: age pyramid in 2025, bottom-right: age pyramid in 2034 <<Return Text>>

Predicted values are from Statistics Canada; simulated values are from EPIC. Satistics Canada's midpoint (as opposed to high and low) projections were used

V 2018.06.08

Figure 4: Comparison of the results of EPIC for COPD prevalence by sex and age groups with the estimations from COLD data <<Return to Text>>

Error bars are 95% confidence intervals. Uncertainty around EPIC output were neglible as simulations were run with large sample sizes.

EPIC: Evaluatoin Platform in COPD, **COLD**: Canadial cohort of Obstructive Lung Disease

Figure 5: Scatterplot of simulated FEV₁s from EPIC versus predicted FEV₁s from LHS equations for up to 11 years <<Return to text>>

Grey bars are 95% prediction intervals around individual FEV$_1$ values predicted from LHS equations. Dark grey dots are instances in which the simulated FEV$_1$ fell within the 95% prediction interval, and light grey dots represent simulated values that fell outside the interval. Graphs are generated based on running the model with 100,000 individuals. For each individual in the model, FEV$_1$ was sampled once every year. Maximum follow-up was 11 years (LHS follow-up time). In line with the structure of the LHS equations, for this validation task, only individuals who were smokers at baseline (a random date) were followed. At random, individuals either remained smokers throughout follow-up (continuous smokers) or stopped smoking at baseline without any recurrence during follow-up time (former smokers). Simulated individuals die when their FEV$_1$ reaches 0.5

***EPIC:** Evaluation Platform In COPD,* **LHS:** *Lung Health Study,* **FEV$_1$:** *Forced Expiratory Volume at one second*

Figure 6: Top panel: Comparison of the results of EPIC for the rate of all (left) and severe (right) exacerbations with the pooled estimates from Hoogendoorn et. al. Bottom panel: comparison between the distribution of exacerbation severity between EPIC and MACRO study for males (left) and females (right) <<Return to text>>

Error bars represent confidence intervals around point estimates. Sampling errors for EPIC-related results are negligible as they are based on a large simulated sample.

EPIC: *Evaluation Platform In COPD.* **MACRO:** *Azithromycin for prevention of exacerbations of COPD study*

Figure 7: Comparison of the results of EPIC and five different COPD models with TORCH (light grey bars) and UPLIFFT (dark grey bars) for total exacerbations (top panel), severe exacerbation rate (middle panel), and mortality rate (bottom panel) <<Return to text>>

After rejection sampling, a total of 3,505 and 5,782 simulated patients remained in the simulated TORCH and UPLIFT groups, respectively.

Error bars are sample confidence intervals for microsimulation models (EPIC and Asukai, first-order uncertainty) while for other models represent uncertainty in outputs due to uncertainty in the model parameters (second-order uncertainty).

EPIC: Evaluation Platform In COPD, TORCH: TOwards a Revolution in COPD Health, UPLIFT: Understanding Potential Long-Term Impacts on Function with Tiotropium

Appendices

Appendix 1: list of events in the model

Event	Affected by	Consequences
Fixed annual (birthday)	None	Updates the following variables: FEV_1, Predicted FEV_1, weight, smoking pack-years
COPD occurrence	Age, sex, smoking history	When COPD occurs, a baseline FEV_1 value is assigned, as well as an individual-specific FEV_1 slope, exacerbation rate, and exacerbation severity probabilities
COPD exacerbation start	COPD status, age, sex, FEV1, smoking status	Sets the exacerbation status to 1 and assigns an exacerbation-related mortality probability
COPD exacerbation end	Exacerbation status	Sets the exacerbation status to 0
Smoking status change	Smoking status, sex, age, calendar year	If non-smoker, sets the status to current smoker, and vice versa. Updates smoking pack-years
Death (background mortality)	Age, sex, calendar year	Removes the individual from the simulation

Appendix 2: Prevalence and incidence equations for COPD

Table A2.1: Regression coefficients for the prevalence of COPD from the COLD study

	Female			Male		
	Estimate	Std. Error	Pr(>|t|)	Estimate	Std. Error	Pr(>|t|)
(Intercept)	-4.37729	0.344947	< 2e-16	-5.338	0.412996	< 2e-16
Age	0.031689	0.005568	1.26E-08	0.044918	0.006461	3.58E-12
Pack-year	0.025183	0.003164	1.73E-15	0.018734	0.002745	8.83E-12
Current smoker	0.636127	0.162765	9.30E-05	1.093586	0.179252	1.06E-09

Estimating incidence equation from prevalence

In the model, the sex-specific probability of having COPD (i.e., the prevalence) is estimated using a logit regression equation with coefficients determined from longitudinal clinical studies:

$$logit\ (P_i) = \beta_0 + \ \beta^T \cdot X_i \tag{1}$$

where i is patient index, and $X = [Age\ ,\ PackYearSmoking]$.

Since the micro-simulation model is a prospective cohort model, the hazard of developing COPD (i.e., the incidence) is required to model incident COPD cases. Let the log-hazard for the i[th] patient be:

$$ln\ (H_i) = \beta'_0 + \ \beta'^T \cdot X \tag{2}$$

As the onset of COPD is gradual, β' is very difficult to estimate in clinical studies. However, the model assumes that the incidence of COPD will remain constant within the strata defined by X_i. Consequently, with the right β' coefficient in place, $logit\ (P)$ should not depend on any factors other than *Age* and *PackYearSmoking, i.e. the coefficients will be constant over the course of the simulation.* This criterion can be used to estimate β' numerically, using an iterative approach. First, an initial value for β' is guessed. In each subsequent step, the initial guess for β' is updated proportional to the difference between prevalence at the end and at the beginning of the simulation, until the difference becomes negligible. Estimated coefficients for incidence hazard regression are shown in Table A2.2. Figures A2.1 and A2.2 show how the coefficients and residual errors converge with iterations.

V 2018.06.08

Figure A2.1 *Convergence of coefficient estimates to estimated values (top) and residual errors (bottom) for male subjects*

Figure A2.2 *Convergence of coefficient estimates to estimated values (top) and residual errors (bottom) for female subjects*

V 2018.06.08

Table A2.2: Regression coefficients for the incidence of COPD from the COLD study. Note that the numerical method used to estimate coefficients does not allow for calculation of standard errors.

	Female	Male
(Intercept)	-7.6681	-7.8024
Age	0.0274	0.0313
Pack-year	0.0401	0.0319

V 2018.06.08

Appendix 3: Lung function (FEV$_1$) equations

For pre-existing COPD, regression equations (with sex, age, and smoking status as independent variables) were developed among individuals in COLD with FEV$_1$/FVC<LLN (n=595). For incident COPD, FEV$_1$ needed to be determined at the exact time when FEV$_1$/FVC equals LLN. To achieve this, a regression model was fitted in the whole sample (n=5,121) with FEV$_1$/FVC as an additional independent variable. The conditional mean and standard deviation of FEV$_1$ at time of COPD incidence could be estimated by calculating predicted values from the regression equation by setting FEV$_1$/FVC=LLN for each individual. Random values corresponding to the distribution of regression residuals were added to simulated FEV$_1$s for both prevalent and incident cases.

Table A3.1: Regression coefficients for FEV$_1$ at creation for prevalent (pre-existing) COPD cases, and initial FEV$_1$ for incident COPD cases

	FEV1 for prevalent COPD				FEV1 for Incident COPD			
	Male		Female		Male		Female	
	Estimate (SD)	P value	Estimate (SD)	P value	Estimate (SD)	P value	Estimate (SD)	P value
Intercept	1.365 (0.584)	0.02	1.456 (0.366)	<0.001	1.489 (0.169)	<0.001	1.573 (0.104)	<0.001
Age	-0.0327 (0.0032)	<0.001	-0.0301 (0.0022)	<0.001	-0.032 (0.001)	<0.001	-0.0275 (0.0007)	<0.001
Smoking pack year	-0.0034 (0.0013)	0.007	-0.00711 (0.0097)	<0.001	-0.00428 (0.0005)	<0.001	-0.00465 (0.0004)	<0.001
Height squared	1.098 (0.167)	<0.001	0.937 (0.116)	<0.001	1.27 (0.04)	<0.001	0.972 (0.033)	<0.001
FEV1/FVC/LLN	--	--	--	--	-0.683 (0.042)	<0.001	-0.425 (0.025)	<0.001

Table A3.3: Characteristics of the LHS sample included in the regression analysis

Follow up time (years)	9.2 (2.9)
Baseline age (years)	48.4 (6.8)
Baseline FEV1* (L)	2.75 (0.63)
FEV1 % predicted	78.47% (9.06)
Weight (kg)	75.9 (15.1)
Height (m)	1.72 (0.09)
Methacholine responsiveness (O'Connor slope) †	-12.73 (23.4)

Pack-years of smoking	40 (19)
Sex	
Male	63%
Female	37%
Smoking status by year 5	
Sustained quitters	17%
Intermittent quitters	28%
Continuous smokers	55%
All values are means (standard deviations) except sex and smoking status. FEV1: forced expiratory volume in 1 second. SD: standard deviation. *: post bronchodilator.	
†: Unit is change in FEV1 per mg/ml change in methacholine concentration	

Table A3.4: Regression coefficients for FEV$_1$ decline over time using the LHS sample

	Baseline FEV1			Rate of FEV1		
Parameters	Effect(ml)	95% CI	P-Value	Effect (ml/year)	95% CI	P-Value
Intercept	1427.5	(-1317.48, 4172.48)	0.3	-154.3	(-435.17, 126.57)	0.3
Baseline age (year)	-5.08	(-17.29, 7.13)	0.4	2.34	(1.08, 3.61)	0.0003
Sex (male=1 vs female=0)	482.5	(456.41, 508.59)	<.0001	-7.62	(-10.31, -4.93)	<.0001
Weight (kg)	-0.49	(-1.26, 0.28)	0.2	0.126	(0.05, 0.21)	0.002
Height (m)	-1872.5	(-4892.27, 1147.27)	0.2	58.35	(-251.13, 367.83)	0.7
Height^2 (m^2)	1951.3	(1080.86, 2821.736)	<.0001	18.07	(-71.27, 107.41)	0.7
Current smoker* (vs. sustained quitters)	-92.21	(-116.95, -67.47)	<.0001	-30.74	(-33.17, -28.31)	<.0001
Baseline age* height^2 ‡	-8.32	(-12.39, -4.25)	<.0001	-0.93	(-1.35, -0.51)	<.0001
Time (year)	NA	NA	NA	-1.46	(-1.58, -1.33)	<.0001

Table A3.5: Estimated G-matrix for random-effect terms of the model

	Intercept	Slope
Intercept	0.1006	0.00087
Slope	0.00087	0.000759

Appendix 4: Exacerbation equations

Table 4.1: Coefficients for the log-rate of total exacerbations.

	Estimate
Intercept	-2.10
Age*	0.004
GOLD II	1.1
GOLD III	1.9
GOLD IV	2.4

* Based on model calibration to correspond to the known relationship between age and exacerbation rate. Other parameters are from the meta-analysis by Hoogendoorn et. al. (see Table 1 of the main text for reference)

Uncertainty around regression coefficients were not considered given that EPIC currently is not a probabilistic model.

Table 4.2: Regression coefficients for the logit model of severity of an exacerbation based on the analysis of MACRO data.

	Estimate	Std. Error
Intercept 1	1.091	1.220
Intercept 2	1.902	1.223
Intercept 3	5.208	1.251
Age	-0.007	0.012
SEX	-0.764	0.201
FEV_1	-0.003	0.002
Pack_years	-0.001	0.002
BMI	0.018	0.016
Current smoker (v. former smoker & never smoker)	0.348	0.255

Table 4.3: Coefficients for mortality due to exacerbations.

	Estimate
Intercept	-13
Age	0.049
Exacerbation severity	
Mild	0
Moderate	-2
Severe	7.4
Very severe	8

V 2018.06.08

Appendix 5: Pay-off values

Table 5.1: Costs and health state utility values associated with events modelled in EPIC.

Parameter(s)	Value(s)	Source/Reference
Direct costs of maintenance COPD therapy (outside exacerbations)	GOLD I: 144 GOLD II: 430 GOLD III: 628 GOLD IV: 628	See Table 1 of the main text
Direct costs associated with exacerbations	Mild: 161 Moderate: 161 Severe: 6501 Very severe: 6501	
Utility values outside exacerbation episodes	GOLD I: 0.81 GOLD II: 0.72 GOLD III & IV: 0.67	
Reduction in QALY due to mild and moderate exacerbation	GOLD I: 0.00246 GOLD II: 0.001698 GOLD III & IV: 0.00534	
Reduction in QALY due to severe and very severe exacerbation	GOLD I: 0.007973 GOLD II: 0.007479 GOLD III & IV: 0.007178	

Appendix 6: External validation results

Rejection sampling:

To create a cohort of simulated patients that mimics the TORCH study in terms of baseline characteristics, the following steps were undertaken:

1) Creating a large simulated sample from EPIC and assigning a random baseline (index) date to each of them.

2) Weighting each simulated individual based on their characteristics (sex, age, current smoking status, FEV_1 percent predicted value) at baseline given the marginal probability density or probability mass function of these characteristics in the external sample. Marginal probability density or mass functions were constructed from the reported baseline characteristics [Table 1] of the original study

3) Randomly including each simulated individual in the final sample according to probabilities that were proportional to these weights.

The same steps were undertaken to create a simulated cohort mimicking the UPLIFT study sample.

Table 6.1: Characteristics of the original UPLIFT® and TORCH® clinical trials and simulated samples from EPIC. After rejection sampling, a total of 3,505 and 5,782 simulated patients remained in the simulated TORCH and UPLIFT groups, respectively.

Characteristics	UPLIFT®		TORCH®	
	Original sample (n=1374138)	EPIC simulation	Original sample (n=23541)	EPIC simulation
N	3006	5782	1524	3505
Male	74	74.3	76	75.8
Age	65±9	65.5±8.5	65±8	66.4±7.8
Current smoker	30	30.5	43	42.6
FEV_1%predicted	47±13	47.9±12.5	44±12	44.6±11.2
GOLD2	45	43.6	35	36
GOLD3	44	46.9	50	48.5
GOLD4	9	9.5	15	15.5

References

1. Soriano JB, Abajobir AA, Abate KH, Abera SF, Agrawal A, Ahmed MB, et al. Global, regional, and national deaths, prevalence, disability-adjusted life years, and years lived with disability for chronic obstructive pulmonary disease and asthma, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. Lancet Respir Med [Internet]. 2017 Aug [cited 2017 Aug 22]; Available from: http://linkinghub.elsevier.com/retrieve/pii/S221326001730293X

2. Evans J, Chen Y, Camp PG, Bowie DM, McRae L. Estimating the prevalence of COPD in Canada: Reported diagnosis versus measured airflow obstruction. Health Rep. 2014 Mar;25(3):3–11.

3. Aaron SD. Management and prevention of exacerbations of COPD. BMJ. 2014;349:g5237.

4. Camp P, Levy R. A snapshot of chronic obstructive pulmonary disease in British Columbia and Canada. BCMJ. 2008;50(2):80.

5. From the Global Strategy for the Diagnosis, Management and Prevention of COPD. Global Initiative for Chronic Obstructive Lung Disease (GOLD) 2017 [Internet]. Available from: http://goldcopd.org

6. American Thoracic Society / European Respiratory Society Task Force. Standards for the Diagnosis and Management of Patients with COPD [Internet]. [Internet]. New York: American Thoracic Society; [cited 2014 Nov 27]. Report No.: Version 1.2. Available from: http://www.thoracic.org/clinical/copd-guidelines/

7. Gruffydd-Jones K, Loveridge C. The 2010 NICE COPD Guidelines: how do they compare with the GOLD guidelines? Prim Care Respir J J Gen Pract Airw Group. 2011 Jun;20(2):199–204.

8. Scotland G, Bryan S. Why Do Health Economists Promote Technology Adoption Rather Than the Search for Efficiency? A Proposal for a Change in Our Approach to Economic Evaluation in Health Care. Med Decis Mak Int J Soc Med Decis Mak. 2017;37(2):139–47.

9. Buxton MJ, Drummond MF, Van Hout BA, Prince RL, Sheldon TA, Szucs T, et al. Modelling in economic evaluation: an unavoidable fact of life. Health Econ. 1997;6(3):217–27.

10. Bellete B, Coberly J, Barnes GL, Ko C, Chaisson RE, Comstock GW, et al. Evaluation of a whole-blood interferon-gamma release assay for the detection of Mycobacterium tuberculosis infection in 2 study populations. Clin Infect Dis Off Publ Infect Dis Soc Am. 2002 Jun;34:1449–1456.

11. Nielsen R, Johannessen A, Benediktsdottir B, Gislason T, Buist AS, Gulsvik A, et al. Present and future costs of COPD in Iceland and Norway: results from the BOLD study. Eur Respir J Off J Eur Soc Clin Respir Physiol. 2009 Oct;34(4):850–7.

12. Najafzadeh M, Marra CA, Lynd LD, Sadatsafavi M, FitzGerald JM, McManus B, et al. Future Impact of Various Interventions on the Burden of COPD in Canada: A Dynamic Population Model. PLoS ONE. 2012 Oct 11;7(10):e46746.

13. Oostenbrink JB, Rutten-van Mölken MPMH, Monz BU, FitzGerald JM. Probabilistic Markov model to assess the cost-effectiveness of bronchodilator therapy in COPD patients in different countries. Value Health J Int Soc Pharmacoeconomics Outcomes Res. 8:32–46.

14. Tappenden P, Chilcott J, Brennan A, Squires H, Stevenson M. Whole disease modeling to inform resource allocation decisions in cancer: a methodological framework. Value Health J Int Soc Pharmacoeconomics Outcomes Res. 2012 Dec;15(8):1127–36.

15. Afzali HHA, Karnon J, Merlin T. Improving the accuracy and comparability of model-based economic evaluations of health technologies for reimbursement decisions: a methodological framework for the development of reference models. Med Decis Mak Int J Soc Med Decis Mak. 2013 Apr;33(3):325–32.

16. Hoogendoorn M, Feenstra TL, Asukai Y, Borg S, Hansen RN, Jansson S-A, et al. Cost-effectiveness models for chronic obstructive pulmonary disease: cross-model comparison of hypothetical treatment scenarios. Value Health J Int Soc Pharmacoeconomics Outcomes Res. 2014 Jul;17(5):525–36.

17. Hoogendoorn M, Rutten-van Mölken MPMH, Hoogenveen RT, van Genugten MLL, Buist AS, Wouters EFM, et al. A dynamic population model of disease progression in COPD. Eur Respir J Off J Eur Soc Clin Respir Physiol. 2005 Aug;26(2):223–33.

18. Birch S, Gafni A. Cost effectiveness/utility analyses. Do current decision rules lead us to where we want to be? J Health Econ. 1992 Oct;11(3):279–96.

19. Roberts M, Russell LB, Paltiel AD, Chambers M, McEwan P, Krahn M, et al. Conceptualizing a model: a report of the ISPOR-SMDM Modeling Good Research Practices Task Force--2. Value Health J Int Soc Pharmacoeconomics Outcomes Res. 2012 Oct;15(6):804–11.

20. Briggs ADM, Wolstenholme J, Blakely T, Scarborough P. Choosing an epidemiological model structure for the economic evaluation of non-communicable disease public health interventions. Popul Health Metr. 2016;14:17.

21. Hennessy DA, Flanagan WM, Tanuseputro P, Bennett C, Tuna M, Kopec J, et al. The Population Health Model (POHEM): an overview of rationale, methods and applications. Popul Health Metr. 2015;13:24.

22. Swanney MP, Ruppel G, Enright PL, Pedersen OF, Crapo RO, Miller MR, et al. Using the lower limit of normal for the FEV1/FVC ratio reduces the misclassification of airway obstruction. Thorax. 2008 Dec;63(12):1046–51.

23. Tan WC, Bourbeau J, Hernandez P, Chapman K, Cowie R, FitzGerald MJ, et al. Canadian prediction equations of spirometric lung function for Caucasian adults 20 to 90 years of age: results from the Canadian Obstructive Lung Disease (COLD) study and the Lung Health Canadian Environment (LHCE) study. Can Respir J J Can Thorac Soc. 2011 Dec;18(6):321–6.

24. Freeman J, Hutchison GB. Prevalence, incidence and duration. Am J Epidemiol. 1980 Nov;112(5):707–23.

25. Bourbeau J, Tan WC, Benedetti A, Aaron SD, Chapman KR, Coxson HO, et al. Canadian Cohort Obstructive Lung Disease (CanCOLD): Fulfilling the Need for Longitudinal Observational Studies in COPD. COPD. 2012 Mar 20;

26. Zafari Z, Sin DD, Postma DS, Löfdahl C-G, Vonk J, Bryan S, et al. Individualized prediction of lung-function decline in chronic obstructive pulmonary disease. CMAJ Can Med Assoc J J Assoc Medicale Can. 2016 Oct 4;188(14):1004–11.

27. Hankinson JL, Odencrantz JR, Fedan KB. Spirometric reference values from a sample of the general U.S. population. Am J Respir Crit Care Med. 1999 Jan;159(1):179–87.

28. Aaron SD, Fergusson D, Marks GB, Suissa S, Vandemheen KL, Doucette S, et al. Counting, analysing and reporting exacerbations of COPD in randomised controlled trials. Thorax. 2008 Feb;63(2):122–8.

29. Hoogendoorn M, Feenstra TL, Hoogenveen RT, Al M, Mölken MR. Association between lung function and exacerbation frequency in patients with COPD. Int J Chron Obstruct Pulmon Dis. 2010 Dec 9;5:435–44.

30. Albert RK, Connett J, Bailey WC, Casaburi R, Cooper JAD, Criner GJ, et al. Azithromycin for prevention of exacerbations of COPD. N Engl J Med. 2011 Aug 25;365(8):689–98.

31. Hurst JR, Vestbo J, Anzueto A, Locantore N, Müllerova H, Tal-Singer R, et al. Susceptibility to exacerbation in chronic obstructive pulmonary disease. N Engl J Med. 2010 Sep 16;363(12):1128–38.

32. Sadatsafavi M, Sin DD, Zafari Z, Criner G, Connett JE, Lazarus S, et al. The Association Between Rate and Severity of Exacerbations in Chronic Obstructive Pulmonary Disease: An Application of a Joint Frailty-Logistic Model. Am J Epidemiol. 2016 Nov 1;184(9):681–9.

33. Statistics canada. Life tables, Canada and provinces and territories [Internet]. Available from: http://www.statcan.gc.ca/pub/84-537-x/4064441-eng.htm

34. Spencer M, Briggs A, Grossman R, Rance L. Development of an economic model to assess the cost effectiveness of treatment interventions for chronic obstructive pulmonary disease. PharmacoEconomics. 2005;23(6):619–37.

35. Mittmann N, Kuramoto L, Seung SJ, Haddon JM, Bradley-Kennedy C, FitzGerald JM. The cost of moderate and severe COPD exacerbations to the Canadian healthcare system. Respir Med. 2008 Mar;102(3):413–21.

36. Statistics Canada. Modgen (Model generator) [Internet]. [cited 2012 Mar 19]. Available from: http://www.statcan.gc.ca/microsimulation/modgen/modgen-eng.htm

37. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2017. Available from: https://www.R-project.org/

38. Calverley PMA, Anderson JA, Celli B, Ferguson GT, Jenkins C, Jones PW, et al. Salmeterol and fluticasone propionate and survival in chronic obstructive pulmonary disease. N Engl J Med. 2007 Feb 22;356(8):775–89.

39. Tashkin DP, Celli B, Senn S, Burkhart D, Kesten S, Menjoge S, et al. A 4-year trial of tiotropium in chronic obstructive pulmonary disease. N Engl J Med. 2008 Oct 9;359(15):1543–54.

40. Asukai Y, Baldwin M, Fonseca T, Gray A, Mungapen L, Price D. Improving clinical reality in chronic obstructive pulmonary disease economic modelling⬚: development and validation of a micro-simulation approach. PharmacoEconomics. 2013 Feb;31(2):151–61.

41. Borg S, Ericsson A, Wedzicha J, Gulsvik A, Lundbäck B, Donaldson GC, et al. A computer simulation model of the natural history and economic impact of chronic obstructive pulmonary disease. Value Health J Int Soc Pharmacoeconomics Outcomes Res. 2004 Apr;7(2):153–67.

42. Briggs AH, Baker T, Risebrough NA, Chambers M, Gonzalez-McQuire S, Ismaila AS, et al. Development of the Galaxy Chronic Obstructive Pulmonary Disease (COPD) Model Using Data from ECLIPSE: Internal Validation of a Linked-Equations Cohort Model. Med Decis Mak Int J Soc Med Decis Mak. 2017 May;37(4):469–80.

43. Menn P, Leidl R, Holle R. A lifetime Markov model for the economic evaluation of chronic obstructive pulmonary disease. PharmacoEconomics. 2012 Sep 1;30(9):825–40.

44. Samyshkin Y, Schlunegger M, Haefliger S, Ledderhose S, Radford M. Cost-effectiveness of roflumilast in combination with bronchodilator therapies in patients with severe and very severe COPD in Switzerland. Int J Chron Obstruct Pulmon Dis. 2013;8:79–87.

45. Celli BR. Counterpoint: Should storefront clinics provide case finding and chronic care for COPD? No. Chest. 2014 Jun;145(6):1193–4.

46. Menzies NA, Soeteman DI, Pandya A, Kim JJ. Bayesian Methods for Calibrating Health Policy Models: A Tutorial. PharmacoEconomics. 2017 Jun;35(6):613–24.

47. Zafari Z, Bryan S, Sin DD, Conte T, Khakban R, Sadatsafavi M. A Systematic Review of Health Economics Simulation Models of Chronic Obstructive Pulmonary Disease. Value Health J Int Soc Pharmacoeconomics Outcomes Res. 2017 Jan;20(1):152–62.

48. Hoogendoorn M, Feenstra TL, Asukai Y, Briggs AH, Borg S, Dal Negro RW, et al. Patient Heterogeneity in Health Economic Decision Models for Chronic Obstructive Pulmonary Disease: Are Current Models Suitable to Evaluate Personalized Medicine? Value Health J Int Soc Pharmacoeconomics Outcomes Res. 2016 Oct;19(6):800–10.

49. Hoogendoorn M, Feenstra TL, Asukai Y, Briggs AH, Hansen RN, Leidl R, et al. External Validation of Health Economic Decision Models for Chronic Obstructive Pulmonary Disease (COPD): Report of the Third COPD Modeling Meeting. Value Health J Int Soc Pharmacoeconomics Outcomes Res. 2017 Mar;20(3):397–403.

50. Vestbo J, Anderson W, Coxson HO, Crim C, Dawber F, Edwards L, et al. Evaluation of COPD Longitudinally to Identify Predictive Surrogate End-points (ECLIPSE). Eur Respir J. 2008 Apr;31(4):869–73.

51. Tabberer M, Gonzalez-McQuire S, Muellerova H, Briggs AH, Rutten-van Mölken MPMH, Chambers M, et al. Development of a Conceptual Model of Disease Progression for Use in Economic Modeling of Chronic Obstructive Pulmonary Disease. Med Decis Mak Int J Soc Med Decis Mak. 2017;37(4):440–52.

52. CEPAC: Cost Effectiveness of Preventing AIDS Complications [Internet]. [cited 2010 Nov 1]. Available from: http://web2.research.partners.org/cepac/

53. Clarke PM, Gray AM, Briggs A, Farmer AJ, Fenn P, Stevens RJ, et al. A model to estimate the lifetime health outcomes of patients with type 2 diabetes: the United Kingdom Prospective Diabetes Study (UKPDS) Outcomes Model (UKPDS no. 68). Diabetologia. 2004 Oct;47(10):1747–59.

54. Schlessinger L, Eddy DM. Archimedes: a new model for simulating health care systems--the mathematical formulation. J Biomed Inform. 2002 Feb;35(1):37–50.

55. Weinstein MC, Coxson PG, Williams LW, Pass TM, Stason WB, Goldman L. Forecasting coronary heart disease incidence, mortality, and cost: the Coronary Heart Disease Policy Model. Am J Public Health. 1987 Nov;77(11):1417–26.

56. Bibbins-Domingo K, Chertow GM, Coxson PG, Moran A, Lightwood JM, Pletcher MJ, et al. Projected effect of dietary salt reductions on future cardiovascular disease. N Engl J Med. 2010 Feb 18;362(7):590–9.

57. Goossens LMA, Nivens MC, Sachs P, Monz BU, Rutten-van Mölken MPMH. Is the EQ-5D responsive to recovery from a moderate COPD exacerbation? Respir Med. 2011 Aug;105(8):1195–202.