Screening Mycobacterium tuberculosis secreted proteins identifies Mpt64 as eukaryotic

membrane-binding virulence factor

Chelsea E. Stamm[a,b], Breanna L. Pasko[a,b], Sujittra Chaisavaneeyakorn[a,c], Luis H.

Franco[a,d]*, Vidhya R. Nair[a], Bethany A. Weigele[b]*, Neal M. Alto[b], Michael U. Shiloh[a,b]#.

[a]Department of Internal Medicine, University of Texas Southwestern Medical Center,

Dallas, Texas, USA

[b]Department of Microbiology, University of Texas Southwestern Medical Center, Dallas,

Texas, USA.

[c]Department of Pediatrics, University of Texas Southwestern Medical Center, Dallas,

Texas, USA.

[d]Center for Autophagy Research, University of Texas Southwestern Medical Center,

Dallas, Texas USA

Running title: Mtb Mp64 is a membrane-binding virulence factor

#Address correspondence to Michael U. Shiloh, michael.shiloh@utsouthwestern.edu

*Present address: Luis Franco, Federal University of Minas Gerais, Belo Horizonte,

Brazil; Bethany Weigele, Arizona State University, Scottsdale, Arizona USA.

Abstract word count: 203

Text word count: 7,901

Abstract

Mycobacterium tuberculosis (Mtb), the causative agent of tuberculosis, is one of the

most successful human pathogens. One reason for its success is that Mtb can reside

within host macrophages, a cell type that normally functions to phagocytose and destroy

infectious bacteria. However, Mtb is able to evade macrophage defenses in order to

survive for prolonged periods of time. Many intracellular pathogens secret virulence

factors targeting host membranes and organelles to remodel their intracellular

environmental niche. We hypothesized that Mtb exported proteins that target host

membranes are vital for Mtb to adapt to and manipulate the host environment for

survival. Thus, we characterized 200 exported proteins from Mtb for their ability to

associate with eukaryotic membranes using a unique temperature sensitive yeast

screen and to manipulate host trafficking pathways using a modified inducible secretion

screen. We identified five Mtb exported proteins that both associated with eukaryotic

membranes and altered the host secretory pathway**.** One of these secreted proteins,

Mpt64, localized to the endoplasmic reticulum during Mtb infection of murine and human

macrophages and was necessary for Mtb survival in primary human macrophages.

These data highlight the importance of exported proteins in Mtb pathogenesis and

provide a basis for further investigation into their molecular mechanisms.

Importance

Advances have been made to identify exported proteins of *Mycobacterium tuberculosis*

during animal infections. These data, combined with transposon screens identifying

genes important for *M. tuberculosis* virulence, have generated a vast resource of

potential *M. tuberculosis* virulence proteins. However, the function of many of these

proteins in *M. tuberculosis* pathogenesis remains elusive. We have integrated three cell

biological screens to characterize nearly 200 *M. tuberculosis* exported proteins for

eukaryotic membrane binding, host subcellular localization and interactions with host

vesicular trafficking. In addition, we observed the localization of one exported protein,

Mpt64, during *M. tuberculosis* infection of macrophages. Interestingly, although Mpt64 is

exported by the Sec pathway, its delivery into host cells was dependent upon the action

of the Type VII Secretion System. Finally, we observed that Mpt64 contributes to the

virulence of *M. tuberculosis* during infection of primary human macrophages.

Introduction

Tuberculosis caused by *Mycobacterium tuberculosis* (Mtb) is a persistent, global

epidemic. While the number of deaths due to Mtb fell below 2 million in 2015, there

were over 9 million new cases (1) and the incidence of multidrug-resistant Mtb is

increasing (1), highlighting the need for new anti-tuberculosis therapies. In addition, the

only currently available vaccine, *Mycobacterium bovis* Bacille-Calmette-Guérin (BCG),

is ineffective in preventing pulmonary tuberculosis infection (2). Thus, understanding the

intracellular survival mechanisms employed by Mtb is vital to developing new anti-

tuberculosis treatments and vaccines.

Macrophages, phagocytic innate immune cells that are generally competent for bacterial

killing, represent the primary intracellular niche for Mtb. Some of the antimicrobial

mechanisms utilized by macrophages include acidification of the phagosome,

production of reactive oxygen and nitrogen species, fusion of lysosomes to bacterial

containing phagosomes and autophagy (3-6). However, despite these robust defenses,

Mtb survives inside macrophages during its infectious life cycle. To facilitate its survival

Mtb has evolved to resist macrophage defenses, either by directly protecting the

bacterial cell from damage (7-9) or by modulating the macrophage's ability to shuttle the

bacteria through the traditional phagolysosomal maturation process (10). In that way,

Mtb prevents its intracellular compartment from acidifying (11) and fusing (12) with the

destructive lysosome. Genetic studies have identified several Mtb proteins important for

remodeling host membrane trafficking (13-15). For example, Mtb *Rv3310* encodes

SapM, a secreted acid phosphatase (16) that converts phosphatidylinositol 3-phosphate

(PI3P) to phosphatidylinositol. Loss of PI3P from the phagosome membrane is sufficient

to prevent fusion of phagosomes with late endosomes (17, 18). Importantly, many

genes reported to be important for Mtb survival inside macrophages remain

uncharacterized (13, 14, 19) and the manipulation of the host cell by Mtb remains poorly

understood.

The problem of intracellular survival faced by Mtb is also shared by other bacterial

pathogens, and many of these organisms utilize specialized secretion systems to

deliver molecules into the host cell to establish a unique intracellular niche (20). For

example, some Gram-negative pathogens use needle-like machines that span the

bacterial and host cell membranes to inject protein cargo into the host (21-24). Encoded

by a Type III Secretion System (TTSS), the needle-like machine shares its structure

with the bacterial flagellum and is expressed by a number of intracellular pathogens

including *Salmonella enterica* subsp. *enterica* serovar Typhimurium, *Shigella flexneri*,

and *Chlamydia trachomatis* (21, 22). Another specialized secretion machine called a

Type IV Secretion System is found in Gram-positive and Gram-negative bacteria and

can used for translocation of nucleic acid through conjugation (23) while in many human

and plant pathogens such as *Legionella pneumophila, Coxiella burnetii* and

Agrobacterium tumefaciens, it also transports effector proteins that promote bacterial

survival (25, 26). The Type VI Secretion System is structurally analogous to the

bacteriophage tail spike and is expressed by pathogenic bacteria including *Francisella*

tularensis, Pseudomonas aeruginosa and Vibrio cholera (24, 27, 28). Finally, Mtb

encodes multiple Type VII secretion systems, discussed below, that are important in

pathogenesis (29, 30). However, the identity and functions of the Type VII dependent

secreted proteins remain unknown. Despite structural differences, a common theme is

that these elaborate secretion systems aid in pathogenesis by delivering virulence

proteins called "effectors" to the host cell.

Effectors are proteins that promote bacterial survival by manipulating vital cellular

processes including signal transduction, vesicular trafficking and the cytoskeleton (31-

33). Like the secretion systems themselves, the repertoire of effectors expressed by

each pathogen can differ, adapted specifically for each unique life cycle. However, a

major common target for effectors are host membranes. For example, SifA from *S.*

typhimurium is prenylated inside the host cell and localizes to the Salmonella containing

vacuole (SCV) (34). SifA recruits lysosomes to maintain SCV membrane integrity and

its membrane interaction is vital to Salmonella pathogenicity (34, 35). The *Legionella*

pneumophila Type IV secreted effector SidM is anchored to the Legionella containing

vacuole and disrupts host vesicle trafficking by sequestering and modifying Rab1 (36).

Some effectors can also function by directly modifying membranes such as IpgD from

S. flexneri that hydrolyzes phosphatidylinositol 4,5-bisphosphate (PI(4,5)P$_2$) to

phosphatidylinositol 5-monophosphate (PI5P) leading to membrane blebbing and

bacterial uptake into host cells (37). Thus, both host membranes themselves and

membrane-dependent processes represent valuable targets for bacterial effectors (31,

33, 38) as we recently showed for a variety of bacterial pathogens (39).

The virulence functions of most Mtb secreted proteins are poorly understood. Although

Mtb does not encode a T3SS, it does contain the conserved general secretion systems

Sec and Tat (40), as well as multiple Type VII (also called ESX) secretion systems (30).

The ESX-1 system is required for Mtb virulence in macrophages and mouse models of

infection (29) through modulation of immune functions such as induction of Type I

interferon via the secreted effectors EsxA and EsxB (41, 42). Similarly, EsxH, a substrate of the ESX-3 system, interacts with a component of the host endosomal sorting complex required for transport (ESCRT) machinery, leading to decreased co-localization of Mtb with lysosomes (43) and inhibition of antigen presentation in infected macrophages (44). Substrates of the accessory SecA2 system such as the protein kinase PknG and the esterase LipO are also important for Mtb virulence by contributing to phagosome maturation arrest (PMA) (13, 18, 45, 46). Thus, because membrane processes are high value targets of many bacterial effectors (39, 47), and Mtb has a large repertoire of exported proteins of unknown function (48-52), we hypothesized that some of the Mtb exported proteins are membrane-binding effectors with virulence activities.

To test our hypothesis we generated a library of 200 exported proteins from Mtb, tested whether they individually bound yeast membranes in a life or death assay, and characterized their ability to alter host protein secretion in an inducible secretion assay. We also determined the subcellular localization of each membrane-binding protein using fluorescence microscopy. By combining data from the cell biological screens, we identified five Mtb exported proteins that localized to eukaryotic membranes and inhibited protein transport through the host secretory pathway. One protein, Mpt64 (Rv1980c), localized to the ER during both heterologous expression in HeLa cells and Mtb infection of macrophages. Though Mpt64 is a Sec substrate, its access to the macrophage cytoplasm was dependent on the ESX-1 secretion system. Finally, bacteria lacking Mpt64 were attenuated for growth in primary human macrophages, demonstrating an important virulence activity for Mpt64.

Results

Categorization of putative effector-like proteins from Mtb

Through the analysis of published datasets we identified Mtb proteins that may function as secreted effectors (Supplemental Table 1). For clarity, we define these proteins as Mycobacterial exported proteins (MEPs), as this encompasses proteins that are secreted as well as those delivered to the mycobacterial surface, which can still modulate interactions with the host (53, 54). We used the following criteria to assemble a library of MEPs: 1) Mtb secreted proteins identified via proteomic approaches (49-52, 55, 56), 2) Mtb proteins known to be involved in manipulation of host vesicular trafficking pathways, such as ones that induce mammalian cell entry (MCE) (57-59) or phagosome maturation arrest (13, 14, 60), 3) a subset of PE/PPE proteins and proteins related to those encoded by Type VII (ESX-1) loci (61-63), and 4) proteins involved in virulence, ranging from defined to unknown functions (19, 54, 64-66). We then used Gateway recombination cloning to subclone MEPs from the freely available Mtb ORFome Gateway compatible library (BEI) into destination vectors for a variety of subsequent assays. The comprehensive list of MEPs is shown in Supplemental Table 1.

Mtb encodes exported proteins that interact with eukaryotic membranes

To identify membrane binding Mtb proteins, we used a system that leverages the signal transduction of the essential yeast GTPase Ras to promote growth and division (67). Ras is lipidated at a unique sequence called the CaaX box that promotes its localization to the plasma membrane, where it can be activated by Cdc25, a guanine nucleotide exchange factor (67, 68). In a yeast strain with a temperature sensitive *CDC25* allele, yeast can only grow at the permissive temperature (25°C) but not the restrictive

temperature (37°C) because Ras activation requires interaction with Cdc25.

Heterologous expression of a non-lipidated, constitutively-active Ras whose activity is

independent of Cdc25 (Ras[mut]) can rescue yeast growth at the restrictive temperature

when Ras is recruited to intracellular membranes by fusion to a membrane binding

protein (Fig. 1A). This system has been used to successfully identify membrane binding

effectors from Gram negative pathogens (39). To identify membrane-localizing proteins

from Mtb, we subcloned MEPs into a destination vector for yeast expression that

generates an in-frame fusion of the MEP to Ras[mut]. We transformed *S. cerevisiae*

cdc25[ts] (39, 67) individually with each of the 200 MEP fused to Ras[mut] and incubated

them at both permissive and restrictive temperatures (Figure 1B). We identified 52 Mtb

proteins that rescued *S. cerevisiae cdc25[ts]* growth at the restrictive temperature (Fig.

1B, C).

We confirmed expression of the Ras[mut]-MEP fusion proteins by Western blotting

(Figure 1D). In addition, we determined the membrane localization of each MEP by

fluorescence microscopy of GFP-MEP fusion proteins in yeast (Fig. 1E). It has been

established that Ras can function from membranes other than the plasma membrane

(69, 70) and Ras[mut] maintains this function (39). Thus, using fluorescence microscopy

we observed GFP-MEP fusion proteins localizing to distinct subcellular compartments

including vacuoles, ER and plasma membrane (Figure 1E). Together these results

show that 25% of the MEPs tested could associate with the membranes of a variety of

organelles in *S. cerevisiae*.

Subcellular localization of membrane-localizing MEPs

While many cellular processes are conserved in eukaryotes, humans represent

the primary natural host for Mtb. Therefore, to confirm that MEPs that rescued *S.*

cerevisiae cdc25^{ts} growth at 37°C also bound mammalian membranes and to determine

their subcellular localization in human cells, we transiently transfected HeLa cells with

vectors for constitutive expression of GFP-MEP fusion proteins and then used

fluorescence microscopy with co-localization markers to identify the specific membrane

to which each MEP localized (Figure 2A). We identified GFP-MEP fusion proteins that

localized to a variety of subcellular compartments including the ER, Golgi, mitochondria

and peroxisomes (Figure 2A and 2B). The largest proportion of the GFP-MEP fusion

proteins expressed in human cells co-localized with the ER marker calreticulin (Figure

2B). In yeast, we observed a similar number of GFP-Mtb fusion proteins which localized

to compact, punctate structures (data not shown). Although there was only moderate

overlap in the subcellular localization identified between yeast and HeLa cells (Figure

2B), we were able to verify that the proteins identified by the Ras rescue assay are

localized to membranes in human cells.

A subset of mycobacterial exported proteins alter the host secretory pathway

Membrane-bound cargo is transported within the host cell and to the extracellular space

in dynamic vesicular trafficking pathways. The membrane-bound and soluble proteins

important for these processes are frequent targets of bacterial effectors (47, 71).

Indeed, Mtb is known to target and manipulate trafficking pathways through

incompletely understood mechanisms (43, 72, 73). To determine if Mtb proteins can

broadly affect the host general secretion pathway as an indicator of interaction with

membranes and intracellular vesicular trafficking, we took advantage of the reverse

dimerization system (74). In this system, a secreted protein of interest is sequestered in

the ER by fusion to a conditional aggregation domain (CAD). Addition of a solubilization

molecule that disrupts the CAD then frees the fusion protein for trafficking and secretion

into the extracellular space. We used a fusion of human growth hormone (hGH) to the

CAD domain of the ligand-reversible crosslinking protein, FKBP F36M. Thus, hGH can

be quantified in cell supernatants by ELISA after addition of the small molecular, D/D

Solubilizer (Figure 3A) (74, 75). We transfected HeLa cells expressing hGH-CAD with

each MEP individually, a negative control protein (GFP), or an enterohemorrhagic

Escherichia coli effector (EspG) that inhibits secretion by promoting the tethering of

vesicles to the Golgi apparatus (75, 76). When we treated transfected cells with D/D

Solubilizer, we observed a spectrum of effects on the secretion of hGH including

increased, decreased and normal hGH secretion (Figure 3B and Supplemental Table

1). Using a cut-off of normalized hGH secretion below 0.25 or above 1.75, we identified

10 proteins that decreased hGH secretion and 10 proteins that increased hGH secretion

as compared to the GFP control (Figure 3C). We next compared the MEPs that altered

host secretion to those that bound eukaryotic membranes, and identified five proteins

with overlapping activities: Rv0594, Rv1646, Rv1810, Rv1980c and Rv2075 (Figure

3D). During expression in HeLa cells, all but one protein localized to the ER and all five

proteins reduced hGH secretion (Figure 3E).

Mpt64 N-terminus is important for membrane binding and secretion inhibition

We focused on the protein Rv1980c, also known as Mpt64, as it is a secreted protein

that is highly antigenic during human tuberculosis infection (77, 78). Furthermore, Mpt64

is a component of the region of difference 2 (RD2) locus, one of the genomic regions

deleted during attenuation of the *M. bovis* BCG vaccine strain (79). Loss of RD2 from

Mtb attenuates its virulence, and complementation with a three gene cluster that

includes Mpt64 can partially restore virulence (80).

Mpt64 is a 25kDa protein with a predicted signal peptidase 1 cleavage site

between amino acids 23 and 24, such that the mature, secreted form of the protein

starts at amino acid 24 (50, 51, 81). While the solution structure of Mpt64 was

previously solved (82), the structure does not align to a known catalytic domain but

does contain a domain of unknown function (DUF3298). This domain is also present in

the lysozyme-binding anti sigma factor RsiV (83). Despite structural homology between

Mpt64 and RsiV (Supplemental Figure 1A) (84), there is little primary sequence

homology. To determine if Mpt64 binds lysoszyme, we purified recombinant Mpt64 from

E. coli and tested binding to human or hen egg white lysozyme in an *in vitro* pull down

assay (83). Using this assay we were unable to demonstrate lysozyme binding by

Mpt64 (Supplemental Figure 1B). We next used the solution structure to guide

truncation analysis of Mpt64 in order to identify the membrane binding sequences of

Mpt64 (Figure 4A and Figure 4B). *S. cerevisiae cdc25^{ts}* expressing a fusion of Ras^{mut}

with either full length Mpt64, mature Mpt64 lacking its predicted signal peptide or the N-

terminal half of Mpt64 also lacking the signal peptide (Mpt64_24-143) were able to grow

at 37°C, whereas *S. cerevisiae cdc25^{ts}* expressing Ras^{mut} fused to the C-terminal half of

Mpt64 (Mpt64_144-228) could not (Figure 4C). We detected expression of Ras^{mut}

fusions of full length Mpt64 and mature Mpt64 by Western blot. In contrast, we could not

detect Ras^{mut} fusions of Mpt64_24-143 or Mpt64_144-228 despite the fact that the

Mpt64_24-143 fusion rescued yeast growth, suggesting that expression of Mpt64_24-

143 below the limit of detection by Western blot was still sufficient to rescue yeast

growth (Figure 4D). However, whether the C-terminal domain plays a role in membrane

binding could not be determined definitively because we were unable to demonstrate

stable fusion protein expression. To test if Mpt64 could interact with lipids directly, we

expressed and purified recombinant Mpt64 and Mpt64 variants from *E. coli* and tested

their ability to bind unique lipid species i*n vitro* using membranes spotted with lipids.

Recombinant   Mpt64_24-228   bound   phosphatidylinositol   4-phosphate   (PI4P),

phosphatidylinositol   5-phosphate   (PI5P),   phosphatidylinositol   4,5-bisphosphate

[PI(4,5)P$_2$] and phosphatidylinositol (3,4,5)-trisphosphate [PI(3,4,5)P$_3$] on PIP strips

membranes (Figure 4E). Similarly, recombinant Mpt64_24-143, the N-terminal portion of

the protein, also bound PI4P and PI5P with additional binding to PI3P, PI(3,4)P$_2$ and

phosphatidylserine (Figure 4E). However, interaction with PI(4,5)P$_2$ and PI(3,4,5)P$_3$ was

weak, suggesting that the C-terminal region of Mpt64 modifies its interactions with host

phospholipids.

Finally, we sought to determine if the N-terminal portion of Mpt64 was also

sufficient to inhibit hGH secretion using the hGH-CAD assay. We co-transfected hGH-

CAD expressing HeLa cells with the same Mpt64 truncations and determined their

ability to inhibit hGH secretion in the presence of drug. Similar to the Ras rescue assay,

full length, mature and Mpt64_24-143 inhibited hGH secretion compared to the GFP

control. In contrast, co-transfection of Mpt64_144-228 with hGH-CAD had no effect on

its secretion (Figure 4F). These data suggest that the ability of Mpt64 to bind

membranes and to inhibit host secretion *in vitro* is dependent on the N-terminus of the

protein.

Mpt64 ER localization depends on its N-terminus

As full-length Mpt64 localized to the ER in yeast and HeLa cells (Fig. 1,3) we next tested the impact of Mpt64 truncations on ER localization. We first determined the phenotypic localization of Mpt64 truncations expressed as GFP fusions in yeast using fluorescence microscopy. Mpt64_1-228 and Mpt64_24-228 localized in a ring indicative of the ER (85, 86)(Figure 5A). In contrast Mpt64_144-228, which did not rescue yeast growth in the Ras Rescue assay (Figure 4C), was diffuse throughout the yeast cell (Figure 5A). Interestingly, Mpt64_24-143 localized to bright puncta within the cells (Figure 5A). To confirm the N-terminal dependence of Mpt64 localization, we transfected HeLa cells with GFP fusions to each Mpt64 truncation and assayed for co-localization with calreticulin through immunofluorescence microscopy. While full length Mpt64, mature Mp64 and Mpt64_24-143 co-localized with calreticulin, Mpt64_144-228 did not (Figure 5B). These results demonstrate that Mpt64 localizes to the ER during exogenous expression in both yeast and mammalian cells and the N-terminal 143 amino acids are sufficient for Mpt64 to localize to membranes.

Secreted Mpt64 localizes to the ER during infection

Although we observed Mpt64 localization to the ER in yeast (Figure 1E and Figure 5A) and HeLa cells (Figure 5B), we wanted to determine if endogenous, untagged Mpt64 localizes to the ER during an Mtb infection of macrophages. To that end, we infected mouse RAW267.4 macrophages with mCherry-labeled Mtb at an MOI of 20:1 and fixed cells at various time points after infection. We then used a rabbit polyclonal antibody developed against recombinant, mature Mpt64 protein to track Mpt64 secretion from Mtb into macrophages using immunofluorescence microscopy. Of

note, this antibody was generated without complete Freund's adjuvant in order to avoid

any cross-reactivity against Mtb antigens generated by the use of this adjuvant. As little

as four hours after infection, endogenous Mpt64 was detected in both the cytoplasm

and ER of host cells (Figure 6A, upper panels). When we performed the same

experiment with MtbΔeccD1, a strain that lacks the Type VII secretion system secretion

pore that cannot secret ESAT-6 (29, 87, 88) and does not result in communication

between the phagosome and cytoplasm (41, 42, 89), Mpt64 appeared to be secreted

but trapped adjacent to the bacteria (Figure 6A, lower panels), suggesting that it could

not escape the phagosome. Importantly, Mpt64 was detected in the culture filtrate

prepared from MtbΔeccD1 (Supplemental Figure 2A). Thus, although Mpt64 is likely

secreted from Mtb by the canonical Sec-dependent pathway, its access to the

macrophage cytoplasm and other targets in the cell was dependent on the Type VII

secretion system.

In order to better understand the role of Mpt64 in Mtb virulence, we used

mycobacteriophage (90-92) to introduce the hygromycin resistance cassette into the

Mpt64 gene to create an in-frame deletion (Figure 6B). We confirmed disruption of

Mpt64 by PCR (Supplemental Figure 2B) and loss of Mpt64 by the absence of protein

on Western blot (Figure 6C). We then complemented MtbΔMpt64 with either full length

Mpt64 (MtbΔMpt64::Mpt64) or Mpt64 lacking its signal peptide (MtbΔMpt64::Mpt64-NS)

under the control of the constitutive mycobacterial strong promoter (93). Both

complemented strains expressed Mpt64 but only full-length Mpt64 (MtbΔMpt64::Mpt64)

could be detected in the supernatant of cultures, confirming that deletion of the signal

peptide inhibits Mpt64 secretion from Mtb (Figure 6C). Furthermore, the

MtbΔMpt64::Mpt64 strain had modestly higher expression of Mpt64 compared to wild-

type Mtb by western blot, consistent with our use of a strong constitutive promoter for

complementation. All four strains grew equally under axenic growth conditions

(Supplemental Figure 2C), and we confirmed that both Mtb and MtbΔMpt64 produced

phthiocerol dimycocerosate by mass spectrometry (Supplemental Figure 2D).

To test if secreted Mpt64 interacts with the ER during infection, we assessed its

colocalization with calreticulin in RAW267.4 cells using confocal immunofluorescence

microscopy. When we infected RAW267.4 macrophages, the Mpt64 signal in Mtb

infected macrophages co-localized with calreticulin, confirming the subcellular

localization of Mpt64 secreted during infection (Figure 6D,E and Supplemental Figure

3). However this co-localization was lost in cells infected with MtbΔMpt64::Mpt64-NS

bacteria (Figure 6D,F and Supplemental Figure 3). As a control for antibody specificity,

no Mpt64 was detected in macrophages infected with MtbΔMpt64 mutant bacteria

(Figure 6D and Supplemental Figure 3). From these data, we can confirm that the signal

peptide of Mpt64 is sufficient for the protein's secretion in vivo and is required (with

concerted action of the Type VII secretion system) for Mpt64 to interact with the ER

during infection.

Mpt64 contributes to early Mtb growth after aerosol infection of mice.

Because Mpt64 is part of the Mtb RD2 locus that partially accounts for the attenuation of

Mtb (80), and our data indicating that Mpt64 may function as a secreted effector, we

investigated the role of Mpt64 in Mtb virulence in a murine model of infection. We

infected BALB/c mice via aerosol with a low-dose of bacteria (~100 CFU Mtb) and

collected lungs at various time points to determine CFU and histopathology. We

compared the infections of four strains: wild type Mtb, MtbΔMpt64, MtbΔMpt64::Mpt64

and MtbΔMpt64::Mpt64-NS (described in Figure 6). While all mice received equal

numbers of bacteria between the four strains at day 0 (data not shown), there were

fewer Mtb isolated from lungs of mice infected with the MtbΔMpt64 mutant compared to

wild type (mean CFU wild type Mtb $2.7x10^6$ vs MtbΔMpt64 $1.7x10^6$, p=0.07) at 21 days

post infection (dpi) but no statistically significant difference by 42dpi (mean CFU wild

type Mtb $5.0x10^5$ vs MtbΔMpt64 $3.4x10^5$). However, by 42dpi there was a statistically

significant decrease in the CFU isolated from lungs of mice infected with

MtbΔMpt64::Mpt64-NS (Figure 7A). At these time points, we observed a reduction in the

area of inflammation in hematoxylin and eosin (H&E) stained lungs of mice infected with

MtbΔMpt64::Mpt64-NS compared to wild type. (Figure 7B and 7D). Despite modest

reductions in CFU in mutant bacteria, there were no differences in mouse survival

comparing the four strains (Figure 7C). Thus, although deletion of *Mpt64* entirely

(MtbΔMpt64) and disruption of Mpt64 secretion (MtbΔMpt64::Mpt64-NS) resulted in a

modest decrease in bacterial growth in the lungs of mice, loss of Mpt64 was not

sufficient to explain the attenuation of the RD2 mutant (80).

Mtb survival in human macrophages requires Mpt64

Next, we assessed whether the localization of Mpt64 in human cells is similar to

that in murine macrophages. To that end, we infected primary human monocyte-derived

macrophages with mCherry-expressing WT Mtb or MtbΔeccD1 and stained for Mpt64.

Consistent with our data in RAW267.4 cells (Figure 6A) the secretion of Mpt64 to extra-

phagosomal sites in primary human macrophages was dependent on the Type VII

Secretion System (Figure 8A). Then we infected primary human macrophages with WT

Mtb, MtbΔMpt64, MtbΔMpt64::Mpt64 or MtbΔMpt64::Mpt64-NS and determined the co-

localization of Mpt64 with calreticulin by fluorescence microscopy (Figure 8B and

Supplemental Figure 4). At 4hpi, we detected co-localization of Mpt64 with calreticulin in

cells infected with WT Mtb and MtbΔMpt64::Mpt64 (top panels) but not in cells infected

with MtbΔMpt64 or MtbΔMpt64::Mpt64-NS (bottom panels, Figure 8B).

To better understand the contribution of Mpt64 in the context of human Mtb

infection, we determined the growth of wild type, MtbΔMpt64 or MtbΔMpt64::Mpt64 in

primary monocyte-derived human macrophages. We recovered CFU from cells directly

after infection (Day 0) and one and three days post infection. While there was no

difference in the CFU isolated from cells infected with each strain after one day of

infection, the MtbΔMpt64 mutant was attenuated for growth three days post infection

compared to wild type (Figure 8C). This growth defect was rescued in the

complemented strain. Together these data indicate a role for Mpt64 in the virulence of

Mtb during acute infection of human macrophages.

Discussion

Numerous efforts have been undertaken to identify Mtb exported proteins, from

lipoproteins that are incorporated into the cell wall, to virulence factors that reach the

extracellular environment such as ESAT-6 (48-50). However, little is known about the

function of this "secretome" as a whole. Here we took a systematic approach towards

characterizing host-dependent interactions of a collated list of putative effector

molecules. We created a library of 200 putative mycobacterial exported proteins and

then through a series of cell biological screens characterized these MEPs for their ability

to bind eukaryotic membranes, their subcellular localization and their ability to modulate

secretion of a model substrate. In addition, we demonstrate that one secreted protein,

Mpt64, localized to the ER during infection and was important for virulence of Mtb in

human macrophages. The cohort of 200 MEPs we generated was large but not

necessarily exhaustive (Supplemental Table 1). For example, the 76 PE/PPE genes we

included represent less than half (45%) of the total number of PE/PPE genes in the Mtb

genome (94). In addition, a recent technology called EXIT identified 593 Mtb proteins

secreted during intravenous infection of mice including 38 proteins that are significantly

enriched only during *in vivo* infection as compared to growth on 7H10 agar, suggesting

a virulence function for these proteins (95). Of the 200 MEPs we characterized, 51

overlap with those identified by EXIT and of the 51 overlapping proteins, 25 are

membrane associated in our study. This emphasizes that host membranes can be

targets of Mtb virulence proteins.

We found 52 Mtb proteins that associated with eukaryotic membranes,

representing nearly 25% of the total screened. When the membrane association of type

III and type IV effectors from several Gram negative pathogens was explored, about

30% of effectors screened also associated with eukaryotic membranes (39). While our

data are in agreement with this value, pathogens that replicate intracellularly in vacuoles

had even higher numbers of membrane-associated effectors (39). This suggests that

there may be additional secreted virulence proteins from Mtb that associate with the

host membranes than our screen identified. Indeed, while we corroborated previously

known membrane-interacting proteins such as the SecA2-secreted PI3P phosphatase

SapM (17, 18), the Rac1-binding protein Ndk (96) and the cholesterol-binding Mce4A

(97), we failed to identify others such as LipY which hydrolyzes extracellular lipids (98)

and the ESX1 substrate ESAT-6 which interacts with the phagosomal membrane (99,

100).

GFP-Mpt64 localized to the ER in both yeast and mammalian cells. Additionally,

endogenous Mpt64 localized to the ER during Mtb infection of macrophages,

suggesting that the observed localization of Mpt64 is not an artifact of heterologous

over-expression. Mpt64 did not co-localize with the ER after infection with a Type VII

secretion system mutant underscoring the importance of the membrane disrupting

properties of ESAT-6 in establishing communication with the host cell (99, 100). This

ESX-1 dependent mechanism of cytoplasmic access is similar to the route taken by the

autotransporter-like protein tuberculosis necrotizing toxin (TNT) (101, 102). Thus, our

data strengthen the argument that the Type VII secretion system facilitates access of

non-ESX-1 substrates beyond the phagosome and into the host cell.

We identified PI4P and PI5P as primary monophosphotidylinositol targets of

recombinant Mpt64, in addition to the di- and tri-phosphotidylinositols $PI(4,5)P_2$ and

$PI(3,4,5)P_3$. While PI4P is thought to be enriched in the Golgi (103), it also has an

established role in mediating protein trafficking from ER exit sites (104, 105). Thus, the

ability of Mpt64 to block secretion may stem from its subcellular localization at the ER,

interaction with PI4P, and interference with ER to Golgi trafficking. Less is known about

PI5P as its basal level is only about 1% of PI4P (106). However, PI5P is increased

during bacterial infection and other stresses, and can be found throughout the cell,

including   the   ER   (106).   Likewise,   while $PI(4,5)P_2$,   the   most   abundant

phosphotidylinositol, is distributed predominantly at the plasma membrane, it can also

be found within the cell at the Golgi and ER (107), and $PI(4,5)P_2$ serves as the

precursor for cellular production of PI(3,4,5)P$_3$ (108). Taken together, while Mpt64 binds

multiple phosphotidylinositols in vitro, the relative contributions of binding to individual

molecules and the role of such binding in mediating the host secretion blockade of

Mpt64 remain unknown.

Disruption of the RD2 locus in Mtb H37Rv leads to decreased bacterial burdens

in the lungs and spleen of aerosol-infected mice at 3 weeks after infection (80). As

Mpt64 is within the RD2 locus, we hypothesized that the single, in-frame deletion of

Mpt64 might explain the attenuation phenotype of the RD2 mutant. In line with this

prediction, we observed decreased bacterial burdens of MtbΔMpt64 compared to WT

Mtb in the lungs of mice at 3 weeks post-infection. Although this decrease was not

statistically significant and was not associated with a survival defect, it does suggest

Mpt64 contributes to the virulence of the RD2 region. Other genes located in the RD2

locus that were not complemented in the RD2 survival study (80) such PE_PGRS35

(Rv1983) and cfp21 (Rv1984) may also contribute alongside Mpt64 to the virulence

defect observed in RD2 deletions. Furthermore, it is possible that one or more of the

other Mtb secreted proteins we identified, including the 27 proteins that also localize to

the ER are able to perform a redundant function to that of Mpt64 during an animal

infection. In a similar vein, whereas *L. pneumophila* encodes over 300 effectors,

individual *L. pneumophila* effector deletion mutants are not defective for growth in cells

or mice (109, 110). Thus, future work disrupting multiple Mtb MEPs simultaneously will

help address the issue of redundancy.

When we infected mice with MtbΔMpt64::Mpt64-NS, a strain of Mtb that still

expresses Mpt64 but cannot secrete it into the host cell, we recovered significantly

fewer CFU compared to WT from the lungs of MtbΔMpt64::Mpt64-NS infected mice. We

hypothesize that this strain suffers from two detrimental consequences. First, blocking

Mpt64 secretion prevents it from exerting its virulence function in the host. Second, non-

secreted Mpt64 can still be cross-presented to the adaptive immune system (111), thus

leading to a cell mediated immune response against Mpt64. This observation is

consistent with data that both human patients with active tuberculosis and their PPD

positive contacts have T-cell responses to Mpt64 (112) and T-cell reactive Mpt64

epitopes have been mapped (113). Furthermore, Mpt64 staining is observed in

granulomas of infected individuals (114, 115). Thus, Mpt64 is highly immunogenic

during human infection with Mtb and suggests an evolutionary tradeoff between the

virulence function of Mpt64 and its antigenicity. In fact, when we explored the

importance of Mpt64 in human disease, we observed that Mpt64 secreted from wild

type bacteria localized to the ER of infected human monocyte-derived macrophages

and its presence enhanced the survival of Mtb within primary human macrophages.

Together these findings suggest that Mpt64 contributes to the virulence of Mtb in human

disease.

Materials and Methods

Bacterial Strains and Growth Conditions

M. tuberculosis Erdman and mutants were grown in Middlebrook 7H9 broth or on Middlebrook 7H11 agar (Difco) supplemented with 10% oleic acid-albumin-dextrose-catalase (OADC, Remel). Liquid medium was also supplemented with 0.05% Tween 80.

Yeast Strains and Assays

The *Saccharomyces cerevisiae* strain INVSc1 (Invitrogen) was grown at 30°C in histidine drop out media (SD/-HIS) or agar plates (Clontech). The construction of cdc25^{ts} was previously described (68). cdc25^{ts} was grown at 25°C in leucine drop out media or agar plates (SD/-LEU) (Clontech).

The cdc25^{ts} strain and INVSc1 strain were transformed using a lithium acetate (LiAc) protocol. Yeast were grown to high density overnight at the appropriate temperature, shaking. The cultures were diluted to an OD$_{600}$=0.2 in 50mL YPD and allowed to reach mid-log phase. Cells were washed, resuspended in 0.1M LiAc and incubated 10 minutes at room temperature. The sample DNA was mixed with an equal volume of pre-boiled Yeastmaker Carrier DNA (Clontech). To the DNA was added 100uL yeast and 500uL of a solution of LiAc+PEG (40% PEG w/v, 0.1M LiAc). This solution was incubated 30 minutes at 25°C (cdc25^{ts}) or 30°C (INVSc1) with agitation every 10 minutes. DMSO was added and the cells were heat shocked at 42°C for 15 minutes. The cells were pelleted, washed in TE and resuspended in 500uL TE. The transformed cells were plated on selective agar plates and incubated at the appropriate temperature for 2-4 days.

To perform the Ras rescue assay, 3-4 fresh colonies were combined in 30µL

SD/-LEU and 3 µl was spotted onto duplicate plates that were subsequently incubated

at either 25°C or at 37°C for 2 days.

INVSc1 yeast were transformed with a galactose-inducible vector

(p413GALGFP) containing GFP-Mtb fusion proteins and selected on SD/-HIS. To

induce protein expression, yeast were inoculated in 3mL galactose/raffinose (Gal/Raf)

base lacking histidine (Clontech) and allowed to grow 16-20 hours at 30°C, shaking.

Cultures were pelleted, resuspended in 30-50uL PBS and immobilized on an agar pad

prior to visualization.

Yeast Lysis and Western Blotting

Yeast (cdc25^{ts}) were inoculated into 5mL SD/-Leu and incubated overnight at room

temperature, shaking (250rpm). To lyse, 1.5mL of each culture was centrifuged at

14,000rpm for one minute. Each pellet was resuspended in 100µL 2.0M LiAc and

incubated on ice for five minutes. Samples were centrifuged at 14,000 for one minute to

pellet, resuspended in 100µL 0.4M NaOH and incubated on ice for five minutes.

Samples were pelleted as before, resuspended in 75µL 1x SDS Laemmli sample buffer

and boiled at 100°C for five minutes. Lysates were centrifuged 14,000rpm for one

minute to remove to debris, separated by SDS- polyacrylamide gel electrophoresis and

transferred to polyvinylidene difluoride membrane for Western blotting. Fusion proteins

were detected by rabbit anti-Ras (1:100) and equal loading was confirmed by detection

with rabbit anti-G6PDH (1:10,000).

Cell Culture

HeLa cells were cultured in Dulbecco's modified Eagle medium (DMEM, Gibco)

supplemented with 10% fetal bovine serum (FBS, Gibco), 100 I.U./mL penicillin, 100

µg/mL streptomycin, 292 µg/mL L-glutamine (Corning). RAW267.4 macrophages were

cultured in RPMI 1640 (Gibco) supplemented with 10% heat-inactivated FBS, 100

I.U./mL penicillin, 100 µg/mL streptomycin, 292 µg/mL L-glutamine, and 10mM HEPES

(Hyclone).

Primary human macrophages were isolated from buffy coats from anonymous

donors provided by a local blood bank. To isolate, 50mL of blood from each donor was

added to an equal volume of PBS then separated by centrifugation over a Ficoll-Paque

Plus gradient at 750 x g for 20 minutes with no brake. The lymphocyte/monocyte layer

was collected and incubated 1-2 minutes with 1mL ACK lysing buffer (Gibco) to remove

red blood cells. The cells were diluted to 50mL in PBS and centrifuged 350 x g for 10

minutes at 4°C. The supernatant was removed and cells were washed in 25mL PBS

and pelleted at 160 x g for 15 minutes at 4°C. Cells were washed again in 25 mL PBS

but centrifuged at 300 x g for 10 minutes at 4°C. This final pellet was resuspended in 5-

10 mL of RPMI 1640 supplemented with 10% human AB serum (Corning). To

differentiate into macrophages, cells were cultured in RPMI 1640 supplemented with

10% human AB serum and 50ng/mL human M-CSF (R&D Systems) for at least 4 hours,

washed in PBS then replaced with RPMI 1640 +10% human AB serum + human M-CSF

(50ng/mL) for 7 days with media changes every 1-2 days.

Antibodies

To generate an antibody against native Mpt64, two rabbits were immunized with

recombinant 6xHIS-tagged Mpt64ΔSP purified from *E. coli* in incomplete Freund's

adjuvant (Pacific Biosciences). The polyclonal rabbit antibody to Antigen 85 and

mouse anti-GroEL2 (CS-44) are from BEI Resources. Chicken, mouse and rabbit anti-

calreticulin were purchased from Abcam and anti-GM130 was purchased from BD

Biosciences. Mouse anti-Tom20 F-10 was purchased from Santa Cruz. Rabbit anti-Ras

and rabbit anti-glucose-6-phosphate dehydrogenase (G6PDH) were purchased from

Cell Signaling Technology and Sigma, respectively. Mouse anti-PMP70 and horse

radish peroxidase (HRP)-conjugated secondary antibodies were purchased from

Thermo Scientific. Alexa fluor-conjugated secondary antibodies were from Life

Technologies.

Molecular Biology

Unless otherwise stated, all Mtb proteins were cloned from the BEI resources Gateway

Mtb ORF library using Gateway cloning technology (Life Technologies). The Mpt64

truncation mutants were PCR amplified (Supplemental Table 2) and cloned into pENTR

(Life Technologies) prior to cloning into subsequent destination vectors.

hGH Secretion Assay and Quantification

HeLa cells were plated in 24-well plates to achieve approximately 50,000 cells/well 24h

prior to transfection. Cells were co-transfected with 1 μg hGH-CAD and 1μg GFP-Mtb

effector or GFP alone using FuGene 6 (Promega) per manufacturer instructions. Cells

were transfected 16-18h at 37°C 5% CO_2. The transfection media was then aspirated

and replaced with DMEM containing 2 µM D/D Solubilizer (Clontech) and incubated for

2 h at 37°C 5% CO_2. The plates were centrifuged at 1500 RPM for 5 minutes to pellet

debris and the culture supernatants were saved at -80°C prior to hGH quantification.

Secreted hGH was quantified by ELISA (Roche, 11585878001). Briefly, samples

were thawed on ice and 20 µL was transferred to each well containing 180µL sample

buffer (1:10). The plate was incubated 1h at 37°C, washed 5 times in 250µL wash buffer

and incubated 1h at 37°C with a polyclonal antibody to hGH conjugated to digoxigenin

(α-hGH-DIG). The plate was washed as described and incubated 1 h at 37°C with a

polyclonal antibody to digoxigenin conjugated to peroxidase (α-DIG-POD). The plate

was   washed   and   developed   in   peroxidase   substrate   (2,2'-Azinobis   [3-

ethylbenzothiazoline-6-sulfonic acid]-diammonium salt). The absorbance was read on a

Biotek plate reader at 405nm.

PIP strips membrane binding

6xHIS-Mpt64_24-228 and 6xHIS-Mpt64_24-143 were purified by cobalt TALON affinity

resin (Clontech). PIP strips (Invitrogen) were blocked for one hour at room temperature

in 3% fatty-acid free bovine serum albumin (BSA) in TBST. Mpt64_24-228 or Mpt64_24-

143 was diluted to 1.5ug/mL in 3mL 3% fatty-acid free BSA and incubated with the PIP

strips for 3h at room temperature with agitation. Membranes were washed three times in

3% fatty-acid free BSA prior to incubation with anti-Mpt64 or pre-immune serum

(1:3,000) overnight at 4°C, with agitation. Membranes were washed three times in 3%

fatty-acid free BSA then incubated with HRP-conjugated donkey anti-rabbit (1:2000) for

30 minutes at room temperature. Membranes were washed three times before detection

of Mpt64 lipid interactions by chemiluminescence.

Transfection and co-localzation of MEPs in HeLa cells

HeLa cells were transfected overnight with GFP fusion proteins using FuGene 6

transfection reagent (Roche). Cells were fixed in 4% paraformaldehyde (PFA) for 15

minutes, washed in PBS and permeabilized in 0.25% Triton X-100 for 3 minutes at room

temperature or in 100% methanol for 10 minutes at -20°C when using Tom20 antibody.

Cells were stained with organelle-specific antibodies for 1h at room temperature.

Antibodies were visualized by secondary antibodies conjugated to Alexa Fluor 594.

Cells were mounted in ProLong Gold + DAPI and z-stacks were collected on an

AxioImager M2 microscope (Zeiss).

Infection and co-localization of Mpt64 in macrophages

Bacteria were washed repeatedly in PBS, then sonicated to create a single cell

suspension. RAW267.4 cells were infected in DMEM+10% horse serum at MOI 20:1

with mycobacteria expressing mCherry. Cells were centrifuged at 1500 rpm for 10

minutes to permit bacterial attachment, then allowed to phagocytose for 1.5h at 37°C

5% CO_2. Cells were fixed after 4 and 24 hours post-infection in 4% PFA for 60 mintues

Cells were permeablized in 0.25% Triton X-100 for 3 minutes at room temperature then

blocked in 5% normal donkey serum (Sigma). Mpt64 was detected with rabbit anti-

Mpt64 antibody (1:500) and an HRP-conjugated goat-anti rabbit secondary antibody

(1:1000, Santa Cruz). Antibody signal was amplified by addition of biotinylated Tyramide

(1:50, PerkinElmer) with detection by Alexa fluor 488-conjugated streptavidin (1:250,

Jackson Immunoresearch) or Cyanine 5 Tyramide (1:50, PerkinElmer) . Z-stack slices

were acquired with an AxioImager M2 microscope (Zeiss).

Primary human macrophages were infected in RPMI + 10% human AB serum at

an MOI 10:1 with mycobacteria expressing mCherry for 2h at 37°C 5% CO_2 to allow for

phagocytosis. Cells were washed and fixed at 4 hours post-infection in 4% PFA for 45-

60 minutes. Cells were permeabilized in 100% ice cold methanol for 10 minutes at -

20°C and blocked in 5% normal goat serum (Sigma). Mpt64 was detected with rabbit

anti-Mpt64 antibody (1:500) and an HRP-conjugated donkey-anti rabbit secondary

antibody (1:500, Thermo Scientific) followed by amplification with cyanine 5 Tyramide

(1:50, PerkinElmer). Co-localization of Mpt64 with the ER was detected with chicken

anti-calreticulin (1:100), followed by goat anti-chicken-488 (Abcam).

Macrophage infections for CFU

Primary human macrophages were seeded in low-evaporation 24-well plates at

approximately 5 x10^5 cells/well. Bacteria were washed repeatedly in PBS, then

sonicated to create a single cell suspension. Macrophages were infected in RPMI +

10% human AB serum at MOI 0.5 for 2 hours at 37°C 5% CO_2 to allow phagocytosis to

occur. The cells were washed in PBS then replaced with RPMI + 10% human AB serum

and cells were washed every day between time points. The cells were lysed at time

zero and subsequent time points in 500 μL 0.5% Triton X-100 in PBS. Serial dilutions

were plated on 7H11 plates and colonies were enumerated after 2-3 weeks.

Construction of the Mtb Mpt64 deletion mutant and complementation

An in-frame Mpt64 deletion in Mtb was made using mycobacteriophage as previously

described (92). Briefly, 500-bp 5' to the Mpt64 start codon and 500-bp 3' to the Mpt64

stop codon were amplified from Erdman genomic DNA (Supplemental Table 2) and

sequentially cloned into the multiple cloning sites of pMSG360HYG. This vector was

linearized with AflII and DraI and transformed into EL350/phAE87 *E. coli* by

electroporation. Phagemid DNA was isolated from pooled colonies and transformed into

M. smegmatis by electroporation. Plaques were isolated and pooled from *M. smegmatis*

lawns and high titer phage was produced. Log phase Mtb Erdman was transduced with

phage at 42°C for 4h. Mutants were selected on 7H11+Hygromycin (100μg/mL). Wild-

type Mpt64 and Mpt64 lacking its secretion signal were cloned into an integrating vector

containing a constitutive promoter (pMV306_MSP), conferring zeocin resistance. The

MtbΔMpt64 was transformed by electroporation and complements were selected on

7H11 + zeocin (25 μg /mL).

To confirm expression and secretion of Mpt64 complements, Mtb strains were

grown to late-log phase and pelleted by centrifugation. The culture supernatants were

saved and passed twice through 0.22 μm filters. Bacterial pellets were boiled 30

minutes in lysis buffer (50mM Tris, pH 7.4, 150mM NaCl) supplemented with Complete

Mini protease inhibitor (Roche), then subjected to bead beating to lyse the cells. Protein

content in lysates was determined by Bradford assay. Mpt64 expression in the lysates

and culture supernatants was detected by Western blotting using a rabbit polyclonal

antibody to Mpt64 (1:10,000). Equal loading of samples in the lysates and supernatants

was confirmed by Western blotting with anti-GroEL2 (1:500) and anti-antigen 85

(1:1000) respectively.

Mouse infections

Female BALBc mice (The Jackson Laboratory) were infected via aerosol as described

previously (116). Briefly, mid-log phase Mtb were washed in PBS repeatedly then

sonicated to create a single-cell suspension. Bacteria were resuspended to yield an

OD_{600}=0.1 in PBS. This suspension was transferred to the nebulizer of a GlassCol

aerosolization chamber calibrated to infect mice with ~100 bacteria per animal. On the

day of infection, whole lungs were collected from 5 mice per group, homogenized and

plated on 7H11 to determine initial inoculum. At subsequent time points, the left lung,

spleen and left lobe of the liver were used to determine CFU, while the right lung was

insulfated with 10% neutral buffered formalin for histopathology. Animal experiments

were reviewed and approved by the Institutional Animal Care and Use Committee at the

University of Texas Southwestern.

Statistical analysis

Statistical analysis was performed using GraphPad Prism software. For in vivo CFU

calculations and area of lung inflammation, the non-parametric Kruskal-Wallis test with

Dunn's multiple comparison was used. Analysis of survival studies was performed by

Kaplan-Meier test. In vitro fold CFU was calculated by dividing the average CFU of

quadruplicates of each Mtb strain at each time point by the average at Day 0.

Propagation of error (z) for each fold calculation was determined by the equation:

$$z = foldD_n \sqrt{\left(\frac{StdError\ D_n}{Avg\ D_n}\right)^2 + \left(\frac{StdError\ D_0}{Avg\ D_0}\right)^2}$$

where D_n represents the time point value for any given strain and D_0 represents the day

zero values for that given strain (117). Statistical analysis on fold CFU was performed

by ANOVA.

Lysozyme pull down

E. coli lysates containing 6x-histidine (6xHIS) tagged Mpt64 or an unrelated protein Cor

were incubated with cobalt affinity resin (TALON, Clontech) to bind histidine-tagged

proteins. After extensive washing, 1mg/mL either hen egg white or human lysozyme

was flowed over the immobilized beads and incubated 5 minutes. Beads were washed

two more times before proteins were eluted with 300mM imidazole.

Mtb genomic DNA isolation

Late exponential phase Mtb was collected by centrifugation and washed once in PBS.

Pellets were boiled 20-30 minutes to sterilize. Pellets were washed once in GTE (25mM

Tris, pH 8.0; 10mM EDTA; 50mM glucose) and incubated overnight in lysozyme

solution (10mg/mL in GTE) at 37°C. Samples were incubated in 10% SDS and

10mg/mL proteinase K for 40minutes at 55°C followed by incubation in NaCl and CTAB

(2.4M NaCl, 274mM cetrimonium bromide (Sigma)) at 60°C for 10 minutes. Genomic

DNA was then isolated using a phenol-chloroform extraction followed by ethanol

precipitation.

Extraction of apolar lipids and PDIM analysis

Log phage Mtb or MtbΔMpt64 were synchronized to OD_{600}=0.2 in 7H9 supplemented

with 0.01% Tween 80 and grown 24 hours. Bacteria were collected by centrifugation at

1,600xg for 10 minutes, resuspended in 1mL 15% isopropanol and transferred to a

glass tube containing 5mL chloroform: methanol (17:1, v/v) and incubated 24 hours at

room temperature. Samples were centrifuged at 1,600xg for five minutes and the apolar

lipids were collected from the bottom, organic layer and dried. Apolar lipids were

resuspended in 1.5mL 100% methanol. Tween 80 was removed by addition of

cobalt/thiocyanate solution and vortexed. Remaining lipids were extracted by addition of

4mL hexane. After centrifugation the organic layer was saved and the aqueous layer

was re-extracted with 4mL hexane. Both hexane fractions were combined, dried and

resuspended in 1mL chloroform: methanol (2:1, v/v). PDIM standard was similarly

resuspended. PDIM standard, or apolar lipids extracted from Mtb or MtbΔMpt64 were

infused into an AbSciex TripleTOF 5600/5600+ mass spectrometer. Samples were

analyzed in the positive mode.

Acknowledgements

We would like to thank Patrick Cherry and Molly Moehlman for their contributions to this

work during their summer internships. We thank BEI resources for providing reagents.

This work was supported by NIH Grants R01 AI099439, R21 AI111023 and U01

AI125939 (to M.U.S.) and T32AI007520 (to C.E.S and B.L.P). M.U.S. is a Disease-

Oriented Clinical Scholar at University of Texas Southwestern Medical Center.

C.E.S., N.M.A. and M.U.S. conceived and designed experiments. B.A.W. provided

reagents and experimental advice. C.E.S., and S.C. performed immunofluorescence

microscopy. C.E.S., B.L.P., L.H.F., and V.R.N. performed mouse aerosol infections with

Mtb. B.L.P. performed mass spectrometry analysis on PDIM. C.E.S. performed the

remaining research include screening experiments, molecular biology, bacterial

genetics and macrophage infections. C.E.S. and M.U.S. drafted the manuscript. All

authors edited and approved of the final manuscript.

References

1.   WHO. 2016. Global tuberculosis report. WHO Press, World Health Organization.

2.   Colditz GA, Brewer TF, Berkey CS, Wilson ME, Burdick E, Fineberg HV,

Mosteller F. 1994. Efficacy of BCG vaccine in the prevention of tuberculosis.

Meta-analysis of the published literature. JAMA 271:698-702.

3.   Huynh KK, Joshi SA, Brown EJ. 2011. A delicate dance: host response to

mycobacteria. Curr Opin Immunol 23:464-72.

4.   Flannagan RS, Cosio G, Grinstein S. 2009. Antimicrobial mechanisms of

phagocytes and bacterial evasion strategies. Nat Rev Microbiol 7:355-66.

5.   Weiss G, Schaible UE. 2015. Macrophage defense mechanisms against

intracellular bacteria. Immunol Rev 264:182-203.

6.   Bradfute SB, Castillo EF, Arko-Mensah J, Chauhan S, Jiang S, Mandell M,

Deretic V. 2013. Autophagy as an immune effector against tuberculosis. Current

opinion in microbiology 16:355-365.

7.   Davis AS, Vergne I, Master SS, Kyei GB, Chua J, Deretic V. 2007. Mechanism of
inducible nitric oxide synthase exclusion from mycobacterial phagosomes. PLoS
Pathog 3:e186.

8.   Vandal OH, Pierini LM, Schnappinger D, Nathan CF, Ehrt S. 2008. A membrane
protein preserves intrabacterial pH in intraphagosomal Mycobacterium
tuberculosis. Nat Med 14:849-54.

9.   Ehrt S, Schnappinger D. 2009. Mycobacterial survival strategies in the
phagosome: defence against host stresses. Cell Microbiol 11:1170-8.

10.   Vergne I, Chua J, Singh SB, Deretic V. 2004. Cell biology of mycobacterium
tuberculosis phagosome. Annu Rev Cell Dev Biol 20:367-94.

11.   Sturgill-Koszycki S, Schlesinger PH, Chakraborty P, Haddix PL, Collins HL, Fok
AK, Allen RD, Gluck SL, Heuser J, Russell DG. 1994. Lack of acidification in
Mycobacterium phagosomes produced by exclusion of the vesicular proton-
ATPase. Science 263:678-81.

12.   Via LE, Deretic D, Ulmer RJ, Hibler NS, Huber LA, Deretic V. 1997. Arrest of
mycobacterial phagosome maturation is caused by a block in vesicle fusion
between stages controlled by rab5 and rab7. J Biol Chem 272:13326-31.

13.   Pethe K, Swenson DL, Alonso S, Anderson J, Wang C, Russell DG. 2004.
Isolation of Mycobacterium tuberculosis mutants defective in the arrest of
phagosome maturation. Proc Natl Acad Sci U S A 101:13642-7.

14.   Stewart GR, Patel J, Robertson BD, Rae A, Young DB. 2005. Mycobacterial
mutants with defective control of phagosomal acidification. PLoS Pathog 1:269-
78.

15.   MacGurn JA, Cox JS. 2007. A genetic screen for Mycobacterium tuberculosis

mutants defective for phagosome maturation arrest identifies components of the

ESX-1 secretion system. Infect Immun 75:2668-78.

16.   Saleh MT, Belisle JT. 2000. Secretion of an acid phosphatase (SapM) by

Mycobacterium tuberculosis that is similar to eukaryotic acid phosphatases. J

Bacteriol 182:6850-3.

17.   Vergne I, Chua J, Lee HH, Lucas M, Belisle J, Deretic V. 2005. Mechanism of

phagolysosome biogenesis block by viable Mycobacterium tuberculosis. Proc

Natl Acad Sci U S A 102:4033-8.

18.   Zulauf KE, Sullivan JT, Braunstein M. 2018. The SecA2 pathway of

Mycobacterium tuberculosis exports effectors that work in concert to arrest

phagosome and autophagosome maturation. PLoS Pathog 14:e1007011.

19.   Sassetti CM, Rubin EJ. 2003. Genetic requirements for mycobacterial survival

during infection. Proc Natl Acad Sci U S A 100:12989-94.

20.   Costa TR, Felisberto-Rodrigues C, Meir A, Prevost MS, Redzej A, Trokter M,

Waksman G. 2015. Secretion systems in Gram-negative bacteria: structural and

mechanistic insights. Nat Rev Microbiol 13:343-59.

21.   Galan JE, Collmer A. 1999. Type III secretion machines: bacterial devices for

protein delivery into host cells. Science 284:1322-8.

22.   Deng W, Marshall NC, Rowland JL, McCoy JM, Worrall LJ, Santos AS,

Strynadka NCJ, Finlay BB. 2017. Assembly, structure, function and regulation of

type III secretion systems. Nat Rev Microbiol 15:323-337.

23. Alvarez-Martinez CE, Christie PJ. 2009. Biological diversity of prokaryotic type IV

secretion systems. Microbiol Mol Biol Rev 73:775-808.

24. Ho BT, Dong TG, Mekalanos JJ. 2014. A view to a kill: the bacterial type VI

secretion system. Cell Host Microbe 15:9-21.

25. Isaac DT, Isberg R. 2014. Master manipulators: an update on Legionella

pneumophila Icm/Dot translocated substrates and their host targets. Future

Microbiol 9:343-59.

26. Backert S, Meyer TF. 2006. Type IV secretion systems and their effectors in

bacterial pathogenesis. Curr Opin Microbiol 9:207-17.

27. Barker JR, Chong A, Wehrly TD, Yu JJ, Rodriguez SA, Liu J, Celli J,

Arulanandam BP, Klose KE. 2009. The Francisella tularensis pathogenicity

island encodes a secretion system that is required for phagosome escape and

virulence. Mol Microbiol 74:1459-70.

28. Russell AB, Peterson SB, Mougous JD. 2014. Type VI secretion system

effectors: poisons with a purpose. Nat Rev Microbiol 12:137-48.

29. Stanley SA, Raghavan S, Hwang WW, Cox JS. 2003. Acute infection and

macrophage subversion by Mycobacterium tuberculosis require a specialized

secretion system. Proc Natl Acad Sci U S A 100:13001-6.

30. Groschel MI, Sayes F, Simeone R, Majlessi L, Brosch R. 2016. ESX secretion

systems: mycobacterial evolution to counter host immunity. Nat Rev Microbiol

14:677-691.

31. Alix E, Mukherjee S, Roy CR. 2011. Subversion of membrane transport

pathways by vacuolar pathogens. J Cell Biol 195:943-52.

32.   Alto NM, Orth K. 2012. Subversion of cell signaling by pathogens. Cold Spring

Harb Perspect Biol 4:a006114.

33.   Ham H, Sreelatha A, Orth K. 2011. Manipulation of host membranes by bacterial

effectors. Nat Rev Microbiol 9:635-46.

34.   Boucrot E, Beuzon CR, Holden DW, Gorvel JP, Meresse S. 2003. Salmonella

typhimurium SifA effector protein requires its membrane-anchoring C-terminal

hexapeptide for its biological function. J Biol Chem 278:14196-202.

35.   Zhao W, Moest T, Zhao Y, Guilhon AA, Buffat C, Gorvel JP, Meresse S. 2015.

The Salmonella effector protein SifA plays a dual role in virulence. Sci Rep

5:12979.

36.   Murata T, Delprato A, Ingmundson A, Toomre DK, Lambright DG, Roy CR. 2006.

The Legionella pneumophila effector protein DrrA is a Rab1 guanine nucleotide-

exchange factor. Nat Cell Biol 8:971-7.

37.   Niebuhr K, Giuriato S, Pedron T, Philpott DJ, Gaits F, Sable J, Sheetz MP,

Parsot C, Sansonetti PJ, Payrastre B. 2002. Conversion of PtdIns(4,5)P(2) into

PtdIns(5)P by the S.flexneri effector IpgD reorganizes host cell morphology.

EMBO J 21:5069-78.

38.   Pizarro-Cerda J, Charbit A, Enninga J, Lafont F, Cossart P. 2016. Manipulation

of host membranes by the bacterial pathogens Listeria, Francisella, Shigella and

Yersinia. Semin Cell Dev Biol 60:155-167.

39.   Weigele BA, Orchard RC, Jimenez A, Cox GW, Alto NM. 2017. A systematic

exploration of the interactions between bacterial effector proteins and host cell

membranes. Nat Commun 8:532.

40. Ligon LS, Hayden JD, Braunstein M. 2012. The ins and outs of Mycobacterium

tuberculosis protein export. Tuberculosis (Edinb) 92:121-32.

41. Manzanillo PS, Shiloh MU, Portnoy DA, Cox JS. 2012. Mycobacterium

tuberculosis activates the DNA-dependent cytosolic surveillance pathway within

macrophages. Cell host & microbe 11:469-480.

42. Stanley SA, Johndrow JE, Manzanillo P, Cox JS. 2007. The Type I IFN response

to infection with Mycobacterium tuberculosis requires ESX-1-mediated secretion

and contributes to pathogenesis. Journal of immunology (Baltimore, Md : 1950)

178:3143-3152.

43. Mehra A, Zahra A, Thompson V, Sirisaengtaksin N, Wells A, Porto M, Koster S,

Penberthy K, Kubota Y, Dricot A, Rogan D, Vidal M, Hill DE, Bean AJ, Philips JA.

2013. Mycobacterium tuberculosis type VII secreted effector EsxH targets host

ESCRT to impair trafficking. PLoS Pathog 9:e1003734.

44. Portal-Celhay C, Tufariello JM, Srivastava S, Zahra A, Klevorn T, Grace PS,

Mehra A, Park HS, Ernst JD, Jacobs WR, Jr., Philips JA. 2016. Mycobacterium

tuberculosis EsxH inhibits ESCRT-dependent CD4+ T-cell activation. Nat

Microbiol 2:16232.

45. Feltcher ME, Gunawardena HP, Zulauf KE, Malik S, Griffin JE, Sassetti CM,

Chen X, Braunstein M. 2015. Label-free Quantitative Proteomics Reveals a Role

for the Mycobacterium tuberculosis SecA2 Pathway in Exporting Solute Binding

Proteins and Mce Transporters to the Cell Wall. Mol Cell Proteomics 14:1501-16.

46. Walburger A, Koul A, Ferrari G, Nguyen L, Prescianotto-Baschong C, Huygen K,

Klebl B, Thompson C, Bacher G, Pieters J. 2004. Protein kinase G from

pathogenic mycobacteria promotes survival within macrophages. Science

304:1800-4.

47. Jimenez A, Chen D, Alto NM. 2016. How Bacteria Subvert Animal Cell Structure

and Function. Annu Rev Cell Dev Biol 32:373-397.

48. Rosenkrands I, Weldingh K, Jacobsen S, Hansen CV, Florio W, Gianetri I,

Andersen P. 2000. Mapping and identification of Mycobacterium tuberculosis

proteins by two-dimensional gel electrophoresis, microsequencing and

immunodetection. Electrophoresis 21:935-48.

49. Mattow J, Schaible UE, Schmidt F, Hagens K, Siejak F, Brestrich G, Haeselbarth

G, Muller EC, Jungblut PR, Kaufmann SH. 2003. Comparative proteome analysis

of culture supernatant proteins from virulent Mycobacterium tuberculosis H37Rv

and attenuated M. bovis BCG Copenhagen. Electrophoresis 24:3405-20.

50. Malen H, Berven FS, Fladmark KE, Wiker HG. 2007. Comprehensive analysis of

exported proteins from Mycobacterium tuberculosis H37Rv. Proteomics 7:1702-

18.

51. de Souza GA, Leversen NA, Malen H, Wiker HG. 2011. Bacterial proteins with

cleaved or uncleaved signal peptides of the general secretory pathway. J

Proteomics 75:502-10.

52. Kelkar DS, Kumar D, Kumar P, Balakrishnan L, Muthusamy B, Yadav AK,

Shrivastava P, Marimuthu A, Anand S, Sundaram H, Kingsbury R, Harsha HC,

Nair B, Prasad TS, Chauhan DS, Katoch K, Katoch VM, Kumar P, Chaerkady R,

Ramachandran S, Dash D, Pandey A. 2011. Proteogenomic analysis of

Mycobacterium tuberculosis by high resolution mass spectrometry. Mol Cell

Proteomics 10:M111 011627.

53. Conrad WH, Osman MM, Shanahan JK, Chu F, Takaki KK, Cameron J,

Hopkinson-Woolley D, Brosch R, Ramakrishnan L. 2017. Mycobacterial ESX-1

secretion system mediates host cell lysis through bacterium contact-dependent

gross membrane disruptions. Proc Natl Acad Sci U S A 114:1371-1376.

54. McCann JR, McDonough JA, Sullivan JT, Feltcher ME, Braunstein M. 2011.

Genome-wide identification of Mycobacterium tuberculosis exported proteins with

roles in intracellular growth. J Bacteriol 193:854-61.

55. Jungblut PR, Schaible UE, Mollenkopf HJ, Zimny-Arndt U, Raupach B, Mattow J,

Halada P, Lamer S, Hagens K, Kaufmann SH. 1999. Comparative proteome

analysis of Mycobacterium tuberculosis and Mycobacterium bovis BCG strains:

towards functional genomics of microbial pathogens. Mol Microbiol 33:1103-17.

56. Rosenkrands I, King A, Weldingh K, Moniatte M, Moertz E, Andersen P. 2000.

Towards the proteome of Mycobacterium tuberculosis. Electrophoresis 21:3740-

56.

57. Marjanovic O, Miyata T, Goodridge A, Kendall LV, Riley LW. 2010. Mce2 operon

mutant strain of Mycobacterium tuberculosis is attenuated in C57BL/6 mice.

Tuberculosis (Edinb) 90:50-6.

58. Gioffre A, Infante E, Aguilar D, Santangelo MP, Klepp L, Amadio A, Meikle V,

Etchechoury I, Romano MI, Cataldi A, Hernandez RP, Bigi F. 2005. Mutation in

mce operons attenuates Mycobacterium tuberculosis virulence. Microbes Infect

7:325-34.

59.   Saini NK, Sharma M, Chandolia A, Pasricha R, Brahmachari V, Bose M. 2008.

917   Characterization of Mce4A protein of Mycobacterium tuberculosis: role in

918   invasion and survival. BMC Microbiol 8:200.

919   60.   Brodin P, Poquet Y, Levillain F, Peguillet I, Larrouy-Maumus G, Gilleron M,

920   Ewann F, Christophe T, Fenistein D, Jang J, Jang MS, Park SJ, Rauzier J,

921   Carralot JP, Shrimpton R, Genovesio A, Gonzalo-Asensio JA, Puzo G, Martin C,

922   Brosch R, Stewart GR, Gicquel B, Neyrolles O. 2010. High content phenotypic

923   cell-based visual screen identifies Mycobacterium tuberculosis acyltrehalose-

924   containing glycolipids involved in phagosome remodeling. PLoS Pathog

925   6:e1001100.

926   61.   Sampson SL. 2011. Mycobacterial PE/PPE proteins at the host-pathogen

927   interface. Clin Dev Immunol 2011:497203.

928   62.   Bottai D, Di Luca M, Majlessi L, Frigui W, Simeone R, Sayes F, Bitter W,

929   Brennan MJ, Leclerc C, Batoni G, Campa M, Brosch R, Esin S. 2012. Disruption

930   of the ESX-5 system of Mycobacterium tuberculosis causes loss of PPE protein

931   secretion, reduction of cell wall integrity and strong attenuation. Mol Microbiol

932   83:1195-209.

933   63.   Shah S, Briken V. 2016. Modular Organization of the ESX-5 Secretion System in

934   Mycobacterium tuberculosis. Front Cell Infect Microbiol 6:49.

935   64.   Sassetti CM, Boyd DH, Rubin EJ. 2003. Genes required for mycobacterial growth

936   defined by high density mutagenesis. Mol Microbiol 48:77-84.

65.   Rengarajan J, Bloom BR, Rubin EJ. 2005. Genome-wide requirements for

Mycobacterium tuberculosis adaptation and survival in macrophages. Proc Natl

Acad Sci U S A 102:8327-32.

66.   Rosas-Magallanes V, Stadthagen-Gomez G, Rauzier J, Barreiro LB, Tailleux L,

Boudou F, Griffin R, Nigou J, Jackson M, Gicquel B, Neyrolles O. 2007.

Signature-tagged transposon mutagenesis identifies novel Mycobacterium

tuberculosis genes involved in the parasitism of human macrophages. Infect

Immun 75:504-7.

67.   Isakoff SJ, Cardozo T, Andreev J, Li Z, Ferguson KM, Abagyan R, Lemmon MA,

Aronheim A, Skolnik EY. 1998. Identification and analysis of PH domain-

containing targets of phosphatidylinositol 3-kinase using a novel in vivo assay in

yeast. EMBO J 17:5374-87.

68.   Aronheim A, Zandi E, Hennemann H, Elledge SJ, Karin M. 1997. Isolation of an

AP-1 repressor by a novel method for detecting protein-protein interactions. Mol

Cell Biol 17:3094-102.

69.   Chiu VK, Bivona T, Hach A, Sajous JB, Silletti J, Wiener H, Johnson RL, 2nd,

Cox AD, Philips MR. 2002. Ras signalling on the endoplasmic reticulum and the

Golgi. Nat Cell Biol 4:343-50.

70.   Yu JW, Mendrola JM, Audhya A, Singh S, Keleti D, DeWald DB, Murray D, Emr

SD, Lemmon MA. 2004. Genome-wide analysis of membrane targeting by S.

cerevisiae pleckstrin homology domains. Mol Cell 13:677-88.

71.   Personnic N, Barlocher K, Finsel I, Hilbi H. 2016. Subversion of Retrograde

Trafficking by Translocated Pathogen Effectors. Trends Microbiol 24:450-62.

72. Sullivan JT, Young EF, McCann JR, Braunstein M. 2012. The Mycobacterium

tuberculosis SecA2 system subverts phagosome maturation to promote growth in

macrophages. Infect Immun 80:996-1006.

73. Bach H, Papavinasasundaram KG, Wong D, Hmama Z, Av-Gay Y. 2008.

Mycobacterium tuberculosis virulence is mediated by PtpA dephosphorylation of

human vacuolar protein sorting 33B. Cell Host Microbe 3:316-22.

74. Rivera VM, Wang X, Wardwell S, Courage NL, Volchuk A, Keenan T, Holt DA,

Gilman M, Orci L, Cerasoli F, Jr., Rothman JE, Clackson T. 2000. Regulation of

protein secretion through controlled aggregation in the endoplasmic reticulum.

Science 287:826-30.

75. Selyunin AS, Sutton SE, Weigele BA, Reddick LE, Orchard RC, Bresson SM,

Tomchick DR, Alto NM. 2011. The assembly of a GTPase-kinase signalling

complex by a bacterial catalytic scaffold. Nature 469:107-11.

76. Selyunin AS, Reddick LE, Weigele BA, Alto NM. 2014. Selective protection of an

ARF1-GTP signaling axis by a bacterial scaffold induces bidirectional trafficking

arrest. Cell Rep 6:878-91.

77. Shenoy VP, Mukhopadhyay C. 2014. Rapid Immunochromatographic Test for the

Identification and Discrimination of Mycobacterium tuberculosis Complex Isolates

from Non-tuberculous Mycobacteria. J Clin Diagn Res 8:DC13-5.

78. Arora J, Kumar G, Verma AK, Bhalla M, Sarin R, Myneedu VP. 2015. Utility of

MPT64 Antigen Detection for Rapid Confirmation of Mycobacterium tuberculosis

Complex. J Glob Infect Dis 7:66-9.

79.   Behr MA, Wilson MA, Gill WP, Salamon H, Schoolnik GK, Rane S, Small PM.

983   1999. Comparative genomics of BCG vaccines by whole-genome DNA

984   microarray. Science 284:1520-3.

985   80.   Kozak RA, Alexander DC, Liao R, Sherman DR, Behr MA. 2011. Region of

986   difference 2 contributes to virulence of Mycobacterium tuberculosis. Infect Immun

987   79:59-66.

988   81.   Sonnenberg MG, Belisle JT. 1997. Definition of Mycobacterium tuberculosis

989   culture filtrate proteins by two-dimensional polyacrylamide gel electrophoresis, N-

990   terminal amino acid sequencing, and electrospray mass spectrometry. Infect

991   Immun 65:4515-24.

992   82.   Wang Z, Potter BM, Gray AM, Sacksteder KA, Geisbrecht BV, Laity JH. 2007.

993   The solution structure of antigen MPT64 from Mycobacterium tuberculosis

994   defines a new family of beta-grasp proteins. J Mol Biol 366:375-81.

995   83.   Hastie JL, Williams KB, Bohr LL, Houtman JC, Gakhar L, Ellermeier CD. 2016.

996   The Anti-sigma Factor RsiV Is a Bacterial Receptor for Lysozyme: Co-crystal

997   Structure Determination and Demonstration That Binding of Lysozyme to RsiV Is

998   Required for sigmaV Activation. PLoS Genet 12:e1006287.

999   84.   Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. 2015. The Phyre2

1000   web portal for protein modeling, prediction and analysis. Nat Protoc 10:845-58.

1001   85.   Breker M, Gymrek M, Schuldiner M. 2013. A novel single-cell screening platform

1002   reveals proteome plasticity during yeast stress responses. J Cell Biol 200:839-

1003   50.

86. Huh WK, Falvo JV, Gerke LC, Carroll AS, Howson RW, Weissman JS, O'Shea
EK. 2003. Global analysis of protein localization in budding yeast. Nature
425:686-91.

87. Champion PA, Stanley SA, Champion MM, Brown EJ, Cox JS. 2006. C-terminal
signal sequence promotes virulence factor secretion in Mycobacterium
tuberculosis. Science 313:1632-6.

88. Hsu T, Hingley-Wilson SM, Chen B, Chen M, Dai AZ, Morin PM, Marks CB,
Padiyar J, Goulding C, Gingery M, Eisenberg D, Russell RG, Derrick SC, Collins
FM, Morris SL, King CH, Jacobs WR, Jr. 2003. The primary mechanism of
attenuation of bacillus Calmette-Guerin is a loss of secreted lytic function
required for invasion of lung interstitial tissue. Proc Natl Acad Sci U S A
100:12420-5.

89. van der Wel N, Hava D, Houben D, Fluitsma D, van Zon M, Pierson J, Brenner
M, Peters PJ. 2007. M. tuberculosis and M. leprae translocate from the
phagolysosome to the cytosol in myeloid cells. Cell 129:1287-98.

90. Bardarov S, Kriakov J, Carriere C, Yu S, Vaamonde C, McAdam RA, Bloom BR,
Hatfull GF, Jacobs WR, Jr. 1997. Conditionally replicating mycobacteriophages:
a system for transposon delivery to Mycobacterium tuberculosis. Proc Natl Acad
Sci U S A 94:10961-6.

91. Glickman MS, Cox JS, Jacobs WR, Jr. 2000. A novel mycolic acid cyclopropane
synthetase is required for cording, persistence, and virulence of Mycobacterium
tuberculosis. Mol Cell 5:717-27.

92. Glickman MS, Cahill SM, Jacobs WR, Jr. 2001. The Mycobacterium tuberculosis

cmaA2 gene encodes a mycolic acid trans-cyclopropane synthetase. J Biol

Chem 276:2228-33.

93. Chan K, Knaak T, Satkamp L, Humbert O, Falkow S, Ramakrishnan L. 2002.

Complex pattern of Mycobacterium marinum gene expression during long-term

granulomatous infection. Proc Natl Acad Sci U S A 99:3920-5.

94. Brennan MJ. 2017. The Enigmatic PE/PPE Multigene Family of Mycobacteria

and Tuberculosis Vaccination. Infect Immun 85.

95. Perkowski EF, Zulauf KE, Weerakoon D, Hayden JD, Ioerger TR, Oreper D,

Gomez SM, Sacchettini JC, Braunstein M. 2017. The EXIT Strategy: an

Approach for Identifying Bacterial Proteins Exported during Host Infection. MBio

8.

96. Sun J, Singh V, Lau A, Stokes RW, Obregón-Henao A, Orme IM, Wong D, Av-

Gay Y, Hmama Z. 2013. Mycobacterium tuberculosis nucleoside diphosphate

kinase inactivates small GTPases leading to evasion of innate immunity. PLoS

pathogens 9:e1003499.

97. Khan S, Islam A, Hassan MI, Ahmad F. 2016. Purification and structural

characterization of Mce4A from Mycobacterium tuberculosis. Int J Biol Macromol

93:235-241.

98. Daleke MH, Cascioferro A, de Punder K, Ummels R, Abdallah AM, van der Wel

N, Peters PJ, Luirink J, Manganelli R, Bitter W. 2011. Conserved Pro-Glu (PE)

and Pro-Pro-Glu (PPE) protein domains target LipY lipases of pathogenic

mycobacteria to the cell surface via the ESX-5 pathway. J Biol Chem 286:19024-

34.

99.   de Jonge MI, Pehau-Arnaudet G, Fretz MM, Romain F, Bottai D, Brodin P,

Honore N, Marchal G, Jiskoot W, England P, Cole ST, Brosch R. 2007. ESAT-6

from Mycobacterium tuberculosis Dissociates from Its Putative Chaperone CFP-

10 under Acidic Conditions and Exhibits Membrane-Lysing Activity. Journal of

Bacteriology 189:6028-6034.

100.   Smith J, Manoranjan J, Pan M, Bohsali A, Xu J, Liu J, McDonald KL, Szyk A,

LaRonde-LeBlanc N, Gao LY. 2008. Evidence for pore formation in host cell

membranes by ESX-1-secreted ESAT-6 and its role in Mycobacterium marinum

escape from the vacuole. Infect Immun 76:5478-87.

101.   Danilchanka O, Sun J, Pavlenok M, Maueroder C, Speer A, Siroy A, Marrero J,

Trujillo C, Mayhew DL, Doornbos KS, Munoz LE, Herrmann M, Ehrt S, Berens C,

Niederweis M. 2014. An outer membrane channel protein of Mycobacterium

tuberculosis with exotoxin activity. Proc Natl Acad Sci U S A 111:6750-5.

102.   Sun J, Siroy A, Lokareddy RK, Speer A, Doornbos KS, Cingolani G, Niederweis

M. 2015. The tuberculosis necrotizing toxin kills macrophages by hydrolyzing

NAD. Nat Struct Mol Biol 22:672-8.

103.   De Matteis MA, Wilson C, D'Angelo G. 2013. Phosphatidylinositol-4-phosphate:

the Golgi and beyond. Bioessays 35:612-22.

104.   Matsuoka K, Orci L, Amherdt M, Bednarek SY, Hamamoto S, Schekman R,

Yeung T. 1998. COPII-coated vesicle formation reconstituted with purified coat

proteins and chemically defined liposomes. Cell 93:263-75.

105.   Blumental-Perry A, Haney CJ, Weixel KM, Watkins SC, Weisz OA, Aridor M.

1072   2006. Phosphatidylinositol 4-phosphate formation at ER exit sites regulates ER

1073   export. Dev Cell 11:671-82.

1074   106.   Sarkes D, Rameh LE. 2010. A novel HPLC-based approach makes possible the

1075   spatial characterization of cellular PtdIns5P and other phosphoinositides.

1076   Biochem J 428:375-84.

1077   107.   Choi S, Thapa N, Tan X, Hedman AC, Anderson RA. 2015. PIP kinases define

1078   PI4,5P(2)signaling specificity by association with effectors. Biochim Biophys Acta

1079   1851:711-23.

1080   108.   Manna P, Jain SK. 2015. Phosphatidylinositol-3,4,5-triphosphate and cellular

1081   signaling: implications for obesity and diabetes. Cell Physiol Biochem 35:1253-

1082   75.

1083   109.   O'Connor TJ, Adepoju Y, Boyd D, Isberg RR. 2011. Minimization of the

1084   Legionella pneumophila genome reveals chromosomal regions involved in host

1085   range expansion. Proc Natl Acad Sci U S A 108:14733-40.

1086   110.   Ensminger AW. 2016. Legionella pneumophila, armed to the hilt: justifying the

1087   largest arsenal of effectors in the bacterial world. Curr Opin Microbiol 29:74-80.

1088   111.   Behar SM, Martin CJ, Nunes-Alves C, Divangahi M, Remold HG. 2011. Lipids,

1089   apoptosis, and cross-presentation: links in the chain of host defense against

1090   Mycobacterium tuberculosis. Microbes Infect 13:749-56.

1091   112.   Roche PW, Triccas JA, Avery DT, Fifis T, Billman-Jacobe H, Britton WJ. 1994.

1092   Differential T cell responses to mycobacteria-secreted proteins distinguish

vaccination with bacille Calmette-Guerin from infection with Mycobacterium

tuberculosis. J Infect Dis 170:1326-30.

113. Roche PW, Feng CG, Britton WJ. 1996. Human T-cell epitopes on the

Mycobacterium tuberculosis secreted protein MPT64. Scand J Immunol 43:662-

70.

114. Mustafa T, Wiker HG, Morkve O, Sviland L. 2007. Reduced apoptosis and

increased inflammatory cytokines in granulomas caused by tuberculous

compared to non-tuberculous mycobacteria: role of MPT64 antigen in apoptosis

and immune response. Clin Exp Immunol 150:105-13.

115. Mustafa T, Wiker HG, Morkve O, Sviland L. 2008. Differential expression of

mycobacterial antigen MPT64, apoptosis and inflammatory markers in

multinucleated giant cells and epithelioid cells in granulomas caused by

Mycobacterium tuberculosis. Virchows Arch 452:449-56.

116. Collins AC, Cai H, Li T, Franco LH, Li XD, Nair VR, Scharn CR, Stamm CE,

Levine B, Chen ZJ, Shiloh MU. 2015. Cyclic GMP-AMP Synthase Is an Innate

Immune DNA Sensor for Mycobacterium tuberculosis. Cell Host Microbe 17:820-

8.

117. Ku HH. 1966. Notes on the Use of Propagation of Error Formulas. Journal Of

Research of the National Bureau of Standards - C Engineering and

Instrumentation 70C:263-273.

1113

Figure legends

Figure 1. *M. tuberculosis* exported proteins interact with yeast membranes. (**A**) Ras rescue assay schematic. (**B**) *S. cerevisiae* (cdc25^{ts}) transformed with Mtb protein fusions to Ras^{mut}, duplicate plated and incubated 48-72 h at the permissive (25°C) and restrictive (37°C) temperatures. Shown are representative images of 20 yeast strains. (**C**) Summary results of Ras rescue screen. (**D**) Western blot of lysates from yeast transformed with the indicated fusion proteins and probed with anti-Ras or anti-G6PDH antibodies. Ras^{mut} fusion proteins are marked by white or black asterisks. PLC (phospholipase C) and DYN (dynamin) are fusions to Ras^{mut} known to be membrane associated (PLC) or cytoplasmic (DYN). (**E**) Representative fluorescence microscopy of *S. cerevisiae* (INVSc1) transformed with GFP-MEP fusion proteins. Scale bars are 3µm.

Figure 2. Host subcellular localization of membrane-binding MEP. (**A**) Representative fluorescent images of HeLa cells transfected with the indicated GFP-MEP fusion proteins (green) and stained with antibodies (red) to calreticulin (ER), GM-130 (Golgi), Tom20 (Mitochondria) or PMP70 (Peroxisomes). Scale bars are 5µm. (**B**) Comparison of the organelle localization of MEP expressed in yeast and HeLa cells.

Figure 3. *M. tuberculosis* exported proteins alter hGH secretion (**A**) Inducible secretion assay schematic. (**B**) Supernatant hGH ELISA from HeLa cells transfected overnight with hGH-CAD and either GFP or GFP-MEP fusion proteins prior to addition of drug to allow for hGH secretion. hGH secretion by GFP-MEP transfected cells was normalized to hGH secretion of cells transfected with GFP alone. (**C**) Summary of results from

1137 inducible secretion screen. (**D**) Venn diagram of MEP that are membrane-localized,

1138 alter host secretion, or both. (**E**) Table summarizing the membrane localization and

1139 degree of hGH secretion in cells transfected with the five overlapping proteins from **D**.

1140

Figure 4. hGH secretion inhibition is dependent on membrane localization of Mpt64. (**A**)

Solution structure of Mpt64. PDB 2hhi. (**B**) Schematic of Mpt64 truncations, colored to

match solution structure in **A**. SP, signal peptide. (**C**) Full length Mpt64 or protein

truncations expressed in the Ras rescue assay. (**D**) Western blot of lysates from cdc25^{ts}

yeast expressing Ras^{mut} fusion proteins to Mpt64 or Mpt64 truncations. Blots were

probed for anti-Ras or anti-G6PDH antibodies. Control protein bands are marked by a

white asterisk and Mpt64 truncations are marked by a black asterisk. (**E**) PIP

membrane strips incubated with recombinant Mpt64_24-228 or Mpt64_24-143. Binding

of Mpt64 to lipids was detected by incubation with α-Mpt64 or pre-immune serum.

Numbers indicate specific lipids as follows: 1, lysophosphatidic acid; 2,

lysophosphatidylcholine; 3, phosphatidylinositol; 4, phosphatidylinositol-3-phopshate

(PI3P); 5, PI4P; 6, PI5P; 7, phosphatidylethanolamine; 8, phosphatidylcholine; 9,

sphingosine 1-phosophate; 10, phosphatidylinositol-3,4-bisphosphate [PI(3,4)P$_2$]; 11,

PI(3,5)P$_2$; 12, PI(4,5)P$_2$; 13, phosphatidylinositol-3,4,5-trisphosphate [PI(3,4,5)P$_3$]; 14,

phosphatidic acid; 15, phosphatidylserine; 16, blank. (**F**) ELISA results of hGH in

supernatants of cells co-expressing full length Mpt64, Mpt64 truncations or controls.

1157

Figure 5. Mpt64 localizes to the endoplasmic reticulum during heterologous expression

in yeast and HeLa cells. (**A**) Immunofluorescence (top panels) and bright field overlay

(bottom panels) images of *S. cerevisiae* transformed with GFP fusion proteins to Mpt64

truncations. Scale bars are 3 µm. (**B**) HeLa cells transfected overnight with GFP fusion

proteins (green) and stained for ER localization with anti-calreticulin antibody (red).

Nuclei are stained with DAPI (blue). Scale bars are 10 µm.

1164

Figure 6. Mpt64 ER localization is ESX1-dependant during *M. tuberculosis* infection of

macrophages. (**A**) RAW267.4 murine macrophages were infected with mCherry-

expressing (red) WT (upper panels) or MtbΔeccD1 (lower panels) for 4 hours at an MOI

20:1. Cells were fixed and stained for Mpt64 (green) and nuclei (blue). Scale bars are

5µm. (**B**) Schematic detailing in-frame deletion of *mpt64* by insertion of a hygromycin

resistance gene. (**C**) Western blot detecting expression of Mpt64, Ag85 and GroEL2 in

either the lysate of the cell pellet (P) or culture supernatant (S) of four Mtb strains. (**D**)

RAW267.4 macrophages were infected with the indicated strains of mCherry-

expressing (cyan) Mtb for 4 hours at an MOI 20:1. Cells were fixed and stained for

Mpt64 (red), calreticulin (green) and nuclei (blue). Images in (**E**) and (**F**) correspond to

box 1 and box 2, respectively. Scale bars are 10µm. (**E**) Enlarged image from box 1 in

(**D**) of an Mtb-infected macrophage stained for Mpt64, calreticulin and DAPI. Insets

show an area of Mpt64-calreticulin co-localization. Scale bars are 5µm. (**F**) Enlarged

image from box 2 in (**D**) of macrophages infected with MtbΔMpt64::Mpt64-NS and

stained for Mpt64, calreticulin and DAPI. Insets show Mpt64 localization in relation to

bacteria. Scale bars are 5 µm.

1181

Figure 7. Mpt64 contributes to early Mtb growth after aerosol infection of mice (**A**)

Bacterial burden in lungs of mice 21 and 42 days after aerosol infection with indicated

strains of Mtb. Results are a combination of three independent experiments, n=25 mice

total per group. Horizontal bar indicates the geometric mean. p values are determined

by nonparametric Kruskal-Wallis analysis. (**B**) Representative images of H&E stained

lungs at 42 days post-infection with the indicated strains of Mtb. Scale bars, 2mm. (**C**)

Ten mice per group were monitored for survival. There were no significant differences in

survival rates between groups by Kaplan–Meier analysis. (**D**) Quantitation of lung

inflammation of mice infected with indicated Mtb strains. Measurement was determined

using ImageJ software (NIH). Bars are colored as in (**A**). Results are the mean ±SEM

for three animals per group. *p<0.02 by Kruskal-Wallis test.

1193

Figure 8. Localization and activity of Mpt64 in human macrophages (**A**) Primary human

monocyte-derived macrophages were infected with mCherry-expressing (red) Mtb or

MtbΔeccD1 for 4 hours at an MOI 10:1. Cells were fixed and stained for Mpt64 (green).

Scale bars are 5µm. (**B**) Primary human monocyte-derived macrophages were infected

with the indicated strains of mCherry-expressing Mtb for 4 hours at an MOI 10:1. Cells

were fixed and stained for Mpt64 (red) and calreticulin (green). Nuclei are stained blue.

Scale bars are 5 µm and scale bars of inset images are 2.5 µm. (**C**) CFU recovered

from primary human macrophages from two separate donors infected with Mtb,

MtbΔMpt64, or MtbΔMpt64::Mpt64 at indicated time points. CFU was normalized to the

bacterial burden at day 0 and is represented as the mean with standard error. Data are

representative of 6 independent experiments.

1205

Supplemental Figure 1. Recombinant Mpt64 does not interact with lysozyme. (**A**)

Sequence and secondary structure alignment of Mpt64 and *B. subtilis* RsiV (template)

performed by Phyre^2 software (http://www.sbg.bio.ic.ac.uk/phyre2/). The Mpt64

DUF3298 is indicated by a red line above the Mpt64 amino acid numbering. (**B**) Mpt64

(M) or an unrelated protein Cor (C) were immobilized on cobalt beads and hen egg

white (HEW) or human lysozyme (HuLYZ) was incubated with these or beads alone (B)

for five minutes. After washes, proteins were eluted with 300mM imidazole.

1213

Supplemental Figure 2. Construction and phenotypic analysis of MtbΔMpt64. (**A**)

Western blot detecting expression of Mpt64, Ag85 and GroEL2 in either the lysate of the

cell pellet (P) or culture supernatant (S) of wild type Mtb or MtbΔeccD1. (**B**) Detection of

hygromycin resistance cassette insertion in place of *mpt64.* Genomic DNA from wild

type Mtb or MtbΔMpt64 was amplified by polymerase chain reaction and products were

analyzed by agarose gel electrophoresis. (**C**) Growth of Mtb, MtbΔMpt64,

MtbΔMpt64::Mpt64 and MtbΔMpt64::Mpt64-NS in 7H9 measured by optical density at

600nm. (**D**) PDIM standard or apolar lipid extracts from Mtb or MtbΔMpt64 were

analyzed on an AbSciex TripleTOF 5600/5600+ mass spectrometer.

1223

1224   Supplemental Figure 3. Secreted Mpt64 co-localizes with calreticulin in murine

1225   macrophages. (A) RAW267.4 murine macrophages were infected with the indicated

1226   strains of mCherry expressing Mtb (cyan) for four hours and subsequently stained for

1227   Mpt64 (red) and calreticulin (green). Nuclei are stained in blue. Scale bars are 10μm.

1228

1229   Supplemental Figure 4. Secreted Mpt64 co-localizes with calreticulin in human

1230   macrophages. (A) Primary human macrophages were infected with the indicated strains

1231   of mCherry expressing Mtb (cyan) or left uninfected for four hours prior to fixation and

1232   staining for Mpt64 (red) and calreticulin (green). Scale bars are 5µm.

Figure 1. M. tuberculosis exported proteins interact with yeast membranes. (A) Ras rescue assay schematic. (B) S. cerevisiae (cdc25ts) transformed with Mtb protein fusions to Rasmut, duplicate plated and incubated 48-72 h at the permissive (25°C) and restrictive (37°C) temperatures. Shown are representative images of 20 yeast strains. (C) Summary results of Ras rescue screen. (D) Western blot of lysates from yeast transformed with the indicated fusion proteins and probed with anti-Ras or anti-G6PDH antibodies. Rasmut fusion proteins are marked by white or black asterisks. PLC (phospholipase C) and DYN (dynamin) are fusions to Rasmut known to be membrane associated (PLC) or cytoplasmic (DYN). (E) Representative fluorescence microscopy of S. cerevisiae (INVSc1) transformed with GFP-MEP fusion proteins. Scale bars are 3μm.

Figure 2. Host subcellular localization of membrane-binding MEP. (A) Representative fluorescent images of HeLa cells transfected with the indicated GFP-MEP fusion proteins (green) and stained with antibodies (red) to calreticulin (ER), GM-130 (Golgi), Tom20 (Mitochondria) or PMP70 (Peroxisomes). Scale bars are 5μm. (B) Comparison of the organelle localization of MEP expressed in yeast and HeLa cells.

Figure 3. M. tuberculosis exported proteins alter hGH secretion (A) Inducible secretion assay schematic. (B) Supernatant hGH ELISA from HeLa cells transfected overnight with hGH-CAD and either GFP or GFP-MEP fusion proteins prior to addition of drug to allow for hGH secretion. hGH secretion by GFP-MEP transfected cells was normalized to hGH secretion of cells transfected with GFP alone. (C) Summary of results from inducible secretion screen. (D) Venn diagram of MEP that are membrane-localized, alter host secretion, or both. (E) Table summarizing the membrane localization and degree of hGH secretion in cells transfected with the five overlapping proteins from D.

Figure 4. hGH secretion inhibition is dependent on membrane localization of Mpt64. (A) Solution structure of Mpt64. PDB 2hhi. (B) Schematic of Mpt64 truncations, colored to match solution structure in A. SP, signal peptide. (C) Full length Mpt64 or protein truncations expressed in the Ras rescue assay. (D) Western blot of lysates from cdc25ts yeast expressing Rasmut fusion proteins to Mpt64 or Mpt64 truncations. Blots were probed for anti-Ras or anti-G6PDH antibodies. Control protein bands are marked by a white asterisk and Mpt64 truncations are marked by a black asterisk. (E) PIP membrane strips incubated with recombinant Mpt64_24-228 or Mpt64_24-143. Binding of Mpt64 to lipids was detected by incubation with α-Mpt64 or pre-immune serum. Numbers indicate specific lipids as follows: 1, lysophosphatidic acid; 2, lysophosphatidylcholine; 3, phosphatidylinositol; 4, phosphatidylinositol-3-phopshate (PI3P); 5, PI4P; 6, PI5P; 7, phosphatidylethanolamine; 8, phosphatidylcholine; 9, sphingosine 1-phosophate; 10, phosphatidylinositol-3,4-bisphosphate [PI(3,4)P2]; 11, PI(3,5)P2; 12, PI(4,5)P2; 13, phosphatidylinositol-3,4,5-trisphosphate [PI(3,4,5)P3]; 14, phosphatidic acid; 15, phosphatidylserine; 16, blank. (F) ELISA results of hGH in supernatants of cells co-expressing full length Mpt64, Mpt64 truncations or controls.

Figure 5. Mpt64 localizes to the endoplasmic reticulum during heterologous expression in yeast and HeLa cells. (A) Immunofluorescence (top panels) and bright field overlay (bottom panels) images of S. cerevisiae transformed with GFP fusion proteins to Mpt64 truncations. Scale bars are 3 μm. (B) HeLa cells transfected overnight with GFP fusion proteins (green) and stained for ER localization with anti-calreticulin antibody (red). Nuclei are stained with DAPI (blue). Scale bars are 10 μm.

Figure 6. Mpt64 ER localization is ESX1-dependant during M. tuberculosis infection of macrophages. (A) RAW267.4 murine macrophages were infected with mCherry-expressing (red) WT (upper panels) or MtbΔeccD1 (lower panels) for 4 hours at an MOI 20:1. Cells were fixed and stained for Mpt64 (green) and nuclei (blue). Scale bars are 5μm. (B) Schematic detailing in-frame deletion of mpt64 by insertion of a hygromycin resistance gene. (C) Western blot detecting expression of Mpt64, Ag85 and GroEL2 in either the lysate of the cell pellet (P) or culture supernatant (S) of four Mtb strains. (D) RAW267.4 macrophages were infected with the indicated strains of mCherry-expressing (cyan) Mtb for 4 hours at an MOI 20:1. Cells were fixed and stained for Mpt64 (red), calreticulin (green) and nuclei (blue). Images in (E) and (F) correspond to box 1 and box 2, respectively. Scale bars are 10μm. (E) Enlarged image from box 1 in (D) of an Mtb-infected macrophage stained for Mpt64, calreticulin and DAPI. Insets show an area of Mpt64-calreticulin co-localization. Scale bars are 5μm. (F) Enlarged image from box 2 in (D) of macrophages infected with MtbΔMpt64::Mpt64-NS and stained for Mpt64, calreticulin and DAPI. Scale bars are 5 μm.

Figure 7. Mpt64 contributes to early Mtb growth after aerosol infection of mice (A) Bacterial burden in lungs of mice 21 and 42 days after aerosol infection with indicated strains of Mtb. Results are a combination of three independent experiments, n=25 mice total per group. Horizontal bar indicates the geometric mean. p values are determined by nonparametric Kruskal-Wallis analysis. (B) Representative images of H&E stained lungs at 42 days post-infection with the indicated strains of Mtb. Scale bars, 2mm. (C) Ten mice per group were monitored for survival. There were no significant differences in survival rates between groups by Kaplan–Meier analysis. (D) Quantitation of lung inflammation of mice infected with indicated Mtb strains. Measurement was determined using ImageJ software (NIH). Bars are colored as in (A). Results are the mean ±SEM for three animals per group. *p<0.02 by Kruskal-Wallis test.

Figure 8

Figure 8. Localization and activity of Mpt64 in human macrophages (A) Primary human monocyte-derived macrophages were infected with mCherry-expressing (red) Mtb or MtbΔeccD1 for 4 hours at an MOI 10:1. Cells were fixed and stained for Mpt64 (green). Scale bars are 5μm. (B) Primary human monocyte-derived macrophages were infected with the indicated strains of mCherry-expressing Mtb for 4 hours at an MOI 10:1. Cells were fixed and stained for Mpt64 (red) and calreticulin (green). Nuclei are stained blue. Scale bars are 5 μm and scale bars of inset images are 2.5 μm. (C) CFU recovered from primary human macrophages from two separate donors infected with Mtb, MtbΔMpt64, or MtbΔMpt64::Mpt64 at indicated time points. CFU was normalized to the bacterial burden at day 0 and is represented as the mean with standard error. Data are representative of 6 independent experiments.