MODELLING THE IMPACT OF DISPERSAL ON A NATAL POPULATION THAT EXHIBITS BOOM-

BUST DYNAMICS

Jacques A. Deere[1,2]*, Ilona van den Berg[3], Gregory Roth[1,4], Isabel M. Smallegange[1]

[1] Institute for Biodiversity and Ecosystem Dynamics (IBED), University of Amsterdam,

P.O. Box 94240, 1090 GE Amsterdam, The Netherlands

[2] Department of Zoology, University of Oxford, Oxford OX1 3SZ, UK

[3] Vrije Universiteit, Faculty of Earth and Life Sciences, Department of Animal Ecology, The

Netherlands

[4] Friedrich Miescher Institute for Biomedical Research, P. O. Box 3775, 4002, Basel

*Correspondence:

Department of Zoology, University of Oxford, Zoology Research and Administrative Building,

11a Mansfield Road, Oxford OX1 3SZ, UK

E-mail addresses:

Jacques A. Deere (corresponding author):   jacques.deere@zoo.ox.ac.uk

Ilona van den Berg: ilonavdberg@hotmail.com

Gregory Roth: gregory.roth@fmi.ch

Isabel M. Smallegange: i.smallegange@uva.nl

Keywords: Life history, Integral Projection Model, bulb mite, *Rhizoglyphus robini*, unsuccessful

dispersal

Abstract

Dispersal, an important form of movement, influences population dynamics, species distribution, and gene flow between populations. In population models, dispersal is often included in a simplified manner by removing a random proportion of the population. Many ecologists now argue that models should be formulated at the individual-level instead of the population-level to accurately capture how dispersal affects population dynamics. This is especially true for populations exhibiting boom-bust dynamics as these often harbour specialised disperser morphs, life histories or behaviours, as dispersal is essential for persistence. Within a management context, such dynamics play a key role in the stability of populations and in turn extinction risk of species. Here we parameterised an integral projection model, which allows studying how individual life histories determine population-level processes. Using bulb mites (*Rhizoglyphus robini*), a species that shows boom-bust dynamics, we assess the extent dispersal expression (frequency of disperser morphs) and dispersal probability (probability to emigrate) affect the proportion of dispersers and natal population growth rate. We find that, if residents and dispersers differ in life history, different combinations of dispersal probability and dispersal expression produce the same proportion of leaving dispersers. Additionally, for a given proportion of dispersing individuals different natal population growth rates occur. Dispersal life histories, and frequency of disperser morphs occurring in the natal population, can thus significantly affect population-level processes. It is therefore important for our understanding of boom-bust dynamics to elucidate how dispersal characteristics of individuals relate to population resilience and potential re-establishment for populations after a bust phase.

Introduction

The movement of individuals is one of the key mechanisms shaping biodiversity (Travis and Dytham 1999; Jeltsch et al. 2013). An important form of movement is dispersal, which is any movement of individuals or propagules with potential for gene flow across space (Ronce 2007; Bonte et al. 2012). Dispersal influences the dynamics and persistence of populations, distribution and abundance of species, community structure, and the level of gene flow between populations (Brown and Kodric-Brown 1977; Dieckmann et al. 1999; Hanski 1999; Bowler and Benton 2005). In so doing, dispersal can fuel evolutionary processes such as local adaptation and speciation (Dieckmann et al. 1999). Currently, understanding dispersal behaviour is important to be able to predict how populations will respond to some of the most important threats to biodiversity, such as climate change, habitat loss and fragmentation, and the invasion of alien species (Bowler and Benton 2005; Clobert et al. 2009).

In the last decades, the drivers of dispersal have been the centre of many theoretical studies (Hamilton and May 1977; Johnson and Gaines 1990; Hanski 1999; Clobert et al. 2004). However, the dispersal process itself has gained less attention. Due to practical problems associated with the study of dispersal in the field, much of dispersal research has taken a theoretical approach (Bélichon et al. 1996; Bowler and Benton 2005); although there has been recent advancement in empirical field studies (Singh et al. 2012; Maag et al. 2018). In population models, dispersal is often considered as a population-level process (Clobert et al. 2004; Bowler and Benton 2005). In such models, individual-level differences are often ignored, which means that any individual within the population has the same probability of dispersing successfully. However, recently it has been argued that population ecology should shift its focus from formulating models at the population level to the level of individual organisms (Clark et al. 2011; Topping et al. 2015; Soudijn and Roos 2017); an argument echoed in the field of dispersal (Bonte et al. 2012). Although ecological patterns can be observed at the population level, it is the behaviour and demographic changes of individuals that shape the dynamics of the population (Clark et al. 2011; Soudijn and Roos 2017). So, in order to fully understand the effects of dispersal on natural systems, it is necessary to incorporate individual-level differences in dispersal behaviour in population models.

Empirical studies have shown that dispersing individuals typically have different demographic properties compared with residents (individuals that remain in the population) (Harrison 1980; Bélichon et al. 1996; O'Riain et al. 1996; Lemel et al. 1997; Zera and Denno 1997; Deere et al. 2015). Dispersers and residents can differ in their morphology, physiology or in their behaviour (Bélichon et al. 1996; Clobert et al. 2009; Bonte et al. 2012). In many species, an individual property, e.g. locomotory movement, which potentially serves many important functions, is enhanced for dispersal (Phillips et al. 2006; Bonte et al. 2012; Cozzi et al. 2018). In other species, dispersers develop a morphology primarily as an adaptation for dispersal, e.g. the development of wings or structures which facilitate attachment to mobile vectors (Harrison 1980; Zera and Denno 1997; Diaz et al. 2000; Bonte et al. 2012). Species with a distinct disperser morphology are ideal study systems to investigate the individual-level effects and adaptive significance of dispersal in natural populations. For many of these species, each genotype has the potential to develop into a disperser or a resident (Harrison 1980; Clobert et al. 2004). The investment in a disperser morphology is costly if it requires resource investment, likely at the expense of investment into body condition or fecundity (Lemel et al. 1997; Bonte et al. 2012; Deere et al. 2015). Such investment costs, defined as "pre-departure costs", arise during development prior to the actual dispersal event, although they may be deferred to later in the life-history as well (Bonte et al. 2012). However, not all individuals that invest in dispersal morphology leave their natal population, e.g. due to a behaviour or physiological component affecting dispersal propensity (Roff and Fairbairn, 2001), reduced patch accessibility (Clobert et al. 2001) or environmental conditions (Johnson 1969). As such, it is important to understand how individuals that invest in dispersal but do not leave natal populations impact population dynamics. This will be especially true in populations that experience boom-bust dynamics. Populations exhibiting boom-bust dynamics predominantly occur in short-lived species (Gamelon et al. 2016) and often in invasive species (Simberloff & Gibbons 2004, Lester & Gruber 2016, Strayer et al. 2017). In addition, these populations are oftentimes isolated and fragmented (Arthington and Balcombe 2011) and so dispersal plays a key role in the long-term persistence of these populations (Boudjemadi et al. 1999; Clobert et al. 2001). Boom-bust dynamics involve an initial outbreak, where populations become large (boom phase), followed by a decline in the population to a lower size (bust phase) (Strayer et al. 2017). Within a management context,

such dynamics will play a key role in the stability of populations and in turn the extinction risk
of species. Indeed, it has been shown that management strategies will differ depending on
the mechanism that produces boom-bust dynamics, so understanding the population
dynamics of species exhibiting large population fluctuation need to be prioritized (Himes Boor
et al. 2018). To incorporate the dispersal process in boom-bust population models in a way
that is more realistic, and its results more biologically relevant, an individual-based method
that takes distinct dispersal life-histories and (un)successful dispersal into account is
necessary. As a first step towards such an approach we focus on the natal population, where
the dispersal process is initiated, and how dispersing individuals impact its dynamics. We
apply the model to the invasive bulb mite (*Rhizoglyphus robini*, Claparède), the bulb mite is
an ideal model system to study dispersal as it has a distinct dispersal stage within its life-
history and can exhibit boom-bust dynamics.

We model the system using an Integral Projection Model (IPM) because such models
comprise individual-level functions that describe demographic rates (thereby tracking
fluctuations in population size and structure), they can be applied to species with complex
demography, require relatively straightforward mathematical techniques from matrix
calculus, and are closely and easily linked to field and experimental data (Easterling et al.
2000; Ellner and Rees 2006). To test our approach, we parameterise an IPM that incorporates
a distinct dispersal life-history, where a dispersal stage is included in the life-cycle, and that
incorporates an emigration component (where dispersers are able to leave the population).
Within the IPM, each individual has the potential to develop into a *disperser* (i.e. develop into
the dispersal stage) or a *resident* (i.e. do not develop into a dispersal stage), and each
disperser has the option to leave the natal population (*successful disperser*) or remain in the
natal population (*unsuccessful disperser*) (Fig. 1). The natal population will thus consist of
residents and unsuccessful dispersers. In this model the individual-level costs of dispersal are
then expressed by dispersers. It is important to note that unsuccessful dispersers in the
context of this study refers to disperser individuals that do not emigrate from the population;
not to disperser individuals that do not survive the transfer phase or fail to establish in a new
habitat. To address our aim of understanding how distinct dispersal life histories play a role
in natal population-level processes during a dispersal event, requires knowing not only the
proportion of residents and dispersers in the natal population, but also the proportion of

successful dispersers. We will quantify the effects of the expression of a distinct disperser

phenotype (frequency of individuals developing into dispersers) and dispersal probability

(probability that dispersing individuals emigrate) on the proportion of successful dispersers

(i.e. the ones that emigrate) and on the growth rate of the natal population (Fig. 1). We then

discuss the potential impact of the outcomes on populations that show boom-bust dynamics.

Methods

Study system and life-history data

Bulb mites consist of five life stages, but develop an additional, sixth dispersal stage

during development (called the deutonymph stage) under unfavourable environmental

conditions (e.g. low temperature, humidity, food quality) (Diaz et al. 2000) (Fig. 2). The

(juvenile) dispersal stage is non-feeding and occurs in both sexes. Bulb mite populations

typically show exponential growth until resources are depleted, after which populations

crash. During this time, as environmental conditions become less favourable, the frequency

of individuals developing into dispersers (i.e. dispersal stage) increases. It is the impact of (i)

the frequency of the dispersal stage, and (ii) the dispersal probability, on the population

growth that we are interested in, and so we only focus on changes in population growth rate

with regard to these two parameters.

We use the same life-history data on female bulb mites as used in Deere et al. (2017)

to parameterise the IPM; these data provide information on individuals that do and do not

develop into the deutonymph stage as life histories differ between individuals that went

through the deutonymph stage and those that did not (for detailed data collection see Deere

et al. (2015). Data can be found in the figshare repository 10.6084/m9.figshare.1312875).

Disperser Phenotype Model (DPM)

Our model has a distinct dispersal stage, the deutonymph stage, which allows for

emigration (henceforth DPM). Dispersal is a two-step process that consists of development

into a disperser (deutonymph) and dispersal out of the population (Fig. 2). Deutonymph

frequency, β, describes the transition probability of developing from a protonymph to a

deutonymph. Dispersal probability, δ, describes the probability of a deutonymph leaving the

population (Fig. 2).

We use the size- and stage- structured IPM developed by Deere et al. (2017) and we

include an emigration component to the model. Briefly, an IPM tracks fluctuations in

population size and structure based on individual-level processes. The IPM projects a new

stage-size joint distribution based on the stage-size joint distribution in the previous time step

(Easterling et al. 2000; Coulson 2012; Deere et al. 2017). For the DPM, in each time step

individuals may survive, disperse, grow and produce new individuals. These processes are

captured in a kernel, *K*, that projects joint distribution $n_t(z,s)$ of body size *z*, and stage *s*, at

time *t*, to the new joint distribution at time *t*+1:

$$n_{t+1}(z',s') = \sum_{s=1}^{8} \int_{\Omega} K(z',s',z,s) n_t(z,s) dz \qquad (1)$$

where Ω designates the range of individual sizes. The kernel, K, is composed of two parts,

$$K = S(z,s)D(z,s)T(s',z,s)G(z',z,s) + R(z,s)O(z',z,s) \qquad (2)$$

The first part describes how individuals grow, survive and move: S(*z*,*s*) is the survival

probability to the next time step of an individual in size *z* and stage *s*; D(*z*,*s*) is the probability

that an individual of size *z* in stage *s* stays in the population in the next time step; T(*s'*,*z*,*s*) is

the probability that an individual in size *z* and stage *s* develops into the new stage *s'* and

G(*z'*,*z*,*s*) is the probability that an individual in size *z* and stage *s* growths to size *z'*. The second

part describes adult reproduction: R(*z*,*s*) is the number of offspring produced by an adult

individual of size *z* in stage *s* and O(*z'*,*z*,*s*) is the probability that offspring produced by adults

of size *z* are of size *z'* (i.e. the parent-offspring association). DPM equations (Table A1

equations 3.1 to 3.12), and parameter values used, can be found in the appendix.

The order of the functions in the kernel K represents the sequence of events in each

time step, and the point at which the population is censused. This order is crucial for the

outcome of the IPM (Rees et al. 2014). The order of events in the DPM is first survival, then

dispersal, then transition and lastly growth. Dispersal is applied prior to transition in order to

track all individuals that develop into the deutonymph stage (the dispersers). Since dispersal

is only possible for individuals in the deutonymph stage (i.e. dispersal stage), the probability

to stay in the population is equal to 1 for all individuals in all stages other than the

deutonymph stage. For all individuals in the deutonymph stage, the probability to stay in the

population is $1 - \delta$.

Models parameterization and analysis

The DPM was parameterised from life-history data of female bulb mites (see *Study*

system and life-history data). Once parameterised, each function is then discretised into a

matrix form by dividing the full size domain into very small-width discrete bins ('mesh points';

see appendix for details). These discretised matrices describe the predicted transition rates

and are used to build a projection matrix that approximates the DPM, so standard methods

for analysing matrix population models can be used (Caswell 2001; Coulson 2012). Details on

how the functions were parameterized can be found in the appendix. All analyses and

simulations were performed in R version 3.0.2 (R Development Core Team 2013).

Combination of disperser expression and dispersal probability

For the DPM we wanted to assess how an increase in disperser expression (β) and

dispersal probability (δ) would affect population-level processes. Within the DPM, β and δ

were increased from 0-1 (at 0.01 increments) and for every combination of β and δ the

proportion of individuals that disperse and population growth rate, λ_0, were calculated. The

proportion of individuals that disperse was calculated by integrating the stable stage

distribution over the range of all sizes in the deutonymph stage and multiplying it with the

dispersal probability δ. The stable stage distribution was calculated as the dominant right

eigenvector of the projection matrix associated with each DPM (Easterling et al. 2000), λ_0 was

calculated as the dominant eigenvalue of the projection matrix associated to each DPM

(Easterling et al. 2000).

Results

When varying β and δ in the DPM there is a non-linear, joint effect of dispersal

expression and dispersal probability on the proportion of dispersing individuals (Fig. 3) and

population growth rate (λ_0) (Fig. 4). If both β and δ increase then, in general, the proportion

of dispersing individuals increases and λ_0 decreases. Reciprocally, if both β and δ decrease

then, in general, the proportion of dispersing individuals decreases and λ_0 increases. When

the changes in β and δ are opposed, any scenario (increase/decrease) in the proportion of

dispersing individuals and λ_0 can occur (Table 1).

The joint effect of β and δ on the proportion of dispersers and λ_0 is not proportional.

As a consequence, for a fixed proportion of dispersing individuals, multiple λ_0 values can occur

(proportion of dispersing individuals: 0.04 (yellow line), 0.08 (red line), 0.12 (blue line); Fig. 3

& 4). Furthermore, with increasing proportions of dispersing individuals (blue line > red line >

yellow line), the values of λ_0 encompassed by each proportion of dispersing individuals,

decreases (Fig. 4). Another consequence of the joint effect of β and δ is unexpected scenarios

in the proportion of dispersing individuals and λ_0 (grey dots in Table 1). For example, if β

increases and δ decreases (in some specific proportion), then both the proportion of

dispersing individuals and λ_0 can decrease. Vice versa, if β decreases and δ increases (in some

specific proportion), then both the proportion of dispersing individuals and λ_0 can increase.

Discussion

The need to incorporate complexity and realism in modelling dispersal (Bonte et al.

2012; Travis et al. 2012; Travis et al. 2013) has prompted a move to incorporate more

individual heterogeneity and individual-level costs when modelling dispersal (Bonte et al.

2012). Here we incorporated individual heterogeneity, in terms of distinct life-histories for

dispersing and non-dispersing individuals, as well as the probability of dispersing individuals

to emigrate using an individual-based method. We applied our method to a system that

shows boom-bust dynamics and assessed the effects of these distinct life-histories and

emigration probability on population-level processes.

When modelling dispersal, dispersal rates are often applied to the whole population

and the potential effect of dispersal to the natal population is then in terms of the proportion

of the individuals that leave. Given that the proportion of individuals that leave the population

during a dispersal event is partly dependent on the number of disperser individuals within the

population, we wanted to investigate the effect on the population of jointly manipulating the

proportion of the dispersers in the population (β) and the dispersal probability (δ). Although

we found a general trend for proportion of dispersers to increase (or decrease) and

population growth rate to decrease (or increase) when both dispersal parameters increase

(or decrease), when the two parameters increase or decrease in *opposite* directions, a

number of different outcomes of the population parameters can occur (Table 1). The

outcomes include scenarios which one would largely expect; an increase in the proportion of

dispersers and a decrease in population growth (or vice versa). For example, when the

proportion of dispersers increase, population growth rate will decrease as these individuals

leave the population and so do not contribute to the population growth. However, there are

also outcomes that we did not expect; scenarios where both the proportion of dispersers and

population growth rate decrease or increase together. The scenario where the proportion of

dispersers and population growth rate decrease occurs when β increases and δ decreases. An

increase in the expression of deutonymphs results in more disperser individuals within the

population, however, with lower dispersal probabilities, this would mean a reduced

proportion of dispersers that emigrate. With fewer dispersing individuals leaving the

population, the individual-level costs of investing in dispersal (i.e. reduced size at maturity,

reduced lifetime egg production, increased development time) are then borne out at the

population level, thereby reducing population growth rate (Deere et al. 2017). The scenario

where both the proportion of dispersers and population growth rate increase, occurs when β

decreases and δ increases. Here, reducing β reduces the expression of deutonymphs within

the population, whereas increasing δ results in a larger proportion of the dispersing

individuals leaving the population (i.e. an increase in the proportion of dispersers). Population

growth rate will increase as fewer dispersing individuals, with their associated individual-level

costs of investing in dispersal, are left in the population and have a reduced effect on

population growth rate. In all scenarios, expected and unexpected, the response is largely

dependent on the ratio of the two dispersal parameters, highlighting that the population

parameters are very responsive to the joint effect of the two dispersal parameters. This

outcome highlights that to fully understand the influence of dispersing individuals on natal

population processes requires a detailed understanding of disperser life-histories and how

the cost of investing in a disperser strategy is borne out at the population level, and to what

extent dispersing individuals are present in the population.

Another key finding is that different population structures (multiple combinations of

β and δ) can give the same proportion of disperser individuals (Fig. 3); this is significant

because population structure can have a large effect on population growth (Cameron et al.

2016; Smallegange et al. 2018). Previously, we have shown that, in the absence of emigration,

distinct dispersal phenotypes within the natal population have different life-histories to non-

dispersers and so do not contribute to population processes in the same way (Deere et al.

2017). Additionally, juveniles and adults do not always respond the same way to changing

environmental and population conditions and so differ in their contribution to populations

dynamics (Coulson et al. 2001; Ozgul et al. 2012). Furthermore, the influence that the

frequency of dispersing individuals within the natal population and dispersal probability has

on the proportion of dispersers can be seen as a natal-habitat induced effect. In this case the

effects of the natal habitat on disperser phenotypes is clear, the expression of dispersing

individuals is dependent on the quality of the natal habitat; however in other instances the

effects may often be more subtle (e.g. maternal effects) (Benard and McCauley 2008). This

effect on the proportion of dispersers can then further influence other processes such as

genetic differentiation between populations (i.e. altering magnitude and/or symmetry of

gene flow) (Benard and McCauley 2008). Indeed, Benard & McCauley (2008) highlight that

environmentally induced asymmetries in the number or quality of dispersing individuals can

lead to asymmetry in patterns of local adaptation. While our models are largely comparable

to phoretic species, our findings are also relevant to actively dispersing species. In species

showing active dispersal, dispersal is often dictated by a combination of dispersal ability (e.g.

structural formation such as wings) and dispersal propensity (e.g. behaviour or physiological

component), which would influence the proportion of dispersers. Moreover, it has been

suggested that interactions between these traits can generate non-linear relationships

between habitat condition and net dispersal rates (Benard and McCauley 2008). All in all, it is

the combination of natal-habitat induced effects on dispersal phenotypes and the population

structure that will determine the proportion of dispersing individuals.

In species that experience large population fluctuations due to environmental

variation, isolated populations experiencing large variation in demographic parameters face

some risk of local extinction. Increased dispersal from a population in a bust phase can lead

to reduced population growth (low λ_0; top right, Fig. 4), so too can a population with high

numbers of dispersers that fail to disperse (low values of proportion dispersing; bottom right,

Fig. 3), in both cases this increases the risk of population extinction. In the latter case this

would be an unexpected outcome (grey dots, Table 1), suggesting that the outcome of a

population in a bust phase, where dispersal is able to occur, will not only be dependent on

the structure of the population, but also on individuals being able to disperse. When the

proportion of dispersing individuals is low, this suggests fewer individuals leaving the

population. In our system, the low proportion of dispersers would not be due to *fewer dispersers* in the population but due to the *reduced probability of dispersers* leaving the population. Such scenarios would be seen in populations where fewer hosts are available for dispersal (e.g. in the case of phoretic species), so too for isolated populations in fragmented landscapes where individuals are prevented to disperse by lack of available habitat. Previous work has shown that a population's recovery potential and changes in key-life history traits (e.g. earlier maturation, increased fecundity) after a bust period will dictate population resilience and re-establishment after environmental variation (e.g. disturbance; depressed resources or pulses of resources) (Ramskov & Forbes 2008, Jacimovic et al 2019). However, we would argue that in species that disperse, disperser propensity (β) and the ability to then disperse (δ) are key traits that would also influence population resilience and potential re-establishment. Indeed, the increase in dispersers within a population reduces reproductive output and together with the additional increase in generation time (Deere at al. 2017) impacts population density negatively. This is then pertinent when considering populations experiencing extreme environmental perturbations that remain in an extremely low state for multiple generations, as such populations are prone to extinction due to stochastic fade-out (Golinski et al. 2008). Understanding these dynamics are key to ensure successful conservation of species that face boom-bust dynamics. Moreover, a better understanding of invasive or pest species showing boom-bust dynamics would aid successful pest management strategies. Management strategies can be implemented to take advantage of a species' dispersal traits when populations are in a bust phase. For example, by reducing the availability of hosts of phoretic pest species reduces the probability of dispersal and increases the negative impact on population growth due to individual-level costs of dispersers that remain in the populations.

The importance of the joint effect of dispersal probability and the frequency of dispersing individuals within the population that we show, indicates the potential importance of accounting for the frequency of dispersing individuals within a population. We appreciate that our focus is on a specific study system, and acknowledge that our models may not generalise beyond arthropod boom-bust life histories. However, our results illustrate that the effects of dispersing individuals on natal populations are more than just a turnover of numbers. Given the importance of dispersal rates in populations, and the fact that natural

dispersal rates are being altered by human activities, the way in which dispersal rates are

applied in population models then becomes vital as models inform conservation management

decisions. Modelling the dynamics of populations is by its very nature a simplified version of

reality. However, as more complexity is added to models in a manner that improves the

biological realism of how dispersal rates are implemented, then not only can model

performance be improved in terms of their realism, but also in terms of the accuracy of their

predictions, which ultimately is more informative when used in management decisions

(Clobert et al. 2001; Peterson and Freeman 2016). Only by identifying and including the

individual-level costs of dispersing individuals in a population, and applying dispersal rates to

those individuals, can the full extent of the effects of dispersal to natal population dynamics

start to be realised.

Acknowledgements

The work was funded by a MEERVOUD grant no. 836.13.001 and VIDI grant no. 864.13.005

from the Netherlands Organisation for Scientific Research to Isabel Smallegange.

Conflict of Interest

None

Author Contributions

JAD and IvdB conceptualized the study. JAD, IvdB and GR developed the models with input

from IMS. JAD, IvdB, GR and IMS wrote and commented on the manuscript.

References

Arthington, A. H., and S. R. Balcombe. 2011. Extreme flow variability and the 'boom and bust'
ecology of fish in arid-zone floodplain rivers: a case history with implications for
environmental flows, conservation and management. Ecohydrology 4:708–720.

Bélichon, S., J. Clobert, and M. Massot. 1996. Are there differences in fitness components
between philopatric and dispersing individuals? Acta Oecologia 17:503–517.

Benard, M. F., and S. J. McCauley. 2008. Integrating across Life-History Stages: Consequences
of Natal Habitat Effects on Dispersal. The American Naturalist 171:553–567.

Bonte, D., H. Van Dyck, J. M. Bullock, A. Coulon, M. Delgado, M. Gibbs, V. Lehouck, et al. 2012.
Costs of dispersal. Biological Reviews 87:290–312.

Boudjemadi, K., J. Lecomte, and J. Clobert. 1999. Influence of connectivity on demography
and dispersal in two contrasting habitats: an experimental approach. Journal of Animal
Ecology 68:1207–1224.

Bowler, D. E., and T. G. Benton. 2005. Causes and consequences of animal dispersal strategies:
relating individual behaviour to spatial dynamics. Biological Reviews 80:205–225.

Brown, J. H., and A. Kodric-Brown. 1977. Turnover Rates in Insular Biogeography: Effect of
Immigration on Extinction. Ecology 58:445–449.

Cameron, T. C., D. O'Sullivan, A. Reynolds, J. P. Hicks, S. B. Piertney, and T. G. Benton. 2016.
Harvested populations are more variable only in more variable environments. Ecology
and Evolution n/a-n/a.

Caswell, H. 2001. Matrix population models: Construction, analysis, and interpretation
(Second.). Sinauer Associates, Sunderland, Massachusetts.

Clark, J. S., D. M. Bell, M. H. Hersh, M. C. Kwit, E. Moran, C. Salk, A. Stine, et al. 2011. Individual-
scale variation, species-scale differences: inference needed to understand diversity.
Ecology Letters 14:1273–1287.

Clobert, J., E. Danchin, A. A. Dhondt, and J. D. Nichols, eds. 2001. Dispersal. Oxford University
Press, New York.

Clobert, J., R. A. Ims, and F. Rousset. 2004. Causes, mechanisms and consequences of
dispersal. Pages 307–335 *in* I. Hanski and O. E. Gaggiotti, eds. Ecology, Genetics and
Evolution of Metapopulations. Elsevier Academic Press Inc, Amsterdam.

Clobert, J., J. F. Le Galliard, J. Cote, S. Meylan, and M. Massot. 2009. Informed dispersal,
heterogeneity in animal dispersal syndromes and the dynamics of spatially structured
populations. Ecology Letters 12:197–209.

Coulson, T. 2012. Integral projections models, their construction and use in posing hypotheses
in ecology. Oikos 121:1337–1350.

Coulson, T., E. A. Catchpole, S. D. Albon, B. J. T. Morgan, J. M. Pemberton, T. H. Clutton-Brock,
M. J. Crawley, et al. 2001. Age, Sex, Density, Winter Weather, and Population Crashes in
Soay Sheep. Science, New Series 292:1528–1531.

Cozzi, G., N. Maag, L. Börger, T. H. Clutton-Brock, and A. Ozgul. 2018. Socially informed
dispersal in a territorial cooperative breeder. Journal of Animal Ecology 87:838–849.

Deere, J. A., T. Coulson, S. Cubaynes, and I. M. Smallegange. 2017. Unsuccessful dispersal
affects life history characteristics of natal populations: The role of dispersal related
variation in vital rates. Ecological Modelling 366:37–47.

Deere, J. A., T. Coulson, and I. M. Smallegange. 2015. Life History Consequences of the
Facultative Expression of a Dispersal Life Stage in the Phoretic Bulb Mite (Rhizoglyphus
robini). PLoS ONE 10:e0136872.

Diaz, A., K. Okabe, C. J. Eckenrode, M. G. Villani, and B. M. Oconnor. 2000. Biology, ecology,
and management of the bulb mites of the genus Rhizoglyphus (Acari : Acaridae).
Experimental & Applied Acarology 24:85–113.

Dieckmann, U., B. O'Hara, and W. Weisser. 1999. The evolutionary ecology of dispersal.
Trends Ecol Evol 14:88–90.

Easterling, M. R., S. P. Ellner, and P. M. Dixon. 2000. Size-specific sensitivity: Applying a new
structured population model. Ecology 81:694–708.

Ellner, S. P., and M. Rees. 2006. Integral Projection Models for Species with Complex
Demography. The American Naturalist 167:410–428.

Hamilton, W. D., and R. M. May. 1977. DISPERSAL IN STABLE HABITATS. Nature 269:578–581.

Hanski, I. 1999. Metapopulation ecology. Oxford University Press, New York.

Harrison, R. G. 1980. DISPERSAL POLYMORPHISMS IN INSECTS. Annual Review of Ecology and
Systematics 11:95–118.

Himes Boor, G. K., C. B. Schultz, E. E. Crone, and W. F. Morris. 2018. Mechanism matters: the

cause of fluctuations in boom–bust populations governs optimal habitat restoration

strategy. Ecological Applications 28:356–372.

Jeltsch, F., D. Bonte, G. Pe'er, B. Reineking, P. Leimgruber, N. Balkenhol, B. Schroder, et al.

2013. Integrating movement ecology with biodiversity research - exploring new avenues

to address spatiotemporal biodiversity dynamics. Movement Ecology 1:6.

Johnson, C. G. 1969. Insect migration and dispersal by flight. Methuen, London.

Johnson, M. L., and M. S. Gaines. 1990. Evolution of Dispersal: Theoretical Models and

Empirical Tests Using Birds and Mammals. Annual Review of Ecology and Systematics

21:449–480.

Lemel, J.-Y., S. Belichon, J. Clobert, and M. E. Hochberg. 1997. The evolution of dispersal in a

two-patch system: some consequences of differences between migrants and residents.

Evolutionary Ecology 11:613–629.

Maag, N., G. Cozzi, T. Clutton-Brock, and A. Ozgul. 2018. Density-dependent dispersal

strategies in a cooperative breeder. Ecology 99:1932–1941.

O'Riain, M. J., J. U. M. Jarvis, and C. G. Faulkes. 1996. A dispersive morph in the naked mole-

rat. Nature 380:619.

Ozgul, A., T. Coulson, A. Reynolds, T. C. Cameron, and T. G. Benton. 2012. Population

Responses to Perturbations: The Importance of Trait-Based Analysis Illustrated through

a Microcosm Experiment. American Naturalist 179:582–594.

Phillips, B. L., G. P. Brown, J. K. Webb, and R. Shine. 2006. Invasion and the evolution of speed

in toads. Nature 439.

Rees, M., D. Z. Childs, and S. P. Ellner. 2014. Building integral projection models: a user's guide.

Journal of Animal Ecology 83:528–545.

Ronce, O. 2007. How does it feel to be like a rolling stone? Ten questions about dispersal

evolution. Annual Review of Ecology Evolution and Systematics 38:231–253.

Singh, N. J., L. Börger, H. Dettki, N. Bunnefeld, and G. Ericsson. 2012. From migration to

nomadism: movement variability in a northern ungulate across its latitudinal range.

Ecological Applications 22:2007–2020.

Smallegange, I. M., R. E. Fernandes, and J. C. Croll. 2018. Population consequences of

individual heterogeneity in life histories: overcompensation in response to harvesting of

alternative reproductive tactics. Oikos 127:738–749.

Strayer, D. L., C. M. D'Antonio, F. Essl, M. S. Fowler, J. Geist, S. Hilt, I. Jarić, et al. 2017. Boom-

bust dynamics in biological invasions: towards an improved application of the concept.

Ecology Letters 20:1337–1350.

Topping, C. J., H. F. Alrøe, K. N. Farrell, and V. Grimm. 2015. Per Aspera ad Astra: Through

Complex Population Modeling to Predictive Theory. The American Naturalist 186:669–

674.

Travis, J. M. J., M. Delgado, G. Bocedi, M. Baguette, K. Barton, D. Bonte, I. Boulangeat, et al.

2013. Dispersal and species' responses to climate change. Oikos 122:1532–1540.

Travis, J. M. J., and C. Dytham. 1999. Habitat persistence, habitat availability and the evolution

of dispersal. Proceedings of the Royal Society of London B: Biological Sciences 266:723–

728.

Travis, J. M. J., K. Mustin, K. a. Bartoń, T. G. Benton, J. Clobert, M. M. Delgado, C. Dytham, et

al. 2012. Modelling dispersal: an eco-evolutionary framework incorporating emigration,

movement, settlement behaviour and the multiple costs involved. Methods in Ecology

and Evolution 3:628–641.

Zera, A. J., and R. F. Denno. 1997. Physiology and ecology of dispersal polymorphism in insects.

Annual Review of Entomology 42:207–230.

Table 1. Response of population parameters to changes in dispersal parameters. Dots represent population parameter outcomes given various dispersal parameter scenarios. Black dots indicate expected outcomes, grey dots indicate unexpected outcomes (see main text for details).

	Population parameters			
Dispersal parameters	PD↑ λ_0↓	PD↓ λ_0↑	PD↓ λ_0↓	PD↑ λ_0↑
β↑, δ↑	●			
β↓, δ↓		●		
β↑, δ↓	●	●	○	
β↓, δ↑	●	●		○

β, disperser expression; δ, dispersal probability; PD, proportion of dispersers; λ_0, population growth rate

 FIGURE LEGENDS

Figure 1. Schematic indicating how a population-level approach and an individual-level

approach differ in terms of the individuals that emigrate from a population. The blue blocks

indicate individuals that stay in the natal population whilst grey blocks indicate individuals

that leave the population.

Figure 2. Life cycle of the bulb mite in in the Disperser Phenotype Model (DPM), indicating

the life stages and the vital rates. From the life cycle we calculated the survival (P) and

fecundity (F) rates and the probability of surviving and growing into the next stage (G).

Dispersal consists of development into a disperser (deutonymph) (β) and dispersal out of the

population (δ); β describes the transition probability of developing from a protonymph to a

deutonymph, δ describes the probability of a deutonymph leaving the population.

Figure 3. Joint effect of increasing deutonymph frequency (β) and increasing dispersal

probability (δ) on the proportion of dispersers. Dispersal probability and deutonymph

frequency increase from 0 to 1 at 0.01 increments. Side bar indicates proportion of dispersers

with proportion increasing from dark to light green. Coloured contour lines highlight a fixed

proportion of dispersers, each proportion can be attained by a number of different δ and β

values: 0.04 (yellow line), 0.08 (red line), 0.12 (blue line).

Figure 4. Joint effect of increasing deutonymph frequency (β) and increasing dispersal

probability (δ) on population growth rate (λ_0). Dispersal probability and deutonymph

frequency increase from 0 to 1 at 0.01 increments. Side bar indicates λ_0, with λ_0 increasing

from black to white. Coloured contour lines highlight a fixed proportion of dispersers (0.04,

yellow line; 0.08, red line; 0.12, blue line), each proportion can result in a number of different

λ_0 values.

	Multi-phase life-history approach	Population-level approach
	Individual	**Individual**
Disperser expression	**Disperser** / **Resident**	**Go** / **Stay**
Emigration proportion	**Successful Disperser** / **Unsuccessful Disperser**	

Disperser Phenotype Model (DPM)

Population growth rate (λ_0)