Activation mechanism of ATP-sensitive K$^+$ channels explored with real-time nucleotide binding

Michael C. Puljung, Natascia Vedovato, Samuel Usher, Frances M. Ashcroft*

Department of Physiology, Anatomy & Genetics, University of Oxford, Oxford UK OX1 3PT

*Address correspondence to Frances Ashcroft, frances.ashcroft@dpag.ox.ac.uk

Author contributions: MCP and FMA designed the experiments. MCP, NV, and SU performed the experiments. MCP and FMA wrote the manuscript.

Keywords:

Ion channel, allostery, binding, gating, diabetes, ligand, metabolism

Abstract

The response of ATP-sensitive K^+ channels (K_{ATP}) to cellular metabolism is coordinated by three classes

of nucleotide binding site (NBS). We used a novel approach involving labeling of intact channels in a

native, membrane environment with a non-canonical fluorescent amino acid and measurement (using

FRET with fluorescent nucleotides) of steady-state and time-resolved nucleotide binding to dissect the

role of NBS2 of the accessory SUR1 subunit of K_{ATP} in channel gating. Binding to NBS2 was Mg^{2+}-

independent, but Mg was required to trigger a conformational change in SUR1. Mutation of a lysine

(K1384A) in NBS2 that coordinates bound nucleotides increased the EC_{50} for trinitrophenyl-ADP binding

to NBS2, but only in the presence of Mg^{2+}, indicating that this mutation disrupts the ligand-induced

conformational change. Comparison of nucleotide-binding with ionic currents suggests a model in which

each nucleotide binding event to NBS2 of SUR1 is independent and promotes K_{ATP} activation by the

same amount.

1   Introduction

ATP-sensitive K^+ channel (K_{ATP}) closure initiates the electrical response of pancreatic β-cells to

metabolic changes induced by increased extracellular glucose.[1,2] K_{ATP}'s metabolic sensitivity is

accomplished through the coordinated activity of three classes of intracellular adenine nucleotide binding

site (NBS), one inhibitory and two stimulatory. Despite the recent publication of several cryo-EM

structures of K_{ATP} showing these NBSs at near atomic resolution, the detailed mechanism by which

energetic contributions from nucleotide binding to each site sum to affect channel gating remains

obscure.[3-6]

K_{ATP} is formed by four inward-rectifier K^+ channel subunits (Kir6.2 in β-cells), each associated with a

modulatory sulfonylurea receptor (SUR1 in β-cells, Figure 1a).[7-10] The inhibitory NBS of K_{ATP} is located

on Kir6.2 and clearly resolved in several of the cryo-EM structures.[3-6,11] Like other ATP-binding cassette

(ABC) proteins, SUR1 has two cytoplasmic nucleotide binding domains (NBDs), which associate to form

two NBSs at the dimer interface (Figure 1a,c).[3] NBS2, formed primarily by the nucleotide-binding

$Walker_A$ and $Walker_B$ motifs of NBD2 and the ABC signature sequence of NBD1, is a consensus binding

site, based on conservation of key catalytic residues with other ABC family members, and is competent to

hydrolyze nucleotides.[12,13] In NBS1, formed by the Walker motifs of NBD1 and the signature sequence of

NBD2, the catalytic $Walker_B$ aspartate (D854) is replaced with a glutamate, rendering this site

"degenerate," i.e. incapable of nucleotide hydrolysis. NBS1 binds nucleotides in the absence of Mg^{2+},

whereas it is thought that Mg^{2+} is required for binding to NBS2.[13] The NBSs of SUR1 mediate K_{ATP}

activation by Mg-nucleotides.[11,14] SUR1 also confers inhibition by antidiabetic sulfonylureas (SUs) and

activation by K_{ATP}-specific K^+ channel openers (KCOs).[11,14-16]

Gating of K_{ATP} is a complex function of the intrinsic opening and closing of the channel pore and the

converging influences of the excitatory and inhibitory NBSs. Mutations that directly disrupt the NBSs or

their ability to transduce nucleotide occupancy to the pore result in diseases of insulin secretion.[2,17] It is

crucial to our understanding of K_{ATP} to be able to independently measure (i) the occupancy of each NBS

in real time, (ii) the effect each NBS has on K_{ATP} conformation, and (iii) how binding to each NBS affects

channel open probability (P_{open}). Electrophysiology, radioligand binding, and ATPase assays have

provided rich mechanistic insight into K_{ATP} gating. However, the ultimate readout of such studies is a

function of nucleotide binding to all three NBSs as well as the conformational equilibria affected by

binding. Photoaffinity labeling studies enable binding to each NBS to be separated, and some

determination of their affinity and specificity, but require partially purified proteins, are performed over

long incubation times, and involve irreversible covalent labeling.[13]

An ideal method to separate the nucleotide interactions at each class of site on K_{ATP} would have high

spatial (specific for a single NBS) and temporal resolution, would require only small amounts of protein,

could be used in a native environment (i.e. the cell or plasma membrane), and would operate under

conditions compatible with electrophysiological experiments. We have developed a novel approach for

studying ligand binding that fits these criteria. K_{ATP} was labeled with the fluorescent non-canonical amino

acid L-3-(6-acetylnaphthalen-2-ylamino)-2-aminopropionic acid (ANAP, Figure 1b) and binding of

fluorescent, trinitrophenyl (TNP, Figure 1b) nucleotide derivatives was measured using FRET.[18] This

enabled us to parse the individual contribution of NBS2 to nucleotide binding/channel gating and has

provided insight into the mechanism by which Mg-nucleotides affect the conformation of SUR1, and the

effect this conformational change has on P_{open}.

Results

Nucleotide binding to NBS2. Binding of nucleotides to NBS2 is believed to stimulate K_{ATP} activation.[14,19]

ANAP was introduced at position Y1353 (Y1353*, Figure 1c) of NBS2 using a transfectable amber

codon suppression system.[18] Briefly, a plasmid encoding SUR1 with an amber (TAG) stop codon

corresponding to position 1353 (SUR1-Y1353[stop]) was co-transfected with an additional plasmid

(pANAP) encoding an ANAP-specific tRNA/tRNA synthetase pair. When cells were cultured in the

presence of ANAP, full-length, ANAP-labeled SUR1 protein (SUR1-Y1353*) was produced (Figure 1—

figure supplement 1d). Fluorescence emission spectra of tagged channels were acquired from "unroofed"

membrane fragments expressing SUR1-Y1353*/Kir6.2 (Figure 1d,e; Figure 1—figure supplement 1a).[20-

22]

In the nucleotide-bound structures of K_{ATP}, the aromatic ring of Y1353 forms a π-stacking interaction

with the adenine ring of ADP (Figure 1c).[3,6] To probe nucleotide binding to K_{ATP}, we measured FRET

between SUR1-Y1353*/Kir6.2 and fluorescent TNP-nucleotides (Figure 1d,e). The absorbance spectra of

TNP nucleotides overlap with the emission spectrum of SUR1-Y1353* (Figure 1—figure supplement 2a),

making them suitable FRET partners. Figure 1—figure supplement 2b shows the calculated distance

dependence of FRET between SUR1-Y1353* and TNP-nucleotides. This steep distance dependence

provides the spatial resolution necessary to discriminate between nucleotides bound directly at NBS2

(FRET efficiency close to 100%) or at other sites.

TNP-nucleotides have been used previously to probe nucleotide binding to several ABC proteins

including NBD1 of SUR2A, as well as the cytoplasmic domain of Kir6.2.[23-25] MgTNP-ADP was able to

activate K_{ATP} formed by wild-type SUR1 with a mutated Kir6.2 (Kir6.2-G334D) in inside-out patches

(Figure 1—figure supplement 2c,d; Table 1). G334D disrupts inhibitory nucleotide binding to Kir6.2,

allowing activation to be measured in isolation.[26,27]

FRET between SUR1-Y1353*/Kir6.2 and bound TNP-ADP produced a concentration-dependent

decrease (quenching) of the donor ANAP peak at 470 nm and a concomitant increase in the TNP-ADP

fluorescence at ~565 nm (Figure 1d,e). We quantified nucleotide binding using the decrease in ANAP

fluorescence, as this signal was specific to K_{ATP} (Figure 1—figure supplement 1b,c) and we observed

non-specific association of TNP-nucleotides in sham-transfected membranes at high concentrations

(Figure 1—figure supplement 2e). No change in fluorescence was observed when saturating

concentrations of MgATP or MgADP were applied to SUR1-Y1353*/Kir6.2 channels (Figure 1—figure

supplement 2f), indicating that quenching of the ANAP signal results from FRET and not an allosteric

change in the environment surrounding 1353* caused by nucleotide binding.

Figure 1f shows concentration-response relationships for binding of MgTNP-ATP and MgTNP-ADP

to SUR1-Y1353*/Kir6.2 channels. Both nucleotides quenched ANAP completely, consistent with direct

binding of TNP-nucleotides at NBS2. MgTNP-ADP bound with ~7-fold lower EC_{50} than MgTNP-ATP

(Table 2), consistent with the lower EC_{50} for activation of SUR1/Kir6.2-G334D channels by MgADP vs.

MgATP (7.7 μM vs. 112 μM).[27]

Quite unexpectedly, we observed binding of both TNP-ATP and TNP-ADP to NBS2 in the *absence* of

Mg^+, i.e. in 1 mM EDTA (Figure 1g). Both nucleotides bound with similar affinities in the absence of

Mg^{2+}, but neither fully quenched SUR1-Y1353*/Kir6.2 fluorescence (Table 2). Although quenching was

not 100%, the amount of FRET obtained in the absence of Mg^{2+} is still consistent with nucleotides

binding at NBS2. The difference in FRET efficiency at saturating nucleotide concentrations in the

presence and absence of Mg^{2+} may reflect a change in orientation between the donor and acceptor, as the

NBDs are expected to change conformation in the presence of Mg^{2+}.[3,4]

Despite the relatively close proximity of NBS1 to Y1353* (39 Å to TNP-ATP aligned at NBS1, Figure

1c), the FRET we observed was primarily the result of quenching by TNP-ATP/ADP bound at NBS2.

When nucleotides are bound directly to NBS2, FRET efficiency is very high and any nucleotide binding

at NBS1 would be expected to have little or no effect on our measurements.[28] Nucleotides at NBS1 would

only contribute to the observed FRET signal if NBS2 were unoccupied. However, with no nucleotide

bound to NBS2, the NBDs would not dimerize and 1353* would be ~56 Å away from TNP-nucleotides

bound to NBS1 (based on PDB 6BAA).[4] At this distance, the predicted FRET efficiency is <20% (Figure

1—figure supplement 2b), much less that the FRET efficiencies we observed for TNP nucleotide

quenching ±Mg^{2+} (Figure 1f,g; Table 2).

In the Mg-nucleotide-bound "quatrefoil" structure of K_{ATP}, the inhibitory NBS of Kir6.2 is within ~33

Å of Y1353 (Figure 1—figure supplement 3a).[3] To eliminate a possible contribution of nucleotide bound

at Kir6.2 to our measured FRET signal, we measured binding of MgTNP-ADP to SUR1-Y1353* co-

expressed with Kir6.2-G334D (Figure 1—figure supplement 3b). The EC_{50} was very similar to that

measured with SUR1-Y1353*/Kir6.2 (Table 2), suggesting that either the NBS on Kir6.2 was too distant

for FRET with Y1353*, or that appreciable binding to Kir6.2 did not occur at concentrations at which

NBS2 was unoccupied.

Mg^{2+} locks nucleotides at NBS2. If nucleotides can bind NBS2 in the absence of Mg^{2+}, why is K_{ATP} only

activated in the presence of Mg^{2+}? The simplest explanation is that binding to NBS2 is Mg^{2+}-independent,

but Mg^{2+} is required to support the conformational change that promotes channel activation. Presumably,

this conformational change involves NBD dimerization, as observed in the cryo-EM structures of K_{ATP}.[3,6]

Because nucleotide dissociation would require opening of the NBD dimer, dimerization is expected to

slow the nucleotide off rate. We therefore measured the time course of nucleotide dissociation from

SUR1-Y1353*/Kir6.2 in the presence and absence of Mg^{2+}.

In the absence of Mg^{2+}, TNP-ADP dissociated very rapidly from SUR1-Y1353*/Kir6.2 (Figure 2a,e;

Table 3). Dissociation of TNP-ADP was greatly slowed by the presence of Mg^{2+} (p = 0.001). Similar, but

less dramatic results were obtained with TNP-ATP (Figure 2b,e; p = 0.0001; Table 3). Whereas the time

courses of dissociation of TNP-ADP ±Mg^{2+} and TNP-ATP in the presence of Mg^{2+} were well described

by single exponential decays, the dissociation of TNP-ATP in the absence of Mg^{2+} appeared bi-

exponential.

Our data are consistent with an activation model for K_{ATP} in which the NBDs of SUR1 dimerize in the

presence of Mg^{2+}, preventing rapid nucleotide dissociation. However, it is formally possible that Mg^{2+}

could directly stabilize nucleotide binding without any conformational change in the NBDs. To test this

hypothesis, we examined nucleotide dissociation rates in the presence of the sulfonylurea (SU)

tolbutamide (Figure 2c,d,e, Table 3). The cryo-EM structure of Kir6.2/SUR1 in the presence of

glibenclamide suggests that SUs prevent NBD dimerization, and electrophysiological studies further

demonstrate that SUs prevent Mg-nucleotide induced current activation.[4,29] In the presence of

tolbutamide, the dissociation rates for MgTNP-ADP and MgTNP-ATP were not significantly different

from those measured for nucleotides in the absence of both tolbutamide and Mg^{2+} (Figure 2e, Table 3; p =

0.1, for TNP-ADP, p = 1 for TNP-ATP). This argues that the slower off rate for TNP-ADP and TNP-ATP

in the presence of Mg^{2+} must result from NBD dimerization, not a direct stabilization of nucleotide

binding by Mg^{2+}.

As further support, we tested the effect of diazoxide (a KCO) on the nucleotide dissociation rate. It has

been proposed that nucleotide binding at the NBDs of SUR2A stabilizes binding of the KCO pinacidil.[30]

The time course of nucleotide dissociation in the presence of diazoxide did not closely follow a single

exponential decay, but the drug greatly reduced the dissociation rates for both MgTNP-ADP and MgTNP-

ATP (Figure 2c,d,e; Table 3). This suggests that KCOs function by stabilizing the activated (NBD-

dimerized) conformation of SUR1.

In the absence of any conformational change (i.e. in Mg^{2+}-free solution), nucleotide binding to NBS2

would be expected to follow a simple binding equilibrium (SCHEME 1), with $K_d = k_{off}/k_{on}$.

13
14 $$\text{TNP-nucleotide} + \text{SUR1} \underset{k_{off}}{\overset{k_{on}}{\rightleftharpoons}} \text{TNP-nucleotide·SUR1} \quad \text{SCHEME 1}$$
15
16

17 A simple activation model like the one proposed by Del Castillo and Katz for acetylcholine receptors

18 (SCHEME 2), in which nucleotides bind SUR1 and change its conformation can be invoked to explain the

19 decrease in the apparent rate of nucleotide dissociation.[31] This would depend on the kinetics of entry into

20 (α) and exit from (β) the activated state (SUR1'), as well as the intrinsic k_{off}. In SCHEME 2, $EC_{50,apparent} =$

21 $K_d / (\alpha/\beta+1)$.

22
23 $$\text{MgTNP} + \text{SUR1} \underset{k_{off}}{\overset{k_{on}}{\rightleftharpoons}} \text{MgTNP·SUR1} \underset{\beta}{\overset{\alpha}{\rightleftharpoons}} \text{MgTNP·SUR1'} \quad \text{SCHEME 2}$$
24
25

In our experiments, the off-rate for TNP-ADP was reduced ~5-fold in the presence of Mg^{2+}, whereas

the EC_{50} for TNP-ADP was largely unaffected (Figure 2e, Figure 1—figure supplement 4a, Table 2,

Table 3). Mg^{2+} reduced the off-rate for TNP-ATP by a factor of 2.2, but actually increased the EC_{50} for

binding by a factor of ~3 (Figure 2e, Figure 1—figure supplement 4b, Table 2, Table 3). In the limiting

case for SCHEME 2, in which β>>α, the apparent EC_{50} would be equal to the actual K_d. (i.e. that measured

in Mg^{2+}-free solution). Therefore, the concentration-response curve for nucleotide binding would not be

expected to shift to the right in the presence of Mg^{2+}. As we observed such a shift for Mg-TNP-ADP, we

conclude that in addition to stabilizing the NBDs in a dimerized state when nucleotides are present, Mg^{2+}

must also decrease the intrinsic on-rate for nucleotide binding (k_{on}).

The nucleotide dissociation rates measured in the presence of Mg^{2+} are slower than the deactivation

rates observed for SUR1/Kir6.2-G334D currents following nucleotide washout.[27] However, it should be

noted that current decay reflects a conformational change in the channel and not the rate of nucleotide

dissociation, and may be affected by the presence of four SUR1 monomers per channel. Furthermore, the

deactivation rates were measured for MgATP/ADP and not TNP-nucleotides. Ultimately, a direct

comparison of TNP-nucleotide dissociation rates and deactivation rates will require simultaneous

measurement of current and binding.

Mutation of the Walker_A lysine of NBS2 affects the conformational change in SUR1. We next sought to

explore the mechanistic consequences of mutating the Walker_A nucleotide binding motif of NBS2. In

other ABC proteins, and in the Mg-nucleotide-bound structure of K_{ATP}, the Walker_A lysine coordinates

the β- and γ-phosphates of bound nucleotides (Figure 3a). Previous electrophysiological studies showed

that mutation of the Walker_A lysine in NBS2 to alanine (K1384A, K2A) reduced the maximal activation

of SUR1/Kir6.2-G334D currents by MgATP and shifted the concentration dependence of MgADP

activation >20-fold to higher concentrations, such that the maximal response could not be determined.[29]

This is consistent with the K2A mutation affecting channel activation.

We mutated the Walker_A lysine of NBS2 to alanine in channels labeled with ANAP at NBS2 (SUR1-

Y1353*,K2A/Kir6.2) and measured nucleotide binding with and without Mg^{2+} (Figure 3). In the absence

of Mg^{2+} we observed no significant difference in steady-state TNP-ATP and TNP-ADP binding to SUR1-

Y1353*,K2A/Kir6.2 compared to SUR1-Y1353*/Kir6.2 (Figure 3b,c; Table 2), suggesting the K2A

mutation has no effect on the intrinsic nucleotide affinity in the absence of Mg^{2+}. The apparent affinity for

MgTNP-ATP was also unaffected by the K2A mutation (Figure 3d; Table 2). However the K2A mutation

disrupted MgTNP-ADP binding (EC_{50} = 18.6 µM for K2A vs. 4.8 µM for SUR1-Y1353*, p = 0.04;

Figure 3e, Table 2). Similar to SUR1-Y1353*/Kir6.2 channels, Mg^{2+} caused a rightward shift in the

concentration dependence of nucleotide binding to SUR1-Y1353*,K2A/Kir6.2 (Figure 3—figure

supplement 1).

As we observed no change in the TNP-ADP concentration-response in the absence of Mg^{2+}, the K2A

mutation may affect MgTNP-ADP binding indirectly, by reducing the ability of the nucleotide to cause

NBD dimerization. To explore this idea further, we measured the rate of MgTNP-ADP dissociation from

SUR1-Y1353*,K2A/Kir6.2 (Figure 3f, Table 3). The K2A mutation increased the off-rate for MgTNP-

ADP compared to SUR1-Y1353*/Kir6.2 channels (Table 3; p = 0.04). There was no apparent change in

the off-rate for TNP-ADP in the absence of Mg^{2+}, compared to SUR1-Y1353*/Kir6.2 (p = 0.8). Taken

together, these data suggest that the K2A mutation destabilized NBD dimerization induced by MgTNP-

ADP. However, these results do not rule out the possibility that the K2A mutation also affected the

intrinsic on-rate for MgTNP-ADP.

Correlation of binding with gating. We next sought to correlate nucleotide binding with changes in

channel activity. Despite our ability to demonstrate a Mg^{2+}- and nucleotide-dependent conformational

change in SUR1-Y1353*/Kir6.2, we were unable to measure activation of SUR1-Y1353*/Kir6.2 channels

by Mg-nucleotides. There are two possible explanations for this observation. The first is that SUR1-

Y1353* disrupts communication between the NBD-dimerized SUR1 and Kir6.2. Inspection of the Mg-

nucleotide-bound "quatrefoil" structure shows that position 1353 is adjacent to residues (e.g. R1352) at

the interface between SUR1 and Kir6.2. Mutation of R1352 disrupts K_{ATP} activation.[3,32,33] Another

possible explanation for the lack of activation of SUR1-Y1353*/Kir6.2 currents is that channels

containing SUR1 subunits truncated at Y1352 (i.e. no ANAP incorporated) traffic to the plasma

membrane. These channels would not be activated by nucleotides as they lack a complete NBS2. K_{ATP}

channels with SUR1 truncated at residue 1330 at the start of NBD2, traffic to the plasma membrane but

are not activated by MgADP.[34]

In fluorescence measurements, we only observe full-length SUR1-Y1353*/Kir6.2 channels as only

these are labeled with ANAP (Figure 1—figure supplement 1). However, in electrophysiological

experiments it may be possible to measure currents from channels with both full-length and truncated

SUR1. Using a surface expression assay, we observed no difference in the ability of full-length and

truncated SUR1-Y1353[stop] (no ANAP included in the culture medium) to chaperone HA-tagged Kir6.2 to

the plasma membrane (Figure 4—figure supplement 1a).[35] Truncated SUR1-Y1353[stop]/Kir6.2 (expressed

without pANAP) formed functional channels that were not activated by MgADP (Figure 4—figure

supplement 1b). 100 μM MgADP increased full-length 3xFLAG_SUR1/Kir6.2 currents 3.6-fold over the

current in nucleotide-free control solutions (Figure 4—figure supplement 1b, Table 4). In contrast, 100

μM MgADP inhibited current from truncated SUR1-Y1353[stop] by 59% (Figure 4—figure supplement 1b,

Table 4). This degree of inhibition is similar to the ~60% reduction in wild-type SUR1/Kir6.2 current by

100 μM ADP in Mg^{2+}-free solutions, in which only inhibition is expected[27], confirming that SUR1-

Y1353[stop]/Kir6.2 channels failed to activate. Currents from SUR1-Y1353[stop]/Kir6.2 co-expressed with

pANAP alone (66% full-length SUR1-Y1353* expected, Figure 1—figure supplement 1d) or with

pANAP and a dominant-negative ribosomal release factor peRF1-E55D (92% full-length SUR-Y1353*

expected, Figure 1—figure supplement 1d) were inhibited by 100 μM MgADP to a similar extent as

currents from cells expressing only truncated channels (Figure 4—figure supplement 1b, Table 4).[36] This

suggests that SUR-Y1353* fails to support nucleotide activation. To confirm this, we also looked at

currents in patches excised from cells expressing SUR1-Y1353*/Kir6.2-G334D (co-expressed with

pANAP and pERF1-E55D). Wild-type SUR1 paired with Kir6.2-G334D produced channels that were

activated by MgADP and inhibited by tolbutamide (Figure 4—figure supplement 1c). When measured

under the same conditions, SUR1-Y1353*/Kir6.2 was inhibited by tolbutamide, but we observed no

activation by MgADP.

We therefore sought to label an additional position near NBS2 of SUR1 that produces functional,

nucleotide-activated channels when paired with Kir6.2. We mutated the codons for 34 different residues

(amino acids 1386-1397, 1400-1408, and 1410-1423) to the amber stop codon and screened the constructs

for their ability to promote surface expression of HA-Kir6.2 in the presence of ANAP. We focused on

SUR1-T1397^{stop}, as this mutation strongly increased the surface expression of HA-Kir6.2 in the presence

of ANAP (Figure 4—figure supplement 1a; p = 0.005). Further, in the absence of ANAP, this construct

decreased the surface expression of HA-Kir6.2 (p = 0.009) compared to no-SUR1 controls (Figure 4—

figure supplement 1a), suggesting that channels with SUR truncated at this position may be selectively

retained within the cell. Western blots of total protein from cells expressing SUR1-T1397^{stop} in the

presence of ANAP and pANAP/peRF1-E55D confirmed that >95% of SUR1-T1397* was full length

(Figure 4—figure supplement 1d). The expected distance between T1397 and TNP-nucleotides bound to

NBS2 is ~18.3 Å (Figure 4a), close enough for >99% FRET efficiency (Figure 1—figure supplement 2b).

Binding of MgTNP-ADP and MgTNP-ATP to SUR1-T1397*/Kir6.2 (Figure 4b) was qualitatively

similar to SUR1-Y1353*/Kir6.2 (Figure 1, Table 2), although there was no obvious difference in binding

affinity between MgTNP-ADP and MgTNP-ATP for SUR1-T1397*/Kir6.2. Both TNP-ADP and TNP-

ATP bound to SUR1-T1397*/Kir6.2 in the absence of Mg^{2+} (Figure 4c; Table 2) as we observed for

SUR1-Y1353*/Kir6.2. Mg^{2+} cause a rightward shift in the concentration dependence of TNP-ATP and

TNP-ADP binding (Figure 4—figure supplement 2).

In excised patches from cells expressing SUR1-T1397*/Kir6.2, we measured robust currents that were

inhibited by ATP (Figure 4—figure supplement 1e). Interestingly, the apparent affinity for ATP inhibition

for SUR1-T1397*/Kir6.2 was lower than for wild-type K_{ATP} (IC_{50} was 53.9 μM vs. 11.2 μM, respectively;

Table 5). More importantly, SUR1-T1397* supported activation of Kir6.2-G334D by both MgADP

(Figure 4d, Table 1) and MgTNP-ADP (Figure 5a,b; Table 1). The EC_{50} for channel activation by

MgTNP-ADP (determined from Hill fits) was 7.3 μM ±1.2 μM as compared to the EC$_{50}$ of 10.8 μM ±0.7

μM obtained for MgTNP-ADP binding to NBS2. Although activation (by necessity) was measured with

Kir6.2-G334D, and our binding measurements were conducted with wild-type Kir6.2, we do not expect

this to have affected our measurements as 1397* is the same distance (~33 Å) as 1353* from the

inhibitory binding site on Kir6.2, and expression of SUR1-Y1353* with Kir6.2-G334D did not affect

binding measurements (Figure 1—figure supplement 3b, Table 2).

Discussion

Taken together, our data suggest a modified model for SUR1/Kir6.2 K_{ATP} activation by nucleotide

binding to SUR1 (Figure 6). ADP and ATP bind to NBS2 in the absence of Mg^{2+}. However, without Mg^{2+}

as a co-factor, adenine nucleotides cannot dimerize the NBDs and initiate the conformational change in

SUR1 required to affect P$_{open}$. As the NBDs fail to dimerize, nucleotides dissociate rapidly. If both Mg^{2+}

and nucleotides are present, the NBDs dimerize, resulting in slower nucleotide dissociation. NBD

dimerization initiates a conformational change in SUR1 that ultimately communicates nucleotide

occupancy to the pore, affecting P$_{open}$. Inhibitory SUs prevent NBD dimerization. Thus, nucleotides

dissociate rapidly even in the presence of Mg^{2+} and P$_{open}$ is not affected by nucleotide occupancy at

NBS2. KCOs have the opposite effect. They stabilize NBD dimerization, prolonging nucleotide

occupancy at NBS2 and channel activation.

Although our data do not offer any direct structural insight into how NBD dimerization propagates to

Kir6.2, our ability to measure MgTNP-ADP binding to NBS2 of SUR1 and channel activation under

similar conditions furnishes some mechanistic insights into the relationship between nucleotide binding

and channel opening. We simultaneously fit the combined data sets for MgTNP-ADP binding to SUR1-

T1397*/Kir6.2 and activation of SUR1-T1397*/Kir6.2-G334D currents with several gating models

(Figure 5, Figure 5—figure supplement1, Figure 5—figure supplement 2). For each model, equations for

binding and P$_{open}$ used common values for K_A (the equilibrium association constant in units of M^{-1},

constrained such that $K_A > 0$), L (the equilibrium constant for the intrinsic opening of unliganded SUR1-

T1397*/Kir6.2-G334D, fixed to 0.52 based on the measured P_{open} of SUR1/Kir6.2-G334D[27]), and E (the

factor by which MgTNP-ADP binding favors activation, constrained to E>1). Goodness of fit was

assessed by comparing the residual sum of the squared deviations (RSS) between the data and fit curves

(Figure 5—figure supplement 2). As K_{ATP} channels open in the absence of Mg^{2+} and nucleotides, we

excluded any gating models that lacked unliganded opening. None of the available cryo-EM

structures.[3,4,6,37,38] show any direct interactions between neighboring SUR1 subunits. Therefore, we

excluded models in which binding to one SUR1 directly affects binding to the other SUR1 subunits (i.e.

with binding cooperativity).

With these assumptions, the best fits to the data were obtained with an extended Monod-Wyman-

Changeux model (Figure 5b; $R^2 = 0.98$, RSS = 0.15).[39] In this model, binding of MgTNP-ADP to each

NBS2 of SUR1 is independent and favors pore opening by the same amount. The data in Figure 5b were

fit with the following expressions for binding and activation:

13
$$\frac{bound}{total} = \frac{LE[ADP]K_A(E[ADP]K_A + 1)^3 + [ADP]K_A([ADP]K_A + 1)^3}{L(E[ADP]K_A + 1)^4 + ([ADP]K_A + 1)^4} \quad (6)$$

15
$$P_{open,norm} = \frac{\left(\frac{L(E[ADP]K_A + 1)^4}{L(E[ADP]K_A + 1)^4 + ([ADP]K_A + 1)^4}\right) - \left(\frac{L}{L + 1}\right)}{\left(\frac{LE^4}{LE^4 + 1} - \frac{L}{L + 1}\right)} \quad (7).$$

Previous attempts to determine the stoichiometry of Mg-nucleotide activation of K_{ATP}, i.e. how many

SUR1 subunits must bind nucleotide in order to activate current, have produced varying results.[40-42] We

also fit MgTNP-ADP binding to NBS2 and current activation for SUR1-T1397* with models in which

fixed stoichiometries of 1-4 nucleotide-bound SUR1 subunits were necessary to increase P_{open} (Figure 5—

figure supplement 1, Figure 5—figure supplement 2). All of these models used the same assumptions as

the MWC model in Figure 5b, but none fit the data as well. An inspection of our model fits (Figure 5—

figure supplement 2) shows that the residuals for the MWC model and the model in which two bound

nucleotide molecules activate K_{ATP} are small and centered around zero. The other three models show

systematic errors. Whereas the model in which two binding events opens K_{ATP} provides a reasonable fit to

the data, this model also predicts $E \approx 1$, i.e. no activation. The MWC fits to our data predicts $E = 2.2$,

which is similar to the value of 1.76 that we previously estimated from electrophysiology data.[27,43] This

would favor the MWC model. However, given the scatter in the data, further experimentation will be

required to truly distinguish between these competing models (see below).

Our binding and activation data were determined in separate experiments. Whereas the equations for

binding and current activation both have a term (L) representing the intrinsic opening equilibrium, there is

no way of knowing the P_{open} of channels in an unroofed membrane. We expect P_{open} to be quite low, as

several minutes may elapse between cell unroofing and the completion of a given experiment, during

which time channel rundown may occur (e.g. due to dissociation/degradation of PIP_2). Ideally, to derive

more quantitative values, experiments should be performed using patch-clamp fluorometry in which

binding and current are measured simultaneously.[44,45] However, we were unable to obtain inside-out

patches from cells expressing SUR1-T1397*/Kir6.2-G334D with sufficient fluorescence intensity to

measure reliable spectra.

In our study, the concentration-response relationship for current activation was normalized to the

minimum and maximum current values, as reflected in equation 7. If the true maximum and minimum

P_{open} in a given experiment could be accurately determined, then the quantity E could be computed

directly from the minimum P_{open} in the absence of ligand, $L/(L+1)$, and the maximum P_{open} at saturating

nucleotide concentrations, $E^4L/(E^4L+1)$, and our models would be further constrained. In practice,

however, these values are difficult to measure for K_{ATP} in the presence of Mg^{2+} as channel rundown from

loss of PIP_2 results in an intrinsic P_{open} (L) that is constantly changing.[46] Nevertheless, our analysis

provides useful qualitative information regarding the mechanism by which nucleotide binding to NBS2 of

SUR1 activates Kir6.2.

The affinities we measured for (Mg)TNP-ADP and (Mg)TNP-ATP binding are higher than those

previously estimated for (Mg)ATP and (Mg)ADP from photoaffinity labeling studies on SUR1 in the

absence of Kir6.2 or from ATPase assays preformed on isolated NBD2 fusion proteins.[12,47] This may

reflect a higher affinity of TNP nucleotides for NBS2 of SUR1 compared to ATP and ADP, as observed

for other ABC transporters or the absence of Kir6.2.[24] Nevertheless, the EC_{50} values obtained from

MgTNP-ATP and MgTNP-ADP binding to SUR1-Y1353*/Kir6.2 were in reasonably good agreement

with those measured for MgATP and MgADP activation of SUR1/Kir6.2-G334D currents.[27] It is unlikely

that the introduction of ANAP modified the binding affinity of NBS2 as the apparent affinity for TNP-

nucleotides in the absence of Mg^{2+} was unaffected by the site at which ANAP was inserted (Table 2). The

difference in apparent affinities between SUR1-Y1353* and SUR1-T1397* in the presence of Mg^{2+} may

indicate a difference in the ability of bound nucleotides to promote NBD dimerization.

MgADP can activate channels to an equal or greater extent than MgATP.[27] Therefore, it is clear that the

act of ATP hydrolysis *per se* is not required for channel activation, even though NBS2 is competent to

hydrolyze ATP.[12,13] It is often stated that activation of K_{ATP} channels by Mg^{2+}-ATP proceeds through

hydrolysis of MgATP to MgADP.[48,49] However, there is no evidence that an irreversible step like ATP

hydrolysis occurs during the gating cycle of the K_{ATP} channel in the presence of MgATP, as is the case for

the closely related ABC protein CFTR.[50,51] Furthermore, high concentrations of ATP have been shown, in

radioligand biding assays, to cause SUR1 to change conformation in the absence of Mg^{2+}, under which

conditions ATP hydrolysis is not expected.[52] Mg^{2+}-free ATP was also recently demonstrated to activate

K_{ATP} formed by SUR1-Q1179R, a gain-of-function SUR1 mutation, paired with Kir6.2-G334D at high

concentrations, as well as SUR constructs in which the $Walker_B$ glutamate was mutated to glutamine.[53] As

lack of Mg^{2+} or mutation of the catalytic $Walker_B$ motif do not support MgATP hydrolysis, this suggests

that ATP can activate K_{ATP} directly. No Mg^{2+}-free ATP activation was demonstrated when wild-type SUR1

was expressed with Kir6.2-G334D and we see no indication in our data that nucleotides caused a SUR1 to

change conformation in the absence of Mg^{2+} (Figure 2).[53]

Our data suggest that ATP hydrolysis did not occur over the course of our experiments, as the off-rate

for MgTNP-ATP was faster than that of MgTNP-ADP ($\tau_{0.5}$ = 10.5 s vs 14.7 s). If MgTNP-ATP were

hydrolysed to MgTNP-ADP before being released (as is thought to be true of many ABC family members,

including CFTR) the dissociation rates should have been similar. The observation that MgTNP-ATP

dissociation was faster than MgTNP-ADP dissociation may reflect a lack of MgTNP-ATP hydrolysis

during the course of our experiments. Alternatively, this may indicate a destabilizing effect of inorganic

phosphate on MgTNP-ADP binding to NBS2 (which would not be present when binding MgTNP-ADP

alone, but would be present as a product of MgTNP-ATP hydrolysis) on binding.

Our ability to directly measure nucleotide binding in real time will be of considerable value for future

investigations. This will allow us to address the role of nucleotide binding at NBS1 and Kir6.2, and

explore the mechanistic ramifications of disease-associated K_{ATP} mutations. Furthermore, simultaneous

binding and current measurements will yield precise information regarding the coupling of stimulatory

and inhibitory nucleotide binding to the channel pore. This coupling of ligand binding to an effector

domain (the channel pore) that lies tens of Å away provides an excellent model for studying long-range

communication within protein complexes. It has also not escaped our notice that our method is readily

adaptable to the study of other ABC proteins, ATP-gated channels like P2X receptors (for which different

subtypes demonstrate distinct selectivity for MgATP vs. ATP) and any protein for which there is a

suitable fluorescent ligand.[54]

1   **Materials and Methods**

2   **Key Resources Table**

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
cell line	HEK-293T	LGC Standards (ATCC® CRL-3216™)		
transfected construct (*Escherichia. coli*)	pANAP	Addgene		
transfected construct	pcDNA4/TO	Addgene		
transfected construct (*Aequorea victoria*)	pCGFP_EU	Gouaux Laboratory (Vollum Institute, Portland OR USA);		
transfected construct (*Homo sapiens*)	peRF1-E55D	Chin Laboratory (MRC Laboratory of Molecular Biology, Cambridge UK);		
antibody	Monoclonal ANTI-FLAG® M2 antibody	Sigma-Aldrich	(Sigma-Aldrich Cat# F3165, RRID:AB_259529)	
antibody	Anti-HA High Affinity; Rat monoclonal antibody (clone 3F10)	Roche	(Roche Cat# 11867423001, RRID:AB_10094468)	
antibody	Sheep Anti-Mouse IgG ECL Antibody, HRP Conjugated	GE Healthcare	(GE Healthcare Cat# NA9310-1ml, RRID:AB_772193)	
antibody	Peroxidase-AffiniPure Goat Anti-Rat IgG (H+L) antibody	Jackson ImmunoResearch Labs	(Jackson ImmunoResearch Labs Cat# 112-035-003, RRID:AB_2338128)	
chemical compound, drug	trinitrophenyl-ATP (TNP-ATP)	Jena Bioscience (Jena, Germany)		
chemical compound, drug	trinitrophenyl-ADP (TNP-ADP)	Jena Bioscience (Jena, Germany)		
chemical compound, drug	L-3-(6-acetylnaphthalen-2-ylamino)-2-aminopropionic acid	Asis Chemicals (Waltham, MA)		

Molecular Biology. *Homo sapiens* SUR1 and human Kir6.2 were subcloned into pcDNA4/TO

(Invitrogen; Carlsbad, CA) for expression in HEK-293T cells. For experiments with GFP-labeled

channels (Figure 1—figure supplement 1b,c), SUR1 was cloned into pCGFP_EU.[55] Mutagenesis,

including insertion of amber (TAG) stop codons, was performed using the QuikChange kit (Agilent;

Santa Clara, CA) or by subcloning synthetic oligonucleotides (Sigma; St. Louis, MO) between convenient

restriction sites. All clones were verified by sequencing (DNA Sequencing and Services; Dundee,

Scotland). pCDNA4/TO and pANAP were obtained from Addgene. peRF1-E55D (*Homo sapiens*) and

pCGFP_EU (*Aequorea victoria*) were kind gifts from the Chin Laboratory (MRC Laboratory of

Molecular Biology, Cambridge UK) and Gouaux Laboratory (Vollum Institute, Portland OR USA),

respectively.

Cell Culture and Expression of Fluorescently Tagged Protein. HEK-293T cells were obtained from and

verified/tested for mycoplasma by LGC Standards (ATTC® CRL-3216™; Middlesex, UK). Frozen

stocks were prepared directly from these cells and stored in liquid nitrogen. Working stocks were

routinely replenished from frozen following 15-25 passages. Our working stock tested negative for

mycoplasma contamination using the MycoAlert® Mycoplasma Detection Kit (Lonza Bioscience; Burton

on Trent, UK). For experiments, cells were grown in 6-well plates (STARLAB; Milton Keynes, UK) or

30 mm culture dishes (STARLAB) on untreated 30 mm borosilicate coverslips (Thickness # 1, VWR

International; Radnor, PA) or on poly-D-lysine coated FluoroDishes (FD35-PDL-100, World Precision

Instruments; Hitchin, UK) in Dulbecco's Modified Eagle Medium (Sigma) supplemented with 10% fetal

bovine serum, 100 U/mL penicillin, and 100 μg/mL streptomycin (Thermo Fisher Scientific; Waltham,

MA) at 37ºC, 5%/95% CO_2/air.

SUR1 constructs were site-specifically tagged with the fluorescent amino acid ANAP using amber

codon suppression as described by Chatterjee *et al.*[18] 30 mm dishes of HEK 293T cells were co-

transfected with pANAP (0.5-1 μg), Kir6.2 (0.5 μg), and SUR1 (1-1.5 μg) constructs containing an amber

stop codon (TAG) at amino acid position 1353 or 1397 (1353[stop] or 1397[stop]) using TransIT transfection

reagent (Mirus Bio LLC; Madison, WI) in a ratio of 3 μL of TransIT per μg of total DNA. Following

transfection the media was supplemented with 20 μM ANAP (free acid or methyl ester). pANAP encodes

a tRNA/tRNA synthetase pair specific for ANAP. In the presence of ANAP, cells transfected with

pANAP generate tRNAs charged with ANAP that recognize the amber stop codon. Thus, full-length

ANAP-tagged SUR can be produced. Proper co-assembly of Kir6.2 subunits with SUR1 subunits is

ensured by the fact that the two subunits rely on one another for proper exit from the ER and trafficking to

the plasma membrane.[35] An exception to this may be SUR1_GFP, which has been shown to traffic

independently of Kir6.2.[56] After transfection, cells were cultured at 33º C, 5%/95% CO_2/air to slow

growth and increase the per-cell protein yield.[57] To enhance expression of full-length, ANAP-labeled

proteins, 1-2 μg of a plasmid containing a dominant negative eukaryotic release factor 1 (peRF1-E55D)

was included in the transfection mix as indicated.[36] Experiments were performed 2-5 days post

transfection.

Preparation of unroofed membrane fragments for imaging. Unroofed membranes were prepared using a

modification of a protocol from the Heuser laboratory.[20-22] Briefly, a coverslip containing adherent cells

was removed from the cell culture medium and a small piece (0.5-1 cm) was broken off using jewelers

forceps (#5) and washed in phosphate buffered saline (PBS, Thermo Fisher Scientific) diluted 1:3 in

deionized water. Cells were exposed 3x10 s to a solution of 0.01% poly-L-lysine (Sigma) with intervening

washes in diluted PBS to adhere them more firmly to the coverslips. The coverslip fragment was then

placed in a 30 mm culture dish filled with 2-3 mL of 1/3X PBS and sonicated very briefly (~100 ms)

using a probe sonicator (Vibra-cell; Newtown, CT), leaving behind adherent plasma membrane and

sometimes additional cellular material. The unroofed fragments in our experiments were sometimes

visible under bright-field illumination, a potential result of our modified "unroofing" procedure.

Morphologically, such cell fragments resemble the "partially unroofed" cells identified by Usukura *et*

al.[21] Cells cultured on poly-D-lysine coated FluoroDishes were unroofed in diluted PBS with no

additional poly-L-lysine treatment. This procedure yielded a higher percentage of completely unroofed

plasma membranes (i.e. nearly invisible with brightfield illumination).

Microscopy/spectroscopy Unroofed membrane fragments were imaged directly on FluoroDishes (FD35-

PDL-100, World Precision Instruments) or on broken coverslips placed in a FluoroDish (FD3510, World

Precision instruments) using a Nikon TE2000-U microscope equipped with a 40x (S Fluor, 1.3 NA;

Nikon, Kingston Upon Thames, UK) or 100x (Apo TIRF, 1.49 NA, Nikon) oil-immersion objective and

low-fluorescence immersion oil (MOIL-30, Olympus; Southend-on-Sea, UK) or a 60x water immersion

objective (Plan Apo VC, 1.20 NA, Nikon). ANAP was excited using a ThorLabs LED source

(LED4D067) at 385 nm connected to the microscope using a liquid light guide with a 390/18 nm band-

pass filter (MF390-18, ThorLabs; Newton, NJ) and MD416 dichroic (ThorLabs). For imaging, emitted

light was filtered through a 479/40 nm band-pass filter (MF479-40, ThorLabs). GFP was imaged using a

similar setup with excitation at 490 nm, a 480/40 band-pass excitation filter (Chroma; Bellows Falls, VT),

a DM505 dichroic mirror (Chroma), and 510 nm long-pass emission filter (Chroma). All images were

acquired using a PIXIS 400B CCD camera (Princeton Instruments; Trenton, NJ). For spectroscopy,

ANAP was excited as above, but emitted light was passed through a 400 nm long-pass filter (FEL0400,

ThorLabs) and directed through the slit of an Isoplane 160 spectrometer (300 g/mm grating; Princeton

Instruments) in series with the camera. The acquired images retained spatial information in the y-

dimension, with the x-dimension replaced by wavelength (Figure 1—figure supplement 1a). Exposure

times were typically 100 ms for brightfield images, 1-10 s for steady-state spectra and 1-5 s for time-

course experiments. The emission spectrum of ANAP-labeled SUR1 showed a peak around 470 nm,

which was used to differentiate labeled channels from autofluorescence (which typically had a broad

emission spectrum), from cytoplasmic ANAP (or potentially ANAP-conjugated to tRNA), which peaks

~480 nm or from bright, fluorescent debris on the coverslip which was morphologically distinct from

unroofed membranes and had spectra peaking at either ~450 nm or 480-485 nm.

To verify that the fluorescence signal we observed in unroofed membrane fragments derived from

ANAP-labeled SUR1, we co-transfected cells with Kir6.2 and GFP-tagged SUR1 with or without an

amber stop codon at position Y1353. After unroofing, membranes from transfected cells could be

identified by the green fluorescence of SUR1_GFP/Kir6.2. Figure 1—figure supplement 1b shows that

membranes expressing SUR1_GFP-Y1353*/Kir6.2 were brightly fluorescent for both GFP and ANAP,

whereas those from SUR1_GFP expressing cells had only GFP fluorescence, even though ANAP was

included in the cell-culture medium. We also acquired emission spectra from membranes expressing

SUR1_GFP-Y1353*/Kir6.2 (Figure 1—figure supplement 1c). The intensity of the ANAP peak in our

fluorescence was linearly proportional to the peak GFP fluorescence, and the intensity of the ANAP peak

extrapolated to zero GFP (i.e. no expression of SUR1_GFP-Y1353*) was zero. Taken together, these data

suggest that non-specific ANAP background was very low. We also performed Western blotting on

FLAG-tagged SUR1-Y1353[stop] constructs to verify that we produced full-length, ANAP-labeled SUR1,

and that cells were unable to read through the amber stop codon to produce full-length protein in the

absence of ANAP (Figure 1—figure supplement 1d).

To prevent accumulation of fluorescent nucleotides in the bath, FluoroDishes were perfused with

either (in mM) 140 KCl, 20 NaCl, 10 MgCl$_2$, 10 HEPES pH 7.2 w/NaOH or 140 KCl, 20 NaCl, 1 EDTA,

10 HEPES pH 7.2 w/NaOH using a peristaltic pump (Ismatec; Wertheim, Germany). Unroofed

membranes were directly perfused with nucleotides using an 8-channel μFlow perfusion system (ALA

Scientific Instruments; Farmingdale, NY). Solution change with the μFlow was rapid ($\tau = 0.73$ s ±0.02 s)

as measured from single exponential fits to wash-out of 50 μM tetramethylrhodamine-5-maleimide

(TMRM) after switching to a solution with 0 TMRM. TMRM was excited with a 565 nm LED with a

D540/25X band-pass filter (Chroma) and DM565 dichroic mirror (Chroma). Emitted light was collected

through a D605/55M band-pass filter (Chroma). Solution exchange was also monitored during

experiments with TNP-nucleotides by examining the emission spectrum of a background region in each

image. However, the detection limit for TNP-nucleotides in solution was low as unbound nucleotides

have a low quantum yield (enhanced by protein binding) and are not excited via FRET, as is the case for

nucleotides bound to ANAP-labeled SUR1.[58]

Images and spectra were acquired using LightField software (Princeton Instruments). The solution

changer, camera, and light source were all controlled using pClamp 10.5 (Molecular Devices; San Jose,

CA) and a DigiData 1440 A/D converter. All data were acquired at room temperature (18º-22º C).

Analysis of images and spectra. Images were adjusted and displayed using Fiji.[59] Spectra were analyzed

using custom code written in Matlab (Mathworks; Natick, MA). The code was deposited on GitHub

(https://github.com/mpuljung/spectra-analysis).[60,61] Regions of interest corresponding to fluorescent

membrane fragments were manually selected from the spectral images and averaged for each wavelength.

A background region of similar size was selected and the average was subtracted from the spectrum of the

region of interest. The peak fluorescence (~470 nm) was automatically selected by determining the

maximum of a boxcar average of 31 points (corresponding to ~10 nm) of the data in the absence of

nucleotides. Data were corrected for photobleaching by fitting an exponential decay ($F = Ae^{-t/\tau} + (1- A)$)

to the normalized peak of 5-6 images acquired prior to washing on TNP-nucleotides and dividing the raw

spectra by the resulting fit at each cumulative exposure time (Figure 1—figure supplement 5).

Photobleaching rates were similar regardless of the position of ANAP in SUR1. For concentration-

response relationships, corrected peak data (at ~470 nm) were normalized by the ANAP fluorescence at 0

[TNP-nucleotide] (F/F_{max}) and displayed as $1-F/F_{max}$. On occasion data were normalized instead to the

fluorescence at 50 nM nucleotide, as we consistently saw no effect at this concentration. TNP-nucleotide

concentration-response relationships were fit with the Hill equation

12
$$1 - \frac{F}{F_{max}} = E_{max} * \frac{[TNP]^h}{EC_{50}{}^h + [TNP]^h} \quad (1)$$

where E_{max} is the FRET efficiency at saturating concentrations, [*TNP*] is the concentration of TNP-

nucleotide, EC_{50} is the half-maximal concentration and h is the Hill slope.

To generate the plot of GFP fluorescence vs. ANAP fluorescence in Figure 1—figure supplement 1c,

spectra of ANAP and GFP tagged SUR1 subunits expressed in unroofed membrane fragments were

acquired as above. We were able to resolve a GFP peak at 510 nm with 385 nm excitation, even though

this wavelength is far from the λ_{max} for GFP. It is possible that GFP emission was enhanced via FRET

between ANAP and GFP. The ANAP peak fluorescence at 470 nm did not overlap with the GFP emission

peak. However, the peak GFP at 510 nm was contaminated by the shoulder of the ANAP peak. Therefore,

the GFP fluorescence was corrected by subtracting the averaged (from 16 membranes) fluorescence of

ANAP incorporated into SUR1, scaled to match the 470 nm peak in the ANAP-GFP spectrum.

To measure the time course of nucleotide unbinding, the fluorescence at the TNP-nucleotide peak

(usually around 560 nm) was plotted as a function of time after exchanging to a zero-nucleotide solution,

as following the time course of the rise in ANAP fluorescence was complicated by photobleaching. After

complete wash out of nucleotides, the fluorescence at 560 nm was non-zero, as the shoulder of the ANAP

emission spectrum extends into the range of wavelengths at which TNP-nucleotides emit. Therefore, the

time course data were corrected by subtracting a linear fit to the data following complete washout. This

corrected for the shoulder of the ANAP spectrum as well as the continued photobleaching of ANAP. Data

were normalized to the maximum fluorescence at t = 0 and fit with single-exponential decays

$$\frac{F}{F_{max}} = e^{-t/\tau} \ (2)$$

10   or double-exponential decays

$$\frac{F}{F_{max}} = A_1 e^{-t/\tau_1} + A_2 e^{-t/\tau_2} \ (3).$$

Wash-out time courses following exchange to [TNP-nucleotide] = 0 should be independent of the

nucleotide concentration before wash. The majority of our time-course experiments were conducted at the

conclusion of our concentration-response experiments, and therefore reflect wash after applying a

concentration of 1 mM. For experiments in diazoxide and tolbutamide, the nucleotide concentration prior

to wash was 50 μM.

In some experiments, there was evidence of cross-contamination of our wash solution or low-nucleotide

solutions via back-flow of solutions with higher concentrations of nucleotide (as evident by accumulation

of yellow TNP-nucleotides). We excluded data from such experiments and maintained/replaced valves on

the μFlow as needed.

Electrophysiology. Currents were recorded from inside-out membrane patches excised from transfected

HEK-293T cells. Pipettes were pulled to a resistance of 1-5 MΩ and filled with an extracellular solution

containing (in mM) 140 KCl, 1.2 $MgCl_2$, 2.6 $CaCl_2$, and 10 HEPES, pH 7.4 or 140 KCl, 1 EGTA, 10

HEPES, pH 7.3. The intracellular (bath) solutions contained either (in mM) 107 KCl, 2 $MgCl_2$, 1 $CaCl_2$,

10 EGTA, and 10 HEPES, pH 7.2 or 140 KCl, 1 EDTA, 1 EGTA, 10 HEPES pH 7.3. Nucleotides were

added as indicated. Patches were perfused with an 8-channel µFlow perfusion system. Data were acquired

at a holding potential of -60 mV using an Axopatch 200B amplifier and Digidata 1322A digitizer with

pClamp 9.0 software (Molecular Devices). Currents were digitized at 10 kHz and low-pass filtered at 1

kHz. K_{ATP} channels run down in excised patches (reviewed by Proks *et al.*).[46] For inhibitory

concentration-response relationships, rundown was corrected by alternating test nucleotide solutions with

nucleotide-free (control) solutions and expressing the test currents as a fraction of the average of the

control currents before and after the test solution. Corrected data were fit to the following expression:

13
$$\frac{I}{I_{max}} = 1 - \frac{[TNP]^h}{IC_{50}{}^h + [TNP]^h} \quad (4).$$

14   With IC_{50} representing the half-maximal inhibitory concentration. For activation concentration-responses,

15   the same protocol was used, but test concentrations were alternated with a saturating concentration (1

16   mM) of MgADP.[27] Data were displayed at the current magnitude in the test concentration minus the

17   current in zero nucleotide (I-I_{min}) divided by the current in saturating nucleotide (1 mM MgADP) minus

18   the current in zero nucleotide ($I_{max} - I_{min}$) and fit to the following expression:

19
$$\frac{I - I_{min}}{I_{max} - I_{min}} = \frac{[TNP]^h}{EC_{50}{}^h + [TNP]^h} \quad (5)$$

Currents were leak corrected by subtracting the remaining current after complete rundown.

Western Blotting. In order to detect SUR1 via Western blot, a 3xFLAG

(MDYKDHDGDYKDHDIDYKDDDDK) tagged construct (tag in an extracellular loop between amino

acid positions T1042 and L1043) was used (3xFLAG_SUR1) with or without amber stop codons at amino

acid positions 1353 or 1397. HEK-293T cells were transfected in 6-well plates using the conditions

described above, but with 1 µg of SUR1 DNA. After 48 h, cells were washed and harvested with gentle

pipetting in 1 mL of PBS (Sigma). Cells were pelleted (4 min at 200 x *g* yielding 20-50 mg pellets) and

solubilized in 50-100 µL of 0.5% Triton X-100 in 110 mM potassium acetate, buffered at pH 7.4. 125 U

of benzonase (Sigma) was added to each sample and samples were incubated at room temperature for 20

min. 7.5 µL of each sample was mixed with 2.5 µL of NuPAGE LDS sample buffer and 1 µL NuPAGE

sample reducing agent (Invitrogen) and run on a NuPAGE 4-12% Bis-Tris gel (Invitrogen) at 200 V for

40 min in 1X MOPS-SDS buffer (Invitrogen). Protein was transferred overnight at 10 V to an Immobilon-

P membrane (Merck Millipore; Burlington, MA) in 25 mM Tris, 192 mM glycine, and 20% methanol +

0.1% SDS. Following transfer, the membrane was rinsed in TBS-Tw (150 mM NaCl, 25 mM Tris, pH

7.2, and 0.05% Tween 20) and incubated (with shaking) for 30 min in TBS-Tw + 5% milk. The

membrane was rinsed three times in TBS-Tw and incubated for 30 min with primary antibody (M2 anti-

FLAG from Sigma, diluted 1:500 in TBS-Tw). The membrane was rinsed three more times in TBS-Tw

and incubated for 30 min with secondary antibody (HRP-conjugated sheep anti-mouse IgG, GE

Healthcare Life Sciences; Freiburg, Germany) diluted 1:20,000 in TBS-Tw + 1% milk. Finally, the

membrane was rinsed 3x10 min in TBS-Tw and developed using SuperSignal West Femto Max

Sensitivity Substrate (Thermo Fisher Scientific). Images were collected using a C-DiGit scanner (Licor

Biosciences; Lincoln NE) and band intensities were quantified using custom code written in MATLAB

(https://github.com/mpuljung/spectra-analysis).[62]

Western blotting was performed (Figure 1—figure supplement 1d) on the total protein from HEK-

293T cells transfected with Kir6.2 and 3xFLAG-tagged SUR1-Y1353[stop] DNA under various conditions.

As the 3xFLAG tag was N-terminal to SUR1, we used it to probe for both full-length and truncated

SUR1. As a positive control, we transfected 3x-FLAG-tagged SUR1 with no stop codon, which was only

expected to produce full-length protein (lane 1). This protein ran above our 140 kDa marker as expected

(predicted MW is ~180 kDa). In the absence of ANAP or in cells transfected without the pANAP

plasmid, only truncated 3xFLAG_SUR1 protein was evident (lanes 4 and 5), confirming that there was no

full-length protein produced by accidental read-through of the amber (TAG) codon. In the presence of

both pANAP and ANAP, we obtained a mixture of full-length (66%) and truncated (34%) SUR1. The

majority of our experiments in unroofed membrane fragments were performed under such conditions as

the signals we measured (fluorescence) were derived solely from full-length (i.e. ANAP-labeled)

channels. Finally, we were able to increase expression of full-length SUR1 as needed by transfecting an

additional plasmid, peRF1-E55D, which encodes a dominant negative ribosomal release factor.[36] With the

additional expression of ERF1-E55D, we obtained 92% full-length protein (Figure 1—figure supplement

1d). A similar approach was taken with SUR1-T1397[stop] (Figure 4—figure supplement 1d).

In parallel experiments, we expressed SUR1_GFP-Y1353[stop] in the absence of ANAP. Under such

conditions, we observed a few cells with diffuse GFP fluorescence throughout the cytoplasm. We believe

that the signal from such cells was derived from soluble GFP generated from an internal methionine after

position 1353 and not read-through of the amber stop codon, as the distribution of this protein differed

from that of SUR1_GFP (which was confined to the plasma membrane and presumably ER/Golgi). We

never observed GFP fluorescence in unroofed membranes of such cells.

Surface Expression Assay. Surface expression of SUR1 constructs was assayed by their ability to

chaperone HA-tagged Kir6.2 subunits to the plasma membrane using an adaptation of the method of

Zerangue *et al.*[35] The HA tag plus linker (YAYMEKGITDLAYPYDVPDY) was inserted in the

extracellular region following helix M1 of Kir6.2 between amino acids L100 and A101. HEK-293T cells

were cultured in 12-well plates (STARLAB) on 19 mm cover slips treated with poly-L-lysine.

Transfections were performed as described above, but DNA amounts were adjusted for the smaller scale

(0.08 μg HA-tagged Kir6.2, 0.14 μg SUR1 constructs, 0.14 μg pANAP, 0.14 μg peRF1-E55D). Cells

were incubated in DMEM (+10% FBS, 100 U/mL penicillin, and 100 μg/mL streptomycin) at 33ºC in a

95% air/5% CO_2 atmosphere for 48 hours in the presence or absence (as indicated) of 20 μM ANAP.

Following incubation, cells were rinsed on ice with PBS and fixed for 30 min in PBS + 10% neutral

buffered formalin. Cells were subsequently washed twice in PBS and blocked with 1% bovine serum

albumin (BSA) in PBS for 30 min at 4º C. Cells were then incubated for 1 h at 4º C in PBS + 1% BSA

with a 1:1000 dilution of rat anti-HA monoclonal antibody (Roche; Basel, Switzerland) and washed 5x10

min on ice with PBS + 1% BSA. HRP-conjugated goat anti-rat polyclonal antibody (diluted 1:2000,

Jackson ImmunoResearch; Ely, UK) was applied in PBS + 1% BSA and cells were incubated an

additional 30 min at 4º C. Finally, cells were washed 4x10 min in PBS + 1% BSA and 5x10 min in PBS

on ice. After the wash, coverslips were removed from PBS and placed in clean, untreated 35 mm culture

dishes. 300 μL of SuperSignal ELISA Femto Maximum Sensitivity Substrate (Thermo Fisher Scientific)

were added and the luminescence was measured using a Glomax 20/20 Luminometer (Promega; Madison,

WI) after a 10 s incubation. For display purposes, data were normalized to the mean of the luminescence

with wild-type SUR1. Background was assessed with cells that were transfected with HA-tagged Kir6.2,

but no SUR1.

Chemicals. ANAP trifluoroacetate salt and methyl ester were obtained from Asis Chemicals (Waltham,

MA). A 1 mM stock of the salt was prepared in 30 mM NaOH and stored at -80º C. The methyl ester was

dissolved at 5 mM in DMSO and stored at -20º C. The two forms were used interchangeably, as the final

product (when ANAP is incorporated into a protein) is identical and in our experience both free acid and

methyl ester were membrane permeant. Trinitrophenyl (TNP) nucleotide analogues were purchased from

Jena Bioscience (Jena, Germany). 1 mM stocks were prepared in buffer and stored at -20º C. Stocks were

verified using UV-Vis spectroscopy (Beckman Coulter DU800 spectrophotometer; Pasadena, CA) as

needed. Paper chromatography (mobile phase 40:10:25 *n*-butanol, glacial acetic acid: H_2O) was used to

verify that older TNP-ATP stocks had not undergone hydrolysis. All other chemicals were from Sigma.

Diazoxide was prepared in a 34 mM stock in 100 mM KOH and stored at -20º C before use. Tolbutamide

was dissolved at 100 mM in DMSO and stored at -20º C.

Data presentation and statistics. Error bars represent ±SEM. The boxes in Figure 2e and Figure 4—

figure supplement 1a,b represent ±SD. The number of experiments (n) represents the number of patches

or membranes used in a given experiment. These can be found in the tables for binding,

electrophysiology, and time course data, and figure legends for other experiments. For surface expression

assays, n represents the number of transfected coverslips. Plots and curve fitting were generated using

Origin 9.1 (OriginLab Corporation; Northampton, MA). Curves in the figures (and parameters reported in

the figure legends) represent fits to the combined data sets from multiple experiments, with errors

representing the standard error of the fits. Shaded regions indicate 95% confidence intervals. Fit

parameters reported in Tables are the mean ±SEM (Tables 1, 2, and 5) or ±SD (Tables 3 and 4) of the fits

to individual experiments. Pairwise comparisons were performed using two-tailed Student's *t*-tests with

the Welch correction (not assuming equal variance) and the Bonferroni correction for multiple

comparisons, as needed. Comparisons of EC_{50} values were performed on log-transformed data as the

untransformed EC_{50} is not normally distributed.

Acknowledgments

We wish to thank Idoia Portillo and Raul Terron Exposito for excellent technical assistance and

Gregor Sachse, Chris Miller, and David Gadsby for helpful discussions. James Cantley provided access to

the Western blot scanner. This work was supported by the European Research Council (grant 332620),

the Biotechnology and Biological Sciences Research Council (BB/R002517/1), the John Fell Fund, and

the Wellcome Trust Oxion graduate program.

Competing Interests

We declare no competing interests.

Figure Legends

Figure 1. Measuring binding to NBS2 of SUR1. **(a)** Cartoon illustrating the topology of the K_{ATP}

channel complex. Two (of four) Kir6.2 subunits and one (of four) SUR1 subunits are shown for clarity.

The locations of the inhibitory (red, on Kir6.2) and stimulatory (green, on SUR1) NBSs are indicated. **(b)**

Chemical structures of TNP-ATP and ANAP. The fluorescent moiety is colored in each. **(c)** Structure of

the ligand-bound NBDs of SUR1 (from PDB accession # 6C3O).[3] TNP-ATP (from PDB accession #

3AR7) was positioned in both NBSs of SUR1 by alignment with MgATP (at NBS1) and MgADP (at

NBS2), which were present in the original structure.[63] Y1353 from the A-loop of NBS2 is highlighted in

yellow to mark the location of ANAP in our experiments. Alignment of TNP-ATP with nucleotides

bound to NBS1 and NBS2 in the cryo-EM structure of SUR1 does not predict any structural clashes. **(d)**

Representative spectral images acquired from an unroofed membrane fragment expressing SUR1-

Y1353*/Kir6.2 channels in the absence or presence of increasing concentrations of TNP-ADP. **(e)**

Averaged spectra from (d). **(f)** Concentration-response relationships for binding of MgTNP-ATP

(magenta) and MgTNP-ADP (cyan) to SUR1-Y1353*/Kir6.2. MgTNP-ATP: Combined data from

multiple experiments in (f) and (g) data were fit to equation 1. $E_{max} = 1.00 \pm 0.05$, $EC_{50} = 29.3$ µM ± 6.5

µM, $h = 0.8 \pm 0.1$. MgTNP-ADP: $E_{max} = 0.99 \pm 0.02$, $EC_{50} = 4.8$ µM ± 0.5 µM, $h = 0.8 \pm 0.1$. **(g)**

Concentration-response relationships for binding of TNP-ATP and TNP-ADP to SUR1-Y1353*/Kir6.2

channels in the absence of Mg^{2+}. TNP-ATP: $E_{max} = 0.84 \pm 0.04$, $EC_{50} = 8.5$ µM ± 2.4 µM, $h = 0.8 \pm 0.1$.

TNP-ADP: $E_{max} = 0.95 \pm 0.02$, $EC_{50} = 6.1$ µM ± 0.7 µM, $h = 0.8 \pm 0.1$. Increasing the concentration of

TNP-ADP to 10 mM did not significantly change the amount of quenching (n = 3, data not shown).

Figure 2. Conditions that favor channel opening slow nucleotide dissociation from NBS2. **(a)** Average

dissociation time course of TNP-ADP from SUR1-Y1353*/Kir6.2 K_{ATP} channels in the presence (orange)

and absence (gray) of Mg^{2+}. TNP-ADP: $\tau = 4.2$ s ± 0.2 s; MgTNP-ADP: $\tau = 21.3$ s ± 0.8 s. **(b)** Dissociation

time course of TNP-ATP from SUR1-Y1353*/Kir6.2 channels in the presence and absence of Mg^{2+}.

TNP-ATP: $\tau_1 = 4.5$ s ± 0.9 s, $A_1 = 0.65 \pm 0.06$, $\tau_2 = 28.2$ s ± 3.7 s, $A_2 = 0.35 \pm 0.06$; MgTNP-ATP: $\tau = 15.0$

s ±0.5 s. **(c,d)** Dissociation time course of MgTNP-ADP (c) or MgTNP-ATP (d) from SUR1-

Y1353*/Kir6.2 in the presence of 500 µM tolbutamide (pink) or 340 µM diazoxide (teal). (c) The orange

curve is the fit to the MgTNP-ADP data from (a). Tolbutamide: τ = 6.4 s ±0.2 s; diazoxide: τ = 52.6 s

±1.4 s. (d) The orange curve is the fit to MgTNP-ATP data from (b). Tolbutamide: τ = 6.3 s ±0.1 s;

diazoxide: τ = 60.5 s ±1.4 s. **(e)** Time to 50% dissociation ($t_{0.5}$) for the individual fits to the data shown in

(a)-(d) for better comparison between single-exponential and more complex time courses. Boxes

represent mean ±SD. Individual data points are shown to the right of the boxes. Combined data from

multiple experiments in (a)-(d) were fit to single exponential decays (equation 2). Dissociation of TNP-

ATP in the absence of Mg^{2+} was better fit with a bi-exponential decay (equation 3).

Figure 3. Mutation of the Walker$_A$ motif of NBS2 affects apparent the affinity for MgTNP-ADP. **(a)**

Interaction of the Walker$_A$ lysine (K1384) with MgADP bound to NBS2 of SUR1 (from PDB accession #

6C3O).[3] The K1384A (K2A) mutation (modeled *below*) disrupts this interaction. **(b)** Binding of TNP-

ATP to SUR1-Y1353*,K2A/Kir6.2 K_{ATP} channels in the absence of Mg^{2+}. Combined data from multiple

experiments in (b)-(e) were fit to equation 1. E_{max} = 0.87 ±0.03, EC_{50} = 4.3 µM ±0.1 µM, h = 0.9 ±0.2.

Gray curves are fits to SUR1-Y1353*/Kir6.2 channels from Figure 1. **(c)** Binding of TNP-ADP to SUR1-

Y1353*,K2A/Kir6.2 in the absence of Mg^{2+}. E_{max} = 0.91 ±0.02, EC_{50} = 4.4 µM ±0.1 µM, h = 0.9 ±0.1. **(d)**

Binding of MgTNP-ATP to SUR1-Y1353*,K2A/Kir6.2. E_{max} = 0.97 ±0.06, EC_{50} = 21.5 µM ±6.7 µM, h =

0.8 ±0.1. **(e)** Binding of MgTNP-ADP to SUR1-Y1353*,K2A/Kir6.2. E_{max} = 0.96 ±0.04, EC_{50} = 12.5 µM

±2.5 µM, h = 0.9 ±0.1. **(f)** Dissociation time course of MgTNP-ADP (green, filled symbols) and TNP-

ADP (green, open symbols) from SUR1-Y1353*,K2A/Kir6.2. Combined data from multiple experiments

were fit to a single exponential; τ = 13.5 s ±0.3 s (for MgTNP-ADP) and τ = 5.4 s ±0.8 s (for TNP-ADP).

Fits to the dissociation of TNP-ADP from SUR1-Y1353*/Kir6.2 in the presence (orange) and absence

(gray) of Mg^{2+} from Figure 2 are shown for comparison.

Figure 4. Binding of TNP-nucleotides to SUR1-T1397*/Kir6.2 K_{ATP} channels. (a) Structure of the

ligand-bound NBDs of SUR1 (from PDB accession # 6C3O) with TNP-ATP aligned as in Figure 1c.[3]

T1397 in NBD2 is highlighted in yellow to mark the placement of ANAP in our experiments. Combined

data from multiple experiments in (c) and (d) were fit to equation 1. **(b)** Concentration-response

relationships for binding of MgTNP-ATP (magenta) and MgTNP-ADP (cyan) to SUR1-T1397*/Kir6.2

channels. MgTNP-ATP: E_{max} = 1.00 ±0.02, EC_{50} = 14.7 μM ±1.8 μM, h = 1.0 ±0.1. MgTNP-ADP: E_{max} =

1.0 ±0.01, EC_{50} = 10.6 μM ±0.07 μM, h = 0.97 ±0.06. **(c)** Concentration-response relationships for

binding of TNP-ATP and TNP-ADP to SUR1-Y1397*/Kir6.2 channels in the absence of Mg^2. TNP-ATP:

E_{max} = 0.94 ±0.02, EC_{50} = 4.8 μM ±0.7 μM, h = 0.8 ±0.1. TNP-ADP: E_{max} = 0.98 ±0.05, EC_{50} = 5.6 μM

±1.7 μM, h = 0.7 ±0.1. **(d)** Activation of SUR1-T1397*/Kir6.2-G334D channels by MgADP in inside-out

patches. Combined data from multiple experiments were fit with equation 5. EC_{50} = 38.6 μM ±5.8 μM, h

= 1.2 ±0.2.

Figure 5. Activation of K_{ATP} by Mg-nucleotide binding at SUR1 proceeds via an MWC-type reaction.

(a) Continuous current trace from an inside out patch expressing SUR1-T1397*/Kir6.2-G334D showing

activation by MgADP and MgTNP-ADP. The dotted line indicates the zero-current level. Downward

deflections indicated increasing current. **(b)** Combined data from multiple experiments for MgTNP-ADP

binding (cyan from Figure 4b; SUR1-T1397*/Kir6.2) and current activation (gray from SUR1-

T1397*/Kir6.2-G334D in separate experiments) were fit simultaneously with MWC expressions for

binding (cyan, equation 6) and activation (gray, equation 7) with shared parameters for the two

expressions.. The schematic on the right describes the MWC model. Closed K_{ATP} channels are drawn as

squares and open channels as circles. Filled symbols represent subunits bound to MgTNP-ADP

(abbreviated as MgADP). Channels open by a concerted flip of all four subunits. L = 0.52, E = 2.2 ±0.2,

and K_A = 5.8x10^4 M^{-1} ±1.0x10^4 M^{-1}. Each MgTNP-ADP binding event is independent and each MgTNP-

ADP-bound NBS2 contributes equally to increase P_{open}.

Figure 6. Schematic for activation of K_{ATP} by nucleotide binding to SUR1. **(top, left)** In the absence of

Mg^{2+} nucleotides bind and rapidly dissociate from the NBSs of SUR1, without affecting its conformation.

Nucleotide occupancy of SUR1 is not communicated to Kir6.2. **(top, right)** In the presence of both Mg^{2+}

and nucleotide, the NBD dimer interface is stabilized. This interface must open in order for nucleotides to

leave, so nucleotide dissociation is greatly slowed. NBD dimerization requires a conformational change in

the transmembrane domain of SUR1. This conformational change is communicated to Kir6.2 to increase

the favorability of channel opening. (***bottom, left***) Sulfonylureas prevent the conformational change in the

transmembrane domain of SUR1 and preclude NBD dimerization. Therefore, nucleotide dissociation is

rapid, even in the presence of Mg^{2+} and no increase in open probability results from Mg-nucleotide

binding to SUR1. (***bottom, right***) Potassium channel openers stabilize the closed-dimer conformation of

the NBDs. As a result, nucleotide dissociation is very slow and channel activation is enhanced.

Figure 1—figure supplement 1. ANAP fluorescence is specific to SUR-Y1353* in unroofed membrane

fragments. **(a)** *above.* Brightfield (BF) and fluorescence images of an unroofed membrane fragment from

an HEK-293T cell expressing SUR1-Y1353*/Kir6.2. The dashed line indicates the position of the

spectrograph slit. *below.* Image of the fluorescence from the same membrane after the light was diffracted

by the gratings of the spectrograph. Spatial information in the *y* dimension is preserved. Spatial

information in the *x* dimension is replaced by wavelength. **(b)** Brightfield (BF) and fluorescence images

of unroofed membrane fragments from HEK-293T cells expressing SUR1_GFP-Y1353*/Kir6.2 (*above*)

and SUR1_GFP/Kir6.2 (*below*). Fluorescence images were acquired and scaled identically for each

fluorophore. Note the absence of ANAP fluorescence in the membrane expressing SUR1_GFP, despite

presence of ANAP in the cell culture medium. **(c)** Intensity of ANAP fluorescence as a function of GFP

fluorescence (n = 34). Data were fit with a straight line with the intercept fixed to 0 and a slope of 0.96

±0.02 (R = 0.99). **(d)** Western blot for 3xFLAG-tagged SUR1 subunits expressed in HEK-293T cells. All

cells were co-transfected with Kir6.2. Unless otherwise indicated, all cells were cultured in 20 μM

ANAP. The first lane (positive control) was 3xFLAG_SUR1 with no amber stop codon at position 1353,

which shows only full-length protein. The next two lanes show 3xFLAG_SUR1-Y1353^{stop} co-transfected

with pANAP and peRF1-E55D (92% full-length SUR1), or with pANAP alone (66% full-length SUR1).

Omission of pANAP (lane 5) or exclusion of ANAP from the cell culture medium (lane 4) resulted in

only truncated protein. Background signal is shown in the last two lanes in which either no SUR1 was

transfected or untagged SUR1 was used. The positions of the full-length and truncated protein (at amino

acid 1352) are marked. Upper bands (~260 kDa and higher) are from aggregated SUR1 protein.

Figure 1—figure supplement 2. TNP-nucleotides are suitable for measuring nucleotide binding to

K_{ATP}. **(a)** Absorbance spectra (dashed lines) and emission spectra (solid lines) for TNP-ADP (orange) and

ANAP (cyan). Note the overlap between the absorbance spectrum of TNP-ADP (orange, dashed) and the

emission spectrum of ANAP (cyan, solid). The ANAP absorption spectrum and the TNP-ADP spectra

were acquired in aqueous solution. ANAP emission was measured from SUR1-Y1353*/Kir6.2 in

unroofed membrane fragments. **(b)** Theoretical FRET efficiency as a function of distance calculated from

the spectral overlap shown in (a). Half maximal FRET efficiency (R_0) occurs at 43 Å. κ^2 was assumed to

be 0.67 (i.e. random orientation between the donor and acceptor was assumed) and the quantum yield of

ANAP is assumed to be 0.22.[22] **(c)** Representative trace showing activation of SUR1/Kir6.2-G334D K_{ATP}

channels in an inside-out patch by MgADP and MgTNP-ADP. **(d)** Activation of SUR1/Kir6.2-G334D

currents by MgTNP-ADP in inside-out patches. Combined data from multiple experiments were fit to

equation 5. *EC_{50}* = 22.1 μM ±3.0 μM, *h* = 0.9 ±0.1. **(e)** MgTNP-ATP "binding" to an unroofed membrane

fragment from an HEK-293 cell transfected with pANAP (but no Kir6.2 or SUR1) and incubated in

media with 20 μM ANAP. No ANAP fluorescence was observed, but non-specific MgTNP-ATP

fluorescence was observed at nucleotide concentrations of 50 μM and higher. **(f)** Spectra of SUR1-

Y1353*/Kir6.2 channels in an unroofed membrane fragment in the absence or presence of 10 mM

MgATP. The intensity in 10 mM MgATP was 98% ±2% of the original peak (n = 7) and 101% ±2% in 10

mM MgADP (n = 8).

Figure 1—figure supplement 3. Mutation of the inhibitory nucleotide site in Kir6.2 does not affect

binding measured at NBS2 of SUR1. **(a)** Proximity of the inhibitory NBS of Kir6.2 to NBS2 of SUR1 in

the Mg-nucleotide-bound structure of K_{ATP} (PDB accession # 6C3O).[3] TNP-ATP (from PDB accession #

3AR7) was aligned with ATP in the original structure.[63] The dashed line represents a distance of 33 Å. **(b)**

Concentration-response relationship for binding of MgTNP-ADP to SUR1-Y1353*/Kir6.2-G334D

channels. Combined data from multiple experiments were fit with equation 1: $E_{max} = 1.00 \pm 0.04$, $EC_{50} =$

7.7 μM ±1.7 μM, $h = 0.82 \pm 0.13$. The fits to the concentration-response relationship for MgTNP-ADP

binding to SUR1-Y1353*/Kir6.2 channels from Figure 1f are shown in gray.

Figure 1—figure supplement 4. Mg^{2+} effects on steady-state nucleotide binding to SUR1-

Y1353/Kir6.2.* **(a)** Concentration-response relationships for TNP-ADP binding to SUR1-Y1353*/Kir6.2

in the presence and absence of Mg^{2+}. **(b)** Concentration-response relationships for TNP-ATP binding to

SUR1-Y1353*/Kir6.2 in the presence and absence of Mg^{2+}. Data and fits are repeated from Figure 1.

Figure 1—figure supplement 5. Photobleaching correction. **(a)** Successive spectra of SUR1-

Y1353*/Kir6.2 before (left) and after (right) correction for photobleaching. The times indicated are

cumulative exposure time. The inset on the right shows the exponential fit to the normalized peak

fluorescence from the image on the left. **(b)** Amplitudes (left) and time constants (right) from exponential

fits to photobleaching data. Y1353*, A= 0.4 ±0.1, τ = 29.6 s ±23.9 s, n = 23; Y1353*,K2A, A = 0.7 ±0.6,

τ = 77.1 s ±92.2 s, n = 24; T1397*, A = 0.6 ±0.1, τ = 46.2 s ±16.6 s, n = 33. Lines indicate means.

Figure 3—figure supplement 1. Mg^{2+} effects on steady-state nucleotide binding to SUR1-

Y1353,K2A/Kir6.2.* **(a)** Concentration-response relationships for TNP-ADP binding to SUR1-

Y1353*,K2A/Kir6.2 in the presence and absence of Mg^{2+}. **(b)** Concentration-response relationships for

TNP-ATP binding to SUR1-Y1353*,K2A/Kir6.2 in the presence and absence of Mg^{2+}. Data and fits are

repeated from Figure 3.

Figure 4—figure supplement 1. SUR1-T1397/Kir6.2 K$_{ATP}$ channels traffic to the plasma membrane*

and form functional channels. **(a)** Chemiluminescence assay for surface expression of HA-tagged Kir6.2

in the presence (n = 11) and absence (n = 11) of SUR1, in SUR1-Y1353[stop]-transfected HEK-293T cells

cultured in the presence (n = 8) or absence (n = 8) of 20 μM ANAP, and in SUR1-T1397[stop] transfected

HEK-293T cells cultured in the presence (n = 7) or absence (n = 6) of 20 μM ANAP. SUR1-Y1353[stop]

and SUR1-T1397[stop] were co-transfected with pANAP/peRF1-E55D. **(b)** Current (normalized to

nucleotide-free solution) from inside-out patches excised from cells expressing 3xFLAG_SUR1/Kir6.2

(SUR1), SUR1-Y1353[stop]/Kir6.2 alone, SUR1-Y1353[stop]/Kir6.2 with pANAP, or SUR1-Y1353[stop]/Kir6.2

with pANAP/peRF1-E55D. All cells expressing SUR1-Y1353[stop] were cultured with ANAP in the media.

A response >1 indicates activation. Responses <1 indicate inhibition. **(c)** Currents from inside out patches

excised from cells expressing either SUR1/Kir6.2-G334D or SUR1-Y1353*/Kir6.2-G334D. MgADP and

tolbutamide applied as indicated. **(d)** Western blot for 3xFLAG-tagged SUR1 subunits expressed in HEK-

293T cells. All cells were co-transfected with Kir6.2 and cultured in the presence of 20 μM ANAP. Two

separate experiments are shown. In each, the positive control was 3xFLAG_SUR1 with no amber stop

codon. The negative control was SUR1 with no FLAG tag. The next two lanes show 3xFLAG_SUR1-

T1397[tag] co-transfected with pANAP and peRF1-E55D, or with peRF1-E55D and no pANAP. The

positions of the full-length and truncated (at amino acid 1396) proteins are marked. **(e)** Inhibition of

SUR1-T1397*/Kir6.2 (gray) and SUR1/Kir6.2 (brown) channels by ATP in the absence of Mg^{2+}

measured in inside-out patches. Combined data from multiple experiments were fit to equation 4. SUR1-

T1397*: IC_{50} = 49.2 μM ±4.7 μM, h = -1.1 ±0.1. SUR1: EC_{50} = 11.1 μM ±6.5 μM, h = -1.3 ±0.1.

***Figure 4—figure supplement 2. Mg^{2+} effects on steady-state nucleotide binding to SUR1-T1397*.* (a)**

Concentration-response relationships for TNP-ADP binding to SUR1-T1397*/Kir6.2 in the presence and

absence of Mg^{2+}. **(b)** Concentration-response relationships for TNP-ATP binding to SUR1-T1397*/Kir6.2

in the presence and absence of Mg^{2+}. Data and fits are repeated from Figure 4.

Figure 5—figure supplement 1. Fits to alternate gating models. Combined binding and gating data from

Figure 5 fit to alternative models. Fit parameters are shown as insets. Goodness of fit was assessed as the

residual sum of squares (RSS). **(a)** MWC model as in Figure 5. **(b)** Model in which binding of MgADP to

a single SUR1 is sufficient to increase channel activity.

24
$$\frac{bound}{total} = \frac{LE[ADP]K_A(1 + 3[ADP]K_A + 3[ADP]^2K_A^2 + [ADP]^3K_A^3) + [ADP]K_A([ADP]K_A + 1)^3}{L(1 + 4E[ADP]K_A + 6E[ADP]^2K_A^2 + 4E[ADP]^3K_A^3 + E[ADP]^4K_A^4) + ([ADP]K_A + 1)^4}$$

$$P_{open,norm} = \frac{\left(\frac{L(1 + 4E[ADP]K_A + 6E[ADP]^2K_A^2 + 4E[ADP]^3K_A^3 + E[ADP]^4K_A^4)}{L(1 + 4E[ADP]K_A + 6E[ADP]^2K_A^2 + 4E[ADP]^3K_A^3 + E[ADP]^4K_A^4) + ([ADP]K_A + 1)^4}\right) - \left(\frac{L}{L+1}\right)}{\left(\frac{LE}{LE+1} - \frac{L}{L+1}\right)}$$

2 **(c)** Model in which binding of MgADP to two SUR1 subunits is required to increase channel activity.

$$\frac{bound}{total} = \frac{L[ADP]K_A(1 + 3E[ADP]K_A + 3E[ADP]^2K_A^2 + E[ADP]^3K_A^3) + [ADP]K_A([ADP]K_A + 1)^3}{L(1 + 4[ADP]K_A + 6E[ADP]^2K_A^2 + 4E[ADP]^3K_A^3 + E[ADP]^4K_A^4) + ([ADP]K_A + 1)^4}$$

$$P_{open,norm} = \frac{\left(\frac{L(1 + 4[ADP]K_A + 6E[ADP]^2K_A^2 + 4E[ADP]^3K_A^3 + E[ADP]^4K_A^4)}{L(1 + 4[ADP]K_A + 6E[ADP]^2K_A^2 + 4E[ADP]^3K_A^3 + E[ADP]^4K_A^4) + ([ADP]K_A + 1)^4}\right) - \left(\frac{L}{L+1}\right)}{\left(\frac{LE}{LE+1} - \frac{L}{L+1}\right)}$$

5 **(d)** Model in which binding of MgADP to three SUR1 subunits is required to increase channel activity.

$$\frac{bound}{total} = \frac{L[ADP]K_A(1 + 3[ADP]K_A + 3E[ADP]^2K_A^2 + E[ADP]^3K_A^3) + [ADP]K_A([ADP]K_A + 1)^3}{L(1 + 4[ADP]K_A + 6[ADP]^2K_A^2 + 4E[ADP]^3K_A^3 + E[ADP]^4K_A^4) + ([ADP]K_A + 1)^4}$$

$$P_{open,norm} = \frac{\left(\frac{L(1 + 4[ADP]K_A + 6[ADP]^2K_A^2 + 4E[ADP]^3K_A^3 + E[ADP]^4K_A^4)}{L(1 + 4[ADP]K_A + 6[ADP]^2K_A^2 + 4E[ADP]^3K_A^3 + E[ADP]^4K_A^4) + ([ADP]K_A + 1)^4}\right) - \left(\frac{L}{L+1}\right)}{\left(\frac{LE}{LE+1} - \frac{L}{L+1}\right)}$$

8 **(e)** Model in which all four SUR1 subunits must be occupied by MgADP in order to increase channel

9 open probability.

$$\frac{bound}{total} = \frac{L[ADP]K_A(1 + 3[ADP]K_A + 3[ADP]^2K_A^2 + E[ADP]^3K_A^3) + [ADP]K_A([ADP]K_A + 1)^3}{L(1 + 4[ADP]K_A + 6[ADP]^2K_A^2 + 4[ADP]^3K_A^3 + E[ADP]^4K_A^4) + ([ADP]K_A + 1)^4}$$

$$P_{open,norm} = \frac{\left(\frac{L(1 + 4[ADP]K_A + 6[ADP]^2K_A^2 + 4[ADP]^3K_A^3 + E[ADP]^4K_A^4)}{L(1 + 4[ADP]K_A + 6[ADP]^2K_A^2 + 4[ADP]^3K_A^3 + E[ADP]^4K_A^4) + ([ADP]K_A + 1)^4}\right) - \left(\frac{L}{L+1}\right)}{\left(\frac{LE}{LE+1} - \frac{L}{L+1}\right)}$$

12 *Figure 5—figure supplement 2. Residuals of fits to alternate gating models.* Simultaneous fits of

13 binding and activation data as in Figure 5—figure supplement 1, with accompanying plots of the residuals

14 of the model fits.

References

1. Ashcroft, F.M. & Rorsman, P. K(ATP) channels and islet hormone secretion: new insights and controversies. *Nat Rev Endocrinol* **9**, 660-9 (2013).

2. Quan, Y., Barszczyk, A., Feng, Z.P. & Sun, H.S. Current understanding of K ATP channels in neonatal diseases: focus on insulin secretion disorders. *Acta Pharmacol Sin* **32**, 765-80 (2011).

3. Lee, K.P.K., Chen, J. & MacKinnon, R. Molecular structure of human KATP in complex with ATP and ADP. *Elife* **6**(2017).

4. Martin, G.M., Kandasamy, B., DiMaio, F., Yoshioka, C. & Shyng, S.L. Anti-diabetic drug binding site in a mammalian KATP channel revealed by Cryo-EM. *Elife* **6**(2017).

5. Puljung, M.C. Cryo-electron microscopy structures and progress toward a dynamic understanding of KATP channels. *J Gen Physiol* **150**, 653-669 (2018).

6. Wu, J.X., Ding, D., Wang, M., Kang, Y., Zeng, X. & Chen, L. Ligand binding and conformational changes of SUR1 subunit in pancreatic ATP-sensitive potassium channels. *Protein Cell* **9**, 553-567 (2018).

7. Inagaki, N., Gonoi, T., Clement, J.P.t., Namba, N., Inazawa, J., Gonzalez, G., Aguilar-Bryan, L., Seino, S. & Bryan, J. Reconstitution of IKATP: an inward rectifier subunit plus the sulfonylurea receptor. *Science* **270**, 1166-70 (1995).

8. Sakura, H., Ammala, C., Smith, P.A., Gribble, F.M. & Ashcroft, F.M. Cloning and functional expression of the cDNA encoding a novel ATP-sensitive potassium channel subunit expressed in pancreatic beta-cells, brain, heart and skeletal muscle. *FEBS Lett* **377**, 338-44 (1995).

9. Aguilar-Bryan, L., Nichols, C.G., Wechsler, S.W., Clement, J.P.t., Boyd, A.E., 3rd, Gonzalez, G., Herrera-Sosa, H., Nguy, K., Bryan, J. & Nelson, D.A. Cloning of the beta cell high-affinity sulfonylurea receptor: a regulator of insulin secretion. *Science* **268**, 423-6 (1995).

10. Inagaki, N., Gonoi, T. & Seino, S. Subunit stoichiometry of the pancreatic beta-cell ATP-sensitive K+ channel. *FEBS Lett* **409**, 232-6 (1997).

11. Tucker, S.J., Gribble, F.M., Zhao, C., Trapp, S. & Ashcroft, F.M. Truncation of Kir6.2 produces ATP-sensitive K+ channels in the absence of the sulphonylurea receptor. *Nature* **387**, 179-83 (1997).

12. de Wet, H., Mikhailov, M.V., Fotinou, C., Dreger, M., Craig, T.J., Venien-Bryan, C. & Ashcroft, F.M. Studies of the ATPase activity of the ABC protein SUR1. *FEBS J* **274**, 3532-44 (2007).

13. Matsuo, M., Kioka, N., Amachi, T. & Ueda, K. ATP binding properties of the nucleotide-binding folds of SUR1. *J Biol Chem* **274**, 37479-82 (1999).

14. Nichols, C.G., Shyng, S.L., Nestorowicz, A., Glaser, B., Clement, J.P.t., Gonzalez, G., Aguilar-Bryan, L., Permutt, M.A. & Bryan, J. Adenosine diphosphate as an intracellular regulator of insulin secretion. *Science* **272**, 1785-7 (1996).

15. Gribble, F.M., Tucker, S.J. & Ashcroft, F.M. The interaction of nucleotides with the tolbutamide block of cloned ATP-sensitive K+ channel currents expressed in Xenopus oocytes: a reinterpretation. *J Physiol* **504 (Pt 1)**, 35-45 (1997).

16. Gribble, F.M., Tucker, S.J. & Ashcroft, F.M. The essential role of the Walker A motifs of SUR1 in K-ATP channel activation by Mg-ADP and diazoxide. *EMBO J* **16**, 1145-52 (1997).

17. Ashcroft, F.M., Puljung, M.C. & Vedovato, N. Neonatal Diabetes and the KATP Channel: From Mutation to Therapy. *Trends Endocrinol Metab* **28**, 377-387 (2017).

18. Chatterjee, A., Guo, J., Lee, H.S. & Schultz, P.G. A genetically encoded fluorescent probe in mammalian cells. *J Am Chem Soc* **135**, 12540-3 (2013).

19. Ueda, K., Komine, J., Matsuo, M., Seino, S. & Amachi, T. Cooperative binding of ATP and MgADP in the sulfonylurea receptor is modulated by glibenclamide. *Proc Natl Acad Sci U S A* **96**, 1268-72 (1999).

20. Heuser, J. The production of 'cell cortices' for light and electron microscopy. *Traffic* **1**, 545-52 (2000).

21. Usukura, J., Yoshimura, A., Minakata, S., Youn, D., Ahn, J. & Cho, S.J. Use of the unroofing technique for atomic force microscopic imaging of the intra-cellular cytoskeleton under aqueous conditions. *J Electron Microsc (Tokyo)* **61**, 321-6 (2012).

22. Zagotta, W.N., Gordon, M.T., Senning, E.N., Munari, M.A. & Gordon, S.E. Measuring distances between TRPV1 and the plasma membrane using a noncanonical amino acid and transition metal ion FRET. *J Gen Physiol* **147**, 201-16 (2016).

23. Lopez-Alonso, J.P., de Araujo, E.D. & Kanelis, V. NMR and fluorescence studies of drug binding to the first nucleotide binding domain of SUR2A. *Biochemistry* **51**, 9211-22 (2012).

24. Oswald, C., Jenewein, S., Smits, S.H., Holland, I.B. & Schmitt, L. Water-mediated protein-fluorophore interactions modulate the affinity of an ABC-ATPase/TNP-ADP complex. *J Struct Biol* **162**, 85-93 (2008).

25. Vanoye, C.G., MacGregor, G.G., Dong, K., Tang, L., Buschmann, A.S., Hall, A.E., Lu, M., Giebisch, G. & Hebert, S.C. The carboxyl termini of K(ATP) channels bind nucleotides. *J Biol Chem* **277**, 23260-70 (2002).

26. Drain, P., Li, L. & Wang, J. KATP channel inhibition by ATP requires distinct functional domains of the cytoplasmic C terminus of the pore-forming subunit. *Proc Natl Acad Sci U S A* **95**, 13953-8 (1998).

27. Proks, P., de Wet, H. & Ashcroft, F.M. Activation of the K(ATP) channel by Mg-nucleotide interaction with SUR1. *J Gen Physiol* **136**, 389-405 (2010).

28. Corry, B., Jayatilaka, D. & Rigby, P. A flexible approach to the calculation of resonance energy transfer efficiency between multiple donors and acceptors in complex geometries. *Biophys J* **89**, 3822-36 (2005).

29. Proks, P., de Wet, H. & Ashcroft, F.M. Sulfonylureas suppress the stimulatory action of Mg-nucleotides on Kir6.2/SUR1 but not Kir6.2/SUR2A KATP channels: a mechanistic study. *J Gen Physiol* **144**, 469-86 (2014).

30. Gribble, F.M., Reimann, F., Ashfield, R. & Ashcroft, F.M. Nucleotide modulation of pinacidil stimulation of the cloned K(ATP) channel Kir6.2/SUR2A. *Mol Pharmacol* **57**, 1256-61 (2000).

31. Del Castillo, J. & Katz, B. Interaction at end-plate receptors between different choline derivatives. *Proc R Soc Lond B Biol Sci* **146**, 369-81 (1957).

32. de Wet, H., Shimomura, K., Aittoniemi, J., Ahmad, N., Lafond, M., Sansom, M.S. & Ashcroft, F.M. A universally conserved residue in the SUR1 subunit of the KATP channel is essential for translating nucleotide binding at SUR1 into channel opening. *J Physiol* **590**, 5025-36 (2012).

33. Magge, S.N., Shyng, S.L., MacMullen, C., Steinkrauss, L., Ganguly, A., Katz, L.E. & Stanley, C.A. Familial leucine-sensitive hypoglycemia of infancy due to a dominant mutation of the beta-cell sulfonylurea receptor. *J Clin Endocrinol Metab* **89**, 4450-6 (2004).

34. Sakura, H., Trapp, S., Liss, B. & Ashcroft, F.M. Altered functional properties of KATP channel conferred by a novel splice variant of SUR1. *J Physiol* **521 Pt 2**, 337-50 (1999).

35. Zerangue, N., Schwappach, B., Jan, Y.N. & Jan, L.Y. A new ER trafficking signal regulates the subunit stoichiometry of plasma membrane K(ATP) channels. *Neuron* **22**, 537-48 (1999).

36. Schmied, W.H., Elsasser, S.J., Uttamapinant, C. & Chin, J.W. Efficient multisite unnatural amino acid incorporation in mammalian cells via optimized pyrrolysyl tRNA synthetase/tRNA expression and engineered eRF1. *J Am Chem Soc* **136**, 15577-83 (2014).

37. Li, N., Wu, J.X., Ding, D., Cheng, J., Gao, N. & Chen, L. Structure of a Pancreatic ATP-Sensitive Potassium Channel. *Cell* **168**, 101-110 e10 (2017).

38. Martin, G.M., Yoshioka, C., Rex, E.A., Fay, J.F., Xie, Q., Whorton, M.R., Chen, J.Z. & Shyng, S.L. Cryo-EM structure of the ATP-sensitive potassium channel illuminates mechanisms of assembly and gating. *Elife* **6**(2017).

39. Monod, J., Wyman, J. & Changeux, J.P. On the Nature of Allosteric Transitions: A Plausible Model. *J Mol Biol* **12**, 88-118 (1965).

40. Babenko, A.P. A novel ABCC8 (SUR1)-dependent mechanism of metabolism-excitation uncoupling. *J Biol Chem* **283**, 8778-82 (2008).

41. Hosy, E. & Vivaudou, M. The unusual stoichiometry of ADP activation of the KATP channel. *Front Physiol* **5**, 11 (2014).

42. Tammaro, P., Girard, C., Molnes, J., Njolstad, P.R. & Ashcroft, F.M. Kir6.2 mutations causing neonatal diabetes provide new insights into Kir6.2-SUR1 interactions. *EMBO J* **24**, 2318-30 (2005).

43. Vedovato, N., Ashcroft, F.M. & Puljung, M.C. The Nucleotide-Binding Sites of SUR1: A Mechanistic Model. *Biophys J* **109**, 2452-60 (2015).

44. Biskup, C., Kusch, J., Schulz, E., Nache, V., Schwede, F., Lehmann, F., Hagen, V. & Benndorf, K. Relating ligand binding to activation gating in CNGA2 channels. *Nature* **446**, 440-3 (2007).

45. Zheng, J. & Zagotta, W.N. Gating rearrangements in cyclic nucleotide-gated channels revealed by patch-clamp fluorometry. *Neuron* **28**, 369-74 (2000).

46. Proks, P., Puljung, M.C., Vedovato, N., Sachse, G., Mulvaney, R. & Ashcroft, F.M. Running out of time: the decline of channel activity and nucleotide activation in adenosine triphosphate-sensitive K-channels. *Philos Trans R Soc Lond B Biol Sci* **371**(2016).

47. Matsuo, M., Tanabe, K., Kioka, N., Amachi, T. & Ueda, K. Different binding properties and affinities for ATP and ADP among sulfonylurea receptor subtypes, SUR1, SUR2A, and SUR2B. *J Biol Chem* **275**, 28757-63 (2000).

48. Zingman, L.V., Alekseev, A.E., Bienengraeber, M., Hodgson, D., Karger, A.B., Dzeja, P.P. & Terzic, A. Signaling in channel/enzyme multimers: ATPase transitions in SUR module gate ATP-sensitive K+ conductance. *Neuron* **31**, 233-45 (2001).

49. Zingman, L.V., Hodgson, D.M., Bienengraeber, M., Karger, A.B., Kathmann, E.C., Alekseev, A.E. & Terzic, A. Tandem function of nucleotide binding domains confers competence to sulfonylurea receptor in gating ATP-sensitive K+ channels. *J Biol Chem* **277**, 14206-10 (2002).

50. Choi, K.H., Tantama, M. & Licht, S. Testing for violations of microscopic reversibility in ATP-sensitive potassium channel gating. *J Phys Chem B* **112**, 10314-21 (2008).

51. Csanady, L., Vergani, P. & Gadsby, D.C. Strict coupling between CFTR's catalytic cycle and gating of its Cl- ion pore revealed by distributions of open channel burst durations. *Proc Natl Acad Sci U S A* **107**, 1241-6 (2010).

52. Ortiz, D., Gossack, L., Quast, U. & Bryan, J. Reinterpreting the action of ATP analogs on K(ATP) channels. *J Biol Chem* **288**, 18894-902 (2013).

53. Sikimic, J., McMillen, T.S., Bleile, C., Dastvan, F., Quast, U., Krippeit-Drews, P., Drews, G. & Bryan, J. ATP-binding without hydrolysis switches sulfonylurea receptor 1 (SUR1) to outward-facing conformations that activate KATP channels. *J Biol Chem* (2018).

54. Li, M., Silberberg, S.D. & Swartz, K.J. Subtype-specific control of P2X receptor channel signaling by ATP and Mg2+. *Proc Natl Acad Sci U S A* **110**, E3455-63 (2013).

55. Kawate, T. & Gouaux, E. Fluorescence-detection size-exclusion chromatography for precrystallization screening of integral membrane proteins. *Structure* **14**, 673-81 (2006).

56. Makhina, E.N. & Nichols, C.G. Independent trafficking of KATP channel subunits to the plasma membrane. *J Biol Chem* **273**, 3369-74 (1998).

57. Lin, C.Y., Huang, Z., Wen, W., Wu, A., Wang, C. & Niu, L. Enhancing Protein Expression in HEK-293 Cells by Lowering Culture Temperature. *PLoS One* **10**, e0123562 (2015).

58. Broglie, K.E. & Takahashi, M. Fluorescence studies of threonine-promoted conformational transitions in aspartokinase I using the substrate analogue 2'(3')-O-(2,4,6-trinitrophenyl)adenosine 5'-triphosphate. *J Biol Chem* **258**, 12940-6 (1983).

59. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.Y., White, D.J., Hartenstein, V., Eliceiri, K., Tomancak, P. & Cardona, A. Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-82 (2012).

60. Puljung, M.C. spectra-analysis. GitHub. https://github.com/mpuljung/spectra-analysis/blob/master/spegui.m. a8fe6af. (2019).

61. Puljung, M.C. spectra-analysis. GitHub. https://github.com/mpuljung/spectra-
analysis/blob/master/wash.m. ed14321. (2019).
62. Puljung, M.C. spectra-analysis. GitHub. https://github.com/mpuljung/spectra-
analysis/blob/master/gelnalyze.m. 9089db3. (2019).
63. Toyoshima, C., Yonekura, S., Tsueda, J. & Iwasawa, S. Trinitrophenyl derivatives bind
differently from parent adenine nucleotides to Ca2+-ATPase in the absence of Ca2+. *Proc Natl*
Acad Sci U S A **108**, 1833-8 (2011).

construct	nucleotide	EC50 (μM)	h	n
SUR1/Kir6.2-G334D	MgTNP-ADP	20.6 ± 0.4	1.1 ± 0.2	12
SUR1-T1397*/Kir6.2-G334D	MgTNP-ADP	7.3 ± 1.2	1.0 ± 0.1	5
	MgADP	44.4 ± 14.5	1.5 ± 0.3	5

Table 1. Mean $\pm$ SEM from fits of equation 5 to individual electrophysiological experiments.

construct	nucleotide	E_{max}	EC_{50} (µM)	h	n
SUR1-Y1353*/Kir6.2	MgTNP-ADP	0.99 ± 0.01	5.5 ± 0.9	0.77 ± 0.03	8
	MgTNP-ATP	0.99 ± 0.02	37.5 ± 11.9	0.86 ± 0.07	7
	TNP-ADP	0.95 ± 0.01	6.8 ± 1.3	0.79 ± 0.04	7
	TNP-ATP	0.83 ± 0.03	12.8 ± 5.6	0.88 ± 0.06	7
SUR1-Y1353*,K2A/Kir6.2	MgTNP-ADP	0.97 ± 0.03	18.6 ± 8.6	0.97 ± 0.07	6
	MgTNP-ATP	0.99 ± 0.02	31.4 ± 9.1	1.1 ± 0.3	6
	TNP-ADP	0.92 ± 0.02	5.0 ± 1.0	0.88 ± 0.05	6
	TNP-ATP	0.90 ± 0.03	8.1 ± 3.8	0.98 ± 0.13	6
SUR1-T1397*/Kir6.2	MgTNP-ADP	1.01 ± 0.005	10.8 ± 0.7	0.98 ± 0.05	6
	MgTNP-ATP	1.00 ± 0.01	17.1 ± 3.9	1.0 ± 0.1	10
	TNP-ADP	0.98 ± 0.02	6.3 ± 2.3	0.84 ± 0.13	9
	TNP-ATP	0.94 ± 0.01	4.7 ± 0.8	0.83 ± 0.09	8
SUR1-Y1353*/Kir6.2-G334D	MgTNP-ADP	1.01 ± 0.04	8.2 ± 1.9	0.90 ± 0.15	5

Table 2. Mean ± SEM from fits of equation 1 to individual nucleotide binding experiments.

SUR1 construct	nucleotide	drug	$T_{0.5}$ (s)	n
Y1353*	TNP-ADP		2.9 ± 0.5	7
	MgTNP-ADP		14.7 ± 4.2	7
	MgTNP-ADP	tolbutamide	4.5 ± 1.5	7
	MgTNP-ADP	diazoxide	36.9 ± 9.4	7
	TNP-ATP		4.8 ± 1.5	7
	MgTNP-ATP		10.5 ± 1.9	7
	MgTNP-ATP	tolbutamide	4.4 ± 1.6	7
	MgTNP-ATP	diazoxide	42.0 ± 9.7	6
Y1353*,K2A	TNP-ADP		3.8 ± 1.0	3
	MgTNP-ADP		9.3±2.1	4

Table 3. Mean ± SD from exponential fits to individual experiments measuring nucleotide wash-out time courses.

construct	ANAP	pANAP	peRF1-E55D	nucleotide	I/I$_{control}$	n
3xFLAG_SUR1/Kir6.2	-	-	-	100 μM MgATP	3.6 ± 1.3	13
SUR1-Y1353^{stop}/Kir6.2	+	-	-	100 μM MgATP	0.41 ± 0.7	7
SUR1-Y1353^{stop}/Kir6.2	+	+	-	100 μM MgATP	0.42 ± 0.05	5
SUR1-Y1353^{stop}/Kir6.2	+	+	+	100 μM MgATP	0.37 ± 0.10	3

Table 4. Mean ± SD response to 100 μM MgADP relative to the current in nucleotide-free solution.

construct	nucleotide	IC50 (µM)	h	n
SUR1/Kir6.2	ATP	11.2 ± 0.1	-1.3 ± 0.1	6
SUR1-T1397*/Kir6.2	ATP	53.9 ± 9.4	-1.2 ± 0.1	6

Table 5. Mean ± SEM from fits of equation 4 to individual electrophysiological experiments.

Figure 1

Figure 1--figure supplement 1

Figure 1--figure supplement 2

Figure 1--figure supplement 3

Figure 1--figure supplement 4

Figure 1--figure supplement 5

Figure 2

Figure 3

Figure 3--figure supplement 1

Figure 4

Figure 4--figure supplement 1

Figure 4--figure supplement 2

Figure 5

Figure 5--figure supplement 1

Figure 5--figure supplement 2

Figure 6