Title: CARDIOMETABOLIC RECOVERY AND LACTATE REMOVAL MAY BE RELATED TO
MUSCULAR ADAPTATIONS

Short Title: CARDIOMETABOLIC RECOVERY AND MUSCULAR ADAPTATIONS

Edgar Ismael Alarcón Meza. Researcher at Universidad Autónoma de Baja California. Email:
eiam@uabc.edu.mx. Baja California, Mexico.

Glauber Lameira de Oliveira. Researcher of LABIMH-UFRJ, Federal University of Rio de
Janeiro. Email: lameiraglauber@gmail.com. Rio de Janeiro, Brazil

Talita Adão Perini de Oliveira. Researcher of LABIMH-UFRJ, Federal University of Rio de
Janeiro. Email: talitap.perini@gmil.com. Rio de Janeiro, Brazil

Marcelo Hubner Moreira. Professor at the North Educational Union (UNINORTE-AC College).
Email: professorhubner@gmail.com. Acre, Brazil.

Angeliete Garcez Militão. Adjunct Professor of Federal University of Rondônia, RO, Brazil.
Email: angeliete@unir.br. Rondônia, Brazil.

José Fernandes Filho. Associated Professor at Federal University of Rio de Janeiro, RJ. E-mail:
jff@eefd.ufrj.br. Rio de Janeiro, Brazil

Vernon Furtado Da Silva. Senior Visiting Professor at Federal University of Rondônia – UNIR.
Rondônia, Brazil.

Estélio Henrique Martin Dantas. Professor at the Tiradentes University, Aracajú, Sergipe, and
in the Nursing Pós-Graduate Doctoral Program at UNIRIO. E-mail: estélio.dantas@unirio.br.
Sergipe, Brazil.

João Rafael Valentim-Silva*, Professor at the North Educational Union (UNINORTE-AC
College) and Federal University of Rondônia. p.jrvalentim@gmail.com; Phone Number: (69) 9-
9211-3015 or 9-8132-9631. Acre, Brazil. **Contact Author.**

All authors contributed equally to this work.

Correspondent author: p.jrvalentim@gmail.com (JRVS)

Abstract

The aim of this study was to measure the differences in cardiorespiratory recovery (CR) and blood lactate removal among young athletes with differences in non-lactic (NP) and lactic anaerobic power (LP) and fatigue index (FI) but with the same degree of cardiorespiratory fitness. Sixteen swimmers from the Brazilian synchronized swimming team (2014) were divided into two groups GAP (Group High Power) (n = 9) , with NP, LP, and FI (p<0.05) compared to GBP (Group Low Power) (n=7). Both groups performed a four-minute routine at competitive intensity. Anaerobic power, maximal heart rate (HR) and blood lactate (BL) were determined before and at 1, 3 and 5 minutes after the routine. Student's t-test was used to analyze the intergroup differences of NP, LP, FI, maximum and lactic HR, and two-way ANOVA followed by Bonferroni was used to analyze HR and BL at 1, 3 and 5 minutes after activity with a significance of 5%. The FI of the GBP group was lower than that of the GAP group (P <0.05). The NP of the GBP group was higher than that of the GAP group (P <0.05). The maximum HR of the GBP group was equal to that of the GAP group (P> 0.05). The GBP group had better HR recovery than did the GAP group (P <0.05). BL had its lowest levels after 1 and 5 minutes of recovery in the GBP group when compared to the GAP group (P <0.05). The GBP group's FI was significantly lower than that of the GAP group, while NP was higher, and CR was better in the GBP group, indicating a relationship between a lower FI and higher NP and LP with CR and suggesting that muscular adaptations have an important influence on CR and BL removal.

Keywords: Post-exercise Blood Lactate Removal. Heart Rate Recovery. Non-lactic Anaerobic Power. Lactic Anaerobic Power.

Introduction

The literature is consistent in affirming that various psychological, biological and performance variables are indicators of fatigue. Investigations have revealed that creatine kinase, lactate dehydrogenase, blood lactate, oxygen consumption and heart rate are associated with exercise intensity and fatigue. Furthermore, the time required to return to pre-exercise levels is considered an indicator of the recovery capacity that is linked to an athlete's physical fitness [1–4].

Heart rate variability (HRV) analysis is an established method used to quantify the extent of autonomic recovery from exercise [5,6]. After completing a given exercise, a rapid decrease in parasympathetic cardiac activity to resting levels suggests a relative and physiological systemic recovery imposed by the workload [7,8] and the amount of time required for parasympathetic reactivation after exercise may be significantly influenced by several factors, including exercise intensity[9] and cardiorespiratory fitness [10,11]. Although many of the factors inherent to HRV have been frequently studied, studies specifically concerning the non-lactic and lactic anaerobic capacities and fatigue index have not yet been conducted.

Another important marker of metabolic activation provided by intense exercise is lactate. Many authors have shown that when the rate of ATP production by oxidative sources becomes insufficient, high rates of glycolytic or glycogenolytic ATP production are required, culminating in the production of pyruvate, a metabolite that can be reduced to lactate or oxidized to CO_2 or H_2O [12–15].

Thus, by increasing the intensity of exercise and muscle load, several tissues begin to produce more lactate and export it to the circulation. Simultaneously, the less active skeletal muscles, the heart, the liver, the renal cortex and the brain remove lactate from circulation, suggesting

that this metabolite acts as an intermediary for transporting carbohydrates from cells and

tissues with relatively low oxidative capacity to cells and tissues with high oxidative capacity

[16,17]. Therefore, it is well established that blood lactate concentration is the result of

production and removal of this metabolite.

Lactate has been a focus of research on skeletal muscle metabolism and a central theme of

the controversy over the relationships between lactate production, acidosis and fatigue,

because it is probably a residual product; lactate is increasingly regarded as a source of fuel,

continuously produced and metabolized in the organism in order to maintain energy

homeostasis at adequate levels in an attempt to maintain cellular activity [12,16,18].

A fact that somewhat contradicts what has been described in the literature up to the present

moment is that high-intensity exercise can produce greater expression and translocation of

MCT-4 (monocarboxylate transporter) [19,20]. This adaptation may have implications for the

export and import of lactate, because the plasma membrane is responsible for this transport;

furthermore, high-intensity exercise, in addition to promoting adaptation related to lactic and

non-lactic anaerobic resistance, is the stimulus for MCT-4 translocation and expression in

muscles [21,22].

These statements support the notion that muscular fitness levels may be related to reduced

blood lactate levels, while cardiorespiratory fitness is seen as being analogous to the velocity

of decrease in heart rate, which is the hypothesis postulated in the present investigation. Thus,

the objective of the present study is based on the possibility that athletes with various levels

of non-lactic and lactic power and fatigue indexes present various behaviors in terms of lactate

removal and heart rate recovery curves.

108 **Materials and Methods**

109 **Volunteer Group**

All 16 swimmers (17 ± 1.4 years old) from the Brazilian synchronized swimming team (2014)

participated in the study at the beginning of the training season (January). The athletes were

divided into two groups – GAP (n= 9) and GBP (n= 7) – based on their non-lactic and lactic

anaerobic capacity. The athletes were instructed not to train the day before the test, to have

their last meal up to three hours ahead of time, to avoid caffeine intake, to avoid medication,

and to sleep at least 6 hours the night before. A brief history was taken to identify any possible

exclusion factors. The volunteer characteristics are summarized in the Table 1.

Table 1: physical data and training experience of the volunteers

Variables	GAP (n=9)	GBP (n=7)
Age (years)	18.8±1.4	15.8± 1.6
Height (cm)	166.31±4.3	162.0±3.1
Body Mass	55.16±4.2	55.28±5.2
Body Mass Index	20.01±3.3	21.06±3.7
Training Time	9.5±1.6	7.9±1.7

118   Values are expressed as the mean ± SD.
119

120 **Procedures**

121 **Exercise Protocol**

The exercises were composed of a competitive routine with a duration of 4 minutes at

maximum competitive intensity. Prior to the tests, heart rates and lactate levels were

measured. Then, the competitive routine was performed with heart rate monitoring in order

to determine maximum heart rate. Finally, after 1, 3 and 5 minutes of recovery, heart rate and

blood samples were collected for later determination of blood lactate levels as summarized in

figure 1.

Figure 1: exercise protocol and data collection.

Functional Capacity

The stress test was performed on a treadmill (ECAFIX), using Bruce's protocol. The Borg scale

was adopted for a subjective measure of effort. The test was interrupted in the presence of

symptoms that impeded its continuity (fatigue) and/or represented a risk to the evaluated

individual. The exercise period was preceded by a period of adaptation and preparation of the

athlete to the equipment, followed by five minutes of monitored active recovery (five km/h).

Gas exchange was measured with a pneumotachograph (Pt, MEDGRAFIC, Wilmington, DE,

USA) coupled to a differential pressure transducer, near the capillary of the gas analyzer in

order to collect a sample of inhaled and exhaled gases. The athlete, whose nose was sealed

by a clip, was connected to the system for acquisition of ventilatory data through a

mouthpiece. Flow signals and gaseous concentrations were sampled (Pentium III computer)

at a rate of 1000 Hz. Cycle-to-cycle gas concentrations were sampled by the cardiorespiratory

diagnostic system MEDGRAFIC, model VO2000. Respiratory flow and electrocardiographic

signals (ECG - ECAFIX) were processed on a personal computer in real time.

The mean values of the maximum load reached during exercise of the last three respiratory

cycles were computed in order to determine the results of the variables representative of the

end of the exercise (peak). The following ventilatory variables were analyzed: ventilation

minute (V_E, l.min^{-1} – BTPS), oxygen consumption (VO_{2peak}, l.min^{-1} – STPD), carbon dioxide

production (VCO_{2peak}, l.min^{-1} – STPD), gas exchange ratio (R=VCO_{2peak}/VO_{2peak}) and ventilatory

anaerobic threshold (LA, 1.min^{-1} – STPD).

Identification of LA was done through a non-invasive method from the analysis of the

ventilatory equivalent of oxygen (V_E/VO_2) and carbon dioxide (V_E/VCO_2), as previously

described [23,24].

The gas analyzer was calibrated daily before beginning the tests with a calibration bullet (AGA

– primary standard) with the following concentrations: 12.1% O_2, 5.0% CO_2 and 83.0% N_2.

Anaerobic Power

Non-lactic and lactic anaerobic power and the fatigue rate were determined by the Wingate

test [25] performed on a mechanical cycle ergometer (MONARK). This test preceded the

ergospirometric test with an interval of at least 60 minutes between them. To compute the

total revolution of the pedal and to calculate the power every five seconds, an optical sensor

was coupled to the cycle ergometer managed by an acquisition program elaborated in

Labview 6.0 (National Instruments, USA).

The load was determined by 0.075 $Kp.kg^{-1}$ of body mass as predicted in the Wingate test

protocol. The test was preceded by a two-minute warm-up period and followed by an active

three-minute recovery period with a 50-watt load. From the individual results, the following

parameters were calculated: non-lactic anaerobic power by means of the absolute (or

maximum) power peak (watts) and relative to body mass ($watts.kg^1$) and the lactic power or

average (watts). The percentage of power decline (%) or rate of fatigue was obtained from the

difference between the highest power and the lowest power achieved by the athlete during

the 30 s test. The bicycle load was checked before starting the test.

Parameter calculations, descriptive statistics of the data and comparisons were performed

using one-way ANOVA and the Bonferroni post-tests in the program "PrismStat 5.0" with a

significance level of $p < 0.05$.

Ethical aspects

The evaluations were carried out in the Ergo-Spirometry Department of the Laboratory of

Exercise Physiology of the School of Physical Education and Sports at the Federal University of

Rio de Janeiro (SE-LABOFISE-EEFD-UFRJ).

All participants were aware and signed a written consent form, including the procedures

adopted and the authorization of the volunteers to investigate the results found in the

scientific study. The anonymity and privacy of the participants were preserved in the study.

The Council of Ethics and Human Research of the Federal Institute of Education of Rondônia

approved this investigation under CAAE number 44907715.2.00005653. All subjects gave their

consent to participate.

Results

Heart rate recovered faster in the GBP group than in the GAP group ($P<0.05$) (Fig. 2A); a similar

phenomenon was observed regarding the behavior of lactate ($P <0.05$) (Fig. 2B). In relation to

physical fitness and respiratory capacities, there were no differences between the groups ($P >$

0.05) (Fig. 2C and 2D); however, the fatigue index, lactic and non-lactic anaerobic power

showed substantial differences between the groups ($P <0.0001$) (Fig. 2E).

Figure 2: heart rate and lactate recovery curve. The two groups, GBP (n=9) and GAP (n=7),

were subjected to a 4-minute routine on different days. **(A)** Comparison of heart rates

between the GBP and GAP groups before, during and after the 4-minute routine. **(B)**

Comparison of lactate between the GBP and GAP groups before, during and after the 4-minute

routine. **(C)** Comparison of the VO_2, VO_2/kg peak, and the V_E peak between the GBP and GAP

groups. **(D)** Comparison of the R Peak, VO_2 and VO_2/kg peak between the GBP and GAP groups.

(E). Comparison of the Fatigue Index, Non-lactic Anaerobic Power and Lactic Anaerobic Power

between the GBP and GAP groups (**A, B, C** and **D** ns = p> 0.05, *p <0.05). (**G** * = p <0.0001 GBP

vs GAP).

Discussion

The present study is the first in the literature to compare the cardiometabolic responses of

elite athletes of synchronized swimming in terms of non-lactic anaerobic, lactic anaerobic

power and fatigue index. To determine the physical characteristics of both groups, some

measures of body composition, functional capacity and anaerobic power were conducted.

Recently, it was shown that lactate removal was partly dependent on the time of recovery

between stimulus sessions, and moderate correlations were observed between post-exercise

blood lactate concentration, peak heart rate and perceived exertion [26]. Nevertheless, the

possible physiological mechanisms underlying these observations were not described. Here,

we propose that the muscle adaptations regarding the fatigue index can be related to lactate

removal and heart hate recovery.

It is well established in the literature that lower fatigue index and higher non-lactic and lactic

anaerobic power are associated with higher level of exercise-related muscle adaptations.

It has been proposed that aerobic training may improve the ability of muscle to recover after

anaerobic exercise, suggesting that an athlete with greater aerobic fitness will use fewer non-

oxidative sources and thus will recover more quickly from exercise. Theoretically, an increase

in aerobic fitness improves the recovery from anaerobic exercise However, this phenomenon

was not completely reproduced in the present study because the aerobic capacity of both

groups was equal while the cardiometabolic recovery was different. This is reasonable

because lactate removal could be linked to uptake by muscle and other tissues that use this

substrate as an energy source [12,27].

A synchronized swimming routine requires a high aerobic effort with fundamental anaerobic

contributions to maintain adequate exercise levels during competition [28,29]. The data

shown here demonstrate that the peaks of non-lactic and lactic anaerobic power have the

same behavior, but the fatigue index was the only factor directly related to physical fitness

that showed differences between the groups.

In the present study, the GAP group had heart rate and lactate curve response behaviors

different from those observed in the GBP group, although the maximum heart rate was equal

in both groups. The GBP group showed a lower lactate peak than did the GAP group. Another

interesting fact is that the non-lactic anaerobic power, lactic anaerobic power and the fatigue

index differed between the two groups, even though VO_2 and VCO_2 were equal. These data

contradict a part of the literature that states that aerobic capacity improves recovery after

intense exercise, a finding that was not seen in the present investigation.

Other investigations have pointed to several possible explanations for the effect of aerobic

conditioning on recovery. Holloszy [30] pointed to the increase in the number and size of

mitochondria in metabolic recovery. However, the increased myoglobin content and blood

volume together would be implicated in the transport of O_2 to the muscle during high-intensity

exercise, and consequently, there would be a decrease in recovery time, since this would also

help to remove the accumulated lactate more quickly [31,32].

These data explain the higher velocity in cardiometabolic recovery, However, the data

described here demonstrate a clear difference in lactate removal between the groups though

both exhibit equal V_E peak, $V_{O2\ Peak}$, $V_{O2\ Peak/body\ mass}$ and peak heart rate, suggesting that the

aerobic capacity of these athletes is unrelated to metabolic recovery after maximal intensity

exercise.

This difference may be related to the use of lactate by other muscles and tissues, since

anaerobic metabolism after a few minutes of intense exercise may activate lactate uptake and

use it as an important energy source during intense exercise by modifying the kinetics of the

use of glucose in the presence of lactate[15]. However, unfortunately, almost all studies

investigating differences in lactate responses depend on blood lactate measurements that

reflect muscle lactate alone, providing indirect evidence of lactate accumulation and removal.

Other authors did not find evidence that associated the aerobic capacity with greater speed

of lactate removal, corroborating the findings of the present study [32–34].

These data, taken together, suggest that the adaptations that led to differences in lactate

responses, but not in heart rate, were linked to the muscular system in greater magnitude

than to the cardiorespiratory system, since the ventilatory capacities between the groups (V_E

$_{Peak}$, $V_{O2\ Peak}$, $V_{O2\ Peak/body\ mass}$, $V_{CO2\ Peak}$, $V_{CO2\ Peak/body\ mass}$) were the same. Therefore, it appears that

the cardiorespiratory adaptations of both groups reached the same levels, suggesting that the

observed difference was related to adaptations of the muscles required during the exercise

routine, although explaining the mechanism that is linked to what was observed here was not

the focus of the present investigation.

A study conducted by Otsuki et al. [35] demonstrated differences in cardiac recovery rates

between athletes trained in endurance and strength. This study may have its results

extrapolated to the data found here, because the non-lactic and lactic anaerobic power and

the fatigue index between the two groups showed differences, a fact that is usually evident

among people who are trained or not in exercises and strength. However, studies that

demonstrate differences in heart rate between groups that have differences in non-lactic and

lactic anaerobic power and the fatigue index are scarce, limiting the discussion of the present

study.

Among the limitations found in the present study is the need for a population with greater

diversity than the one tested here and in previous studies; for example, diversity in terms of

age, BMI and health status is the next logical step in order to refine the interpretation of the

data described here. There is also a scarcity of data regarding metabolic and cardiac recovery

in elite athletes with the same level of aerobic conditioning for comparison between our

findings and those of other authors.

Conclusions

Lactate removal and HR characteristics showed differences between the groups, with the best

performance and the group with the lowest fatigue index, although cardiorespiratory fitness,

demonstrated by the maximum HR reached, and the measure of pulmonary capacity and O_2

uptake were equal between the two groups. These data suggest that muscular adaptations

may have a critical role in the behavior of cardiometabolic recovery, in contrast to the findings

in the literature that focused on muscle metabolism in exercise.

Practical Applications

This paper challenges the idea that only the cardiovascular training is related to recovery after

an exercise training session. Therefore, an intense neuromuscular training could help in

recovery, especially in sports in which there is more than one competition on the same day.

This may be applicable to athletes in sports with fundamentally aerobic components. This

knowledge could be crucial for improving performance during the several trials during the

competition day.

Acknowledgments

The authors wish to thank the athletes who participated in the study for their effort and dedication for

their help and availability.

References

1.   Rider BC, Coughlin AM, Hew-Butler TD, Goslin BR. Effect of compression stockings on

physiological responses and running performance in division III collegiate cross-

country runners during a maximal treadmill test. *J Strength Cond Res*.

2014;28(6):1732-1738. doi:10.1519/JSC.0000000000000287.

2.   Elias GP, Wyckelsma VL, Varley MC, McKenna MJ, Aughey RJ. Effectiveness of water

immersion on postmatch recovery in elite professional footballers. *Int J Sports Physiol*

Perform. 2013;8(3):243-253. doi:10.1123/ijspp.8.3.243.

3.   Higgins TR, Climstein M, Cameron M. Evaluation of hydrotherapy, using passive tests

and power tests, for recovery across a cyclic week of competitive rugby union. *J*

Strength Cond Res. 2013;27(4):954-965. doi:10.1519/JSC.0b013e318260ed9b.

4.   Higgins T, Cameron M, Climstein M. Evaluation of passive recovery, cold water

immersion, and contrast baths for recovery, as measured by game performances

markers, between two simulated games of rugby union. *J Strength Cond Res*. 2012:1.

doi:10.1519/JSC.0b013e31825c32b9.

5.   Michael S, Graham KS, Davis GM. Cardiac Autonomic Responses during Exercise and

Post-exercise Recovery Using Heart Rate Variability and Systolic Time Intervals—A

Review. *Front Physiol*. 2017;8. doi:10.3389/fphys.2017.00301.

6.   Molina GE, Fontana KE, Porto LGG, Junqueira LF. Post-exercise heart-rate recovery

correlates to resting heart-rate variability in healthy men. *Clin Auton Res*.

2016;26(6):415-421. doi:10.1007/s10286-016-0378-2.

7.   Cinaz B, Arnrich B, La Marca R, Tröster G. Monitoring of mental workload levels during

an everyday life office-work scenario. *Pers Ubiquitous Comput*. 2013;17(2):229-239.

doi:10.1007/s00779-011-0466-1.

8.   Buchheit M, Chivot A, Parouty J, et al. Monitoring endurance running performance

using cardiac parasympathetic function. *Eur J Appl Physiol*. 2010;108(6):1153-1167.

doi:10.1007/s00421-009-1317-x.

9.   Stuckey MI, Tordi N, Mourot L, et al. Autonomic recovery following sprint interval

exercise. *Scand J Med Sci Sports*. 2012;22(6):756-763. doi:10.1111/j.1600-

0838.2011.01320.x.

10.   Sandercock G, Hurtado V, Cardoso F. Changes in cardiorespiratory fitness in cardiac

rehabilitation patients: A meta-analysis. *Int J Cardiol*. 2013;167(3):894-902.

doi:10.1016/j.ijcard.2011.11.068.

11.   Da Silva DF, Bianchini JAA, Antonini VDS, et al. Parasympathetic cardiac activity is

associated with cardiorespiratory fitness in overweight and obese adolescents.

Pediatr Cardiol. 2014;35(4):684-690. doi:10.1007/s00246-013-0838-6.

12.   Brooks GA. Cell-cell and intracellular lactate shuttles. *J Physiol*. 2009;587(23):5591-

5600. doi:10.1113/jphysiol.2009.178350.

13.   Brooks GA. Lactate shuttles in nature. *Biochem Soc Trans*. 2002;30(2):258-264.

doi:10.1042/.

14.   Brooks GA. Current concepts in lactate exchange. *Med Sci Sports Exerc*.

1991;23(8):895-906. doi:10.1249/00005768-199108000-00003.

15.   Brooks GA. Lactate production under fully aerobic conditions: the lactate shuttle

during rest and exercise. *Fed Proc*. 1986;45(13):2924-2929.

http://www.ncbi.nlm.nih.gov/pubmed/3536591.

16.   van Hall G, Lundby C, Araoz M, Calbet JAL, Sander M, Saltin B. The lactate paradox

revisited in lowlanders during acclimatization to 4100 m and in high-altitude natives. *J*

Physiol. 2009;587(5):1117-1129. doi:10.1113/jphysiol.2008.160846.

17.   van Hall G, Strømstad M, Rasmussen P, et al. Blood lactate is an important energy

source for the human brain. *J Cereb Blood Flow Metab*. 2009;29(6):1121-1129.

doi:10.1038/jcbfm.2009.35.

18.   Faude O, Kindermann W, Meyer T. Lactate threshold concepts: How valid are they?

Sport Med. 2009;39(6):469-490. doi:10.2165/00007256-200939060-00003.

19.   Mogi A, Koga K, Aoki M, et al. Expression and role of GLUT-1, MCT-1, and MCT-4 in

malignant pleural mesothelioma. *Virchows Arch*. 2013;462(1):83-93.

doi:10.1007/s00428-012-1344-6.

20.   Luo F, Zou Z, Liu X, et al. Enhanced glycolysis, regulated by HIF-1α via MCT-4,

promotes inflammation in arsenite-induced carcinogenesis. *Carcinogenesis*.

2017;38(6):615-626. doi:10.1093/carcin/bgx034.

21.   Ideno M, Kobayashi M, Sasaki S, et al. Involvement of monocarboxylate transporter 1

(SLC16A1) in the uptake of l -lactate in human astrocytes. *Life Sci*. 2017.

doi:10.1016/j.lfs.2017.10.022.

22.   Takimoto T, Tsue H, Takahashi H, Tamura R, Sasaki H. Synthesis of p-tert-

butylcalix[4]thiacrowns exhibiting sulfur number-dependent complexation with

mercury(II) ion. *Heterocycles*. 2014;88(2):911-917. doi:10.3987/COM-13-S(S)61.

23.   Beaver WL, Wasserman K, Whipp BJ. A new method for detecting anaerobic threshold

by gas exchange. *J Appl Physiol*. 1986;60(6):2020-2027.

http://www.ncbi.nlm.nih.gov/pubmed/3087938.

24.   Dickstein K, Barvik S, Aarsland T, Snapinn S, Karlsson J. A comparison of

methodologies in detection of the anaerobic threshold. *Circulation*. 1989;81(1

Suppl):II38-II46.

http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=2295151&

retmode=ref&cmd=prlinks%5Cnpapers2://publication/uuid/C7D07161-D4E6-4953-

A5AD-12B432B33AD8.

25.   Bar-Or O. The Wingate Anaerobic Test An Update on Methodology, Reliability and

Validity. *Sport Med An Int J Appl Med Sci Sport Exerc*. 1987;4(6):381-394.

doi:10.2165/00007256-198704060-00001.

26.   Lessard SJ, Rivas DA, Alves-Wagner AB, et al. Resistance to aerobic exercise training

causes metabolic dysfunction and reveals novel exercise-regulated signaling networks.

Diabetes. 2013;62(8):2717-2727. doi:10.2337/db13-0062.

27.   Rotstein A, Dotan R, Bar-Or O, Tenenbaum G. Effect of training on anaerobic

threshold, maximal aerobic power and anaerobic performance of preadolescent boys.

Int J Sports Med. 1986;7(5):281-286. doi:10.1055/s-2008-1025775.

28.   Yamamura C, Zushi S, Takata K, Ishiko T, Matsui N, Kitagawa K. Physiological

characteristics of well-trained synchronized swimmers in relation to performance

scores. *Int J Sports Med*. 1999;20(4):246-251. doi:10.1055/s-2007-971125.

29.   Dos-Santos JW, de Mello MA. Responses of Blood Lactate Concentration in Aerobic

and Anaerobic Training Protocols at Different Swimming Exercise Intensities in Rats. *J*

Exerc Physiol Online. 2011;14(3):34-42. http://proxy.lib.ohio-

state.edu/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=s3h&AN

=65237273&site=ehost-live.

30.   Holloszy JO, Coyle EF. Adaptations of skeletal muscle to endurance exercise and their

metabolic consequences. *J Appl Physiol*. 1984;56(4):831-838.

31.   Kanda K, Sugama K, Hayashida H, et al. Eccentric exercise-induced delayed-onset

muscle soreness and changes in markers of muscle damage and inflammation. *Exerc*

Immunol Rev. 2013;19:72-85.

32.   Tomlin DL, Wenger H a. The relationship between aerobic fitness and recovery from

high intensity intermittent exercise. *Sport Med*. 2001;31(1):1-11.

doi:10.2165/00007256-200131010-00001.

33.   Freund H, Lonsdorfer J, Oyono-Enguéllé S, Lonsdorfer A, Bogui P. Lactate exchange

and removal abilities in sickle cell patients and in untrained and trained healthy

humans. *J Appl Physiol*. 1992;73(6):2580-2587.

http://www.ncbi.nlm.nih.gov/pubmed/1490972.

34.   Bassett DR, Merrill PW, Nagle FJ, Agre JC, Sampedro R. Rate of decline in blood lactate

after cycling exercise in endurance-trained and -untrained subjects. *J Appl Physiol*.

1991;70(4):1816-1820. http://www.ncbi.nlm.nih.gov/pubmed/2055859.

35.   Otsuki T, Maeda S, Iemitsu M, et al. Postexercise heart rate recovery accelerates in

strength-trained athletes. *Med Sci Sports Exerc*. 2007;39(2):365-370.

doi:10.1249/01.mss.0000241647.13220.4c.

THE CARDIOMETABOLIC RECOVERY MAY BE RELATED TO MUSCULAR ADAPTATIONS

Figure 1: exercise protocol and data collection.

CARDIOMETABOLIC RECOVERY AND LACTATE REMOVAL MAY BE RELATIONSHIP TO MUSCULAR ADAPTATIONS

Figure 2: heart rate and lactate recovery curve. The two groups, GBP (n=9) and GAP (n=7), were submitted to a 4-minute routine on different days. **(A)** Comparison of heart rate between GBP and GAP before, during and after the 4-minute routine. **(B)** Comparison of lactate between GBP and GAP before, during and after the 4-minute routine. **(C)** Comparison of the Vo2, Vo2/kg peak, and the VE peak between GBP and GAP **(D)** Comparison of the R Peak, Vo2 and Vo2/Kg peak between GBP and GAP **(E)**. Comparison of the Fatigue Index, Non-lactic Anaerobic Power and Lactic Anaerobic Power between GBP and GAP (**A, B, C** and **D** ns = p> 0.05, *p <0.05). (**G** * = p <0.0001 GBP vs GAP).