Authors: Anderson Matos Medina

Title: Following the dung or the climate? Species richness pattern of dung beetles

(Scarabaeinae) on the Americas

Authors affiliation: Programa de Pós☐Graduação em Ecologia e Evolução, Instituto de

Ciências Biológicas, Universidade Federal de Goiás, Goiânia, Goiás, Brazil

E-mail address: bojaum@gmail.com

Abstract: Climatic conditions are a main driver of species richness. Specifically,

increase on climatic instability may reduce species richness directly and indirectly by

reducing resources available. This hypothesis is evaluated here using a producer-

consumer interaction to explain dung beetle richness on a continental scale (America)

using mammal richness as resource proxy and temperature and precipitation seasonality

as proxy for climatic instability. A spatial path analysis was built in order to evaluate

this hypothesis while controlling for spatial autocorrelation and differences on the

sampling effort and abundance of each study (n=115) gathered from the literature. Dung

beetle richness was directly explained by temperature seasonality, precipitation

seasonality and mammal richness, whereas only precipitation seasonality had an effect

modulated by mammal richness. This result reinforces the notion that species richness

can be explained by climatic conditions, but also reveals the importance of biotic

interactions in order to understand the mechanisms behind such patterns.

Keywords: climatic instability, resource diversity, dung beetle diversity, mammal

richness, Scarabaeidae

Acknowledgements

I received a scholarship from Coordenação de Aperfeiçoamento de Pessoal de Nível

Superior (CAPES). I am also grateful for the effort of all authors that sampled dung

beetle   communities   used   as   source   data   for   this   study.

Introduction

Latitudinal diversity gradients are a widespread pattern in ecology (Willig et al. 2003; Hillebrand 2004). Since latitude is not an explanation *per se* (Hawkins and Diniz-Filho 2004), many causal explanations, such as productivity, environmental heterogeneity, area, historical factors and biotic interactions were proposed for this geographic pattern (Willig et al. 2003; Field et al. 2009). Among these, climate and productivity are between the most used predictors of latitudinal diversity gradients (Field et al. 2009).

On the other hand, variation on temperature and rainfall may play a role in generating resource bottlenecks that can limit the number of species coexisting (Williams and Middleton 2008). In fact, the climatic stability hypothesis predicts that sites with higher stability possesses higher richness than sites with lesser stability (Pianka 1966) because species can be more specialized on stable environments (Moles and Ollerton 2016). Another factor potentially playing a role in structuring latitudinal richness pattern is resource availability. For example, the number of species of woody plants has been used to explain the richness of arboreal or frugivorous mammals, and figs richness has been demonstrated to determine frugivorous birds richness (Andrews and O'Brien 2000, Kissling et al. 2007; but see Hawkins and Pausas 2004). The rationale behind this hypothesis is that an increase on species richness at basal trophic levels should increase the number of species at higher trophic levels.

Here, I focus on how biotic interactions between fecal detritus producers (mammals) and consumers (dung beetles) (Nichols et al. 2016) can help explain dung beetle diversity pattern along with climatic stability. Dung beetles (Scarabaeinae) is a rich group of detritivores beetles that probably diversified after the transition from saprophagy to coprophagy (Halffter 1991). In this study, two hypothesis will be

evaluated. First, if resources are important for richness patterns of dung beetles, than it is expected that dung beetles diversity should be driven by mammals' diversity, since mammals dungs are the primordial source of food resource for dung beetles (Halffter and Matthews 1966). Thus, a reduction on mammals richness because of hunting should lead to a reduction on dung beetles richness (Nichols et al. 2009). Besides that, a shift on mammals species composition may explain changes on dung beetle composition (Bogoni et al. 2016). Second, if variations on climatic stability should affect dung beetles, than I expect to find that an increase on climatic stability should have positive effects on dung beetle richness. In general, many insects display seasonality patterns of diversity (Wolda 1988), and specifically studies found changes on richness of dung beetle communities as result of temporal rainfall variation (Andresen 2005; Liberal et al. 2011). Climatic stability should also allow dung beetles to be more specialized on feeding resources and present more species because of niche partitioning (Larsen et al. 2006), especially in tropical areas where higher mammal richness are also expected (Davies and Buckley 2011; Safi et al. 2011).

In this study, I test the hypothesis that dung beetle latitudinal richness is explained by 1) an increase feeding resources available because of biotic interactions with mammals, and 2) an increase on climatic stability. For this, I used a spatialized path analysis. I predict that dung beetle richness should increase due increases on mammals' richness and decreases in precipitation seasonality and temperature seasonality. In the latter, I predict increases on dung beetle richness on two ways: directly and indirectly (via increases on mammal richness) (See Figure 1 for more details on proposed path analysis).

Materials and Methods

Dung beetle database

I performed a search on the Scopus database for dung beetles inventories using the following keywords: "*dung beetle"* or "*Scarabaeinae"*. I only included the period between 1980 and 2016. This search retrieved a total of 1443 articles and after applying the following criteria, the database comprised 115 studies and 213 sites (Figure 2): 1) sampled dung beetles should belong to the subfamily Scarabaeinae (in another words, studies with only Aphodinae, Geotrupinae and Troginae were excluded); 2) the study employed a standardized sampling protocol and clearly stated the number of pitfalls, sites and temporal replication used while sampling the dung beetles. Sampling protocol information was used to the measure sampling effort using in each study. The sampling effort was the total number of pitfalls measured as pitfalls multiplied per area multiplied per temporal replications. Differences on   use of different baits (types of dung, decomposing meat or rotten fruits) were ignored 3) provided a list of sampled species and their respective abundances 4) sampled in the American continent and presented geographic coordinates or specified the municipality where the sampled occurred. If one study sampled more than one site and provided a geographic coordinate for each site, than each site was included on the database. However, species lists were combined in cases where there was only one geographic coordinate for multiples sites. Studies included on the analysis are available on the Online Resource 1.

Explanatory variables

I used mammal richness as a proxy for resource availability because of biotic interactions between dung beetles and mammals (Nichols et al. 2009, 2013). In order to

measure mammal richness, I downloaded the terrestrial mammals red list range

shapefile (IUCN 2016) to estimate the number of mammals richness in the locations

where dung beetles were sampled. For each dung beetle locality, I counted the number

of mammal species' range shapefiles that overlap their coordinates using package

maptools (Bivand and Lewin-Koh 2017).

Climatic stability was measured as precipitation seasonality and temperature

seasonality obtained from the worldclim with 5 arc-minutes resolution (approximately

10 km at equator line) (Hijmans et al. 2005). I opted for a coarser resolution because

many localities had their geographic coordinates manually assigned. These variables are

important drivers of mammals distribution and are commonly used on studies with

species distribution of mammals (e.g.: Moura et al. 2016, Ribeiro et al. 2016). Values of

precipitation seasonality and temperature seasonality were extracted for each locality

using package raster (Hijmans 2016).

Both sampling effort and dung beetle abundance were used to control the effects

of different sampling protocols and efforts on dung beetle richness. Furthermore, dung

beetle abundance was also included as responding negatively to climatic stability and

resource availability.

Data Analysis

A path analysis was built using the piecewise structural equation modelling

approach that allows incorporation of different models by building each model

separately (Shipley 2009; Lefcheck 2016). Additionally, path analysis helps to

disentangles direct and indirect effects of climatic and resources on the richness patterns

(Kissling et al. 2007; Moura et al. 2016). First, I built three generalized least squares

models (GLS) in order to account spatial autocorrelation using the package *nlme*

(Pinheiro et al. 2014). Spatial correlation structure present on the three models was

reduced by selecting the best among four spatial correlations: Spheric, Exponential,

Gaussian and Rational quadratic. The first model was built using mammal richness as

response variable, precipitation seasonality and temperature seasonality as explanatory

variables, and using a rational quadratic spatial correlation structure. The Second model

was built using dung beetle abundance as response variable, precipitation seasonality,

temperature seasonality, mammal richness and sampling effort as explanatory variables,

and using an exponential spatial correlation structure. The third model was built using

dung beetle richness as response variable and precipitation seasonality, temperature

seasonality, mammal richness and dung beetle abundance as explanatory variables.

These three models were assembled in a path analysis using the package *piecewiseSEM*

(Lefcheck 2016). Significance of the piecewise SEM was evaluated using Fisher's C

statistic in which a p-value above 0.05 is an indication that the model fits well to the

data.

In all models dung beetle abundance and dung beetle richness were log10

transformed in order to achieve residuals with normal distribution. All analyses were

carried out on R environment (R Core Team 2017).

Results

Dung beetle richness ranged from one to 101 species (mean $\pm$ SD = 21.5 $\pm$ 17.2),

whereas mammals richness ranged from three to 193 species (113.9 $\pm$ 39.5). Dung

beetle abundance ranged from seven to 93,274 individuals (5,226.8 $\pm$ 11,606.66).

I found that the proposed path analysis had a good fit to the observed data

(Fisher's C = 5.05; d.f. =4; P=0.282; Figure 3). Mammal richness was negatively

affected both by temperature seasonality and precipitation seasonality (R^2=71.31%)

whereas dung beetle abundance variation was explained only by sampling effort

(R^2=9.03%).

In the model 56.32% of the variation on dung beetle richness was explained. An

increase on four units of dung beetle abundance resulted on an increase of one unit on

dung beetle richness (b = 0.2472), and an increase of mammal richness resulted on an

increase of dung beetle richness (b_{log10}=0.0023; $b_{antilog}$= 1.01). As expected, negative

effects on dung beetle richness included a direct and an indirect effect, modulated

through mammal richness, of temperature seasonality (b_{direct}=-0.0001and $b_{indirect}$= -

0.00003), but only indirect effects of precipitation seasonality dung beetle richness

($b_{indirect}$= -0.0005).

Discussion

Here I have corroborated my hypothesis that dung beetle richness was higher on

areas with higher climatic stability and higher resource availability. Despite this, dung

beetle abundance was poorly explained on the model and that could be a consequence of

lack of standardization between sampling protocols of studies (Larsen and Forsyth

2005; da Silva and Hernández 2015). As far as I known, this is among the firsts

empirical assessments of dung beetle richness pattern (see: Frank et al. 2018), since

only anecdotal evidence and local scale assessments were previously made (Halffter and

Matthews 1966; Hernández and Vaz-de-Mello 2009; Liberal et al. 2011).

Two non-exclusive explanations help to understand why there are more dung

beetle species on areas with more resources available (i.e. higher mammals' richness).

First, an increase on resource availability should allow dung beetles to specialize on

certain types of dung (e.g.: Larsen et al. 2006, Jacobs et al. 2008) and reduce

interspecific competition between dung beetles. However, a recent study have shown an

opposite trend – dung beetles have low specificity on resource use even on communities

with high richness (Frank et al. 2018). Second, an increase on mammal richness could

result on an increase of different traits and lineages of mammals (Safi et al. 2011). That

in turn can increase the number of mammals with different activity times (e.g.: diurnal

or nocturnal) and with different feeding and digestive systems (e.g.: herbivores or

carnivores). Indeed, dung beetle community structure changes between night and day

(Lopes et al. 2011) or depending on dung type used as bait (Filgueiras et al. 2009).

Climatic instability affects directly the time of activity when dung beetles can be

active due to physiological constraints, mostly of small dung beetles (<2g) that are

thermoconformers (Verdú et al. 2006). Considering that areas with more climatic

instability should have higher variation on temperature during the day and during the

year, this could in turn limit the number of dung beetles species active. Additionally,

climatic instability should also increase the number of generalist dung beetle species in

order to deal with decrease on resource availability, climatic instability also reduces the

number of mammals' species, which reduces dung beetle species due to interspecific

competition (Finke and Snyder 2008). Surprisingly, dung beetle richness was not

directly affected by precipitation seasonality. This, once again points towards climatic

instability having an indirect effect through resource availability.

Here I had shown the importance of incorporating biotic interactions in order to

decouple direct and indirect effects of climatic on species richness patterns.

Furthermore, I present analytical evidence that dung beetle richness is largely affected

by mammals' richness and that mammal's richness can mediate climatic instability

effects on dung beetles.

Conflict of Interest: The author declares that he have no conflict of interest.

References

Andresen E (2005) Effects of Season and Vegetation Type on Community Organization

of Dung Beetles in a Tropical Dry Forest. Biotropica 37:291–300. doi:

10.1111/j.1744-7429.2005.00039.x

Andrews P, O'Brien EM (2000) Climate, vegetation, and predictable gradients in

mammal species richness in southern Africa. J Zool 251:205–231. doi:

10.1111/j.1469-7998.2000.tb00605.x

Bivand R, Lewin-Koh N (2017) maptools: Tools for Reading and Handling Spatial

Objects. R package version 0.9-2. https://cran.r-project.org/package=maptools

Bogoni JA, Graipel ME, de Castilho PV, et al (2016) Contributions of the mammal

community, habitat structure, and spatial distance to dung beetle community

structure. Biodivers Conserv 25:1661–1675. doi: 10.1007/s10531-016-1147-1

da Silva PG, Hernández MIM (2015) Spatial Patterns of Movement of Dung Beetle

Species in a Tropical Forest Suggest a New Trap Spacing for Dung Beetle

Biodiversity Studies. PLoS One 10:e0126112. doi: 10.1371/journal.pone.0126112

Davies TJ, Buckley LB (2011) Phylogenetic diversity as a window into the evolutionary

and biogeographic histories of present-day richness gradients for mammals. Philos

Trans R Soc B Biol Sci 366:2414–2425. doi: 10.1098/rstb.2011.0058

Field R, Hawkins B a., Cornell H V., et al (2009) Spatial species-richness gradients

across scales: A meta-analysis. J Biogeogr 36:132–147. doi: 10.1111/j.1365-

2699.2008.01963.x

Filgueiras BKC, Liberal CN, Aguiar CDM, et al (2009) Attractivity of omnivore,

carnivore and herbivore mammalian dung to Scarabaeinae (Coleoptera,

Scarabaeidae) in a tropical Atlantic rainforest remnant. Rev Bras Entomol 53:422–

427. doi: 10.1590/S0085-56262009000300017

Finke DL, Snyder WE (2008) Niche Increases Resource Partitioning by Diverse

Communities Exploitation. Science (80-) 321:1488–1490. doi:

10.1126/science.1160854

Frank K, Krell F-T, Slade EM, et al (2018) Global dung webs: high trophic generalism

of dung beetles along the latitudinal diversity gradient. Ecol Lett 21:1229–1236.

doi: 10.1111/ele.13095

Halffter G (1991) Historical and Ecological Factors Determining the Geographical

Distribution of Beetles (Coleoptera Scarabaeidae: Scarabaeinae). Folia

Entomológica Mex 82:194–238

Halffter G, Matthews E (1966) The Natural History of Dung Beetles of the Subfamily

Scarabaeinae (Coleoptera, Scarabaeidae). Folia Entomol Mex 12–14:1–312

Hawkins BA, Diniz-Filho JAF (2004) "Latitude" and Geographic Patterns in Species

Richness. Ecography (Cop) 27:268–272. doi: 10.1111/j.0906-7590.2004.03883.x

Hawkins BA, Pausas JG (2004) Does plant richness influence animal richness?: The

mammals of Catalonia (NE Spain). Divers Distrib 10:247–252. doi:

10.1111/j.1366-9516.2004.00085.x

Hernández MIM, Vaz-de-Mello FZ (2009) Seasonal and spatial species richness

variation of dung beetle (Coleoptera, Scarabaeidae s. str.) in the Atlantic Forest of

southeastern Brazil. Rev Bras Entomol 53:607–613. doi: 10.1590/S0085-

56262009000400010

Hijmans RJ (2016) raster: Geographic Data Analysis and Modeling. R package version

2.5-8. https://CRAN.R-project.org/package=raster

Hijmans RJ, Cameron SE, Parra JL, et al (2005) Very high resolution interpolated

climate surfaces for global land areas. Int J Climatol 25:1965–1978. doi:

10.1002/joc.1276

Hillebrand H (2004) On the Generality of the Latitudinal Diversity Gradient. Am Nat

163:192–211. doi: 10.2307/2678832

IUCN (2016) The IUCN Red List of Threatened Species. Version 2016-1.

http://www.iucnredlist.org

Jacobs J, Nole I, Palminteri S, Ratcliffe B (2008) First come, first serve: "sit and wait"

behavior in dung beetles at the source of primate dung. Neotrop Entomol 37:641–5

Kissling WD, Rahbek C, Bohning-Gaese K (2007) Food plant diversity as broad-scale

determinant of avian frugivore richness. Proc R Soc B Biol Sci 274:799–808. doi:

10.1098/rspb.2006.0311

Larsen TH, Forsyth A (2005) Trap Spacing and Transect Design for Dung Beetle

Biodiversity Studies. Biotropica 37:322–325

Larsen TH, Lopera A, Forsyth A (2006) Extreme trophic and habitat specialization by

Peruvian dung beetles (Coleoptera: Scarabaeidae: Scarabaeinae). Coleopt Bull

60:315–324

Lefcheck JS (2016) piecewiseSEM□: Piecewise structural equation modelling in r for

ecology, evolution, and systematics. Methods Ecol Evol 7:573–579. doi:

10.1111/2041-210X.12512

Liberal CN, Isidro de Farias Â., Meiado MV, et al (2011) How habitat change and

rainfall affect dung beetle diversity in Caatinga, a Brazilian semi-arid ecosystem. J

Insect Sci 11:1–11

Lopes J, Korasaki V, Catelli LL, et al (2011) A comparison of dung beetle assemblage

structure (Coleoptera: Scarabaeidae: Scarabaeinae) between an Atlantic forest

fragment and adjacent abandoned pasture in Paraná, Brazil. Zool (Curitiba,

Impresso) 28:72–79. doi: 10.1590/S1984-46702011000100011

Moles AT, Ollerton J (2016) Is the notion that species interactions are stronger and

more specialized in the tropics a zombie idea? Biotropica 48:141–145. doi:

10.1111/btp.12281

Moura MR, Villalobos F, Costa GC, Garcia PCA (2016) Disentangling the Role of

Climate, Topography and Vegetation in Species Richness Gradients. PLoS One

11:e0152468. doi: 10.1371/journal.pone.0152468

Nichols E, Gardner T a., Peres C a., Spector S (2009) Co-declining mammals and dung

beetles: an impending ecological cascade. Oikos 118:481–487. doi:

10.1111/j.1600-0706.2008.17268.x

Nichols E, Peres CA, Hawes JE, Naeem S (2016) Multitrophic diversity effects of

network degradation. Ecol Evol 6:4936–4946. doi: 10.1002/ece3.2253

Nichols E, Uriarte M, Peres C a, et al (2013) Human-Induced Trophic Cascades along

the Fecal Detritus Pathway. PLoS One 8:e75819. doi:

10.1371/journal.pone.0075819

Pianka ER (1966) Latitudinal Gradients in Species Diversity: A Review of Concepts.

Am Nat 100:33. doi: 10.1086/282398

Pinheiro J, Bates D, DebRoy S, et al (2014) nlme: Linear and Nonlinear Mixed Effects

Models_. R package version 3.1-117, URL: http://CRAN.R-

project.org/package=nlme

R Core Team (2017) R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-

project.org/.

Ribeiro BR, Sales LP, De Marco P, Loyola R (2016) Assessing Mammal Exposure to

Climate Change in the Brazilian Amazon. PLoS One 11:e0165073. doi:

10.1371/journal.pone.0165073

Safi K, Cianciaruso M V., Loyola RD, et al (2011) Understanding global patterns of

303   mammalian functional and phylogenetic diversity. Philos Trans R Soc B Biol Sci

304   366:2536–2544. doi: 10.1098/rstb.2011.0024

305  Shipley B (2009) Confirmatory path analysis in a generalized multilevel context.

306   Ecology 90:363–368. doi: 10.1890/08-1034.1

307  Verdú JR, Arellano L, Numa C (2006) Thermoregulation in endothermic dung beetles

308   (Coleoptera: Scarabaeidae): effect of body size and ecophysiological constraints in

309   flight. J Insect Physiol 52:854–60. doi: 10.1016/j.jinsphys.2006.05.005

310  Williams SE, Middleton J (2008) Climatic seasonality, resource bottlenecks, and

311   abundance of rainforest birds: Implications for global climate change. Divers

312   Distrib 14:69–77. doi: 10.1111/j.1472-4642.2007.00418.x

313  Willig MR, Kaufman DM, Stevens RD (2003) Latitudinal Gradients of Biodiversity:

314   Pattern, Process, Scale, and Synthesis. Annu Rev Ecol Evol Syst 34:273–309. doi:

315   10.1146/annurev.ecolsys.34.012103.144032

316  Wolda H (1988) Insect seasonality: why? Annu Rev Ecol Syst 19:1–18

Figures legends

Figure 1. Path analysis representing the theoretical relationships between predictors of

dung beetle richness. Black arrows are negative effects while gray arrows are

positive relationships.

Figure 2. Localities of the 115 dung beetle inventories performed on the America

continent and used on the analysis.

Figure 3. Observed path analysis used to predict dung beetle richness. Dashed arrows

are non-significant relationships between variables ($p>0.05$) while continuous arrows

are significant relationships ($p<0.05$). Black arrows are negative effects while dark gray

arrows are positive effects. Double arrow represents correlations between exogenous

variables. Note that both dung beetle richness and dung beetle abundance was log

transformed.