Coding of whisker motion across the mouse face

Kyle S. Severson[1], Duo Xu[1], Hongdian Yang[1,2], Daniel H. O'Connor[1,*]

[1]The Solomon H. Snyder Department of Neuroscience, Kavli Neuroscience Discovery Institute, Brain Science Institute, The Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

[2]Current address: Department of Molecular, Cell and Systems Biology, University of California, Riverside, Riverside, CA 92521, USA

*Correspondence: dan.oconnor@jhmi.edu

Abstract

Haptic perception synthesizes touch with proprioception, or sense of body position. Humans and mice alike experience rich active touch of the face. Because most facial muscles lack proprioceptor endings, the sensory basis of facial proprioception remains unsolved. Facial proprioception may instead rely on mechanoreceptors that encode both touch and self-motion. In rodents, whisker mechanoreceptors provide a signal that informs the brain about whisker position. Whisking involves coordinated orofacial movements, so mechanoreceptors innervating facial regions other than whiskers could also provide information about whisking. To define all sources of sensory information about whisking available to the brain, we recorded spikes from mechanoreceptors innervating diverse parts of the face. Whisker motion was encoded best by whisker mechanoreceptors, but also by those innervating whisker pad hairy skin and supraorbital vibrissae. Redundant self-motion responses may provide the brain with a stable proprioceptive signal despite mechanical perturbations such as whisker growth and active touch.

Introduction

Proprioception is the sense of where the body or its parts are in space. To interpret touch, it is critical that the brain also knows where in space the touched body part was at the time of contact. Thus, touch and proprioception are intimately linked during normal sensory-motor function. Touch begins with the activation of low-threshold mechanoreceptors (LTMRs) in the skin. Information about body position can come from efference copy signals that report the motor commands ultimately used to control muscles. However, the nervous system contains dedicated mechanoreceptive proprioceptor endings to provide feedback about actual, rather than intended, position. Classical proprioceptors include muscle spindle and Golgi tendon organ afferents.

Many rodents use rapid motions of their mystacial vibrissae (whiskers) to explore the tactile world (Carvell and Simons, 1990; Welker, 1964; Wineski, 1983). Curiously, the muscles controlling these "whisking" motions, as with other facial muscles, lack classical proprioceptor endings (Moore et al., 2015). Therefore, feedback about whisker position must occur via self-motion-triggered ("reafferent") activity of peripheral mechanoreceptors other than classical muscle proprioceptors, such as the cutaneous LTMRs responsible for sensing touch. Neurons throughout the whisker somatosensory system respond to whisker self-motion in a manner that depends on the relative position of the whisker within the current whisk cycle, or whisk "phase" (Campagner et al., 2016; Crochet and Petersen, 2006; Curtis and Kleinfeld, 2009; Fee et al., 1997; Hires et al., 2015; Khatri et al., 2009; Leiser and Moxon, 2007; Moore et al., 2015; Severson et al., 2017; Szwed et al., 2003; Wallach et al., 2016; Yu et al., 2006; Yu et al., 2016). Whisk phase is thought to be a key coordinate system for whisker-based sensation (Curtis and Kleinfeld, 2009; Kleinfeld and Deschenes, 2011; Szwed et al., 2003).

LTMRs that innervate the whisker follicles encode whisk phase and other aspects of whisker motion and touch during active sensing (Bush et al., 2016; Campagner et al., 2016; Khatri et al., 2009; Leiser and Moxon, 2007; Severson et al., 2017; Szwed et al., 2003; Szwed

et al., 2006; Wallach et al., 2016). These LTMRs include Merkel-type endings, which are slowly

adapting and thought to play a major role in perception of object shape and texture. Individual

Merkel and unidentified slowly adapting whisker afferents respond both to touch and to self-

motion (Severson et al., 2017). Self-motion (reafferent) responses arise from diverse

mechanical sensitivities of whisker afferents (Campagner et al., 2016; Severson et al., 2017;

Wallach et al., 2016), and may be used by the brain for whisker proprioception (for behavioral

studies addressing whisker proprioception, see: Knutsen et al., 2006; Mehta et al., 2007;

O'Connor et al., 2013). Neurons in the brainstem (one synapse downstream from

mechanoreceptors) with tactile receptive fields on parts of the face other than whiskers can also

respond during whisking in a manner that reports whisk phase (Moore et al., 2015). This

suggests that mechanoreceptors innervating facial parts other than whiskers could encode

whisker motion, including whisk phase, but this has not been tested. What is the full set of

possible mechanoreceptor sources of information that could tell the mouse brain about whisker

motion, and how do they compare to one another?

Here, we addressed this question by recording whisker motion and electrophysiological

responses from primary mechanoreceptor afferents innervating several distinct structures on the

face, including regions of hairy skin, vibrissae other than the mystacial whiskers, and jaw

muscles (Figure 1A). We compared the encoding of whisker motion among these different

populations of mechanoreceptors to that of whisker mechanoreceptors. We find that a subset of

hairy skin mechanoreceptors encodes whisker motion at levels comparable to whisker

mechanoreceptors. However, as a population, whisker and other non-whisker vibrissae

mechanoreceptors encode the most information about whisker motion. Our results suggest that

information about whisking arises from multiple sensory sources, providing the brain with a

robust basis for facial proprioception.

Results

Self-motion encoding by whisker mechanoreceptors

We obtained electrophysiological and high-speed video recordings (500 Hz) from head-fixed mice as they ran on a treadmill and whisked freely in air (Figure 1B; Supplemental Video 1). From these video frames, we measured several kinematic variables derived from the whisker's angular position (θ) (Figure 1C). During whisking, θ can be decomposed into three quantities that the brain appears to process differently (Hill et al., 2011): midpoint (θ_{mid}), amplitude (θ_{amp}), and phase (Φ) (Figure 1D; Materials and Methods). Whisker primary motor cortex (wM1) robustly encodes θ_{mid} and θ_{amp} (Hill et al., 2011; Huber et al., 2012), and sends this information along cortico-cortical pathways to primary somatosensory cortex (wS1) (Petreanu et al., 2012). This suggests that the brain could use efference copy to keep track of θ_{mid} and θ_{amp}. A small fraction of neurons in wM1 does encode Φ, including after transection of the infraorbital nerve (Hill et al., 2011) that carries sensory information from the whisker pad (Dorfl, 1985). However, the encoding of Φ by neurons across all levels of the ascending somatosensory system, as well as the elimination of phase signals in wS1 after peripheral block of whisking (Fee et al., 1997), indicate a major reafferent contribution to phase coding in the brain. For this reason, and because whisk phase is thought to be a key coordinate scheme for whisker sensation, we focused analysis of the encoding of whisk phase. However, we also analyzed encoding of θ_{mid} and θ_{amp} to determine whether they too could be directly sensed, and encoding of whisker angle (θ), angular velocity (θ'), and angular acceleration (θ''), as these quantities give insight into the mechanical basis of what makes mechanoreceptors spike (Campagner et al., 2018; Severson et al., 2017; Wallach et al., 2016). We aligned these kinematic quantities with simultaneously recorded spikes from different classes of facial mechanoreceptor afferents (Figure 1E).

We first analyzed units in the trigeminal ganglion (TG) with touch receptive fields confined to single whiskers (Figure 2A, n = 67). Many of these afferents were direction-selective, preferring manual deflections in either the protraction or retraction direction (not

shown). A subset of these whisker afferents was more active during whisking (n = 42 "whisking-

sensitive" units, defined in Glossary), consistent with prior work (see Materials and Methods for

comparison to Szwed et al., 2003) and strongly modulated by phase, preferring to fire at a

particular phase of the whisk cycle (Figure 2B,C). This sharp phase tuning largely reflects

sensitivity to inertial stresses (Severson et al., 2017). We used information theory analyses to

quantify how well spiking of single mechanoreceptors encoded phase and other variables

related to whisker kinematics. Specifically, we calculated the mutual information (MI; Cover and

Thomas, 2006), a measure of association between two random variables derived from their joint

probability distribution (Figure 2D), between (1) spike counts obtained during 2 ms video

frames, and (2) binned values (Materials and Methods) of kinematic variables extracted from the

video frames, including θ, θ', θ'', θ_{amp}, θ_{mid}, and Φ. Mutual information between phase and spike

count for whisker afferents, expressed as an information rate via multiplying by the 500 Hz

sampling frequency, was 9.1 ± 23.8 bits/s (median ± interquartile range [IQR]; n = 42 whisking-

sensitive units). To determine which kinematic variable best accounted for the spiking of whisker

afferents, we calculated a "normalized mutual information" by dividing MI by the spike count

entropy (Jamali et al., 2016). This quantity gives the fraction of spike count uncertainty

accounted for by a given kinematic variable. Whisker afferent spike counts were better

explained by phase (Figure 2E; normalized MI = 0.096 ± 0.121; median ± IQR) than by θ (0.034

± 0.064), θ' (0.046 ± 0.086), θ'' (0.030 ± 0.076), θ_{amp} (0.012 ± 0.024), or θ_{mid} (0.0089 ± 0.013; p <

0.0031 for all 5 comparisons, two-tailed K-S tests).

Whisker motion coding by mechanoreceptors innervating hairy skin

While whisker mechanoreceptors showed strong phase coding, our goal was to put this coding

into context by comparing the information provided by these whisker afferents to that of any

other types of mechanoreceptor we could find that responded during whisking in air. We began

by recording from TG units with touch receptive fields on hairy skin (n = 85) rather than a

vibrissa (Figure 3A). Afferents responded to manual deflections of all small hairs or a small

number of guard hairs within the mapped receptive field, were rapidly adapting, and responded

to touch in all directions (not shown). Remarkably, activity of a large number of facial hairy skin

afferents was modulated during whisking in air (Figure 3B; 58 of 85 were whisking-sensitive).

Consistent with the lack of direction-selectivity, many facial hairy skin afferents fired at multiple

phases (e.g. both protraction and retraction phases) of the whisk cycle (Figure 3C). Phase

coding by hairy skin afferents varied by receptive field location, such that units with receptive

fields closer to the whisker pad tended to encode phase more strongly than those distant from

the pad (Figure 3D; overall MI rate = 1.3 ± 4.5 bits/s; median ± IQR).

We next grouped the hairy skin receptive fields into six different "zones" of the face

(Supplemental Figure 1), including the pad, cheek, snout, eye, lip, and jaw. Units with receptive

fields on the whisker pad (n = 14) were particularly modulated by phase (13 of 14 were

whisking-sensitive). We found several receptive fields comprised of small hairs surrounding

whisker follicles, in between whisker arcs or rows, or flanking the outer whiskers. Receptive

fields on the pad were smaller in area than other regions of the face (Figure 3D). Pad hairy skin

afferent encoding of phase (MI rate = 9.26 ± 8.53 bits/s, median ± IQR, n = 13) was comparable

to whisker afferents and significantly higher than afferents innervating hairy skin on the cheek

(0.73 ± 2.10 bits/s, n = 18), eye (1.11 ± 3.57 bits/s, n = 10), lip (0.75 ± 1.76 bits/s, n = 5), snout

(1.63 ± 1.23 bits/s, n = 9), and jaw (0.08 ± 0.09 bits/s, n = 3; Figure 3E; p < 0.032 for all 5 two-

tailed K-S tests). Across all facial hairy skin afferents, normalized MI (Figure 3F) was

significantly higher for phase (0.0165 ± 0.0348) compared to θ' (0.0062 ± 0.0098), θ'' (0.0056 ±

0.0098), θ_{amp} (0.0085 ± 0.012), and θ_{mid} (0.0094 ± 0.011; p<0.003 for all 4 two-tailed K-S tests),

but similar to θ (0.014 ± 0.023; p = 0.77, two-tailed K-S test).

Video capturing facial motion and whisker position suggested that widespread patterns

of skin strain likely occur in a manner correlated with whisking, with stronger correlations

between skin and whisker displacements occurring for facial regions on or near the whisker pad

(Supplemental Figure 1; Supplemental Video 2). In addition to skin movements, we observed that the vibrissae above the eye whisk in phase with the whiskers. We next focused our attention on vibrissae outside of the whisker pad.

Non-mystacial vibrissa movement correlates with whisking

Mice have several vibrissae outside of the mystacial pad, including two supraorbital vibrissae above the eye, one genal vibrissa on the cheek (Danforth, 1925), and several microvibrissae on the upper lip (Figure 4A; Brecht et al., 1997). These sinus follicle structures are highly conserved within strains of mice (Dun and Fraser, 1958) and are present in many mammals (Danforth, 1925; Grant et al., 2013; Wineski, 1983). Surprisingly, in our high-speed videos we noticed periodic movement of supraorbital vibrissae apparently locked to whisking (Supplemental Video 2). To quantify these movements, we simultaneously tracked non-mystacial vibrissae and whiskers (i.e. mystacial vibrissae) using high-speed videography. Supraorbital vibrissae (Figure 4B-D; Supplemental Video 3) and the genal vibrissa (Figure 4B-D; Supplemental Video 4) moved in phase with the whiskers. We observed some instances of "missed" whisk cycles, in which the whisker moved, but the supraorbital or genal vibrissae remained still (Supplemental Figure 2A-F). Microvibrissa barely moved (Figure 4B-D; Supplemental Video 5). Small translations we observed could be due to passive pulling of lip tissue during whisking, rather than active rotation of the microvibrissa follicle.

We computed cross-correlations to quantify the phase lag and degree of correlation between whiskers and the supraorbital and genal vibrissae, and microvibrissae (Figure 4E). We first analyzed pairs of mystacial whiskers to set an "upper bound" on correlations, as whiskers in the same row have highly correlated movements (Wallach et al., 2016). Adjacent whiskers correlated almost perfectly (Pearson's correlation coefficient, r = 0.98 ± 0.03, n = 14 recordings from 11 mice) with no phase lag (0.00 ± 0.03 radians). These strong correlations among whiskers also validated the use of adjacent or nearby whiskers—chosen for their convenience in

obtaining high-speed videos of both whiskers and other vibrissae—when quantifying

correlations between whiskers and other vibrissae.

Supraorbital vibrissae movements correlated strongly with whisker movements (r = 0.78

± 0.07, mean ± SD, n = 6 recordings from 6 mice), but with a short phase lag (Figure 4C-E; Φ_{lag}

= 0.27 ± 0.05 radians, mean ± SD; p = 4.9e-12, one-tailed t-test). Whisking amplitude (θ_{amp}) was

smaller for supraorbital vibrissae (4.7 ± 2.1°, mean ± SEM) compared to whiskers (Figure 4F;

12.5 ± 2.5°, mean ± SEM; p = 1.1e-6, one-tailed t-test).

Genal vibrissa motion also correlated strongly with that of the whiskers (r = 0.83 ± 0.08,

mean ± SD), but with a longer phase lag (Figure 4C-E; Φ_{lag} = 0.61 ± 0.18 radians, mean ± SD; p

= 9.9e-11, one-tailed t-test). Whisking amplitude for genal vibrissae was also smaller (2.2 ± 1.4°,

mean ± SEM) compared to the tracked whiskers (Figure 4F; p = 6.9e-9, one-tailed t-test) and

supraorbital vibrissae (p = 0.016, one-tailed t-test).

Microvibrissae motion correlated with whisker motion less well (r = 0.27 ± 0.29, mean ±

SD, n = 3 recordings from 3 mice) with a short delay (Figure 4C-E; Φ_{lag} = 0.44 ± 0.11 radians,

mean ± SD; p = 4.8e-10, one-tailed t-test). Whisking amplitude for microvibrissae (Figure 4F,

0.84 ± 0.23°, mean ± SEM) was smaller than for whiskers (p = 3.5e-7, one-tailed t-test) and

supraorbital vibrissae (p = 0.0086, one-tailed t-test). With smaller amplitude movements and

smaller sizes, the mechanical stresses generated at the base of microvibrissae during whisking

are likely smaller than those at the bases of other vibrissae types.

The motion of supraorbital and genal vibrissae could occur under active neuromuscular

control, or passively due to mechanical coupling with the moving mystacial pad. To distinguish

between these possibilities, we performed experiments in which we cut different branches of the

facial nerve, which supplies motor innervation for whisking (Dorfl, 1985; Fee et al., 1997), and

observed the effects on motion of the whiskers and non-mystacial vibrissae (Figure 4G;

Supplemental Figure 2G-K). If the supraorbital or genal vibrissae move actively, abolishing

whisking of the mystacial whiskers should not impact their movement. First, in a new group of

mice (n = 3), we again measured movement of whiskers and non-mystacial vibrissae, but using

dual-view videography such that we could simultaneously measure motion of whiskers on both

the left and right sides of the face, together with motion of the non-mystacial vibrissae on the left

side (Supplemental Figure 2H; Supplemental Videos 6-7). We then lesioned the buccal branch

of the left facial motor nerve, which innervates the mystacial pad (Dorfl, 1985; Fee et al., 1997).

As expected, this manipulation abolished whisking of the mystacial whiskers on the left side

(mean whisker θ_{amp} reduced from 11.2 ± 4.6° to 0.4 ± 0.2°, mean ± SEM, n = 6 videographic

recordings), while leaving whisking intact on the right side (Figure 4G; Supplemental Figure 2I;

Supplemental Video 6). Whisking of the supraorbital vibrissae was intact after buccal cut

(supraorbital θ_{amp}: 4.2 ± 2.4° vs 4.9 ± 2.8°, n = 3 recordings), despite the abolished whisker

motion (Figure 4G; Supplemental Figure 2I; Supplemental Video 6). In contrast, genal vibrissae

showed a reduction in θ_{amp} after buccal cut (genal θ_{amp}: 2.2 ± 1.3° vs 0.8 ± 0.5°, n = 3

recordings; Figure 4G; Supplemental Figure 2J; Supplemental Video 7). Subsequent cut of the

left facial nerve at the junction of the temporal and zygomatic branches (Supplemental Figure

2G) eliminated whisking of the left supraorbital vibrissae (supraorbital θ_{amp}: 0.08 ± 0.05°, n = 3

recordings; Figure 4G; Supplemental Figure 2I). Thus, genal vibrissae motion is passive or

dependent on the buccal innervation of the whisker pad, whereas whisking by the supraorbital

vibrissae is active and under neuromuscular control separate from the mystacial whiskers.

Non-mystacial vibrissa mechanoreceptors encode information about whisking

The motion of non-mystacial vibrissae was correlated with whisker motion during whisking.

Therefore, mechanoreceptors with receptive fields on these vibrissae could show activity

patterns that encode whisker self-motion. To test this possibility, we recorded from TG units with

receptive fields on non-mystacial vibrissae. We found units, some of which were active during

whisking, on supraorbital vibrissae (8 of 17 whisking-sensitive), genal vibrissae (3 of 8 whisking-

sensitive), and microvibrissae (8 of 10 whisking-sensitive). Non-mystacial vibrissa afferent

spiking aligned with whisk phase (Figure 4H). Similar to whisker afferents, we observed

examples of sharp phase tuning (Figure 4I).

Supraorbital afferents encoded similar amounts of information about phase (MI rate =

16.2 ± 14.9 bits/s, median ± IQR) as genal afferents (5.2 ± 8.6 bits/s, median ± IQR; p = 0.23,

two-tailed K-S test), and more than microvibrissa afferents (Figure 4J; 0.74 ± 1.60 bits/s, median

± IQR; p = 0.0014, two-tailed K-S test). The spike counts of non-mystacial vibrissa afferents

overall were better explained by Φ (Figure 4K; normalized MI = 0.062 ± 0.17, median ± IQR)

and θ (0.047 ± 0.042) compared to θ_{amp} (0.019 ± 0.034), and θ_{mid} (0.021 ± 0.033; p < 0.049 for

all 4 two-tailed K-S tests).

252   Information encoded by jaw proprioceptors

The trigeminal mesencephalic nucleus (MeV) resides in the brainstem and contains

mechanoreceptor neurons that innervate the masseter muscles involved in mastication.

Recently, it has been suggested that MeV neurons respond to aspects of whisker motion

(Mameli et al., 2017; Mameli et al., 2010; Mameli et al., 2014), which necessitates their inclusion

in a full account of possible sources of peripheral information about whisker motion available to

the brain (Bosman et al., 2011). We thus recorded the activity of single neurons using 32-

channel tetrode microdrives implanted in MeV (Figure 5A).

As with TG recordings, head-fixed mice were placed on a treadmill to elicit running and

whisking. Mice also licked at a lickport for water rewards. We used this preparation to identify

jaw muscle proprioceptors, as their activity was strongly modulated by the licking associated

with reward consumption (Figure 5B,C). For analysis we considered both these putative jaw

muscle proprioceptors (n = 23 units), plus units that were recorded on the same tetrode as a

putative proprioceptor (n = 20 units) and therefore also presumably in MeV. We did not observe

obvious phasic modulation of MeV activity during whisking (Figure 5B; periods of licking

excluded from this analysis). MeV units (n = 33 whisking-sensitive) were not tuned to whisk

phase (Figure 5D) and thus did not encode much information about phase (0.04 ± 0.04 bits/s,

median ± IQR). Overall, only 2 of 33 whisking-sensitive MeV units (both putative jaw

proprioceptors) had MI confidence intervals for Φ above chance levels (Figure 5E).

However, we did observe a correlation between MeV activity and whisking midpoint

(Figure 5D-F). Eighteen of 33 whisking-sensitive MeV units (including 15 out of 23 putative

proprioceptors) had MI confidence intervals for θ_{mid} above chance levels (Figure 5E). Several

units increased or decreased spiking with increasing θ_{mid} (Figure 5F). Units that most strongly

encoded θ_{mid} were putative jaw proprioceptors (Figure 5G). The kinematic variables that

associated best with MeV spike counts were midpoint, amplitude, and position (Figure 5H).

However, MI values between spike count and these slowly varying quantities were low (e.g. θ_{mid}:

0.17 ± 0.38 bits/s, median ± IQR). Thus, MeV activity is not correlated with whisk phase and

appears only weakly correlated with whisking midpoint. We speculate that this weak correlation

may be explained by slight changes in jaw position associated with whisking around more or

less protracted midpoints.

Comparison of whisker motion coding across facial mechanoreceptor classes

So far, we have described how well whisker afferents and other types of facial

mechanoreceptors encode whisk phase and other variables related to whisker motion. A major

goal was to compare whisker self-motion coding by whisker afferents with that of other classes

of afferents. We therefore directly compared coding of whisk phase and midpoint, given the

importance of these variables in describing whisking behavior and neural activity (Curtis and

Kleinfeld, 2009; Hill et al., 2011; Kleinfeld and Deschenes, 2011; Severson et al., 2017; Wallach

et al., 2016). Overall, as a population the non-mystacial vibrissae afferents best encoded θ_{mid}

(Figure 6A; MI rate = 0.81 ± 1.80 bits/s, median ± IQR), with similar encoding by whisker

afferents (0.68 ± 1.08 bits/s, median ± IQR; p = 0.28, two-tailed K-S test vs non-mystacial

afferents) and facial hairy skin afferents (0.72 ± 1.16 bits/s, median ± IQR; p = 0.61, two-tailed

K-S test). MeV spike counts encoded θ_{mid} less well than all other afferent classes (0.17 ± 0.38

bits/s, median ± IQR; p < 6.7e-4 for all three two-tailed K-S tests). Similarly, normalized mutual

information values showed that θ_{mid} explained spike counts less well for MeV than for all other

afferent classes (Figure 6A; MeV: 0.0019 ± 0.0077; whiskers: 0.0089 ± 0.0134; non-mystacial

vibrissae: 0.021 ± 0.033; facial hairy skin: 0.0094 ± 0.0110; median ± IQR; p < 6.2e-4 for all

three two-tailed K-S tests of MeV vs other classes).

Putative jaw muscle proprioceptors within the set of MeV units tended to show stronger

encoding of θ_{mid} (Figure 5G). Therefore, for a more stringent comparison we compared this

subset of MeV units against whisker afferents and the subgroups of hairy skin and non-

mystacial vibrissae afferents that were most informative about θ_{mid}. Mutual information between

spike count and θ_{mid} was also lower for this subset of MeV jaw afferents (Figure 6B; n = 23; 0.20

± 0.40 bits/s, median ± IQR) compared with the whisker afferents (n = 42 units; 0.66 ± 1.04

bits/s; p = 0.0059, K-S test), hairy skin afferents (n = 31 pad + cheek units; 1.23 ± 1.23 bits/s; p

= 1.41e-4, K-S test), and non-mystacial vibrissae afferents (n = 11 supraorbital + genal units;

1.51 ± 3.69 bits/s; p = 0.0049, K-S test).

Phase was best encoded by whisker (Figure 6C; MI rate = 9.1 ± 23.8 bits/s, median ±

IQR) and non-mystacial vibrissae afferents (2.9 ± 11.9 bits/s, median ± IQR; p = 0.15, two-tailed

K-S test vs whisker) compared with facial hairy skin afferents (1.3 ± 4.5 bits/s, median ± IQR; p

= 3.0e-5, two-tailed K-S test vs whisker) and MeV afferents (0.04 ± 0.04 bits/s, median ± IQR; p

= 8.9e-16, two-tailed K-S test vs whisker). Similarly, normalized mutual information values

showed that phase better explained the spike count of whisker afferents (Figure 6C; 0.096 ±

0.121, median ± IQR) compared with facial hairy skin (0.017 ± 0.035, median ± IQR; p = 1.0e-

10, two-tailed K-S test), non-mystacial vibrissae (0.062 ± 0.166, median ± IQR; p = 0.02, two-

tailed K-S test), and MeV afferents (6e-4 ± 10e-4, median ± IQR; p = 8.9e-17, two-tailed K-S

test).

While whisker afferents as a group were overall best at encoding phase, other

mechanoreceptor classes included more or less informative subgroups. For a more stringent

comparison, we considered the best encoding subgroup from each class: whisker pad

mechanoreceptors within facial hairy skin, supraorbital vibrissa mechanoreceptors within non-

mystacial vibrissae, and putative jaw muscle proprioceptors within MeV. Mutual information

between phase and spike count was similar for whisker (Figure 6D; n = 42; 9.1 ± 23.8 bits/s),

pad (n = 13; 9.3 ± 8.5 bits/s, median ± IQR), and supraorbital mechanoreceptors (n = 8; 16.2 ±

14.9 bits/s; p > 0.11 for all three two-tailed K-S tests), and negligible for jaw proprioceptors (n =

23; 0.04 ± 0.04 bits/s; p < 8.4e-6 for all three two-tailed K-S tests). Thus, while whisker

mechanoreceptors as a group best encode whisk phase, mechanoreceptors with receptive

fields on whisker pad hairy skin and on the supraorbital vibrissae also send to the brain a signal

that encodes whisk phase (Figure 6E).

Discussion

Here we surveyed primary mechanoreceptive afferents that innervate multiple regions of the

face to quantify correlations between spiking activity of these mechanoreceptors and whisker

motion. Our specific goal was to provide a comprehensive account of the possible sources of

reafferent information sent to the brain about whisking. This quantitative survey provides

important context to interpret the encoding of whisker motion—and in particular, whisk phase—

previously observed among whisker afferents (Campagner et al., 2016; Severson et al., 2017;

Szwed et al., 2003; Wallach et al., 2016), and more generally to investigate the hypothesis that

facial proprioception relies on the reafferent activity of cutaneous LTMRs. We found that whisker

afferents as a group encoded whisk phase best, together with supraorbital and genal vibrissae

afferents. Thus, our results support the hypothesis that the strong phase coding observed in

prior work with whisker afferents (Campagner et al., 2016; Severson et al., 2017; Szwed et al.,

2003; Wallach et al., 2016) could serve as a basis for whisker proprioception.

We found that a large number of mechanoreceptors with receptive fields on the hairy

skin of the face responded in a phasic manner during whisking. While passive or active touch of

the whiskers did not strongly activate facial hairy skin mechanoreceptors, passive stretch of the

skin within or near their receptive fields was sufficient to cause spiking (not shown). This

suggests that skin strain occurring within the receptive field, and in a pattern correlated with

whisker motion, underlies the self-motion responses of these afferents.

Mechanoreceptors with receptive fields on the whisker pad were especially informative

about whisker kinematics, presumably because of their proximity to the whiskers and the fact

that motion of the whisker pad itself is an integral part of whisking (Hill et al., 2008). Pad

mechanoreceptor responses could provide the brain with reference signals to use in interpreting

incoming touch responses from whisker afferents. For example, pad afferents encoded the

angular position and velocity of the whiskers, kinematic variables that can be used in

combination with stresses induced by whisker touch to compute both the azimuthal and radial

location of objects (Birdwell et al., 2007; Knutsen and Ahissar, 2009; Mehta et al., 2007;

Solomon and Hartmann, 2011; Szwed et al., 2003). The radial distance of an object contacted

by the whiskers can be determined using ratios of different contact stresses (Pammer et al.,

2013; Solomon and Hartmann, 2011). However, alternative schemes compute radial distance

using knowledge of whisker angle or velocity to disambiguate changes in whisker bending

moment due to object location from those due to self-motion (Birdwell et al., 2007; Solomon and

Hartmann, 2011). Pad afferents may thus play a role in supporting redundant methods for object

localization.

Cutaneous afferents have been reported to be active during self-motion in systems other

than the whisker system. During jaw movements in rabbits, non-direction-selective hairy skin

afferents were found to respond to self-motion in a manner proportional to movement speed

(Appenteng et al., 1982). In humans, microneurography studies have reported activity in

cutaneous afferents related to active movement of the face (Johansson et al., 1988), ankle

(Aimonetti et al., 2007), knee (Edin, 2001), and finger (Edin and Abbs, 1991; Hulliger et al.,

1979). Thus, "cutaneous" (reafferent) signals of potential use for proprioception occur across a

wide variety of body parts and animal species. While our focus was on whisking-related

proprioception, in future work it will be important to understand the degree to which the

mechanoreceptors we recorded encode other aspects of facial motion.

Using high-speed videography, we found correlated motions of the non-mystacial

vibrissae and the mystacial whiskers. In rodents, major aspects of the structure and innervation

(Fundin et al., 1995; Wineski, 1985) of the supraorbital and genal vibrissae closely resemble

those of mystacial vibrissae (Fundin et al., 1994). Motions of these non-mystacial vibrissae were

assessed in the golden hamster, and they were found to be relatively immobile (Wineski, 1983).

In opossum, the genal vibrissae were observed to be mobile, contain intrinsic protractor

muscles, and move in phase with mystacial whiskers (Grant et al., 2013). Here, we show that in

mice the supraorbital and genal vibrissae are indeed mobile and move in phase with mystacial

whiskers. In the case of the supraorbital vibrissae, we confirm that this whisking motion is under

active neuromuscular control and persists after cutting the motor nerve that drives whisking of

the mystacial whiskers. The observation of tight coupling of whisker and non-mystacial vibrissa

movements adds to our understanding of the exquisitely coordinated orofacial motor actions in

rodents (Kurnikova et al., 2017; Welker, 1964) and suggests that their premotor circuits are

linked (Deschenes et al., 2016; Kleinfeld et al., 2014; McElvain et al., 2018; Moore et al., 2013).

Afferents with receptive fields on these structures, especially the supraorbital and genal

vibrissae, displayed strong phase tuning and carried information about phase comparable to

that of whisker afferents.

While the supraorbital and genal vibrissae afferents encoded the phase of the whiskers

in the whisk cycle, these non-mystacial vibrissae are unlikely to contact objects that are in reach

of the whiskers. Thus, an interesting possibility is that afferents with non-mystacial vibrissae

receptive fields could provide the brain with a phase signal that is, unlike that of touch-sensitive

whisker afferents we report here and in past work (Severson et al., 2017), unperturbed by

contacts between whiskers and objects in the world. It will be important to determine whether

the strong correlations in phase we observed between non-mystacial vibrissae and whiskers

generalize across behavioral conditions, such as during behavioral tasks in freely moving

animals. Moreover, tasks that require highly precise information about the phase of a particular

whisker to be combined with touch signals may require direct sensing of phase by afferents

innervating that whisker itself. Whisker afferents that respond to whisking in air but not touch

have been found (Szwed et al., 2003), and could serve the role of separating contact from

phase signals with high precision.

Recent reports have found that neurons in the trigeminal mesencephalic nucleus can be

activated during periods of whisker motion, leading to the suggestion that MeV neurons encode

whisking kinematics (Mameli et al., 2017; Mameli et al., 2010; Mameli et al., 2014). However,

these studies were limited to anesthetized animals. To clarify whether MeV must be considered

as a source of information about whisking kinematics (Bosman et al., 2011) during behavior, we

recorded extracellularly from MeV units during periods of active whisking and during periods of

licking. We found that MeV units did not encode whisk phase nor other rapid aspects of whisker

motion. MeV units did encode the midpoint of whisking, albeit very modestly relative to other

afferent classes. MeV houses the muscle spindles of jaw muscles, which spike during jaw

movements (Goodwin and Luschei, 1975). We therefore speculate that these weak correlations

with midpoint occur due to coordinated motion of the jaw and whisker pad, perhaps with subtle

jaw muscle changes occurring at more protracted whisking midpoints (which occur at higher

locomotion speeds; Sofroniew et al., 2014). However, we identified MeV units in our

extracellular recordings based on responses to licking (presumably jaw-motion-correlated), or

based on a unit being recorded on the same tetrode (nearby location) as a licking-correlated

unit. It is possible that MeV houses neurons that we did not sample and that encode other
aspects of whisking.

Together, our results provide a quantitative survey of how much information
mechanoreceptors in the face can provide the mouse brain about whisking. Our data reveal that
non-mystacial vibrissae can whisk in phase with the whiskers, and that mechanoreceptors
innervating these non-mystacial vibrissae, as well as a subset of mechanoreceptors innervating
facial hairy skin, can provide the brain with information about whisker motion comparable to
mechanoreceptors that innervate the whiskers. Whisker mechanoreceptors provided the best,
but not the only, source of information about whisking for the brain to use in whisker
proprioception. We conclude that the coding of whisker self-motion occurs via a multitude of
sensory signals arising from distinct classes of facial mechanoreceptors.

Author Contributions

KSS and HY performed experiments. KSS and DX developed analysis code. KSS analyzed
data. KSS and DHO wrote the paper with input from all authors.

Acknowledgements

We thank Ernst Niebur for discussion of data analysis and Bilal Bari for demonstration of MeV
recording. We thank Ernst Niebur, Kathleen Cullen, Jeremiah Cohen and William Olson for
comments on the manuscript. The authors were supported by NIH grants R01NS089652 and
1R01NS104834-01.

References

Aimonetti, J.M., Hospod, V., Roll, J.P., and Ribot-Ciscar, E. (2007). Cutaneous afferents provide a
neuronal population vector that encodes the orientation of human ankle movements. J Physiol *580*,
649-658.

Appenteng, K., Lund, J.P., and Seguin, J.J. (1982). Behavior of cutaneous mechanoreceptors recorded in
mandibular division of Gasserian ganglion of the rabbit during movements of lower jaw. J Neurophysiol
47, 151-166.
Birdwell, J.A., Solomon, J.H., Thajchayapong, M., Taylor, M.A., Cheely, M., Towal, R.B., Conradt, J., and
Hartmann, M.J. (2007). Biomechanical models for radial distance determination by the rat vibrissal
system. J Neurophysiol *98*, 2439-2455.
Bosman, L.W., Houweling, A.R., Owens, C.B., Tanke, N., Shevchouk, O.T., Rahmati, N., Teunissen, W.H.,
Ju, C., Gong, W., Koekkoek, S.K.*, et al.* (2011). Anatomical pathways involved in generating and sensing
rhythmic whisker movements. Front Integr Neurosci *5*, 53.
Brecht, M., Preilowski, B., and Merzenich, M.M. (1997). Functional architecture of the mystacial
vibrissae. Behav Brain Res *84*, 81-97.
Bush, N.E., Schroeder, C.L., Hobbs, J.A., Yang, A.E., Huet, L.A., Solla, S.A., and Hartmann, M.J. (2016).
Decoupling kinematics and mechanics reveals coding properties of trigeminal ganglion neurons in the
rat vibrissal system. Elife *5*.
Campagner, D., Evans, M.H., Bale, M.R., Erskine, A., and Petersen, R.S. (2016). Prediction of primary
somatosensory neuron activity during active tactile exploration. Elife *5*.
Campagner, D., Evans, M.H., Loft, M.S.E., and Petersen, R.S. (2018). What the whiskers tell the brain.
Neuroscience *368*, 95-108.
Carvell, G.E., and Simons, D.J. (1990). Biometric analyses of vibrissal tactile discrimination in the rat. J
Neurosci *10*, 2638-2648.
Clack, N.G., O'Connor, D.H., Huber, D., Petreanu, L., Hires, A., Peron, S., Svoboda, K., and Myers, E.W.
(2012). Automated tracking of whiskers in videos of head fixed rodents. PLoS Comput Biol *8*, e1002591.
Cohen, J.Y., Haesler, S., Vong, L., Lowell, B.B., and Uchida, N. (2012). Neuron-type-specific signals for
reward and punishment in the ventral tegmental area. Nature *482*, 85-88.
Cover, T.M., and Thomas, J.A. (2006). Elements of information theory, 2nd edn (Hoboken, N.J.: Wiley-
Interscience).
Crochet, S., and Petersen, C.C. (2006). Correlating whisker behavior with membrane potential in barrel
cortex of awake mice. Nat Neurosci *9*, 608-610.
Curtis, J.C., and Kleinfeld, D. (2009). Phase-to-rate transformations encode touch in cortical neurons of a
scanning sensorimotor system. Nat Neurosci *12*, 492-501.
Danforth, C.H. (1925). Hair in its relation to questions of homology and phylogeny. American Journal of
Anatomy *36*, 47-68.
Deschenes, M., Takatoh, J., Kurnikova, A., Moore, J.D., Demers, M., Elbaz, M., Furuta, T., Wang, F., and
Kleinfeld, D. (2016). Inhibition, Not Excitation, Drives Rhythmic Whisking. Neuron *90*, 374-387.
Dorfl, J. (1985). The innervation of the mystacial region of the white mouse: A topographical study. J
Anat *142*, 173-184.
Dun, R.B., and Fraser, A.S. (1958). Selection for an invariant character; vibrissa number in the house
mouse. Nature *181*, 1018-1019.
Edin, B. (2001). Cutaneous afferents provide information about knee joint movements in humans. J
Physiol *531*, 289-297.
Edin, B.B., and Abbs, J.H. (1991). Finger movement responses of cutaneous mechanoreceptors in the
dorsal skin of the human hand. J Neurophysiol *65*, 657-670.
Fee, M.S., Mitra, P.P., and Kleinfeld, D. (1997). Central versus peripheral determinants of patterned
spike activity in rat vibrissa cortex during whisking. J Neurophysiol *78*, 1144-1149.
Fundin, B.T., Arvidsson, J., and Rice, F.L. (1995). Innervation of nonmystacial vibrissae in the adult rat. J
Comp Neurol *357*, 501-512.
Fundin, B.T., Rice, F.L., Pfaller, K., and Arvidsson, J. (1994). The innervation of the mystacial pad in the
adult rat studied by anterograde transport of HRP conjugates. Exp Brain Res *99*, 233-246.

Goodwin, G.M., and Luschei, E.S. (1975). Discharge of spindle afferents from jaw-closing muscles during
chewing in alert monkeys. J Neurophysiol *38*, 560-571.
Grant, R.A., Haidarliu, S., Kennerley, N.J., and Prescott, T.J. (2013). The evolution of active vibrissal
sensing in mammals: evidence from vibrissal musculature and function in the marsupial opossum
Monodelphis domestica. J Exp Biol *216*, 3483-3494.
Hill, D.N., Bermejo, R., Zeigler, H.P., and Kleinfeld, D. (2008). Biomechanics of the vibrissa motor plant in
rat: rhythmic whisking consists of triphasic neuromuscular activity. J Neurosci *28*, 3438-3455.
Hill, D.N., Curtis, J.C., Moore, J.D., and Kleinfeld, D. (2011). Primary motor cortex reports efferent control
of vibrissa motion on multiple timescales. Neuron *72*, 344-356.
Hires, S.A., Gutnisky, D.A., Yu, J., O'Connor, D.H., and Svoboda, K. (2015). Low-noise encoding of active
touch by layer 4 in the somatosensory cortex. Elife *4*.
Huber, D., Gutnisky, D.A., Peron, S., O'Connor, D.H., Wiegert, J.S., Tian, L., Oertner, T.G., Looger, L.L., and
Svoboda, K. (2012). Multiple dynamic representations in the motor cortex during sensorimotor learning.
Nature *484*, 473-478.
Hulliger, M., Nordh, E., Thelin, A.E., and Vallbo, A.B. (1979). The responses of afferent fibres from the
glabrous skin of the hand during voluntary finger movements in man. J Physiol *291*, 233-249.
Jamali, M., Chacron, M.J., and Cullen, K.E. (2016). Self-motion evokes precise spike timing in the primate
vestibular system. Nat Commun *7*, 13229.
Johansson, R.S., Trulsson, M., Olsson, K.A., and Abbs, J.H. (1988). Mechanoreceptive afferent activity in
the infraorbital nerve in man during speech and chewing movements. Exp Brain Res *72*, 209-214.
Khatri, V., Bermejo, R., Brumberg, J.C., Keller, A., and Zeigler, H.P. (2009). Whisking in air: encoding of
kinematics by trigeminal ganglion neurons in awake rats. J Neurophysiol *101*, 1836-1846.
Kleinfeld, D., and Deschenes, M. (2011). Neuronal basis for object location in the vibrissa scanning
sensorimotor system. Neuron *72*, 455-468.
Kleinfeld, D., Moore, J.D., Wang, F., and Deschenes, M. (2014). The Brainstem Oscillator for Whisking
and the Case for Breathing as the Master Clock for Orofacial Motor Actions. Cold Spring Harb Symp
Quant Biol *79*, 29-39.
Knutsen, P.M., and Ahissar, E. (2009). Orthogonal coding of object location. Trends Neurosci *32*, 101-
109.
Knutsen, P.M., Pietr, M., and Ahissar, E. (2006). Haptic object localization in the vibrissal system:
behavior and performance. J Neurosci *26*, 8451-8464.
Kurnikova, A., Moore, J.D., Liao, S.-M., Deschenes, M., and Kleinfeld, D. (2017). Coordination of orofacial
motor actions of the nose, head, and vibrissae into exploratory behavior. Curr Biol.
Leiser, S.C., and Moxon, K.A. (2007). Responses of trigeminal ganglion neurons during natural whisking
behaviors in the awake rat. Neuron *53*, 117-133.
Mameli, O., Caria, M.A., Biagi, F., Zedda, M., and Farina, V. (2017). Neurons within the trigeminal
mesencephalic nucleus encode for the kinematic parameters of the whisker pad macrovibrissae. Physiol
Rep *5*.
Mameli, O., Stanzani, S., Mulliri, G., Pellitteri, R., Caria, M.A., Russo, A., and De Riu, P. (2010). Role of the
trigeminal mesencephalic nucleus in rat whisker pad proprioception. Behav Brain Funct *6*, 69.
Mameli, O., Stanzani, S., Russo, A., Pellitteri, R., Manca, P., De Riu, P.L., and Caria, M.A. (2014).
Involvement of trigeminal mesencephalic nucleus in kinetic encoding of whisker movements. Brain Res
Bull *102*, 37-45.
McElvain, L.E., Friedman, B., Karten, H.J., Svoboda, K., Wang, F., Deschenes, M., and Kleinfeld, D. (2018).
Circuits in the rodent brainstem that control whisking in concert with other orofacial motor actions.
Neuroscience *368*, 152-170.
Mehta, S.B., Whitmer, D., Figueroa, R., Williams, B.A., and Kleinfeld, D. (2007). Active spatial perception
in the vibrissa scanning sensorimotor system. PLoS Biol *5*, e15.

Moore, J.D., Deschenes, M., Furuta, T., Huber, D., Smear, M.C., Demers, M., and Kleinfeld, D. (2013).
Hierarchy of orofacial rhythms revealed through whisking and breathing. Nature *497*, 205-210.
Moore, J.D., Mercer Lindsay, N., Deschenes, M., and Kleinfeld, D. (2015). Vibrissa self-motion and touch
are reliably encoded along the same somatosensory pathway from brainstem through thalamus. PLoS
Biol *13*, e1002253.
O'Connor, D.H., Hires, S.A., Guo, Z.V., Li, N., Yu, J., Sun, Q.Q., Huber, D., and Svoboda, K. (2013). Neural
coding during active somatosensation revealed using illusory touch. Nat Neurosci *16*, 958-965.
Pammer, L., O'Connor, D.H., Hires, S.A., Clack, N.G., Huber, D., Myers, E.W., and Svoboda, K. (2013). The
mechanical variables underlying object localization along the axis of the whisker. J Neurosci *33*, 6726-
6741.
Petreanu, L., Gutnisky, D.A., Huber, D., Xu, N.L., O'Connor, D.H., Tian, L., Looger, L., and Svoboda, K.
(2012). Activity in motor-sensory projections reveals distributed coding in somatosensation. Nature *489*,
299-303.
Severson, K.S., Xu, D., Van de Loo, M., Bai, L., Ginty, D.D., and O'Connor, D.H. (2017). Active Touch and
Self-Motion Encoding by Merkel Cell-Associated Afferents. Neuron *94*, 666-676 e669.
Sofroniew, N.J., Cohen, J.D., Lee, A.K., and Svoboda, K. (2014). Natural whisker-guided behavior by head-
fixed mice in tactile virtual reality. J Neurosci *34*, 9537-9550.
Solomon, J.H., and Hartmann, M.J. (2011). Radial distance determination in the rat vibrissal system and
the effects of Weber's law. Philos Trans R Soc Lond B Biol Sci *366*, 3049-3057.
Suter, B.A., O'Connor, T., Iyer, V., Petreanu, L.T., Hooks, B.M., Kiritani, T., Svoboda, K., and Shepherd,
G.M. (2010). Ephus: multipurpose data acquisition software for neuroscience experiments. Front Neural
Circuits *4*, 100.
Szwed, M., Bagdasarian, K., and Ahissar, E. (2003). Encoding of vibrissal active touch. Neuron *40*, 621-
630.
Szwed, M., Bagdasarian, K., Blumenfeld, B., Barak, O., Derdikman, D., and Ahissar, E. (2006). Responses
of trigeminal ganglion neurons to the radial distance of contact during active vibrissal touch. J
Neurophysiol *95*, 791-802.
Wallach, A., Bagdasarian, K., and Ahissar, E. (2016). On-going computation of whisking phase by
mechanoreceptors. Nat Neurosci *19*, 487-493.
Welker, W.I. (1964). Analysis of Sniffing of the Albino Rat. Behaviour *22*, 223-244.
Wineski, L.E. (1983). Movements of the cranial vibrissae in the Golden hamster (*Mesocricetus auratus*). J
Zool *200*, 261-280.
Wineski, L.E. (1985). Facial morphology and vibrissal movement in the golden hamster. J Morphol *183*,
199-217.
Yang, H., Kwon, S.E., Severson, K.S., and O'Connor, D.H. (2016). Origins of choice-related activity in
mouse somatosensory cortex. Nat Neurosci *19*, 127-134.
Yu, C., Derdikman, D., Haidarliu, S., and Ahissar, E. (2006). Parallel thalamic pathways for whisking and
touch signals in the rat. PLoS Biol *4*, e124.
Yu, J., Gutnisky, D.A., Hires, S.A., and Svoboda, K. (2016). Layer 4 fast-spiking interneurons filter
thalamocortical signals during active somatosensation. Nat Neurosci *19*, 1647-1657.

Supplemental Video Legends

Supplemental Video 1: Whisker mechanoreceptor activity during whisking.

Raw video (slowed 20-fold) showing mouse whiskers during whisking. The tracked γ whisker is indicated with the black overlay, and its angular position (θ) is shown in the trace at bottom. Spike times from a simultaneously recorded afferent responsive to the C2 whisker (third from right) are indicated as black ticks above the θ trace. Audio is the playback of the spike waveform at the corresponding spike time.

Supplemental Video 2: Whisking is accompanied by widespread motion of the face.
Raw video (slowed 20-fold) showing side-view of a mouse face during whisking. All whiskers except the A1 and A2 whiskers and much of the facial fur have been trimmed, for a better view of the skin. The skin moves in complex patterns during whisking.

Supplemental Video 3: Supraorbital vibrissa movement during whisking.
Raw video (slowed 20-fold) showing mouse whiskers (A1 and A2) and supraorbital (SO) vibrissae during whisking. The A1 whisker (black overlay) and caudal SO vibrissa (blue overlay) are tracked. The angular positions of the whisker (θ_{A1}, black trace) and SO vibrissae (θ_{SO}, blue trace) are shown at bottom. The SO vibrissae whisk in phase with the whiskers.

Supplemental Video 4: Genal vibrissa movement during whisking.
Raw video (slowed 20-fold) showing mouse whiskers (C-row) and genal vibrissa moving in free air. The C1 whisker (black overlay) and genal vibrissa (red overlay) are tracked. The angular positions of the whisker (θ_{C1}, black trace) and genal vibrissa (θ_G, red trace) are shown at bottom. The genal vibrissa whisks in phase with the whiskers.

Supplemental Video 5: Microvibrissa movement during whisking.
Raw video (slowed 20-fold) showing mouse whiskers (D2 and D3) and microvibrissae during whisking. The D3 whisker (black overlay) and one of the larger, more dorsal and caudal microvibrissae (green overlay) are tracked. The angular positions of the whisker (θ_{D3}, black trace) and microvibrissa (θ_μ, green trace) are shown at bottom. The microvibrissa does not whisk.

Supplemental Video 6: Supraorbital vibrissa movement persists after buccal but not temporal/zygomatic nerve cut.
Raw video (slowed 20-fold) showing split-screen view of mouse whiskers (A1 and A2) and supraorbital (SO) vibrissae on the left side, and whiskers (C-row) on the right side, during

whisking. The left A1 whisker (black overlay), left caudal SO vibrissa (blue overlay), and right γ

or C1 whisker (black overlay) are tracked. The angular positions of the whiskers (θ_L and θ_R,

black traces) and SO vibrissae (θ_{SO}, blue trace) are shown at bottom. The first episode shows

that the SO vibrissae whisk in phase with the ipsilateral whiskers prior to nerve cut. The second

episode occurs after cut of the left buccal branch of the facial nerve. Whisking of the left

whiskers is abolished, but the SO vibrissae continue to whisk. The third episode occurs after

subsequent cut of the left facial motor nerve at the junction of the temporal and zygomatic

branches (Supplemental Figure 2G). SO vibrissae whisking is abolished.

Supplemental Video 7: Genal vibrissa movement reduced after buccal nerve cut.

Raw video (slowed 20-fold) showing split-screen view of mouse whiskers (C-row) and genal

vibrissa on the left side, and whiskers (C-row) on the right side during whisking. The left γ

whisker (black overlay), left genal vibrissa (red overlay), and right γ whisker are tracked. The

angular positions of the whiskers (θ_L and θ_R, black traces) and genal vibrissa (θ_G, red trace) are

shown at bottom. The first episode shows that the genal vibrissa moves in phase with the

ipsilateral whiskers prior to nerve cut. The second episode occurs after cut of the left buccal

branch of the facial nerve. Whisking of the left whiskers is abolished, and the genal vibrissa

motion is greatly reduced. Whisking of the supraorbital vibrissae can be seen in the background.

Source Data Files

Figure 2–Source Data 1. MATLAB R2016b "table" data structure with Normalized MI values

shown in Figure 2E.

Figure 3–Source Data 1. MATLAB R2016b "table" data structure with MI and Normalized MI

values shown in Figure 3E,F.

Figure 4–Source Data 1. MATLAB R2016b "table" data structure with cross-correlation and

mean amplitude values shown in Figure 4E,F.

Figure 4–Source Data 2. MATLAB R2016b "table" data structure with mean amplitude values

before and after nerve cuts shown in Figure 4G.

Figure 4–Source Data 3. MATLAB R2016b "table" data structure with MI and Normalized MI

values shown in Figure 4J,K.

Figure 5–Source Data 1. MATLAB R2016b "table" data structure with MI values and

confidence intervals shown in Figure 5E.

Figure 5–Source Data 2. MATLAB R2016b "table" data structure with MI values shown in
Figure 5G.

Figure 5–Source Data 3. MATLAB R2016b "table" data structure with Normalized MI values
shown in Figure 5H.

Figure 6–Source Data 1. MATLAB R2016b "table" data structure with MI and Normalized MI
values shown in Figure 6A,C.

Figure 6–Source Data 2. MATLAB R2016b "table" data structure with MI values shown in
Figure 6B.

Figure 6–Source Data 3. MATLAB R2016b "table" data structure with MI values shown in
Figure 6D.

Materials and Methods

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers
Software, algorithm	WaveSurfer	HHMI Janelia Research Campus	http://wavesurfer.janelia.org/
Software, algorithm	StreamPix 5 or 7	Norpix	RRID:SCR_015773
Software, algorithm	Janelia Whisker Tracker	Clack et al. 2012	N/A
Software, algorithm	MATLAB 2014a, 2016b, or 2018a	MathWorks	RRID: SCR_001622
Software, algorithm	Adobe Illustrator	Adobe	RRID: SCR_010279
Other	High speed CMOS camera	PhotonFocus	DR1-D1312-200-G2-8

Other	Telecentric lens	Edmund Optics	Cat #: 55-349
Other	Tungsten microelectrode 2 MΩ	World Precision Instruments	Cat #: TM33A20
Other	Tetrode microdrive (custom-built)	Cohen et al. 2012	N/A
Other	Fine 0.3 mm tip water-based marker	Platinum Art Supplies	Cat #: B01FWIE032
Other	Suture thread 8/0	Fine Science Tools	Cat #: 12051-08

670 All procedures were in accordance with protocols approved by the Johns Hopkins University
671 Animal Care and Use Committee. Sample sizes were not predetermined.
672
673 **Mice.** Mixed background mice were housed singly in a vivarium with reverse light-dark cycle (12
674 hours each phase). Behavior experiments were conducted during the dark (active) cycle. The
675 sex and line of each mouse used for recordings is detailed in Supplemental Table S1.
676
Surgical preparation – TG recordings. Adult mice (6-18 weeks old) were implanted with
titanium headcaps (Yang et al., 2016). Prior to electrophysiological recordings, two small
openings (0.5 mm anterior-posterior, 2 mm medial-lateral) in the skull were made centered at 0
and 1.0 mm anterior and 1.5 mm lateral to Bregma, with dura left intact. Craniotomies were
covered acutely with hemostatic gelatin sponge (VetSpon, Ferrosan Medical Devices) or
chronically with silicone elastomer (Kwik-Cast, WPI) followed by a layer of dental acrylic (Jet
Repair Acrylic).
684
Surgical preparation – MeV recordings. Custom microdrives with eight tetrodes (Cohen et al.,
2012) were built to make extracellular recordings from MeV neurons. Each tetrode comprised
four recording wires (100-300 kΩ). A ~1 mm diameter craniotomy was made (centered at -5.4
mm caudal to bregma, 0.9 mm lateral to midline) for implanting the microdrive to a depth of 2
mm, ~0.5 mm dorsal to MeV. Adult mice (9-18 weeks old) were implanted with a titanium
headcap for head-fixation. The microdrive was advanced in steps of ~100 μm each day until
reaching MeV, identified by the presence of clear high-frequency firing responses to jaw
opening and/or closing. Putative MeV jaw proprioceptors were identified post hoc by clear
modulations of spike rate aligned to lick times (Figure 5C).
694
Behavioral training and apparatus. Mice received 1 ml water per day for ≥ 7 days prior to
training. Mice were head-fixed and placed on a linear treadmill to promote whisking, as mice
whisk during running. Voluntary bouts of running were encouraged by providing subsequent
water rewards via a custom lickport. On training days (2-10 days total), mice were weighed

before and after each session to determine the volume of water consumed. If mice consumed <
1 ml, additional water was given to achieve 1 ml total. During recordings, treadmill position was
tracked with a custom optical rotary encoder comprised of a 3D printed encoder disk (2 cm
diameter, 20 holes) and a commercial photointerrupter (1A51HR, Sharp).
703
Whisker and other hair trimming. One day prior to electrophysiological recording, non-
mystacial hairs on the left side of the face were trimmed short with fine forceps and
microdissection scissors (Fine Science Tools), during isoflurane (1.5%) anesthesia. For TG
recordings, all whiskers and microvibrissae were trimmed short except β, γ, δ, B1-4, C1-4, and
D1-4. For improved tracking of whiskers, we minimized obstruction of the field of view by hairs
that were not whiskers intended to be tracked. We did not use chemical hair remover. Fur
between the whiskers was manually removed by plucking or trimming. Non-whisker hairs were
maintained at this short length by repeating this procedure as necessary. Receptive fields on
facial hairy skin were always on fur cut <1 mm by trimming. Whisker and non-mystacial vibrissa
afferents were recorded while the vibrissa in the receptive field was at or near its intact length.
714
Trigeminal ganglion electrophysiology. Recordings from TG afferents were performed as
described (Severson et al., 2017). Briefly, awake mice were head-fixed and allowed to run on
the treadmill. The craniotomy was exposed and covered with PBS. A single tungsten recording
electrode (2 MΩ nominal, Parylene coated; WPI) was lowered ~5.5 mm until it reached the TG.
The tissue was allowed to relax at least 10 min to stabilize recordings. An identical reference
electrode was lowered to a similar depth or placed outside the craniotomy in the PBS. The
differential electrophysiological signal between recording and reference electrodes was
amplified 10,000x, bandpass filtered between 300 Hz and 3,000 Hz (DAM80, WPI), and
acquired at 20 kHz in 5 second sweeps. Electrophysiology, high-speed video, and other
measurements were synchronized by Ephus (Suter et al., 2010) or WaveSurfer
(http://wavesurfer.janelia.org) software. A micromanipulator (Sutter Instruments) advanced the
recording electrode until a well-isolated unit responsive to manual touch stimulation was
encountered. The unit's receptive field, response type (RA or SA), and direction selectivity were
manually classified. All whiskers except the row containing the whisker-of-interest and/or
surrogate tracking whisker were trimmed short. Small manual movements of the treadmill
encouraged the mouse to run and whisk. After recordings, the craniotomy was covered with
silicone elastomer and a thin layer of dental acrylic. Spike waveforms were obtained by
thresholding high-pass filtered (500 Hz) traces and clustered using MClust-4.1 or MClust-4.4
(AD Redish et al.). A subset of TG whisker afferent recordings is reanalyzed from a previous
report (Severson et al., 2017), as detailed in Supplemental Table 1. This subset includes 33 that
were selected based on responses to either touch or whisking. Fifteen of these 33 (45%) were
whisking-sensitive, similar to the fraction of whisking-sensitive units reported previously (36%;
(Szwed et al., 2003)). The remaining whisker afferents reported here were selected based on
responses to whisking and therefore cannot be used to estimate the fraction of whisking-
sensitive neurons.
740
MeV electrophysiology. Water was intermittently delivered via a lickport tube placed below the
animal's snout. Lick signals were recorded by a custom electrical circuit designed to detect
when the tongue contacted the lickport. Lick traces and broadband voltage traces from
individual tetrode wires were acquired continuously at 30 kHz (Intan Technologies). Signals
were bandpass filtered online between 0.1 Hz and 10 kHz, highpass filtered offline below 500
Hz, and spikes were detected using a threshold of 4-6 standard deviations of the filtered signal.
The timestamp of the peak of each detected spike, as well as a 1-ms waveform centered at the
peak, were extracted from each channel of the tetrode for spike sorting, and clustered using
MClust (AD Redish et al.).

Mapping facial hairy skin receptive fields. The touch receptive fields of TG units were identified with a hand-held probe, while monitoring activity using an audio monitor (Model 3300, A-M Systems). When a whisker receptive field could not be found, the receptive field could often be located after probing hairy skin on the entire face. In these cases, before recording began, the extent of the receptive field was mapped by determining the region of hair and skin in which gentle touch with fine forceps (Dumont AA, tip dimensions 0.4 mm x 0.2 mm; FST, #11210-10) evoked spikes and marked with a fine, water-based color marker (0.3 mm tip, Micro-Line, Platinum Art Supplies). Following the recording, the mouse's head with marked receptive fields and a micro-ruler (Electron Microscopy Sciences, #62096-08) were photographed (13 megapixel camera, LG Stylo 2) from the side, above, and/or below. The receptive fields were then compiled on a template "face map". The template image was drawn by outlining the profile and fiducial marks (e.g. eye, whisker follicles, nostrils) of a side view image of a mouse's face in Adobe Illustrator CS 6 (Adobe Systems). The approximate shape, location, and relative size of each imaged receptive field were mapped onto the template by: outlining the receptive field, locating nearby fiducial marks in the original image, applying a fixed scaling to match receptive field and template image dimensions, and translating to align to fiducial marks in the template image. Using the SVG Interactivity Panel in Illustrator, receptive fields were tagged with unique identifier text and their coordinates exported to a text file subsequently read into MATLAB. Borders of each zone of the face (e.g. pad, cheek) were drawn by outlining and connecting fiducial marks (Supplemental Figure 1). Receptive fields were designated to the zone in which the center of mass was located.

High-speed videography. Video frames (640 pixels x 480 pixels, 32 µm/pixel) were acquired at 500 Hz using a PhotonFocus DR1-D1312-200-G2-8 camera (90 µs exposure time) and Streampix 5 software (Norpix). Light from a 940 nm LED (Roithner Laser) was passed through a condenser lens (Thorlabs), through the whisker field, reflected off a mirror (Thorlabs), and directed into a 0.25X telecentric lens (Edmund Optics). Ephus or WaveSurfer triggered individual camera frames (5 seconds, 2,500 frames per sweep) synchronized with electrophysiological recordings. To record microvibrissa movement, whiskers were trimmed, except for the D-row whiskers used for tracking whisker movement. The LED was rotated 30° to capture an oblique view of the profile of the mouse's face, thus maximizing the apparent length of the microvibrissae to enable tracking. To record facial and supraorbital vibrissa movements, the mouse's fur was trimmed to <1 mm, as described above. Whiskers and microvibrissae were trimmed to the base, except for the A-row whiskers used for tracking whisker movement. An additional mirror was placed in the light path to capture a side view of the mouse's face.

Video analysis. All whisker tracking was computing using the Janelia Whisker Tracker (Clack et al., 2012). X-Y coordinates of the whisker objects for each frame were computed by tracing the backbone of each whisker at subpixel-resolution. To reduce noise in measurement of θ, we truncated the tracked whisker trace at its intersection with that frame's "facemask", a curve offset from an outline of the face profile. The facemask was drawn for each frame, briefly, by fitting a smoothing spline to the contour of the face and performing several other image processing steps in MATLAB (Severson et al., 2017). The whisker's follicle location was then estimated by extrapolating past the facemask along the angle of the whisker base (Pammer et al., 2013; Severson et al., 2017). A simple "linking" algorithm was used to ensure the same whisker was tracked across frames. Traced objects outside of the expected region of interest, e.g. whisker pad, and outside of the expected length range were excluded. Whisker identity was then determined based on its follicle X-coordinate in either ascending or descending order.

799 Finally, a number of events could render individual videos ineligible for further processing.
800 These events included objects placed in or entering the video frame or grooming behavior.
801 Using a custom GUI, every sweep was inspected to determine if an exclusion event had
802 occurred.
803
Processing kinematics. We used the Hilbert transform to quantify the instantaneous phase
(Φ), amplitude (θ_{amp}) and midpoint (θ_{mid}) of bandpass (8-30 Hz, Butterworth) filtered θ (Hill et al.,
2011). Instantaneous whisking frequency (f_{whisk}) was calculated by taking the time derivative of
the unwrapped Φ signal. We first smoothed θ with a Savitzky-Golay filter (3rd order, span of 9
frames) and interpolated missing frames when possible. Angular velocity, θ', the time derivative
of θ, was calculated using central differences and smoothed with the same Savitzky-Golay filter.
Sweeps with more than 2% of frames having missing θ data were excluded. For θ, θ', θ_{amp}, f_{whisk},
θ_{mid}, observations outside of the 0.25 and 99.75 percentiles were excluded. No outlier removal
was performed on Φ. We calculated cross-correlation values (MATLAB 'xcorr' with 'coeff'
option) on pairs of traces for whiskers and non-mystacial vibrissae (Figure 4E) after converting
the sampling intervals from equally spaced time intervals to equally spaced phase intervals,
using linear interpolation separately for each whisk cycle. Whisk cycles containing any non-
whisking frames were removed. For cross-correlation analysis, we included between 79 and
333 videos for each session, including 195-591 seconds of whisking data.
818
Tracking facial movement. We acquired epochs of facial movement with high-speed video
(500 Hz, 480 pixels x 640 pixels, 32 μm/pixel) to analyze correlations between facial skin
movement and whisker kinematics (Figure S1). Two mirrors were placed in the light path to
capture a side view of the mouse's face. Facial hair and whiskers were trimmed short except
two A-row whiskers for tracking whisker movement. Displacement of each pixel for each frame
was estimated by applying an image registration algorithm (MATLAB "imregdemons" with
pyramid level iterations 32, 16, 8, and 4) that aligns each "moving" frame with a "fixed" template
frame. First, fixed and moving frames were resized by half on each dimension (to 320 pixels x
240 pixels; MATLAB "imresize" with bicubic smoothing) to reduce compute time and file size.
Next, pixel values outside of the face and in the eye were set to zero. Image registration was
then applied to every video frame in the session. We then calculated Pearson's correlation
coefficients between the time series of x-dimension pixel displacement values (Δx) and whisker
position (θ) time series. Y-displacement values were not used for calculating correlations
because they could not be estimated as accurately from 2D images, due to substantial out-of-
image-plane curvature of the mouse face that varies along the y-dimension. Mean Pearson's r
values for each facial region (Supplemental Figure 1E) were obtained by averaging r values
across all pixels within each facial region. These regions were determined for the fixed template
image using fiducial marks as described above.
837
Facial nerve lesion experiments. Mice were injected preoperatively with ketoprofen (5 mg/ml,
s.c., Ketofen) to reduce inflammation and lidocaine (5%, s.c., Vedco) to anesthetize the area
around the incision. A small incision was made 3 mm caudal to whisker delta on the left side. To
abolish movement of the mystacial pad, the buccal branch of the facial nerve was cut
(Supplemental Figure 2G; (Dorfl, 1985; Fee et al., 1997)) using microdissection scissors (Fine
Science Tools). To further assay control of movement of the supraorbital vibrissae, a small
incision was made dorsal and caudal to the left eyelid. The facial nerve was cut at the junction
of the temporal and zygomatic branches. Following nerve cut, the incision was closed with fine
suture thread (8/0, Fine Science Tools #12051-08) and coated with antibiotic ointment (Pac-Kit).
Mice were injected postoperatively with buprenorphine HCL (0.1 mg/ml, s.c., Par
Pharmaceutical). Behavioral recordings were conducted after at least one day of postoperative
recovery. Whisking was recorded before and after surgical intervention using high-speed video

(500 Hz; 90 or 250 μs exposure time for genal and supraorbital recordings, respectively) to
assess the effect of motor nerve lesions on non-mystacial vibrissa movements. Whisking
periods (defined in Glossary) were determined by tracking a whisker on the right, unaffected,
whisker pad. Movements of mystacial and non-mystacial vibrissae on the left side and mystacial
whiskers on the right side were recorded simultaneously with a single camera using mirrors to
split the image. Nerve cut experiments were performed on 3 mice, with 15 total videographic
recording sessions: for each mouse, first 2 sessions under control conditions, then 2 sessions
after buccal branch cut, then 1 session after the temporal/zygomatic cut.

Data analysis – tuning curves. To calculate tuning curves, kinematic variables were
processed by performing outlier removal, restricting observations to whisking periods, and
binning into 30 equally spaced bins, unless otherwise noted. Bins with fewer than 25
observations were set to NaN.

Data analysis – mutual information. Mutual information (MI) was calculated between the
distributions of spike counts and kinematic variable values across individual 2 ms frames. The
distribution of spike counts was $P(X = x), x \in \{0,1,2,...n\}$, where n is the maximum number of
spikes observed during a single 2 ms frame across the duration of the recording. For the
recordings presented here, n was ≤ 3. For each kinematic variable, Y, the distribution $P(Y)$ was
estimated after binning Y into 16 equally spaced bins ranging from max(Y) to min(Y) after
removing outliers as described above. Uniform count binning of kinematic variables yielded
similar results for θ_{mid} and Φ (Supplemental Figure 3). The joint distribution $P(X = x, Y = y)$ was
estimated similarly. MI (Cover and Thomas, 2006) was then computed as:

$$MI(X;Y) = \sum_{x \in X, y \in Y} P(x,y) log_2 \frac{P(x,y)}{P(x)P(y)}$$

To obtain the "MI rate", we multiplied MI by the sampling frequency, which was 500 Hz for the 2
ms time bins.

Confidence intervals on the MI values for each unit (Figure 5E, Supplemental Figure 4) were
obtained by bootstrap after resampling frames with replacement and recalculating MI for 1000
iterations. MI values and confidence intervals that would be obtained by chance under the null
hypothesis of no true correlation ("Shuffled" data in Figure 5E and Supplemental Figure 4) were
obtained after shuffling spike counts with respect to either θ_{mid} or Φ, via sampling without
replacement, for 1000 iterations.

MI values for Φ were not strongly affected by our use of a 2 ms time bin, showing stable values
at bin sizes from 1-8 ms, then declining with larger bin sizes (Supplemental Figure 5A-B). As
expected due to the relatively slow variation in θ_{mid}, MI values for θ_{mid} increased with bin size up
to the largest bin size tested (64 ms; Supplemental Figure 5D-E). Our goal was not to determine
the maximum MI values that could be obtained based on each kinematic variable, but rather to
compare identically calculated MI values across different classes of facial mechanoreceptors.
The relative abilities of different mechanoreceptor classes to encode Φ and θ_{mid} were similar
across choices of window sizes (Supplemental Figure 5C,F). Therefore, for simplicity, we used
the 2 ms bin size that corresponded to an individual video frame for all kinematic variables.

Similarly, while MI values could be maximized for each unit by shifting the window for spike
count relative to that for kinematic variables, the delay at which peak MI occurred was
heterogenous across units of the same mechanoreceptor class (Supplemental Figure 5G), and
for simplicity we therefore chose to use temporally aligned (simultaneous) windows. However,

the exposure time for each video frame occurred in the first 90 µs of the 2 ms frame period, whereas spike counts were measured over the full 2 ms frame period. There was thus an effective delay between the mean windows used for kinematic variable and spike count measurements of ~955 µs.

To calculate "normalized MI" for each recording, we first calculated the entropy of the spike count distribution:

$$H_X = -\sum_{x \in X} P(x) \log_2 P(x).$$

Normalized MI was then computed as:

$$Normalized\ MI = \frac{MI(X;Y)}{H_X}$$

Glossary:

"Whiskers": macrovibrissae located on the mystacial pad.

"Non-mystacial vibrissae": vibrissae that are not whiskers; includes supraorbital and genal macrovibrissae, and the microvibrissae.

"Whisking" periods: Frames with θ_{amp} > 2.5° and f_{whisk} > 1 Hz for the tracked whisker.

"Non-whisking" periods: Frames with θ_{amp} < 1° for the tracked whisker.

"Whisking-sensitive": Applies to a unit with 95% confidence interval (CI) on mean spike rate during whisking in air non-overlapping with 95% CI for mean spike rate during non-whisking and with mean spike rate > 1 Hz during whisking.

"Whisker afferents" or **"whisker mechanoreceptors"**: LTMRs with single-whisker receptive fields, presumably which innervate the whisker follicle.

"Proprioceptors": Mechanoreceptors presumed to associate with muscle spindle or Golgi tendon organ structures.

Figure 1. Recording whisking and facial mechanosensory afferent spiking.
(**A**) Schematic of types of afferents (open circles with dotted lines) recorded, grouped by type of receptive field: trigeminal ganglion (TG, beige) low threshold mechanoreceptors (LTMRs) with receptive fields localized to (1) a mystacial whisker follicle (filled black dots; e.g. black whisker), (2) hairy skin (e.g. gray patch on cheek), or (3) a non-mystacial vibrissa (red dots; e.g. black supraorbital vibrissa), and trigeminal mesencephalic nucleus (MeV) proprioceptors innervating facial muscles. (**B**) Schematic of experimental setup. A head-fixed mouse ran on a treadmill and whisked in air. Single neurons were recorded simultaneously with high-speed (500 Hz) video of the whiskers. (**C**) Example video frame, capturing the silhouette of the whiskers and profile of the mouse face, and illustrating measurement of the angular position of a whisker (θ) relative to the mediolateral axis. (**D**) Example traces showing one second of whisker angular position (θ), whisking midpoint (θ_{mid}), amplitude (θ_{amp}; bottom of scale bar indicates 0°), and phase (Φ; gray vertical lines, times when the whisker is fully retracted, $\Phi = \pi$). (**E**) Overview of dataset, including the number of units recorded and the number of each type that were whisking-sensitive (Glossary).

Figure 2. Self-motion responses from mechanoreceptors innervating whisker follicles.
(**A**) Schematic of a unit with whisker receptive field. (**B**) Spike times (black ticks) for two example whisker afferent units, each aligned with whisker position traces (gray lines: fully retracted positions). Unit 1 (top) responded during protracting phases. Unit 2 (bottom) responded during retracting phases. (**C**) Phase tuning curves (mean ± SEM; SEM here and in some subsequent panels narrower than line width) for unit 1 and unit 2. (**D**) Joint probability distributions for spike count and whisk phase (Φ), obtained from 2 ms periods corresponding to individual video frames, for unit 1 (top) and unit 2 (bottom). Mutual information (MI) between spike count and phase for each unit is shown at the top of each panel. Per 2 ms period, unit 1 spiked up to once and unit 2 up to twice. (**E**) Heatmap of normalized mutual information values for all whisking-sensitive whisker mechanoreceptors (n = 42), measured between spike count and each kinematic quantity (•): phase (Φ), position (θ), velocity (θ'), acceleration (θ''), amplitude (θ_{amp}), and midpoint (θ_{mid}). Units (rows) are sorted by increasing normalized MI averaged across the kinematic quantities. A subset of whisker afferent recordings was previously reported (Severson et al., 2017) and is reanalyzed here (see Supplemental Table 1 for details). Data for panel E are given in Figure 2–Source Data 1.

Figure 3. Self-motion responses from mechanoreceptors innervating facial hairy skin.
(**A**) Receptive fields on facial hairy skin (n = 85). Approximate size, shape, and location of receptive fields (gray) were compiled onto a template image of a mouse face (scale bar: 2 mm). Colored receptive fields show examples from whisker pad (green), rostral cheek (orange), and caudal cheek (blue). Whisker follicles and non-mystacial vibrissae (filled black dots) are included as fiducial marks. (**B**) Example one second traces showing spike times (colored ticks) aligned with whisker position, from recordings corresponding to the examples in (A). (**C**) Phase tuning curves (mean ± SEM) for example pad unit (green, top), rostral cheek unit (orange, middle), and caudal cheek unit (blue, bottom). Units with similar mean spike rates during whisking can differ in their phase modulation. (**D**) Mutual information rate between spike count and phase overlaid on outlines of receptive fields (scale bar: 2 mm). Many but not all receptive fields with large MI rates were located near whiskers. (**E**) Cumulative histograms of MI rate between spike count and phase for whisking-sensitive neurons with receptive fields in each region of the face, including whisker pad (n = 13), cheek (n = 18), eye (n = 10), lip (n = 5), snout (n = 9), and jaw (n = 3). (**F**) Heatmap of normalized MI values for all whisking-sensitive facial hairy skin units (n = 58), measured between spike count and each kinematic quantity (•, columns): phase (Φ), position (θ), velocity (θ'), acceleration (θ''), amplitude (θ_{amp}), and midpoint (θ_{mid}). Units (rows) are sorted by receptive field location (labeled at right) and within each face region by increasing normalized MI averaged across the kinematic quantities. Data for panels E, F are given in Figure 3–Source Data 1.

Figure 4. Non-mystacial vibrissae move in phase with whiskers, and their mechanoreceptors encode motion.
(**A**) Schematic of non-mystacial vibrissae included in these experiments: supraorbital (SO, blue), genal (G, red), and microvibrissae (μ, green) on the upper lip. (**B**) Correlated motion between whiskers and non-mystacial vibrissae. Example one second traces showing angular positions of a supraorbital vibrissa (top), genal vibrissa (middle) or microvibrissa (bottom), each with simultaneously tracked whisker angle. (**C**) Scatter plot of phase for whisker vs. non-mystacial vibrissae (n = 1000 random frames; top, SO; middle, G; bottom, μ; dashed lines: unity). (**D**) Example whisk cycles for whisker and non-mystacial vibrissae pairs, normalized based on whisker angle (n = 500 random whisks; top, A1 and SO ; middle, C1 and G; bottom, D2 and μ). (**E**) Peak cross-correlation (Pearson's r) and phase lag (open circles) between angular positions of a tracked whisker and either an adjacent whisker (W, n = 14 whisker pairs from 12 mice), a supraorbital vibrissa (n = 6 recordings from 6 mice), a genal vibrissa (G, red; n = 6 recordings from 6 mice), or microvibrissa (μ, green; n = 3 recordings from 3 mice). (**F**) Mean whisk amplitude (± SD across recordings) for whisker, supraorbital, genal, or microvibrissa. (**G**) Result of unilateral facial nerve lesions on whisk amplitude for whiskers and non-mystacial vibrissae (n = 3 mice). Open circles: mean θ_{amp} for whiskers and supraorbital vibrissa during whisking (determined based on contralateral whiskers) before and after facial nerve cuts, performed sequentially at the buccal and then temporal/zygomatic branches. Closed circles: mean θ_{amp} obtained from separate videographic sessions for whiskers and genal vibrissae. (**H**) Example one second traces showing spike times from SO (top), G (middle), and μ units (bottom), each aligned with position of a tracked whisker (θ). (**I**) Phase tuning curves (mean ± SEM) for the example units in (H): top, SO; middle, G; bottom, μ. (**J**) Histograms of mutual information rate between spike count and phase for whisking-sensitive SO (n = 8), G (n = 3), and μ (n = 8) units. (**K**) Heatmap of normalized mutual information for all whisking-sensitive non-mystacial vibrissae units (n = 19). Conventions as in Figure 3F. Data for panels E, F are given in Figure 4–Source Data 1. Data for panels J, K are given in Figure 4–Source Data 2.

Figure 5. Responses of proprioceptors in the trigeminal mesencephalic nucleus (MeV) during licking and whisking.
(**A**) Schematic of MeV tetrode recordings during licking and whisking. (**B**) Example traces for an MeV unit showing spike times (black ticks) and lick times (blue ticks) aligned with position (θ) of a tracked whisker, for one second periods with (top) and without (bottom) licks. (**C**) Top, spike raster aligned to lick times (n = 1000 random licks) for example unit in (B). Bottom, peri-event time histogram (± SEM) aligned to lick times across all licks. (**D**) Top, spike raster aligned to whisk cycles (n = 1000 random whisks) for unit in (B). Whisks are ordered by mean θ_{mid}. Bottom, phase tuning curve for same unit (mean ± SEM) across all whisk cycles. (**E**) Top, MI rate between spike count and Φ for each unit (± 95% bootstrap CI). Cyan: results of same calculation but after randomly shuffling spike counts with respect to phase. Bottom, same as top but for MI rate between spike count and θ_{mid}. (**F**) Midpoint tuning curve (mean ± SEM) for unit in (B). (**G**) Histograms of MI rate between spike count and θ_{mid} for whisking-sensitive MeV units, separately for those that showed modulation by licking or passive jaw movement (orange, n = 23), and others recorded on the same tetrode (purple, n = 10). (**H**) Heatmap of normalized mutual information for all whisking-sensitive MeV units (n = 33). Conventions as in Figure 2E. Data for panel H are given in Figure 5–Source Data 1.

Figure 6. Coding of self-motion by diverse classes of facial mechanoreceptors.
(**A**) Left, summary cumulative histograms of MI rate between spike count and θ_{mid} for whisking-sensitive MeV (n = 33), face (n = 58), non-mystacial vibrissae (n = 19), and whisker units (n = 42). Right, summary histograms of normalized MI between spike count and θ_{mid} for same units. (**B**) Summary histograms of MI rate between spike count and θ_{mid} for the best encoding subgroups of whisking-sensitive units: putative jaw proprioceptors in MeV (n = 23), pad and cheek units (n = 31), supraorbital and genal units (n = 11), and whisker units (replotted from A). (**C**) Left, same as (A) but for MI rate between spike count and Φ. (**D**) Same as (B) but for the subgroups that best encoded Φ: putative proprioceptors in MeV (n = 23), pad units (n = 13), supraorbital units (n = 8), and whisker units (replotted from C). (**E**) Schematic depicting flow of information about whisking kinematics from various peripheral mechanoreceptors to the brain: whisker follicle (black), supraorbital vibrissa (blue), and whisker pad hairy skin afferents (green). (**B**-**D**) Panels include data from Figures 2-5 plotted together for comparison. Data for panels A and C are given in Figure 6–Source Data 1. Data for panels B and D are given in Figure 6–Source Data 2.

Supplemental Figure 1. Widespread facial movement correlated with whisker motion.
(**A**) Six facial regions on template mouse face were used for categorizing afferent receptive fields. Zones were drawn based on fiducial marks, including the mystacial whiskers (filled circles). Locations of non-mystacial vibrissa follicles (open circles) are also indicated. Receptive fields were scaled to the template image. (**B**) "Fixed" template image for image registration method of tracking facial motion. Overlaid are facial regions drawn using image fiducial marks. Pixels outside the edge of the face and within the eye, outlined in white, were effectively excluded from image registration by setting their values to zero. (**C**) Example "moving" frame overlaid with a subset of the estimated x- and y-displacements, plotted as vectors (red arrows), that would align it to the fixed template image. (**D**) Template images from each of two mice overlaid with color scale showing Pearson's correlation coefficients (r) between the time series of each pixel's x-displacement (Δx) and the A1 whisker position (θ). (**E**) Pearson's r values averaged ($\pm$ SEM) across all pixels within each facial region, separately for two mice. (**B-D**) Scale bars: 2 mm.

Supplemental Figure 2. Non-mystacial vibrissae movement and its dependence on facial nerve innervation.
(**A**) Example traces of simultaneously recorded A1 whisker position (θ_{A1}, top black trace), A2 whisker position (θ_{A2}, bottom black trace) and supraorbital vibrissa position (θ_{SO}, blue; gray lines: fully retracted phase of A1). Whisks by A1 that we not matched by whisks of SO vibrissa (missed whisks) are marked by blue arrows. (**B**) Histogram of ratio of SO vibrissa amplitude ($\theta_{amp,SO}$) over A1 whisker amplitude ($\theta_{amp,A1}$) for whisking periods (330,085 frames) in the example recording in (A). Ratios above 1.2 are not shown on plot for clarity (0.09% of frames). The peak with ratio near zero, indicated by blue arrow, indicates a substantial fraction of missed whisks in this example recording. (**C**) Same as (B) but for A2 vs A1 whiskers. There is no histogram peak indicative of missed whisks (0.16% of values are above the axis limit of 1.2). (**D**) Example traces of θ_{C1} and θ_{C2} (black traces) and θ_G (red). Missed whisks by the genal vibrissa are marked by red arrows. (**E**) Histogram of ratio of genal vibrissa amplitude ($\theta_{amp,G}$) versus C1 whisker amplitude ($\theta_{amp,C1}$) for whisking periods (96,429 frames) in the example recording in (C). Ratios above 1.2 not shown (0.36% of frames). Missed whisks are indicated by red arrow. (**F**) Same as (E) but for C2 vs C1 whiskers (0.75% of values above the axis limit). (**G**) Diagram of facial nerve innervation of whisking muscles in mouse, adapted from Dörfl, 1985. Motor nerve lesions were performed by cutting the facial nerve (*FN*) sequentially at the buccolabialis branch (*b*), which innervates the mystacial pad whisking muscles, followed by cutting at the junction of the temporalis (*t*) and zygomatico-orbitalis (*z*) branches (cuts approximately at dashed lines). Whisker and non-mystacial vibrissa movement was measured before and after each cut. (**H**) Left, example frame capturing movement of left side SO vibrissae, left side A-row whiskers, and right side C-row whiskers. Right, example frame capturing movement of left side G vibrissa, left side C-row whiskers, and right side C-row whiskers. (**I**) Left, example 1.5 second trace of SO angle (θ_{SO}, blue), left whisker (θ_L, black, middle), and right whisker (θ_R, black, top) before nerve cut. Middle, same as left, but after left buccolabialis nerve cut. Left whisker movement was abolished, but SO movement persisted. Right, same as middle, but after left temporalis/zygomatico-orbitalis nerve cut. SO movement was abolished. (**J**) Left, same as (I), but for G angle (θ_{SO}, red) before nerve cut. Right, same as left, but after buccolabialis cut. G movement was mostly abolished. (**K**) Summary of nerve cut results, reproduced for convenience from Figure 4G.

Supplemental Figure 3. Alternative binning methods for mutual information calculation.
(**A**) Joint distribution for spike count and Φ comparing binning of Φ with linear spacing (LS, left, same calculation reported throughout paper) and uniform count (UC, right) for an example unit. Marginal probability distributions are plotted for Φ (top) and spike count (right). Note that the LS distribution of Φ is nearly uniform, except fewer bins are observed during retraction phases due to more rapid whisker retraction. MI rate values calculated using each method of binning are reported at top. (**B**) Same as (A), but for spike count and θ_{mid} comparing LS (left) and UC (right) for an example whisker afferent. (**C**) Cumulative histograms of MI rate between spike count and Φ, calculated using LS (solid lines) or UC (dotted lines) for the different afferent groups. (**D**) Same as (A) but for MI rate between spike count and θ_{mid}. (**E**) Same as (C) but for best subgroups. (**F**) Same as (D) but for best subgroups. (**C-F**) LS values are repeated from Figure 6 to allow comparison with UC values. For Φ and θ_{mid}, MI values calculated with LS or UC binning are almost identical.

Supplemental Figure 4. Confidence intervals for mutual information and dependence on sample size.
(**A**) MI rate (± 95% bootstrap CI) between spike count and Φ for all whisking-sensitive units (n = 152). Cyan: results of same calculation but after after randomly shuffling spike counts with respect to Φ. Neurons are sorted by group label (right) and MI rate. (**B**) Mean number of frames in each bin (n = 16 bins) for marginal distribution of Φ, plotted against the width of the 95% CI for MI rate for Φ. Cyan: results of same calculation after randomly shuffling spike times with respect to Φ. (**C**) Cumulative histogram showing mean number of frames per bin for all whisking-sensitive units (n = 152). (**D**) Same as (A), but with MI rate calculated between spike count and θ_{mid}. (**E**) Same as (B), but for θ_{mid}. (**A**,**D**) MeV data are repeated from Figure 5E.

Supplemental Figure 5. Mutual information calculated using varying windows for spike count.
(**A**) MI values for spike count and Φ calculated using different spike count window durations, ranging from 1 to 64 ms for all whisking-sensitive units (rows). Values are normalized from min to max MI across time windows for each unit. Rows are sorted by afferent group (labels on right) and within group by index of max. (**B**) Mean MI values (± SEM) for spike count and Φ, for each afferent group and spike count window, after normalizing each unit as in (A). (**C**) Summary cumulative histograms of MI rate between spike count and Φ for all whisking-sensitive MeV, face, non-mystacial vibrissae, and whisker units, for 2 ms (solid lines, taken from Figure 6C) and 8 ms windows (dashed lines). (**D-F**), Same as A-C but for θ_{mid}. Data depicting 2 ms windows (solid lines) in panels C and F are taken from Figure 6. (**G**) MI rates between spike count and Φ, calculated for varying spike time shifts relative to Φ, in 0.5 ms increments for all whisking-sensitive units (rows). Conventions as in (A). (**H**) Same as (B) but for spike time shifts.

Mouse ID	Sex	Mouse Line	Birth Date	Recording Dates	Receptive Fields/ Regions of Interest	Unit IDs *Prev. incl. in Severson, et al. 2017 ‡ Whisking-Sensitive	Fig. 1	Fig. 2	Fig. 3	Fig. 4	Fig. 5	Fig. 6	Fig. S1	Fig. S2	Fig. S3	Fig. S4	Fig. S5	Other Notes
Recordings of trigeminal ganglion neurons																		
KSt66	M	$TrKC^{CreER};Rosa^{Ai32}$	05/09/14	07/11-07/18/14	whisker	125B*, 128D*‡	E						A-D		C-F	A-E	A-H	
KSt69	M	$TrKC^{CreER};Rosa^{Ai32}$	05/09/14	07/28/14	whisker	132A*‡	E						A-D		C-F	A-E	A-H	
KSt76	M	$TrKC^{CreER};Rosa^{Ai32}$	06/29/14	09/27/14	whisker	145B*‡	E						A-D		C-F	A-E	A-H	
KSt80	F	$TrKC^{CreER};Rosa^{Ai32}$	06/29/14	10/25/14	whisker	154B*												
KSt77	M	$TrKC^{CreER};Rosa^{Ai32}$	06/29/14	11/10-11/12/14	whisker	161A*‡, 163B*	E						A-D		C-F	A-E	A-H	
KSt82	M	$TrKC^{CreER};Rosa^{Ai32}$	07/20/14	12/08/14	whisker	167A*‡	E						A-D		C-F	A-E	A-H	
KSt88	M	$TrKC^{CreER};Rosa^{Ai32}$	10/28/14	03/20-03/30/15	whisker; micro	194A*; 197C‡			J,K				A,C		C,D	A-E	A-H	
KSt89	F	$TrKC^{CreER};Rosa^{Ai32}$	10/25/14	03/31/15	whisker	198B*												
KSt92	M	$TrKC^{CreER};Rosa^{Ai32}$	10/28/14	04/01-04/05/15	whisker; cheek	202A*; 199A‡		A,D-F					A-C		C,D,F	A-E	A-H	
KSt91	M	$TrKC^{CreER};Rosa^{Ai32}$	10/28/14	04/13/15	whisker	207A*												
KSt96	F	$TrKC^{CreER};Rosa^{Ai32}$	01/28/15	04/20/15	whisker	211A*												
KSt94	F	$TrKC^{CreER};Rosa^{Ai32}$	10/28/14	04/23-04/24/15	whisker; eye	214A*; 213A‡		A,D-F					A,C		C,D	A-E	A-H	
KSt97	M	$CCK^{Cre};Rosa^{Ai32}$	01/03/15	04/27/15	whisker	216D*												
KSt95	M	$TrKC^{CreER};Rosa^{Ai32}$	10/25/14	04/29/15	whisker; SO	217F*‡; 217A‡	E		J,K				A-D		C-F	A-E	A-H	
KSt99	M	$TrKC^{CreER};Rosa^{Ai32}$	02/07/15	05/06/15	whisker	220B*‡, 220C*	E						A-D		C-F	A-E	A-H	
KSt106	F	$TrKC^{CreER};Rosa^{Ai32}$	02/07/15	05/08/15	whisker	221B*‡, 221C*	E						A-D		C-F	A-E	A-H	
KSt101	F	$TrKC^{CreER};Rosa^{Ai32}$	02/23/15	05/11/15	whisker	223B*												
KSt107	F	$TrKC^{CreER};Rosa^{Ai32}$	02/07/15	05/14-05/15/15	whisker	225D*‡, 227A*‡, 227C*	E						A-D		C-F	A-E	A-H	
KSt100	M	$TrKC^{CreER};Rosa^{Ai32}$	10/25/14	05/18/15	cheek	228A‡		A,D-F					A-C		C,D,F	A-E	A-H	
KSt109	F	$TrKC^{CreER};Rosa^{Ai32}$	02/07/15	05/21/15	whisker	231A*‡	E						A-D		C-F	A-E	A-H	
KSt111	M	$TrKC^{CreER};Rosa^{Ai32}$	02/07/15	05/29/15	whisker	234A*												
KSt112	M	$TrKC^{CreER};Rosa^{Ai32}$	02/07/15	06/06-06/10/15	whisker; micro; G	240A*‡, 243B*; 242B‡; 242E‡	E		J,K				A-D		C-F	A-E	A-H	
KSt115	F	$TrKC^{CreER};Rosa^{Ai32}$	02/07/15	06/19-06/22/15	whisker; SO	250A*‡; 247A‡	E		J,K				A-D		C-F	A-E	A-H	
KSt117	M	$CCK^{Cre};Rosa^{Ai32}$	01/03/15	06/29/15	whisker	251B*‡	E						A-C		C-F	A-E	A-H	
KSt119	M	$TrKC^{CreER};Rosa^{Ai32}$	03/25/15	07/02-07/06/15	whisker; cheek; snout	253A*‡, 254C*, 256B*‡; 254A‡; 255B‡	E	A,D-F					A-C		C-F	A-E	A-H	
KSt122	M	$TrKC^{CreER};Rosa^{Ai32}$	03/25/15	07/17-07/19/15	whisker; cheek	265A*; 263A‡		A,D-F					A-C		C,D,F	A-E	A-H	
KSt124	F	$TrKC^{CreER};Rosa^{Ai32}$	03/25/15	07/21/15	whisker; cheek	266B*; 266A‡		A,D-F					A-C		C,D,F	A-E	A-H	
KSt123	F	$TrKC^{CreER};Rosa^{Ai32}$	03/25/15	07/25-07/29/15	whisker; cheek; snout; eye	271A*; 268A‡; 270A‡; 270B‡		A,D-F					A-C		C,D,F	A-E	A-H	
KSt125	F	$TrKC^{CreER};Rosa^{Ai32}$	03/25/15	08/04/15	eye	274A‡		A,D-F					A-C		C,D	A-E	A-H	
KSm31	M	$TrKC^{CreER};Rosa^{Ai32}$	05/10/16	08/20-08/22/16	whisker; eye; pad; SO	275A, 276A‡; 276B‡; 277C		A,D-F					A-D		C-F	A-E	A-H	
KSm30	M	$Rosa^{Ai32}$	05/10/16	08/25-08/26/16	whisker; cheek	278B*‡; 279A‡	E	A,D-F					A-D		C-F	A-E	A-H	
KSt126	M	$TrKC^{CreER};Rosa^{Ai32}$	05/10/16	08/29-08/31/16	whisker; SO	282A*; 280B‡			J,K				A-D		C-F	A-E	A-H	
KSt127	F	$TrKC^{CreER};Rosa^{Ai32}$	05/10/16	09/07-09/09/16	whisker	285A‡, 286A*‡	E						A-D		C-F	A-E	A-H	
KSt128	M	$TrKC^{CreER};Rosa^{Ai32}$	01/20/16	09/13/16	whisker; pad	288C*‡; 288B‡	E	A,D-F					A-D		C-F	A-E	A-H	
KSt130	M	$TrKC^{CreER};Rosa^{Ai32}$	09/27/15	09/23/16	whisker; cheek	295A*‡; 295B	E	A,D					A-D		C-F	A-E	A-H	
KSt131	M	$TrKC^{CreER};Rosa^{Ai32}$	09/27/15	09/23/16	whisker	295C*‡	E						A-D		C-F	A-E	A-H	
KSt137	M	$Rosa^{Ai32}$	05/10/16	10/15/16	whisker	303A*‡, 304B*‡, 305B*‡	E						A-D		C-F	A-E	A-H	
KSt135	M	$TrKC^{CreER};Rosa^{Ai32}$	05/10/16	10/24-10/27/16	whisker; snout; pad	309B*‡, 311C*‡; 309A; 311A‡	E	A,D-F					A-D		C-F	A-E	A-H	
KSt138	M	$TrKC^{CreER};Rosa^{Ai32}$	05/10/16	11/20/16	whisker	313A*‡	E						A-D		C-F	A-E	A-H	
KSt140	M	$TrKC^{CreER};Rosa^{Ai32}$	05/10/16	12/01-12/05/16	whisker	318A*‡, 321A*‡, 321B*‡	E						A-D		C-F	A-E	A-H	
KSt143	M	$TrKC^{CreER};Rosa^{Ai32}$	07/04/16	02/17-02/22/17	whisker; snout; eye; pad	332A; 329B, 333B‡; 331A; 332C‡		A,D-F					A-D		C-F	A-E	A-H	
KSt145	M	$CCK^{Cre};Rosa^{Ai32}$	07/04/16	02/24-02/28/17	eye; G; micro	335A‡, 338A; 336A‡; 338B‡		A,D-F	H-K				A-C		C,D,F	A-E	A-H	338B example micro unit in Fig. 3
KSt148	F	$TrKC^{CreER};Rosa^{Ai32}$	10/24/16	03/02-03/05/17	cheek; eye; pad; micro	341A, 342A; 341B; 341C‡; 343A		A,D-F					A-D		C-F	A-E	A-H	
KSt147	M	$TrKC^{CreER};Rosa^{Ai32}$	10/24/16	03/07-03/08/17	whisker; eye; micro	345B‡; 345A‡, 346A; 346B‡	E	A,D-F	J,K				A-D		C-F	A-E	A-H	
KSt150	M	$TrKC^{CreER};Rosa^{Ai32}$	10/24/16	03/16-03/27/17	whisker; micro; SO; eye; cheek; lip; pad; snout	350C‡; 349A‡, 350D‡; 349B‡; 350A, 350B‡, 352C, 356A‡; 351C‡, 352A‡; 351E‡, 352B‡, 354C‡, 355A‡; 353B‡, 354A, 354B	E	A,D-F	H-K				A-D		C-F	A-E	A-H	349B example SO unit in Fig. 3
KSt149	F	$TrKC^{CreER};Rosa^{Ai32}$	10/24/16	03/30/17	SO	358A‡			J,K				A-D		C-F	A-E	A-H	
KSt151	M	$TrKC^{CreER};Rosa^{Ai32}$	10/24/16	04/17-04/18/17	micro; jaw; cheek	359A‡; 360A, 363A; 361A, 361E‡; 363B‡		A-F	J,K				A-C		C,D,F	A-E	A-H	361E example caudal cheek unit in Fig. 3
KSt152	M	$TrKC^{CreER};Rosa^{Ai32}$	10/24/16	04/25-04/28/17	lip; cheek; snout	364A‡; 365A‡; 364B; 367A		A,D-F					A-C		C,D	A-E	A-H	
KSt155	M	$TrKC^{CreER};Rosa^{Ai32}$	12/07/16	07/10-07/17/17	whisker; cheek; jaw; micro; SO; pad	375B‡, 377A‡; 372A; 373A, 373B; 374B; 375C, 376B‡; 376A	E	A,D	J,K				A-D		C-F	A-E	A-H	
KSt154	M	$TrKC^{CreER};Rosa^{Ai32}$	12/07/16	07/20-07/26/17	whisker; jaw	379C, 380A‡; 383A‡	E	A,D-F					A-D		C-F	A-E	A-H	
KSt157	F	$TrKC^{CreER};Rosa^{Ai32}$	12/07/16	08/07-08/12/17	whisker; eye; cheek; jaw	387A, 387C‡; 386A‡; 387B‡; 390B; 390A, 391A	B-E	A,D-F					A-D		A-F	A-E	A-H	387C example unit 2 in Fig. 2 and example in S3
KSt158	M	$TrKC^{CreER};Rosa^{Ai32}$	12/07/16	08/16-08/29/17	whisker; jaw; cheek	402A‡; 394A‡, 401A, 401B‡; 394B‡, 395A‡; 397A‡	E	A-F					A-D		C-F	A-E	A-H	394B example rostral cheek unit in Fig. 3
KSt161	M	$CCK^{Cre};Rosa^{Ai32}$	07/04/16	09/22-09/25/17	whisker; eye; pad	405A‡, 406C‡; 406B‡; 406A‡	E	A,D-F					A-D		C-F	A-E	A-H	
KSt163	M	$TrKC^{CreER};Rosa^{Ai32}$	08/08/17	01/12-01/16/18	SO; snout; pad; lip	408A, 409A, 410B‡; 409B‡, 410A, 411C; 409C‡, 411B‡; 411A‡		A,D-F	J,K				A-D		C-F	A-E	A-H	
KSt164	M	$Rosa^{Ai32}$	08/08/17	01/29-02/07/18	eye; snout; SO; cheek; pad	415A‡, 420A‡; 416A‡, 420B‡; 416B; 417B‡; 418A‡		A-F					A-D		C-F	A-E	A-H	418A example pad unit in Fig. 3
KSt166	M	$TrKC^{CreER};Rosa^{Ai32}$	08/09/17	02/08-02/16/18	whisker; cheek; snout; SO; G	422C‡, 422D‡; 421B‡, 424A‡; 421C‡, 421D‡; 422A, 422B‡, 424B, 425A‡, 425D; 425C	C,D	B-E	A,D-F	J,K			A-D		C-F	A-E	A-H	422D example unit 2 in Fig. 2
KSt165	F	$TrKC^{CreER};Rosa^{Ai32}$	08/09/17	02/20-02/21/18	G	427A, 428B												
KSt167	F	$TrKC^{CreER};Rosa^{Ai32}$	08/09/17	02/24/18	G	430A‡, 430B, 430C			H-K				A-C		C,D,F	A-E	A-H	430A example G unit in Fig. 3
Recordings of trigeminal mesencephalic nucleus neurons																		
HYLC44	M	$TH^{Cre};Rosa^{Ai32}$	12/07/15	05/24-05/30/16	MeV	44B-2‡,3‡,4‡; 44C-15‡,16‡						B-H	A-D		D-F	A-E	A-H	HY44C-15 example in Fig. 5
HYLC46	M	$DBH^{Cre};Rosa^{Ai32}$	12/30/15	06/08/16	MeV	46A-6‡,7‡						E,G,H	A-D		D-F	A-E	A-H	
HYLC48	M	$DBH^{Cre};Rosa^{Ai32}$	01/10/16	08/17/16	MeV	48D-5‡,6‡,7‡,8‡						E,G,H	A-D		D-F	A-E	A-H	
HYLC52	M	$DBH^{Cre};Rosa^{Ai32}$	05/12/16	08/31/16	MeV	52A-16‡,17,18‡						E,G,H	A-D		D-F	A-E	A-H	
HYLC53	M	$DBH^{Cre};Rosa^{Ai32}$	05/12/16	09/02/16	MeV	53A-9‡,22‡,23‡,24,25‡,26						E,G,H	A-D		D-F	A-E	A-H	
HYLC55	M	$DBH^{Cre};Rosa^{Ai32}$	06/06/16	09/16/16	MeV	55A-3‡,4‡,5‡,6,7,8‡,9‡,10‡,11‡,12‡,13,14‡,19‡,20						E,G,H	A-D		D-F	A-E	A-H	
HYLC56	M	$DBH^{Cre};Rosa^{Ai32}$	06/06/16	09/26/16	MeV	56A-5‡,6,7,8‡,12‡,13‡,14,15‡,16						E,G,H	A-D		D-F	A-E	A-H	
Recordings of facial and whisker movement																		
KSt159	F	$TrKC^{CreER};Rosa^{Ai32}$	12/07/16	10/20/17	SO; face; whisker	N.A.					E,F	D,E						
KSt160	M	$TrKC^{CreER};Rosa^{Ai32}$	12/07/16	10/14, 10/20/17	SO; whisker	N.A.					E,F							face video registration failed
KSt164	M	$TrKC^{CreER};Rosa^{Ai32}$	08/08/17	11/08/17	G; SO; face; whisker	N.A.					B-F		B-E					Example in Videos S2, S3
KSt165	F	$TrKC^{CreER};Rosa^{Ai32}$	08/09/17	02/20/18	G; whisker	N.A.					B-F			D-F				Example in Video S4
KSt166	M	$TrKC^{CreER};Rosa^{Ai32}$	08/09/17	02/09/18	G; whisker	N.A.					E,F							Example in Video S1
KSt167	F	$TrKC^{CreER};Rosa^{Ai32}$	08/09/17	02/24/18	G; whisker	N.A.					E,F							
KSt168	F	$TrKC^{CreER};Rosa^{Ai32}$	08/09/17	05/04, 05/06/18	SO; micro; whisker	N.A.					E,F							
KSt169	F	$TrKC^{CreER};Rosa^{Ai32}$	08/09/17	05/05, 05/06/18	SO; micro; whisker	N.A.					B-F		A-C					Example in Video S5
KSt170	M	$Ret^{CreER};Rosa^{Ai32}$	12/05/17	05/05, 05/06/18	SO; micro; whisker	N.A.					E,F							
KSt173	M	$TrKC^{CreER};Rosa^{Ai32}$	02/01/18	12/02-12/08/18	G; SO	N.A.					G	K						
KSt176	M	$TrKC^{CreER};Rosa^{Ai32}$	02/01/18	12/02-12/08/18	G; SO	N.A.					G		H,I,K					Example in Video S6
KSt178	M	$TrKC^{CreER};Rosa^{Ai32}$	02/01/18	12/02-12/08/18	G; SO	N.A.					G		H,J,K					Example in Video S7

Supplemental Table 1: Meta-data and mouse information