Sleep regulates visual selective attention in *Drosophila*

Leonie Kirszenblat[1], Deniz Ertekin[1], Joseph Goodsell[1], Yanqiong Zhou[1], Paul J Shaw[2], Bruno van Swinderen[1].

[1] Queensland Brain Institute, The University of Queensland, Brisbane, QLD, 4072, Australia.

[2] Department of Anatomy and Neurobiology, Washington University in St. Louis, 660 South Euclid Avenue, St. Louis, MO 63110, USA.

Corresponding author: b.vanswinderen@uq.edu.au

Key words: attention, sleep, *Drosophila*, flies, behaviour

Summary statement: Sleep deprivation specifically impairs visual selective attention in fruit flies, without affecting behavioural responses to simple visual stimuli.

Abstract

Although sleep-deprivation is known to impair attention in humans and other mammals, the underlying reasons are not well understood, and whether similar effects are present in non-mammalian species is not known. We therefore sought to investigate whether sleep is important for optimising attention in an invertebrate species, the genetic model *Drosophila melanogaster*. We developed a high-throughput paradigm to measure visual attention in freely-walking *Drosophila*, using competing foreground/background visual stimuli. We found that whereas sleep-deprived flies could respond normally to either stimulus alone, they were more distracted by background cues in a visual competition task. Other stressful manipulations such as starvation, heat exposure, and mechanical stress had no effects on visual attention in this paradigm. In contrast to sleep-deprivation, providing additional sleep using the GABA-A agonist 4,5,6,7-tetrahydroisoxazolo-[5,4-c]pyridine-3-ol (THIP) did not affect attention in wild-type flies, but significantly improved attention in the learning mutant *dunce*. Our results reveal a key function of sleep in optimising attention processes in *Drosophila*, and establish a behavioural paradigm that can be used to explore the molecular mechanisms involved.

Introduction

The restorative effect of a night's sleep seems obvious, yet we still know very little about the function of sleep and how it impacts our behaviour. Studies in humans and other animals suggest that a fundamental function of sleep is to preserve cognitive functions such as learning, memory and attention (Stickgold 2005). This implies that sleep promotes brain plasticity, for example by strengthening neuronal circuits to consolidate memories (Diekelmann & Born 2010), or by maintaining optimal levels of neuronal functions by globally altering synaptic strengths (Tononi & Cirelli 2013). If a major function of sleep is to promote plasticity, a brain process that may be particularly vulnerable to sleep loss is selective attention. Given that selective attention requires precise temporal coordination between different neural populations (Fries et al. 2001; Elhilali et al. 2009), it may be most vulnerable to the changes in neural processing that may accrue when sleep homeostasis mechanisms are not in place (Kirszenblat & van Swinderen 2015). Consistent with this idea,

sleep-deprivation in humans leaves basic sensory processing intact (Casagrande et al. 2006;

Killgore 2010; Kendall et al. 2006), whereas tasks that involve high attentional load are

impaired (Chee & Chuah 2007; Kong et al. 2011). For this reason, sleep and attention may

share a deeper relationship than previously thought, and one that has not been thoroughly

investigated in genetic models such as *Drosophila*.

In *Drosophila*, sleep-deprivation has been associated with deficits in a variety of behaviours.

Sleep-deprivation impacts olfactory and visual memory in *Drosophila* (Glou et al. 2012;

Seugnet et al. 2008), as well as courtship memory (Ganguly-Fitzgerald et al. 2006) and

aggression in male flies (Kayser et al. 2015). In contrast to sleep-deprivation, induction of

sleep has been shown to reverse memory deficits in flies with genetic lesions and in a fly

model of Alzheimer's disease (Dissel, Melnattur, et al. 2015; Dissel, Angadi, et al. 2015). This

suggests that sleep can be used as a powerful therapeutic to enhance memory. As the

ability to pay attention is a prerequisite for all of the aforementioned behaviours, we

questioned whether optimising selective attention may be a key function of sleep in

Drosophila.

In this study, we developed a simple, high-throughput method to study visual selective

attention in flies. We found that whereas sleep-deprivation did not affect simple visual

behaviours (optomotor and fixation), sleep-deprived flies were more distractible in a visual

attention task involving competing stimuli. Normal attention was restored following sleep,

indicating that sleep promotes behavioural plasticity. In contrast to sleep-deprivation, we

found that inducing additional sleep using the GABA-A agonist 4,5,6,7-tetrahydroisoxazolo-

[5,4-c]pyridine-3-ol (THIP) had no effect on wild-type flies but could restore normal

attention to *dunce* learning mutants. Together, our results suggest that sleep optimises

selective attention processes in *Drosophila*.

Methods

Fly stocks

Strains used in this study were *Canton-S* (*CS*) wild-type flies, *dunce[1]* mutants (outcrossed to

CS), *rutabaga[2080]*, *pdf-gal4* flies (Bloomington Stock Centre, Indiana), and *UAS-Chrimson*

(P[20xUAS-IVS-CsChrimson.mVenus]attp18) flies (a gift from Vivek Jayaraman, Janelia Farm

Research Campus, Virginia, U.S.A).

Sleep and visual behaviour

Our visual arena was adapted from Buridan's paradigm (Götz 1980). Flies had their wings

clipped on CO_2, at least 2 days prior to the experiment. During the experiment, flies walked

freely on a round platform, 86mm in diameter, surrounded by a water-filled moat to

prevent escape. An individual fly was only tested once in each experiment, such that it was

not influenced by previous visual stimuli.   The temperature of the arena was 24-26 °C

during experiments. Each experiment lasted 3 minutes, and the visual stimuli were

presented on the horizontal or the vertical axes in alternation. Optomotor experiments

were conducted with clockwise and anticlockwise gratings for 1.5 minutes each. A camera

(Sony Hi Resolution Colour Video Camera CCD-IRIS SSC-374) placed above the arena was

used to detect the fly's movement on the platform at 30 frames per second, and open-

source tracking software was used to record the position of the fly (Colomb et al. 2012).

Sleep was quantified using the DART system as previously described (Faville et al. 2015)

using the 5 minute criterion for sleep (Shaw et al. 2000; Hendricks et al. 2000).

Visual stimuli

Each LED panel comprised 1024 individual LED units (32 rows by 32 columns) and was

computer-controlled with LED Studio software (Shenzen Sinorad, Medical Electronics,

Shenzen, China).   The LEDs had a refresh rate of 200 Hz, ensuring there was no background

flicker visible to the flies. All visual stimuli were created in Vision Egg software (Maimon et

al. 2008), written in Python programming language (Ferguson et al. 2017). The walls of the

arena consisted of 6 LED panels of green (520nm) and blue (468nm) LEDs that formed a

hexagon surrounding the moat (29cm diameter, 16cm height), and onto which the visual

stimuli were presented. Fixation stimuli were two dark stripes 180° apart, each 9° in width

and 45° in height from the centre of the arena. The fixation stripes ranged from 38 - 55° in

height and 4 – 14° in width depending on the fly's position in the arena.   For visual

competition experiments, 7 Hz flickering stripes (targets) were overlaid on top of a 3 Hz

grating (speed 54° s^{-1}, luminance 402 Lux) in the background. For visual experiments in

which stationary stripes were used as a distractor, the target stripes were flickered at 3 Hz,

and the distractor stripes were non-flickering. For experiments in which the luminance

contrast of the grating was increased, the increments used were 0, 75, 146, 224, 402, 473,

649, 730 Lux.

Sleep-deprivation

3-5 day old female CS flies with clipped wings were sleep-deprived for 24 hours from 11am

until 11am the next day, and then tested in the visual arena immediately afterwards.

Mechanical sleep-deprivation was achieved using a sleep nullifying apparatus (SNAP) device,

which has been shown to sleep deprive flies without triggering stress responses (Shaw et al.

2002). The device tilted back and forth, forcefully knocking and displacing flies every 20s (or

10s where indicated). Flies were contained in vials (20 females, 5 males) during the sleep-

deprivation, because containing them individually in small tubes affected their performance

in the attention assay. During the experiments, flies waiting to be tested were gently

handled at least every 3 minutes so that they could not fall asleep prior to testing.

Optogenetics

105   Flies were grown on standard (agar/yeast) medium until 2-3 days prior to an experiment,

106   after which they were transferred to standard medium supplemented with 0.5µM all-*trans*-

107   retinal. Chrimson was activated using Red-Orange LEDS (Luxeon Rebel, 617nm, 700mA,

108   Phillips LXM2-PH021-0070) as previously reported (Klapoetke et al. 2014). Flies were

109   illuminated with >1000 Lux from 4 LED arrays.

Pharmacology

111   THIP was administered to flies in standard food at 0.1mg/ml concentration for 2 days, and

112   removed 1 hour prior to measuring performance, as described previously (Dissel et al.,

113   2015).

Data Analyses

Analyses of visual responses were performed using CeTran (3.4) software (Colomb et al,

2012), as well as custom-made scripts in R programming language. Stripe deviation was

calculated as the smallest angle between the fly's trajectory and either of the vertical stripes

(Ferguson et al. 2017; Colomb et al. 2012). For optomotor responses, the angular velocity

(turning angle in $° s^{-1}$) in the direction of the moving grating was calculated. All sleep and

arousal metrics were obtained through the DART software. Statistical analyses were

performed using Prism, R and MATLAB software. Lillifors tests were performed to confirm

the normality of the data, and *t* tests or one-way ANOVAs were used to detect significant

differences between groups.

Results

A visual attention paradigm for freely walking flies.

Selective attention allows us to focus on a single object or group of objects, while ignoring

less salient information (La Berge 1983; Eriksen & St James 1986). To examine attention-like

behaviour in freely-walking *Drosophila*, we designed a paradigm to measure behavioural

responses of flies to competing visual stimuli, involving 'targets' and 'distractors'. We took

advantage of two robust visual behaviours in *Drosophila*: fixation – a fly's tendency to orient

and walk towards a visually salient object, and optomotor behaviour, whereby a fly will turn

in the same direction as wide-field motion to stabilize its visual surroundings (Heisenberg et

al. 1984). Fixation has been previously measured using Buridan's paradigm, where flies with

clipped wings walk back and forth between two opposing vertical black stripes (Götz 1980;

Ferguson et al. 2017). We modified this paradigm to measure behavioural responses to

competing visual stimuli, using two opposing stripes as 'targets' in the foreground and

adding a wide-field motion stimulus as the 'distractor' in the background (Fig. 1A). A similar

configuration has been used for 'Fig./ground' discrimination in tethered flight experiments

(Heisenberg et al. 1984; Fox et al. 2014; Fenk et al. 2014; Aptekar et al. 2015; Aptekar & Frye

2013), but this complex stimulus has not been tested in walking flies.

We measured fixation on the targets by calculating the deviation angle between the fly's

heading direction and the target stripe, with smaller angles indicating greater attention to

the target (Fig. 1B) (Colomb et al. 2012). Response to the distractor, the motion stimulus,

was measured by the angular velocity in the direction of motion (the turning angle s^{-1} (γ):

Fig. 1B). First, we investigated visual responses to both stimuli alone, across a range of

frequencies (Fig. S1A,B). Flies fixated best on targets flickering in the range of 3 – 7 Hz (Fig.

S1A), while optomotor responses were highest at 16 Hz (Supplementary Fig. 1B). For visual

competition experiments, we used a target frequency of 7 Hz, as this stimulus was very

salient to the flies and has previously been found to evoke strong fixation behaviour (Paulk

et al. 2015). The distractor was a lower contrast grating that spanned the entire arena and

moved "behind" the flickering targets. We used a grating frequency of 3 Hz (54 ° s^{-1}) at a

mid-luminance contrast, since we found responses under these conditions were robust and

consistent among flies, eliciting circular walking responses (Fig. 1E), and provided a low

enough level of distraction for the flies such that they could still continue to fixate on the

targets. Adding this low-contrast motion distractor significantly reduced flies' fixation on

the target stripes: the straight paths evident when the targets were presented alone (Fig.

1C) were replaced by a combination of straight paths and circular paths, often angled in the

direction of motion (Fig. 1D). We modified the salience of the grating distractor by adjusting

the luminance contrast in linear increments (see methods) (Fig. 1F,G). As expected, when

the grating had a higher contrast (more salient), the distraction effect increased (Fig. 1F).

Motion responses to the distractor also increased (Fig. 1G) but appeared to plateau earlier

than target deviation in response to increasing the distractor salience. This suggests that

increased target deviation (Fig. 1F) may be a more sensitive measure of the distractor effect

than increased optomotor behaviour (Fig. 1G). In summary, our results show that attention-

like responses towards fixed targets can be effectively modulated (and titrated) by a motion

distractor stimulus in freely walking flies.

Sleep-deprived flies are more distractible.

Studies in humans and other animals have shown that sleep loss can impair a variety of

cognitive behaviours, such as learning, memory and attention (Alhola & Polo-kantola 2007;

Drummond et al. 2012; Lim & Dinges 2010). Similarly, studies in *Drosophila* have indicated a

role for sleep in visual and olfactory memory (Glou et al. 2012; Li et al. 2009; Seugnet et al.

2008; Seugnet et al. 2011). However, it is not known whether sleep modulates attention-like

behaviour in *Drosophila*. To investigate whether sleep-deprivation affects visual attention,

we sleep-deprived flies for 24 hours using the Sleep Nullifying Apparatus (SNAP) according

to previous methods (Shaw et al. 2002). We first confirmed that flies were sleep-deprived

by examining their sleep rebound (*i.e.* an increase in sleep need following sleep-

deprivation). Following sleep-deprivation, sleep was monitored using the DART system

(Faville et al. 2015) across three consecutive days, while another group of flies had their

attention tested (Fig. 2A). As expected, sleep-deprived flies slept more than control flies the

day after sleep-deprivation (day 1) particularly during the daytime, whereas by days 2 and 3

there was no difference in sleep duration between the sleep-deprived and control flies (Fig.

2B).

Following sleep-deprivation we measured fixation to stationary targets and optomotor

responses to a moving grating, in scenarios in which the stimuli were presented alone or in

competition (with the grating at three different levels of luminance contrast of the

distractor) (Fig. 2D,E). Optomotor responses to the moving grating in all visual scenarios

were largely unaffected by sleep-deprivation (Fig. 2E), as were fixation responses in the

absence of the grating (Fig. 2D, first point from the left). Interestingly, the effect of sleep-

deprivation on fixation behaviour was only evident when the two visual stimuli were

combined (Fig. 2C), as indicated by significantly increased target deviation for all three

combined conditions (Fig. 2D). This suggests that sleep-deprived flies are specifically

affected in their response to visual competition. Finally, we tested whether flies could

recover from sleep-deprivation by allowing them to rest for 24 hours. For this experiment

(and all subsequent experiments involving visual competition), we used a grating of mid-

luminance contrast (indicated by the red arrow heads in Fig. 2E,F). Again, we found that

sleep-deprived flies were more distractible, but their attention returned to normal following

24 hours rest (Fig. 2F). Overall, our results suggest that sleep-deprivation alters visual

attention, and that this effect is reversible.

Although we saw no alterations to optomotor responses of sleep-deprived flies when the

grating was presented alone, we considered the possibility that the sleep-deprivation

phenotype may be caused by an increased sensitivity to motion, which could potentially

arise from the preceding 24hours of constant motion they experience in the SNAP

apparatus. However, flies sleep-deprived in 24hours of darkness showed a similar

impairment of attention compared to controls, suggesting that visual experience did not

play a role in the sleep-deprivation phenotype (Fig. S2). We next asked whether the

increased distractibility of sleep-deprived flies was specific to the motion stimulus, or

whether it could apply to other types of visual distractors. We used a stationary stripe as a

distractor that was identical to the target but less salient as it was non-flickering (Fig. 3A).

As with the moving grating in the preceding experiments, attention to the target could be

titrated by modulating the salience (*i.e.,* contrast) of the distractor (Fig. 3B,C). Interestingly,

increasing the salience of the distractor correspondingly increased distractibility (Fig. 3C).

Furthermore, consistent with our previous attention experiments, we found that sleep-

deprived flies similarly paid less attention to the target in the presence of this alternative

distractor object (Fig. 3D). Our observation that sleep-deprivation is similarly disrupted by

different types of distractors suggests that a common attention mechanism is affected,

which is not dependent on lower level visual responses.

Given that sleep-deprivation reduces overall activity, we wondered whether this may

explain the reduced orientation to salient features in the visual attention assay. To

investigate this, we performed a correlation analysis on all data, to see if walking speed of

individual flies may predict selective attention behaviour (target deviation). However, we

found no correlation between attention and walking speed for either sleep-deprived or

control groups (Fig. 4A,B).

Considering that the sleep-deprivation protocol might be stressful (it involves waking the

flies every 20 seconds by mechanical stimulation), we further examined the effects of

different types of stress on the flies' visual attention. This allowed us to answer two

questions: 1) whether attention phenotypes could be modulated by environmental stressors

(e.g. heat, mechanical stress and starvation), which would indicate that the sleep-

deprivation effect on attention may simply be caused by stressful situations, and 2) whether

the sleep-deprivation phenotype occurs under different environmental conditions *(i.e.,* how

robust the effect is).   For all these experiments we measured flies' visual responses in the

combined stimulus paradigm (in which flies fixate on targets in the presence of distractors)

(Fig. 1A).

First, we measured the effect of heating flies to 31°C for 24 hours (Fig. 5A). Prolonged

exposure to heat had no effect on attention, whereas the increase in distractibility caused

by sleep-deprivation was still present under conditions of heat stress (Fig. 5A). This

indicates that heat does not influence attention in this paradigm, or the sleep-deprived

effects on attention. Next, we measured the effects of mechanical stress using a stimulation

protocol that would provide the same amount of stimulation experienced during the sleep-

deprivation protocol, but including rest periods for sleep. This was achieved by subjecting

flies to mechanical stimulation for one hour followed by rest for one hour, repeated across

24 hours, but with double the rate of stimulation (to keep the actual number of stimuli the

same as for the sleep-deprivation regime) (Fig. 5B). Unlike sleep-deprived flies, flies

subjected to this stimulation regime containing rest periods had normal attention, indicating

that the sleep-deprivation effect was not due to mechanical stress of perturbing the flies

(Fig. 5B). We next assessed the effects of starvation, as sleep is known to be affected by

food availability (Siegel et al, 2009). In *Drosophila*, food deprivation has been found to

suppress sleep (Keene et al. 2010). Feeding was assessed during the sleep-deprivation by

measuring the intake of food containing blue food dye and, as previously reported, sleep-

deprived flies showed similar food intake to controls (Thimgan et al. 2010). We then tested

whether depriving flies of food, with or without sleep-deprivation, affected visual attention.

The visual attention scores of starved controls were not different from those of fed controls,

suggesting that starvation *per se* does not impair visual attention (Fig. 5C). As expected,

sleep-deprivation impaired visual attention in both starved and fed conditions but,

interestingly, starvation was able to partially suppress this effect (Fig. 5C, "SD fed" vs "SD

starved"), suggesting that starvation may protect against the detrimental effects of sleep-

deprivation. In summary, our results show that sleep-deprivation consistently affects visual

attention, even under conditions of heat and starvation, but that visual attention is resistant

to the effects of a variety of stresses in the absence of sleep-deprivation. Therefore, sleep is

more important than lack of stress for maintaining optimal levels of visual attention.

Having found that sleep-deprivation affects visual attention, we also sought alternative

methods to sleep deprive flies, to confirm that the sleep-deprivation effect on attention was

not specific to our mechanical stimulation protocol. Pigment Dispersing Factor (PDF)

neurons in the optic lobes of the fly brain are known to promote wake (Parisky et al. 2008).

We hypothesised that activating these neurons for 24 hours would have wake-promoting

effects that may also impair visual attention. To test this, we transiently expressed a red-

light activated channelrhodopsin, Chrimson (Klapoetke et al. 2014), in the Pigment

Dispersing Factor (PDF) neurons, such that exposing flies to red light could trigger

wakefulness. We noticed that red light exposure caused a general decrease in sleep among

all flies (Fig. 6A). However, sleep loss was greater in flies that expressed the

channelrhodopsin (*pdf-gal4/+>uas-chrimson/+*) and were exposed to red light for 24 hours,

as they lost more than 55% of their sleep compared to non-exposed controls. When we

subsequently tested these flies for visual attention, we found that this method of sleep-

deprivation resulted in an increase in target deviation specifically in the flies that were kept

awake genetically, *pdf-gal4/+>uas-chrimson/+* (Fig. 6B,C). This provides further support for

the idea that sleep-deprivation increases distractibility, and is not specific to the method of

sleep-deprivation.

Additional sleep improves attention in *dunce* mutants, but not wild-type flies.

Considering that sleep-deprivation made flies more distractible, we next asked whether

increasing sleep had the opposite effect on visual attention, making flies *less* distractible.

The GABA-A agonist (THIP) has recently been used to induce sleep in *Drosophila* (Dissel,

Angadi, et al. 2015; Yap et al. 2017). This increase in sleep was shown to reverse memory

deficits in *Drosophila* learning mutants *dunce* and *rutabaga* (Dissel, Angadi, et al. 2015) and

in a *Drosophila* model of Alzheimer's disease (Dissel et al. 2017). As *dunce* and *rutabaga*

mutants have previously been found to have defective attention processes (van Swinderen

& Brembs 2010; van Swinderen 2007), we wondered whether their attention deficits could

also be rescued by induced sleep.

We first quantified sleep in *dunce* and *rutabaga* mutants, and found that both these

mutants slept significantly less than wild-type flies, with *dunce* flies sleeping less during the

day and the night, and *rutabaga* mutants sleeping less during the day (Fig. 7A,B). THIP could

then be effectively used to increase sleep in both wild-type and mutant flies, to a similar

level (Fig. 7C,D). We next tested whether inducing sleep affected the visual attention of

wild-type flies in our free-walking paradigm. THIP was fed to wild-type (CS) flies for two

days and removed one hour prior to testing their behaviour (the same procedure used by

(Dissel, Melnattur, et al. 2015). We performed a within-group experiment (the same flies

tested before and after induced sleep), and a between-group experiment (aged-matched

flies, with or without induced sleep). In wild-type flies we observed that distractibility

(deviation away from the targets) decreased slightly for those flies that that had been

induced to sleep more, in both experiments; however, this effect was not significant (Fig. 7E,

p=0.084 and p=0.187 for within group and between group comparisons, by t-test.).

We next tested whether increasing sleep altered attention in *dunce* and *rutabaga* mutants.

Under normal conditions (without increasing sleep), *dunce* and *rutabaga* flies were

significantly less attentive towards the visual targets compared to wild-type flies as

measured by increased target deviation (Fig. 7F, black bar (WT) compared to light blue bar

(*dnc* (-)) and yellow bar (*rut* (-))). Interestingly, THIP administration was able to significantly

reduce target deviation in *dunce* mutants but not *rutabaga* mutants (Fig. 7F), such that

attention in *dunce* mutants with induced sleep was not different from that of wild-type flies

(Fig. 7F, dark blue bar (*dnc* (+)) compared to black bar (WT(-))). This suggests that increasing

sleep may improve attention in some learning mutants but not others.

Discussion

Selective attention is crucial for discriminating important from irrelevant stimuli in our

environment, even for insects such as *Drosophila*. The selection and suppression dynamics

required for attention seem to have emerged in brains at the same time at which sleep

became important for maintaining attention-related behaviours (Kirszenblat & van

Swinderen 2015). In other words, some sleep functions may have evolved to optimise

attention. Here, we have developed a free-walking attention paradigm which allowed us to

obtain a functional readout of the effects of sleep-deprivation and sleep induction on visual

attention in *Drosophila*.

Selective attention must require a certain level of brain coordination to deal with competing

stimuli. This is because different stimuli may be processed by different brain regions; in our

attention paradigm, the optomotor and fixation responses that compete with each other

have been found to be driven to some degree by independent visual circuits (Bahl et al.

2013; Fenk et al. 2014). Furthermore, attention-like behaviour in the fly is associated with

increased coherence between brain regions (Paulk et al. 2015), suggesting greater synaptic

coordination. If sleep is required to maintain synaptic coordination across the brain, it

follows that tasks involving greater cognitive load may be more vulnerable to sleep loss.

Indeed, we found that although sleep-deprived flies showed normal responses to simple

visual stimuli, they had altered responses to visual competition.

One might still ask, whether sleep has a privileged role in regulating selective attention

relative to other high order cognitive functions. Sleep loss is known to affect a variety of

complex cognitive behaviours in humans, such as learning and memory, creative thinking,

and even the ability to speak clearly or to appreciate humour (Harrison & Horne 1997;

Killgore 2010; Kendall et al. 2006). Attention is probably integral to these complex processes

because it allows us to filter out irrelevant information and to select the right actions. In

light of this, we would suggest that selective attention is a key *mechanism* that is affected by

sleep loss, which disrupts the brain's ability to prioritize competing information.

The sleep-deprivation effects we observed on attention seemed to be remarkably robust

under different environmental conditions, as flies were still more distractible following

sleep-deprivation in conditions of heat stress, darkness and starvation. Interestingly,

although starvation and heat alone can promote waking in flies (Ishimoto et al. 2012; Keene

et al. 2010), they did not appear to affect our attention readout, unlike the mechanical

perturbation method of sleep-deprivation which is in standard use for *Drosophila*. It is

possible that because starvation and heat are environmental stimuli that a fly encounters in

natural situations, it has already evolved physiological protective mechanisms to cope with

these kinds of stresses. This is supported by reports that heat stress response factors can

protect against sleep-deprivation-induced lethality (Shaw et al., 2002), and that starvation

disrupts sleep homeostasis (Keene et al. 2010; Thimgan et al. 2010). *Foraging* mutants of

Drosophila, which tend to explore further for food, also appear to have reduced sleep need

and are resistant to memory impairments caused by sleep loss (Donlea et al. 2012). More

recently, it was also discovered that sexual arousal in male flies can suppress the need for

sleep (Beckwith et al. 2017), and vice versa, sleep can inhibit male sexual behaviour (Chen et

al. 2017). Overall, these studies suggest that sleep need is flexible and may compete with

other survival needs such as food, sexual reproduction and the need to escape unfavourable

environments (e.g. high temperatures). In future it would be interesting to further

investigate how a fly's environment influences its attention.

Interestingly, *increasing* sleep was able to improve the attention of *dunce* mutants, but not

wild-type flies or *rutabag*a mutants. The finding that wild-type flies did not show improved

attention following induced sleep is consistent with previous reports that increasing sleep

does not improve learning and memory in wild-type flies (Dissel, Angadi, et al. 2015). One

interpretation is that attention is already optimal in wild-type *Drosophila*. In contrast, both

rutabaga and *dunce* mutants have previously been identified as having attention deficits

(van Swinderen 2007), and our result confirmed this finding in a free-walking attention

paradigm. However, it is not clear why inducing sleep would improve attention specifically

in *dunce* mutants, and not in *rutabaga* mutants. It is possible that this may relate to their

different sleep phenotypes – *dunce* mutants appear more severely sleep-deficient in our

DART system (including at night, which *rutabaga* mutants are not) meaning that THIP had a

greater ability to restore sleep to *dunce* mutants. Related to this, it is possible that the

attention deficits of these mutants are due to different underlying causes. For example, the

mutants may have poor attention because they sleep less, or they may sleep less because

they have poor attention.

How inducing sleep could improve attention, or other aspects of cognition such as learning

and memory, also remains unclear. There is some evidence that the sleep-promoting agent

THIP can interact with specific GABA-receptors to promote sleep in flies, via known sleep-

promoting neurons of the dorsal fan-shaped body (Dissel, Angadi, et al. 2015), but can also

modulate dopaminergic pathways to facilitate memories (Berry et al. 2015). Interestingly, a

recent study found that increasing sleep promoted survival of wild-type flies exposed to

oxidative stress (Hill et al. 2018). Whether increasing sleep in flies improves brain function

through specific circuits, or by general cellular mechanisms like reducing reactive oxygen

species (ROS) needs to be further investigated. Our study opens up an opportunity to

understand the molecules, circuits and environmental factors involved in the role of sleep in

optimising attention.

Acknowledgements

We would like to thank Rowan Tweedale for comments on the manuscript. We would like

to thank John John for help with behavioural experiments, and Richard Faville and Ben

Kottler for help with DART software. This work was supported by an NIH grant RO1

NS076980-01 to PJS and BVS and an Australian Research Council grant DP140103184 to BVS.

Competing Interests

The authors declare no competing interests.

<u>References</u>

Alhola, P. & Polo-kantola, P., 2007. Sleep deprivation : Impact on cognitive performance.

Neuropsychiatric Disease and Treatment, 3(5), pp.553–567.

Aptekar, J.W. et al., 2015. Neurons forming optic glomeruli compute figure – ground

discriminations in Drosophila. *The Journal of Neuroscie*, 35(19), pp.7587–7599.

Aptekar, J.W. & Frye, M.A., 2013. Higher-order figure discrimination in fly and human vision.

Current Biology, 23(16), pp.694–700. Available at:

http://dx.doi.org/10.1016/j.cub.2013.07.022.

Bahl, A. et al., 2013. Object tracking in motion-blind flies. *Nature Neuroscience*, pp.1–11.

Available at: http://dx.doi.org/10.1038/nn.3386.

Beckwith, E.J. et al., 2017. Regulation of sleep homeostasis by sexual arousal. *eLife*, 6, pp.1–

19.

La Berge, D., 1983. Spatial extent of attention to letters and words. *Journal of experimental*

psychology. Human perception and performance, 9(3), pp.371–379.

Berry, J.A. et al., 2015. Sleep facilitates memory by blocking dopamine neuron-mediated

forgetting. *Cell*, 161(7), pp.1656–1667. Available at:

http://www.sciencedirect.com/science/article/pii/S0092867415005772.

Casagrande, M. et al., 2006. Orienting and alerting: Effect of 24 h of prolonged wakefulness.

Experimental Brain Research, 171(2), pp.184–193.

Chee, M.W.L. & Chuah, Y.M.L., 2007. Functional neuroimaging and behavioral correlates of

capacity decline in visual short-term memory after sleep deprivation. *Proceedings of*

the National Academy of Sciences of the United States of America, 104(22), pp.9487–

9492.

Chen, D. et al., 2017. Genetic and neuronal mechanisms governing the sex-specific

interaction between sleep and sexual behaviors in Drosophila. *Nature Communications*.

8 (154).

Colomb, J. et al., 2012. Open source tracking and analysis of adult Drosophila locomotion in

Buridan's paradigm with and without visual targets. *PLoS ONE*, 7(8), pp.1–12.

Diekelmann, S. & Born, J., 2010. The memory function of sleep. *Nature Reviews*
Neuroscience, 11, p.114. Available at: http://dx.doi.org/10.1038/nrn2762.

Dissel, S. et al., 2017. Enhanced sleep reverses memory deficits and underlying pathology in
drosophila models of Alzheimer's disease. *Neurobiology of Sleep and Circadian*
Rhythms.

Dissel, S., Angadi, V., et al., 2015. Sleep restores behavioral plasticity to drosophila mutants.
Current Biology, 25(10), pp.1270–1281. Available at:
http://dx.doi.org/10.1016/j.cub.2015.03.027.

Dissel, S., Melnattur, K. & Shaw, P.J., 2015. Sleep, Performance, and Memory in Flies.
Current Sleep Medicine Reports, 1(1), pp.47–54.

Donlea, J. et al., 2012. foraging alters resilience/vulnerability to sleep disruption and
starvation in Drosophila. *Proceedings of the National Academy of Sciences*, 109(7),
pp.2613–2618. Available at: http://www.pnas.org/cgi/doi/10.1073/pnas.1112623109.

Drummond, S.P.A. et al., 2012. The effects of two types of sleep deprivation on visual
working memory capacity and filtering efficiency. *PLoS ONE*, 7(4), pp.1–8.

Elhilali, M. et al., 2009. Interaction between Attention and Bottom-Up Saliency Mediates the
Representation of Foreground and Background in an Auditory Scene. *PLOS Biology*,
7(6), p.e1000129. Available at: https://doi.org/10.1371/journal.pbio.1000129.

Eriksen, C.W. & St James, J.D., 1986. Visual attention within and around the field of focal
attention: a zoom lens model. *Perception & psychophysics*, 40(4), pp.225–240.

Faville, R. et al., 2015. How deeply does your mutant sleep? Probing arousal to better
understand sleep defects in Drosophila. *Scientific Reports*, 5, p.8454. Available at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4326961&tool=pmcentrez
&rendertype=abstract.

Fenk, L.M., Poehlmann, A. & Straw, A.D., 2014. Asymmetric processing of visual motion for
simultaneous object and background responses. *Current Biology*, 24(24), pp.2913–
2919. Available at: http://dx.doi.org/10.1016/j.cub.2014.10.042.

Ferguson, L. et al., 2017. Transient dysregulation of dopamine signaling in a developing

Drosophila arousal circuit permanently impairs behavioral responsiveness in adults.

Frontiers in Psychiatry, 8(February), p.22.

Fox, J.L. et al., 2014. Figure-ground discrimination behavior in Drosophila. I. Spatial

organization of wing-steering responses. *Journal of Experimental Biology*, 217(4),

pp.558–569. Available at: http://jeb.biologists.org/cgi/doi/10.1242/jeb.097220.

Fries, P. et al., 2001. Modulation of oscillatory neuronal synchronization by selective visual

attention. *Science*, 291(5508), pp.1560–1563. Available at:

http://science.sciencemag.org/content/291/5508/1560.abstract.

Ganguly-Fitzgerald, I., Donlea, J. & Shaw, P.J., 2006. Waking experience affects sleep need in

Drosophila. *Science*, 313(5794), pp.1775–81. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/16990546.

Glou, E. Le et al., 2012. Circadian modulation of consolidated memory retrieval following

sleep deprivation in Drosophila. *Sleep*, 35(10), pp.2–11.

Götz, K.G., 1980. Visual Guidance in Drosophila. *Basic. Life. Sci. 16*, 391-407.

Harrison, Y. & Horne, J.A., 1997. Sleep deprivation affects speech. *Sleep*, 20(10), pp.871–

877.

Heisenberg, M. et al., 1984. Vision in Drosophila. Genetics of Microbehavior. , pp.1–250.

Hendricks, J.C. et al., 2000. Rest in Drosophila Is a Sleep-like State. *Neuron*, 25(1), pp.129–

138. Available at: http://linkinghub.elsevier.com/retrieve/pii/S0896627300808776.

Hill, V.M. et al., 2018. A bidirectional relationship between sleep and oxidative stress in

Drosophila P. Taghert, ed. *PLOS Biology*, 16(7), p.e2005206. Available at:

http://dx.plos.org/10.1371/journal.pbio.2005206 [Accessed August 6, 2018].

Ishimoto, H., Lark, A. & Kitamoto, T., 2012. Factors that differentially affect daytime and

nighttime sleep in Drosophila melanogaster. , 3(February), pp.1–5.

Kayser, M.S. et al., 2015. Sleep deprivation suppresses aggression in Drosophila. *eLife*,

4(JULY 2015).

Keene, A.C. et al., 2010. Clock and cycle limit starvation-induced sleep loss in drosophila.
Current Biology, 20(13), pp.1209–1215.

Kendall, A.P. et al., 2006. Effects of sleep deprivation on lateral visual attention. *The*
International journal of neuroscience, 116(10), pp.1125–38. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/16923682.

Killgore, W.D.S., 2010. *Effects of sleep deprivation on cognition*, Elsevier B.V. Available at:
http://dx.doi.org/10.1016/B978-0-444-53702-7.00007-5.

Kirszenblat, L. & van Swinderen, B., 2015. The Yin and Yang of Sleep and Attention. *Trends in*
Neurosciences, 38(12), pp.776–786.

Klapoetke, N.C. et al., 2014. Independent optical excitation of distinct neural populations.
Nature methods, 11(3), pp.338–46. Available at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3943671&tool=pmcentrez
&rendertype=abstract.

Kong, D., Soon, C.S. & Chee, M.W.L., 2011. Reduced visual processing capacity in sleep
deprived persons. *NeuroImage*, 55(2), pp.629–634. Available at:
http://dx.doi.org/10.1016/j.neuroimage.2010.12.057.

Li, X., Yu, F. & Guo, A., 2009. Sleep deprivation specifically impairs short-term olfactory
memory in Drosophila. *Sleep*, 32(11), pp.1417–1424.

Lim, J. & Dinges, D.F., 2010. A meta-analysis of the impact of short-term sleep deprivation
on cognitive variables. *Psychological bulletin*, 136(3), pp.375–89. Available at:
/pmc/articles/PMC3290659/?report=abstract.

Maimon, G., Straw, A.D. & Dickinson, M.H., 2008. A Simple Vision-Based Algorithm for
Decision Making in Flying Drosophila. *Current Biology*, 18(6), pp.464–470.

Parisky, K.M. et al., 2008. PDF Cells Are a GABA-Responsive Wake-Promoting Component of
the Drosophila Sleep Circuit. *Neuron*, 60(4), pp.672–682.

Paulk, A.C. et al., 2015. Closed-Loop Behavioral Control Increases Coherence in the Fly Brain.
The Journal of neuroscience : the official journal of the Society for Neuroscience, 35(28),
pp.10304–15. Available at: http://www.jneurosci.org/content/35/28/10304.abstract.

Seugnet, L. et al., 2008. D1 receptor activation in the mushroom bodies rescues sleep-loss-
induced learning impairments in Drosophila. *Current Biology*, 18(15), pp.1110–1117.

Seugnet, L. et al., 2011. Sleep deprivation during early-adult development results in long-
lasting learning deficits in adult Drosophila. *Sleep*, 34(2), pp.137–146.

Shaw, P.J. et al., 2000. Correlates of sleep and waking in *Drosophila melanogaster*. *Science*,
287(2000), pp.1834–1837.

Shaw, P.J. et al., 2002. Stress response genes protect against lethal effects of sleep
deprivation in Drosophila. 417(May), pp.287–291.

Stickgold, R., 2005. Sleep-dependent memory consolidation. *Nature*, 437(7063), pp.1272–
1278.

van Swinderen, B., 2007. Attention-like processes in Drosophila require short-term memory
genes. *Science*, 315(5818), pp.1590–1593.

van Swinderen, B. & Brembs, B., 2010. Attention-like deficit and hyperactivity in a
Drosophila memory mutant. *The Journal of neuroscience*, 30(3), pp.1003–1014.

Thimgan, M.S. et al., 2010. The Perilipin homologue, Lipid storage droplet 2, regulates sleep
homeostasis and prevents learning impairments following sleep loss. *PLoS Biology*,
8(8), pp.29–30.

Tononi, G. & Cirelli, C., 2013. Perspective Sleep and the Price of Plasticity : From Synaptic
and Cellular Homeostasis to Memory Consolidation and Integration. *Neuron*, 81(1),
pp.12–34. Available at: http://dx.doi.org/10.1016/j.neuron.2013.12.025.

Yap, M.H.W. et al., 2017. Oscillatory brain activity in spontaneous and induced sleep stages
in flies. *Nature Communications*, 8(1), p.1815. Available at:
http://www.nature.com/articles/s41467-017-02024-y.

Figure Legends:

Figure 1. A visual attention paradigm. A) Diagram of set up. Flies walk on a platform surrounded by a moat of water, with six LED panels forming walls onto which visual cues are presented. Visual attention is measured by fixation on either of two opposing stripes ('fixation targets') in the presence of a motion stimulus ('distractor'). B) Calculating fixation and optomotor responses (adapted from Colomb et al, 2013). Target fixation is measured by the stripe deviation (β), the angle between the fly's heading direction (e.g. red line from A to B) and the centre of the target stripe that is in the direction of movement (dotted black line).   Response to the motion stimulus (optomotor) is determined by the fly's angular velocity, i.e. the fly's turning angle (g)/second. C-E) Example traces of a fly's walking path in response to flickering target stripes (C), a motion stimulus (D) or competing stimuli (E). F,G) Quantification of target deviation (F) or optomotor response (G) in wild type Canton-S flies when different visual cues were presented: fixation targets alone ('no grating') or fixation targets combined with gratings of increasing salience (luminance contrast), or grating alone. n>10 flies for each condition, *p<0.05,**p,0.01,***p<0.001 by t-tests between adjacent conditions.

Figure 2. Sleep deprivation makes flies more distractible. A) Sleep deprivation protocol. Flies were sleep deprived for 24 hours and visual attention was tested the following day (day 1), and compared to control (non-sleep deprived flies). B) Flies subjected to the sleep deprivation protocol in (A) showed a sleep rebound (increase in sleep) measured by the difference in sleep compared to control (non-sleep deprived) flies. C) Example traces of walking behaviour in a control and sleep deprived fly responding to competing visual cues. D,E) Quantification of target deviation (D) or optomotor response (E) in control vs sleep deprived flies responding to target objects alone (target only), competing visual cues (objects and grating together, with three levels of grating salience), and grating alone (motion only). F) Attention returns to normal levels in flies rested for 24hrs following sleep deprivation (tested with objects and mid-salience grating). n= 60 flies in (A,B) n>16 flies for each condition in (D,E) and n>12 flies in (F). Red arrowheads indicate the grating contrast that was used for all subsequent experiments *p<0.05 (one-way Anova in F, t-test in E) and **p <0.01, ***p <0.001 by t test between control and sleep deprived. Error bars show the s.e.m.

Figure 3. Sleep deprivation increases distractibility to stationary objects. A) Diagram of visual stimuli, with flickering target stripes and non-flickering distractor stripes. B) Example traces of a flies responding to targets only, or targets with distractor stripes. C) Target deviation scores with increasing distractor salience (luminance contrast to background). D) Target deviation for controls and sleep deprived flies in the presence of distractor objects. n<20 flies in (C) and 50 flies in (D). **p<0.001 by t-tests between adjacent groups in (C) and sleep deprived and controls in (D).

Figure 4. Target deviation in the attention paradigm is not correlated with walking speed or sleep following sleep deprivation. A,B) Correlation of walking speed and target deviation in control (A) and sleep deprived (B) flies. n = 48 flies. Pearson's correlation coefficient ('R') were not significant (shown on graphs, alongside p values), indicating no correlations were found.

Figure 5. Starvation, heat, and mechanical stress effects on visual attention. Visual attention was measured by orientation towards 7hz flickering object ('targets') with a mid-luminance contrast grating distractor (see methods). A) Target deviation in flies that underwent heat stress (31 degrees) for 24 hours with or without sleep deprivation. B) Target deviation following 24hrs mechanical stimulation. C) Target deviation in flies that were starved and/or sleep deprived for 24hrs. "Control fed" vs "control starved" groups were not significantly different. n>17 flies in (A), n>28 flies in (B) n = 20 flies in (C). **p<0.01, ***p<0.001 by One-way ANOVA. Error bars indicate the s.e.m.

Figure 6. Activation of wake promoting circuits for 24 hours increases distractibility. A) Change in sleep caused by red-light exposure. n= 17 B) Change in attention (target deviation) following 24 hrs red light exposure. n = 10. C) Representative traces of a control fly (CS) compared to a sleep deprived fly (pdf-gal4>uas-chrimson) following red light exposure. t tests compared red-light activated to non-activated controls. **p=0.002, ***p<0.001, one-way ANOVA. Error bars show the s.e.m.

Figure 7. Increasing sleep improved attention in *dunce* mutants. A) Hourly sleep profiles

averaged across 24 hours for wild-type (Canton-S), dunce and rutabaga flies (n>50 flies). B)

Quantification of sleep duration averaged across the day and the night, from data in (A). C)

Sleep across 24 hours in wild-type flies, dunce and rutabaga mutants treated with the sleep

promoting agent, THIP. Wild-type sleep without THIP is shown for comparison (black trace).

n>20 flies. D) Quantification of day-time and night-time sleep in wild-type, dunce and

rutabaga flies with and without THIP. n>20 flies. E) Target deviation of wild-type (Canton-S)

flies on control (vehicle) or THIP (0.1mg/ml) food. THIP was administered for 2 days and flies

were removed from THIP 1 hr prior to the visual attention test. 'Within group' compared

flies before and after THIP administration (n=11 flies), and 'between group' compared age-

matched flies fed control (vehicle) or THIP food (n = 8 flies). F) Stripe deviation in dunce

mutants before and after 2 days of THIP (n>30 for dunce and CS, n> 16 for rutabaga).

*p<0.05, ***p<0.0001 by one-way ANOVA (A-D) and t-test (E-F). Error bars show the s.e.m.

Figure S1. Visual responses are modulated by flicker and grating frequency. (A) Visual

fixation in wild-type flies responding to fixation targets that were static (0 Hz) or flickering at

0.5 Hz, 3 Hz, 7 Hz, or 16 Hz. Lower target deviation indicates greater fixation. n = 14 flies for

each condition. Asterisks indicate significance between adjacent data points (2-way ANOVA

with Tukey's correction). (B) Optomotor response in wild-type flies responding to gratings

that were stationary (0 Hz) or rotating at 0.5 Hz, 3 Hz, 7 Hz, or 16 Hz. Higher optomotor

index (OI) indicates greater optomotor response. n = 10. Asterisks indicate significance

between adjacent points (A), significance via 2-way ANOVA with Tukey's correction. * p

< .05; ** p< .01; *** p< .001. Error bars show S.E.M.

Figure S2. Sleep deprivation effects on attention are independent of visual experience.

Target deviation during figure/ground discrimination in flies that were sleep deprived for 24

hrs in normal circadian conditions (light) or in 24 hrs of darkness (dark). The visual stimuli

were a 7 Hz flickering object ('target') and a mid luminance contrast grating (see methods).

n>19 flies per data set. *p=0.03,**p=0.01 by t-test. Error bars indicate the s.e.m.

Figure 1

A

target
platform
moat
target
motion
distractor

B

target
moat
motion
distractor
β
γ
A
B
C
target

β : stripe deviation
γ : turning angle

C Fixation
(targets only)

D Competing Stimuli

E Motion responses
(grating only)

F

increasing distractibility

Fixation
(Competing stimuli)

target deviation (degrees)

**
***
*

no
grating

grating
only

increasing grating salience

G

Motion responses
(Competing Stimuli)

optomotor (degrees/sec)

***
***

no
grating

grating
only

increasing grating salience

Figure 2

A Control

day 0   day 1   day 2   day 3

Attention
test

Sleep measured

Sleep deprivation

day 0   day 1   day 2   day 3

B Sleep rebound

Δday sleep duration (mins/hr)
Sleep Deprived - Control

day   night

***   **   n.s. n.s.   n.s. n.s.

day 1   day 2   day 3

C Control

Sleep Deprived

D Fixation on targets

more
distractible

target deviation (degrees)

more
focussed

**   **   ***

SD
control

target
only   motion
only

increasing grating salience

E Responses to motion

optomotor (degrees/sec)

*

SD
control

target
only   motion
only

increasing grating salience

F Recovery from SD

target deviation (degrees)

n.s.

*

Control   SD   SD
+24hr rest

Figure 3

A distractor / target (flickering) / target (flickering) / distractor

B targets only / targets + distractors

C target deviation (degrees)

(Increasing Distractor Salience)

D target deviation (degrees)

**

Control   Sleep Deprived

Figure 4

Figure 5

A **Heat Stress**

control (No SD, 25°C)

heated (No SD, 31°C)

heated (31°C) + SD (every 20 seconds, 24hrs)

n.s.

**

target deviation (degrees)

25°C   31°C   31°C
                 +
                 SD

B **Mechanical Stress**

control (unstimulated)

stimulated (every 10 seconds,
every second hr for 24 hrs)

n.s.

target deviation (degrees)

control   Stim

C **Starvation**

***

***

**

target deviation (degrees)

Control   SD   Control   SD
fed   fed   starved   starved

target

target

Figure 6

Activation of wake promoting circuits

Wild type and genetic controls

A ΔSleep duration

Δ sleep %
(Red light activated - Control)

*** ***   ***

CS (wildtype)
pdf-gal4/+
uas-chrimson/+
pdf-gal4>uas-chrimson

B Attention test
(Δdistractibility)

Δ target deviation (degrees)
(Red light activated - Control)

n.s.   **

CS (wildtype)
pdf-gal4/+
uas-chrimson/+
pdf-gal4>uas-chrimson

C

target

target

Control

Sleep deprived

Figure 7

Figure S1

A

Target deviation (deg)

20 — n.s.

10 — ***

0 — n.s.

0   0.5   3   7   16

Flicker frequency (Hz)

flickering target

flickering target

B

Optomotor (deg/sec)

300 — ***

200 — ***

100 — *

0 — n.s.

0   0.5   3   7   16

Grating frequency (Hz)

moving grating

Figure S2