FRMPD2: a novel GluN2A-interacting scaffold protein in synaptic excitatory transmission

Xi Lu[1], Haiqing Zhang[1], Xin Tian[1], Xuefeng Wang[1, 2*].

Affiliations:

1: Department of Neurology, Chongqing Key Laboratory of Neurology, The First Affiliated Hospital of Chongqing Medical University, 1 Youyi Road, Chongqing 400016, China

2: Center of Epilepsy, Beijing Institute for Brain Disorders, 10 Xitoutiao, Youanmen, Fengtai District, Beijing 100069, China.

*: The corresponding author: Xuefeng Wang, Email: wxfcqmu@163.com

Address correspondence to Xuefeng Wang, Department of Neurology, The First Affiliated Hospital, Chongqing Medical University, 1 Youyi Road, Chongqing 400016, China.

Abstract

The scaffold proteins FRMPD2 is localized at the basolateral membranes of polarized epithelial cells and associated with tight junction formation. But the expression and function of FRMPD2 in vivo remain unknown. Here, we found that postsynaptic FRMPD2 effectively alters the neuronal excitability by regulating NMDARs-component excitatory synaptic transmission in adult hippocampus. FRMPD2 anchors the NMDAR subunit GluN2A at the synapses without changing the total NMDAR expression. By using of Co-IP, Pull Down and SPR, the second PDZ (PDZ2) domain of FRMPD2 direct and binds to the C-terminus of GluN2A. Our study discover that a novel scaffold protein FRMPD2 regulates synaptic excitatory in a PDZ2 domain-dependent manner in adult hippocampus.

Keywords: excitatory synapses, NMDAR, PDZ domain, scaffold protein

Introduction

Scaffold proteins contain several modular protein interaction domains within their structure, and they bind to various interacting proteins, such as ion channels, neurotransmitter receptors, adhesion molecule, microtubule, filamentous actin (F-actin)(Anzai et al., 2002; Durand et al., 2012; Elias and Nicoll, 2007; Hruska-Hageman et al., 2002; Kim and Sheng, 2004; Sala et al., 2001; Schultz et al., 1998). They strongly contribute to the organization of synaptic signalling complexes, and the clustering of surface receptors and ion channels, and they participate in the dynamic trafficking of vesicle transport and coordinate cytoskeletal dynamics(Feng and Zhang, 2009). FRMPD2 (also named Gm626 in mice) is a novel scaffold protein that consists of an N-terminal KIND domain, a Ferm (band 4.1, ezrin, radixin, and moesin) domain and three C-terminal PDZ (PSD-95/Discs-large/ZO1) domains(Lipinski et al., 2012; Stenzel et al., 2009). FRMPD2 is reported to localize in the basolateral membranes of polarized epithelial cells and is associated with tight junction formation and immune response(Lipinski et al., 2012; Stenzel et al., 2009). However, there are few studies investigating FRMPD2 in vivo and the expression and function of FRMPD2 in nervous system remain unknown.

Inspired by multiple domains in its structure, FRMPD2 is predicted to play a critical role in neuronal excitation. PDZ domain-containing proteins are involved in neuronal excitatory by trafficking neurotransmitter receptors to plasma membrane(Kim and Sheng, 2004). Thus, FRMPD2 may be implicated in the regulation of neuronal excitation in a PDZ- dependent manner.

In this paper, we first reported that postsynaptic scaffold protein FRMPD2

activated neuronal excitability in adult hippocampus by mediating the

N-methyl-d-aspartic acid (NMDA) receptors-component excitatory synaptic

transmission. FRMPD2 anchored the GluN2A subunit of NMDA receptors

(NMDARs) to the synapses by directly binding the second PDZ (PDZ2) domain to

the C-terminus of GluN2A. Our study shed light on a novel scaffold protein

FRMPD2 mediate synaptic excitatory via directly anchoring synapses GluN2A

subunits.

Results

1. Expression patterns of the FRMPD2 protein.

The FRMPD2 protein was expressed in a restricted manner in some organs including the brain, eyes, testes and lungs (Fig 1A), and was enriched in a more widespread manner in various brain regions, including the cortex, hippocampus and cerebellum, in addition to the ependymal cells lining the ventricles (Fig 1B, C).

When immunofluorescence staining was performed in mouse cortex and the hippocampus Cornu Ammonis 1 (CA1) region, FRMPD2 was localized in microtubule-associated protein 2 (MAP2)-positive neuronal dendrites and another major scaffold protein, PSD95 (postsynaptic marker) (Fig 1D, E). Synaptosomes in the hippocampus were isolated by ultracentrifugation, and the FRMPD2 protein was detected in the postsynaptic density (PSD) fraction, but not in the presynaptic fraction or non-synaptic synaptosomal fraction (Fig 1F).

2. Effects of FRMPD2 on NMDA receptor-mediated excitatory synaptic transmission

PDZ domain-containing proteins are reported to be involved in neuronal excitatory (Kim and Sheng, 2004). We tested that whether postsynaptic FRMPD2 containing three PDZ domains affects neuronal excitatory. Whole cell patch-clamp electrophysiology was performed in CA1 pyramidal neurons in mouse hippocampal slices at 14 days after lentivirus vector (LV) injection of mice that was used to knockdown or overexpress FRMPD2. LV-encoding green fluorescent protein (GFP) was detected in the mouse hippocampus (Fig 2A). At 2 and 6 weeks post-LV injection,

the FRMPD2 protein levels of the hippocampus in FRMPD2 knockdown mice

(shFRMPD2) was almost 40 percent less than in scramble controls (shFRMPD2-Scr).

FRMPD2 protein levels were significantly higher in shFRMPD2-Res mice that were

injected with shRNA plus overexpression LV in the hippocampus than in FRMPD2

knockdown mice. The FRMPD2 protein in FRMPD2 overexpression mice

(FRMPD2-OE) was 1.9-2.1 times higher than in negative controls (FRMPD2-OE NC)

(Fig 2B). These findings suggest that FRMPD2 protein levels in the hippocampus

were effectively modulated by LV injection.

We firstly analyzed the effect of FRMPD2 on spontaneous action potential (AP)

in CA1 pyramidal neurons. The frequency of spontaneous AP in FRMPD2

knockdown mice reduced by 71 percent compared with the scramble control. AP

frequency of FRMPD2 overexpression mice was 2.28 times faster than in the

negative control (Fig 3A). To further explore the contribution of glutamate excitatory

receptors and/or inhibitory synaptic inputs contribute to the increase in neuronal

hyperactivity induced by FRMPD2, □-amino-3-hydroxy-5-methyl-4-isoxazole

propionic acid (AMPA) receptor (AMPARs) component or NMDAR component

mEPSCs, and miniature inhibitory postsynaptic currents (mIPSCs) of pyramidal

neurons in the CA1 region of the hippocampus were recorded. The two types of

ionotropic glutamate receptors, AMPARs and NMDARs, present at postsynaptic

membrane of excitatory synapses mediate almost all synaptic transmission in the

brain(Bredt and Nicoll, 2003; Valtschanoff and Weinberg, 2001). There was no

significant difference in AMPAR mEPSCs (Fig 3B) or mIPSCs (Fig 3C) after

FRMPD2 knockdown or overexpression. However, the mEPSC amplitude, but not

the frequency, of NMDAR mEPSCs was decreased in FRMPD2 knockdown mice

compared with scramble controls, and increased in FRMPD2 overexpression mice

compared with negative controls (Fig 4A).

To verify the changes of mEPSCs, NMDAR-mediated evoked EPSCs (eEPSCs)

were measured. Firstly, the input-output curve of NMDAR eEPSCs (Fig 4B) verified

that FRMPD2 enhanced the postsynaptic efficacy of NMDAR responses (Shen et al.,

2010). As shown in Fig 4C, the amplitudes of NMDAR currents were significantly

lower in the FRMPD2 knockdown than in the scramble controls. There was no

significant difference in evoked AMPAR currents between the FRMPD2 knockdown

and overexpression groups, in line with data of AMPAR component mEPSCs. In

particular, to determine whether GluN2A and/or GluN2B component currents

contributed to the observed changes in NMDAR eEPSCs, selective antagonist for

GluN2A (NVP-AAM077) and GluN2B (Ro25-6981) were used (Fig 4D). The

amplitude of GluN2A-sensitive currents of the FRMPD2 knockdowns was decreased

by 65 percent compared with the scramble groups, and amplitude of GluN2A currents

of overexpressing FRMPD2 mice was 2.38 times higher than of the negative controls.

Interesting, the GluN2B component-mediated currents were not changed after

FRMPD2 knockdown or overexpression. These results suggested that FRMPD2

strengthens GluN2A -mediated NMDARs synaptic transmission.

Because presynaptic release at postsynaptic glutamate receptors also contribute

to synaptic excitatory transmission, we analyzed the paired-pulse ratio (PPR), a

reliable index of the contribution of presynaptic vesicles release to excitatory

postsynaptic currents(Zucker and Regehr, 2002). There was no significant difference

in PPR values between the FRMPD2 knockdown and overexpression results and

those in the corresponding control groups (Fig 4E), indicating that presynaptic

vesicles release is not responsible for the FRMPD2-mediated changes in mEPSCs

3. Effects of FRMPD2 anchoring GluN2A at synapses.

To correlate electrophysiological changes with biochemical alterations in

synaptic NMDAR, we performed immunoblotting using hippocampal synaptosomal

and total lysates. As shown in Fig 5A, the synaptic GluN2A and GluN2B were

significantly fewer in the FRMPD2 knockdown mice than in the scramble mice, and

higher in the FRMPD2 overexpression mice than in the negative controls. The total

protein levels of GluN2A, GluN2B and GluN1 were not affected by FRMPD2.

Immunofluorescence staining was also used to test for co-localization between

FRMPD2 and GluN2A at postsynaptic membranes, which were marked by PSD-95 in

hippocampal CA1 areas (Fig 5B).

The extreme C-terminus of the NMDAR subunits provides a mechanism for

trafficking NMDAR subunits in postsynaptic membrane by interacting with PDZ

proteins (Bard et al., 2010; Kim and Sheng, 2004). We performed

co-immunoprecipitation (Co-IP) and GST pull-down assays to detect whether the

three PDZ domains of FRMPD2 interact with NMDAR subunits. In vivo,

co-immunoprecipitation of FRMPD2 with GluN1, GluN2A and GluN2B were

detected in the lysates of mice hippocampus (Fig 5C). A GST fusion protein bearing

the PDZ2 domain but not the first PDZ or third PDZ domain pulled down

endogenous GluN1, GluN2A and GluN2B from mice hippocampus in vivo. The

full-length GluN1, GluN2A and GluN2B proteins tagging with GFP were generated

in translation experiments in vitro, GST-PDZ2 pulled down full-length GluN2A and

GluN2B, but not GluN1 (Fig 5D). However, the GST-PDZ2 fusion protein did not

pull down GluN2A proteins from the cells transfected with a form of GFP-GluN2A

that lacked a set of ten C-terminal amino acids (Fig 5D).

The binding abilities of FRMPD2-isolated PDZ2 domain to the GluN2A and

GluN2B C-terminus was measured using surface plasmon resonance (SPR) analyses.

The binding capacity of the FRMPD2 PDZ2 domain to bind to the wild type GluN2A

C-terminus (KKMPSIESDV) and wild type GluN2B C-terminus (EKLSSIESDV)

yielded a K_D (dissociation constant) value of 0.31μM and 0.25mM, respectively (Fig

5E,F), whereas no binding was observed between the mutant GluN2A C terminus

(KKMPSIEADK) (Fig 5G) and the FRMPD2 PDZ2. These results supported the

direct interaction of C terminus of GluN2A to PDZ2 of FRMPD2.

To further explore potential sites for binding between GluN2A and FRMPD2,

we first obtained the crystal X-ray structure of the PDZ2 domain(Fig 6A) of the

mouse homologue of FRMPD2 (PDB ID: 5ZDS) (Table 1). The crystal structure was

solved by PHENIX with the template PDZ2 domain structure of tyrosine phosphatase

PTP-BL (PDB ID: 1VJ6) (Fig 6B) with 63% identity(Adams et al., 2010; Gianni et

al., 2006). In the template structure (Fig 6C), there was a peptide that is a C-terminal

(HSGSYLVTSV) of tumor suppressor protein APC, binding to β2 strand and α2 helix.

The residues V(0) and T(-2) of the peptide formed critical hydrophobic interactions

with residues L25, I27, L85, V29, and V65 of the template PDZ2 structure(Gianni et

al., 2006). These residues were highly conserved in the FRMPD2 PDZ2 crystal

structure, as shown by the alignment (Fig 6D).

The results shown so far reveal that FRMPD2 affects NMDAR-mediated

excitatory postsynaptic currents by anchoring GluN2A subunits at postsynaptic

membranes. This activity is achieved by an interaction of the FRMPD2 PDZ2 domain

to the GluN2A C-terminus.

Discussion

In our report, we identified a novel scaffold protein FRMPD2, and explored its

physiological contribution to the modulation of excitatory synaptic

electrophysiological function.

FRMPD2 was essential for NMDA-component, but not AMPA-component

excitatory transmission in adult hippocampus. It is seems that the binding of

glutamate receptors by FRMPD2 appears to change with glutamate receptor-specific

manner(Beique et al., 2006; Elias et al., 2008). Our results showed that synaptic

FRMPD2 mediated neurotransmission by trafficking GluN2A subunits on synapses in

adult hippocampus. NMDA receptors are composed of obligatory GluN1 subunit

co-assembled with GluN2A or GluN2B subunit (GluN1/GluN2A or GluN1/GluN2B).

The identity of the GluN2A and/or GluN2B subunits determines the biophysical and

pharmacological properties of NMDARs (Bard et al., 2010; Sultan et al., 2015). Our

results of Co-IP, GST-pull-down and SPR assays identified the second PDZ domain

of FRMPD2 as a novel binding partner of GluN2A and GluN2B *in vivo* and *in vitro*.

Although the FRMPD2 PDZ2 did not directly bind to GluN1, the obligatory GluN1

subunit of GluN1/2A complex was, expectedly, detectable by Co-IP of FRMPD2.

Meanwhile, the GluN1 protein of synaptosome was changed after FRMPD2

knockdown or overexpression, which was accompanied by the alteration of GluN2A

subunits. In addition, the binding capacity of PDZ2 to GluN2A subunit was 1000

times higher than that to GluN2B subunit. It is probably that the binding of PDZ2 to

GluN2B is flexible and instant. In addition, the GluN2B-containing NMDARs are

located at extrasynaptic sites and involved in neurotoxicity in adult hippocampal

neurons, GluN2A-containing NMDARs mainly resided at synaptic sites and involved

in neurotransmission (Parsons and Raymond, 2014).

In summary, the results presented here suggest a critical role of novel scaffold

protein FRMPD2 in regulating NMDA receptors anchoring, which is critical for

regulation of glutamatergic synapses transmission. The dysregulation of this scaffold

protein contributed to the impaired excitatory conductance.

Methods and materials

DNA constructs

To construct the following GST fusion proteins, the PDZ domain of mouse FRMPD2 (Uniprot: A0A140LI67) was subcloned into pGEX4T-1 with a GST tag: GST-PDZ1 (amino acids 786-872) and GST-PDZ2 (amino acids 934-1027), GST-PDZ3 (amino acids 1071-1159).

The lentivirus vector (LV) constructs were manufactured by Genechem Co. Biotechnology (Shanghai City, China). The mouse FRMPD2 (NCBI Gene: 268729) short hairpin RNA (shRNA) knockdown construct targeted 5'-3' CTGAATCTTTCTCTTGCAT and was subcloned it into the LV GV248 (GeneChem, China), which contained green fluorescent protein (GFP). A scramble shRNA (5'-3' CTTTTCTAACGTGTCGACT) was used as a negative control for the FRMPD2 shRNA. The complete mouse cDNA sequence for FRMPD2 (XM_017316244) (FRMPD2 overexpression, FRMPD2 OE) and the GFP sequence were inserted into the LV GV358 (GeneChem, China).

Antibodies

The following antibodies were used: FRMPD2 (rabbit, Sigma;rabbit, Origene), Flag(mouse, Sigma), GST(rabbit, Invitrogen), GFP (rabbit, Invitrogen), GluN2A(rabbit, Proteintech), GluN2B(mouse, Proteintech; rabbit, Abcam), GluN1(rabbit, Abcam), MAP2(guinea pig, SYSY), GFAP(mouse, Sigma), PSD-95(guinea pig, SYSY; mouse, Millipore), VGLUT1(guinea pig, SYSY),

synaptophysin(rabbit, Abcam), syntaxin 1a(rabbit, Abcam).

Cell line cultures and transfection

HEK293T and SH-SY5Y cells were maintained in DMEM (Gibco)

supplemented with 10% fetal bovine serum (Gibco), 100 U/mL penicillin and 100

μg/mL streptomycin (Gibco) at 37°C in 5% CO2.   The cells were transfected using

Lipofectamine 2000 (Invitrogen)(Klein et al., 2017). At 4 h after transfection, the

cells were rinsed with warm PBS and incubated in culture medium containing 10%

fetal bovine serum.

Intrahippocampal injection of LV

Adult C57/BL6 male mice were obtained from the Experimental Animal

Center of Chongqing Medical University. Intrahippocampal injection of LV was

performed using a stereotaxic apparatus (Stoelting Co. Ltd Wood Dale, IL, USA).

Briefly, the mice were anesthetized with pentobarbital (80mg/kg, intraperitoneal) and

placed on stereotaxic apparatus. The reference points were bregma for the

anterior-posterior axis, the midline for the medial-lateral axis and the dura mater for

the dorsal-ventral axis with a tooth bar. LV particles were injected bilaterally into the

dorsal hippocampus through a glass pipette (0.2 μl/min) attached to a glass

microsyringe(Shangguan et al., 2017). The mice were allowed to recover for 2 weeks

after the vector injections.

Patch clamp recordings

Brain slices were prepared as previously reported(Chen et al., 2006; Kim et al., 2001). Briefly, C57/BL6 mice were anesthetized with pentobarbital. Brain slices (300 μm) were prepared with a Leica (Germany) VP1200S Vibratome and incubated in artificial cerebral spinal fluid (ACSF) (119 mM NaCl, 26 mM $NaHCO_3$, 2.5 mM KCl, 1 mM $MgCl_2$, 1.25 mM NaH_2PO_4, 2 mM $CaCl_2$ and 25 mM glucose, pH 7.4, 310 mOsm) bubbled with 5% CO_2 and 95% O_2 for at least 1 h at room temperature before recording. Whole-cell recording was performed as described previously(Liu et al., 2005). Glass microelectrodes (Sutter, USA) were shaped by a pipette puller (P-97, Sutter, USA) to a final resistance of 3–5 MΩ when the pipette was filled with internal solution. A multi-clamp 700B amplifier (Axon, USA) was used for the recordings. Signals were sampled at 10 kHz and filtered at 2 kHz. A stable baseline was obtained for at least 5 min prior to recording and data were discarded when the access resistance (15–20 MΩ) was changed by 20% at the end of recording. Mini Analysis 6.0.1 (Synaptosoft) and Clamp Fit 10.3 software (Axon) were used to analyze the recorded data.

For action potential (AP) recording, glass pipettes were filled with the following internal solution: 17.5 mM KCl, 0.5 mM EGTA, 122.5 Mm K-gluconate, 10 mM HEPES, and 4 mM ATP, pH adjusted to 7.2 with KOH [20]. The internal solution used to record excitatory post-synaptic currents (EPSCs) contained: 17.5 mM CsCl, 10 mM HEPES, 4 mM ATP, 0.5 mM EGTA, 132.5 mM Cs-gluconate, and 5 mM QX-314. The internal solution used to record inhibitory post-synaptic currents (IPSCs)

contained: 100 mM CsCl, 1 mM $MgCl_2$, 1 mM EGTA, 30 mM

N-methyl-D-glucamine, 10 mM HEPES, 5 mM MgATP 0.5 mM Na_2GTP and 12 mM

phosphocreatine.

Spontaneous activity was assessed by observing the neuron under current clamp

for 1 min. Tetrodotoxin (TTX, 1 μM), bicuculline (10 μM),

6,7-dinitroquinoxaline-2,3(1H,4H)-dione (DNQX, 20 μM) and D-serine (10 μM)

were added to Mg-free-ACSF to record NMDAR mEPSCs that held at −60 mV. TTX

(1 μM), bicuculline (10 μM) and amino-5-phosphonopentanoate (AP-V, 40 μM) were

added to ACSF to record AMPA mEPSCs. TTX(1 μM), DNQX(20 μM) and APV(40

μM) were added to ACSF to record mIPSCs that held at −70 mV (Shangguan et al.,

2017).

Evoked EPSCs (eEPSCs) currents were generated with a 40μs pulse (0.1 Hz)

from a stimulated isolation unit controlled by an AMPI generator (Master-8, USA). A

bipolar stimulation electrode (FHC) was located ~100 μm rostral to the recording

electrode in the same layer(Chen et al., 2008; Tang et al., 2015). The eEPSCs at −70

mV holding potential in the present of PTX (100 μM) was considered as the

AMPAR-mediated current. Neurons were then held at +40 mV and the amplitude of

the eEPSCs at 50 ms post-stimulus was interpreted as the NMDAR-mediated

current(Etherton et al., 2009; Etherton et al., 2011). Input-output curve of NMDAR

eEPSCs were recorded in response to a series of stimulation intensities (pA).

Paired-pulse ratio (PPR) recordings were voltage-clamped at +40 mV and obtained

using a paired-pulse protocol of two stimuli at an inter-pulse interval of 50 ms. PPR

values were defined as the ratio of the second peak amplitude to the first peak

amplitude. The highly selective GluN2B subunit antagonist Ro25-6981 (Ro, 0.5μM)

was applied to assess GluN2A-sensitive eEPSCs, and the selective GluN2A

antagonist NVP-AAM077(NVP, 0.4μM) was added to record GluN2B component

eEPSCs(Bartlett et al., 2007).

Western blotting

Samples boiled with 4X sample buffer at 95°C for 5 min, and then separated on

8-10% SDS-PAGE gels and transferred onto polyvinylidene difluoride membranes

(Millipore). The target proteins were immunoblotted with primary antibodies

overnight at 4°C and then incubated with HRP-conjugated secondary antibodies. The

blots were imaged and quantified using a Fusion Imaging System. The quantitative

densitometric values of the proteins were normalized to that of GAPDH or beta-actin.

Immunofluorescence

For animal tissues, brain were post-fixed at 4°C for 24 h. Floating slices (50 mm

in thickness) were permeabilized with 0.5% Triton X-100 and blocked in goat serum.

The slices were then incubated with primary antibodies overnight at 4°C followed by

incubation with secondary fluorescent conjugated antibodies at room temperature for

2 h. Images were captured using laser-scanning confocal microscopy (Nikon A1+R

Microsystems) on an Olympus IX 70 inverted microscope (Olympus) equipped with

a Fluoview FVX confocal scanning head. Areas of overlap and fluorescence intensity

were analyzed using Image Pro Plus 6.0.

Sub-synaptic fractionation

Sub-synaptic fractionation was performed as previously described(Phillips et al.,

2001). To obtain synaptosomes, adult mouse hippocampus were homogenized in

solution buffer (0.32 M sucrose, 0.1 mM $CaCl_2$, 1 mM $MgCl_2$, 0.1 mM PMSF) and

brought to a final sucrose concentration of 1.25 M. Protease inhibitors were used in

all purification steps. The homogenate was overlaid with 1.0 M sucrose and

centrifuged (100,000 g, 3 h, 4°C). Synaptosomes were collected at the 1 M/1.25 M

interface.

For sub-synaptic fractionation, synaptosomes were sequentially extracted with

1% Triton X-100 first at pH 6.0 and then at pH 8.0. Synaptosomes (described above)

were diluted 1:10 with 0.1 mM $CaCl_2$ to a final concentration of 20 mM Tris, pH 6.0

and 1% Triton X-100. After the solution was incubated for 30 min and then

centrifuged 40,000 g for 30 min at 4°C. The supernatant (presynaptic fraction) were

precipitated with 10 volumes of acetone at −20°C and recovered by centrifugation

(20,000 g; 30 min, 4°C). The pellet containing insoluble postsynaptic density (PSD)

was resuspended in 10 volumes solution buffer (1% Triton X-100, 20 μM Tris, pH

8.0, 4°C), incubated for 30 min, and centrifuged (40,000 g, 30 min, 4°C).

GST-pull-down and co-immunoprecipitation assays.

The methods used to produce the GST fusion proteins in bacteria have been

previously reported (Anzai et al., 2002; Anzai et al., 2004). GST-pull-down assays

were carried out using a GST Protein Interaction Pull-Down Kit (Pierce) according to

the manufacturer's instructions. Briefly, the bacterial lysate containing the fusion

proteins was sonicated and centrifuged, and the supernatant was used for protein

binding assays. In vitro translation was facilitated using a plasmid carrying

full-length GluN2A, GluN2B or GluN1 (Addgene) that was transfected into

HEK293T cells. The translated total proteins obtained from cells and the GST fusion

proteins were incubated together with GST glutathione agarose resin. The protein

complexes were then eluted by glutathione and analyzed by Western blotting using

GST, GFP, GluN2A, GluN2B or GluN1 antibodies.

For co-immunoprecipitation assays, the extracts contained in the crude

synaptosomal fraction obtained from adult mouse hippocampal tissues were prepared

with anti-GluN2A, anti-GluN2B, anti-GluN1 or anti-FRMPD2 antibodies or IgG

(rabbit, Abcam) (Control) at 4°C for 12 h before they were incubated with protein

A/G agarose beads (Beyotime). The immunoprecipitated mixture was immunoblotted

with GluN2A, GluN2B, GluN1 or FRMPD2 antibodies.

Surface Plasmon Resonance (SPR) assays.

SPR biosensor analysis was used to verify binding between peptides and the

PDZ2 domain of FRMPD2 protein using a Biacore 8K instrument (GE Healthcare),

as previously described(Zhang et al., 2014). Proteins were immobilized on S CM5

chips using amine-coupling chemistry. Binding experiments were performed in

running buffer (10 mM PBS, pH 7, 0.05% P20) at 25°C with a flow rate of 30 μL/min.

Gradient concentrations of PDZ2 proteins were diluted to 50 ug/ml in sodium

acetate(10 mM, pH 5.5) and injected into the channel for 120 s, followed by

disassociation for 120 s. RU values were collected at 10 Hz and all the experimental

data were globally analyzed in a steady-state model within Biacore 8K Evaluation

software.

Study approval

All protocols of animal studies were approved by the Commission of Chongqing

Medical University for the ethics of experiments on animals and conducted in

accordance with international standards.

Crystallization

GST-PDZ2 (amino acids 934-1027) was expressed in Escherichia coli,

BL21(DE3) cells. Cells pellet was sonicated and centrifuged at 5000 rpm. The

supernatant was subjected to GST column purification (10 mM Tris, 10 mM GSH,

pH 7.4). Pooled fractions were dialyzed with thrombin and passed through a Q-HP

column (20 mM Tris, 0-1 M NaCl, 5% glycerol, pH 8.0). Finally, protein fractions

were further purified by Superdex 75 columns (20 mM Tris, 5% glycerol, 150 mM

NaCl, 2 mM DTT pH 7.4) for crystallization.

Screen kits, including the JCSG-CORE-I-IV (Qiagen) and HR2-144 Index

(Hampton), were used to perfrom crystallization screen with the hanging drop

method at 18 ℃. The protein (15 mg/ml) was mixed with an equivalent volume (1 µl)

of screen buffer. The crystal was grown in 20% PEG3350 and 0.2 M sodium

thiocyanate. A single crystal was transferred to the crystallization buffer containing an

additional 20% glycerol and flash frozen with liquid nitrogen. The crystal diffracted

well at BL19U (Shanghai Synchrotron Radiation Facility, SSRF, China). In all,

images for 180° were collected. The data were processed by HKL3000 (Table 1). The

crystal structure was refined by PHENIX and Coot. The structure was refined to 1.80

Å with R factor 19.76% and R free 24.76%.

Statistical analysis

All averaged data are presented as the means ± SD, and all graphs were prepared

using GraphPad Prism 4 software (La Jolla, CA). For independent-samples, Student's

t-test was used. Repeated measures ANOVA followed by post hoc t-tests was used to

measure differences between two groups at multiple time points. All tests were

two-sided. n indicates the number of cells/slices or independent experiments and was

used to calculate the degree of freedom. All samples included in each experiment

were analyzed in triplicate.

Acknowledgements

402   This work was supported by grants from the National Natural Science

403 Foundation of China (No. 81271445, 81471319, 81301109, and 81301110) and the

404 National Clinical Key Specialty Construction Foundation of China.

Author contributions

406 X. L. and XF. W. conceived and designed the experiments. HQ. Z. and X.

407 L. performed the experiments. X. L. collected and analyzed the data. X. L. and X.T.

408 wrote the manuscript. All authors contributed to preparation of the paper and

409 approved the final contributions.

410 The authors have declared that no conflict of interest exists.

References

Adams, P.D., Afonine, P.V., Bunkoczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J., Hung, L.W., Kapral, G.J., Grosse-Kunstleve, R.W., et al. (2010). PHENIX: a comprehensive Python-based system for macromolecular structure solution. Acta crystallographica Section D, Biological crystallography 66, 213-221.

Anzai, N., Deval, E., Schaefer, L., Friend, V., Lazdunski, M., and Lingueglia, E. (2002). The multivalent PDZ domain-containing protein CIPP is a partner of acid-sensing ion channel 3 in sensory neurons. The Journal of biological chemistry 277, 16655-16661.

Anzai, N., Miyazaki, H., Noshiro, R., Khamdang, S., Chairoungdua, A., Shin, H.J., Enomoto, A., Sakamoto, S., Hirata, T., Tomita, K., et al. (2004). The multivalent PDZ domain-containing protein PDZK1 regulates transport activity of renal urate-anion exchanger URAT1 via its C terminus. The Journal of biological chemistry 279, 45942-45950.

Bard, L., Sainlos, M., Bouchet, D., Cousins, S., Mikasova, L., Breillat, C., Stephenson, F.A., Imperiali, B., Choquet, D., and Groc, L. (2010). Dynamic and specific interaction between synaptic NR2-NMDA receptor and PDZ proteins. Proceedings of the National Academy of Sciences of the United States of America 107, 19561-19566.

Bartlett, T.E., Bannister, N.J., Collett, V.J., Dargan, S.L., Massey, P.V., Bortolotto, Z.A., Fitzjohn, S.M., Bashir, Z.I., Collingridge, G.L., and Lodge, D. (2007). Differential roles of NR2A and NR2B-containing NMDA receptors in LTP and LTD in the CA1 region of two-week old rat hippocampus. Neuropharmacology 52, 60-70.

Beique, J.C., Lin, D.T., Kang, M.G., Aizawa, H., Takamiya, K., and Huganir, R.L. (2006). Synapse-specific regulation of AMPA receptor function by PSD-95. Proceedings of the National Academy of Sciences of the United States of America 103, 19535-19540.

Bredt, D.S., and Nicoll, R.A. (2003). AMPA receptor trafficking at excitatory synapses. Neuron 40, 361-379.

Chen, G., Chen, P., Tan, H., Ma, D., Dou, F., Feng, J., and Yan, Z. (2008). Regulation of the NMDA receptor-mediated synaptic response by acetylcholinesterase inhibitors and its impairment in an animal model of Alzheimer's disease. Neurobiology of aging 29, 1795-1804.

Chen, G., Kittler, J.T., Moss, S.J., and Yan, Z. (2006). Dopamine D3 receptors regulate GABAA receptor function through a phospho-dependent endocytosis mechanism in nucleus accumbens. The Journal of neuroscience : the official journal of the Society for Neuroscience 26, 2513-2521.

Durand, C.M., Perroy, J., Loll, F., Perrais, D., Fagni, L., Bourgeron, T., Montcouquiol, M., and Sans, N. (2012). SHANK3 mutations identified in autism lead to modification of dendritic spine morphology via an actin-dependent mechanism. Molecular psychiatry 17, 71-84.

Elias, G.M., Elias, L.A., Apostolides, P.F., Kriegstein, A.R., and Nicoll, R.A. (2008). Differential trafficking of AMPA and NMDA receptors by SAP102 and PSD-95 underlies synapse development. Proceedings of the National Academy of Sciences of the United States of America 105, 20953-20958.

Elias, G.M., and Nicoll, R.A. (2007). Synaptic trafficking of glutamate receptors by MAGUK scaffolding proteins. Trends in cell biology 17, 343-352.

Etherton, M.R., Blaiss, C.A., Powell, C.M., and Sudhof, T.C. (2009). Mouse neurexin-1alpha deletion causes correlated electrophysiological and behavioral changes consistent with cognitive impairments. Proceedings of the National Academy of Sciences of the United States of America 106, 17998-18003.

Etherton, M.R., Tabuchi, K., Sharma, M., Ko, J., and Sudhof, T.C. (2011). An autism-associated point

mutation in the neuroligin cytoplasmic tail selectively impairs AMPA receptor-mediated synaptic
transmission in hippocampus. The EMBO journal *30*, 2908-2919.
Feng, W., and Zhang, M. (2009). Organization and dynamics of PDZ-domain-related supramodules in
the postsynaptic density. Nature reviews Neuroscience *10*, 87-99.
Gianni, S., Walma, T., Arcovito, A., Calosci, N., Bellelli, A., Engstrom, A., Travaglini-Allocatelli, C.,
Brunori, M., Jemth, P., and Vuister, G.W. (2006). Demonstration of long-range interactions in a PDZ
domain by NMR, kinetics, and protein engineering. Structure (London, England : 1993) *14*, 1801-1809.
Hruska-Hageman, A.M., Wemmie, J.A., Price, M.P., and Welsh, M.J. (2002). Interaction of the synaptic
protein PICK1 (protein interacting with C kinase 1) with the non-voltage gated sodium channels BNC1
(brain Na+ channel 1) and ASIC (acid-sensing ion channel). The Biochemical journal *361*, 443-450.
Kim, C.H., Chung, H.J., Lee, H.K., and Huganir, R.L. (2001). Interaction of the AMPA receptor subunit
GluR2/3 with PDZ domains regulates hippocampal long-term depression. Proceedings of the National
Academy of Sciences of the United States of America *98*, 11725-11730.
Kim, E., and Sheng, M. (2004). PDZ domain proteins of synapses. Nature reviews Neuroscience *5*,
771-781.
Klein, Z.A., Takahashi, H., Ma, M., Stagi, M., Zhou, M., Lam, T.T., and Strittmatter, S.M. (2017). Loss of
TMEM106B Ameliorates Lysosomal and Frontotemporal Dementia-Related Phenotypes in
Progranulin-Deficient Mice. Neuron *95*, 281-296.e286.
Lipinski, S., Grabe, N., Jacobs, G., Billmann-Born, S., Till, A., Hasler, R., Aden, K., Paulsen, M., Arlt, A.,
Kraemer, L., *et al.* (2012). RNAi screening identifies mediators of NOD2 signaling: implications for
spatial specificity of MDP recognition. Proceedings of the National Academy of Sciences of the United
States of America *109*, 21426-21431.
Liu, Q.S., Pu, L., and Poo, M.M. (2005). Repeated cocaine exposure in vivo facilitates LTP induction in
midbrain dopamine neurons. Nature *437*, 1027-1031.
Parsons, M.P., and Raymond, L.A. (2014). Extrasynaptic NMDA receptor involvement in central
nervous system disorders. Neuron *82*, 279-293.
Phillips, G.R., Huang, J.K., Wang, Y., Tanaka, H., Shapiro, L., Zhang, W., Shan, W.S., Arndt, K., Frank, M.,
Gordon, R.E., *et al.* (2001). The presynaptic particle web: ultrastructure, composition, dissolution, and
reconstitution. Neuron *32*, 63-77.
Sala, C., Piech, V., Wilson, N.R., Passafaro, M., Liu, G., and Sheng, M. (2001). Regulation of dendritic
spine morphology and synaptic function by Shank and Homer. Neuron *31*, 115-130.
Schultz, J., Hoffmuller, U., Krause, G., Ashurst, J., Macias, M.J., Schmieder, P., Schneider-Mergener, J.,
and Oschkinat, H. (1998). Specific interactions between the syntrophin PDZ domain and voltage-gated
sodium channels. Nature structural biology *5*, 19-24.
Shangguan, Y., Xu, X., Ganbat, B., Li, Y., Wang, W., Yang, Y., Lu, X., Du, C., Tian, X., and Wang, X. (2017).
CNTNAP4 Impacts Epilepsy Through GABAA Receptors Regulation: Evidence From Temporal Lobe
Epilepsy Patients and Mouse Models. Cerebral cortex (New York, NY : 1991), 1-14.
Shen, G., Van Sickle, B.J., and Tietz, E.I. (2010). Calcium/calmodulin-dependent protein kinase II
mediates hippocampal glutamatergic plasticity during benzodiazepine withdrawal.
Neuropsychopharmacology : official publication of the American College of
Neuropsychopharmacology *35*, 1897-1909.
Stenzel, N., Fetzer, C.P., Heumann, R., and Erdmann, K.S. (2009). PDZ-domain-directed basolateral
targeting of the peripheral membrane protein FRMPD2 in epithelial cells. Journal of cell science *122*,
3374-3384.

Sultan, S., Li, L., Moss, J., Petrelli, F., Casse, F., Gebara, E., Lopatar, J., Pfrieger, F.W., Bezzi, P.,

Bischofberger, J., *et al.* (2015). Synaptic Integration of Adult-Born Hippocampal Neurons Is Locally

Controlled by Astrocytes. Neuron *88*, 957-972.

Tang, B., Luo, D., Yang, J., Xu, X.Y., Zhu, B.L., Wang, X.F., Yan, Z., and Chen, G.J. (2015). Modulation of

AMPA receptor mediated current by nicotinic acetylcholine receptor in layer I neurons of rat

prefrontal cortex. Scientific reports *5*, 14099.

Valtschanoff, J.G., and Weinberg, R.J. (2001). Laminar organization of the NMDA receptor complex

within the postsynaptic density. The Journal of neuroscience : the official journal of the Society for

Neuroscience *21*, 1211-1217.

Zhang, C., Wan, X., Zheng, X., Shao, X., Liu, Q., Zhang, Q., and Qian, Y. (2014). Dual-functional

nanoparticles targeting amyloid plaques in the brains of Alzheimer's disease mice. Biomaterials *35*,

456-465.

Zucker, R.S., and Regehr, W.G. (2002). Short-term synaptic plasticity. Annual review of physiology *64*,

355-405.

Figures

Fig 1. Expression patterns of the FRMPD2 protein.

A, FRMPD2 protein is located in the mouse brain, eyes, testes and lungs. B, The

brain-specific expression of the FRMPD2 protein was revealed in adult mouse tissue

homogenates. C, The FRMPD2 protein has a widespread distribution pattern in adult

mouse brain slices. Bar, 500 μm. D, FRMPD2 was present in MAP2-positive

neuronal dendrites but not in GFAP-positive glial cells as determined by

immunofluorescence staining. Bar, 40 μm. E, FRMPD2 co-localized with the

postsynaptic marker PSD95 but not with the presynaptic marker VGLUT1 in the

mouse hippocampus and cortex. Bar, 40 μm. F, Immunoblot of the synapse fraction

(Syn) showing that FRMPD2 was preferentially distributed in the postsynaptic

fraction (Post) rather than in the presynaptic (Pre) or non-synaptic synaptosomal

proteins (NSSP) fraction. Anti-synapphysin, anti-syntaxin 1a and anti-PSD95 were

used as markers for ultrasynaptic fraction to verify that the purity of the preparation

was higher than 90%. The blots were representative of 4-6 analyses.

Fig 2. Alteration of FRMPD2 protein levels in the hippocampus by LV injection.

A, Lentiviral vector (LV)-encoding green fluorescent protein (GFP) was detected in

the mouse hippocampus following the injection of LV. Bar, 250 μm. B, Hippocampal

FRMPD2 protein expression levels were detected in FRMPD2 knockdown or

overexpression mice on the 2 and 6 weeks after LV injection. Data were shown as

mean±SD and analyzed by Student's t test, n=5 mice, *p< 0.05.

Fig 3. Effects of FRMPD2 on electrophysiology of CA1 neurons in mouse hippocampal brain slices.

A, The frequency of spontaneous AP was compared after FRMPD2 knockdown or overexpression. Bar, 1 s, 25 mV. B, C, FRMPD2 had no significant effect on the amplitude or frequency of AMPAR mEPSCs (B) and mIPSCs (C). Bar, 1 s, 20 pA. Data were shown as mean±SD and analyzed by Student's *t* test, n=6-7 neurons of 3 mice, **p< 0.01.

Fig 4. Effects of FRMPD2 on NMDA receptor-mediated excitatory synaptic

transmission.

A, Quantitative analysis of amplitude and frequency of NMDAR mEPSCs in CA1

pyramidal cells of hippocampus after FRMPD2 knockdown or overexpression. Bar, 1

s, 20 pA. B, The input-output curves of NMDAR eEPSCs were recorded with gradual

increasing intensities. Bar, 25 ms, 50 pA. C, Amplitude of NMDAR, AMPAR

component eEPSCs and AMPA/NMDA ratio were compared. Bar, 25 ms, 25 pA. D,

Quantitative analysis of the amplitude of GluN2A- and GluN2B-sensitive eEPSCs.

Bar, 25 ms, 25 pA. E, Knocking down or overexpressing FRMPD2 did not alter the

paired-pulse ratio (PPR) values. Bar, 25 ms, 50 pA. Data of were presented as the

mean±SD or median with interquartile range. Data of A, C, D, E, were analyzed by

Student's t test, data of B was analyzed by repeated measures ANOVA, n=5-7

neurons from 3 mice. **p<0.01. *p<0.05.

Fig 5. Effects of FRMPD2 anchoring GluN2A at synapses.

A, Quantitative analysis of GluN2A, GluN2B and GluN1 expressions of total

hippocampus tissue and synaptic homogenates after FRMPD2 knockdown or

overexpression. The data were shown as the mean±SD and analyzed by Student's *t*

test, *p< 0.05. B, Immunofluorescence staining in the mouse hippocampus CA1 area

showed that FRMPD2 co-localizes with GluN2A at the postsynaptic density, which

was marked by PSD95. Bar, 40 μm. C, FRMPD2 and GluN2A, GluN2B and GluN1

were co-immunoprecipitated from mouse hippocampal tissues. D, GST pull-down

assay showing that the GST-PDZ2 fusion protein but not the PDZ1 or PDZ3 fusion

proteins pulled down endogenous GluN2A from mouse hippocampal tissues, and

exogenous full-length GFP-GluN2A from HEK293 cells but did not pull down the

deletion variant GFP-GluN2A from HEK293 cells transfected with GFP-GluN2A that

lacked ten amino acids at the C-terminus. E-G, The ability of the isolated FRMPD2

PDZ2 domain to bind to the GluN2A and GluN2B C-terminus was measured using

SPR. Peptides sequences of C-termini of wild type GluN2A (E), wild type GluN2B

(F) and mutant GluN2A (G) were used in this study. Sensorgrams were shown to

illustrate the binding of increasing concentrations of the PDZ2 domain to peptides

and steady-state equilibrium responses from the sensorgrams were shown plotted

against the injected PDZ2 concentrations. K_D: dissociation constant.

Fig 6. Crystal structure of FRMPD2 PDZ2 domain.

A, The crystal structure of a compact globular fold in the ligand-free FRMPD2 PDZ2

domain (PDB ID: 5ZDS) was shown. B, The crystal structure of the FRMPD2 PDZ2

domain (green) was shown aligned to that of the PDZ2-peptide template structure

(cyan and magenta) (PDB ID: 1VJ6). C, A template of PDZ2-peptide (cyan and

magenta) interaction (PDB ID: 1VJ6). D, The binding pocket alignment between

FRMPD2 PDZ2 crystal structure (green) and PDZ2 template (cyan).

Table 1. Data collection and refinement statistics of PDZ2 of FRMPD2.

	PDZ2
Data collection	
Wavelength(Å)	0.9786
Resolution range(Å)	50-1.58 (1.64-1.58)[a]
Space group	P 2_1 2_1 2
Unit cell dimensions	
a, b, c(Å)	32.200, 48.179, 56.845
α, β, γ(°)	90, 90, 90
No. of reflections	80345
No. unique reflections	12699 (1248)
Multiplicity	6.3 (6.4)
Completeness (%)	99.7 (100)
Mean I/σ (I)	22.52(7.70)
R-meas	0.065(0.268)
Refinement	
Resolution range(Å)	50-1.8 (1.86-1.8)
No. of reflections	8610 (849)
No. of reflections used for R-free	860 (85)
R-work	0.1980 (0.2202)
R-free	0.2347 (0.2630)
No. of non-hydrogen atoms	810

Macromolecules	658
Protein residues	88
R.m.s.d.	
Bond lengths(Å)	0.005
Bond angles(°)	0.83
Ramachandranplot(%)	
Allowed regions	1.1
Outliers	0
Average B-factor	13.71
Macromolecules	11.25
Solvent	24.37

595 [a]Statistics for the highest-resolution shell are shown in parentheses. No., number,

596 r.m.s.d., root mean squared deviation.