FRMPD2: a novel GluN2A-interacting scaffold protein in synaptic excitatory transmission

Xi Lu[1], Haiqing Zhang[1], Yun Li[1], Ruijiao Zhou[1], Miaoqing He[1], Yuanlin Ma[1], Xin Tian[1], Xuefeng Wang[1, 2*].

Affiliations:

1: Department of Neurology, Chongqing Key Laboratory of Neurology, The First Affiliated Hospital of Chongqing Medical University, 1 Youyi Road, Chongqing 400016, China

2: Center of Epilepsy, Beijing Institute for Brain Disorders, 10 Xitoutiao, Youanmen, Fengtai District, Beijing 100069, China.

*: The corresponding author: Xuefeng Wang, Email: wxfcqmu@163.com

Address correspondence to Xuefeng Wang, Department of Neurology, The First Affiliated Hospital, Chongqing Medical University, 1 Youyi Road, Chongqing 400016, China.

Abstract

The scaffold proteins FRMPD2 is localized at the basolateral membranes of polarized epithelial cells and associated with tight junction formation. But the expression and function of FRMPD2 in vivo remain unknown. Here, we found that postsynaptic FRMPD2 effectively alters the neuronal excitability by regulating NMDARs-component excitatory synaptic transmission in adult hippocampus. FRMPD2 anchors the NMDAR subunit GluN2A at the synapses without changing the total NMDAR expression. By using of Co-IP, Pull Down and SPR, the second PDZ (PDZ2) domain of FRMPD2 direct and binds to the C-terminus of GluN2A. Moreover, Ferm domain of FRMPD2 directly interacts with actin filaments and modifies excitatory synapse morphogenesis, further strengthen neuronal excitatory transmission. In particular, FRMPD2 mediates the behavioral and field potential phenotypes of epilepsy, a brain disorder of excitatory/inhibitory imbalance. Our study discover that a novel scaffold protein FRMPD2 regulates synaptic excitatory in a PDZ2 domain- and Ferm domain-dependent manner in adult hippocampus.

Keywords: dendritic spines/ excitatory synapses/ NMDAR/ PDZ domain/ scaffold protein

Introduction

Scaffold proteins contain several modular protein interaction domains within their structure, and they bind to various interacting proteins, such as ion channels, neurotransmitter receptors, adhesion molecule, microtubule, filamentous actin (F-actin)(Anzai et al, 2002; Durand et al, 2012; Elias et al, 2007; Hruska-Hageman et al, 2002; Kim et al, 2004; Sala et al, 2001; Schultz et al, 1998). They strongly contribute to the organization of synaptic signalling complexes, and the clustering of surface receptors and ion channels, and they participate in the dynamic trafficking of vesicle transport and coordinate cytoskeletal dynamics(Feng et al, 2009). FRMPD2 (also named Gm626) is a novel scaffold protein that consists of an N-terminal KIND domain, a Ferm (band 4.1, ezrin, radixin, and moesin) domain and three C-terminal PDZ (PSD-95/Discs-large/ZO1) domains(Lipinski et al, 2012; Stenzel et al, 2009). FRMPD2 is reported to localize in the basolateral membranes of polarized epithelial cells and is associated with tight junction formation and immune response(Lipinski et al, 2012; Stenzel et al, 2009). However, there are few studies investigating FRMPD2 in vivo and the expression and function of FRMPD2 in nervous system remain unknown.

Inspired by multiple domains in its structure, FRMPD2 is predicted to play a critical role in neuronal excitation. PDZ domain-containing proteins are involved in neuronal excitatory by trafficking neurotransmitter receptors to plasma membrane(Kim & Sheng, 2004). Ferm domain-containing proteins are reported to recruit actin-regulatory proteins(Yang et al, 2009), and the F-actin cytoskeleton is

essential for neuronal morphogenesis and synaptic connectivity(Bourne et al, 2008;

Lei et al, 2001). Thus, FRMPD2 may be implicated in the regulation of neuronal

excitation or morphogenesis in a PDZ- or Ferm-dependent manner.

In this paper, we first reported that postsynaptic scaffold protein FRMPD2

activated neuronal excitability in adult hippocampus by mediating the

N-methyl-d-aspartic acid (NMDA) receptors-component excitatory synaptic

transmission. FRMPD2 anchored the GluN2A subunit of NMDA receptors

(NMDARs) to the synapses by directly binding the second PDZ (PDZ2) domain to

the C-terminus of GluN2A. FRMPD2 further regulated maturation of dendritic spines.

At last, FRMPD2 mediated the susceptibility of epilepsy, a brain disorder of

excitability(Kwan et al, 2011). Our study shed light on a novel scaffold protein

FRMPD2 mediate synaptic excitatory and modulate epileptic seizures via directly

anchoring synapses GluN2A subunits.

Results

1. Expression patterns of the FRMPD2 protein.

The FRMPD2 protein was expressed in a restricted manner in some organs including the brain, eyes, testes and lungs (Supplementary Fig 1A), and was enriched in a more widespread manner in various brain regions, including the cortex, hippocampus and cerebellum, in addition to the ependymal cells lining the ventricles (Supplementary Fig 1B, C).

When immunofluorescence staining was performed in mouse cortex and the hippocampus Cornu Ammonis 1 (CA1) region, FRMPD2 was localized in microtubule-associated protein 2 (MAP2)-positive neuronal dendrites and another major scaffold protein, PSD95 (postsynaptic marker) (Supplementary Fig 1D, E). Synaptosomes in the hippocampus were isolated by ultracentrifugation, and the FRMPD2 protein was detected in the postsynaptic density (PSD) fraction, but not in the presynaptic fraction or non-synaptic synaptosomal fraction (Supplementary Fig 1F).

2. Effects of FRMPD2 on NMDA receptor-mediated excitatory synaptic transmission

PDZ domain-containing proteins are involved in neuronal excitatory by interacting and trafficking neurotransmitter receptors(Kim & Sheng, 2004). We tested that whether postsynaptic FRMPD2 containing three PDZ domains affects neuronal excitatory. Whole cell patch-clamp electrophysiology was performed in CA1 pyramidal neurons in mouse hippocampal slices at 14 days after lentivirus vector (LV)

injection of mice that was used to knockdown or overexpress FRMPD2. LV-encoding

green fluorescent protein (GFP) was detected in the mouse hippocampus

(Supplementary Fig 2A). At 2 and 6 weeks post-LV injection, the FRMPD2 protein

levels of the hippocampus in FRMPD2 knockdown mice (shFRMPD2) was almost 40

percent less than in scramble controls (shFRMPD2-Scr). FRMPD2 protein levels

were significantly higher in shFRMPD2-Res mice that were injected with shRNA

plus overexpression LV in the hippocampus than in FRMPD2 knockdown mice. The

FRMPD2 protein in FRMPD2 overexpression mice (FRMPD2-OE) was 1.9-2.1 times

higher than in negative controls (FRMPD2-OE NC) (Supplementary Fig 2B). These

findings suggest that FRMPD2 protein levels in the hippocampus were effectively

modulated by LV injection.

We firstly analyzed the effect of FRMPD2 on spontaneous action potential (AP)

in CA1 pyramidal neurons. The frequency of spontaneous AP in FRMPD2

knockdown mice reduced by 71 percent compared with the scramble control. AP

frequency of FRMPD2 overexpression mice was 2.28 times faster than in the

negative control (Supplementary Fig 3A). To further explore the contribution of

glutamate excitatory receptors and/or inhibitory synaptic inputs contribute to the

increase   in   neuronal   hyperactivity   induced   by   FRMPD2,

α-amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) receptor (AMPARs)

component or NMDAR component mEPSCs, and miniature inhibitory postsynaptic

currents (mIPSCs) of pyramidal neurons in the CA1 region of the hippocampus were

recorded. The two types of ionotropic glutamate receptors, AMPARs and NMDARs,

present at postsynaptic membrane of excitatory synapses mediate almost all synaptic

transmission in the brain(Bredt et al, 2003; Valtschanoff et al, 2001). There was no

significant difference in AMPAR mEPSCs (Supplementary Fig 3B) or mIPSCs

(Supplementary Fig 3C) after FRMPD2 knockdown or overexpression. However, the

mEPSC amplitude, but not the frequency, of NMDAR mEPSCs was decreased in

FRMPD2 knockdown mice compared with scramble controls, and increased in

FRMPD2 overexpression mice compared with negative controls (Fig 1A).

To verify the changes of mEPSCs, NMDAR-mediated evoked EPSCs (eEPSCs)

were measured. Firstly, the input-output curve of NMDAR eEPSCs (Fig 1B) verified

that FRMPD2 enhanced the postsynaptic efficacy of NMDAR responses (Shen et al,

2010). As shown in Fig 1C, the amplitudes of NMDAR currents were significantly

lower in the FRMPD2 knockdown than in the scramble controls. There was no

significant difference in evoked AMPAR currents between the FRMPD2 knockdown

and overexpression groups, in line with data of AMPAR component mEPSCs. In

particular, to determine whether GluN2A and/or GluN2B component currents

contributed to the observed changes in NMDAR eEPSCs, selective antagonist for

GluN2A (NVP-AAM077) and GluN2B (Ro25-6981) were used (Fig 1D). The

amplitude of GluN2A-sensitive currents of the FRMPD2 knockdowns was decreased

by 65 percent compared with the scramble groups, and amplitude of GluN2A currents

of overexpressing FRMPD2 mice was 2.38 times higher than of the negative controls.

Interesting, the GluN2B component-mediated currents were not changed after

FRMPD2 knockdown or overexpression. These results suggested that FRMPD2

strengthens GluN2A -mediated NMDARs synaptic transmission.

Because presynaptic release at postsynaptic glutamate receptors also contribute

to synaptic excitatory transmission, we analyzed the paired-pulse ratio (PPR), a

reliable index of the contribution of presynaptic vesicles release to excitatory

postsynaptic currents(Zucker et al, 2002). There was no significant difference in PPR

values between the FRMPD2 knockdown and overexpression results and those in the

corresponding control groups (Fig 1E), indicating that presynaptic vesicles release is

not responsible for the FRMPD2-mediated changes in mEPSCs

3. Effects of FRMPD2 anchoring GluN2A at synapses.

To correlate electrophysiological changes with biochemical alterations in

synaptic NMDAR, we performed immunoblotting using hippocampal synaptosomal

and total lysates. As shown in Fig 2A, the synaptic GluN1 and GluN2A were

significantly fewer in the FRMPD2 knockdown mice than in the scramble mice, and

higher in the FRMPD2 overexpression mice than in the negative controls. The

synaptic GluN2B and total protein levels of GluN2A, GluN2B and GluN1 were not

affected by FRMPD2. Immunofluorescence staining was also used to test for

co-localization between FRMPD2 and GluN2A at postsynaptic membranes, which

were marked by PSD-95 in hippocampal CA1 areas (Fig 2B).

The extreme C-terminus of the NMDAR subunits provides a mechanism for

trafficking NMDAR subunits in postsynaptic membrane by interacting with PDZ

proteins (Bard et al, 2010; Kim & Sheng, 2004). We performed

co-immunoprecipitation (Co-IP) and GST pull-down assays to detect whether the

three PDZ domains of FRMPD2 interact with NMDAR subunits. In vivo,

co-immunoprecipitation of FRMPD2 with GluN1 and GluN2A, but not GluN2B were

detected in the lysates of mice hippocampus (Fig 2C). A GST fusion protein bearing

the PDZ2 domain but not the first PDZ or third PDZ domain pulled down

endogenous GluN1 and GluN2A, but not GluN2B from mice hippocampus in vivo.

The full-length GluN1, GluN2A and GluN2B proteins tagging with GFP were

generated in translation experiments in vitro, GST-PDZ2 pulled down full-length

GluN2A, but not GluN1 and GluN2B (Fig 2D). However, the GST-PDZ2 fusion

protein did not pull down GluN2A proteins from the cells transfected with a form of

GFP-GluN2A that lacked a set of ten C-terminal amino acids (Fig 2D).

The binding abilities of FRMPD2-isolated PDZ2 domain to the GluN2A and

GluN2B C-terminus was measured using surface plasmon resonance (SPR) analyses.

The binding capacity of the FRMPD2 PDZ2 domain to bind to the wild type GluN2A

C-terminus (KKMPSIESDV) and wild type GluN2B C-terminus (EKLSSIESDV)

yielded a K_D (dissociation constant) value of 0.31μM and 0.25mM, respectively (Fig

2E,F), whereas no binding was observed between the mutant GluN2A C terminus

(KKMPSIEADK) (Fig 2G) and the FRMPD2 PDZ2. These results supported the

direct interaction of C terminus of GluN2A to PDZ2 of FRMPD2.

To further explore potential sites for binding between GluN2A and FRMPD2,

we first obtained the crystal X-ray structure of the PDZ2 domain(Supplementary Fig

4A) of the mouse homologue of FRMPD2 (PDB ID: 5ZDS) (Supplementary Table 1).

The crystal structure was solved by PHENIX with the template PDZ2 domain

structure of tyrosine phosphatase PTP-BL (PDB ID: 1VJ6) (Supplementary Fig 4B)

with 63% identity(Adams et al, 2010; Gianni et al, 2006). In the template structure

(Supplementary Fig 4C), there was a peptide that is a C-terminal (HSGSYLVTSV) of

tumor suppressor protein APC, binding to β2 strand and α2 helix. The residues V(0)

and T(-2) of the peptide formed critical hydrophobic interactions with residues L25,

I27, L85, V29, and V65 of the template PDZ2 structure(Gianni et al, 2006). These

residues were highly conserved in the FRMPD2 PDZ2 crystal structure, as shown by

the alignment (Supplementary Fig 4D).

The results shown so far reveal that FRMPD2 affects NMDAR-mediated

excitatory postsynaptic currents by anchoring GluN2A subunits at postsynaptic

membranes. This activity is achieved by an interaction of the FRMPD2 PDZ2 domain

to the GluN2A C-terminus.

4. Effects of FRMPD2 on the formation of excitatory synapses in hippocampus

CA1 area.

Ferm domain-containing proteins are reported to recruit actin-regulatory

proteins(Yang et al, 2009), and the F-actin cytoskeleton is essential for neuronal

morphogenesis and synaptic connectivity(Bourne & Harris, 2008; Lei et al, 2001).

We further evaluated the effect of FRMPD2 on the synaptic ultrastructure of CA1

hippocampal neurons by electron micrograph (Fig 3A). First, we assessed the number

of excitatory (asymmetric) synapses, inhibitory (symmetric) synapses and presynaptic

vesicles(Weeks et al, 2007). There were no differences in the numbers of excitatory

synapses (Fig 3B), inhibitory synapses (Fig 3C) and presynaptic vesicles (Fig 3D)

after FRMPD2 knockdown or overexpression, consistent with our earlier

electrophysiology data, indicating that FRMPD2 does not affect the frequency of

mEPSC, mIPSCs or PPR values(Beique et al, 2003; Beique et al, 2006).

The length and thickness of the postsynaptic density (PSD) are primarily

correlated with the number of glutamate receptors anchoring at individual synapses

and this also closely related to the magnitude of mEPSCs (Sheng et al, 2007). Hence,

PSD (measured along the postsynaptic membrane) length and thickness were

calculated in mice after FRMPD2 knockdown or overexpression. The PSD was

shorter and thinner in the FRMPD2 knockdowns mice but longer and thicker in

FRMPD2 overexpression mice than in the corresponding controls (Fig 3E, F). These

results suggested that FRMPD2 is critically requirement for PSD size enlargement.

We next quantitatively assessed the contact length of synapses which is closely

related to spine head morphology in 2D electron micrographs. Contact length was

defined as the length of the postsynaptic membrane in register with the presynaptic

terminal was determined from the set of synapses used to measure PSD lengths

(Harris et al, 2012). The contact length was shorter in FRMPD2 knockdown mice but

longer in FRMPD2 overexpression mice than in the corresponding controls (Fig 3G).

These results indicate that FRMPD2 is probably associated with dendritic spine

morphology modifications.

Dendritic spines of neurons are the location of 90% excitatory synapses and

contain the postsynaptic machinery, including glutamate receptors, scaffold proteins

and actin cytoskeleton(Bourne & Harris, 2008). Hence, we tested that whether

FRMPD2 regulates the spines density and morphogenesis. Golgi staining was

performed to determine the density and morphology of both apical and basal

dendritic spines in hippocampal CA1 pyramidal neurons obtained from FRMPD2

overexpression or knockdown mice (Fig 3H). Spines are notably classified as "thin",

"stubby", "mushroom" and " filopodia" based on various morphological

characteristics (as illustrated in Fig 3I). Mature spines ("mushroom") have an

enlarged head and a larger, more complex PSD that contains a higher density of

glutamate receptors, resulting in a higher sensitivity to glutamate than in immature

spines ("thin", "stubby" and " filopodia")(Bourne & Harris, 2008). We measured the

numbers of mature and immature spines. There was an obvious reduction in mature

spines numbers and an increase in immature spines numbers in CA1 hippocampal

neurons in FRMPD2 knockdown mice. Conversely, the density of mature spines in

mice overexpressing FRMPD2 was higher than in the negative controls, while the

number of immature spines was reduced (Fig 3J). These data supported the view that

FRMPD2 plays a role in spines maturation and density modulations.

5. Binding of the FRMPD2 Ferm domain to F-actin.

Ferm domain proteins have been reported to interact with F-actin that is closely

associated with the NMDAR trafficking and dendritic spines morphogenesis(Bourne

& Harris, 2008; Lei et al, 2001; Yang et al, 2009). Then, we tried to find whether the

Ferm domain of FRMPD2 binds to F-actin. We transfected Flag-FRMPD2 constructs

with or without a Ferm domain into HEK293T (Fig 4A) and SYHY-5Y (Fig 4B) cells,

and then stained the cells for F-actin using rhodamine phalloidin. The full length of

FRMPD2 constructs with a Ferm domain (WT) colocalized with F-actin. Deleting the

FRMPD2 Ferm domain (ΔFerm) eliminated its colocalization with actin, but the

Ferm domain (Ferm) alone was required for actin colocalization. A quantification

analysis showed that F-actin colocalization was significantly lower in the FRMPD2

ΔFerm-transfected cells than in those transfected with either FRMPD2 WT or Ferm

(Fig 4C). Finally, we used an F-actin co-sedimentation assay, and the results showed

that the Ferm domain of FRMPD2 binds directly to F-actin (Fig 4D). These results

suggested that the FRMPD2-mediated morphological changes in dendritic spines

probably depend on an interaction between the FRMPD2 Ferm domain and F-actin.

6. Modulation of FRMPD2 in epilepsy susceptibility.

The NMDAR subunit GluN2A has been powerfully implicated in the

development of epilepsy, a disease that manifests as strong recurrent excitatory

connections in the hippocampus(Kwan et al, 2011; Lemke et al, 2013). Because

FRMPD2 was proved to mediate GluN2A-containing excitatory synaptic

transmission in the hippocampus, we further explored whether FRMPD2 is involved

in the regulation of epilepsy susceptibility.

Chronic pentylenetetrazol (PTZ) and kainic acid (KA)-induced epilepsy models

were used. FRMPD2 protein levels were significantly higher in epileptic patients (Fig

5A, B) as well as in both the hippocampus and neocortex of epileptic mice than in the

corresponding controls (Fig 5C-F). To further explore whether FRMPD2 modulates

epilepsy susceptibility, behavioral and local field potential (LFP) studies were

performed in two chronic epileptic animal models. Behavioral observation performed

in PTZ induced models showed that daily seizure scores were lower in FRMPD2

knockdown mice than in scramble controls (Fig 5G). More specifically, the FRMPD2

knockdown mice exhibited a lower total number of seizures and longer induced

latency than were observed in the scramble controls (Fig 5H, I). In a behavioral

analysis of KA-induced mice, the frequency of spontaneous recurrent seizures (SRSs)

was lower and SRS latency was longer in FRMPD2 knockdown mice than in

scramble controls(Fig 5J, K). Seizure-like events (SLEs) were recorded on EEGs

performed in KA-induced mice (Fig 5O, P), while the FRMPD2 knockdown mice

presented less numerous SLEs and had a shorter duration of SLE and a longer SLE

interval time than were observed in the scramble controls(Fig 5L-N).

These behavioral results suggested that FRMPD2 modulates susceptibility to

epilepsy by changing the duration, frequency and number of seizures.

Discussion

In our report, we identified a novel scaffold protein FRMPD2, and explored its physiological contribution to the modulation of excitatory synaptic electrophysiological function as well as excitatory synaptic formation (Fig 6). Scaffold proteins traffic and target of glutamate receptors(Elias & Nicoll, 2007; Lau et al, 2007) or sodium channels(Hruska-Hageman et al, 2002; Schultz et al, 1998) to the plasma membrane, and are involved in neuronal excitatory. Scaffolding proteins are also well-characterized for the regulation of dendritic spine morphology through coordinating cytoskeleton dynamic(Durand et al, 2012; Lee et al, 2016). Mature dendritic spines with numerous postsynaptic glutamate receptors are tightly correlated with strength of excitatory synapses(Chen et al, 2011). In general, there is little scaffolding proteins not only directly bind to receptors to govern excitatory synaptic function, but also directly bind to F-actin to control excitatory synaptic formation. For example, Shank proteins can bind to several actin-regulatory proteins, but they indirectly associate with type I metabotropic glutamate receptors via Homer protein(Durand et al, 2012). However, FRMPD2 interdependently enhance neuronal excitatory by the two isolate pathways.

While NMDAR-mediated and AMPAR-mediated transmission are both affected by PDZ proteins, but the FRMPD2 was not essential for AMPA-component excitatory transmission in adult hippocampus. It is seems that the binding of glutamate receptors by PDZ proteins appears to change with development in a glutamate receptor-specific manner(Beique et al, 2006; Elias et al, 2008). Knockout

of PSD-95 reduces AMPAR synaptic transmission with little effects on NMDARs in

mature neurons(Beique et al, 2006). Interesting, mice deficient MAGUKs family

members PSD-93, PSD-95 and SAP-102 had modestly reduced NMDAR EPSCs at

CA1 neurons(Elias et al, 2006), suggesting the potential existence of other PDZ

proteins that act independently of the MAGUKs family that regulate the targeting of

PDZ-dependent synaptic NMDARs(Lau & Zukin, 2007). FRMPD2 is supposed as

the potential PDZ protein to be capable of replenishing NMDAR-mediated excitatory

synaptic transmission, providing a more comprehensive understanding of the roles of

PDZ proteins in NMDAR targeting and function.

Our results showed that synaptic FRMPD2 mediated neurotransmission by

trafficking GluN2A subunits on synapses in adult hippocampus. NMDA receptors are

composed of obligatory GluN1 subunit co-assembled with GluN2A or GluN2B

subunit (GluN1/GluN2A or GluN1/GluN2B). The identity of the GluN2A and/or

GluN2B subunits determines the biophysical and pharmacological properties of

NMDARs (Bard et al, 2010; Sultan et al, 2015). Our results of Co-IP, GST-pull-down

and SPR assays identified the second PDZ domain of FRMPD2 as a novel binding

partner of GluN2A *in vivo* and *in vitro*. Although the FRMPD2 PDZ2 did not directly

bind to GluN1, the obligatory GluN1 subunit of GluN1/2A complex was, expectedly,

detectable by Co-IP of FRMPD2. Meanwhile, the GluN1 protein of synaptosome was

changed after FRMPD2 knockdown or overexpression, which was accompanied by

the alteration of GluN2A subunits. It is interesting that the FRMPD2 did not pull

down the GluN2B subunit *in vivo* or *in vitr*o. It might be because the binding of

PDZ2 to GluN2B is flexible and instant. In addition, only four amino acids were

different between peptides sequences of C-termini of GluN2A and GluN2B in SPR

experiments, the binding capacity of PDZ2 to GluN2A subunit was 1000 times higher

than that to GluN2B subunit. It is probably that the GluN2B-containing NMDARs are

located at extrasynaptic sites and involved in neurotoxicity in adult hippocampal

neurons, GluN2A-containing NMDARs mainly resided at synaptic sites and involved

in neurotransmission (Parsons et al, 2014). The low binding capacity of PDZ2 to

GluN2B may partially explain that the FRMPD2 did not pull down the GluN2B

subunit *in vivo* or *in vitr*o. Further investigation will be needed to explore that the

binding of PDZ2 of FRMPD2 to GluN2A was stronger than that to GluN2B.

FRMPD2 was involved in excitatory synapses formation possibly in a Ferm

domain-dependent manner. Ferm domain-containing proteins could function as a

bridge between cytoskeletal structures and specific locations on the plasma

membrane. The GST-Ferm domain of FRMPD2 was shown to bind to

phosphatidylinositol PtdIns(3,4)P2, a key cofactor in signaling to the actin

cytoskeleton(Stenzel et al, 2009). In our study, a direct interaction between the Ferm

domain of FRMPD2 and F-actin was firstly reported. These data are consistent with

previous reports showing that F-actin cosediments with other Ferm

domain-containing proteins, including merlin and myo1(Gotesman et al, 2010; Sun et

al, 2002). Ferm domain of FRMPD2 is capable of interacting with F-actin to link

F-actin to the synaptic plasma membrane, especially the PSD, a growing ends of

F-actin resulting in changes in the morphology of spines(Rao et al, 2000). Moreover,

the F-actin cytoskeleton plays a pivotal role in NMDAR trafficking(Lei et al, 2001),

and dramatic increases in the proportion of NMDAR, especially GluN1/GluN2A

subunits trafficking at the postsynaptic membrane can also efficiently facilitate

dendritic spines maturation, which is tightly correlated with the strength of excitatory

synapses in the hippocampus(Chen et al, 2011; Meyer et al, 2014).

In addition, although Ferm-containing proteins have been reported to bind to the

C-terminal tail of metabotropic glutamate receptor 5 (mGluR5) by the Ferm domain

(Hu et al, 2012b), in our study, the results of Co-IP and GST-pull-down assays did not

reveal any interaction between the FRMPD2 Ferm domain and mGluR1 or mGluR5

(data not shown). The mechanisms underlying these interactions remain unclear.

Our study indicated that FRMPD2 increases susceptibility to epilepsy, probably

by increasing the NMDARs, as well as excitatory synapses formation, which

contribute to excitatory neurotransmission integratedly. Multiple characteristics

associated with excitatory neurotransmission, from receptor trafficking to dendritic

spine morphology, contribute to the etiology underlying epileptogenesis (Noebels,

2015). The NMDAR subunit GluN2A is powerfully implicated in the development of

epilepsy, and mutations in the gene encoding GluN2A subunit cause idiopathic

focal epilepsy (Endele et al, 2010; Lemke et al, 2013). Selectively

blocking GluN2A-containing NMDARs impaired epileptogenesis in pilocarpine

induced epileptic rat models (Chen et al, 2007), suggesting that the GluN2A subunit

is a therapeutic target for epilepsy treatment. In addition, epileptic mice showed an

increase in mature dendritic spines of excitatory neurons (Oberheim et al, 2008), and

disrupting of F-actin polymerization in dendritic spines induced epileptic seizures and

a long-term increase in neuronal excitability (Sierra-Paredes et al, 2006).

In summary, the results presented here suggest a critical role of novel scaffold

protein FRMPD2 in regulating NMDA receptors anchoring and dendritic spine

maturation, both of which is critical for regulation of glutamatergic synapses

transmission. The dysregulation of this scaffold protein contributed to the impaired

excitatory conductance which has been associated with neurological disease epilepsy.

Methods and materials

DNA constructs

To make the Flag-tagged FRMPD2 expression constructs, the following human FRMPD2 sequences (uniprot: Q68DX3-1) were cloned into pcDNA3.1-3xFlag (Sigma): Flag-FRMPD2 (amino acids1-1309), Flag-ΔFerm (full length with a deletion at amino acids342-642), and Flag-Ferm (amino acids 342-642).

To construct the following GST fusion proteins, the PDZ or Ferm domain of mouse FRMPD2 (Uniprot: A0A140LI67) was subcloned into pGEX4T-1 with a GST tag: GST-PDZ1 (amino acids 786-872), GST-PDZ2 (amino acids 934-1027), GST-PDZ3 (amino acids 1071-1159) and GST-Ferm (amino acids 339-639).

The lentivirus vector (LV) constructs were manufactured by Genechem Co. Biotechnology (Shanghai City, China). The mouse FRMPD2 (NCBI Gene: 268729) short hairpin RNA (shRNA) knockdown construct targeted 5'-3' CTGAATCTTTCTCTTGCAT and was subcloned it into the LV GV248 (GeneChem, China), which contained green fluorescent protein (GFP). A scramble shRNA (5'-3' CTTTTCTAACGTGTCGACT) was used as a negative control for the FRMPD2 shRNA. The complete mouse cDNA sequence for FRMPD2 (XM_017316244) (FRMPD2 overexpression, FRMPD2 OE) and the GFP sequence were inserted into the LV GV358 (GeneChem, China).

Antibodies

The following antibodies were used: FRMPD2 (rabbit, Sigma;rabbit, Origene),

Flag(mouse, Sigma), GST(rabbit, Invitrogen), GFP (rabbit, Invitrogen),

GluN2A(rabbit, Proteintech), GluN2B(mouse, Proteintech; rabbit, Abcam),

GluN1(rabbit, Abcam), MAP2(guinea pig, SYSY), GFAP(mouse, Sigma),

PSD-95(guinea pig, SYSY; mouse, Millipore), VGLUT1(guinea pig, SYSY),

synaptophysin(rabbit, Abcam), syntaxin 1a(rabbit, Abcam).

Cell line cultures and transfection

HEK293T and SH-SY5Y cells were maintained in DMEM (Gibco)

supplemented with 10% fetal bovine serum (Gibco), 100 U/mL penicillin and 100

μg/mL streptomycin (Gibco) at 37°C in 5% CO_2. The cells were transfected with

Flag-FRMPD2, Flag-Ferm or Flag- ΔFerm using Lipofectamine 2000

(Invitrogen)(Klein et al, 2017). At 4 h after transfection, the cells were rinsed with

warm PBS and incubated in culture medium containing 10% fetal bovine serum.

Intrahippocampal injection of LV

Adult C57/BL6 male mice were obtained from the Experimental Animal

Center of Chongqing Medical University. Intrahippocampal injection of LV was

performed using a stereotaxic apparatus (Stoelting Co. Ltd Wood Dale, IL, USA).

Briefly, the mice were anesthetized with pentobarbital (80mg/kg, intraperitoneal) and

placed on stereotaxic apparatus. The reference points were bregma for the

anterior-posterior axis, the midline for the medial-lateral axis and the dura mater for

the dorsal-ventral axis with a tooth bar. LV particles were injected bilaterally into the

dorsal hippocampus through a glass pipette (0.2 μl/min) attached to a glass

microsyringe(Shangguan et al, 2017). The mice were allowed to recover for 2 weeks

after the vector injections.

Patch clamp recordings

Brain slices were prepared as previously reported(Chen et al, 2006; Kim et al,

2001). Briefly, C57/BL6 mice were anesthetized with pentobarbital. Brain slices (300

μm) were prepared with a Leica (Germany) VP1200S Vibratome and incubated in

artificial cerebral spinal fluid (ACSF) (119 mM NaCl, 26 mM $NaHCO_3$, 2.5 mM KCl,

1 mM $MgCl_2$, 1.25 mM NaH_2PO_4, 2 mM $CaCl_2$ and 25 mM glucose, pH 7.4, 310

mOsm) bubbled with 5% CO_2 and 95% O_2 for at least 1 h at room temperature before

recording. Whole-cell recording was performed as described previously(Liu et al,

2005). Glass microelectrodes (Sutter, USA) were shaped by a pipette puller (P-97,

Sutter, USA) to a final resistance of 3–5 MΩ when the pipette was filled with internal

solution. A multi-clamp 700B amplifier (Axon, USA) was used for the recordings.

Signals were sampled at 10 kHz and filtered at 2 kHz. A stable baseline was obtained

for at least 5 min prior to recording and data were discarded when the access

resistance (15–20 MΩ) was changed by 20% at the end of recording. Mini Analysis

6.0.1 (Synaptosoft) and Clamp Fit 10.3 software (Axon) were used to analyze the

recorded data.

For action potential (AP) recording, glass pipettes were filled with the following

internal solution: 17.5 mM KCl, 0.5 mM EGTA, 122.5 Mm K-gluconate, 10 mM

HEPES, and 4 mM ATP, pH adjusted to 7.2 with KOH [20]. The internal solution

used to record excitatory post-synaptic currents (EPSCs) contained: 17.5 mM CsCl,

10 mM HEPES, 4 mM ATP, 0.5 mM EGTA, 132.5 mM Cs-gluconate, and 5 mM

QX-314. The internal solution used to record inhibitory post-synaptic currents (IPSCs)

contained: 100 mM CsCl, 1 mM $MgCl_2$, 1 mM EGTA, 30 mM

N-methyl-D-glucamine, 10 mM HEPES, 5 mM MgATP 0.5 mM Na_2GTP and 12 mM

phosphocreatine.

Spontaneous activity was assessed by observing the neuron under current clamp

for 1 min. Tetrodotoxin (TTX, 1 μM), bicuculline (10 μM),

6,7-dinitroquinoxaline-2,3(1H,4H)-dione (DNQX, 20 μM) and D-serine (10 μM)

were added to Mg-free-ACSF to record NMDAR mEPSCs that held at −60 mV. TTX

(1 μM), bicuculline (10 μM) and amino-5-phosphonopentanoate (AP-V, 40 μM) were

added to ACSF to record AMPA mEPSCs. TTX(1 μM), DNQX(20 μM) and APV(40

μM) were added to ACSF to record mIPSCs that held at −70 mV (Shangguan et al,

2017).

Evoked EPSCs (eEPSCs) currents were generated with a 40μs pulse (0.1 Hz)

from a stimulated isolation unit controlled by an AMPI generator (Master-8, USA). A

bipolar stimulation electrode (FHC) was located ~100 μm rostral to the recording

electrode in the same layer(Chen et al, 2008; Tang et al, 2015). The eEPSCs at −70

mV holding potential in the present of PTX (100 μM) was considered as the

AMPAR-mediated current. Neurons were then held at +40 mV and the amplitude of

the eEPSCs at 50 ms post-stimulus was interpreted as the NMDAR-mediated

current(Etherton et al, 2009; Etherton et al, 2011). Input-output curve of NMDAR

eEPSCs were recorded in response to a series of stimulation intensities (pA).

Paired-pulse ratio (PPR) recordings were voltage-clamped at +40 mV and obtained

using a paired-pulse protocol of two stimuli at an inter-pulse interval of 50 ms. PPR

values were defined as the ratio of the second peak amplitude to the first peak

amplitude. The highly selective GluN2B subunit antagonist Ro25-6981 (Ro, 0.5μM)

was applied to assess GluN2A-sensitive eEPSCs, and the selective GluN2A

antagonist NVP-AAM077(NVP, 0.4μM) was added to record GluN2B component

eEPSCs(Bartlett et al, 2007).

Western blotting

Samples boiled with 4X sample buffer at 95°C for 5 min, and then separated on

8-10% SDS-PAGE gels and transferred onto polyvinylidene difluoride membranes

(Millipore). The target proteins were immunoblotted with primary antibodies

overnight at 4°C and then incubated with HRP-conjugated secondary antibodies. The

blots were imaged and quantified using a Fusion Imaging System. The quantitative

densitometric values of the proteins were normalized to that of GAPDH or beta-actin.

Immunofluorescence

For animal tissues, brain were post-fixed at 4°C for 24 h. Floating slices (50 mm

in thickness) were permeabilized with 0.5% Triton X-100 and blocked in goat serum.

The slices were then incubated with primary antibodies overnight at 4°C followed by

incubation with secondary fluorescent conjugated antibodies at room temperature for

2 h. Images were captured using laser-scanning confocal microscopy (Nikon A1+R

Microsystems) on an Olympus IX 70 inverted microscope (Olympus) equipped with

a Fluoview FVX confocal scanning head. Areas of overlap and fluorescence intensity

were analyzed using Image Pro Plus 6.0.

Sub-synaptic fractionation

Sub-synaptic fractionation was performed as previously described(Phillips et al,

2001). To obtain synaptosomes, adult mouse hippocampus were homogenized in

solution buffer (0.32 M sucrose, 0.1 mM $CaCl_2$, 1 mM $MgCl_2$, 0.1 mM PMSF) and

brought to a final sucrose concentration of 1.25 M. Protease inhibitors were used in

all purification steps. The homogenate was overlaid with 1.0 M sucrose and

centrifuged (100,000 g, 3 h, 4°C). Synaptosomes were collected at the 1 M/1.25 M

interface.

For sub-synaptic fractionation, synaptosomes were sequentially extracted with

1% Triton X-100 first at pH 6.0 and then at pH 8.0. Synaptosomes (described above)

were diluted 1:10 with 0.1 mM $CaCl_2$ to a final concentration of 20 mM Tris, pH 6.0

and 1% Triton X-100. After the solution was incubated for 30 min and then

centrifuged 40,000 g for 30 min at 4°C. The supernatant (presynaptic fraction) were

precipitated with 10 volumes of acetone at −20°C and recovered by centrifugation

(20,000 g; 30 min, 4°C). The pellet containing insoluble postsynaptic density (PSD)

was resuspended in 10 volumes solution buffer (1% Triton X-100, 20 μM Tris, pH

8.0, 4°C), incubated for 30 min, and centrifuged (40,000 g, 30 min, 4°C).

GST-pull-down and co-immunoprecipitation assays.

The methods used to produce the GST fusion proteins in bacteria have been

previously reported (Anzai et al, 2002; Anzai et al, 2004). GST-pull-down assays

were carried out using a GST Protein Interaction Pull-Down Kit (Pierce) according to

the manufacturer's instructions. Briefly, the bacterial lysate containing the fusion

proteins was sonicated and centrifuged, and the supernatant was used for protein

binding assays. In vitro translation was facilitated using a plasmid carrying

full-length GluN2A, GluN2B or GluN1 (Addgene) that was transfected into

HEK293T cells. The translated total proteins obtained from cells and the GST fusion

proteins were incubated together with GST glutathione agarose resin. The protein

complexes were then eluted by glutathione and analyzed by Western blotting using

GST, GFP, GluN2A, GluN2B or GluN1 antibodies.

For co-immunoprecipitation assays, the extracts contained in the crude

synaptosomal fraction obtained from adult mouse hippocampal tissues were prepared

with anti-GluN2A, anti-GluN2B, anti-GluN1 or anti-FRMPD2 antibodies or IgG

(rabbit, Abcam) (Control) at 4°C for 12 h before they were incubated with protein

A/G agarose beads (Beyotime). The immunoprecipitated mixture was immunoblotted

with GluN2A, GluN2B, GluN1 or FRMPD2 antibodies.

Golgi staining

Golgi staining was performed using a Hito Golgi-Cox OptimStain Kit according

to the manufacturer's protocol (Hitobiotec Inc.). Spines were counted on secondary

and tertiary branches of apical and basal dendrites in the CA1 hippocampal. 5

neurons were chosen from 3 mice per group, and 5 counting segments (10 μm in

length) of either secondary or tertiary dendrites were randomly selected from each

neuron for statistical comparison. Dendrite length was measured using Image pro

plus 6.0.

Actin cosedimentation assays.

F-actin binding assays were performed as described by the manufacturer

(Cytoskeleton). Briefly, α-actinin and purified GST-Ferm (amino acids 339-639) of

mouse FRMPD2 were incubated with freshly polymerized actin (F-actin) for 30 min

at room temperature, and then subjected to ultracentrifugation (150,000 x g) for 1.5 h

to pellet the proteins that had bound to F-actin. The pellet and supernatant were

analyzed by Coomassie blue staining.

Surface Plasmon Resonance (SPR) assays.

SPR biosensor analysis was used to verify binding between peptides and the

PDZ2 domain of FRMPD2 protein using a Biacore 8K instrument (GE Healthcare),

as previously described(Zhang et al, 2014a). Proteins were immobilized on S CM5

chips using amine-coupling chemistry. Binding experiments were performed in

running buffer (10 mM PBS, pH 7, 0.05% P20) at 25°C with a flow rate of 30 μL/min.

Gradient concentrations of PDZ2 proteins were diluted to 50 ug/ml in sodium

acetate(10 mM, pH 5.5) and injected into the channel for 120 s, followed by

disassociation for 120 s. RU values were collected at 10 Hz and all the experimental

data were globally analyzed in a steady-state model within Biacore 8K Evaluation

software.

Transmission electron microscopy

Tissues were prepared and sectioned for transmission electron microcscopy as

previously described(Hu et al, 2012a). Briefly, mice were transcardially perfused with

2.5% glutaraldehyde and 4% paraformaldehyde. Brain tissue (1 mm^3 size) was cut

from the hippocampus CA1 area, fixed in 2.5% glutaraldehyde for 2 h, post-fixed in

1% osmium tetroxide for 2 h and then dehydrated through ethanol gradients. After the

tissues were embedded in Epon 618 epoxy resin, the samples were cut into ultrathin

sections by an ultramicrotome (LKB-V, LKB Produkter AB, Sweden), and

double-stained by uranyl acetate/lead citrate. Ultra-structures were observed by

transmission electron microscope (JEM-1400PLUS, JEOL, Japan). In the present

study, excitatory synapses were identified as those with round vesicles and

asymmetric pre- and postsynaptic dense material, while inhibitory synapses were

identified as those with flattened vesicles and symmetric dense material(Larramendi

et al, 1967; Weeks et al, 2007). Sampling fields were randomly selected for imaging

in each section, and measured using the software Image Pro Plus 6.0.

Human samples

Temporal lobe tissues were randomly obtained from twenty post-surgery patients treated for TLE for whom tissues were available in our brain tissue bank (Shangguan et al, 2017; Xu et al, 2017; Zhang et al, 2014b). A diagnosis of TLE was determined based on the 2001 International Classification of Epileptic Seizures by the International League Against Epilepsy (ILAE)(Engel, 2001). The temporal cortical tissues used as control samples were obtained from twelve randomized patients who underwent surgery for trauma-induced intracranial hypertension. The control patients had no history of epilepsy, no exposure to antiepileptic drugs, and no history of neurological or psychiatric disorders. A neuropathological examination confirmed that the temporal neocortex specimens obtained from the control patients were normal. There was no significant difference in age or gender between the TLE and control patients (Supplementary Table 2).

Animal behaviors in epilepsy

Two weeks after the intrahippocampal injection of LV, the mice were kindled by pentylenetetrazol (PTZ) or kainic acid (KA) to measure epilepsy susceptibility. This PTZ induced model was performed according to the methods described in a previously study(Gelfuso et al, 2013). C57/BL6 mice were administered an intraperitoneal injection of PTZ (35 mg/kg) every other day and then observed for 60 min after each injection. Behavioral seizure scores were assessed based on the Racine standard scale, as follows: Grade 0 indicated arrest, wet dog shaking, and normal

behavior; Grade 1 indicated facial twitches (nose, lips, and eyes); Grade 2 indicated

chewing and head nodding; Grade 3 indicated forelimb clonus; Grade 4 indicated

rearing and falling on forelimbs; and Grade 5 indicated imbalance and falling on the

side or back(Racine, 1972). Only fully induced seizures (Grades 4 and 5) were

evaluated to determine the number of seizures and the latency period. The controls

was not performed at least 3 consecutive seizures with a score of 4 or 5.

Mice were injected with KA as previously described (Bitsika et al, 2016). Briefly,

two weeks after the intrahippocampal injection of LV, the mice were anesthetized and

slowly injected with KA (1 nmol in 50 nl) into dorsal hippocampus via a guide

cannula under stereotaxic guidance. At one hour after the onset of KA-induced status

epilepticus(SE), diazepam (10 mg/kg, i.p.) was administered to suppress SE. Then,

the animals were video recorded for twenty-four hours to monitor spontaneous

recurrent seizures (SRSs) for 1 month. Seizure scores were evaluated on the Racine

scale (see above), and only seizures with a score of 4 or 5 were recorded to determine

the number of SRSs and the latency period that between the SE termination and the

first SRS. Multichannel electroencephalogram (EEG) recording was performed to

quantify the EEG seizures using the OmniPlex® D Neuronal Data Acquisition

System (Plexon, Dallas, TX) as previously described(Shangguan et al, 2017; Xu et al,

2017). The signals were preamplified (1000×), filtered (0.1-1000 Hz), and digitized at

4 kHz. EEG seizures like events were characterized by high frequencies (>5 Hz),

high amplitudes (>2 times the baseline), and durations of >5 s (Jimenez-Mateos et al,

2012).

Study approval

All protocols of animal studies were approved by the Commission of Chongqing Medical University for the ethics of experiments on animals and conducted in accordance with international standards. All procedures of human samples collections were formally consented to by the patients or their lineal relatives and approved by the Ethics Committee of Chongqing Medical University according to the tenets of the Declaration of Helsinki.

Statistical analysis

All averaged data are presented as the means ± SD, and all graphs were prepared using GraphPad Prism 4 software (La Jolla, CA). For independent-samples, Student's t-test was used. Repeated measures ANOVA followed by post hoc t-tests was used to measure differences between two groups at multiple time points. The $\chi 2$ test was used to compare gender differences between the patients in the TLE and control groups. $p < 0.05$ and 0.01 were considered to indicate statistical significance and are indicated as $*p < 0.05$ or $**p < 0.01$, respectively. All tests were two-sided. n indicates the number of cells/slices or independent experiments and was used to calculate the degree of freedom. All samples included in each experiment were analyzed in triplicate.

Acknowledgements

The authors are sincerely grateful for the support of Beijing Tiantan Hospital, Beijing Xuanwu Hospital, and the Daping Hospital of the Third Military Medical University for supplying the brain tissues used in this study. In addition, we thank all of the patients and their families for their participation in this study. This work was supported by grants from the National Natural Science Foundation of China (No. 81271445, 81471319, 81301109, and 81301110) and the National Clinical Key Specialty Construction Foundation of China.

Author contributions

648 X. L. and XF. W. conceived and designed the experiments. HQ. Z., Y.L.,

649 RJ.Z., MQ.H. and YL. M. performed the experiments. X. L. collected and analyzed

650 the data. X. L. wrote the manuscript. All authors contributed to preparation of the

651 paper and approved the final contributions.

652 The authors have declared that no conflict of interest exists.

References

Adams PD, Afonine PV, Bunkoczi G, Chen VB, Davis IW, Echols N, Headd JJ, Hung LW, Kapral GJ, Grosse-Kunstleve RW, McCoy AJ, Moriarty NW, Oeffner R, Read RJ, Richardson DC, Richardson JS, Terwilliger TC, Zwart PH (2010) PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta crystallographica Section D, Biological crystallography* **66:** 213-221

Anzai N, Deval E, Schaefer L, Friend V, Lazdunski M, Lingueglia E (2002) The multivalent PDZ domain-containing protein CIPP is a partner of acid-sensing ion channel 3 in sensory neurons. *The Journal of biological chemistry* **277:** 16655-16661

Anzai N, Miyazaki H, Noshiro R, Khamdang S, Chairoungdua A, Shin HJ, Enomoto A, Sakamoto S, Hirata T, Tomita K, Kanai Y, Endou H (2004) The multivalent PDZ domain-containing protein PDZK1 regulates transport activity of renal urate-anion exchanger URAT1 via its C terminus. *The Journal of biological chemistry* **279:** 45942-45950

Bard L, Sainlos M, Bouchet D, Cousins S, Mikasova L, Breillat C, Stephenson FA, Imperiali B, Choquet D, Groc L (2010) Dynamic and specific interaction between synaptic NR2-NMDA receptor and PDZ proteins. *Proceedings of the National Academy of Sciences of the United States of America* **107:** 19561-19566

Bartlett TE, Bannister NJ, Collett VJ, Dargan SL, Massey PV, Bortolotto ZA, Fitzjohn SM, Bashir ZI, Collingridge GL, Lodge D (2007) Differential roles of NR2A and NR2B-containing NMDA receptors in LTP and LTD in the CA1 region of two-week old rat hippocampus. *Neuropharmacology* **52:** 60-70

Beique JC, Andrade R (2003) PSD-95 regulates synaptic transmission and plasticity in rat cerebral cortex. *The Journal of physiology* **546:** 859-867

Beique JC, Lin DT, Kang MG, Aizawa H, Takamiya K, Huganir RL (2006) Synapse-specific regulation of AMPA receptor function by PSD-95. *Proceedings of the National Academy of Sciences of the United States of America* **103:** 19535-19540

Bitsika V, Duveau V, Simon-Areces J, Mullen W, Roucard C, Makridakis M, Mermelekas G, Savvopoulos P, Depaulis A, Vlahou A (2016) High-Throughput LC-MS/MS Proteomic Analysis of a Mouse Model of Mesiotemporal Lobe Epilepsy Predicts Microglial Activation Underlying Disease Development. *Journal of proteome research* **15:** 1546-1562

Bourne JN, Harris KM (2008) Balancing structure and function at hippocampal dendritic spines. *Annual review of neuroscience* **31:** 47-67

Bredt DS, Nicoll RA (2003) AMPA receptor trafficking at excitatory synapses. *Neuron* **40:** 361-379

Chen BS, Thomas EV, Sanz-Clemente A, Roche KW (2011) NMDA receptor-dependent regulation of dendritic spine morphology by SAP102 splice variants. *The Journal of neuroscience : the official*

journal of the Society for Neuroscience **31:** 89-96

Chen G, Chen P, Tan H, Ma D, Dou F, Feng J, Yan Z (2008) Regulation of the NMDA receptor-mediated
synaptic response by acetylcholinesterase inhibitors and its impairment in an animal model of
Alzheimer's disease. *Neurobiology of aging* **29:** 1795-1804

Chen G, Kittler JT, Moss SJ, Yan Z (2006) Dopamine D3 receptors regulate GABAA receptor function
through a phospho-dependent endocytosis mechanism in nucleus accumbens. *The Journal of*
neuroscience : the official journal of the Society for Neuroscience **26:** 2513-2521

Chen Q, He S, Hu XL, Yu J, Zhou Y, Zheng J, Zhang S, Zhang C, Duan WH, Xiong ZQ (2007) Differential
roles of NR2A- and NR2B-containing NMDA receptors in activity-dependent brain-derived
neurotrophic factor gene regulation and limbic epileptogenesis. *The Journal of neuroscience : the*
official journal of the Society for Neuroscience **27:** 542-552

Durand CM, Perroy J, Loll F, Perrais D, Fagni L, Bourgeron T, Montcouquiol M, Sans N (2012) SHANK3
mutations identified in autism lead to modification of dendritic spine morphology via an
actin-dependent mechanism. *Molecular psychiatry* **17:** 71-84

Elias GM, Elias LA, Apostolides PF, Kriegstein AR, Nicoll RA (2008) Differential trafficking of AMPA and
NMDA receptors by SAP102 and PSD-95 underlies synapse development. *Proceedings of the National*
Academy of Sciences of the United States of America **105:** 20953-20958

Elias GM, Funke L, Stein V, Grant SG, Bredt DS, Nicoll RA (2006) Synapse-specific and developmentally
regulated targeting of AMPA receptors by a family of MAGUK scaffolding proteins. *Neuron* **52:**
307-320

Elias GM, Nicoll RA (2007) Synaptic trafficking of glutamate receptors by MAGUK scaffolding proteins.
Trends in cell biology **17:** 343-352

Endele S, Rosenberger G, Geider K, Popp B, Tamer C, Stefanova I, Milh M, Kortum F, Fritsch A, Pientka
FK, Hellenbroich Y, Kalscheuer VM, Kohlhase J, Moog U, Rappold G, Rauch A, Ropers HH, von Spiczak S,
Tonnies H, Villeneuve N et al (2010) Mutations in GRIN2A and GRIN2B encoding regulatory subunits
of NMDA receptors cause variable neurodevelopmental phenotypes. *Nature genetics* **42:** 1021-1026

Engel J, Jr. (2001) Intractable epilepsy: definition and neurobiology. *Epilepsia* **42 Suppl 6:** 3

Etherton MR, Blaiss CA, Powell CM, Sudhof TC (2009) Mouse neurexin-1alpha deletion causes
correlated electrophysiological and behavioral changes consistent with cognitive impairments.
Proceedings of the National Academy of Sciences of the United States of America **106:** 17998-18003

Etherton MR, Tabuchi K, Sharma M, Ko J, Sudhof TC (2011) An autism-associated point mutation in
the neuroligin cytoplasmic tail selectively impairs AMPA receptor-mediated synaptic transmission in
hippocampus. *The EMBO journal* **30:** 2908-2919

Feng W, Zhang M (2009) Organization and dynamics of PDZ-domain-related supramodules in the
postsynaptic density. *Nature reviews Neuroscience* **10:** 87-99

Gelfuso EA, Liberato JL, Cunha AO, Mortari MR, Beleboni RO, Lopes NP, Dos Santos WF (2013)
Parawixin2, a novel non-selective GABA uptake inhibitor from Parawixia bistriata spider venom,
inhibits pentylenetetrazole-induced chemical kindling in rats. *Neuroscience letters* **543:** 12-16

Gianni S, Walma T, Arcovito A, Calosci N, Bellelli A, Engstrom A, Travaglini-Allocatelli C, Brunori M,
Jemth P, Vuister GW (2006) Demonstration of long-range interactions in a PDZ domain by NMR,
kinetics, and protein engineering. *Structure (London, England : 1993)* **14:** 1801-1809

Gotesman M, Hosein RE, Gavin RH (2010) A FERM domain in a class XIV myosin interacts with actin
and tubulin and localizes to the cytoskeleton, phagosomes, and nucleus in Tetrahymena thermophila.
Cytoskeleton (Hoboken, NJ) **67:** 90-101

Harris KM, Weinberg RJ (2012) Ultrastructure of synapses in the mammalian brain. *Cold Spring Harbor*
perspectives in biology **4**

Hruska-Hageman AM, Wemmie JA, Price MP, Welsh MJ (2002) Interaction of the synaptic protein
PICK1 (protein interacting with C kinase 1) with the non-voltage gated sodium channels BNC1 (brain
Na+ channel 1) and ASIC (acid-sensing ion channel). *The Biochemical journal* **361:** 443-450

Hu B, Chen H, Yang L, Tao ZF, Yan J, Zhang YH, Zhu ZR, Sun WZ, Huang W, Huang WQ, Sui JF (2012a)
Changes of synaptic ultrastructure in the guinea pig interpositus nuclei associate with response
magnitude and timing after trace eyeblink conditioning. *Behavioural brain research* **226:** 529-537

Hu JH, Yang L, Kammermeier PJ, Moore CG, Brakeman PR, Tu J, Yu S, Petralia RS, Li Z, Zhang PW, Park
JM, Dong X, Xiao B, Worley PF (2012b) Preso1 dynamically regulates group I metabotropic glutamate
receptors. *Nature neuroscience* **15:** 836-844

Jimenez-Mateos EM, Engel T, Merino-Serrais P, McKiernan RC, Tanaka K, Mouri G, Sano T, O'Tuathaigh
C, Waddington JL, Prenter S, Delanty N, Farrell MA, O'Brien DF, Conroy RM, Stallings RL, DeFelipe J,
Henshall DC (2012) Silencing microRNA-134 produces neuroprotective and prolonged
seizure-suppressive effects. *Nature medicine* **18:** 1087-1094

Kim CH, Chung HJ, Lee HK, Huganir RL (2001) Interaction of the AMPA receptor subunit GluR2/3 with
PDZ domains regulates hippocampal long-term depression. *Proceedings of the National Academy of*
Sciences of the United States of America **98:** 11725-11730

Kim E, Sheng M (2004) PDZ domain proteins of synapses. *Nature reviews Neuroscience* **5:** 771-781

Klein ZA, Takahashi H, Ma M, Stagi M, Zhou M, Lam TT, Strittmatter SM (2017) Loss of TMEM106B
Ameliorates Lysosomal and Frontotemporal Dementia-Related Phenotypes in Progranulin-Deficient

Mice. *Neuron* **95:** 281-296.e286

Kwan P, Schachter SC, Brodie MJ (2011) Drug-resistant epilepsy. *The New England journal of medicine*
365: 919-926

Larramendi LM, Fickenscher L, Lemkey-Johnston N (1967) Synaptic vesicles of inhibitory and
excitatory TERMINALS IN THE CEREBELLUM. *Science (New York, NY)* **156:** 967-969

Lau CG, Zukin RS (2007) NMDA receptor trafficking in synaptic plasticity and neuropsychiatric
disorders. *Nature reviews Neuroscience* **8:** 413-426

Lee SE, Kim Y, Han JK, Park H, Lee U, Na M, Jeong S, Chung C (2016) nArgBP2 regulates excitatory
synapse formation by controlling dendritic spine morphology.   **113:** 6749-6754

Lei S, Czerwinska E, Czerwinski W, Walsh MP, MacDonald JF (2001) Regulation of NMDA receptor
activity by F-actin and myosin light chain kinase. *The Journal of neuroscience : the official journal of*
the Society for Neuroscience **21:** 8464-8472

Lemke JR, Lal D, Reinthaler EM, Steiner I, Nothnagel M, Alber M, Geider K, Laube B, Schwake M,
Finsterwalder K, Franke A, Schilhabel M, Jahn JA, Muhle H, Boor R, Van Paesschen W, Caraballo R,
Fejerman N, Weckhuysen S, De Jonghe P et al (2013) Mutations in GRIN2A cause idiopathic focal
epilepsy with rolandic spikes. *Nature genetics* **45:** 1067-1072

Lipinski S, Grabe N, Jacobs G, Billmann-Born S, Till A, Hasler R, Aden K, Paulsen M, Arlt A, Kraemer L,
Hagemann N, Erdmann KS, Schreiber S, Rosenstiel P (2012) RNAi screening identifies mediators of
NOD2 signaling: implications for spatial specificity of MDP recognition. *Proceedings of the National*
Academy of Sciences of the United States of America **109:** 21426-21431

Liu QS, Pu L, Poo MM (2005) Repeated cocaine exposure in vivo facilitates LTP induction in midbrain
dopamine neurons. *Nature* **437:** 1027-1031

Meyer D, Bonhoeffer T, Scheuss V (2014) Balance and stability of synaptic structures during synaptic
plasticity. *Neuron* **82:** 430-443

Noebels J (2015) Pathway-driven discovery of epilepsy genes. *Nature neuroscience* **18:** 344-350

Oberheim NA, Tian GF, Han X, Peng W, Takano T, Ransom B, Nedergaard M (2008) Loss of astrocytic
domain organization in the epileptic brain. *The Journal of neuroscience : the official journal of the*
Society for Neuroscience **28:** 3264-3276

Parsons MP, Raymond LA (2014) Extrasynaptic NMDA receptor involvement in central nervous system
disorders. *Neuron* **82:** 279-293

Phillips GR, Huang JK, Wang Y, Tanaka H, Shapiro L, Zhang W, Shan WS, Arndt K, Frank M, Gordon RE,

Gawinowicz MA, Zhao Y, Colman DR (2001) The presynaptic particle web: ultrastructure, composition,
dissolution, and reconstitution. *Neuron* **32:** 63-77
830

831   Racine RJ (1972) Modification of seizure activity by electrical stimulation. II. Motor seizure.
832   *Electroencephalography and clinical neurophysiology* **32:** 281-294
833

834   Rao A, Craig AM (2000) Signaling between the actin cytoskeleton and the postsynaptic density of
835   dendritic spines. *Hippocampus* **10:** 527-541
836

837   Sala C, Piech V, Wilson NR, Passafaro M, Liu G, Sheng M (2001) Regulation of dendritic spine
838   morphology and synaptic function by Shank and Homer. *Neuron* **31:** 115-130
839

840   Schultz J, Hoffmuller U, Krause G, Ashurst J, Macias MJ, Schmieder P, Schneider-Mergener J, Oschkinat
841   H (1998) Specific interactions between the syntrophin PDZ domain and voltage-gated sodium
842   channels. *Nature structural biology* **5:** 19-24
843

844   Shangguan Y, Xu X, Ganbat B, Li Y, Wang W, Yang Y, Lu X, Du C, Tian X, Wang X (2017) CNTNAP4
845   Impacts Epilepsy Through GABAA Receptors Regulation: Evidence From Temporal Lobe Epilepsy
846   Patients and Mouse Models. *Cerebral cortex (New York, NY : 1991)***:** 1-14
847

848   Shen G, Van Sickle BJ, Tietz EI (2010) Calcium/calmodulin-dependent protein kinase II mediates
849   hippocampal glutamatergic plasticity during benzodiazepine withdrawal. *Neuropsychopharmacology :*
850   *official publication of the American College of Neuropsychopharmacology* **35:** 1897-1909
851

852   Sheng M, Hoogenraad CC (2007) The postsynaptic architecture of excitatory synapses: a more
853   quantitative view. *Annual review of biochemistry* **76:** 823-847
854

855   Sierra-Paredes G, Oreiro-Garcia T, Nunez-Rodriguez A, Vazquez-Lopez A, Sierra-Marcuno G (2006)
856   Seizures induced by in vivo latrunculin a and jasplakinolide microperfusion in the rat hippocampus.
857   *Journal of molecular neuroscience : MN* **28:** 151-160
858

859   Stenzel N, Fetzer CP, Heumann R, Erdmann KS (2009) PDZ-domain-directed basolateral targeting of
860   the peripheral membrane protein FRMPD2 in epithelial cells. *Journal of cell science* **122:** 3374-3384
861

862   Sultan S, Li L, Moss J, Petrelli F, Casse F, Gebara E, Lopatar J, Pfrieger FW, Bezzi P, Bischofberger J, Toni
863   N (2015) Synaptic Integration of Adult-Born Hippocampal Neurons Is Locally Controlled by Astrocytes.
864   *Neuron* **88:** 957-972
865

866   Sun CX, Robb VA, Gutmann DH (2002) Protein 4.1 tumor suppressors: getting a FERM grip on growth
867   regulation. *Journal of cell science* **115:** 3991-4000
868

Tang B, Luo D, Yang J, Xu XY, Zhu BL, Wang XF, Yan Z, Chen GJ (2015) Modulation of AMPA receptor
mediated current by nicotinic acetylcholine receptor in layer I neurons of rat prefrontal cortex.
Scientific reports **5:** 14099

Valtschanoff JG, Weinberg RJ (2001) Laminar organization of the NMDA receptor complex within the

postsynaptic density. *The Journal of neuroscience : the official journal of the Society for Neuroscience*

21: 1211-1217

Weeks AC, Connor S, Hinchcliff R, LeBoutillier JC, Thompson RF, Petit TL (2007) Eye-blink conditioning

is associated with changes in synaptic ultrastructure in the rabbit interpositus nuclei. *Learning &*

memory (Cold Spring Harbor, NY) **14:** 385-389

Xu X, Shangguan Y, Lu S, Wang W (2017) Tubulin beta-III modulates seizure activity in epilepsy.   **242:**

297-308

Yang Y, Baboolal TG, Siththanandan V, Chen M, Walker ML, Knight PJ, Peckham M, Sellers JR (2009) A

FERM domain autoregulates Drosophila myosin 7a activity. *Proceedings of the National Academy of*

Sciences of the United States of America **106:** 4189-4194

Zhang C, Wan X, Zheng X, Shao X, Liu Q, Zhang Q, Qian Y (2014a) Dual-functional nanoparticles

targeting amyloid plaques in the brains of Alzheimer's disease mice. *Biomaterials* **35:** 456-465

Zhang X, Chen G, Lu Y, Liu J, Fang M, Luo J, Cao Q, Wang X (2014b) Association of mitochondrial letm1

with epileptic seizures. *Cerebral cortex (New York, NY : 1991)* **24:** 2533-2540

Zucker RS, Regehr WG (2002) Short-term synaptic plasticity. *Annual review of physiology* **64:** 355-405

Fig 1. Effects of FRMPD2 on NMDA receptor-mediated excitatory synaptic transmission.

A, Quantitative analysis of amplitude and frequency of NMDAR mEPSCs in CA1 pyramidal cells of hippocampus after FRMPD2 knockdown or overexpression. Bar, 1 s, 20 pA. B, The input-output curves of NMDAR eEPSCs were recorded with gradual increasing intensities. Bar, 25 ms, 50 pA. C, Amplitude of NMDAR, AMPAR component eEPSCs and AMPA/NMDA ratio were compared. Bar, 25 ms, 25 pA. D, Quantitative analysis of the amplitude of GluN2A- and GluN2B-sensitive eEPSCs. Bar, 25 ms, 25 pA. E, Knocking down or overexpressing FRMPD2 did not alter the paired-pulse ratio (PPR) values. Bar, 25 ms, 50 pA. Data of were presented as the mean±SD or median with interquartile range. Data of A, C, D, E, were analyzed by Student's *t* test, data of B was analyzed by repeated measures ANOVA, n=5-7 neurons from 3 mice. **p<0.01. *p<0.05.

Fig 2. Effects of FRMPD2 anchoring GluN2A at synapses.

A, Quantitative analysis of GluN2A, GluN2B and GluN1 expressions of total hippocampus tissue and synaptic homogenates after FRMPD2 knockdown or overexpression. The data were shown as the mean±SD and analyzed by Student's t test, *$p < 0.05$. B, Immunofluorescence staining in the mouse hippocampus CA1 area showed that FRMPD2 co-localizes with GluN2A at the postsynaptic density, which was marked by PSD95. Bar, 40 μm. C, FRMPD2 and GluN2A, GluN2B and GluN1

were co-immunoprecipitated from mouse hippocampal tissues. D, GST pull-down

assay showing that the GST-PDZ2 fusion protein but not the PDZ1 or PDZ3 fusion

proteins pulled down endogenous GluN2A from mouse hippocampal tissues, and

exogenous full-length GFP-GluN2A from HEK293 cells but did not pull down the

deletion variant GFP-GluN2A from HEK293 cells transfected with GFP-GluN2A that

lacked ten amino acids at the C-terminus. E-G, The ability of the isolated FRMPD2

PDZ2 domain to bind to the GluN2A and GluN2B C-terminus was measured using

SPR. Peptides sequences of C-termini of wild type GluN2A (E), wild type GluN2B

(F) and mutant GluN2A (G) were used in this study. Sensorgrams were shown to

illustrate the binding of increasing concentrations of the PDZ2 domain to peptides

and steady-state equilibrium responses from the sensorgrams were shown plotted

against the injected PDZ2 concentrations. K_D: dissociation constant.

Fig 3. Effects of FRMPD2 on the formation of excitatory synapses in hippocampus CA1 area.

A, Electron micrograph of a spine synapse showing a typical continuous electron dense PSD and contact length. Scale bar, 400 nm; n=10-12 images from 3 mice. B-D, The number of excitatory synapses(B), inhibitory synapse(C) and vesicles per synapse(D) were compared after FRMPD2 knockdown or overexpression. E-F, Quantitative analysis and distribution of PSD length (E) and thickness (F) were

shown. G, Quantitative analysis and distribution of contacted length of the spine with

the presynaptic terminal were represented. H, Representative images of Golgi

staining to showed spine morphology in basal and apical dendrites of CA1

hippocampal neurons. Bar, 10 μm. I, Illustration of different spine morphologies:

mushroom, thin, stubby, and filopodia. J, Quantification analysis showed the densities

of total, mature and immature spines in both the apical and basal dendrites of CA1

hippocampal neurons in the FRMPD2 knockdown or overexpression mice. The data

were shown as the mean±SD or median with interquartile range; *p< 0.05, **p< 0.01.

Fig 4. Binding of the FRMPD2 Ferm domain with F-actin.

A, B, The Ferm domain of FRMPD2 was sufficient and required for F-actin colocalization. HEK293 (A) and SYHY-5Y (B) cells were transfected with Flag-tagged FRMPD2 (WT), a deletion variants (ΔFerm) or only the Ferm domain (Ferm) and then stained for Flag and F-actin (rhodamine-phalloidin). Scale bar, 50 μm. C, Quantification of colocalization in FRMPD2 variants between Flag and F-actin based on the results shown in A-B. The area of overlap between Flag and F-actin was compared with the total area expressing Flag to obtain the F-actin-positive Flag/total Flag ratio. Mean±SD (n=6 cells, respectively, **p< 0.01). D, Ferm of FRMPD2 co-precipitated with F-actin in a co-sedimentation assay. GST -Ferm or α-actinin (a positive control) were mixed with F-actin, co-precipitated by ultracentrifugation, and stained with Coomassie blue. P, pellet; S, supernatant.

Fig 5. Modulation of FRMPD2 in epilepsy susceptibility.

A, B, FRMPD2 expression of temporal neocortex of TLE patients(n=20) and control

patients(n=12) by immunoblot (A) and immunofluorescence staining(B). Bar, 40 μm.

C, D, FRMPD2 expression in the hippocampus(C, D) and neocortex(C) in the PTZ

induced mice. n=10-11 mice in each group(C). Bar, 40 μm. E, F, FRMPD2 expression

in the hippocampus(E, F) and neocortex(E) were compared between the KA-treated

mice and control mice without SRS. n=8-11 mice in each group. Bar, 40μm. G-I,

Quantitative analysis of daily seizure scores (G), total number of seizures with a

score of 4 or 5 (H) during and latency of the first seizure with a score of 4 or 5 (I) in

PTZ induced mice after FRMPD2 knockdown and overexpression. n=9-12 mice in

each group. J-K, Quantitative analysis of total number (J) and latency (K) of the first

SRSs with a score of 4 or 5 in KA induced mice. n=7-9 mice in each group. L-N,

Quantitative analysis of number (L), duration (M) and interval times (N) of seizure

like event (SLE) in KA induced mice. n=7-9 mice in each group. O, P, Representative

images of SLEs obtained in each group during a 30 min EEG recording. Bar, 100 s,

0.25 mV (O). EEG showed normal baseline, sharp waves in the interictal stage and an

SRS in the KA-induced epilepsy model. Bar, 2.5 s, 0.25 mV (P). OD, optical density.

Data were shown as the mean±SD or median with interquartile range; *p< 0.05, **p<

0.01.

Fig 6. Model demonstrating regulation of FRMPD2 at excitatory synaptic

transmission.

Supplementary Methold

Crystallization

GST-PDZ2 (amino acids 934-1027) was expressed in Escherichia coli,

BL21(DE3) cells. Cells pellet was sonicated and centrifuged at 5000 rpm. The

supernatant was subjected to GST column purification (10 mM Tris, 10 mM GSH,

pH 7.4). Pooled fractions were dialyzed with thrombin and passed through a Q-HP

column (20 mM Tris, 0-1 M NaCl, 5% glycerol, pH 8.0). Finally, protein fractions

were further purified by Superdex 75 columns (20 mM Tris, 5% glycerol, 150 mM

NaCl, 2 mM DTT pH 7.4) for crystallization.

Screen kits, including the JCSG-CORE-I-IV (Qiagen) and HR2-144 Index

(Hampton), were used to perfrom crystallization screen with the hanging drop

method at 18 ºC. The protein (15 mg/ml) was mixed with an equivalent volume (1 μl)

of screen buffer. The crystal was grown in 20% PEG3350 and 0.2 M sodium

thiocyanate. A single crystal was transferred to the crystallization buffer containing an

additional 20% glycerol and flash frozen with liquid nitrogen. The crystal diffracted

well at BL19U (Shanghai Synchrotron Radiation Facility, SSRF, China). In all,

images for 180° were collected. The data were processed by HKL3000

(Supplemental Table 1). The crystal structure was refined by PHENIX and Coot.

The structure was refined to 1.80 Å with R factor 19.76% and R free 24.76%.

1009

1010

1011 **Supplementary Figures**

1012

Supplementary Fig 1. Expression patterns of the FRMPD2 protein.

A, FRMPD2 protein is located in the mouse brain, eyes, testes and lungs. B, The

brain-specific expression of the FRMPD2 protein was revealed in adult mouse tissue

homogenates. C, The FRMPD2 protein has a widespread distribution pattern in adult

mouse brain slices. Bar, 500 μm. D, FRMPD2 was present in MAP2-positive

neuronal dendrites but not in GFAP-positive glial cells as determined by

immunofluorescence staining. Bar, 40 μm. E, FRMPD2 co-localized with the

postsynaptic marker PSD95 but not with the presynaptic marker VGLUT1 in the

mouse hippocampus and cortex. Bar, 40 μm. F, Immunoblot of the synapse fraction

(Syn) showing that FRMPD2 was preferentially distributed in the postsynaptic

fraction (Post) rather than in the presynaptic (Pre) or non-synaptic synaptosomal

proteins (NSSP) fraction. Anti-synapphysin, anti-syntaxin 1a and anti-PSD95 were

used as markers for ultrasynaptic fraction to verify that the purity of the preparation

was higher than 90%. The blots were representative of 4-6 analyses.

Supplementary Fig 2. Alteration of FRMPD2 protein levels in the hippocampus

by LV injection.

A, Lentiviral vector (LV)-encoding green fluorescent protein (GFP) was detected in

the mouse hippocampus following the injection of LV. Bar, 250 μm. B, Hippocampal

FRMPD2 protein expression levels were detected in FRMPD2 knockdown or

overexpression mice on the 2 and 6 weeks after LV injection. Data were shown as

mean±SD and analyzed by Student's t test, n=5 mice, *p< 0.05.

1036

1037

Supplementary Fig 3. Effects of FRMPD2 on electrophysiology of CA1 neurons in mouse hippocampal brain slices.

A, The frequency of spontaneous AP was compared after FRMPD2 knockdown or overexpression. Bar, 1 s, 25 mV. B, C, FRMPD2 had no significant effect on the amplitude or frequency of AMPAR mEPSCs (B) and mIPSCs (C). Bar, 1 s, 20 pA. Data were shown as mean±SD and analyzed by Student's *t* test, n=6-7 neurons of 3 mice, **p< 0.01.

1046

Supplementary Fig 4. Crystal structure of FRMPD2 PDZ2 domain.

A, The crystal structure of a compact globular fold in the ligand-free FRMPD2 PDZ2

domain (PDB ID: 5ZDS) was shown. B, The crystal structure of the FRMPD2 PDZ2

domain (green) was shown aligned to that of the PDZ2-peptide template structure

(cyan and magenta) (PDB ID: 1VJ6). C, A template of PDZ2-peptide (cyan and

magenta) interaction (PDB ID: 1VJ6). D, The binding pocket alignment between

FRMPD2 PDZ2 crystal structure (green) and PDZ2 template (cyan).

1054

1055

 Supplementary Table 1. Data collection and refinement statistics of PDZ2 of

 FRMPD2.

	PDZ2
Data collection	
Wavelength(Å)	0.9786
Resolution range(Å)	50-1.58 (1.64-1.58)[a]
Space group	P 2_1 2_1 2
Unit cell dimensions	
a, b, c(Å)	32.200, 48.179, 56.845
α, β, γ(°)	90, 90, 90
No. of reflections	80345
No. unique reflections	12699 (1248)
Multiplicity	6.3 (6.4)
Completeness (%)	99.7 (100)
Mean I/σ (I)	22.52(7.70)
R-meas	0.065(0.268)
Refinement	
Resolution range(Å)	50-1.8 (1.86-1.8)
No. of reflections	8610 (849)
No. of reflections used for R-free	860 (85)
R-work	0.1980 (0.2202)
R-free	0.2347 (0.2630)

No. of non-hydrogen atoms	810
Macromolecules	658
Protein residues	88
R.m.s.d.	
Bond lengths(Å)	0.005
Bond angles(°)	0.83
Ramachandranplot(%)	
Allowed regions	1.1
Outliers	0
Average B-factor	13.71
Macromolecules	11.25
Solvent	24.37

1058 [a]Statistics for the highest-resolution shell are shown in parentheses. No., number,

1059 r.m.s.d., root mean squared deviation.

1060

1061

 Supplementary Table 2. Clinical details of TLE patients and control patients

Cases	Sex	Age (years)	Duration (years)	Seizure type	Preoperative AEDs consumption	Resected tissue	Pathological diagnosis
E1	F	27	14	SGS	PHT, VAP, CZP	RTN	G
E2	F	13	4	SGS	CBZ, TPM, VPA	LTN	NL, ND, G
E3	F	15	3	CPS, SE	CBZ, TPM, LTG	RTN	G
E4	M	47	29	SGS	VPA, PHT, TPM, PB,LTG	RTN	G
E5	M	28	3	SGS	PB, CBZ, VPA	LTN	NL, ND, G
E6	F	38	12	SGS	CBZ, TPM, VPA	LTN	NL, ND, G
E7	F	29	27	SGS	CBZ, VPA, LTG	RTN	NL, G
E8	F	37	7	CPS, GTCS	CBZ, TPM, VPA	LTN	NL, G
E9	M	32	19	CPS	CBZ, TPM, PB,VPA	LTN	ND, G
E10	M	30	10	SGS	CBZ, VPA, PHT	LTN	NL, ND, G
E11	F	11	4	SGS	VPA, CBZ, TPM, LTG	RTN	G
E12	M	22	7	SGS, SPS	VPA, TPM, LTG	LTN	NL, G
E13	F	44	12	CPS	VPA, PHT, LTG	RTN	NL, G
E14	M	33	9	CPS	VPA, PHT, LTG	RTN	G
E15	M	17	3	SGS	PHT, PB, CBZ, LTG	LTN	NL, G
E16	F	14	5	SGS, CPS	VPA, CBZ, PHT, PB	RTN	NL, ND
E17	M	24	3	SGS, CPS	VPA, TPM, OXC	RTN	G

E18	F	33	10	SGS	CBZ, PHT	RTN	G
E19	M	29	14	CPS	VPA, TPM, CBZ, LTG,PHT	LTN	NL, ND, G
E20	M	24	11	CPS	VPA, PHT, CBZ	RTN	NL, G
C1	F	14	0	None	None	LTN	N
C2	F	25	0	None	None	RTN	N
C3	F	32	0	None	None	RTN	N
C4	M	28	0	None	None	RTN	N
C5	M	21	0	None	None	LTN	N
C6	M	54	0	None	None	LTN	N
C7	F	38	0	None	None	RTN	N
C8	M	50	0	None	None	RTN	N
C9	M	43	0	None	None	LTN	N
C10	F	33	0	None	None	LTN	N
C11	M	31	0	None	None	RTN	N
C12	F	15	0	None	None	LTN	N

AEDs, antiepileptic drugs; E, epilepsy; C, control; F, female; M, male; CPS, complex

partial seizure; SPS, simple partial seizure; SGS, secondarily generalized seizure;

GTCS, generalized tonic clonic seizure; SE, status epilepticus; CBZ, carbamazepine;

CZP, clonazepam; LTG, lamotrigine; OXC, oxcarbazepine; PB, phenobarbital; PHT,

phenytoin; TPM, topiramate; VPA, valproate; LTN, left temporal neocortex; RTN,

right temporal neocortex; NL, neuron loss; ND, neuron degeneration; G, gliosis; N,

normal.

1070
1071