Methodologic Issues in Doubly Labeled Water Measurements of Energy Expenditure

During Very Low-Carbohydrate Diets

Kevin D. Hall[1][†], Juen Guo[1], Kong Y. Chen[1], Rudolph L. Leibel[2], Marc L. Reitman[1], Michael

Rosenbaum[2], Steven R. Smith[4], Eric Ravussin[3]

[1]National Institute of Diabetes and Digestive and Kidney Diseases; [2]Columbia University;

[3]Pennington Biomedical Research Center; [4]The Translational Research Institute for Metabolism

and Diabetes

Running Title: Dietary Carbohydrate and Doubly Labeled Water

Names for PubMed indexing: Hall, Guo, Chen, Leibel, Reitman, Rosenbaum, Smith, Ravussin.

ClinicalTrials.gov Identifier: NCT01967563

Funding: Nutrition Sciences Initiative (NuSI). This work was also supported, in part, by the

Intramural Research Program of the National Institutes of Health, National Institute of Diabetes

and Digestive and Kidney Diseases (KDH, KYC, MLR), NIH UL1 TR00040 (Columbia CTSA,

MR and RL), and NORC Center Grant # P30DK072476 (ER).

†To whom correspondence should be addressed:

Kevin D. Hall, Ph.D.

National Institute of Diabetes & Digestive & Kidney Diseases,

12A South Drive, Room 4007

Bethesda, MD 20892

kevinh@niddk.nih.gov

Conflict of interest disclosure statement: None of the authors have conflicts of interest

WORD COUNT: 2994

REVISION DATE: October 23, 2018

Abstract

Background: Very low-carbohydrate diets have been reported to substantially increase human

energy expenditure as measured by doubly labeled water (DLW) but not by respiratory

chambers. Do the DLW data reflect true physiological differences that are undetected by

respiratory chambers? Alternatively, are the apparent DLW energy expenditure a consequence of

failure to fully account for respiratory quotient (RQ) differences between diets?

Objective: To examine energy expenditure differences between diets varying drastically in

carbohydrate and to quantitatively compare DLW data with respiratory chamber and body

composition measurements within an energy balance framework.

Design: DLW measurements were obtained during the final two weeks of month-long baseline

(BD; 50% carbohydrate, 35% fat, 15% protein) and isocaloric ketogenic diets (KD; 5%

carbohydrate, 80% fat, 15% protein) in 17 men with BMI 25-35 kg/m^2. Subjects resided 2d/week

in respiratory chambers to measure energy expenditure ($EE_{chamber}$). DLW expenditure was

calculated using chamber-determined respiratory quotients (RQ) either unadjusted (EE_{DLW}) or

adjusted ($EE_{DLW\Delta RQ}$) for net energy imbalance using diet-specific coefficients. Accelerometers

measured physical activity. Body composition changes were measured by dual-energy X-ray

absorptiometry which were combined with energy intake measurements to calculate energy

expenditure by balance (EE_{bal}).

Results: After transitioning from BD to KD, neither $EE_{chamber}$ nor EE_{bal} were significantly

changed ($\Delta EE_{chamber}$=24±30 kcal/d; p=0.43 and ΔEE_{bal}=-141±118 kcal/d; p=0.25). Similarly,

physical activity (-5.1±4.8%; p=0.3) and exercise efficiency (-1.6±2.4%; p=0.52) were not

significantly changed. However, EE_{DLW} was 209±83 kcal/d higher during the KD (p=0.023) but

was not significantly increased when adjusted for energy balance ($EE_{DLW\Delta RQ}$ =139±89 kcal/d;

p=0.14). After removing 2 outliers whose EE_{DLW} were incompatible with other data, EE_{DLW} and

$EE_{DLW\Delta RQ}$ were marginally increased during the KD by 126±62 kcal/d (p=0.063) and 46±65

kcal/d (p=0.49), respectively.

Conclusions: DLW calculations failing to account for diet-specific energy imbalance effects on

RQ erroneously suggest that very low carbohydrate diets substantially increase energy

expenditure.

Introduction

There is a great deal of interest in whether very low-carbohydrate diets offer a "metabolic

advantage" for weight loss by increasing total energy expenditure (1, 2) – a phenomenon

predicted by the carbohydrate-insulin model of obesity (3, 4). In support of this concept, a

highly-cited controlled feeding study in humans using the doubly labeled water (DLW) method

found a substantial increase in average daily energy expenditure during a very low-carbohydrate

diet compared with an isocaloric high-carbohydrate diet (5). However, such results are

inconsistent with several controlled feeding studies employing respiratory chambers that failed

to detect important energy expenditure differences between isocaloric diets varying in

carbohydrate content (6-8).

Our recent controlled feeding inpatient study was the first to compare daily energy expenditure

between isocaloric diets widely varying in carbohydrate content using both respiratory chambers

and DLW (9). Our primary result was a small but statistically significant increase daily energy

expenditure as measured by respiratory chamber ($EE_{chamber}$) of 57 ± 13 kcal/d (P = 0.0004) after

17 men ($25<BMI<35$ kg/m^2) transitioned from a one-month inpatient run-in period consuming a

moderate-carbohydrate baseline diet (BD) (50% carbohydrate, 35% fat, 15% protein) to a

subsequent month-long inpatient period consuming an isocaloric ketogenic diet (KD) (5%

carbohydrate, 80% fat, 15% protein). An exploratory aim of our study was to measure changes in

average daily energy expenditure by the DLW method (EE_{DLW}) and we reported a more

substantial increase of 151 ± 63 kcal/d (P = 0.03 uncorrected for multiple comparisons)

following the KD (9).

Some investigators who promote the carbohydrate-insulin model of obesity have discounted

these $EE_{chamber}$ results and have reinterpreted our exploratory EE_{DLW} data as underestimating the

true physiological effect of very low carbohydrate diets which have been suggested to increase

energy expenditure by 200-300 kcal/d (10, 11). However, there are methodological reasons why

the energy expenditure calculated by the DLW method is impacted by dietary carbohydrate

content that do not reflect true physiological differences. Specifically, the DLW method requires

an estimate of the overall metabolic fuel utilization of the body as quantified by the average daily

respiratory quotient (RQ). While it is widely known that RQ depends on the composition of the

diet as well as the overall state of energy balance, it is less well appreciated that the energy

imbalance adjustment of RQ also depends on dietary carbohydrate (12) and such adjustments are

rarely employed in DLW calculations.

Here, we reanalyzed our DLW data taking account of the diet-specific energy imbalance

adjustments of RQ used in the DLW method and highlight some key challenges to interpreting

DLW data when comparing energy expenditure differences between diets varying in

carbohydrate.

Methods

Details of the study and its approval by the Institutional Review Boards were reported previously

(9). Briefly, 17 men with BMI between 25-35 kg/m^2 provided informed consent and were

admitted as inpatients to metabolic wards where they consumed a standard baseline diet (BD)

composed of 50% energy from carbohydrate, 35% fat, and 15% protein for 4 weeks immediately

followed by 4 weeks of an isocaloric very low-carbohydrate, ketogenic diet (KD) composed of

5% carbohydrate, 80% fat, 15% protein. Body weight and height were measured to the nearest

0.1 kg and 0.1 cm, respectively, with subjects wearing a hospital gown and undergarments and

following an overnight fast. Body fat was measured using dual-energy X-ray absorptiometry

(DXA) scanners (Lunar iDXA, GE Healthcare, Madison, WI, USA).

Subjects spent two consecutive days each week residing in respiratory chambers to measure

energy expenditure ($EE_{chamber}$). As described previously (9), during the BD period, the daily

energy expenditure was calculated as follows:

$EE_{chamber} (\text{kcal}) = 3.88 \times VO_2(\text{L}) + 1.08 \times VCO_2(\text{L}) - 1.57 \times N(\text{g})$

where VO_2 and VCO_2 were the volumes of oxygen consumed and carbon dioxide produced,

respectively, and N was the 24hr urinary nitrogen excretion measured by chemiluminescence

(Antek MultiTek Analyzer, PAC, Houston, TX). During the KD period, the equations were

adjusted to account for 24-hour urinary ketone excretion, K_{excr}:

$EE_{chamber} (\text{kcal}) = 3.88 \times [VO_2(\text{L}) - 0.32\ (\text{L/g}) \times K_{excr}(\text{g})] + 1.08 \times VCO_2(\text{L}) - 1.57 \times N(\text{g}) + 1.39 \times K_{excr}(\text{g})$
119

Energy efficiency of physical activity was measured in the respiratory chamber with subjects

exercising at a constant, self-selected, level of moderate-intensity cycle ergometry.

Energy expenditure was calculated by energy balance (EE_{bal}) using the daily metabolizable

energy intake (EI) and the measured rates of change of the body energy storage pools determined

from measurements of fat mass (FM) and fat-free mass (FFM) by DXA at the beginning and end

of each two-week BD and KD period coincident with the DLW measurements as follows:

$EE_{bal} = EI - \rho_{FFM} \dfrac{dFFM}{dt} - \rho_{FM} \dfrac{dFM}{dt}$

where $\rho_{FM} = 9300$ kcal/kg is the energy density of body fat mass, $\rho_{FFM} = 1100$ kcal/kg is the

energy density of fat-free mass.

Energy expenditure was measured by DLW during the final two weeks of the BD and KD

periods to allow sufficient time for fluid shifts as subjects adjusted to each diet. Subjects drank

from a stock solution of ^2H_2O and $H_2{}^{18}O$ water in which 1 g of ^2H_2O (99.99% enrichment) was

mixed with 19 g of $H_2{}^{18}O$ (10% enrichment). An aliquot of the stock solution was saved for

dilution to be analyzed along with each set of urine samples. The water was weighed to the

nearest 0.1 g into the dosing container. The prescribed dose was 1.0 g per kg body weight and

the actual dose amounts were entered in the dose log. Spot urine samples were collected daily.

Isotopic enrichments of urine samples were measured by isotope ratio mass spectrometry. The

average CO_2 production rate (rCO_2) was corrected for previously administered isotope doses

(13), can be estimated from the rate constants describing the exponential disappearances of the

labeled ^{18}O and deuterated water isotopes (k_O and k_D) in repeated spot urine samples collected

over several days. We used the parameters of Racette et al. (14) with the weighted dilution space

calculation, R_{dil}, proposed by Speakman (15):

$$rCO_2 = (N/2.078)(1.007k_O - 1.007R_{dil}k_D) - 0.0246r_{GF}$$

144    $$r_{GF} = 1.05(1.007k_O - 1.007R_{dil}k_D)$$

$$R_{dil} = \left[(N_D/N_O)_{ave} \times n + 1.034 \times 255\right]/(n+255)$$

where r_{GF} accounts for the fractionation of the isotopes and $(N_D/N_O)_{ave}$ is the mean of the $N_D/$

N_O values from the $n=17$ subjects.

In our previous report (9), we used the 24hr respiratory quotient, RQ, measured during the

respiratory chamber stays to calculate energy expenditure (EE_{DLW}) during the baseline period

was calculated as:

$$EE_{DLW}(\text{kcal}) = \left[\frac{3.85}{RQ} + 1.07\right] \times rCO_2(\text{L})$$

During the KD period, EE_{DLW} was calculated as:

$$EE_{DLW}(\text{kcal}) = \left[\frac{3.85}{RQ} + 1.07\right] \times rCO_2(\text{L}) - [3.85 \times 0.32 + 1.39] \times K_{excr}(\text{g})$$

However, because the energy expended outside the chamber was significantly greater than inside

the chamber (9), the chamber RQ values did not accurately represent the overall RQ values

during the DLW period. In other words, the relative magnitude of energy imbalance (EB) during

the DLW period was different than the energy imbalance (EI-$EE_{chamber}$) measured during the

chamber stays. Therefore, we used the overall energy imbalance during the DLW period, defined

by the rate of change of body energy stores, to adjust the chamber RQ measurements by an

amount ΔRQ as described in the Online Supplemental Materials:

$$\Delta RQ = \lambda \times EB$$

where $\lambda = 5.28 \times 10^{-5}$ d/kcal for the BD and $\lambda = 9.17 \times 10^{-6}$ d/kcal for the KD.

Importantly, the value of λ is affected by dietary carbohydrate content, and, consequently, the

energy equivalence of CO_2 produced in the DLW calculations differs between diets varying in

carbohydrate for the same degree of energy imbalance.

Thus, the ΔRQ adjustment for each individual's diet-specific state of energy imbalance results in

the following calculation of DLW energy expenditure ($EE_{DLW\Delta RQ}$) during the baseline period:

$$EE_{DLW\Delta RQ}(\text{kcal}) = \left[\frac{3.85}{(RQ + \Delta RQ)} + 1.07\right] \times rCO_2(\text{L})$$

During the KD period, $EE_{DLW\Delta RQ}$ was calculated as:

$$EE_{DLW\Delta RQ}(\text{kcal}) = \left[\frac{3.85}{(RQ + \Delta RQ)} + 1.07\right] \times rCO_2(\text{L}) - \left[3.85 \times 0.32 + 1.39\right] \times K_{excr}(\text{g})$$

As opposed to our previous study (9) that reported results during the entire six-week period when

EI was held constant (i.e. the last two weeks of the BD phase and the entirety of the KD phase),

we now report results based upon data obtained only during the two-week DLW phase of both

BD and KD periods. In addition to allowing direct comparison with the coincident DLW

measures, focusing on the last two weeks of each diet allowed for two weeks to adapt to the BD

from the usual diet as well as to the KD from BD.

Statistical analysis was performed using a paired, two-sided t-test with significance declared at

the $p < 0.05$ threshold. The data are reported as mean±SE.

Results

During the final two weeks of the BD, EI was 2738±107 kcal/d which was significantly higher

than $EE_{chamber}$ = 2626±104 kcal/d ($p < 0.0001$). EE_{DLW} was 2964±126 kcal/d and significantly

higher than EI ($p = 0.011$). After adjusting for the energy imbalance, $EE_{DLW\Delta RQ}$ = 3045±135

kcal/d was significantly greater than EE_{DLW} ($p = 0.0003$). Energy expenditure calculated using EI

and the changes in body energy stores was $EE_{bal} = 3136\pm171$ kcal/d and was not significantly

different from EE_{DLW} (p=0.23) or $EE_{DLW\Delta RQ}$ (p=0.47). Compared to $EE_{chamber}$, EE_{DLW} was

338 ± 77 kcal/d higher (p=0.0005), $EE_{DLW\Delta RQ}$ was 419 ± 76 kcal/d higher (p<0.0001), and EEbal

was 509 ± 100 kcal/d higher (p<0.0001) indicating that subjects expended significantly more

energy outside the chamber during the BD phase.

During the final two weeks of the KD phase, EI was 2730 ± 110 kcal/d which, by design, was not

significantly different from the BD phase (p=0.16). Whereas we previously reported a transient

increase in $EE_{chamber}$ during the first two weeks after introducing the KD (9), neither $EE_{chamber}$

(2650 ± 89 kcal/d; p=0.43) nor EE_{bal} (2995 ± 160 kcal/d; p=0.25) were significantly different

during the last two weeks of the KD compared to the last two weeks of the BD coincident with

the DLW measurement periods (Figure 1A). Likewise, physical activity measured using an

accelerometer mounted on the hip was not significantly different (KD relative to BD, $-5.1\pm4.8\%$;

p=0.3); and energy efficiency of physical activity measured in the respiratory chamber with

subjects exercising at a constant level of moderate-intensity cycle ergometry was not

significantly different ($-1.6\pm2.4\%$; p=0.52) between the BD and KD phases.

Despite no significant differences in EI, $EE_{chamber}$, EE_{bal}, physical activity, or exercise efficiency

between the KD and BD phases (Figure 1B), EE_{DLW} was 209 ± 83 kcal/d higher during the KD

phase (p=0.023). The transition from the BD to KD coincided with increases in EE_{DLW} that were

in the opposite direction to EE_{bal}, indicating that the DLW calculations during the KD were

incommensurate with the changes in body weight and fat mass. After adjusting for the degree of

energy imbalance, $EE_{DLW\Delta RQ}$ was 139 ± 89 kcal/d higher during the KD, but this difference was

no longer significant (p=0.14) and was still in a direction opposite to the changes in body energy

stores.

There were two clear DLW outliers. The first outlier, "subject A", had an EE_{DLW} that was 1220

kcal/d greater than EI during the BD, and was 1751 kcal/d greater than EI during the KD despite

slight body weight and fat mass gains during these periods. In contrast, the $EE_{chamber}$

measurements for this subject were 173 kcal/d less than EI during the BD, and 65 kcal/d less

than EI during the KD. The second outlier, "subject B", had an EE_{DLW} during the BD that was

only 123 kcal/d higher than $EE_{chamber}$, but during the KD his EE_{DLW} increased by 1136 kcal/d

which was ~3 standard deviations greater than the mean increase in $EE_{DLW,}$ suggesting severe

negative energy balance despite the subject gaining weight during this period and an $EE_{chamber}$

increasing by only 72 kcal/d. Tables S1-S4 in the Online Supplemental Materials provide

summary data on the energy expenditure comparisons between BD and KD phases with and

without the exclusion of these subjects. After excluding these subjects, the increase in EE_{DLW}

during the KD was 126±62 kcal/d (p=0.063) and $EE_{DLW\Delta RQ}$ increased by only 46±65 kcal/d

(p=0.49), neither of which were significant (Table S4)

Discussion

As previously reported (9), our inpatient isocaloric feeding study showed that four days of

respiratory chamber measurements that were coincident with each two-week DLW period did

not detect significant changes in energy expenditure that were attributable to diet composition.

However, the increase in EE_{DLW} after transitioning to the KD was substantially greater than

could be accounted for by changes in body energy stores. In the present analysis, after employing

diet-specific RQ adjustments in the DLW calculations to account for energy imbalance (12), we

found that the calculated increase in DLW energy expenditure during the KD was significantly

attenuated and was no longer statistically significant. After removal of two clear DLW outliers,

the RQ-adjusted increase in DLW energy expenditure was further reduced to a marginal ~50

kcal/d higher during the KD.

We previously hypothesized that the discrepancy between the respiratory chamber and DLW

measurements was due to increased physical activity outside the respiratory chamber during the

KD (9). However, this potential explanation does not agree with the lack of increase in

objectively measured physical activity. Also, the energy expended to perform the same low-

intensity exercise in the respiratory chamber was not significantly changed by the KD, making it

unlikely that the energy efficiency of skeletal muscle contraction had been altered after

transitioning to the KD. Thus, the EE_{DLW} measurements were discordant with several

independent measures indicating that the KD did not result in significant energy expenditure

changes. The appropriately adjusted RQ values resulted in DLW data more commensurate with

the other data, especially after removing the pair of outliers.

Another methodological concern with the DLW method is the theoretical possibility that CO_2

production rates can be influenced by the fluxes through biosynthetic pathways that likely vary

substantially depending on the carbohydrate content of the diet, especially the de novo

lipogenesis pathway (16-18). However, the magnitude of this potential bias in humans is thought

to be relatively small, amounting to an energy expenditure difference of only about 30-60 kcal/d

(see Online Supplemental Materials). Interestingly, even this small systematic bias would have

been sufficient to nullify the statistical significance of the observed increased EE_{DLW} in our study

and the effect size was similar to the nonsignificant increase in the $EE_{DLW\Delta RQ}$ difference after

removal of the outliers. Individual variability of de novo lipogenesis changes after transitioning

to the KD might explain the diminished correlations during the KD between the DLW

measurements with $EE_{chamber}$ and EE_{bal}. To directly address this question, we need a validation

study in humans with several days of simultaneous respiratory chamber and DLW measurements

high- and low-carbohydrate diets, ideally with simultaneous measurements of de novo

lipogenesis.

Ebbeling et al. (5) reported significant ~200-300 kcal/d differences in EE_{DLW} during the last two

weeks in each of three consecutive month-long periods during which outpatient subjects

consumed a very low-carbohydrate diet (10% of energy) compared to isocaloric diets with

moderate (40% of energy) or high (60% of energy) carbohydrate in random order. However,

resting energy expenditure differences were much smaller (~29-67 kcal/d) and objectively

measured physical activity was not significantly different among the diets. Despite claims of

weight stability, reported energy intake was several hundred calories below the measured EE_{DLW}

but RQ was estimated without adjustment for the subjects' state of energy balance. Therefore,

similar to the current study, the diet specific effects of energy imbalance on RQ likely

contributed to the observed differences in EE_{DLW}. Thus, the data of Ebbeling et al. could be

interpreted as supporting the possibility that the EE_{DLW} calculations using RQ values unadjusted

for energy imbalance in a diet-specific way were systematically biased such that the isocaloric

very low-carbohydrate diet resulted in systematically increased EE_{DLW}.

In contrast, Bandini et al. (19) found in an outpatient study that EE_{DLW} was lower during the very

low-carbohydrate diet (~7% of energy) as compared to a high-carbohydrate diet (~83% of

energy), but this reduction was attributed to decreased physical activity because the subjects

reported nausea and lethargy on the low-carbohydrate diet. No significant differences in REE

were found. Stubbs et al. (20) found no significant difference between EE_{DLW} using a narrower

range of diets with 29-67% of energy as carbohydrate. But the diets were fed *ad libitum* and

energy intake on the low-carbohydrate diet was greater than the moderate-carbohydrate diet

which was greater than the high-carbohydrate diet. Thus, the variation in total carbohydrate

content of the test diets was attenuated such that daily carbohydrate intake varied by only ~22%

of the mean energy intake between diets which may not have been a sufficient range to observe a

systematic bias of the DLW method. Furthermore, the positive energy balance with the low-

carbohydrate diet increased RQ whereas the negative energy balance with the high-carbohydrate

diet decreased RQ. Thus, the RQ differences during consumption of these diets were attenuated

by the differences in energy balance.

It is important to emphasize that our study was not intended to be a DLW validation study and

there were several limitations. The DLW measurements were not pre-specified as either primary

or secondary endpoints of the study. Whereas respiratory chamber measurements have high-

precision, with an intrasubject coefficient of variation of $EE_{chamber}$ ~2-3% (21), the DLW method

is less precise, with an intrasubject coefficient of variation of energy expenditure of ~ 8-15%

(22). Therefore, the relatively large inherent variability of the DLW method may have led to an

apparent increase in EE_{DLW} during the KD simply by chance (type-1 error). However, we cannot

definitively exclude the possibility of a real increase in energy expenditure, especially at an

effect size of ~50-140 kcal/d after excluding two likely DLW outliers or using diet-specific

adjustments of the DLW calculations to account for the energy imbalance.

The DLW method has been validated during 30% caloric restriction with a 55% carbohydrate

diet (23) and agrees with our result that EE_{DLW} and $EE_{DLW\Delta RQ}$ were not significantly different

from EE_{bal} during the BD diet phase. Nevertheless, the calculated EE_{bal} values are somewhat

uncertain because DXA has a limited ability to precisely and accurately detect small changes in

body energy stores (24). We cannot rule out the possibility that the KD resulted in undetected

increases in activity-related energy expenditure that were undetected by accelerometers. Finally,

the order of the diets was not randomized, and it is possible that the elevated EE_{DLW} occurred

simply because the KD followed the BD. Indeed, others have reported a greater metabolic rate

during a low-carbohydrate diet when it followed a high-carbohydrate diet as compared with the

reverse order (25).

In summary, our data illustrate the challenges of using the DLW method to estimate energy

expenditure differences between diets varying widely in the proportion of carbohydrate. We urge

caution when interpreting such data, especially if the DLW calculations do not appropriately

adjust RQ for energy imbalance or if the DLW results are not corroborated by quantitatively

commensurate observations of energy intake and body composition change.

Acknowledgements

Nutrition Sciences Initiative (NuSI), convened the research team, helped to formulate the

hypothesis and provided partial funding. NuSI and its scientific advisors were given the

opportunity to comment on the study design and the manuscript, but the investigators retained

full editorial control. The authors' contributions were as follows: KDH, KC, RLL, MLR, MR,

SRS, and ER designed the study and conducted the research; KDH and JG analyzed the data;

KDH, KC, RLL, MLR, MR, SRS, and ER wrote the manuscript; KDH had primary

responsibility for the final content. The authors have no conflicts of interest. We thank Jim

DeLany, Herman Pontzer, and John Speakman for helpful discussions regarding the DLW

method and its interpretation. The clinical study protocol and deidentified individual data are

currently available for download on the Open Science Framework website at https://osf.io/h4xju/

for any purpose by anyone.

References

1.   Fine EJ, Feinman RD. Thermodynamics of weight loss diets. Nutr Metab (Lond)
2004;1(1):15.
2.   Manninen AH. Is a calorie really a calorie? Metabolic advantage of low-carbohydrate
diets. J Int Soc Sports Nutr 2004;1(2):21-6.
3.   Ludwig DS, Ebbeling CB. The Carbohydrate-Insulin Model of Obesity: Beyond
"Calories In, Calories Out". JAMA internal medicine 2018;178(8):1098-103. doi:
10.1001/jamainternmed.2018.2933.
4.   Ludwig DS, Friedman MI. Increasing adiposity: consequence or cause of overeating?
JAMA 2014;311(21):2167-8. doi: 10.1001/jama.2014.4133.
5.   Ebbeling CB, Swain JF, Feldman HA, Wong WW, Hachey DL, Garcia-Lago E, Ludwig
DS. Effects of dietary composition on energy expenditure during weight-loss
maintenance. Jama 2012;307(24):2627-34.
6.   Hall KD. A review of the carbohydrate-insulin model of obesity. Eur J Clin Nutr
2017;71(3):323-6. doi: 10.1038/ejcn.2016.260.
7.   Hall KD, Guo J. Obesity Energetics: Body Weight Regulation and the Effects of Diet
Composition. Gastroenterology 2017;152(7):1718-27 e3. doi:
10.1053/j.gastro.2017.01.052.
8.   Hall KD, Guyenet SJ, Leibel RL. The Carbohydrate-Insulin Model of Obesity Is Difficult
to Reconcile With Current Evidence. JAMA internal medicine 2018;178(8):1103-5. doi:
10.1001/jamainternmed.2018.2920.
9.   Hall KD, Chen KY, Guo J, Lam YY, Leibel RL, Mayer LE, Reitman ML, Rosenbaum
M, Smith SR, Walsh BT, et al. Energy expenditure and body composition changes after
an isocaloric ketogenic diet in overweight and obese men. Am J Clin Nutr
2016;104(2):324-33. doi: 10.3945/ajcn.116.133561.

10. Friedman MI, Appel S. Energy expenditure and body composition changes after an
isocaloric ketogenic diet in overweight and obese men: a secondary analysis of energy
expenditure and physical activity. bioRxiv 2018. doi: 10.1101/383752.
11. Ludwig DS, Ebbeling CB. Raising the bar on the low-carbohydrate diet. Am J Clin Nutr
2016;104(5):1487-8. doi: 10.3945/ajcn.116.142182.
12. Elia M. Energy equivalents of CO2 and their importance in assessing energy expenditure
when using tracer techniques. Am J Physiol 1991;260(1 Pt 1):E75-88. doi:
10.1152/ajpendo.1991.260.1.E75.
13. Bhutani S, Racine N, Shriver T, Schoeller DA. Special Considerations for Measuring
Energy Expenditure with Doubly Labeled Water under Atypical Conditions. Journal of
obesity & weight loss therapy 2015;5(Suppl 5). doi: 10.4172/2165-7904.S5-002.
14. Racette SB, Schoeller DA, Luke AH, Shay K, Hnilicka J, Kushner RF. Relative dilution
spaces of 2H- and 18O-labeled water in humans. Am J Physiol 1994;267(4 Pt 1):E585-
90.
15. Speakman JR. Doubly labelled water: Theory and practice. London: Chapman & Hall,
1997.
16. Coward WA. Measurement of energy expenditure: the doubly-labelled-water method in
clinical practice. Proc Nutr Soc 1991;50(2):227-37.
17. Haggarty P. Effect of isotope sequestration and exchange. Edtion ed. In: Prentice AM, ed.
The doubly labeled water method for measuring energy expenditure: Technical
recommendations for use in humans: A concensus report by the IDECG working group.
Vienna: International atomic energy agency, 1990.
18. Haggarty P, McGaw BA, Fuller MF, Christie SL, Wong WW. Water hydrogen
incorporation into body fat in pigs: effect on double/triple-labeled water method. Am J
Physiol 1991;260(3 Pt 2):R627-34. doi: 10.1152/ajpregu.1991.260.3.R627.
19. Bandini LG, Schoeller DA, Dietz WH. Metabolic differences in response to a high-fat vs.
a high-carbohydrate diet. Obes Res 1994;2(4):348-54.
20. Stubbs RJ, Ritz P, Coward WA, Prentice AM. Covert manipulation of the ratio of dietary
fat to carbohydrate and energy density: effect on food intake and energy balance in free-
living men eating ad libitum. Am J Clin Nutr 1995;62(2):330-7.
21. Ravussin E, Lillioja S, Anderson TE, Christin L, Bogardus C. Determinants of 24-hour
energy expenditure in man. Methods and results using a respiratory chamber. J Clin
Invest 1986;78(6):1568-78.
22. Black AE, Cole TJ. Within- and between-subject variation in energy expenditure
measured by the doubly-labelled water technique: implications for validating reported
dietary energy intake. Eur J Clin Nutr 2000;54(5):386-94.
23. de Jonge L, DeLany JP, Nguyen T, Howard J, Hadley EC, Redman LM, Ravussin E.
Validation study of energy expenditure and intake during calorie restriction using doubly
labeled water and changes in body composition. Am J Clin Nutr 2007;85(1):73-9.
24. Pourhassan M, Schautz B, Braun W, Gluer CC, Bosy-Westphal A, Muller MJ. Impact of
body-composition methodology on the composition of weight loss and weight gain. Eur J
Clin Nutr 2013;67(5):446-54. doi: 10.1038/ejcn.2013.35.
25. Hron BM, Ebbeling CB, Feldman HA, Ludwig DS. Relationship of insulin dynamics to
body composition and resting energy expenditure following weight loss. Obesity (Silver
Spring) 2015;23(11):2216-22. doi: 10.1002/oby.21213.
408
409

Figure 1. A) Neither energy expenditure by respiratory chamber (EE$_{chamber}$), nor energy

expenditure by balance (EE$_{bal}$) were significantly different during the baseline diet (BD) versus

the ketogenic diet (KD). However, energy expenditure by doubly labeled water (EE$_{DLW}$) was

significantly greater during the KD, but not after adjustment of the respiratory quotient (RQ) to

account for the differential diet effect of energy imbalance (EE$_{DLW\Delta RQ}$) or after removal of two

outliers whose DLW data were incomparible with other measurements (EE$_{DLW\Delta RQ}$ - Outliers). B)

417   Differences in $EE_{chamber}$, EE_{bal}, EE_{DLW}, $EE_{DLW\Delta RQ}$, and $EE_{DLW\Delta RQ}$ - Outliers between KD and BD

418   phases. NS = not significant, $p > 0.05$.