**Molecular evolution of non-fertilizing sperm in Lepidoptera suggests minimal direct**
**involvement in sperm competition**

Keywords: sperm dimorphism, population genetics, *Manduca sexta*, *Danaus plexippus*

Andrew J. Mongue[1*], Megan E. Hansen[1], Liuqi Gu[1], Clyde E. Sorenson[2], and James R. Walters[1]

[1]University of Kansas, Department of Ecology and Evolutionary Biology

[2]North Carolina State University, Department of Entomology

[*]Corresponding author contact: ajmongue@gmail.com

**Abstract**

Sperm are among the most variable cells in nature. Some of this variation results from non-adaptive

errors in spermatogenesis, but many species consistently produce multiple sperm morphs, the adaptive

significance of which remains unknown. Here, we investigate the evolution of dimorphic sperm in

Lepidoptera, the butterflies and moths. Males of this order produce both fertilizing sperm and a

secondary, non-fertilizing type that lacks DNA. Previous organismal studies suggested a role for non-

fertilizing sperm in sperm competition, but this hypothesis has never been evaluated from a molecular

framework. We combined published datasets with new sequencing in two species, the monandrous

Carolina sphinx moth and the highly polyandrous monarch butterfly. Based on population genetic

analyses, we see evidence for increased adaptive evolution in fertilizing sperm, but only in the

polyandrous species. This signal comes primarily from a decrease in non-synonymous polymorphism in

sperm proteins compared to the rest of the genome, suggesting stronger purifying selection, consistent

with selection via sperm competition. Non-fertilizing sperm proteins, in contrast, do not show an effect

of mating system and do not appear to evolve differently from the background genome in either species,

arguing against the involvement of non-fertilizing sperm in direct sperm competition. Based on our

results and previous work, we suggest that non-fertilizing sperm may be used to delay female remating

in these insects and decrease the risk of sperm competition rather than directly affect its outcome.

**Introduction**

Sperm cells display remarkable diversity throughout the animal kingdom (Pitnick, Hosken, & Birkhead,

2009), from small and plentiful to gigantic and few (Pizzari, 2006) to super-structure-forming (Higginson,

Miller, Segraves, & Pitnick, 2012). This variation exists at every level, from fixed differences between

species to variability within individual males (John Buckland-Nicks, 1998; Marks, Biermann, Eanes, &

Kryvi, 2008; Sasakawa, 2009; Swallow & Wilkinson, 2002; Tavares-Bastos, Teixeira, Colli, & Báo, 2002). In

many independently evolved cases, males consistently produce two different sperm types, a

phenomenon known as sperm dimorphism. In all cases examined, only one of the two sperm morphs is

capable of fertilization (Bressac et al., 1991; Carcupino, Baldacci, Fausto, Scapigliati, & Mazzini, 1999;

Eckelbarger, Young, & Cameron, 1989; Sasakawa, 2009; Wilms, 1986).  The evolutionary causes and

consequences of variation in sperm morphology, both within and between morphs, are immediately

intriguing. As gametes, these cells are the final step in the long chain of events leading to reproductive

success or failure. Why should such important components of fitness be so variable?

Much of this morphological diversity within morphs can be attributed to deleterious variation, *e.g.*

genetic defects (Chenoweth, 2005) or age-related decline in sperm quality (Preston, Saint Jalme, Hingrat,

Lacroix, & Sorci, 2015). This deleterious variation has been shown to be inversely correlated with rates

of sperm competition between species; taxa that experience more sperm competition tend to have less

morphologically variable sperm at both population and individual levels (Kleven, Laskemoen, Fossøy,

Robertson, & Lifjeld, 2008). In other words, sperm often vary *in spite of* constraint imposed by their

reproductive importance. In species with high rates of polyandry, postcopulatory selection through

sperm competition and cryptic female choice weeds out the suboptimal sperm variants, at least for

fertilizing sperm (Birkhead, 1998; Immler, Calhim, & Birkhead, 2008).

Production of multiple sperm morphs, conversely, is often posited to be adaptive in some way. The very

fact that sperm dimorphism has repeatedly evolved suggests that it has some fitness benefit. Most

commonly, non-fertilizing sperm in dimorphic systems are proposed to be specialized agents of male-

male competition, acting as final combatants in the struggle for reproductive success (J Buckland-Nicks,

Bryson, Hart, & Partridge, 2010; John Buckland-Nicks, 1998; Swallow & Wilkinson, 2002). Indeed, some

have suggested that sperm dimorphism allows specialization in the non-fertilizing sperm for a

competitor-inhibiting function, sometimes called "kamikaze sperm" (Baker & Bellis, 1989). Although this

hypothesis has fallen out of favor, it was proposed and mainly evaluated in the context of mammalian

sperm (A. Harcourt, 1991; A. H. Harcourt, 1989; Moore, Martin, & Birkhead, 1999), where non-fertilizing

sperm are not usually differentiated from fertilizing sperm in a sophisticated way.

One of the most extreme cases of sperm dimorphism occurs in butterflies and moths (Lepidoptera). In

nearly all species of this order, males produce both fertilizing (**eupyrene**) sperm and a second type

(**apyrene**) that lacks a nucleus and nuclear DNA (Meves, 1902). The function of apyrene sperm is poorly

understood, but because it lacks DNA, it is clearly incapable of fertilizing eggs. Nevertheless, it does not

appear to be the result of errors in spermatogenesis; apyrene sperm production is hormonally regulated

and occurs in a developmentally predictable way, implying a novel gain of function in these insects

(Friedlander, 1997). Organismal studies have demonstrated that males can control the ratio of the two

sperm types in their ejaculate and typically transfer to females 10 to 20 times as much apyrene sperm as

eupyrene sperm, depending in part on the female's past mating history (Oberhauser, 1988). These

observations have led some to suggest that apyrene sperm play a specialized role in sperm competition

(Silberglied, Shepherd, & Dickinson, 1984), yet there remain several other competing hypotheses for

apyrene sperm function that have not been resolved through organismal observations and experiments

(Swallow & Wilkinson, 2002).

Recently, characterizations of the proteins found in lepidopteran sperm has opened a new avenue to

assess their evolution and function (Whittington et al., 2017; Whittington, Zhao, Borziak, Walters, &

Dorus, 2015). Proteomic studies have revealed distinct protein profiles for these two cell types

(Whittington, Karr, Mongue, Walters, & Dorus, in press). In both morphs, these proteins are retained

through maturation, and, in the case of apyrene sperm, the discarding of the nucleus. Because distinct

cellular functions are ultimately the product of their expressed protein complement, the class of

proteins uniquely found in apyrene sperm make logical targets for understanding the function of these

cells from a molecular perspective.

At the molecular level, sperm and other reproductive proteins are often observed to evolve rapidly

(Civetta & Singh, 1995; Dorus, Evans, Wyckoff, Sun, & Lahn, 2004; Willie J. Swanson & Vacquier, 2002).

For certain reproductive proteins, like sperm-egg interaction pairs, there is compelling evidence that

adaptive co-evolution drives this accelerated change (Herberg, Gert, Schleiffer, & Pauli, 2018; W J

Swanson & Vacquier, 1998). Yet there are also many instances of reproductive proteins that diverge

quickly because of relaxed purifying selection owing to expression in a single sex instead of the whole

population (Barker, Demuth, & Wade, 2005; Wade, Priest, & Cruickshank, 2008).  Many other factors,

including number of protein-protein interactions or importance of reproductive role, can also act to

shape the intensity of positive or purifying selection on reproductive proteins (Schumacher, Rosenkranz,

& Herlyn, 2014; Schumacher, Zischler, & Herlyn, 2017). Recent theoretical work has formalized the

prediction that strong purifying selection on sperm proteins should depend on high rates of polyandry to

generate sperm competition (Dapper & Wade, 2016). Thus, with the appropriate datasets, the degree of

each sperm morph's role in sperm competition can be assessed via molecular tests of evolution.

In this study, we report the first molecular evolutionary analyses of dimorphic sperm. We assessed

patterns of both polymorphism and divergence among sperm proteins from both eupyrene and apyrene

sperm using proteomic datasets of two species: the monarch butterfly, *Danaus plexippus*, and the

Carolina sphinx moth, *Manduca sexta* (Whittington et al., in press). North American monarchs spend

time at incredibly high density in overwintering colonies in Mexico and California (Urquhart, 1976) and,

owing to these unique population dynamics, have some of the highest female remating rates observed

in Lepidoptera. Female monarchs mate an average of 2.6 times (and up to 14 times) in overwintering

colonies in the wild (Hill Jr., Wenner, & Wells, 1976; Smith, 1984), creating ample opportunity for sperm

competition. In contrast, Carolina sphinx moths are typically monandrous (Snow et al., 1974), making

sperm competition rarely relevant as a selective force. Taking advantage of this contrast, we investigate

the differences in patterns of selection between the two sperm morphs in each species to assess the

role of apyrene sperm in sperm competition. If apyrene sperm are involved in sperm competition, their

proteins should show evidence of stronger purifying selection in the monarch butterfly. To complete

these analyses, we have generated the first published set of whole-genome resequencing data for

*Manduca sexta* from a wild population. To test the general predictions for relaxed selection in sex-

limited proteins, we used RNA-seq gene expression datasets from previously published data for Carolina

sphinx moths (Cao & Jiang, 2017) and newly generated data for the monarch butterfly.

**Materials and Methods**

*Sources of data*

We used gene sets from the published genomes of each species (Kanost et al., 2016; Zhan & Reppert,

2013) with sperm genes identified from their respective proteomes (Whittington, Karr, Mongue, Walters,

& Dorus, in press) . We inferred selection from patterns of polymorphism and divergence from

congeners using whole genome Illumina resequencing data for both species: a previously published

dataset for North American monarch butterflies (Zhan et al., 2014) and a new dataset of North

Carolinian sphinx moths. Focal moths were collected with a mercury vapor light trap in July of 2017 in

Rocky Mount, North Carolina (see supplemental table S1 for sequencing summary statistics and

accessions). Divergences were called by comparison to the queen butterfly (*Danaus gilippus*, previously

published in Zhan et al. (2014)) for monarchs, and the five-spotted hawkmoth *(Manduca*

*quinquemaculata*, sequenced for this project) for the Carolina sphinx moth.

In both focal species, we used twelve wild-caught individuals for sampling of polymorphism. In the case

of Carolina sphinx moths, these were twelve males caught over the course of three nights. The sex-

biased sampling reflects a sex bias in dispersal and collection at the light trap. In the case of monarchs,

samples were selected based on depth of sequencing coverage in the published dataset and included 8

females and 4 males from the panmictic North American migratory population. This mixed-sex sampling

added the complication of unequal sampling between the autosomes (n = 24) and Z sex chromosome (n

= 16). Despite the male-biased gene accumulation on the Z chromosome, the vast majority of sperm

genes (92% in the Carolina sphinx, 90% in the monarch) are autosomal in both species (Mongue &

Walters, 2017). Due to the sampling complication and limited inference to be gained from Z-linked

genes, we focused on the autosomal genes in both species in subsequent analyses.

*SNP-based methods*

We aligned sequenced reads with bowtie2 (Langmead & Salzberg, 2012)  for conspecifics to their

reference genome or with stampy (Lunter & Goodson, 2011) with an increased allowance for

substitution for heterospecific alignments. Alignments were taken through GATK's best practices

pipeline (McKenna et al., 2010), including hard filtering, to yield a set of high quality variants both within

and between species. Effect-class of each variable site (synonymous, non-synonymous, intergenic, etc.)

was determined using custom databases for the two species created with SnpEff (Cingolani et al., 2012).

Annotated SNPs were curated to remove false divergences (ancestral polymorphism) and then

differences in adaptive evolution were calculated using an estimator of the neutrality index to calculate

α, the proportion of substitutions driven by adaptive evolution (Stoletzki & Eyre-Walker, 2011). This

form of α corrects the inherent bias in a ratio of ratios while also allowing summation across multiple

genes to reduce noise associated with small numbers in count data. For any set of *i* genes with non-zero

counts of synonymous (s) polymorphism (P) and divergence (D):

$$\alpha = \ 1 - \frac{\sum (Dsi * Pni)/(Psi + Dsi)}{\sum (Dni * Psi)/(Psi + Dsi)}$$

This statistic was calculated with custom scripts in R (R Core Team, 2017).

*Assessment of adaptive evolution and statistical significance*

In each analysis, we calculated α for a biologically meaningful set of genes, *e.g.* the sperm proteome and

the background genome, and generated a test statistic from the absolute difference of the two point-

estimates. To determine significance, we combined the two sets and randomly assigned genes into two

new sets of sizes equal to the originals. The difference of these two datasets was determined and the

process was repeated for 50,000 permutations to build a distribution of differences between the point

estimates of two gene sets of these relative sizes. The p-value was taken as the proportion of times a

greater absolute difference was observed between the two random data sets than between the original

sets.

We used this permutation approach to make within-species comparisons of α for several different

groupings of genes. We first examined differences between the whole sperm proteome and background

genome (i.e. all autosomal non-sperm proteins). Next, we considered differences between sperm

homologs and sperm proteins unique to one species to assess how selection acted on the same genes in

different species. We identified sperm homologs as predicted orthologs that are present in the sperm of

both species, with orthology predicted via the proteinOrtho pipeline, as previously reported in Mongue

& Walters (2017). Unique sperm proteins may or may not have an ortholog in the other species but are

present in the sperm of only one species. Finally, we compared among proteins grouped by their

presence in apyrene versus eupyrene sperm. To do so, we classified sperm proteins into three subsets:

specific to eupyrene sperm, specific to apyrene sperm, or shared in both types. Pairwise comparisons

were made between each subset. For these analyses, we did not consider orthology status owing to the

reduction in power that would accompany multiple layers of subdivision of the dataset. For the whole

proteome and morph subset comparisons, we further assessed the relative contributions of

synonymous and non-synonymous polymorphism and divergence to the α calculation, using a Wilcoxon-

Mann-Whitney test to assess significant differences.

*Site-frequency-based methods*

We also investigated molecular evolution by leveraging site-frequency-spectrum-based approaches as

complimentary evidence. Owing to the redundancy in results, we have included these analyses in the

supplement rather the main text. In brief, we used the population genetics software suite ANGSD

(Korneliussen, Albrechtsen, & Nielsen, 2014) to generate site frequency spectra at putatively neutral

(four-fold degenerate) and selected (zero-fold-degenerate) sites in the genome. We unfolded site

frequency spectra and analyzed these spectra with the software polyDFE (Tataru, Mollion, Glémin, &

Bataillon, 2017) to examine rates of adaptive evolution in the whole sperm proteomes and background

genomes with a more complex likelihood model that corrects for effects of demography and potential

misattribution of ancestral state.

*Investigation of sex-limited and tissue-specific expression*

Next, we used RNA-seq data to assess whether or not differences in tissue specificity of expression

impacted our results from the sperm proteomes in these taxa. For *Manduca sexta*, there already existed

a wealth of tissue-specific data at multiple developmental timepoints (Cao & Jiang, 2017). Because we

were primarily interested in sperm involvement, we focused on data from adult males, specifically RNA

from the testes, head, thorax, and gut. Expression (measured as fragments per kilobase of transcript per

million mapped reads, FPKM) was averaged across biological replicates where available in this species.

Monarchs had no comparable published data, so we generated separate RNA-seq data sets from the

head, thorax, gut, testes, and accessory gland of three adult males (summarized in Table S2 with

accessions).

We quantified tissue-specificity of expression using the *specificity metric* (SPM) statistic, a ratio ranging

from 0 to 1 indicating the proportion of gene expression occurring in a given focal tissue (Kryuchkova-

Mostacci & Robinson-Rechavi, 2017). For instance, a gene with SPM = 0.8 for the testes shows 80% of its

total expression across all sampled tissues in the testes. This same gene would have a much lower SPM

value in head, thorax, or other tissues. We observed a bimodal distribution of tissue specificities, which

allowed us to bin genes into one of two classes: those that displayed low levels of specificity (SPM < 0.5)

and those that displayed high levels (SPM > 0.5). After separating genes by specificity, we calculated α

for three classes of genes in these two specificity bins.

We had two goals with these analyses: (1) to determine if patterns of adaptive evolution between

classes remained the same at both low- and high-specificities and (2) if α increased within a class of

genes at higher specificity compared to low. First, we considered background genome genes (i.e. non-

sperm genes) ranked by maximum specificity observed in the head, thorax, or gut for each of these

genes. Next, we considered only genes identified in the sperm proteome and ranked them by SPM in the

testes. Finally, for putatively male-limited non-sperm genes, we excluded sperm proteome genes and

considered again those ranked by specificity in the testes (or testes and accessory glands for monarchs).

As with our other α calculations, we used non-parametric bootstrapping to generate 95% confidence

intervals. For cases in which confidence intervals overlapped, we assessed significance with permutation

testing. These analyses were completed with custom R scripts.

*Demographic estimates*

Finally, to contextualize the previous analyses and take full advantage of our newly-generated data, we

characterized present and historical population sizes of our study species from genomic data. Using

folded four-fold degenerate site frequency spectra, we estimated neutral coalescence patterns with

Stairway Plot (Liu & Fu, 2015). For estimated generation time, we used four generations per year for

monarchs and three for the Carolina sphinx moth. For mutation rate, we chose the estimate $2.9*10^{-9}$

from the butterfly *Heliconius melpomene*, the closest relative with a spontaneous mutation rate

estimate (Keightley et al., 2015).

**Results**

*Differences Between Sperm Proteins and the Background Genome*

First, we considered the sperm proteome as a whole (*i.e.* all apyrene, shared, and eupyrene proteins)

and compared adaptive evolution of genes found in sperm to those in the background genome, defined

as all autosomal protein coding genes not present in the sperm proteome. Z-linked genes were excluded

from the analysis. We counted and classified synonymous and non-synonymous single nucleotide

polymorphisms within species and divergences to a congener (*Danaus gilippus* for the monarch, and

*Manduca quinquemaculata* for the Carolina sphinx). These quantities were used to generate an estimate

of the proportion of adaptive substitutions (α) per gene-class for both the sperm proteome and the

background genome. We found no difference in α between the sperm proteome and the rest of the

genome in the Carolina sphinx (p = 0.40892 by permutation testing, Figure 1A, left); for monarchs,

however, the sperm proteome showed a significantly greater proportion of adaptive substitutions than

the rest of the genome (p = 0.00006, Figure 1A, right). Note that in the strict sense, negative α values are

not biologically meaningful and likely point to an abundance of weakly deleterious variants within

populations or complex demographic histories (Eyre-Walker & Keightley, 2009); nevertheless, these

confounding variables should not differentially affect genes within species, so our observed differences

point to true differences in selection in gene sets.

To better understand the relative roles of polymorphism and divergence in sperm and background

genes, we investigated the individual components of α: counts of non-synonymous polymorphism (Pn),

synonymous polymorphism (Ps), non-synonymous divergence (Dn), and synonymous divergence (Ds).

We compared the scaled estimates of each (*e.g.* non-synonymous polymorphisms per non-synonymous

site) to the background genome within each species using a Wilcoxon-Mann-Whitney test (Figure 1B).

We found no differences between sperm and the background for any class of variants in *M. sexta* (Pn: W

= 3014100, p = 0.5964; Ps: W = 2879300, p = 0.1830; Dn: W = 3068300, p = 0.2009; Ds: W = 2895700, p =

0.2686). The signal for elevated α in monarch sperm primarily reflects non-synonymous polymorphism,

which was greatly depressed (W = 3062400; p = $3.224 * 10^{-11}$), as would be expected under strong

purifying selection, while other classes were comparable between sperm and the background genome

(Ps: W = 2684200, p = 0.2720; Dn: W = 2506400, p = 0.1300; Ds: W = 2544400, p = 0.3437).

Next, we leveraged orthology, as established by Whittington *et al.* (2017), to test for differences in

mating system while controlling for the effects of sperm proteome content. Substantial numbers of

orthologous proteins are found in the sperm proteomes of both species, which we hereafter referred to

as *sperm homologs*. Sperm homologs offer the opportunity to directly assess the selective pressures

experienced by the same genes with putatively conserved function but found in species with different

levels of postcopulatory selection. Nearly half of the monarch sperm proteome (~42%, 216 genes, Figure

2A) shares an ortholog in the sperm proteome of *M. sexta*; reciprocally, there are 236 genes (37%) in the

Carolina sphinx sperm proteome that share an ortholog in the monarch sperm proteome; these

numbers are not equal due to lineage-specific duplications among sperm homologs creating a few cases

of one-to-many orthology. We tested for differences in adaptive evolution between sperm homologs

and sperm proteins unique to one species (orthology outside of sperm or no detectable orthology). In

Carolina sphinx moths, genes of these two classes did not differ in the proportion of adaptive

substitutions with permutation testing (p = 0.6174, Figure 2B). In monarchs, we detected an increased

proportion of adaptive substitution in the sperm homologs compared to unique proteins (p = 0.0372,

Figure 2B). Comparing between species, sperm homologs had much higher α values in monarchs than in

Carolina sphinx moths (p = 0.00008), while genes with unique expression in either species did not show

differences between species (p = 0.5922). Thus, the same sperm proteins appear to be evolving more

adaptively in the polyandrous species.

*Site-frequency based methods*

We also took a likelihood approach to modeling adaptive evolution using site frequency spectra

generated from the same samples we used for SNP-counting. These results are detailed in the

supplement. In short though, we found a shift in the predicted distribution of fitness effects of new

mutations in monarch sperm proteins compared to the background consistent with stronger purifying

selection (Figure S1) and drastically higher α in sperm genes in monarchs alone (Figure S2).

*Patterns of Adaptive Evolution in Sex-Specific Tissues*

Next, we used RNA-seq data to examine the effect of tissue-specificity on selection in these insects.

With these data, we calculated the tissue specificity metric, SPM (Kryuchkova-Mostacci & Robinson-

Rechavi, 2017), which ranges from ubiquitous expression (near 0) to single-tissue specific (1). Although

the sperm proteomes of both of our species were enriched for gene products specifically expressed in

testes, they also contained broadly expressed gene products (Figure 3B). To assess the effect of these

broadly expressed genes on our inference of selection, we recalculated the α statistic for two bins of

genes (Figure 3C): those with broad expression (SPM < 0.5) and those with high tissue-specificity (SPM >

0.5).

In Carolina sphinx moths, there were no significant changes in α between low- and high-specificity genes

in any part of the genome (background genes: p = 0.3868, sperm proteome genes: p = 0.3248, male-

limited genes: p = 0.5579; Figure 3C, left), nor did any of the gene classes differ from each other within a

specificity bin. In monarchs, however, both sperm proteome genes (p = 0.0242) and testes genes (p =

0.0137) showed higher α in the high-specificity group than the low-specificity group, though somatically

expressed genes did not (p = 0.6831). Additionally, we found that sperm genes showed much greater α

than the background genome or other genes expressed in the testes at both low- and high-specificities

(Figure 3C, right). This result indicates that our initial results (considering the whole sperm proteome)

are not dependent on the underlying specificity of sperm genes.

*Molecular evolution in dimorphic sperm*

Having verified the patterns of evolution in the whole sperm proteomes with several approaches, we

turned to our primary question, assessing apyrene sperm function through analysis of molecular

evolution. We considered the different subsets of the sperm proteomes based on the two sperm types.

The two datasets consisted of three classes of sperm proteins: unique to eupyrene sperm, unique to

apyrene sperm, or found in both cell types (henceforth "shared", Figure 4A). We assessed differences in

selective pressures between the sperm morphs with another series of permutation tests, both

comparing parts of the sperm proteome to the background genome and comparing parts of the

proteome to each other.

As expected based on the whole-proteome results from Carolina sphinx moth, neither eupyrene-specific

(p = 0.55912), shared (p = 0.4647), nor apyrene-specific proteins (p = 0.96496) differed from the

background genome (Figure 4B). α did not vary between apyrene-specific and eupyrene-specific

proteins (p = 0.7271), between apyrene-specific and shared (p = 0.7176) or eupyrene-specific and

shared proteins (p = 0.9979). In monarchs, both eupyrene-specific proteins (p =0.00018) and shared

proteins (p = 0.01038) showed elevated α, but apyrene-specific proteins did not evolve differently from

the background genome (p = 0.55934). Neither apyrene nor eupyrene sperm differed significantly from

the shared set in monarchs (p = 0.6332 & p = 0.6234, respectively), but there was a trend towards

significantly increased α in eupyrene-specific proteins compared to apyrene-specific proteins (p =

0.0986).

As with the whole sperm proteome, we investigated which classes of variants contributed to our

observed differences in α (Figure 4C). Consistent with the results above, none of the variant classes

significantly differed from the genome background in the sphinx moth eupyrene-specific proteins (Pn: W

= 995190, p = 0.0857; Ps: W = 943550, p = 0.6782; Dn: W = 966630, p = 0.3183; Ds: W = 963410, p =

0.3596). Shared proteins also showed the same level of variation as the background across all variants

(Pn: W = 1470300, p = 0.3277; Ps: W = 1444400, p = 0.1369; Dn: W = 1540700, p = 0.6883; Ds: W =

1437100, p = 0.1030). And finally, apyrene-specific proteins were not significantly different either (Pn: W

= 548570, p = 0.4974; Ps: W = 491410, p = 0.2149; Dn: W = 560910, p = 0.2741; Ds: W = 495180, p =

0.2653). In summary, there was no evidence for stronger selection on either sperm morph in Carolina

sphinx moths.

For monarchs, we found that the elevated α in the eupyrene-specific and shared subsets was driven

primarily by a decrease in non-synonymous polymorphism compared to the background genome (W =

1291700, p = 0.0003 for eupyrene; W = 1486100, p = $1.167*10^{-8}$ for shared). Apyrene-specific proteins

did not show a reduction in non-synonymous polymorphism (W = 284620, p = 0.1684). Synonymous

polymorphism did not significantly differ from the background in any subset of the sperm proteome

(eupyrene: W = 1164200, p = 0.4492; shared: W = 1249900, p = 0.5570, apyrene: W = 270160, p =

0.4927). Nor did synonymous divergence (eupyrene: W = 1056000, p = 0.0928; shared: W = 1209000, p =

0.7665, apyrene: W = 279420, p = 0.2594). Intriguingly, non-synonymous divergence was elevated

compared to the background in eupyrene-specific proteins (W = 1021800, p = 0.0151), but not the

shared (W = 121800, p = 0.9185) or apyrene-specific portions of the proteome (W = 266580, p = 0.6042).

This suggests periodic sweeps of positively selected variants in fertilizing sperm proteins.

We did not examine orthology within dimorphic sperm owing to small gene counts giving reduced

statistical power. Nor could we could examine tissue specificity here because apyrene and eupyrene

sperm are produced at different developmental timepoints and we did not have suitable expression

data in both species. Nonetheless, the consistency of results in the whole proteome datasets gives us no

reason to expect that within-proteome results would be idiosyncratic to our methodology.

*Demographic estimates*

Finally, to contextualize our results with population dynamics, we estimated population size history

using site frequency from 4-fold degenerate sites in the two species' genomes (Figure S3). Both have

effective population sizes near 2,000,000, as expected of herbivorous invertebrates with high dispersal

potential, numerous host plants, and a large range over North America. We also recovered a population

size increase in monarch butterflies in the recent past, which has been previously reported with genomic

data (Zhan et al., 2014). We note that our inferred timing of this event differs from that of the previous

authors, who used mutation rate estimates from *Drosophila melanogaster*. Such input parameter

differences affect the estimated time of events, but not the trajectories.

**Discussion**

We investigated the molecular evolution of eupyrene (fertilizing) and apyrene (non-fertilizing) sperm,

the ubiquitous lepidopteran cell-type of unknown functional significance.  These sperm have long been

posited to interfere with competitors' sperm, in part because their quantity varies with levels of male-

male competition (Silberglied et al., 1984; Solensky & Oberhauser, 2009; Swallow & Wilkinson, 2002). In

contrast to these organismal observations, the results of our molecular analyses cast doubt on this

hypothesis. If apyrene sperm played an active role in sperm competition, we would expect evidence for

stronger selection in apyrene sperm compared to the background genome in monarchs. We found a

signal for elevated adaptive evolution ($\alpha$) in the sperm proteome compared to the background genome

in these polyandrous butterflies, but this signal did *not* include apyrene-sperm-specific proteins. Instead,

genes encoding apyrene sperm proteins evolve similarly to the background genome in both monarchs

and Carolina sphinx moths. This result is unlikely to have arisen from a lack of power in our

methodologies, as eupyrene-specific and shared sperm proteins showed patterns in line with

expectations for a role of sperm competition in molecular evolution in monarchs.

*Selection consistent with sperm competition, but only in fertilizing sperm*

The source of the apparently elevated $\alpha$ in the monarch sperm proteome came mainly from a dearth of

non-synonymous polymorphisms in sperm proteins compared to the background genome, indicating the

action of purifying selection to remove many variants before fixation in monarchs. Strong purifying

selection has been similarly observed in genes expressed in pollen, the main male-male competitors in

flowering plants (Arunkumar, Josephs, Williamson, & Wright, 2013). A similar pattern can also be

observed in passerine birds, in which species with higher rates of sperm competition show less

intraspecific and intra-male variation in sperm length compared to sperm of less polyandrous species

(Immler et al., 2008; Kleven et al., 2008).

Moreover, the elevated $\alpha$ in sperm homologs in monarchs suggests that genes that have had conserved

sperm function since the divergence of the two species some 100 million years ago (Heikkila, Kaila,

Mutanen, Pena, & Wahlberg, 2012) are under stronger purifying selection in the polyandrous species.

According to recent gene ontology analyses, such genes are enriched for core traits in sperm, such as

mitochondrial function, respiration, and flagellar structure. Similarly, proteins shared between the two

sperm types and those unique to eupyrene sperm show an elevated $\alpha$ compared to the background

genome in monarchs. Sperm proteins shared between morphs are enriched for structural proteins that

give rise to the sperm tail and thus impact motility (Whittington et al., in press), while those expressed

only in eupyrene sperm doubtless include important mediators of fertilization. At the cellular level,

variation in sperm traits like swimming ability, longevity, and overall viability affects sperm competition

outcomes (Burness, Casselman, Schulte-Hostedde, Moyes, & Montgomerie, 2004; Kim et al., 2017) and

has a polygenic basis in other taxa (Hering, Olenski, & Kaminski, 2014). For traits like longevity and

motility there is a threshold below which fertilization becomes significantly impaired, but in the absence

of competitor alleles, there is a larger range of effectively-neutral trait-values, allowing for more

variation to be maintained in the population. In the presence of competitor alleles, however, marginal

differences in fertilization success come under selection, leading to the removal of deleterious variants

through sperm competition.

Stronger selection from competition may include even the event of fertilization itself. Lepidopteran eggs

are known to possess multiple micropyle openings for sperm (Kumar, Kariappa, Babu, & Dandin, 2007)

and eupyrene sperm possess structures resembling an acrosome (while their apyrene counterparts do

not) (Friedlander, 1997). This rare combination of male and female gamete structures is also found in

sturgeon, in which the multiple micropyles give several sperm potential access to the egg nucleus and

there is competition among sperm to initiate karyogamy via the acrosome reaction (Psenicka, Rodina, &

Linhart, 2010). Consistent with micropyle-mediated competition, it has been shown that more

polyandrous species of Lepidoptera tend to have more micropyles on their egg surfaces than

monandrous species (Iossa, Gage, & Eady, 2016). If this truly does extend the opportunity for male-male

competition and cryptic choice, then acrosomal proteins in eupyrene sperm would be likely targets for

selection in polyandrous systems.

Whatever the mechanics of fertilization are, paternity outcomes in polyandrous species are often

bimodally distributed (Simmons & Siva-Jothy, 1998; Wedell & Cook, 1998), including in monarch

butterflies (Mongue, Ahmed, Tsai, & De Roode, 2015). For females that mate twice, one of the two

males typically fathers most, if not all, of the observed offspring produced by the female, but there is

little consistency in whether it is the first or second male. With these dynamics, fitness differences

between winning and losing sperm phenotypes are large and selection can reliably remove less

successful genotypes.

Evidence of this can be seen in the estimated distribution of fitness effects of new mutations in monarch

sperm proteins. Compared to the background genome, we see a decrease in the proportion of

effectively neutral and weakly deleterious mutations and an increase in both strongly deleterious and

beneficial mutations. In the absence of competition, not only are mildly suboptimal variants effectively

neutral, but novel, more efficient competitors should have no selective advantage in monandrous

species unless they also markedly increase fitness in a single mating. This reasoning is supported by the

estimated distribution of fitness effect for the complimentary gene sets in the Carolina sphinx moth; in

this species, we see little variation in the DFE between the background genome and the sperm

proteome. Moreover, there is no decrease (and indeed) an increase in non-synonymous divergence of

eupyrene sperm proteins in monarchs compared to the rest of the genome. This pattern suggests that in

addition to strong purifying selection there must be periodic sweeps of beneficial alleles. Without a

broader, phylogenetically controlled study, these results between a single pair of species are not

conclusive, but they fit well with the prediction that sperm protein evolution depends on the rates of

polyandry in a species (Dapper & Wade, 2016).

*Evolution of tissue-specific and male-limited genes*

Other studies have demonstrated that tissue specificity of expression can strongly influence the

molecular evolution of reproductive proteins (Schumacher & Herlyn, 2018), in some cases more than

mating system (Carnahan-Craig & Jensen-Seaman, 2014). Because our proteomic data did not contain

information on tissue specificity on their own, we examined this dynamic with RNA-seq data. We found

increased adaptive evolution in monarch sperm genes with higher specificity compared to low-

specificity sperm genes. Furthermore, while not significantly different from background genome, $\alpha$ for

non-sperm genes expressed in the testes increased with greater specificity in monarchs, suggesting that

they too may be subject to stronger sexual selection in this polyandrous species. Neither of these

patterns held for Carolina sphinx moths, which showed no differences based on tissue specificity. This

consistency further suggests the difference in mating system as an explanation for differences in

intensity of selection.

Finally, we did not observe relaxed constraint in reproductive proteins predicted due to the smaller

effective population size of males or females compared to the population as a whole, as predicted by

theory (Dapper & Wade, 2016; Wade et al., 2008). Specifically, we did not observe a difference in the

adaptive evolution of genes with testes-specific expression, our proxy for sex-limited expression,

compared to the background genome. To explain this discrepancy between theory and observation, we

turn to Nearly Neutral Theory. Large populations have more efficient selection than small populations

and a smaller range of slightly deleterious mutations that behave neutrally (Ohta, 1992). Mutations with

a selective effect less than $1/N_e$ are expected to behave neutrally. For instance, one commonly cited

estimate for human population size is Ne ≈ 10,000 over evolutionary history (Zhao et al., 2000). Based

on this, mutations with selective effects less than 0.0001 should behave neutrally for alleles expressed in

both sexes, while those with effects of 0.0002 are effectively neutral for alleles only expressed in one

sex. And indeed, there is evidence that genes expressed only in men have a higher mutational load than

those expressed in both sexes (Gershoni & Pietrokovski, 2014). Chimpanzees, another species with a

similar effective population size (Won & Hey, 2005), also show increased non-synonymous divergence in

reproductive proteins (Wong, 2010). Broadly, male reproductive protein evolution appears to depend

more on effective population sizes than intensity of sperm competition in the great apes in general

(Good et al., 2013), as one would expect for species with relatively small effective population sizes.

In contrast to mammals, the effective population sizes of most insect species are orders of magnitude

higher. Using neutral site frequency spectra, we estimated effective populations near 2,000,000 for both

North American monarchs and Carolina sphinx moths. Selection is much more effective in these massive

populations; mutations with effects above $5*10^{-7}$ should be subject to selection in both sexes and those

above $1*10^{-6}$ should be subject to selection if expression is sex-limited. Thus, even selection on alleles

with sex-limited expression in these insects should be 100 times stronger than selection on the entire

human population. Even if there is a relative two-fold difference in selection, the absolute magnitude of

the difference should be miniscule, and the effects of mating system more apparent.

*Advancing understanding of apyrene sperm*

Previous morphological work found that eupyrene sperm traits (like sperm length) but not apyrene

sperm traits, varied with risk of sperm competition in other butterflies (Gage, 1994). Similarly, from a

molecular perspective, none of the patterns of increased purifying and positive selection that we

observed for monarch sperm proteins applied to the apyrene-specific proteins. That we also do not see

evidence for the action of sperm competition on apyrene-specific protein evolution is itself informative,

however. Research to-date has proposed four main hypotheses for apyrene sperm (Swallow &

Wilkinson, 2002): active sperm competition agents, passive competition agents, nutrient nuptial gifts, or

necessary facilitators of fertilization. Our molecular analyses argue against apyrene sperm as active

agents of sperm competition, but it is worth considering predictions for molecular evolution of apyrene

sperm under the other hypotheses.

Indeed, apyrene sperm may still have adaptive significance without specialized molecular function,

especially under the filler hypothesis. This proposed function also relates to sperm competition, but

posits that apyrene sperm are employed proactively, to fill the female's sperm storage organ and delay

remating, thus decreasing the risk of sperm competition, rather than impacting its outcome (Swallow &

Wilkinson, 2002). Both in monarchs and the butterfly *Pieris napi*, female time to remating increases with

the number of apyrene sperm received from males (Cook & Wedell, 1999; Oberhauser, 1988). Such

observations are somewhat confounded by the size of the spermatophore nuptial gift that males

provide during mating, but apyrene sperm themselves have been proposed as a form of nutritional

nuptial gift (He, Tanaka, & Miyata, 1995; Lamunyon, 2000). Under both the nutrient and filler

hypotheses, the actual sequence of apyrene sperm proteins should be less important than their physical

presence and abundance, so factors affecting the rate of apyrene sperm production would be more

likely targets for selection in polyandrous species than the proteins sequences themselves.

Finally, apyrene sperm appear to capacitate fertilization in *Bombyx mori* (Takemura, Sahara, Mochida, &

Ohnuma, 2006); the mechanism here is unclear and the phenomenon is untested in other taxa, but it

could conceivably involve proteins that modulate female reproductive physiology to make conditions

more favorable for eupyrene sperm or induce oviposition. In such a case, these proteins would behave

more akin to the broader class of reproductive proteins and evolve independently of rates of polyandry

in a species.  If there is an evolutionarily conserved capacitation effector in our study taxa, it is possible

that this function is governed by a small subset of apyrene-specific proteins. Because our methods

aggregate signal for selection across multiple genes or sites to counteract high variance in variant counts

within genes (Stoletzki & Eyre-Walker, 2011), the importance of one or a few genes could be lost in the

heterogeneous selection on different proteins.

Conclusions

Variation in reproductive traits has long been studied at the morphological and molecular level,

generally. Yet sperm dimorphism, one of the most striking and enigmatic reproductive traits, has not

previously been assessed using population genetic analyses. Our investigation of the sperm proteome in

two Lepidoptera demonstrates a pattern of stronger purifying selection on fertilizing-sperm genes in a

species with higher rates of sperm competition. In this polyandrous species, these genes experience a

strikingly different selective environment than the rest of the genome, with strong purifying selection

reducing variation in sperm genes. In contrast, fertilizing-sperm genes in the monandrous species hold

as much deleterious variation as other parts of their genome. Our new molecular findings fit well with

established studies on sperm morphology which show that sperm competition results in decreased

variation in sperm traits.

The evolution of non-fertilizing sperm, however, does not show a strong influence of sperm competition.

This lack of pattern itself argues against apyrene sperm as active agents of sperm competition, one of

the long-held hypotheses for non-fertilizing sperm function. Instead, apyrene sperm may play a passive

role in reducing the risk of competition by delaying female remating. The method by which apyrene

sperm capacitate fertilization in some species remains unclear based solely on genomic approaches and

will likely require functional experiments to completely understand.

**Acknowledgments**

This project was funded by the NSF DDIG (DEB-1701931) and Kansas Idea Network of Biomedical
research (NIH P20 GM103418). The authors wish to acknowledge Wesley Mason and Michael Hulet and
the rest of the Information and Telecommunication Technology Center (ITTC) staff at the University of
Kansas for their support with our high-performance computing. Thank you to Jacobus de Roode for use
of the monarch image, Elizabeth Moore for facilitating collaboration between Kansas and North Carolina,
Tawny Scanlan for comments on sperm biology, and Amanda Pierce and Tom de Man for housing during
field collection.

**References**

Arunkumar, R., Josephs, E. B., Williamson, R. J., & Wright, S. I. (2013). Pollen-specific, but not sperm-

specific, genes show stronger purifying selection and higher rates of positive selection than

sporophytic genes in *Capsella grandiflora*. *Molecular Biology and Evolution*, *30*(11), 2475–2486.

doi:10.1093/molbev/mst149

Baker, R. R., & Bellis, M. A. (1989). Elaboration of the kamikaze sperm hypothesis: a reply to Harcourt.

*Animal Behaviour*, *37*(5), 865–867.

Barker, M. S., Demuth, J. P., & Wade, M. J. (2005). Maternal expression relaxes constraint on innovation

of the anterior determinant, bicoid. *PLoS Genetics*, *1*(5), e57.

Birkhead, T. R. (1998). Cryptic female choice: criteria for establishing female sperm choice. *Evolution*,

1212–1218.

Bressac, C., Joly, D., Devaux, J., Serres, C., Feneux, D., & Lachaise, D. (1991). Comparative kinetics of

short and long sperm in sperm dimorphic *Drosophila* species. *Cell Motility and the Cytoskeleton*,

*19*(4), 269–274. doi:10.1002/cm.970190405

Buckland-Nicks, J, Bryson, I., Hart, L., & Partridge, V. (2010). Sex and a snail's sperm: on the transport,

storage and fate of dimorphic sperm in Littorinidae. *Invertebrate Reproduction and*

*Development*, *36*, 145–152.

Buckland-Nicks, John. (1998). Prosobranch parasperm: Sterile germ cells that promote paternity? In

*Micron* (Vol. 29, pp. 267–280). doi:10.1016/S0968-4328(97)00064-4

Burness, G., Casselman, S. J., Schulte-Hostedde, A. I., Moyes, C. D., & Montgomerie, R. (2004). Sperm

swimming speed and energetics vary with sperm competition risk in bluegill (*Lepomis*

*macrochirus*). *Behavioral Ecology and Sociobiology*, *56*(1), 65–70. doi:10.1007/s00265-003-0752-

7

Cao, X., & Jiang, H. (2017). An analysis of 67 RNA-seq datasets from various tissues at different stages of

a model insect, *Manduca sexta*. *BMC Genomics*, *18*(1). doi:10.1186/s12864-017-4147-y

Carcupino, M., Baldacci, A., Fausto, A. M., Scapigliati, G., & Mazzini, M. (1999). Sperm dimorphism in

Chilopoda: comparison of Scolopendromorpha and Geophilomorpha. *Invertebrate Reproduction*

*& Development*, *35*(1), 45–53.

Carnahan-Craig, S., & Jensen-Seaman, M. (2014). Rates of evolution of hominoid seminal proteins are

correlated with function and expression, rather than mating system. *Journal of Molecular*

*Evolution*, *78*(1), 87–99.

Chenoweth, P. J. (2005). Genetic sperm defects. In *Theriogenology* (Vol. 64, pp. 457–468).

doi:10.1016/j.theriogenology.2005.05.005

Cingolani, P., Platts, A., Wang, L. L., Coon, M., Nguyen, T., Wang, L., … Ruden, D. M. (2012). A program

for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in

the genome of *Drosophila melanogaster* strain w 1118; iso-2; iso-3. *Fly*, *6*(2), 80–92.

doi:10.4161/fly.19695

Civetta, A., & Singh, R. S. (1995). High divergence of reproductive tract proteins and their association

with postzygotic reproductive isolation in *Drosophila melanogaster* and *Drosophila virilis* group

species. *Journal of Molecular Evolution*, *41*(6), 1085–1095. doi:10.1007/BF00173190

Cook, P. A., & Wedell, N. (1999). Non-fertile sperm delay female remating. *Nature*, *397*(6719), 486.

doi:10.1038/17257

Dapper, A. L., & Wade, M. J. (2016a). The evolution of sperm competition genes: The effect of mating

system on levels of genetic variation within and between species. *Evolution*, *70*(2), 502–511.

doi:10.1111/evo.12848

Dorus, S., Evans, P. D., Wyckoff, G. J., Sun, S. C., & Lahn, B. T. (2004). Rate of molecular evolution of the

seminal protein gene SEMG2 correlates with levels of female promiscuity. *Nature Genetics*,

*36*(12), 1326–1329. doi:10.1038/ng1471

Eckelbarger, K. J., Young, C. M., & Cameron, J. L. (1989). Ultrastructure and development of dimorphic

583   sperm in the abyssal echinoid *Phrissocystis multispina* (Echinodermata: Echinoidea): implications

584   for deep sea reproductive biology. *The Biological Bulletin*, *176*(3), 257–271.

585 Eyre-Walker, A., & Keightley, P. D. (2009). Estimating the rate of adaptive molecular evolution in the

586   presence of slightly deleterious mutations and population size change. *Molecular Biology and*

587   *Evolution*, *26*(9), 2097–2108. doi:10.1093/molbev/msp119

588 Friedlander, M. (1997). Control of the eupyrene-apyrene sperm dimorphism in Lepidoptera. *J Insect*

589   *Physiol*, *43*(12), 1085–1092. doi:10.1016/s0022-1910(97)00044-9

590 Gage, M. J. (1994). Associations between body size, mating pattern, testis size and sperm lengths across

591   butterflies. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *258*(1353),

592   247–254.

593 Gershoni, M., & Pietrokovski, S. (2014). Reduced selection and accumulation of deleterious mutations in

594   genes exclusively expressed in men. *Nature Communications*, *5*. doi:10.1038/ncomms5438

595 Good, J. M., Wiebe, V., Albert, F. W., Burbano, H. A., Kircher, M., Green, R. E., … Pääbo, S. (2013).

596   Comparative population genomics of the ejaculate in humans and the great apes. *Molecular*

597   *Biology and Evolution*, *30*(4), 964–976. doi:10.1093/molbev/mst005

598 Harcourt, A. (1991). Sperm competition and the evolution of nonfertilizing sperm in mammals. *Evolution*,

599   *45*(2), 314–328.

600 Harcourt, A. H. (1989). Deformed sperm are probably not adaptive. *Animal Behaviour*, *37*(5), 863–864.

601 He, Y. B., Tanaka, T., & Miyata, T. (1995). Eupyrene and apyrene sperm and their numerical fluctuations

602   inside the female reproductive tract of the armyworm, *Pseudaletia separata*. *Journal of Insect*

603   *Physiology*, *41*(8), 689–694. doi:10.1016/0022-1910(95)00020-U

Heikkila, M., Kaila, L., Mutanen, M., Pena, C., & Wahlberg, N. (2012). Cretaceous origin and repeated

tertiary diversification of the redefined butterflies. *Proceedings of the Royal Society B: Biological*

*Sciences*, *279*(1731), 1093–1099. doi:10.1098/rspb.2011.1430

Herberg, S., Gert, K. R., Schleiffer, A., & Pauli, A. (2018). The Ly6/uPAR protein Bouncer is necessary and

sufficient for species-specific fertilization. *Science*, *361*(6406), 1029–1033.

Hering, D. M., Olenski, K., & Kaminski, S. (2014). Genome-wide association study for poor sperm motility

in Holstein-Friesian bulls. *Animal Reproduction Science*, *146*(3–4), 89–97.

doi:10.1016/j.anireprosci.2014.01.012

Higginson, D. M., Miller, K. B., Segraves, K. A., & Pitnick, S. (2012). Convergence, recurrence and

diversification of complex sperm traits in diving beetles (Dytiscidae). *Evolution*, *66*(5), 1650–

1661. doi:10.1111/j.1558-5646.2011.01532.x

Hill Jr., H. F., Wenner, A. M., & Wells, P. H. (1976). Reproductive Behavior in an Overwintering

Aggregation of Monarch Butterflies. *American Midland Naturalist*, *95*(1), 10–19.

doi:10.2307/2424229

Immler, S., Calhim, S., & Birkhead, T. R. (2008). Increased postcopulatory sexual selection reduces the

intramale variation in sperm design. *Evolution*, *62*(6), 1538–1543.

Iossa, G., Gage, M. J., & Eady, P. E. (2016). Micropyle number is associated with elevated female

promiscuity in Lepidoptera. *Biology Letters*, *12*(12), 20160782.

Kanost, M. R., Arrese, E. L., Cao, X., Chen, Y.-R., Chellapilla, S., Goldsmith, M., … Jiang, H. (2016).

Multifaceted biological insights from a draft genome sequence of the tobacco hornworm moth,

*Manduca sexta*. *Insect Biochemistry and Molecular Biology*, *76*, 118–147.

Keightley, P. D., Pinharanda, A., Ness, R. W., Simpson, F., Dasmahapatra, K. K., Mallet, J., … Jiggins, C. D.

(2015). Estimation of the Spontaneous Mutation Rate in *Heliconius melpomene*. *Molecular*

*Biology and Evolution*, *32*(1), 239–243. doi:10.1093/molbev/msu302

Kim, K. W., Bennison, C., Hemmings, N., Brookes, L., Hurley, L. L., Griffith, S. C., ... Slate, J. (2017). A sex-

linked supergene controls sperm morphology and swimming speed in a songbird. *Nature*

*Ecology and Evolution*, *1*(8), 1168–1176. doi:10.1038/s41559-017-0235-2

Kleven, O., Laskemoen, T., Fossøy, F., Robertson, R. J., & Lifjeld, J. T. (2008). Intraspecific variation in

sperm length is negatively related to sperm competition in passerine birds. *Evolution*, *62*(2),

494–499. doi:10.1111/j.1558-5646.2007.00287.x

Korneliussen, T. S., Albrechtsen, A., & Nielsen, R. (2014). ANGSD: Analysis of Next Generation

Sequencing Data. *BMC Bioinformatics*, *15*(1). doi:10.1186/s12859-014-0356-4

Kryuchkova-Mostacci, N., & Robinson-Rechavi, M. (2017). A benchmark of gene expression tissue-

specificity metrics. *Briefings in Bioinformatics*, *18*(2), 205–214. doi:10.1093/bib/bbw008

Kumar, V., Kariappa, B. K., Babu, A. M., & Dandin, S. B. (2007). Surface ultrastructure of the egg chorion

of eri silkworm, *Samia ricini* (Donovan) (Lepidoptera: Saturniidae). *Journal of Entomology*, *4*(2),

68–81. doi:10.3923/je.2007.68.81

Lamunyon, C. (2000). Sperm storage by females of the polyandrous noctuid moth *Heliothis virescens*.

*Animal Behaviour*, *59*(2), 395–402. doi:10.1006/anbe.1999.1294

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat Methods*, *9*(4),

357–359. doi:10.1038/nmeth.1923

Liu, X., & Fu, Y.-X. (2015). Exploring population size changes using SNP frequency spectra. *Nature*

*Genetics*, *47*(5), 555–559. doi:10.1038/ng.3254

Lunter, G., & Goodson, M. (2011). Stampy: A statistical algorithm for sensitive and fast mapping of

Illumina sequence reads. *Genome Research*, *21*(6), 936–939. doi:10.1101/gr.111120.110

Marks, J. A., Biermann, C. H., Eanes, W. F., & Kryvi, H. (2008). Sperm polymorphism within the sea urchin

*Strongylocentrotus droebachiensis*: Divergence between pacific and atlantic oceans. *Biological*

*Bulletin*, *215*(2), 115–125.

McKenna, A. H., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., … Depristo, M. (2010).

The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA

sequencing data. *Genome Research*, *20*(9), 1297–1303. doi:10.1101/gr.107524.110

Meves, F. (1902). Über oligopyrene und apyrene Spermien und über ihre Entstehung, nach

Beobachtungen an Paludina und Pygaera. *Archiv Für Mikroskopische Anatomie*, *61*(1), 1–84.

Mongue, A. J., Ahmed, M. Z., Tsai, M. V., & De Roode, J. C. (2015). Testing for cryptic female choice in

monarch butterflies. *Behavioral Ecology*, *26*(2), 386–395. doi:10.1093/beheco/aru196

Mongue, A. J., & Walters, J. (2017). The Z chromosome is enriched for sperm proteins in two divergent

species of Lepidoptera. *Genome*, *61*(4), 248–253.

Moore, H. D. M., Martin, M., & Birkhead, T. R. (1999). No evidence for killer sperm or other selective

interactions between human spermatozoa in ejaculates of different males in vitro. *Proceedings*

*of the Royal Society of London. Series B: Biological Sciences*, *266*(1436), 2343–2350.

Oberhauser, K. S. (1988). Male monarch butterfly spermatophore mass and mating strategies. *Animal*

*Behaviour*, *36*, 1384–1388. doi:10.1016/s0003-3472(88)80208-2

Ohta, T. (1992). The Nearly Neutral Theory of Molecular Evolution. *Annual Review of Ecology and*

*Systematics*, *23*(1), 263–286. doi:10.1146/annurev.es.23.110192.001403

Pitnick, S., Hosken, D. J., & Birkhead, T. R. (2009). Sperm morphological diversity. In *Sperm Biology* (pp.

69–149). doi:10.1016/B978-0-12-372568-4.00003-3

Pizzari, T. (2006). Evolution: The Paradox of Sperm Leviathans. *Current Biology*.

doi:10.1016/j.cub.2006.05.031

Preston, B. T., Saint Jalme, M., Hingrat, Y., Lacroix, F., & Sorci, G. (2015). The sperm of aging male

bustards retards their offspring's development. *Nature Communications*, *6*, 6146.

doi:10.1038/ncomms7146

Psenicka, M., Rodina, M., & Linhart, O. (2010). Ultrastructural study on the fertilisation process in

sturgeon (*Acipenser*), function of acrosome and prevention of polyspermy. *Animal Reproduction*

*Science*, *117*(1–2), 147–154. doi:10.1016/j.anireprosci.2009.03.013

R Core Team. (2017). R Core Team (2017). R: A language and environment for statistical computing. *R*

*Foundation for Statistical Computing, Vienna, Austria. URL Http://Www.R-Project.Org/.*, R

Foundation for Statistical Computing.

Sasakawa, K. (2009). Marked sperm dimorphism in the ground beetle *Scarites terricola*: A novel type of

insect sperm polymorphism. *Physiological Entomology*, *34*(4), 387–390. doi:10.1111/j.1365-

3032.2009.00694.x

Schumacher, J., & Herlyn, H. (2018). Correlates of evolutionary rates in the murine sperm proteome.

*BMC Evolutionary Biology*, *18*(1), 35.

Schumacher, J., Rosenkranz, D., & Herlyn, H. (2014). Mating systems and protein–protein interactions

determine evolutionary rates of primate sperm proteins. *Proceedings of the Royal Society of*

*London B: Biological Sciences*, *281*(1775), 20132607.

Schumacher, J., Zischler, H., & Herlyn, H. (2017). Effects of different kinds of essentiality on sequence

evolution of human testis proteins. *Scientific Reports*, *7*, 43534.

Silberglied, R. E., Shepherd, J. G., & Dickinson, J. L. (1984). Eunuchs - the role of apyrene sperm in

Lepidoptera. *American Naturalist*, *123*(2), 255–265. doi:10.1086/284200

Simmons, L. W., & Siva-Jothy, M. T. (1998). Sperm competition in insects: mechanisms and the potential

for selection. *Sperm Competition and Sexual Selection*, 341–434.

Smith, R. L. (1984). *Sperm competition and the evolution of animal mating systems*. Elsevier.

Snow, J. W., Copeland, W. W., Goodenough, J. L., Baumhover, A. H., Lingren, P. D., & Haile, D. G. (1974).

tobacco hornworm: notes on morphoology and mating habits. *Journal of the Georgia*

*Entomological Society*.

Solensky, M. J., & Oberhauser, K. S. (2009). Male monarch butterflies, *Danaus plexippus*, adjust

ejaculates in response to intensity of sperm competition. *Animal Behaviour*, *77*(2), 465–472.

doi:10.1016/j.anbehav.2008.10.026

Stoletzki, N., & Eyre-Walker, A. (2011). Estimation of the neutrality index. *Molecular Biology and*

*Evolution*, *28*(1), 63–70. doi:10.1093/molbev/msq249

Swallow, J. G., & Wilkinson, G. S. (2002). The long and short of sperm polymorphisms in insects.

*Biological Reviews of the Cambridge Philosophical Society*, *77*, 153–182.

doi:10.1017/S146479310100585

Swanson, W J, & Vacquier, V. D. (1998). Concerted evolution in an egg receptor for a rapidly evolving

abalone sperm protein. *Science*, *281*(5377), 710–712. doi:10.1126/science.281.5377.710

Swanson, Willie J., & Vacquier, V. D. (2002). Reproductive Protein Evolution. *Annual Review of Ecology*

*and Systematics*, *33*(1), 161–179. doi:10.1146/annurev.ecolsys.33.010802.150439

Takemura, Y., Sahara, K., Mochida, Y., & Ohnuma, A. (2006). Apyrene sperm from the triploid donors

restore fecundity of cryopreserved semen in *Bombyx mori*. *Journal of Insect Physiology*, *52*(10),

1021–1026. doi:10.1016/j.jinsphys.2006.06.010

Tataru, P., Mollion, M., Glémin, S., & Bataillon, T. (2017). Inference of distribution of fitness effects and

proportion of adaptive substitutions from polymorphism data. *Genetics*, *207*(3), 1103–1119.

doi:10.1534/genetics.117.300323

Tavares-Bastos, L., Teixeira, R. D., Colli, G. R., & Báo, S. N. (2002). Polymorphism in the sperm

ultrastructure among four species of lizards in the genus *Tupinambis* (Squamata: Teiidae). *Acta*

*Zoologica*, *83*(4), 297–307. doi:10.1046/j.1463-6395.2002.00119.x

Urquhart, F. A. (1976). Found at Last - Monarchs Winter Home. *National Geographic*, *150*(2), 161–173.

Wade, M., Priest, N., & Cruickshank, T. (2008). A theoretical overview of maternal genetic effects:

evolutionary predictions and empirical tests using sequence data within and across mammalian

taxa. *Maternal Effects in Mammals, University of Chicago Press, Chicago*.

Wedell, N., & Cook, P. A. (1998). Determinants of paternity in a butterfly. *Proceedings of the Royal

Society of London. Series B: Biological Sciences*, *265*(1396), 625–630.

doi:10.1098/rspb.1998.0340

Whittington, E., Forsythe, D., Borziak, K., Karr, T. L., Walters, J. R., & Dorus, S. (2017). Contrasting

patterns of evolutionary constraint and novelty revealed by comparative sperm proteomic

analysis in Lepidoptera. *BMC Genomics*, *18*(1). doi:10.1186/s12864-017-4293-2

Whittington, E., Karr, T. L., Mongue, A. J., Walters, J. R., & Dorus, S. (in press). Evolutionary proteomics

reveals distinct patterns of complexity and divergence between Lepidopteran sperm morphs.

*Genome Biology and Evolution*.

Whittington, E., Zhao, Q., Borziak, K., Walters, J. R., & Dorus, S. (2015). Characterisation of the *Manduca

sexta* sperm proteome: Genetic novelty underlying sperm composition in Lepidoptera. *Insect

Biochemistry and Molecular Biology*, *62*, 183–193. doi:10.1016/j.ibmb.2015.02.011

Wilms, H. J. (1986). Dimorphic sperm cells in the pollen grain of *Spinacia*.

Won, Y. J., & Hey, J. (2005). Divergence population genetics of chimpanzees. *Molecular Biology and

Evolution*, *22*(2), 297–307. doi:10.1093/molbev/msi017

Wong, A. (2010). Testing the effects of mating system variation on rates of molecular evolution in

primates. *Evolution*, *64*(9), 2779–2785. doi:10.1111/j.1558-5646.2010.01038.x

Zhan, S., & Reppert, S. M. (2013). MonarchBase: The monarch butterfly genome database. *Nucleic Acids

Research*, *41*(D1), D758–D763. doi:10.1093/nar/gks1057

Zhan, S., Zhang, W., Niitepõld, K., Hsu, J., Haeger, J. F., Zalucki, M. P., ... Kronforst, M. R. (2014). The

genetics of monarch butterfly migration and warning colouration. *Nature*, *514*(7522), 317–321.

Zhao, Z., Jin, L., Fu, Y. X., Ramsay, M., Jenkins, T., Leskinen, E., … Li, W. H. (2000). Worldwide DNA

sequence variation in a 10-kilobase noncoding region on human chromosome 22. *Proceedings of*

*the National Academy of Sciences of the United States of America*, *97*(21), 11354–8.

doi:10.1073/pnas.200348197

**Author Contributions:** AJM designed the experiments, collected samples, performed analyses, and

wrote the manuscript. MEH collected samples and conducted analyses. LG provided data and performed

analyses. CES planned and facilitated sample collection and edited the manuscript. JRW assisted in

experiment design and manuscript editing.

**Data accessibility:** *Manduca sexta* whole genome resequencing data can be found on NCBI's Sequence

Read Archive with the following accession: SRP144217. *Danaus plexippus* RNA sequencing data

can be retrieved with accessions: SRR8580831 - SRR8580842. Analysis scripts can be found at

https://github.com/WaltersLab/DimorphicSpermMolEvo.

Figure 1. A. In the Carolina sphinx moth (*M. sexta*), there is no difference between the sperm proteome and the rest of the genome (left); conversely, genes in the sperm proteome of monarch butterflies (*D. plexippus*) show a significantly higher proportion of adaptive substitutions (α) than the rest of the genome (right). P-values come from permutation tests. Error bars represent 95% bootstrapped confidence intervals from the point estimates. B. Decomposing α into its components: Pn, Ps, Pn, and Ds and comparing the sperm proteome (filled boxes) to the background genome (open boxes). There were no strong differences between sperm genes and the background genome in Carolina sphinx moths. In monarch butterflies, the signal for increased adaptive substitution comes from a marginal increase in non-synonymous divergence (bottom left) combined with a great reduction in non-synonymous polymorphism in sperm genes compared to the rest of the genome (top left). P-values reflect Wilcoxon-Mann-Whitney tests, with * < 0.05, ** < 0.005, *** < 0.0005, etc.

# Sperm Proteome Overlap

**A**

*Manduca sexta*

The Carolina sphinx moth

401  236│216  294

*Danaus plexippus*

The monarch butterfly

**B**  Adaptive evolution accounting for orthology in the sperm proteins

**Figure 2. A.** Composition of the portion of the sperm proteomes analyzed in this study. Numbers indicate counts of proteins unique to one species' sperm or with an ortholog in the other species' sperm (sperm homologs). Note that the overlap number varies between species due to the presence of a few one-to-many-orthologs. **B.** Sperm homologs show evidence for a greater proportion of adaptive substitutions ($\alpha$) in monarch butterflies, but not in Carolina sphinx moths. P-values are based on permutation tests comparing the difference between two sets of genes randomly assigned from the sperm proteome in each species; error bars are 95% bootstrap confidence intervals.

**Figure 3.** Investigating how tissue specificity of gene expression impacts adaptive evolution in both Carolina sphinx moths (**left column**) and monarchs (**right column**). **A.** Maximum specificity of all genes across all studied tissues using the RNA-seq data considered in these analyses. **B.** Observed distribution of specificity of sperm proteome genes expressed in the testes. Based on these distributions, we separated genes into one of two categories, low-specificity (SPM < 0.5) or high-specificity (SPM > 0.5), divided by the dashed line. **C.** Inferred proportion of adaptive substitutions ($\alpha$) in background genes (squares), sperm proteome genes (circles), and male-limited genes (as defined by testes expression). Bars represent 95% confidence intervals from non-parametric bootstrapping. Non-overlapping confidence intervals imply significant differences generally, but we have also highlighted two significant differences that are less visibly apparent. Monarchs show evidence for increasing $\alpha$ with increasing tissue specificity in sperm and testes genes, but sphinx moths do not. Moreover, sperm proteome genes evolve more adaptively than background or testes-specific genes in both specificity groups for monarchs but not sphinx moths.

Figure 4. A. Composition of the sperm proteome with respect to dimorphic sperm. The majority of identified proteins were shared between the two cell types, followed by the set unique to eupyrene sperm, and finally the smallest set was the proteins found only in apyrene sperm. B. None of the sets of sperm proteins evolved either differently from each other or distinctly from the background genome (shaded regions represent 95% confidence intervals of the background) in the Carolina sphinx (left). In the monarch however (right), the signal for elevated α was localized to the eupyrene-specific and shared proteins. There was also a trend for increased α in eupyrene-specific proteins as compared to apyrene-specific. Error bars represent 95% confidence intervals from bootstrapping. C. Decomposing α into Pn, Ps, Dn, and Ds for dimorphic sperm. Plotting of variation follows the coloring and order in parts A and B; from left to right in each panel: background genome, eupyrene, shared, and apyrene sperm. Asterisks denote significant differences from the background genome based on a Wilcoxon-Mann-Whitney test, with * < 0.05, ** < 0.005, *** < 0.0005, etc.