Title: The capsule regulatory network of *Klebsiella pneumoniae* defined by density-

TraDISort

Running title: Density-TraDISort for *K. pneumoniae* capsule.

Matthew J. Dorman[a], Theresa Feltwell[a], David A. Goulding[a], Julian Parkhill[a], & Francesca L.

Short[a,b#]

a.   Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridgeshire,

10   CB10 1SA, United Kingdom.

11   b.   Department of Medicine, University of Cambridge, Cambridge CB2 0QQ, United

12   Kingdom.

[#] Correspondence to F.L.S.: fs13@sanger.ac.uk

Word count: 5,312

Abstract/importance word count: 239/120

Two-part abstract

Abstract: *Klebsiella pneumoniae* infections affect infants and the immunocompromised, and

the recent emergence of hypervirulent and multi-drug resistant *K. pneumoniae* lineages is a

critical healthcare concern. Hypervirulence in *K. pneumoniae* is mediated by several factors,

including the overproduction of extracellular capsule. However, the full details of how *K.*

pneumoniae capsule biosynthesis is achieved or regulated are not known. We have developed

a robust and sensitive procedure to identify genes influencing capsule production, density-

TraDISort, which combines density gradient centrifugation with transposon-insertion

sequencing. We have used this method to explore capsule regulation in two clinically-

relevant *Klebsiella* strains; *K. pneumoniae* NTUH-K2044 (capsule type K1), and *K.*

pneumoniae ATCC43816 (capsule type K2). We identified multiple genes required for full

capsule production in *K. pneumoniae*, as well as putative suppressors of capsule in NTUH-

K2044, and have validated the results of our screen with targeted knockout mutants. Further

investigation of several of the *K. pneumoniae* capsule regulators identified – ArgR,

MprA/KvrB, SlyA/KvrA and the Sap ABC transporter – revealed effects on capsule amount

and architecture, serum resistance and virulence. We show that capsule production in *K.*

pneumoniae is at the centre of a complex regulatory network involving multiple global

regulators and environmental cues, and that the majority of capsule regulatory genes are

located in the core genome. Overall our findings expand our understanding of how capsule is

regulated in this medically-important pathogen, and provide a technology that can be easily

implemented to study capsule regulation in other bacterial species.

Importance: Capsule production is essential for *K. pneumoniae* to cause infections, but its

regulation and mechanism of synthesis are not fully understood in this organism. We have

developed and applied a new method for genome-wide identification of capsule regulators.

Using this method, many genes that positively or negatively affect capsule production in *K.*

pneumoniae were identified, and we use these data to propose an integrated model for

capsule regulation in this species. Several of the genes and biological processes identified

have not previously been linked to capsule synthesis. We also show that the methods

presented here can be applied to other species of capsulated bacteria, providing the

opportunity to explore and compare capsule regulatory networks in other bacterial strains and

species.

Introduction

Klebsiella pneumoniae is an ubiquitous Gram-negative bacterium, found both in the

environment and as an asymptomatic coloniser of the mucosal surfaces of mammals (1). *K.*

pneumoniae is also an opportunistic pathogen, and can express a multitude of virulence

factors which enable it to cause infections in humans (1–4). Historically associated with

infections in the immunocompromised and in neonates (1, 5), focus has been directed to *K.*

pneumoniae following the emergence of antimicrobial-resistant and hypervirulent lineages

(6, 7). Hypervirulent lineages are a particular concern in a clinical setting, because they have

the potential to cause infection in immunocompetent hosts (6, 8, 9), to metastasise (10), and

to cause infections in unusual infection sites (11).

Numerous factors contribute to *K. pneumoniae* virulence, such as the ability to produce

siderophores, fimbriae, lipopolysaccharide, and extracellular polysaccharide capsule (6, 12–

14). Hypervirulence is associated with these factors, particularly with the overproduction of

capsular polysaccharide (6, 12, 15, 16), and in the absence of these virulence factors, *K.*

pneumoniae virulence is reduced or abolished (14, 17, 18). The ~200 kilobase *K. pneumoniae*

virulence plasmid, which is also associated with the hypervirulent phenotype (6), encodes

siderophores such as aerobactin and salmochelin, and positive regulators of capsule

biosynthesis (6, 19–21). More than 100 capsule locus types have been identified in *K.*

pneumoniae (17), though the majority of clinical cases of *K. pneumoniae* infection are caused

by strains of capsule types K1 and K2 (12, 22).

Excessive capsule production is strongly associated with hypervirulence in *K. pneumoniae*

(23–25) and several studies have sought to identify genetic determinants of hypervirulence.

For example, the mucoviscosity-associated gene *magA* (now named *wzy_K1* (26, 27)) was

originally identified by transposon mutagenesis screening (28). The *rmpA* and *rmpA2* genes

also affect hypermucoviscosity, and encode transcription factors that positively regulate the

K. pneumoniae capsule biosynthesis locus (6, 15, 16, 29). These regulators can be either

chromosomally-encoded or plasmid-borne. Although *rmpA* is correlated with hypervirulence,

and strains lacking *rmpA* and aerobactin are avirulent in mice (12, 30), it has been shown that

this increased virulence is a consequence of the hypermucoviscous phenotype conferred by

rmpA, rather than the gene itself (24).

Capsule is also a potential therapeutic target in *K. pneumoniae*. Capsule-targeting monoclonal

antibodies increased the killing of *K. pneumoniae* ST258 (an outbreak lineage) by human

serum and neutrophils (31, 32) and limited the spread of a respiratory *K. pneumoniae* ST258

infection in mice. Specific capsule-targeting bacteriophage have been shown to clear or limit

infections caused by *K. pneumoniae* strains of capsule types K1, K5 and K64 (33–35), and in

some cases protection could also be achieved by treatment with the capsule depolymerase

enzymes produced by these phage, rather than the phage itself. Translating these early-stage

findings to the clinic is a priority as extensively drug-resistant *K. pneumoniae* strains become

more prevalent (36).

Despite the absolute requirement for capsule in *Klebsiella* infections, its promise as a

therapeutic target, and the connection between capsule overproduction and hypervirulence,

the biosynthetic and regulatory mechanisms governing *K. pneumoniae* capsule have not yet

been fully explored. Genetic screens and reverse genetics (28, 37–41), including transposon

mutagenesis approaches (38), have been used to identify biosynthetic genes and activators of

capsule production (including the *rmpA* and *magA* genes), and some of the cues that elevate

capsule expression above basal levels have also been described – these include temperature,

iron availability and the presence of certain carbon sources (12, 42). Defining the capsule

regulatory network of *Klebsiella pneumoniae* in more detail would not only deepen our

understanding of this pathogen's interaction with the host environment, but could also inform

efforts to target this factor with new therapeutics.

Here we employed transposon directed insertion sequencing (TraDIS) (43, 44) to identify

genes influencing *K. pneumoniae* capsule production. We performed density-based selection

on mutant libraries of both K1 and K2 capsule type *K. pneumoniae*, using a discontinuous

density gradient (45). This approach allowed for the simultaneous selection of both

capsulated and non-capsulated mutants from mutant libraries, without requiring growth of

bacteria under a selective pressure. We have identified 78 genes which, when mutagenised by

transposon insertion, reduce the ability of one or both of these strains to manufacture capsule.

These included multiple genes not previously associated with capsule production in *K.*

pneumoniae. We have also identified 26 candidate genes in NTUH-K2044 which cause an

increase in capsulation when disrupted. Our results allow us to present an integrated model

for capsule regulation in *K. pneumoniae*, and establish a technology for study of capsule

production that is applicable to other bacterial species.

Results

Density gradient centrifugation separates bacterial populations based on capsule

phenotype

K. pneumoniae capsule influences the centrifugation period required to pellet cells, and this

property has been used to semi-quantitatively compare capsule production between different

strains (38). Based on this observation, we speculated that *Klebsiella* cells of different

capsulation states could be separated by density-based centrifugation, and that this method

could be combined with TraDIS to screen for capsule-regulating genes. Discontinuous

Percoll density gradients are routinely used to purify macrophages from complex samples

(46), and have also been used to examine capsulation in *Bacteroides fragilis* and

Porphyromonas gingivalis (45, 47). Tests with *K. pneumoniae* NTUH-K2044 (capsule type

K1, hypermucoid, Fig. 1A), *K. pneumoniae* ATCC43816 (K2, hypermucoid) and a non-

capsulated *E. coli* control showed that these strains differed in density and migrated to above

15%, above 35% and below 50% Percoll, respectively (Fig. S1A). Growth of the *Klebsiella*

strains at 25 °C (which reduces capsule production) decreased their density, with NTUH-

K2044 localising to above 35% Percoll, and ATCC43816 showing limited migration into the

35% layer. A third *K. pneumoniae* strain, RH201207 (non-hypermucoid, capsule type K106),

was retained above 50% Percoll (Fig. S1A). We also examined the proportion of *Klebsiella*

bacteria migrating to different locations within the density gradient (Fig. S1B). While the

majority of *K. pneumoniae* NTUH-2044 bacteria were located above the 15% layer, a small

fraction (4% viable count of the top) migrated to below the interface of this layer, suggesting

some heterogeneity in capsule production. In both NTUH-K2044 and ATCC43816, only a

very small number of cells were present below the 50% layer (Fig. S1B); these numbers

could be partially due to cross-contamination during extraction of this part of the gradient,

which was recovered by pipetting from the top.

We then tested this method with a high-density transposon insertion library of *K. pneumoniae*

NTUH-K2044 (generated as described in Materials and Methods). This mutant library gave

rise to three fractions on a 15%-35%-50% Percoll gradient (Fig. 1B). The middle and bottom

fractions separated from the NTUH-K2044 library contained less capsule than the input

library culture, as shown by quantification of capsular uronic acids (Fig. 1C) and the

hypermucoidy centrifugation test (Fig. S1C). To determine whether altered migration in a

Percoll gradient was the result of stable mutant phenotypes, the fractions of the *K.*

pneumoniae NTUH-K2044 gradient were cultured overnight and centrifuged on a fresh

gradient. The "bottom" and "top" fractions migrated to the same position as before, while the

"middle" fraction showed a partially heritable phenotype and was distributed across the top

and middle positions (Fig. S1D).

These data indicated that centrifugation on a discontinuous density gradient successfully

separated *K. pneumoniae* populations based on their capsule production, and that variations

in density among library mutants were, in part, due to stable phenotypes. Random-prime PCR

of several colonies from the bottom fraction of the *K. pneumoniae* NTUH-K2044 library

identified insertions in known capsule biosynthetic or regulatory genes (Supplementary text

S1), and a *wza* insertion mutant was retained as a capsule-negative control for future

experiments. We then wished to determine whether our method could be used for capsule-

based separations of other species, for which we tested capsulated and non-capsulated

Streptococcus pneumoniae 23F. These strains were separated reproducibly on a Percoll

gradient (Fig. 1D), indicating that our method is applicable to other bacterial species.

***Density-TraDISort identifies multiple capsule-associated genes in* Klebsiella pneumoniae**

TraDISort is a term for transposon sequencing screens based on the physical, rather than

survival-based, enrichment of insertion mutants from saturated libraries (48), which allows

examination of phenotypes that are not linked to survival. We call our approach density-

TraDISort to distinguish it from the original method using fluorescence-based flow sorting as

the physical selection method. The final density-TraDISort approach is shown (Fig. 1B).

Briefly, transposon mutant libraries were grown overnight at 37 °C in LB, applied to the top

of a Percoll gradient and centrifuged at a moderate speed for 30 minutes (see Materials and

Methods). Following centrifugation, each bacteria-containing fraction was extracted and

subject to TraDIS, as was a sample of the input culture. Data were analysed using the Bio-

TraDIS pipeline (see Materials and Methods). The *K. pneumoniae* NTUH-K2044 library

contained approximately 120,000 unique transposon insertion sites (equivalent to an insertion

every 45 bp), with a median of 14 insertion sites per gene. Statistics summarising the

sequencing results from each fraction are reported in Table S2.

Unlike traditional growth-based transposon-insertion screens, our density-TraDISort setup

combines positive and negative selection within a single experiment: mutants with reduced

capsule can be identified through their loss from the top fraction, or by virtue of their

enrichment in another fraction (see Fig. 2A for an example). We applied stringent cutoffs

based on both selections to identify putative capsule-related genes. Briefly, a gene was only

counted as a hit if it was (i) lost from the top fraction ($\log_2$FC < -1, q-value < 0.001) and (ii)

enriched in another fraction ($\log_2$FC > 1, q-value < 0.001) (see Tables S3, S4). The presence

of a "middle" fraction also enabled us to identify genes which increased capsulation when

disrupted (example in Fig. 2B). These were apparent in the raw TraDIS plot files as genes

where all or nearly all the mutants localised to the top fraction following centrifugation

(mutants with wild-type capsule were distributed between the top and middle fractions, in

keeping with the migration pattern of wild-type NTUH-K2044 (Fig. S1A)). We defined

putative "capsule up" genes as those which showed a marked depletion in the middle fraction

relative to the input ($\log_2$FC < -3; q-value < 0.001), with genes with very low initial read

counts ($\log_2$CPM < 4) excluded (Tables 1, S4). Examples of TraDISort data for several genes

known to affect capsule production are shown in Figure S2C.

In total, we identified 62 genes required for full capsule production in *K. pneumoniae* NTUH-

K2044 (Tables 1, S4). We also identified 26 putative "capsule up" mutants in this strain

(Tables 1, S4). The biological roles of the genes determined to influence capsule production

are discussed in more detail below.

***Global regulators, metabolic genes and cell surface components affect capsule in* K.**

pneumoniae

Our "capsule down" mutants included almost every gene of the capsule biosynthesis locus of

K. pneumoniae NTUH-K2044 (Fig. 2A), further confirming that our experimental strategy

had successfully isolated capsule-deficient mutants. The *manB* and *ugd* genes were not called

as hits, however these contained very few insertions in the input sample (visible in the plot

files above these genes in Fig. 2A). In addition to biosynthetic genes, the known capsule

regulators *rmpA*, *rcsB* and *rfaH* were identified (Fig. S3C).

Other "capsule down" hits included multiple cell surface components, metabolic genes and

genes of global regulatory systems. Extended functional information for all of our hits,

including reported links to capsule production in *Klebsiella* and other proteobacteria, is

included in Table S4. Gene enrichment analysis using the TopGO package (49) indicated that

genes for polysaccharide metabolic processes ($p = 1.7 \times 10^{-5}$), molecular transducers ($p =$

0.0042) and protein kinases ($p = 0.005$) were overrepresented in our hits (all probabilities are

from Fisher's exact test, parent-child algorithm (50)).

Several global regulators were identified as affecting capsule production. These included the

anaerobic/redox-sensitive sensor kinase ArcB, the osmotic stress response system OmpR-

EnvZ, the BarA-UvrY system, the arginine repressor ArgR, and the carbon storage regulatory

small RNA CsrB. The transcription regulators MprA and SlyA were also needed for full

capsule production. Metabolic genes were also identified, including genes of the electron

transport chain, glycolytic enzymes, and the nitrogen source regulatory protein GlnB.

The "capsule up" mutants of NTUH-K2044 included nucleoid proteins, membrane-bound

transporters such as SapBCDF (Fig. 2B), and several metabolic genes (Table S4).

Interestingly, some of the "capsule up" hits are known antagonists or regulators of genes in

the "capsule down" hits: CsrD (which targets CsrB for degradation), H-NS (reported to be

antagonised by SlyA), and GlnD (modifies and controls activity of GlnB).

Finally, our results also included many genes for other cell surface components; genes for the

synthesis and modification of LPS were identified, along with enterobacterial common

antigen genes, and the lipoprotein Lpp.

Distinct and overlapping capsule regulators in two* K. pneumoniae *strains

We wished to determine the extent of conservation of our capsule-regulatory hits across the

Klebsiella species, and the overlap between the capsule-influencing genes in different

Klebsiella strains. To explore this question, we applied our density-TraDISort method to a

second strain, *K. pneumoniae* ATCC43816, a capsule type K2 strain which is commonly used

in *Klebsiella* infection studies (Fig. S3A). We constructed a *K. pneumoniae* ATCC43816

saturated transposon insertion library with ~250,000 unique insertion sites (or an insertion

every 22 bp) and a median of 36 insertion sites per gene (see Materials and Methods; Fig.

S3C-D). This library resolved into two fractions on a 35%-50% Percoll gradient, with no

obvious heritable "middle" fraction, and fractions were collected and their transposon

insertion junctions sequenced as for *K. pneumoniae* NTUH-K2044. Genes were counted as

hits in this strain if they were lost from the top fraction relative to the input ($\log_2$FC < -1, q-

value < 0.001) and enriched in the bottom fraction ($\log_2$FC > 1, q-value < 0.001). The lack of

a "middle" fraction in this experiment means there is less sensitivity for identifying reduced-

capsule mutants, particularly using the second, positive selection-based filtering criterion. We

used a pan-genome generated from 263 annotated *K. pneumoniae* genomes (Table S5) to

define the common genes in NTUH-K2044 and ATCC43816 and determine their prevalence

across the *K. pneumoniae* population (Tables 1, S4).

We identified 34 candidate capsule-regulatory genes in *K. pneumoniae* ATCC43816. As

observed for NTUH-K2044, the genes of the capsule biosynthetic locus were nearly all called

as hits (Fig. S3E). Of the three genes that were not, two had very low initial insertion counts,

and the third met our first selection criterion of being lost from the top fraction but was not

enriched in the bottom fraction. Putative capsule-influencing genes of ATCC43816 fell into

diverse functional categories, with cell surface components, metabolic genes, transporters and

known regulators implicated.

Although the majority of capsule hits are encoded in the core genome (48 of 62 in *K.*

pneumoniae NTUH-K2044; 23 of 32 in *K. pneumoniae* ATCC43816, Fig. 2C), only sixteen

genes were called as hits in both strains. These shared genes included five shared components

of the two strains' capsule biosynthesis loci, the transcription antiterminator *rfaH*, two

enterobacterial common antigen genes, three genes of the *arn* operon responsible for

modification of LPS lipid A with L-Ara4N, and the glutathione reductase *gor*. Strain-specific

differences were also identified, with the caveat that NTUH-K2044 produces more capsule

(see Fig S3B) and therefore afforded us greater sensitivity in our experiment. Genes that were

identified as required for ATCC43816 capsule production but not NTUH-K2044 included the

yrbCDEF (*mlaCDEF*) ABC transporter and the *mlaA* gene, which are involved in

maintaining outer membrane asymmetry through the cycling of phospholipids (51). Thirty-

two genes common to both strains had a "capsule down" phenotype in NTUH-K2044 but

were not called as hits in ATCC43816 (note that "capsule up" hits are not included in our

comparison as these were not resolved in ATCC43816). These included the electron transport

pump *rnf*, global regulators such as *arcB* and *ompR*, the transcription factor *mprA*, and

several additional cell surface component biosynthetic genes. It appears that for at least some

conserved genes their influence on capsule is strain-specific.

Phenotypes of single-gene mutants confirm results of density-TraDISort

We generated a set of ten single-gene deletion mutants in *K. pneumoniae* NTUH-K2044, and

three in *K. pneumoniae* ATCC43816, in order to validate the results of our density-TraDISort

screen. Genes selected for mutagenesis were the known capsule and LPS regulator *rfaH* in

both strains, the LPS O-antigen ligase *waaL*, the aerobic respiration control sensor *arcB*, and

multiple transcriptional regulators (*ompR*, *argR*, *slyA*, *mprA*, *uvrY*). We also deleted the *arnF*

gene in ATCC43816. The *sapBCDF* ABC transporter in NTUH-K2044 was examined in

order to validate our assignment of "capsule up" hits.

Single gene knockout mutants were grown under the same conditions as in the original screen

and subjected to density gradient centrifugation. Every mutant showed a banding pattern

consistent with the results of density-TraDISort (Fig. 3A-B). The $\Delta arcB$ mutant appeared to

have two populations, one with wild-type and one with reduced capsule, most other NTUH-

K2044 mutants migrated to the position of the middle fraction, while the $\Delta waaL$, i-*wza* and

$\Delta rfaH$ mutants migrated to the bottom of the gradient. The $\Delta sapBCDF$ mutant showed higher

density than wild-type and remained above the 15% Percoll layer, with no movement into the

gradient itself. Mutants of NTUH-K2044 were also tested for hypermucoidy and uronic acid

production, and these experiments showed increased capsule in NTUH-K2044 $\Delta sapBCDF$

and reduced capsule in all other mutants (Fig. 3C, Fig. S4B). Of the *K. pneumoniae*

ATCC43816 mutants (Fig. 3B), $\Delta rfaH$ migrated to the bottom of the gradient and $\Delta arnF$ to

the middle and bottom, while $\Delta mprA$ showed the same pattern as wild-type. This is consistent

with the TraDISort assignment of this gene as having a strain-specific effect on capsule, at

least in these growth conditions (Table 1).

Our hits included several genes with roles in LPS biosynthesis, which raised the possibility

that LPS O-antigen may affect cell density independently of capsule. To test this, we

constructed a $\Delta waaL$ deletion in our *wza* transposon insertion strain (see Materials and

Methods). The resulting double mutant had no detectable reduction in density compared to

the *wza* mutant on the standard 15%-35%-50% gradient or on 70% Percoll (the minimum

concentration required to exclude the *wza* mutant, Fig. S4A), indicating that the LPS O-

antigen alone does not affect the density of *K. pneumoniae*, at least within the resolution of

this experiment.

Virulence and capsule architecture of* K. pneumoniae *NTUH-K2044 ΔargR, ΔmprA,

Δ*sapBCDF* and Δ*slyA*

We selected the transcription factors ArgR, SlyA and MprA, along with the ABC transporter

SapBCDF, for further characterisation. ArgR represses arginine synthesis and transport as

well as other genes (52), SlyA is an antagonist of H-NS (known to suppress capsule in *K.*

pneumoniae) (53, 54), and MprA is a transcriptional regulator with an effect on capsule in

UPEC (55). Both SlyA and MprA were very recently also shown to be virulence and capsule

regulators in *K. pneumoniae*, and were renamed KvrA and KvrB (56). SapBCDF has been

reported to mediate resistance to antimicrobial peptides (AMPs) in *H. influenzae* by

importing them for degradation (57) and was presumed to have this activity in

Enterobacteriaceae as well, though it has recently been reported that this pump functions as a

putrescine exporter in *E. coli* and has no role in AMP resistance (58). ArgR and SapBCDF

have not previously been linked to capsule regulation.

To confirm that the alterations in capsule production observed in the NTUH-K2044 Δ*argR*,

Δ*slyA*, Δ*mprA* and Δ*sapBCDF* mutants were due to the deleted genes, each mutant was

complemented by reintroducing the wild-type gene on the chromosome (see Materials and

Methods). Complementation caused a complete restoration of wild-type capsule production,

as measured by the hypermucoidy and uronic acid assays (Fig. 4A-B). To define changes in

capsule architecture, each mutant strain was examined by transmission electron microscopy

(Fig. 4C). Wild-type *K. pneumoniae* NTUH-K2044 had a thick, filamentous capsule of

roughly half the cell diameter. The Δ*argR* and Δ*slyA* mutants had capsules with slightly

reduced thickness and finer filaments, while the Δ*mprA* mutant had extremely fine and

diffuse filaments such that the boundary of the capsule was not clear. The Δ*sapBCDF* capsule

had some thick filaments but at lower density than NTUH-K2044, with an additional gel-like

layer visible outside these filaments. The virulence of the Δ*argR*, Δ*slyA*, Δ*mprA* and

Δ*sapBCDF* mutants, and their complements, was assessed by infection of research-grade

Galleria mellonella larvae, an established invertebrate model for *Klebsiella* infections (59).

Each of the reduced-capsule strains showed a virulence defect relative to the wild-type strain

which was restored on complementation (Fig. 5A), while the Δ*sapBCDF* mutant did not have

changed virulence compared to the wild-type.

The Sap transporter alters serum survival but does not affect antimicrobial peptide

resistance

We then examined the effect of ArgR, MprA, SlyA and SapBCDF on resistance to human

serum. After two hours, NTUH-K2044 wild type showed full survival with a slight increase

in viable count, while the *wza* mutant was reduced in viable count by ~25-fold. The Δ*argR*,

Δ*mprA* and Δ*slyA* mutants did not change significantly (Fig. 5B). The Δ*sapBCDF* showed

increased viable count compared to the wild type strain, with a 7-fold increase over the

course of the experiment. This increase was unexpected, as the wild type strain is already

fully serum resistant. To determine if the increased serum survival was dependent on capsule

production, a double Δ*sapBCDF cps* mutant was constructed (see Materials and Methods).

This strain showed a drastic reduction in survival, suggesting that the NTUH-K2044

Δ*sapBCDF* mutant can grow better in serum because of its increased capsule (Fig. 5B).

We also tested the Δ*sapBCDF* mutant for resistance to the peptide antibiotics colistin and

polymyxin B. The MIC was the same as that of the wild type, at 1 µg/ml for colistin, and

0.75 µg/ml for polymyxin B, indicating that the Sap transporter in *K. pneumoniae* does not

contribute to antimicrobial peptide resistance.

Sap mutation increases transcription of capsule middle genes without activating the Rcs

system

We then wished to determine whether the mutation of the Sap transporter increased capsule

production by acting on transcription. RNA was extracted from late exponential-phase wild

type and Δ*sapBCDF* cells, and the abundance of three capsule locus transcripts – *manC*,

wcaG and *wza* – was measured by qRT-PCR. These genes are transcribed from separate

promoters. The Δ*sapBCDF* mutant showed elevated expression of *wcaG*, at 2.5 times wild

type levels, while expression of *wza* and *manC* was not significantly changed (Fig. 5C).

The Rcs phosphorelay system regulates capsule expression in *E. coli* and other

enterobacteriaciae, and is induced by cues such as membrane stress (60, 61). RcsA is a

component of the system which autoregulates and increases its own expression when

activated. We measured *rcsA* transcript levels to determine whether loss of the Δ*sapBCDF*

genes induces *rcsA* (Fig. 5C). Unexpectedly, levels of *rcsA* were much lower in the mutant

than the wild-type, indicating that Sap-dependent induction of capsule expression does not

occur through *rcsA*. Note, however, that RcsA is not required for all permutations of Rcs

signalling as RcsB can interact with a number of partner proteins to regulate transcription

(60).

***Capsule is at the centre of a complex regulatory network in* Klebsiella pneumoniae**

We identified numerous putative capsule regulators by density-TraDISort, and validated the

results of our screen with single-gene deletion mutants. We propose an integrated model for

how the genes we identified may collectively control *K. pneumoniae* NTUH-K2044 capsule.

This model is based on our results and previous published work in *K. pneumoniae* and other

Enterobacteria (particularly *Escherichia coli*). Full details of the literature relevant to each hit

are in Table S4.

Major nodes for transcriptional control are the CsrB carbon source utilisation system and the

Rcs phosphorelay system. Each of these systems is itself regulated by multiple genes

identified in our study – CsrB integrates signals from the UvrY-BarA two component system

(a "capsule down" hit) and various carbon metabolic genes, is activated by DksA, and is

targeted for degradation by CsrD (a "capsule up" hit); the Rcs system is induced by MdoGH

mutation, can cooperate with RmpA and RmpA2 to induce capsule, and also responds to

carbon metabolism and some forms of enterobacterial common antigen. SlyA/KvrA and

MprA/KvrB both promote capsule transcription (56). The SlyA/KvrA protein acts as a

temperature-dependent switch which acts by relieving H-NS mediated transcriptional

silencing; H-NS has been reported to operate on both the Rcs system and the capsule operon

in *E. coli*, though the precise target of its activity in *Klebsiella* is not known. Capsule is also

affected by the composition of the cell envelope, and mutations in *lpp* or various LPS-related

genes can reduce the retention of capsule at the cell surface. Note that several genes related to

cell envelope composition and membrane stress have been shown to regulate the Rcs system

(61), therefore some of the cell envelope component genes identified in our study may act

through RcsB. Our work has also uncovered novel regulators of capsule that, at this stage,

cannot be tied to the wider regulatory network, such as *argR*, and the ABC transporter Sap.

We intend to define the mechanisms by which these genes affect capsule in future studies.

Discussion

We have developed a simple, robust technology for genome-wide studies of bacterial

capsule, density-TraDISort, and applied it to identify capsule regulators in two strains of *K.*

pneumoniae. In doing so, we have identified multiple positive and negative regulators of

capsule production, including several genes not previously linked to capsule in this species.

To our knowledge, this is one of the first studies employing a physical selection independent

of bacterial survival and growth to separate TraDIS libraries, and is the first time that density-

based physical selection has been applied to studying capsule regulation in *K. pneumoniae*.

TraDISort/FAST-INSeq technology with fluorescence-based sorting has to date been used to

identify genes affecting efflux of ethidium bromide, and mutations influencing expression of

a *Salmonella* Typhi toxin reporter (48, 62). We have expanded the utility of this method by

adding a selection step based on cell density, allowing us to resolve different capsulation

states. We envisage that, in addition to facilitating genome-wide screens for altered

capsulation in other bacterial species, density-TraDISort could be used to identify genes

affecting cell size and shape, or aggregation.

Our study is the second application of TraDIS to screen for genes affecting bacterial capsule

production, following a recent study focused on UPEC (55). The UPEC study utilised a

capsule-specific phage to positively select transposon insertion mutants lacking capsule; two

novel capsule regulators were identified in this way. Compared with phage-based selection,

our method offers increased sensitivity – mutants with a range of capsule phenotypes can be

identified, in addition to capsule-null mutations. In addition, there is the option for very

stringent selection of hits, as cutoffs can be applied based on both negative selection (loss

from the top fraction) and positive selection (enrichment in another fraction). However,

density-based selection is less specific to capsule than phage infection, and there is the

possibility that mutations could affect cell density in a capsule-independent manner.

Interestingly, one of the novel *Klebsiella* capsule regulators identified in this study, MprA,

was also shown to regulate capsule in UPEC. In *Klebsiella pneumoniae* this gene increases

capsule production above a baseline in hypermucoid strains (Fig 3A, S4B) and (56)), while in

UPEC an Δ*mprA* mutant did not produce capsule at all.

We have shown that capsule production in *K. pneumoniae* NTUH-K2044 is controlled by

many different global regulatory systems, allowing us to describe a detailed snapshot of the

control of capsule in this strain (Fig. 5). Note that our assay was performed on bacterial cells

at late stationary phase growth, in LB medium, under microaerophilic conditions. This

condition was used in this study because capsule production is high, offering good resolution

for capsule-based selection of mutants. Additional regulators, linked to different cues and

stresses, are likely to be involved in different environments. Many of the regulators identified

in this study were only called as hits in the hypermucoid strain, *K. pneumoniae* NTUH-

K2044. It remains to be seen whether these same regulators control capsule (though to a

degree outside the resolution of our gradient) in other *K. pneumoniae* strains, though note that

several genes of *K. pneumoniae* ATCC43816 (including *uvrY*, *barA*, *csrB*, *rcsA* and *rcsB*)

met our first screening criterion of being lost from the top fraction but not the second of being

enriched in the bottom fraction. We speculate that capsule production is subject to complex

environmental control across the *Klebsiella* species, but that the hypermucoid phenotype is

more expensive to maintain, and more sensitive to disruptions in its regulatory network.

Many of our hits are involved in the synthesis of other cell surface polysaccharides; these

included genes for enterobacterial common antigen (ECA), as well as genes for the synthesis

or modification of LPS. ECA is a non-immunogenic surface glycolipid found in various

forms in Enterobacteriaceae, and structural modifications in this moiety can induce the Rcs

system (63, 64). LPS is a major contributor to *K. pneumoniae* pathogenesis in sepsis, though

to a lesser extent in pneumonia (12, 18), and various LPS modifications have roles in immune

modulation during infection (65, 66). We are confident that the LPS mutations identified in

our study affect capsule retention or biosynthesis, rather than density *per se*, because 1)

deletion of the O-antigen ligase *waaL* did not reduce cell density in an acapsular *K.*

pneumoniae NTUH-K2044 strain (Fig. 3A); 2) some LPS biosynthesis genes, but not all,

were hits in our screen; and 3) the glucuronic acid moieties on the core LPS polysaccharide

are required for capsule retention in *K. pneumoniae* (67, 68). In both strains we studied,

disrupting genes of the *arn* operon reduced capsulation (Table 1, S4). The *arn* operon is

responsible for modifying lipid A of LPS with 4-amino-4-deoxy-L-arabinose to mediate

resistance to peptide antibiotics (69), but has not previously been linked to capsule. The

arnEF genes encode a flippase thought to translocate the modified arabinose across the cell

membrane (70), while *arnD* is involved in its biosynthesis (71). We hypothesise that the

reduced capsule of mutants of *arnD* and *arnEF* on capsule is independent of Lipid A

modification, because other genes in this operon did not affect capsule, and because a

previous study showed that Lipid A modification with L-Ara4N does not occur in cells

grown in LB (66). Overall, our results hint at a high degree of interdependence between the

three major surface polysaccharides of *K. pneumoniae*.

The *sap* ABC transporter was found to promote capsule production when mutated, by

increasing the expression of capsule middle genes (Fig. 3A, Fig. 4A, Tables 1, S3, S4). To

our knowledge our study is the first to implicate *sapABCDF* in capsule regulation, though its

full functions, or indeed the substrate of this transporter in *Klebsiella*, are not known. The *H.*

influenzae Sap homologue mediates resistance to antimicrobial peptides by importing them

for degradation, and is also required for haem uptake (57, 72), while in *E. coli* the Sap pump

is reported to export putrescine and facilitate potassium import through TrkGH (58, 73). We

found that the Sap transporter did not affect antimicrobial peptide resistance, which was also

observed in *E. coli*. It is unclear how Sap mutation induces *wcaG* while suppressing an

important component of the Rcs system – more work will be needed to define the role and

mechanism of this transporter in *K. pneumoniae*.

We have developed a simple, broadly-applicable method for studies of capsulation, and used

it to define the regulatory network that controls capsule in *K. pneumoniae* NTUH-K2044. We

have also identified genes required for full production of capsule in a K2 strain. Although the

majority of regulators are located in the core genome of *K. pneumoniae*, there are differences

in the specific regulators deployed in the two strains we investigated, and it would be

interesting to determine whether this pattern of strain-specific regulatory networks

comprising primarily core genes holds across the *Klebsiella* phylogeny. This intra-species

comparison, together with our data showing that density-based capsule selection can be used

in other capsulated bacteria, also opens the possibility for robust interspecies comparisons of

capsule regulation.

Materials and Methods

Culture conditions and microscopy

K. pneumoniae strains were cultured routinely in LB media supplemented with 1.5% w/v

agar as appropriate. Cultures were supplemented with 12.5 µg/ml chloramphenicol and 12.5

µg/ml tetracycline when required. *S. pneumoniae* strains were grown on blood agar plates

(Oxoid, CM02718) in microaerobic candle jars containing Campygen sachets at 37 °C, or in

static BHI liquid media (Oxoid, SR0050C). The list of strains, plasmids, and oligonucleotides

used in this study is reported in Table S1.

Generation of transposon insertion libraries

TraDIS libraries were generated using the mini-Tn*5* transposon delivery plasmid pDS1028

(74), introduced into the recipient strain by conjugation. Full details are described in

Supplementary methods.

Mutant library fractionation on Percoll gradients

Bacterial mutant libraries were separated on the basis of their capsule expression by

centrifugation on a discontinuous Percoll (GE Healthcare) density gradient for 30 minutes at

3,000 x *g* (Fig. 1B). Full details are described in Supplementary methods.

Identification of transposon insertion sites by random-prime PCR

Genomic DNA was prepared from overnight cultures of single reduced-capsule mutants

using the DNeasy® Blood and Tissue kit (Qiagen). Random-prime PCR to identify the

transposon insertion site in each gDNA template was performed as described (75) using

primers FS57-59 and FS109 and Herculase II polymerase (Agilent). Amplicons were

sequenced using primer FS107.

DNA extraction and next-generation sequencing

Genomic DNA (gDNA) was prepared from each Percoll-resolved fraction by phenol-

chloroform extraction. From each gDNA preparation, 2 µg DNA was used to prepare TraDIS

transposon-specific sequencing libraries as described previously, using primer FS108 for

specific amplification of transposon junctions (43). Sequencing was carried out on the

Illumina MiSeq platform using primer FS107.

Analysis of TraDIS data

The analysis of TraDIS sequencing results was carried out using the Bio-TraDIS pipeline as

described previously (43, 44), with minor modifications (see Supplementary methods). All

scripts used in this study are available at https://github/sanger-pathogens/Bio-Tradis and

https://github/francesca-short/tradis_scripts. Comparisons between fractions were based on

normalised read counts per gene. Genes with (i) reduced mutant abundance in the top

fraction, and (ii) increased mutant abundance in the middle or bottom fraction were called as

decreased capsule hits, with thresholds of an absolute change in $\log_2 FC > 1$ and q-value <

0.001. Increased capsule hits in NTUH-K2044 were defined as those that with severely

reduced mutant abundance in the middle fraction ($\log_2 FC < -3$; q-value < 0.001) without

enrichment in the bottom fraction ($\log_2 FC < 1$), with genes containing very few reads in any

fraction excluded (*i.e.*, $\log_2$(counts per million) in the top fraction was greater than 4).

Generation of the pan-genome, and enrichment analysis, is described in Supplementary

methods.

Construction of single-gene deletion strains

Single gene knockout mutants were constructed in *K. pneumoniae* by allelic exchange.

Upstream and downstream sequences (> 500bp) for each target gene were amplified and

joined by overlap PCR, cloned into pKNG101-Tc, and introduced into the recipient strain by

conjugation with the *E. coli* β2163 donor strain. All primers used, and the resulting

constructs, are listed in Table S1. Conjugation patches were incubated for one hour at 37 ℃,

then for 16 hours at 20 ℃. Single-crossover mutants were selected on LB agar + 15 µg/ml

tetracycline. Double-crossover mutants were selected on low-salt LB agar + 5% sucrose at

room temperature, and were subsequently patched onto LB + sucrose and LB + tetracycline

plates to confirm loss of the vector. Mutants were confirmed by PCR across the deleted

region. Mutants were complemented by introduction of the relevant gene back into its

original location on the chromosome by allelic exchange as described above, using a vector

encoding the gene and its flanking region. The Δ*sapBCDF cps (KP1_3713)* double mutant

was generated by random transposon mutagenesis of the Δ*sapBCDF* strain with the pDS1028

vector, followed by selection of acapsular mutants from the pool by density-gradient

centrifugation and random-prime PCR to identify the insertion site.

Quantification of capsule by uronic acid assay

Capsule extraction and quantification of uronic acids was performed as described previously

(14, 76), with modifications (see Supplementary methods).

Hypermucoviscosity assay

Cultures of *K. pneumoniae* were grown overnight in 5 ml LB medium at 37 °C. These

cultures were sedimented at 1,000 x *g* or 2,500 x *g* for 5 minutes (room temperature). The

OD_{600} of the top 500 µl of supernatant was determined by spectrophotometry. Results were

expressed as a ratio of the supernatant OD_{600} relative to the input culture.

Electron microscopy

Colonies were taken directly from an agar plate, high-pressure frozen in a Balzers HP010,

and freeze-substituted for 8 hours in acetone containing 0.1% tannic acid and 0.5%

glutaraldehyde at -90 ˚C followed by 1% osmium tetroxide in acetone for 24 hours at -50°C.

They were then embedded in Lowicryl HM20 monostep resin. Ultrathin sections were cut on

a Leica UC6 ultramicrotome and contrasted with uranyl acetate and lead citrate. Images of

bacteria were taken on an FEI Spirit Biotwin 120 kV TEM with a Tietz F4.15 CCD.

Serum resistance assay

Bacteria were grown in LB to an OD_{600} of 1, pelleted and resuspended in sterile PBS. Human

sera (400 µl; Sigma-Aldrich S7023) was pre-warmed to 37 °C and added to 200 µl bacterial

suspension, and the mixture was incubated at 37 °C for two hours. Viable bacterial counts

were determined before and after incubation.

Galleria mellonella *infection*

Larvae of *G. mellonella* were purchased from BioSystems Technology UK (research-grade

larvae) and used within one week. Bacteria were grown overnight, subcultured and grown to

an OD_{600} of 1, then resuspended in sterile PBS. Larvae were infected by injecting the

bacterial suspension (10^5 cells) into the right hind proleg of the larvae using a Hamilton

syringe. Infected larvae were incubated at 37 °C and monitored every 24 h, and were scored

as dead when they were unresponsive to touch. Thirty larvae were used per strain, and these

were infected in three batches of 10 using replicate cultures.

Antimicrobial peptide resistance tests

Strains were grown overnight, subcultured and grown to an OD_{600} of 1.0. This culture (100

µl) was spread on the surface of an LB agar plate, dried, and an E-test (BioMerieux) strip was

placed on the surface of the plate. Plates were incubated face-up at 37 °C and the result was

read after 6 hours to avoid overgrowth of the capsule interfering with the reading.

RNA extraction and qRT-PCR

Bacteria were grown in LB medium at 37 °C to an OD_{600} of 1.0. Cultures were mixed with 2

x volumes RNAProtect reagent, centrifuged, and RNA extracted using the Masterpure

Complete DNA and RNA Purification kit (Epicentre) according to the manufacturer's

instructions. Samples were then subject to in-solution DNaseI digest (Qiagen), and cleaned

up using the Qiagen RNeasy Mini kit. Reverse transcription of 200 ng RNA was performed

using the ProtoScriptII enzyme (NEB) with supplied instructions.

Transcripts were quantified using a StepOne real-time PCR instrument with the KAPA

SYBR FAST qPCR kit. Relative abundance was determined using the relative standard curve

method with *K. pneumoniae* NTUH-K2044 gDNA as a standard, and *recA* as the reference

gene (77).

Accession numbers

Sequences generated during this study have been deposited into the European Nucleotide

Archive (ENA; http://www.ebi.ac.uk/ena) under study accession number ERP105653.

Acknowledgements

We thank Matt Mayho, Jacqui Brown, and the sequencing teams at the Wellcome Trust

Sanger Institute for TraDIS sequencing and the Pathogen Informatics team for advice on the

analysis. We also thank Susannah Salter, Nick Croucher, George Salmond, Rita Monson,

Jose Bengoechea and Jin-Town Wang for supplying strains and plasmids. This work was

supported by Wellcome (grant 206194). M.J.D. is supported by a Wellcome Trust Sanger

Institute PhD Studentship. F.L.S. is supported by a Sir Henry Wellcome postdoctoral

fellowship (grant 106063/A/14/Z).

References

1.   Podschun R, Ullmann U. 1998. *Klebsiella* spp. as nosocomial pathogens:

Epidemiology, taxonomy, typing methods, and pathogenicity factors. Clin Microbiol

Rev 11:589–603.

2.   Holt KE, Wertheim H, Zadoks RN, Baker S, Whitehouse CA, Dance D, Jenney A,

Connor TR, Hsu LY, Severin J, Brisse S, Cao H, Wilksch J, Gorrie C, Schultz MB,

Edwards DJ, Nguyen K Van, Nguyen TV, Dao TT, Mensink M, Minh V Le, Nhu

NTK, Schultsz C, Kuntaman K, Newton PN, Moore CE, Strugnell RA, Thomson NR.

2015. Genomic analysis of diversity, population structure, virulence, and antimicrobial

resistance in *Klebsiella pneumoniae* , an urgent threat to public health. Proc Natl Acad

Sci 201501049.

3.   Broberg CA, Palacios M, Miller VL. 2014. *Klebsiella*: a long way to go towards

understanding this enigmatic jet-setter. F1000Prime Rep 6:64.

4.   Maroncle N, Rich C, Forestier C. 2006. The role of *Klebsiella pneumoniae* urease in

intestinal colonization and resistance to gastrointestinal stress. Res Microbiol 157:184–

193.

5.   Sahly H, Podschun R, Ullmann U. 2002. *Klebsiella* infections in the

immunocompromised host, p. 237–249. *In* Keisari, Y, Ofek, I (eds.), The biology and

pathology of innate immunity mechanisms. Springer US, Boston, MA.

6.   Shon AS, Bajwa RPS, Russo TA. 2014. Hypervirulent (hypermucoviscous) *Klebsiella*

pneumoniae. Virulence 4:107–118.

7.   Gomez-Simmonds A, Uhlemann A-C. 2017. Clinical implications of genomic

adaptation and evolution of carbapenem-resistant *Klebsiella pneumoniae*. J Infect Dis

215:S18–S27.

8.   Wang JH, Liu YC, Lee SS, Yen MY, Chen YS, Wang JH, Wann SR, Lin HH. 1998.

Primary liver abscess due to *Klebsiella pneumoniae* in Taiwan. Clin Infect Dis

26:1434–1438.

9.   Pomakova DK, Hsiao C-B, Beanan JM, Olson R, MacDonald U, Keynan Y, Russo

TA. 2012. Clinical and phenotypic differences between classic and hypervirulent

Klebsiella pneumonia: an emerging and under-recognized pathogenic variant. Eur J

Clin Microbiol Infect Dis 31:981–989.

10.   Cheng D-L, Liu Y-C, Yen M-Y, Liu C-Y, Wang R-S. 1991. Septic metastatic lesions

of pyogenic liver abscess: Their association with *Klebsiella pneumoniae* bacteremia in

diabetic patients. Arch Intern Med 151:1557–1559.

11.   Fang C-T, Lai S-Y, Yi W-C, Hsueh P-R, Liu K-L, Chang S-C. 2007. *Klebsiella*

pneumoniae genotype K1: An emerging pathogen that causes septic ocular or central

nervous system complications from pyogenic liver abscess. Clin Infect Dis 45:284–

293.

12.   Paczosa MK, Mecsas J. 2016. *Klebsiella pneumoniae*: Going on the offense with a

strong defense. Microbiol Mol Biol Rev 80:629–61.

13.   Struve C, Forestier C, Krogfelt KA. 2003. Application of a novel multi-screening

signature-tagged mutagenesis assay for identification of *Klebsiella pneumoniae* genes

essential in colonization and infection. Microbiology 149:167–176.

14.   Favre-Bonté S, Licht TR, Forestier C, Krogfelt KA. 1999. *Klebsiella pneumoniae*

capsule expression is necessary for colonization of large intestines of streptomycin-

treated mice. Infect Immun 67:6152–6156.

15.   Hsu C-R, Lin T-L, Chen Y-C, Chou H-C, Wang J-T. 2011. The role of *Klebsiella*

pneumoniae rmpA in capsular polysaccharide synthesis and virulence revisited.

Microbiology 157:3446–3457.

16.   Cheng HY, Chen YS, Wu CY, Chang HY, Lai YC, Peng HL. 2010. RmpA regulation

of capsular polysaccharide biosynthesis in *Klebsiella pneumoniae* CG43. J Bacteriol

192:3144–3158.

17.   Wyres KL, Wick RR, Gorrie C, Jenney A, Follador R, Thomson NR, Holt KE. 2016.

Identification of *Klebsiella* capsule synthesis loci from whole genome data. Microb

Genomics 2:e000102.

18.   Cortés G, Borrell N, de Astorza B, Gómez C, Sauleda J, Albertí S. 2002. Molecular

analysis of the contribution of the capsular polysaccharide and the lipopolysaccharide

O side chain to the virulence of *Klebsiella pneumoniae* in a murine model of

pneumonia. Infect Immun 70:2583–2590.

19.   Patel PK, Russo TA, Karchmer AW. 2014. Hypervirulent *Klebsiella pneumoniae*.

Open Forum Infect Dis 1:ofu028-ofu028.

20.   Nassif X, Sansonetti PJ. 1986. Correlation of the virulence of *Klebsiella pneumoniae*

K1 and K2 with the presence of a plasmid encoding aerobactin. Infect Immun 54:603–

608.

21.   Chen Y-TT, Chang H-YY, Lai Y-CC, Pan C-CC, Tsai S-FF, Peng H-LL. 2004.

Sequencing and analysis of the large virulence plasmid pLVPK of *Klebsiella*

pneumoniae CG43. Gene 337:189–198.

22.   Mizuta K, Ohta M, Mori M, Hasegawa T, Nakashima I, Kato N. 1983. Virulence for

mice of *Klebsiella* strains belonging to the O1 group: relationship to their capsular (K)

types. Infect Immun 40:56–61.

23.   Nassif X, Fournier JM, Arondel J, Sansonetti PJ. 1989. Mucoid phenotype of

Klebsiella pneumoniae is a plasmid-encoded virulence factor. Infect Immun 57:546–

552.

24.   Yeh K-M, Kurup A, Siu LK, Koh YL, Fung C-P, Lin J-C, Chen T-L, Chang F-Y, Koh

T-H. 2007. Capsular serotype K1 or K2, rather than *magA* and *rmpA*, is a major

virulence determinant for *Klebsiella pneumoniae* liver abscess in Singapore and

Taiwan. J Clin Microbiol 45:466–471.

25.   Russo TA, Shon AS, Beanan JM, Olson R, MacDonald U, Pomakov AO, Visitacion

MP. 2011. Hypervirulent *K. pneumoniae* secretes more and more active iron-

acquisition molecules than "classical" *K. pneumoniae* thereby enhancing its virulence.

PLoS One 6:e26734.

26.   Yeh K-M, Lin J-C, Yin F-Y, Fung C-P, Hung H-C, Siu L-K, Chang F-Y. 2010.

Revisiting the importance of virulence determinant *magA* and its surrounding genes in

Klebsiella pneumoniae causing pyogenic liver abscesses: Exact role in serotype K1

capsule formation. J Infect Dis 201:1259–1267.

27.   Fang C-T, Lai S-Y, Yi W-C, Hsueh P-R, Liu K-L. 2010. The function of *wzy_K1*

(*magA*), the serotype K1 polymerase gene in *Klebsiella pneumoniae cps* gene cluster. J

Infect Dis 201:1268–1269.

28.   Fang C-T, Chuang Y-P, Shun C-T, Chang S-C, Wang J-T. 2004. A novel virulence

gene in *Klebsiella pneumoniae* strains causing primary liver abscess and septic

metastatic complications. J Exp Med 199:697–705.

29.   Lai Y-C, Peng H-L, Chang H-Y. 2003. RmpA2, an activator of capsule biosynthesis in

Klebsiella pneumoniae CG43, regulates K2 *cps* gene expression at the transcriptional

level. J Bacteriol 185:788–800.

30.   Yu W-L, Ko W-C, Cheng K-C, Lee C-C, Lai C-C, Chuang Y-C. 2008. Comparison of

prevalence of virulence factors for *Klebsiella pneumoniae* liver abscesses between

isolates with capsular K1/K2 and non-K1/K2 serotypes. Diagn Microbiol Infect Dis

62:1–6.

31.   Kobayashi SD, Porter AR, Freedman B, Pandey R, Chen L, Kreiswirth BN, Deleo R.

2018. Antibody-mediated killing of carbapenem-resistant ST258 *Klebsiella*

pneumoniae by human neutrophils. mBio 9:e00297-18.

32.   Diago-Navarro E, Motley M, Ruiz-Perez G, Yu W, Austin J, Seco B, Xiao G,

Chikhalya A, Seeberger PH, Fries BC. 2018. Novel, broadly reactive anticapsular

antibodies against carbapenem-resistant *Klebsiella pneumoniae* protect from infection.

mBio 9:e00091-18.

33.   Lin TL, Hsieh PF, Huang YT, Lee WC, Tsai YT, Su PA, Pan YJ, Hsu CR, Wu MC,

Wang JT. 2014. Isolation of a bacteriophage and its depolymerase specific for K1

capsule of *Klebsiella pneumoniae*: Implication in typing and treatment. J Infect Dis

210:1734–1744.

34.   Pan YJ, Lin TL, Lin YT, Su PA, Chen CT, Hsieh PF, Hsu CR, Chen CC, Hsieh YC,

Wang JT. 2015. Identification of capsular types in carbapenem-resistant *Klebsiella*

pneumoniae strains by *wzc* sequencing and implications for capsule depolymerase

treatment. Antimicrob Agents Chemother 59:1038–1047.

35.   Hsieh PF, Lin HH, Lin TL, Chen YY, Wang JT. 2017. Two T7-like bacteriophages,

K5-2 and K5-4, each encodes two capsule depolymerases: Isolation and functional

characterization. Sci Rep 7:1–13.

36.   Navon-Venezia S, Kondratyeva K, Carattoli A. 2017. *Klebsiella pneumoniae*: A major

worldwide source and shuttle for antibiotic resistance. FEMS Microbiol Rev 41:252–

275.

37. Yu W-L, Ko W-C, Cheng K-C, Lee H-C, Ke D-S, Lee C-C, Fung C-P, Chuang Y-C.

2006. Association between *rmpA* and *magA* genes and clinical syndromes caused by

Klebsiella pneumoniae in Taiwan. Clin Infect Dis 42:1351–1358.

38. Bachman MA, Breen P, Deornellas V, Mu Q, Zhao L, Wu W, Cavalcoli JD, Mobley

HLT. 2015. Genome-wide identification of *Klebsiella pneumoniae* fitness genes

during lung infection. mBio 6:1–9.

39. Gottesman S, Trisler P, Torres-Cabassa A. 1985. Regulation of capsular

polysaccharide synthesis in *Escherichia coli* K-12: characterization of three regulatory

genes. J Bacteriol 162:1111–1119.

40. Stout V, Torres-Cabassa A, Maurizi MR, Gutnick D, Gottesman S. 1991. RcsA, an

unstable positive regulator of capsular polysaccharide synthesis. J Bacteriol 173:1738–

1747.

41. McCallum KL, Whitfield C. 1991. The *rcsA* gene of *Klebsiella pneumoniae* O1:K20 is

involved in expression of the serotype-specific K (capsular) antigen. Infect Immun

59:494–502.

42. Lin CT, Wu CC, Chen YS, Lai YC, Chi C, Lin JC, Chen Y, Peng HL. 2011. Fur

regulation of the capsular polysaccharide biosynthesis and iron-acquisition systems in

Klebsiella pneumoniae CG43. Microbiology 157:419–429.

43. Barquist L, Mayho M, Cummins C, Cain AK, Boinett CJ, Page AJ, Langridge GC,

Quail MA, Keane JA, Parkhill J. 2016. The TraDIS toolkit: Sequencing and analysis

for dense transposon mutant libraries. Bioinformatics 32:1109–1111.

44. Langridge GC, Phan MD, Turner DJ, Perkins TT, Parts L, Haase J, Charles I, Maskell

DJ, Peters SE, Dougan G, Wain J, Parkhill J, Turner KA. 2009. Simultaneous assay of

every *Salmonella* Typhi gene using one million transposon mutants. Genome Res

19:2308–2316.

45.   Patrick S, Reid JH. 1983. Separation of capsulate and non-capsulate *Bacteroides*

fragilis on a discontinuous density gradient. J Med Microbiol 16:239–241.

46.   Menck K, Behme D, Pantke M, Reiling N, Binder C, Pukrop T, Klemm F. 2014.

Isolation of human monocytes by double gradient centrifugation and their

differentiation to macrophages in Teflon-coated cell culture bags. J Vis Exp 1–10.

47.   Brunner J, Scheres N, El Idrissi NB, Deng DM, Laine ML, van Winkelhoff AJ,

Crielaard W. 2010. The capsule of *Porphyromonas gingivalis* reduces the immune

response of human gingival fibroblasts. BMC Microbiol 10:5.

48.   Hassan KA, Cain AK, Huang T, Liu Q, Elbourne LDH, Boinett CJ, Brzoska AJ, Li L,

Ostrowski M, Nhu NTK, Nhu TDH, Baker S, Parkhill J, Paulsen IT, Thi N, Nhu K, Do

T, Nhu H, Baker S, Parkhill J, Paulsen IT. 2016. Fluorescence-based flow sorting in

parallel with transposon insertion site sequencing identifies multidrug efflux systems

in *Acinetobacter baumannii*. mBio 7:e01200-16.

49.   Alexa A, Rahnenfuhrer J. 2016. topGO: Enrichment analysis for gene ontology. R

package version 2.32.0.

50.   Grossmann S, Bauer S, Robinson PN, Vingron M. 2007. Improved detection of

overrepresentation of Gene-Ontology annotations with parent-child analysis.

Bioinformatics 23:3024–3031.

51.   Malinverni JC, Silhavy TJ. 2009. An ABC transport system that maintains lipid

asymmetry in the Gram-negative outer membrane. Proc Natl Acad Sci 106:8009–

8014.

52.   Cho B, Federowicz S, Park Y, Kim D, Palsson BØ. 2013. Deciphering the regulatory

logic of a stimulon. Nat Chem Biol 8:65–71.

53.   Ares MA, Fernández-Vázquez JL, Rosales-Reyes R, Jarillo-Quijada MD, von Bargen

K, Torres J, González-y-Merchand JA, Alcántar-Curiel MD, De la Cruz MA. 2016. H-

NS nucleoid protein controls virulence features of *Klebsiella pneumoniae* by

regulating the expression of type 3 pili and the capsule polysaccharide. Front Cell

Infect Microbiol 6:13.

54.   Corbett D, Bennett HJ, Askar H, Green J, Roberts IS. 2007. SlyA and H-NS regulate

transcription of the *Escherichia coli* K5 capsule gene cluster, and expression of *slyA* in

Escherichia coli is temperature-dependent, positively autoregulated, and independent

of H-NS. J Biol Chem 282:33326–33335.

55.   Goh KGK, Phan M-D, Forde BM, Chong TM, Yin W-F, Chan K-G, Ulett GC, Sweet

MJ, Beatson SA, Schembri MA. 2017. Genome-wide discovery of genes required for

capsule production by uropathogenic *Escherichia coli*. mBio 8:1–16.

56.   Palacios M, Miner TA, Frederick DR, Sepulveda VE, Quinn JD, Walker KA, Miller

VL. 2018. Identification of two regulators of virulence that are conserved in *Klebsiella*

pneumoniae classical and hypervirulent strains. mBio 9:e01443-18.

57.   Gruenheid S, Le Moual H. 2012. Resistance to antimicrobial peptides in Gram-

negative bacteria. FEMS Microbiol Lett 330:81–89.

58.   Sugiyama Y, Nakamura A, Matsumoto M, Kanbe A, Sakanaka M, Higashi K, Igarashi

K, Katayama T, Suzuki H, Kurihara S. 2016. A novel putrescine exporter SapBCDF of

Escherichia coli. J Biol Chem 291:26343–26351.

59.   Insua JL, Llobet E, Moranta D, Pérez-Gutiérrez C, Tomás A, Garmendia J,

Bengoechea JA. 2013. Modeling *Klebsiella pneumoniae* pathogenesis by infection of

the wax moth *Galleria mellonella*. Infect Immun 81:3552–3565.

60.   Wall E, Majdalani N, Gottesman S. 2018. The complex Rcs regulatory cascade 111–

812   139.

813   61.   Majdalani N, Gottesman S. 2005. The Rcs phosphorelay: A complex signal

814   transduction system. Annu Rev Microbiol 59:379–405.

815   62.   Fowler CC, Galán JE. 2018. Decoding a *Salmonella* Typhi regulatory network that

816   controls typhoid toxin expression within human cells. Cell Host Microbe 23:65–76.e6.

817   63.   Castelli ME, Véscovi EG. 2011. The Rcs signal transduction pathway is triggered by

818   enterobacterial common antigen structure alterations in *Serratia marcescens*. J

819   Bacteriol 193:63–74.

820   64.   Gozdziewicz TK, Lugowski C, Lukasiewicz J. 2014. First evidence for a covalent

821   linkage between enterobacterial common antigen and lipopolysaccharide in *Shigella*

822   *sonnei* phase II ECALPS. J Biol Chem 289:2745–2754.

823   65.   Mills G, Dumigan A, Kidd T, Hobley L, Bengoechea JA. 2017. Identification and

824   characterization of *Klebsiella pneumoniae* two *lpxL* lipid A late acyltransferases and

825   their role in virulence. Infect Immun 85:IAI.00068-17.

826   66.   Llobet E, Martínez-Moliner V, Moranta D, Dahlström KM, Regueiro V, Tomás A,

827   Cano V, Pérez-Gutiérrez C, Frank CG, Fernández-Carrasco H, Insua JL, Salminen TA,

828   Garmendia J, Bengoechea JA. 2015. Deciphering tissue-induced *Klebsiella*

829   *pneumoniae* lipid A structure. Proc Natl Acad Sci 112:E6369–E6378.

830   67.   Fresno S, Jiménez N, Izquierdo L, Merino S, Corsaro MM, De Castro C, Parrilli M,

831   Naldi T, Requé M, Tomás JM, Fresno S, Jime N, Corsaro MM, Castro C De, Parrilli

832   M, Toma JM, Naldi T, Regue M, Toma JM, Ii F, Universitario C. 2006. The ionic

833   interaction of *Klebsiella pneumoniae* K2 capsule and core lipopolysaccharide.

834   Microbiology 152:1807–1818.

68.   Fresno S, Jiménez N, Canals R, Merino S, Corsaro MM, Lanzetta R, Parrilli M,

Pieretti G, Regué M, Tomás JM. 2007. A second galacturonic acid transferase is

required for core lipopoylsaccharide biosynthesis and complete capsule association

with the cell surface in *Klebsiella pneumoniae*. J Bacteriol 189:1128–1137.

69.   Olaitan AO, Morand S, Rolain JM. 2014. Mechanisms of polymyxin resistance:

Acquired and intrinsic resistance in bacteria. Front Microbiol 5:1–18.

70.   Yan A, Guan Z, Raetz CRH. 2007. An undecaprenyl phosphate-aminoarabinose

flippase required for polymyxin resistance in *Escherichia coli*. J Biol Chem

282:36077–36089.

71.   Breazeale SD, Ribeiro AA, McClerren AL, Raetz CRH. 2005. A formyltransferase

required for polymyxin resistance in *Escherichia coli* and the modification of lipid A

with 4-amino-4-deoxy-L-arabinose: Identification and function of UDP-4-deoxy-4-

formamido-L-arabinose. J Biol Chem 280:14154–14167.

72.   Mason KM, Raffel FK, Ray WC, Bakaletz LO. 2011. Heme utilization by nontypeable

Haemophilus influenzae is essential and dependent on Sap transporter function. J

Bacteriol 193:2527–2535.

73.   Harms C, Domoto Y, Celik C, Rahe E, Stumpe S, Schmid R, Nakamura T, Bakker EP.

2001. Identification of the ABC protein SapD as the subunit that confers ATP

dependence to the K^+-uptake systems TrkH and TrKG from *Escherichia coli* K-12.

Microbiology 147:2991–3003.

74.   Monson R, Smith DS, Matilla MA, Roberts K, Richardson E, Drew A, Williamson N,

Ramsay J, Welch M, Salmond GPC. 2015. A plasmid-transposon hybrid mutagenesis

system effective in a broad range of Enterobacteria. Front Microbiol 6:1–13.

75.   Fineran PC, Everson L, Slater H, Salmond GPC. 2005. A GntR family transcriptional

regulator (PigT) controls gluconate-mediated repression and defines a new,

independent pathway for regulation of the tripyrrole antibiotic, prodigiosin, in

Serratia. Microbiology 151:3833–3845.

76.   Lawlor MS, Hsu J, Rick PD, Miller VL. 2005. Identification of *Klebsiella pneumoniae*

virulence determinants using an intranasal infection model. Mol Microbiol 58:1054–

1073.

77.   Érika A, Gomes I, Stuchi LP, Maria N, Siqueira G, Henrique JB, Vicentini R, Ribeiro

ML, Darrieux M. 2018. Selection and validation of reference genes for gene

expression studies in *Klebsiella pneumoniae* using reverse transcription quantitative

real-time PCR. Sci Rep 9001:s1–14.

 Figure legends

Figure 1 Summary of density-dependent TraDISort strategy. (A) Electron microscope image

of capsulated *K. pneumoniae* NTUH-K2044. (B) Schematic of the TraDISort strategy to

identify capsule regulators. A high-density transposon library is applied to the top of a

discontinuous Percoll gradient, which is then centrifuged at moderate speed to separate

capsulated and non-capsulated mutants. The separate fractions are sequenced to identify

transposon-gDNA junctions. (C) Validation of the Percoll gradient method for separating

cells by capsule phenotype. Individual fractions immediately following separation on a

Percoll gradient were adjusted to $OD_{600} = 4$ in sterile PBS, and assayed for uronic acid

content. Statistical significance was evaluated by one-way ANOVA followed by Tukey's

Honest Significant Differences (HSD) test, and is reported for each fraction relative to the

input sample (** $p < 0.01$; *** $p < 0.001$). (D) Application of density separation method to *S.*

pneumoniae strain 23F.

Figure 2 (A) Results of TraDIS mapping at the capsule locus of *K. pneumoniae* NTUH-

K2044. Transposon insertions in capsule genes were very abundant in the input sample.

These majority of these mutants were not found in the top fraction but were instead enriched

in the middle fraction (e.g., *gnd*, *galF*) or the bottom fraction (all genes from *wcaJ* to *wzi*).

Genes defined as hits in our screen are shown in yellow, others in grey. (B) Insertion mutants

in genes of the *sap* ABC transporter locus were not found in the middle fraction, suggesting

higher capsule production than wild-type. Genes which are putative increased capsule hits are

shown in blue, with others in grey. Note that *sapD* had a very low number of reads in all

fractions. The flanking genes, *codB* and KP1_2336, are examples of genes where transposon

insertion does not change capsule, and the insertion mutants are found in both the top and the

middle fractions. (C) Common and strain-specific genes required for full capsule production

in *K. pneumoniae* NTUH-K2044 and ATCC43816. Putative increased capsule mutants are

not considered in this chart. These two strains share approximately 4,200 genes, and the

majority of genes required for full capsule production are present in both strains. Sixteen

genes were defined as capsule-related in both strains. Note that the gene content of the *cps*

locus differs between these two strains, which accounts for the majority of hits in strain-

specific genes.

Figure 3 Validation of putative capsule regulators with single-gene deletion mutants. (A)

Percoll gradient centrifugation of clean deletion mutants in selected NTUH-K2044 genes. All

of the genes tested showed reduced density compared to wild-type, with the exception of the

putative increased capsule mutant, Δ*sapBDEF*, which stayed above the 15% Percoll layer.

(B) Validation of ATCC43816 deletion mutant phenotypes on 35%-50% Percoll gradients.

The Δ*arnF* and Δ*rfaH* mutants showed reduced density compared to wild-type, while the

Δ*mprA* mutant did not, in contrast to its phenotype in NTUH-K2044. (C) Hypermucoidy tests

with Kpn NTUH-2044 mutants. Strains were grown to late stationary phase and cultures

centrifuged for 5 min at 1,000 x *g*. The OD_{600} of the supernatant was measured and is

presented here as a proportion of the starting OD_{600}. *** $p < 0.001$, One-way ANOVA

followed by Tukey's HSD test, relative to wild-type.

Figure 4 Complementation and electron microscopy of *K. pneumoniae* NTUH-K2044

Δ*argR*, Δ*mprA*, Δ*slyA* and Δ*sapBCDF* mutants. (A) Hypermucoidy assay. Strains were

centrifuged at 1,000 x *g* for 5 min to define decreased hypermucoidy relative to wild type, or

2,500 x *g* to identify increases in hypermucoidy relative to wild type. Significant differences

are indicated by ** $p < 0.01$ or *** $p < 0.001$, one-way ANOVA and Tukey's HSD test. This

shows an independent experiment to Figure 3C. (B) Uronic acid assay to confirm the capsule

phenotype of each strain. Differences relative to wild type were evaluated by pairwise one-

way ANOVA with Benjamini-Hochberg correction for multiple testing * $p < 0.05$, ** $p <$

0.001, *** $p < 0.0001$. (C) Transmission electron microscopy images of *K. pneumoniae*

NTUH-K2044 and its Δ*argR*, Δ*mprA*, Δ*slyA* and Δ*sapBCDF* mutants.

Figure 5. Virulence of selected mutants. (A) Killing of research-grade *Galleria mellonella*

larvae by infection with *K. pneumoniae* NTUH-K2044 wild type or mutant strains. Larvae

were infected at an inoculum of 10^5. Differences in killing compared to wild type were

evaluated using the Kaplan-Meier log rank test and are indicated by * $p < 0.05$, ** $p < 0.01$ or

*** $p < 0.001$. (B) Survival in human serum. Differences relative to wild type are indicated

by ** $p < 0.01$, *** $p < 0.001$, pairwise one-way ANOVA. (C) Expression of several

capsule-related genes in NTUH-K2044 Δ*sapBCDF*. Transcript abundance was measured

using the relative standard curve method with *recA* as a reference gene, and normalised to

WT. * $p < 0.05$, *** $p < 0.001$, one-way ANOVA.

Figure 6. Overview of capsule regulation in NTUH-K2044. Products are coloured red for

mutants with low capsule, blue for mutants with high capsule, and those genes that were

validated in clean deletion knockouts are indicated with bold labels and outlines. Likely

modes of action are indicated by green or red arrows for predicted positive and negative

effects on transcription of the capsule locus. Grey arrows indicate inputs that may affect

capsule synthesis without modulating transcription. Omitted are individual capsule

biosynthetic genes, ECA biosynthetic genes, and components of the transcription and

translation machinery.

Tables

Table 1. Genes in which mutants have altered capsule production. A list of all

statistically-significant genes from this TraDIS screen which, when disrupted by transposon

increase capsule production in *K. pneumoniae*. Cutoff criteria are described in

Methods. Gene names and functional annotations are taken from the pan-genome consensus

file (see Materials and Methods). The complete dataset, including statistical data, is provided

in Tables S3 and S4.

| | | | Locus tag** | |
| | | | NTUH- | |
Gene	Annotation	Proportion of strains	K2044	ATCC43816
wzxE	WzxE protein	0.94	KP1_0154	VK055_3183
wzy	ECA polymerase	1.00	KP1_0156	VK055_3181
rfaH	transcriptional activator RfaH	1.00	KP1_0199	VK055_3141
seqA	replication initiation regulator SeqA	0.99	KP1_1663	VK055_1821
pgm	phosphoglucomutase	1.00	KP1_1664	VK055_1820
gnd	6-phosphogluconate dehydrogenase	0.69	KP1_3704	VK055_5026
wcaJ	colanic acid biosynthsis UDP-glucose lipid carrier transferase WcaJ	0.20	KP1_3705	VK055_5025
wza	polysaccharide export protein	0.98	KP1_3720	VK055_5015
wzi	outer membrane protein	0.99	KP1_3721	VK055_5014
orf2	PAP2 family protein	0.93	KP1_3725	VK055_5013
mlaD	ABC transporter	1.00	KP1_4915	VK055_3874
arnF	4-amino-4-deoxy-L-arabinose-phosphoundecaprenol flippase subunit ArnF	1.00	KP1_5178	VK055_3630
arnE	SMR family multidrug resistance protein	0.95	KP1_5179	VK055_3628
arnD	Polymyxin resistance protein PmrJ	1.00	KP1_5181	VK055_3626
gor	glutathione reductase	1.00	KP1_5206	VK055_3604
wabN	deacetylase	1.00	KP1_5319	VK055_3502
wecA	Undecaprenyl-phosphate N-acetylglucosaminyl 1-phosphate transferase	0.98	KP1_0146	*VK055_3191*
yjeA	translation elongation factor P Lys34:lysine transferase	1.00	KP1_0426	*VK055_2911*
miaA	tRNA delta(2)-isopentenylpyrophosphate transferase	1.00	KP1_0439	*VK055_2895*
dksA	RNA polymerase-binding transcription factor	0.99	KP1_0973	*VK055_2423*
htpG	chaperone protein HtpG	0.97	KP1_1331	*VK055_2094*
manA	mannose-6-phosphate isomerase	0.65	KP1_2524	*VK055_0991*
ydgI	Arginine/ornithine antiporter ArcD	1.00	KP1_2534	*VK055_0982*
rnfA	electron transport complex protein RnfA	0.98	KP1_3036	*VK055_0514*
rnfC_2	electron transport complex protein RnfC	0.38	KP1_3038	*VK055_0512*
nqrB_1 (rnfD)	electron transport complex protein RnfD	0.99	KP1_3039	*VK055_0511*
rnfE	electron transport complex protein RnfE	0.78	KP1_3041	*VK055_0509*
slyA_1	transcriptional regulator SlyA	1.00	KP1_3054	*VK055_0496*
lpp	major outer membrane lipoprotein	1.00	KP1_3230	*VK055_0326*
galU	UTP-glucose-1-phosphate uridylyltransferase	0.97	KP1_3315	*VK055_0250*
uvrY	BarA-associated response regulator UvrY (GacA, SirA)	1.00	KP1_3542	*VK055_0032*
rmpA	regulator of mucoid phenotype	0.05	KP1_3619	*VK055_5097*

Low capsule mutants

Gene	Annotation			
manC	mannose-1-phosphate guanylyltransferase	0.24	KP1_3703	*VK055_5027*
galF	UDP-glucose pyrophosphorylase	0.88	KP1_3726	*VK055_5012*
rcsB	DNA-binding capsular synthesis response regulator RcsB	0.77	KP1_3872	*VK055_4883*
glnB	nitrogen regulatory protein P-II	1.00	KP1_4132	*VK055_4623*
rluD	23S rRNA pseudouridine synthase D	0.96	KP1_4172	*VK055_4583*
emrR	transcription repressor	1.00	KP1_4277	*VK055_4504*
barA	BarA sensory histidine kinase	0.92	KP1_4400	*VK055_4386*
greA	transcription elongation factor GreA	0.99	KP1_4900	*VK055_3886*
arcB	Aerobic respiration control sensor protein arcB	1.00	KP1_4931	*VK055_3858*
argR	Arginine pathway regulatory protein ArgR	1.00	KP1_4961	*VK055_3832*
envZ_2	Osmolarity sensory histidine kinase EnvZ	0.90	KP1_5105	*VK055_3697*
ompR	osmolarity response regulator	1.00	KP1_5106	*VK055_3696*
rfaZ	lipopolysaccharide core biosynthesis protein RfaZ	0.76	KP1_5316	*VK055_3505*
waaL	O-antigen polymerase	0.78	KP1_5317	*VK055_3504*
rfaQ (waaQ)	lipopolysaccharide heptosyltransferase III	1.00	KP1_5320	*VK055_3501*
wabH	glycosyltransferase	1.00	KP1_5322	*VK055_3499*
yjeK	EF-P beta-lysylation protein EpmB	NA	KP1_0415	NA
wcaI	putative glycosyl transferase	0.11	KP1_3706	NA
gmm (wcaH)	GDP-mannose mannosyl hydrolase	0.05	KP1_3708	NA
wcaG	GDP-fucose synthetase	0.11	KP1_3709	NA
gmd	GDP-mannose 4,6-dehydratase	0.13	KP1_3711	NA
KP1_3712	galactoside O-acetyltransferase	0.00	KP1_3712	NA
group_19979	glycosyltransferase	0.06	KP1_3713	NA
magA	mucoviscosity-associated protein	0.06	KP1_3714	NA
KP1_3715	polysaccharide pyruvyl transferase	0.05	KP1_3715	NA
wzx	repeat unit exporter	0.06	KP1_3716	NA
wzc	tyrosine-protein kinase Wzc	0.06	KP1_3718	NA
wzb	putative protein tyrosine phosphatase	0.00	KP1_3719	NA
*csrB**	carbon storage regulatory sRNA	NA	KP1_6106	NA
*rmpA_2**	regulator of mucoid phenotype	NA	KP1_p020	NA
pgi	glucose-6-phosphate isomerase	1.00	*KP1_0264*	VK055_3061
mioC_2	flavodoxin	0.99	*KP1_0001*	VK055_3326
glpD	Aerobic glycerol-3-phosphate dehydrogenase	0.99	*KP1_5126*	VK055_3679
yrbF	putative ABC transporter ATP-binding protein YrbF	1.00	*KP1_4917*	VK055_3872
mlaE (yrbE)	ABC transporter	1.00	*KP1_4916*	VK055_3873
mlaC (yrbC)	ABC transporter	1.00	*KP1_4914*	VK055_3875
mlaA	lipoprotein	1.00	*KP1_3977*	VK055_4786
wzb	putative acid phosphatase Wzb	0.01	NA	VK055_5016
wzc (etk)	tyrosine autokinase	0.08	NA	VK055_5017
mshA	group 1 glycosyl transferase	0.08	NA	VK055_5018
orf8	group 1 glycosyl transferase	0.08	NA	VK055_5019
VK055_5020	group 1 glycosyl transferase	0.08	NA	VK055_5020
VK055_5021	Lipid A core - O-antigen ligase and related enzymes	0.08	NA	VK055_5021
wzxC	colanic acid exporter	0.08	NA	VK055_5022
orf12	hypothetical protein	0.08	NA	VK055_5023
orf13	putative lipopolysaccharide biosynthesis O-acetyl transferase WbbJ	0.06	NA	VK055_5024

High-capsule mutants

Gene	Annotation	Proportion of strains	Locus tag NTUH-K2044
polA	DNA polymerase I	0.79	KP1_0024
cyaA	adenylate cyclase	0.99	KP1_0164
trkH	potassium transport protein	1.00	KP1_0206
purA	adenylosuccinate synthetase	1.00	KP1_0448

apaH	diadenosinetetraphosphatase	0.96	KP1_0859
ace	pyruvate dehydrogenase E1 component	0.98	KP1_0941
glnD	PII uridylyl-transferase	0.94	KP1_1019
hha	haemolysin expression modulating protein	0.99	KP1_1317
tolR	putative inner membrane protein involved in the tonB-independent uptake of group A colicins	1.00	KP1_1700
tolB	translocation protein TolB precursor	1.00	KP1_1702
mdoG	periplasmic glucans biosynthesis protein	1.00	KP1_2050
mdoH	glucosyltransferase	0.93	KP1_2051
sapF	ABC-type peptide transport system ATP-binding component	0.97	KP1_2331
sapC	ABC-type peptide transport system permease component	1.00	KP1_2333
sapB	ABC-type peptide transport system permease component	0.99	KP1_2334
sapA	ABC-type peptide transport system periplasmic component	0.68	KP1_2335
pykF	pyruvate kinase	0.99	KP1_3229
hns	DNA-binding protein HLP-II/pleiotropic regulator	0.99	KP1_3314
prc	carboxy-terminal protease for penicillin-binding protein 3	0.11	KP1_3473
ackA	acetate/propionate kinase	1.00	KP1_3933
pta	phosphate acetyltransferase	0.80	KP1_3934
smpB	SsrA tmRNA-binding protein	1.00	KP1_4198
ptsN	sugar-specific PTS family enzyme IIA component	1.00	KP1_4926
KP1_4976 *(csrD)*	csrB regulatory protein csrD	0.95	KP1_4976
fis	DNA-binding protein	1.00	KP1_4989
pitA	putative low-affinity inorganic phosphate transporter	0.84	KP1_5198

953
*Note that the pan-genome only includes protein-coding sequences located on the chromosome, so the small RNA *csrB* and
plasmid-encoded *rmpA2* are not included in our pan-genome analysis
**Locus tags are shown in black non-italic for hits, and grey italics where the gene was not called as a hit

Supplementary material legends

Figure S1. Validation of density gradient centrifugation for capsule-based separations. (A) *K.*

pneumoniae NTUH-K2044 (K1), *K. pneumoniae* ATCC43816 (K2) and *E. coli* DH5α grown

at 37 °C each localised to a different location on a 15%-35%-50% Percoll gradient, and the

two *Klebsiella* strains migrated further down the gradient when grown at 25 °C. Locations

equivalent to the Top, Middle and Bottom fractions used in (B) are indicated by red letters

'T', 'M' and 'B'. A non-hypermucoid capsule type K106 strain, *K. pneumoniae* RH201207,

was distributed throughout the 35% Percoll layer following centrifugation. (B) Viable counts

of gradient fractions of wild type *K. pneumoniae* NTUH-K2044 and *K. pneumoniae*

ATCC43816. The NTUH-K2044 strain localised primarily above the 15% Percoll layer, with

a smaller proportion of cells in the middle fraction (4% of the viable count of the top

fraction). The viable count of the bottom fraction was close to the detection limit (indicated

by a dashed line) at 0.0005% of that of the top fraction. The ATCC43816 strain migrated

almost solely to the 15%-35% interface of the gradient, with 0.01% of cells located in the

bottom fraction in this experiment. *** $p < 0.001$, one-way ANOVA followed by Tukey's

HSD performed on log-transformed viable counts. (C) Hypermucoidy test of individual

fractions of the NTUH-K2044 TraDIS library following separation. (D) Heritability of cell

density phenotype following overnight growth of fractions. The "middle" fraction had a

partially heritable phenotype and was distributed across the previous locations of the "top"

and "middle" fractions. The top and bottom fractions localised to the same place following

overnight growth.

Figure S2. (A) Relationship between sequencing depth and number of insertion sites in the

Klebsiella pneumoniae NTUH-K2044 library, sequenced input sample 1. The plot was

generated using the **seq_saturation_test.py** script available at https://github.com/francesca-

short/tradis_scripts. (B) Gene-wise insertion index values along the chromosome of *K.*

pneumoniae NTUH-K2044. (C) Density-TraDISort results for NTUH-K2044 genes with

known capsule phenotypes. Mutants with unchanged capsule production were located in the

top and middle fractions, as shown for *fadB*. Severe capsule reduction resulted in migration

to the bottom fraction, shown for *rfaH*, while mutants in genes required for the hypermucoidy

phenotype (*rcsB* and *rmpA*) were located primarily in the middle fraction. Genes where

mutation results in increased capsule were evident as those that were absent from the middle

fraction, but enriched in the top fraction, as shown for *hns*.

Figure S3. TraDIS analysis of capsule regulation in *Klebsiella pneumoniae* ATCC43816 (A)

Electron microscope image of ATCC43816. (B) Uronic acid assay to validate density-based

separation for this strain and compare its capsule production to NTUH-K2044. ** $p < 0.01$,

one-way ANOVA and Tukey's HSD. (C) Relationship between sequencing depth and

number of unique insertion sites identified in the *Klebsiella pneumoniae* ATCC43816 mutant

library. (D) Distribution of insertion sites across the chromosome. (E) TraDIS plot files at the

capsule locus of *K. pneumoniae* ATCC43816; almost all mutants were found in the bottom

fraction. Genes called as capsule-regulatory hits are shown in yellow, those not called as hits

are in grey.

1002

Figure S4. (A) Mutation of *waaL* does not further reduce density in a *wza* mutant strain. The

indicated mutants of NTUH-K2044 were centrifuged on a 70% Percoll layer, which was

determined to be the concentration required to retain the *wza* mutant. (B) Uronic acid assay

with mutants of NTUH-K2044. These data are from the same experiment as that shown in

Figure 4B. Differences relative to wild type were evaluated by pairwise one-way ANOVA

with Benjamini-Hochberg correction for multiple testing (* $p < 0.05$, ** $p < 0.01$ or *** $p <$

0.001).

1010

1011   **Supplementary Text S1.** Extended methods, supplementary references, and random-prime

1012   PCR identification of transposon insertion sites in acapsular clones.

1013

1014   **Supplementary Table S1**. Strains, plasmids, and oligonucleotides used in this study**.**

1015

1016   **Supplementary Table S2.** Summary statistics and reference numbers for TraDIS samples

1017   sequenced in this study.

1018

1019   **Supplementary Table S3.** Raw comparison data for *K. pneumoniae* ATCC43816 and

1020   NTUH-K2044 capsule gradient fractions.

1021

1022   **Supplementary Table S4.** Combined capsule hits in *K. pneumoniae* ATCC43816 and

1023   NTUH-K2044 with pan-genome information and references to relevant literature.

1024

1025   **Supplementary Table S5.** List of genomes used to generate the *K. pneumoniae* pan-genome.

1026

1027

1028

(A)

NTUH-K2044 (K1)

(B)

Overnight culture of TraDIS library → "Input"; DNA isolation and transposon sequencing

% Percoll [15 / 35 / 50] —Centrifuge→ "Top" / "Middle" / "Bottom" → DNA isolation and transposon sequencing

(C)

Uronic acid per 10^9 cells (µg)

Input, Top, Middle (**), Bottom (***)

(D)

% Percoll [30% / 50% / 80%]   WT   Δcps   WT + Δcps

S. pneumoniae 23F

(A) Kpn NTUH-K2044

% Percoll

15%
35%
50%

WT ΔslyA ΔuvrY ΔarcB

ΔargR ΔmprA ΔompR ΔrfaH

ΔwaaL i-wza i-wza - ΔwaaL

WT ΔsapBCDF

(B) Kpn ATCC43816

% Percoll

35%
50%

WT ΔarnF ΔrfaH ΔmprA

(C)

(A)

1,000 x g, 5 min (left panel)

OD_{600} (supernatant) / OD_{600} (culture)

WT, ΔargR ***, ΔargR comp, ΔmprA ***, ΔmprA comp, ΔslyA ***, ΔslyA comp, i-wza ***, ΔsapBCDF i-cps ***

2,500 x g, 5 min (right panel)

OD_{600} (supernatant) / OD_{600} (culture)

WT, ΔsapBCDF **, ΔsapBCDF comp

(B)

μg Uronic acid/OD_{600}

WT, ΔargR **, ΔargR comp, ΔmprA **, ΔmprA comp, ΔsapBCDF *, ΔsapBCDF comp, ΔslyA ***, ΔslyA comp

(C)

WT   ΔargR   ΔmprA   ΔslyA   ΔsapBCDF

500 nm

(A)

(B)

(C)