**Genetic continuity in the last seven Millennia in human hepatitis B viruses**

Xiaoyun Lei[1,2], Ye Zhang[1], and Shi Huang[1]*

[1]Center for Medical Genetics, School of Biological Sciences, Central South University, 110 Xiangya Road, Changsha, Hunan 410078, China

[2] Institute of Molecular Precision Medicine, Xiangya Hospital, Central South University, 110 Xiangya Road, Changsha, Hunan 410078, China

*Corresponding author: huangshi@sklmg.edu.cn
The first two authors contributed equally to this work.

**Keywords:**
Ancient DNA, hepatitis B virus HBV, slow clock

**Abstract**
Hepatitis B virus (HBV) is a major human pathogen and yet the evolution history of HBV has largely remained uncertain. With a better theoretical understanding of genetic diversity, we here used a new method to examine the previously published ancient and present day HBV genomes. We identified an informative region in the HBV polymerase that is slow evolving and used it to study genetic distances among HBVs. Three ancient human HBV isolates from 4488-7074 years ago in Germany were identified as genotype G that is also presently common in the same country. We constructed a new phylogenetic tree of HBVs that placed genotype D as the most basal branch with an inferred age of ~20500 years, which is remarkably consistent with the worldwide distribution and a most parsimonious migration route of HBV genotypes today. These results help resolve the evolutionary history of HBV and provide a useful method for studying the phylogenetics of HBV and other viruses in general.

**Introduction:**

Hepatitis B virus (HBV) is a major cause of human hepatitis and related diseases (http://www.who.int/mediacentre/factsheets/fs204/en/). The origin and evolution of HBV has largely remained uncertain, like most viruses. HBV has a circular, partially double-stranded DNA genome of about 3.2kbp that encodes four overlapping open reading frames (P polymerase , pre-S/S envelope, pre-C/C core protein, and X). At least 8 genotypes (A–H) based on nucleotide sequence similarity are classified for human HBV and they have a heterogeneous global distribution (Castelhano, Araujo, Arenas 2017). The putative basal genotypes F and H are found exclusively in the Americas, thus inconsistent with the notion that HBV co-evolved with modern humans as part of the Recent Out of Africa hypothesis. Yet, HBVs in non-human primates (NHP), such as chimpanzees and gorillas, are phylogenetically related to human HBV isolates, seemingly supporting the idea of an Africa origin of the virus (Locarnini et al. 2013; Souza et al. 2014).

Recently, a number of ancient HBV genomes have been uncovered from human skeletons found in Europe and Asia that are between approximately 500-7000 years ago (Kahila Bar-Gal et al. 2012; Krause-Kyora et al. 2018; Muhlemann et al. 2018). While most of the relatively younger HBV genomes (<4488 years ago) were closest to present day human HBVs, all three oldest HBV samples found in Germany (between 4488-7074 years ago) were unexpectedly closest to chimpanzee or gorilla HBVs and hence considered extinct today. The finding challenges expectations as HBV today must have an ancient ancestor which must have infected a large population in the past to have a chance to survive to the present. As large populations have a greater probability of having some of its remains discovered today, the probability of discovering an ancestor of today's human HBVs should be much greater than that of finding a now extinct ancient human HBV sample. Thus, the unusually high rate of discovering ancient human HBV samples that are now extinct (3 independent samples in 3 different archaeological sites) indicates potential flaws in the phylogenetic method employed, especially given that existing methods have yet to produce a consistent evolutionary history of the HBVs. Importantly, the theoretical framework underlying the existing methods, the neutral theory, has been widely known to be inadequate as an explanatory theory of the observed genetic diversity patterns (Kreitman 1996; Ohta, Gillespie 1996; Hahn 2008; Leffler et al. 2012; Hu et al. 2013; Kern, Hahn 2018). It is unfortunate that existing phylogenetic methods have relied heavily on the neutral theory being a valid interpretation of nature.

Different positions in a viral genome are known to have different mutation rates. The fast changing sites in influenza virus play adaptive roles in escaping host immune defense and undergo constant and quick turnovers (Shih et al. 2007). The antigenic sites in human influenza A virus mutate and turn over quickly (several times within a 30 year period), which is critical for the virus to escape host immune defense and hence for flu epidemics. In contrast, other sites stayed largely unchanged within the same period. The influenza results illustrate two general points with regard to evolutionary dynamics of a genome that have so far been overlooked. First, fast evolving or less conserved DNAs are also functional rather than neutral as they are

essential for quick adaptive needs in response to fast changing environments.
Second, fast evolving DNAs turn over quickly and can be shown to violate the infinite
sites model. Hence, they cannot be used for phylogenetic inference involving
evolutionary timescales. If one uses the fast changing sites in a flu virus to infer the
phylogenetic relationship of the virus isolates responsible for different epidemics in a
past period of say 10 years, one would have reached the erroneous conclusion that
each epidemic was caused by a distinct type of flu virus with no genetic continuity
among them rather than just minor variations of the same type.
For short term lineage divergence that has yet to reach saturation for the fast
changing sites, both fast and slow changing sequences could be informative to
phylogeny. However, for evolutionary timescale where divergence in fast changing
sites have reached saturation, only the slow sites (the slow clock method) could be
informative, as has been previously shown and explained by the maximum genetic
diversity (MGD) hypothesis (Huang 2012; Hu et al. 2013; Huang 2016; Yuan, Huang
2017; Yuan et al. 2017). The MGD hypothesis has recently solved the longstanding
puzzle of genetic diversity (Huang 2009; Huang 2016) and made it now possible for
the first time to realistically infer phylogenetic relationships based on genetic diversity
data. It has now been demonstrated that genetic diversities are mostly at saturation
level (Yuan et al. 2012; Yuan et al. 2014; Zhu et al. 2015a; Zhu et al. 2015b; Gui, Lei,
Huang 2017; He et al. 2017; Lei, Huang 2017; Lei et al. 2018; Teske et al. 2018), which
therefore renders most of the past molecular results invalid since those results were
based on mistreating saturated phases of genetic distance as linear phases. Only
slow evolving nuclear sequences are still at linear phase and hence informative to
phylogenetic relationships.
Here, we investigated the genetic relationships among the ancient HBVs and
present day human HBVs using the slow clock method. We found that all three
ancient HBV samples that were thought to group with NHP isolates in fact grouped
with human HBVs. We also constructed a new phylogenetic tree of the human HBV
genotypes, which is remarkably consistent with their distribution patterns.
111
**Results**
**Identity analyses in nucleotide and amino acid sequences**
We selected 3 ancient HBV genomes from Germany for analyses here, 7074 year
old Karsdorf from LBK culture in Lower-Saxony, 5353 year old Sorsum from
Funnelbeaker culture from Lower-Saxony, and 4488 year old RISE563 from Bell
Beaker culture in Osterhofen-Altenmarkt (Krause-Kyora et al. 2018; Muhlemann et al.
2018). The other Bronze age samples were not studied as one (RISE254) was very
close to RISE563 and the others had many sequence gaps. In nucleotide identity, the
three ancient samples were all closer to each other than to any present day samples,
and the highest identity was between Sorsum and RISE563 (Table 1).
122
123
124
125

126 **Table 1. Nucleotide identity among the three ancient HBV genomes**

| Samples | Nucleotide identity | | |
| --- | --- | --- | --- |
| | Karsdorf | Sorsum | RISE563 |
| Karsdorf | | 94.8% | 93.0% |
| Sorsum | | | 95.1% |
| RISE563 | | | |

Gaps not included in the identity calculation.

We searched the Genbank protein database to identify the closest present day
HBV genome to the ancient HBVs in amino acid identity in the polymerase, the largest
open reading frame in the HBV genome (832-845 amino acids). Upon identifying the
closest, we also examined its identity to the ancient HBV in other proteins, pre S, X,
and core proteins (Table 2). Present day HBV isolates closest to the ancient HBVs in
nucleotide identity were not found to be the closest in amino acid identity in the
polymerase. For example, the Karsdorf sample was closest to a chimpanzee HBV
(accession AB032433) in nucleotide sequence but a human HBV (HE981175,
genotype G) in amino acid sequence in the polymerase. However, in other proteins,
the closest to the ancient HBVs were all present day NHP HBVs (Table 2).

**Table 2. Amino acid identities in the polymerase among ancient and present day**
**HBV genomes.**

| Proteins | Species | Karsdorf (7074) | | Sorsum (5353) | | RISE563 (4488) | |
| --- | --- | --- | --- | --- | --- | --- | --- |
| | | Access. | Identity | Access. | Identity | Access. | Identity |
| Polymerase | Human | HE981175 | 92.9% | EU239218 | 92.5% | EU239218 | 90.8% |
| | Chimp. | AB032433 | 92.6% | AB032433 | 92.5% | AB032433 | 90.9% |
| | Gorilla | AJ131567 | 90.6% | AJ131567 | 90.6% | AJ131567 | 89.4% |
| | Orangutan | AF193863 | 88.2% | AF193863 | 88.3% | AF193863 | 87.4% |
| | Gibbon | AJ131571 | 91.3% | AJ131571 | 89.8% | AJ131571 | 88.7% |
| | Karsdorf | Karsdorf | | Karsdorf | 95.5% | Karsdorf | 93.5% |
| | Sorsum | Sorsum | 95.5% | Sorsum | | Sorsum | 94.8% |
| | RISE563 | LT992443 | 93.5% | LT992443 | 94.8% | LT992443 | |
| Pre S | Human | HE981175 | 91.1% | EU239218 | 92.3% | EU239218 | 91.7% |
| | Ape | AB032433 | 92.6% | AB032433 | 95.6% | AJ131567 | 91.5% |
| X prot. | Human | HE981175 | 83.1% | EU239218 | 87.4% | EU239218 | 85.3% |
| | Ape | AB032433 | 84.4% | AB032433 | 90.9% | AJ131567 | 91.7% |
| Core prot. | Human | HE981175 | 94.2% | EU239218 | 91.7% | EU239218 | 92.6% |
| | Ape | AB032433 | 99.3% | AJ131567 | 96.1% | AJ131567 | 98.7% |

Note: Approximate age in years of ancient HBV samples is indicated next to sample
name. The closest human HBV to the ancient HBV in polymerase identity was
selected for comparison. Gaps were excluded in the identity calculation. The

underlined accession numbers indicate the closest modern HBV to the ancient HBV in
nucleotide identity as previously published.

The 3 ancient HBV genomes were also closest to each other in polymerase
amino acid identity than to any other present day samples (Table 2). However,
different from the nucleotide result, the highest amino acid identity in polymerase was
between Karsdorf and Sorsum. As Sorsum differs from RISE563 in both time periods
and locations while only in time periods from Karsdorf, Sorsum is expected to be a
closer relative of Karsdorf and hence to have more similarity in slow changing sites
(amino acid) with Karsdorf. On the other hand, as Sorsum was 1721 years apart from
Karsdorf, ~2 fold more than its time difference with RISE563 (865 years), Sorsum is
expected to have more genetic distance from Karsdorf due to fast changing sites as
may be reflected in the nucleotide sequence. Together, these results showed
significant disconnect between amino acid and nucleotide sequences in revealing
genetic relationships among HBVs.
While results in Table 2 showed clear affinity of Karsdorf with human HBV,
Sorsum showed equal affinity with human and chimpanzee HBVs and RISE563
showed slightly more affinity to chimpanzee than to human HBV, indicating some
uncertainty regarding the informative nature of the full length polymerase protein. The
polymerase is composed of 4 domains, terminal protein, non-conserved spacer,
reverse transcriptase, and RNase H. Upon examining the HBVdb database
(https://hbvdb.ibcp.fr/) (Hayer et al. 2013), together with our own alignment analyses,
we found that the amino acid region corresponding to the reverse transcriptase and
RNase H domains are more conserved or slow evolving (343-844 aa for genotype G
starting with VNL). We therefore tested this 501 aa region to see if it may show better
results than the full length polymerase in linking ancient HBVs with human rather than
NHP (Table 3). Again, all three ancient HBVs showed closer identity, but to a greater
degree, to human HBVs than to NHP HBVs. While Karsdorf was again closest to
Sorsum, it was closer to a present day human HBV (HE981175) than to the ancient
HBV RISE563, indicating clear HBV genetic continuity from the time of Karsdorf to
present time and the more informative nature of the 501 aa slow region of the
polymerase.
As slow evolving DNAs are more likely to be in linear phase and hence more
informative to phylogenetic relationships as explained by the MGD theory, we
examined whether the 501 aa slow region of the polymerase is the slowest evolving
among the protein genes in the HBV genome by comparing amino acid identity
between human and orangutan HBV proteins (Supplementary Table S1). The 501 aa
slow region of the polymerase was found to be the second most conserved, just
slightly less conserved than the core protein. However, because the core protein was
relatively short (178 aa), it is expected to be less informative than a longer protein with
similar degree of conservation. Together with outgroup analyses (see below), we have
found the 501 aa slow region of the polymerase to be the most informative to
phylogenetic inferences of HBV strains.

194 **Table 3. Amino acid identity among ancient and present day HBV genomes in**
195 **the slow region (501 aa)**

| Species | Karsdorf (7074) | | Sorsum (5353) | | RISE563 (4488) | |
|---------|-----------------|---------|---------------|---------|----------------|---------|
| | Access. | Identity | Access. | Identity | Access. | Identity |
| **Human** | HE981175 | 96.0% | HE981175 | 95.4% | EU239218 | 93.9% |
| **Chimp.** | AB032433 | 93.8% | AB032433 | 95.2% | AB032433 | 93.5% |
| **Gorilla** | AJ131567 | 93.8% | AJ131567 | 95.2% | AJ131567 | 93.7% |
| **Orangutan** | AF193863 | 92.0% | AF193863 | 93.2% | AF193863 | 92.1% |
| **Gibbon** | AJ131571 | 93.2% | AJ131571 | 94.0% | AJ131571 | 92.2% |
| **Karsdorf** | Karsdorf | | Karsdorf | 97.4% | Karsdorf | 94.6% |
| **Sorsum** | Sorsum | 97.4% | Sorsum | | Sorsum | 96.2% |
| **RISE563** | LT992443 | 94.6% | LT992443 | 96.2% | LT992443 | |

HBV genomes selected for comparison were the same as those in Table 2. The
underlined accession numbers indicate the closest modern HBV to the ancient HBV in
nucleotide identity as previously published.

**Outgroup inferences based on amino acid mutations**
The above results raise the important question of which type of sequences may
be most informative to HBV phylogeny. For viruses, different hosts may confer
different physiological selection pressures which may result in viruses from different
hosts to have drastic or non-conservative amino acid changes. Taking into account of
non-conservative changes may thus be informative to phylogenetic relationships
where an outgroup NHP HBV to two sister strains of human HBV is expected to show
more non-conservative amino acid changes from the human HBVs.
To confirm the human rather than NHP affinity of the ancient HBV isolates as
shown by the polymerase, we therefore performed protein alignment involving 3
strains, an ancient HBV, its closest human HBV, and its closest NHP HBV. It is
expected that an outgroup should have a higher fraction of non-conservative or
drastic amino acid changes among all mutations that led to differences between the
outgroup and the other two sister strains. We examined those positions where the two
sisters had the same residue while the outgroup was different. We tested each of the
three compared HBV viruses as the candidate outgroup and obtained the fraction of
drastic changes among all positions where the two sisters were the same while the
outgroup was different (Table 4 and Supplementary Materials for details of this
analysis). For the full length polymerase protein, RISE563 as the outgroup had a
significantly smaller fraction of drastic changes than the gorilla HBV as the outgroup
(0.33 vs 0.69, P = 0.004). This indicates that RISE563 and present day human HBV
(EU239218) were sister strains while the gorilla HBV was the outgroup. Similar
analyses showed that for Karsdorf and Sorsum samples, the NHP HBVs all showed
the highest fraction of drastic changes (Table 4). We also performed the combined

analysis where we first add up all the drastic changes of an outgroup (with the
outgroup being ancient HBV, present day human HBV, or NHP HBV) and then
calculated the fraction of drastic changes. The fraction of drastic changes in the NHP
HBV when tested as the outgroup was significantly higher than either that in the
ancient HBVs or the present day human HBVs when they were tested as the outgroup.
We also obtained similar results for the slow region of the polymerase (Table 4).
These results confirmed that ancient HBVs isolates grouped with human rather than
NHP HBVs.

**Table 4. Outgroup inferences from non-conservative (drastic) amino acid changes.**

| Outgroup | | Polymerase | | | | Slow region (aa 343-844) | | | |
|---|---|---|---|---|---|---|---|---|---|
| | | Mutations | | Fraction | P value | Mutations | | Fraction | P value |
| Isolates | Accession | Drastic | All | Drastic | Hu. v Ape | Drastic | All | drastic | Hu. v Ape |
| **Karsdorf** | NA | 5 | 16 | 0.31 | 0.02 | 5 | 10 | 0.50 | n.s. |
| **Human** | HE981175 | 9 | 24 | 0.38 | 0.02 | 4 | 9 | 0.44 | n.s. |
| **Chimpanzee** | AB032433 | 16 | 22 | 0.73 | | 11 | 15 | 0.73 | |
| **Sorsum** | NA | 7 | 16 | 0.44 | n.s. | 2 | 6 | 0.33 | n.s. |
| **Human** | EU239218 | 18 | 39 | 0.46 | n.s. | 5 | 18 | 0.28 | n.s. |
| **Chimpanzee** | AJ131575 | 26 | 39 | 0.67 | | 4 | 7 | 0.57 | |
| **RISE563** | LT992443 | 8 | 24 | 0.33 | 0.004 | 2 | 6 | 0.44 | n.s. |
| **Human** | EU239218 | 16 | 32 | 0.50 | n.s. | 2 | 13 | 0.24 | n.s. |
| **Gorilla** | AJ131567 | 27 | 39 | 0.69 | | 12 | 16 | 0.75 | |
| **Sum** | | | | | | | | | |
| **Ancient human** | | 20 | 56 | 0.36 | <0.0001 | 9 | 22 | 0.41 | 0.03 |
| **Present human** | | 43 | 95 | 0.45 | 0.0003 | 11 | 40 | 0.28 | 0.0002 |
| **Ape** | | 69 | 100 | 0.69 | | 27 | 38 | 0.71 | |

Drastic or non-conservative and conservative changes were according to the
designation by the blastp algorithm. Numbers of these changes can be found in the
Supplementary Table S2. P value was from Fisher's test.

In contrast, for the other three smaller size proteins of HBV genome, the Pre S
protein, the X protein and the pre core protein, none was found informative in
identifying an outgroup (Supplementary Table S2). When we did the same analysis by
using these three proteins as concatenated single molecule, we also failed to identify
any clear outgroup. The fractions of drastic changes in either ancient or present day
human HBVs were similar to that of NHP and showed enrichment of non-conservative
amino acid changes, which was unlike the case for the polymerase. Thus, the ancient
HBVs did not group with either present day NHP or human in any of these proteins.
That the observed changes were enriched for non-conservative amino acid mutations
indicated functional adaptation or selection. Although ancient HBVs all showed
slightly closer identities in these proteins to NHP HBVs, such weak affinity may be

253 fortuitous.

256 **Table 5. Average pairwise identities in the slow region of the HBV polymerase**
257 **(501 aa)**

| Genotype | Identical number of amino acids | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | A | B | C | D | E | F | G | H | All[1] | Lowest[2] |
| A | | 460.75 | 455.75 | 458.25 | 460.5 | 454 | 467 | 451.5 | 458.25 | 468 |
| B | 460.75 | | 467.25 | 461.5 | 463 | 463.5 | 468.5 | 461.5 | 463.71 | >467 |
| C | 455.75 | 467.25 | | 458.25 | 461.5 | 455.5 | 465.5 | 454 | 459.68 | >467 |
| D | 458.25 | 461.5 | 458.25 | | 460.75 | 459 | 461.5 | 458 | 459.61 | 461 |
| E | 460.5 | 463 | 461.5 | 460.75 | | 460 | 471.25 | 453.75 | 461.54 | >471 |
| F | 454 | 463.5 | 455.5 | 459 | 460 | | 461.25 | 474 | 461.04 | >474 |
| G | 467 | 468.5 | 465.5 | 461.5 | 471.25 | 461.25 | | 460.75 | 465.11 | 482 |
| H | 451.5 | 461.5 | 454 | 458 | 453.75 | 474 | 460.75 | | 459.07 | 488 |

[1]Average identities of a genotype to all other genotypes. [2]Lowest pairwise identities
within the genotype.

**Phylogenetic relationships among HBV genotypes**
The above results suggest that the slow region of the polymerase (aa 343-844)
may be the most informative with regard to HBV phylogenetic relationships. We next
used this region to reconstruct the phylogenetic tree for the 8 HBV genotypes by using
the reference genomes for these genotypes (2 genomes for each genotype) as
indexed by the HBVdb database. We first obtained the pairwise identities in the slow
region among the 8 genotypes or 16 genomes and the average identity of each
genotype to the other 7 genotypes (Table 5 and Supplementary Table S3). We also
determined the lowest pairwise identity within each genotype by searching the
Genbank database using the reference genomes and found D to have the lowest
within genotype identity (461 aa), indicating that D has the largest within genotype
genetic diversity among all genotypes. As D was also among the lowest in identity to
all other genotypes (459.61, just slightly greater than the lowest H), D qualifies as the
most basal genotype. We constructed a schematic diagram of the phylogenetic tree
that best fits the data in Table 5 (Figure 1A). Relative to the existing tree (Figure 1B),
the new tree is more consistent with the present day distribution pattern of the HBV
genotypes (Figure 1C). In particular, consistent with expectations, the basal type D is
the most widely distributed and most common in Northeast Asia and Russia/Siberia,
and well positioned to split the first branch, genotype F and H, specific to the New
World.

**Figure 1. Phylogenetic tree of HBV genotypes.** Branch lengths are relatively and roughly to scale. The tree is meant to be more of a schematic diagram. A. Tree built by using the slow region. B. Tree built by using nucleotide sequences as found in Muhlemann et al (Muhlemann et al. 2018). C. Distribution map of HBV genotypes.

Using the slow region of the polymerase, we also examined the relationships of selected NHP HBVs with the human reference genomes and found closer relative affinity of African NHP HBVs with genotype G and of Asian NHP HBVs with genotype C (Table 6). Although we only looked at one HBV genome each for each NHP species, the closest chimpanzee and gorilla in nucleotide sequence to the ancient human HBVs and an arbitrarily chosen orangutan and gibbon HBV, the relative affinity should hold for other NHP HBVs. We also examined the identity of ancient HBVs with the human HBV reference genomes and found all three to be most related to genotype G, with Sorsum and RISE563 relatively closer to genotype E than Karsdorf. The results suggest that Sorsum and RISE563 may be on their way diverging from genotype G to E. Based on the amino acid difference between Karsdorf and genotype G (501-479 = 22 amino acid) and the age of Karsdorf, we inferred the mutation rate in the slow region to be 2.0 x 10$^{-6}$ aa per aa per year and the age of G at ~14500 years (14500 x

$501 \times 2.0 \times 10^{-6} + 7500 \times 501 \times 2.0 \times 10^{-6} = 22$ aa). Given the distance between the

basal genotype D and the other genotypes to be ~41 as well as the largest within

genotype distance of 40 for D, we inferred an origination of human HBV at ~20500

years ago followed soon by the divergence of genotype F and H as people crossed

the Bering Strait into the New World.

**Table 6. Identities between NHP and ancient HBV isolates with the reference**

**genomes of human HBV genotypes**

| Genotype | Identical number of amino acids | | | | | | |
|---|---|---|---|---|---|---|---|
| | AB032433 | AJ131567 | AF193863 | AJ131571 | | | |
| | Chimp. | Gorilla | Orangutan | Gibbon | Karsdorf | Sorsum | RISE563 |
| A | 458 | 459 | 449 | 452 | 470 | 467 | 463 |
| B | 463 | 461.5 | 457 | 462 | 467 | 473.5 | 468.5 |
| C | 462.5 | 459 | 463 | 461.5 | 463.5 | 470 | 467 |
| D | 459.5 | 460 | 454.5 | 455.5 | 463 | 468.5 | 461 |
| E | 463.5 | 461 | 455.5 | 456.5 | 461 | 469 | 470.5 |
| F | 458.5 | 459 | 454 | 458 | 461.5 | 466.5 | 471.5 |
| G | 468 | 466 | 458 | 465 | 479 | 476 | 475.5 |
| H | 454 | 454.5 | 452.5 | 462 | 455.5 | 460.5 | 466 |

Values represent the average of two genomes of each genotype.

**Discussion**

Our results suggest genetic continuity in the last seven Millennia for human HBV

and establish a more informative method for building phylogenetic trees for HBVs.

Our findings on the ancient HBV isolates here, rather than the conclusions from

previous analyses (Krause-Kyora et al. 2018; Muhlemann et al. 2018), appear more

consistent with a priori expectation on the ancient ancestors. That the oldest Karsdorf

sample (7074 years ago) found in Germany was closest to present day genotype G

that is also common in Germany, France and the United States is also consistent with

expectation and further supports the conclusion of genetic continuity in HBVs in the

last 7000 years (Sunbul 2014). That Sorsum and RISE563 samples were closely

related to Karsdorf indicates also an affinity of these two isolates with genotype G

although they were relatively more related to genotype E than Karsdorf. E is the most

closely related to G among present HBV isolates. As E is found today only in Africa

(Andernach, Hubschen, Muller 2009), our finding here is in line with the known

migration of Europeans into Africa during the last 5000 year period (Fregel et al.

2018).

The closer nucleotide distance of Sorsum with RISE563 than with Karsdorf may

reflect linear phase distance whereas the closer amino acid distance of Sorsum with

Karsdorf may reflect continuity in housekeeping functions. That Karsdorf was most

similar to gorilla HBV in nucleotide sequence but to present day human HBV in amino

acid sequence in the polymerase or the slow region may reflect continuation in the
housekeeping physiology of the viruses and merely fortuitous similarities in nucleotide
sequences reflecting some shared environmental adaptation. The closer similarity of
ancient HBVs with NHP HBVs in non-polymerase proteins may reflect
non-housekeeping adaptive roles in these proteins: the ancient HBV non-polymerase
proteins in the ancient human hosts are expected to be more similar to NHP proteins
given that the primitive life style of ancient humans may be less different from that of
NHP.
The three ancient samples differ from genotype G however in that they lacked the
36 nucleotide insertion in the core gene region that is specific to HBV-G. This
indicates that the insertion of this 36 bp sequence may have taken place later than
7000 years ago.
Genotype D is known to have the largest worldwide distribution, which is
consistent with its basal position in the new tree here and its widest within genotype
genetic diversity as found here. D shares the 33 nucleotide deletion in the pre S
region with the NHP HBVs and the ancient HBVs, again indicating its more primitive
nature. As all present day genotypes other than D lacks the 33 nucleotide deletion, we
can infer that this deletion occurred only in the last few thousand years and
independently in each genotype. As African NHP HBVs were closer to G while Asian
NHP HBVs were closer to C, there was likely transmission of human HBVs into NHPs
via migration of G from Europe into Africa and migration of C in the Asia mainland
where it is most common today into the islands in Southeast Asia.
The new HBV phylogenetic tree appears more consistent with the reality
genotype distribution around the globe than the existing ones and has implications for
the origin of the virus. That the first split was between D and the rest followed by the
split of F and H indicates that HBV may have originated in a region covering Northeast
Asia and Siberia where D may have originated. It then went to either the West to give
rise to G, E, and A or the South to give rise to B and C. The estimated age of the
human HBV of ~20500 years corresponds to the period of Last Glacial Maximum
(LGM), indicating that human population expansion post LGM may have played a role
in the origin of HBV (Tallavaara et al. 2015).
The study here demonstrates the informative nature of the slow clock method in
resolving the longstanding uncertainty on HBV origin and evolution. While it is widely
known that different regions of a virus genome may produce different phylogenetic
trees, it remains uncertain as to which region or whether the whole genome is the
most informative to phylogenetic inferences. The MGD theory and the study here
illustrate the informative nature of the slow evolving regions. The method should be
generally applicable to evolutionary studies of other viruses.
377
378
**Materials and Methods:**
Present day HBV sequences were retrieved from Genbank and HBVdb database.
Ancient HBV genomes were downloaded using previously published accession
numbers. Alignments in nucleotide and amino acid sequences were done by blast.

Fisher's exact test was used to estimate p values, 2 tailed.

**Acknowledgements:**

We thank Ben Krause-Kyora and Julian Susat for sharing the sequences of ancient

human HBV samples. This work is supported by the National Natural Science

Foundation of China grant 81171880, the National Basic Research Program of China

grant 2011CB51001, and the Furong Scholars program (S. H.).

**Author contributions**: XL, YZ, SH designed the study and performed data analyses.

SH wrote the first draft and all authors contributed to the final version.

**Financial Interest statements:** The authors declare that they have no competing

interests that might be perceived to influence the results and/or discussion reported in

this paper.

**Tables:**

**Figure legends:**

**Figure 1. Phylogenetic tree of HBV genotypes.** Branch lengths are relatively and

roughly to scale. The tree is meant to be more of a schematic diagram. A. Tree built

by using the slow region. B. Tree built by using nucleotide sequences as found in

Muhlemann et al (Muhlemann et al. 2018). C. Distribution map of HBV genotypes.

**References:**

Andernach, IE, JM Hubschen, CP Muller. 2009. Hepatitis B virus: the genotype E

puzzle. Rev Med Virol 19:231-240.

Castelhano, N, NM Araujo, M Arenas. 2017. Heterogeneous recombination among

Hepatitis B virus genotypes. Infect Genet Evol 54:486-490.

Fregel, R, FL Mendez, Y Bokbot, et al. 2018. Ancient genomes from North Africa

evidence prehistoric migrations to the Maghreb from both the Levant and

Europe. Proc Natl Acad Sci U S A 115:6774-6779.

Gui, Y, X Lei, S Huang. 2017. Collective effects of common SNPs and genetic risk

prediction in type 1 diabetes. Clin Genet.

Hahn, MW. 2008. Toward a selection theory of molecular evolution. Evolution

62:255-265.

Hayer, J, F Jadeau, G Deleage, A Kay, F Zoulim, C Combet. 2013. HBVdb: a

knowledge database for Hepatitis B Virus. Nucleic Acids Res 41:D566-570.

He, P, X Lei, D Yuan, Z Zhu, S Huang. 2017. Accumulation of minor alleles and risk

prediction in schizophrenia. Sci Rep 7:11661.

Hu, T, M Long, D Yuan, Z Zhu, Y Huang, S Huang. 2013. The genetic equidistance

result, misreading by the molecular clock and neutral theory and

427              reinterpretation nearly half of a century later. Sci China Life Sci 56:254-261.

428    Huang, S. 2009. Inverse relationship between genetic diversity and epigenetic

429              complexity. Preprint available at Nature

430              Precedings:doi.org/10.1038/npre.2009.1751.1032.

431    Huang, S. 2012. Primate phylogeny: molecular evidence for a pongid clade excluding

432              humans and a prosimian clade containing tarsiers. Sci China Life Sci

433              55:709-725.

434    Huang, S. 2016. New thoughts on an old riddle: What determines genetic diversity

435              within and between species? Genomics 108:3-10.

436    Kahila Bar-Gal, G, MJ Kim, A Klein, et al. 2012. Tracing hepatitis B virus to the 16th

437              century in a Korean mummy. Hepatology 56:1671-1680.

438    Kern, AD, MW Hahn. 2018. The Neutral Theory in Light of Natural Selection. Mol Biol

439              Evol 35:1366-1371.

440    Krause-Kyora, B, J Susat, FM Key, et al. 2018. Neolithic and medieval virus genomes

441              reveal complex evolution of hepatitis B. Elife 7.

442    Kreitman, M. 1996. The neutral theory is dead. Long live the neutral theory. Bioessays

443              18:678-683; discussion 683.

444    Leffler, EM, K Bullaughey, DR Matute, WK Meyer, L Segurel, A Venkat, P Andolfatto,

445              M Przeworski. 2012. Revisiting an old riddle: what determines genetic diversity

446              levels within species? PLoS Biol 10:e1001388.

447    Lei, X, S Huang. 2017. Enrichment of minor allele of SNPs and genetic prediction of

448              type 2 diabetes risk in British population. PLoS ONE 12:e0187644.

449    Lei, X, J Yuan, Z Zhu, S Huang. 2018. Collective effects of common SNPs and risk

450              prediction in lung cancer. Heredity:doi:10.1038/s41437-41018-40063-41434.

451    Locarnini, S, M Littlejohn, MN Aziz, L Yuen. 2013. Possible origins and evolution of

452              the hepatitis B virus (HBV). Semin Cancer Biol 23:561-575.

453    Muhlemann, B, TC Jones, PB Damgaard, et al. 2018. Ancient hepatitis B viruses from

454              the Bronze Age to the Medieval period. Nature 557:418-423.

455    Ohta, T, JH Gillespie. 1996. Development of Neutral and Nearly Neutral Theories.

456              Theor Popul Biol 49:128-142.

457    Shih, AC, TC Hsiao, MS Ho, WH Li. 2007. Simultaneous amino acid substitutions at

458              antigenic sites drive influenza A hemagglutinin evolution. Proc Natl Acad Sci U

459              S A 104:6283-6288.

460    Souza, BF, JF Drexler, RS Lima, O Rosario Mde, EM Netto. 2014. Theories about

461              evolutionary origins of human hepatitis B virus in primates and humans. Braz J

462              Infect Dis 18:535-543.

463    Sunbul, M. 2014. Hepatitis B virus genotypes: global distribution and clinical

464              importance. World J Gastroenterol 20:5427-5434.

465    Tallavaara, M, M Luoto, N Korhonen, H Jarvinen, H Seppa. 2015. Human population

466              dynamics in Europe over the Last Glacial Maximum. Proc Natl Acad Sci U S A

467              112:8232-8237.

468    Teske, PR, TR Golla, J Sandoval-Castillo, A Emami-Khoyi, CD van der Lingen, S von

469              der Heyden, B Chiazzari, B Jansen van Vuuren, LB Beheregaray. 2018.

470              Mitochondrial DNA is unsuitable to test for isolation by distance. Sci Rep

471    8:8448.

472 Yuan, D, S Huang. 2017. On the peopling of the Americas: molecular evidence for the

473    Paleoamerican and the Solutrean models. bioRxiv bioRxiv 130989; doi:

474    https://doi.org/10.1101/130989

475 Yuan, D, X Lei, Y Gui, Z Zhu, D Wang, J Yu, S Huang. 2017. Modern human origins:

476    multiregional evolution of autosomes and East Asia origin of Y and mtDNA.

477    bioRxiv:doi: https://doi.org/10.1101/106864.

478 Yuan, D, Z Zhu, X Tan, et al. 2012. Minor alleles of common SNPs quantitatively affect

479    traits/diseases and are under both positive and negative selection.

480    arXiv:1209.2911.

481 Yuan, D, Z Zhu, X Tan, et al. 2014. Scoring the collective effects of SNPs: association

482    of minor alleles with complex traits in model organisms. Sci China Life Sci

483    57:876-888.

484 Zhu, Z, Q Lu, J Wang, S Huang. 2015a. Collective effects of common SNPs in

485    foraging decisions in Caenorhabditis elegans and an integrative method of

486    identification of candidate genes. Sci. Rep.:doi:10.1038/srep16904.

487 Zhu, Z, D Yuan, D Luo, X Lu, S Huang. 2015b. Enrichment of Minor Alleles of

488    Common SNPs and Improved Risk Prediction for Parkinson's Disease. PLoS

489    ONE 10:e0133421.