Title:

Eye-position error influence over "open-loop" smooth pursuit

initiation

Abbreviated title:

Transient disruption of smooth pursuit initiation

Authors:

Antimo Buonocore[1,2], Julianne Skinner[1,2,3], Ziad M. Hafed[1,2]

1. Werner Reichardt Centre for Integrative Neuroscience, Tuebingen University,

Tuebingen, Germany, 72076

2. Hertie Institute for Clinical Brain Research, Tuebingen University, Tuebingen,

Germany, 72076

3. International Max Planck Graduate School of Behavioral and Neural Sciences,

Tuebingen University, Tuebingen, 72076

Corresponding author:

Antimo Buonocore

Werner Reichardt Centre for Integrative Neuroscience

Otfried-Mueller Str. 25

Tuebingen, 72076, Germany

antimo.buonocore@cin.uni-tuebingen.de

Number of pages: 54

Number of figures: 9

Number of words in abstract: 248

Number of words in introduction: 643

Number of words in discussion: 1591

Conflicts of interest: None

Acknowledgements:

We were funded by the Hertie Institute for Clinical Brain Research and by the

Deutsche Forschungsgemeinschaft (DFG) (Grant: FOR1847; HA6749/2-1). We also

received support from the Werner Reichardt Centre for Integrative Neuroscience

(CIN). The CIN is an Excellence Cluster (EXC 307) funded by the DFG.

Abstract

The oculomotor system integrates a variety of visual signals into appropriate motor plans, but such integration can have widely varying time scales. For example, unlike with rapid saccades, it has been suggested that initiating a smooth pursuit eye movement to follow a moving target involves an obligatory open-loop interval, in which new visual motion signals presumably cannot influence the ensuing motor plan for up to 100 ms after movement initiation. However, this view runs directly contrary to the idea that the oculomotor periphery has privileged access to short-latency visual signals. Here we show that smooth pursuit initiation is highly sensitive to visual inputs, even in open-loop intervals. We instructed male rhesus macaque monkeys to initiate saccade-free smooth pursuit eye movements, and we injected a transient, instantaneous eye position error signal at different times relative to movement initiation. We found highly robust short-latency modulations in eye velocity and acceleration, starting only ~50 ms after transient signal occurrence, and even during open-loop pursuit initiation. Critically, the spatial direction of the injected position error signal had predictable effects on smooth pursuit initiation, with forward errors increasing eye acceleration and backwards errors reducing it. These effects persisted during sustained pursuit, and also affected catch-up saccades in a similar manner. Importantly, catch-up saccade kinematics were altered ~50 ms after transient signal occurrence, much like microsaccade kinematic alterations during simple fixation. Our results demonstrate that smooth pursuit initiation is highly sensitive to visual signals, and that catch-up saccades behave like regular microsaccades and saccades.

Introduction

A highly intriguing property of motor circuits is that they have privileged access to sensory signals. For example, neck muscles and related upper body motor synergies, which are normally recruited to re-orient the upper body towards behaviorally-relevant stimuli, exhibit very early stimulus-locked responses after visual flashes (Corneil, 2004; Corneil et al., 2013; Corneil and Munoz, 2014). These stimulus-locked responses, arriving <100 ms after stimulus onset, occur almost as early as neural responses in primary visual areas (Corneil, 2004; Boehnke and Munoz, 2008; Corneil and Munoz, 2014). Similarly, oculomotor control circuitry in the brainstem are riddled with eye-movement related neurons exhibiting both visual and eye-movement related discharge in the very same neurons. In fact, at least in the superior colliculus (SC), it is only such visual-motor neurons in particular that exhibit the most perceptual and behavioral relevance to task success in some cases (Wurtz and Mohler, 1976; Chen and Hafed, 2017). This concomitance of visual and motor signals in the very same neurons occurs even in pre-motor nuclei (Evinger and Kaneko, 1982; Everling et al., 1998; Missal and Keller, 2002), a mere single synapse away from the motor neurons innervating the ocular muscles.

Intuitively, access to the sensory periphery by motor systems makes sense. For example, in the case of vision and the oculomotor system, such access can allow very rapid resetting of oculomotor plans in order to make appropriate orienting reflexes (Reingold and Stampe, 1999, 2002; Buonocore and McIntosh, 2008; Rolfs et al., 2008; Hafed and Ignashchenkova, 2013). Such resetting becomes evident in eye movement behavior after a stimulus flash, and it can also have large repercussions on eye movement kinematics (Edelman and Xu, 2009; Guillaume, 2012; Buonocore et al., 2016; Buonocore et al., 2017), even before most higher-level cortical visual areas finalize processing of the appearing stimuli (Schmolesky et al., 1998). That is,

when one carefully analyzes the properties of eye movements a mere ~50-100 ms

after stimulus onset, one discovers that a full processing cycle from visual input all

the way to eye movement output has taken place (Fischer and Boch, 1983; Boch et

al., 1984; Fischer and Ramsperger, 1984; Tian et al., 2018). This rapid access to

visual inputs imparts on the oculomotor system a highly efficient pathway for

behavioral re-orienting.

However, even within a single motor output modality, different classes of

behaviors can have very different time scales. For example, saccadic eye

movements occur on a much more rapid time scale when compared to much slower

smooth pursuit eye movements used to track a moving target. Even though recent

research has shown that saccades and smooth pursuit eye movements do share

neural resources (Krauzlis, 2004), the different time scales between these two types

of eye movements have led to descriptions of categorical differences between them.

One such difference concerns the initiation of the two types of eye movements. While

saccade initiation may be independent of visual stimulation, smooth pursuit eye

movements require motion signals (Rashbass, 1961). More importantly, smooth

pursuit eye movements are overwhelmingly described as possessing an "open-loop"

interval of ~100 ms, before which new visual motion signals cannot influence the

motor behavior (Lisberger and Westbrook, 1985). This directly contradicts the idea

that the oculomotor periphery, which ultimately influences both saccades and smooth

pursuit, has privileged access to visual signals.

Here we found that smooth pursuit initiation is indeed sensitive, and with short

latencies, to visual stimuli within the same time window as saccades. We introduced

instantaneous visually-defined eye position errors to the smooth pursuit system

during the initiation of tracking eye movements. We found a highly systematic

directional influence of such signals on smooth pursuit initiation, and with latencies of

only 50-80 ms. These results, directly paralleling saccadic ones (Reingold and

Stampe, 2002; Buonocore and McIntosh, 2008; Buonocore et al., 2017), suggest that

the concept of "open-loop pursuit" may be too coarse to accurately describe the

mechanisms of smooth pursuit initiation.

Materials and Methods

Animal preparation and laboratory setup

We recorded eye movements with high spatial and temporal precision from two male

rhesus macaque monkeys (monkey M and monkey A) aged 7 years. The monkeys

were implanted, under general anesthesia and using sterile surgical techniques, with

a titanium head-post attached to the skull with titanium skull screws (all implanted

under the skin and muscle surrounding the skull). The head-post allowed stabilizing

the animals' head position during the experiments. In a subsequent surgery, one eye

in each animal was implanted with a scleral search coil to allow tracking eye

movements using the magnetic induction technique (Fuchs and Robinson, 1966;

Judge et al., 1980). The animals are part of a larger neurophysiology experiment for

which the behavioral training needed for the current study was a necessary

prerequisite. The experiments were approved by ethics committees at the regional

governmental offices of the city of Tuebingen, and these experiments were in

accordance with European Union guidelines on animal research, as well as the

associated implementations of these guidelines in German law.

During the experiments, the animals were seated in a primate chair 74 cm

from a CRT computer monitor in an otherwise dark room. The monitor had a pixel

resolution of 34 pixels/deg and a refresh rate of 120 Hz, and stimuli were presented

over a uniform gray background (29.7 Cd/m^2). A small white spot (5 x 5 min arc

square) having 86 Cd/m^2 luminance was used as either a fixation spot or the moving

target for smooth pursuit eye movements (see *Behavioral tasks* below). Other stimuli

that we used consisted of white flashes of different sizes and positions, as described

in *Behavioral tasks* below; they all had the same luminance as the spot (86 Cd/m^2).

The CRT monitor viewed by the monkeys was controlled by a computer running

Matlab's Psychophysics Toolbox (Brainard, 1997; Pelli, 1997; Kleiner et al., 2007).

This computer was in turn receiving commands from a real-time I/O system from

National Instruments (Austin, USA), which was maintaining control of the display on a

frame-by-frame basis. The system was programmed in-house and described recently

(Chen and Hafed, 2013; Tian et al., 2016).

Behavioral tasks

Experiment 1: Saccade-free smooth pursuit initiation with transient position error

signals

We employed a paradigm eliciting smooth (i.e. saccade-free) ocular following of a

moving target while we introduced a transient eye position error signal in a specific

direction. Trials started with a white fixation spot presented at the center of the

display over a uniform gray background (spot and background luminance were

described above). After the monkey aligned gaze with the fixation spot by 700-1000

ms, the spot jumped by 4 deg either to the right or left of fixation and simultaneously

started moving horizontally (at a speed of 20.4 deg/s) in the opposite direction (i.e.

there was a leftward step followed by rightward motion or a rightward step followed

by leftward motion). The step size was chosen such that by the time a smooth pursuit

eye movement (in the target motion direction) was initiated, no initial catch-up

saccade was needed (Rashbass, 1961). This allowed us to achieve saccade-free

smooth pursuit initiation velocities and, therefore, to explore how instantaneous eye

position error signals could modify such smooth velocity initiation.

Instantaneous eye position error signals were introduced by interleaving trials

in which the target motion above was accompanied, for a single display frame (~8

ms), by a white flash of the same luminance as the pursuit target spot. The flash,

appearing 50-167 ms (in steps of ~8 ms) after target motion onset, could be a full-

screen flash or a square of 1 x 1 deg dimensions centered on a position either 2.1

deg in front of or 2.1 deg behind the instantaneous pursuit target position (along the

motion trajectory; in monkey M, we also ran the experiment with the square

appearing at 6 deg, instead of 2.1 deg, in front of or behind the instantaneous pursuit

target position). Thus, with gaze being near the pursuit target, the "forward" flash was

introducing an instantaneous eye position error signal in the direction of the smooth

pursuit eye movement that was being initiated, and the "backward" flash was

introducing an instantaneous eye position error signal in the opposite direction. The

full-screen flash was expected to act more like the backward flash, since there was

more stimulus energy behind the pursuit target than ahead of it given flash timing in

the trials (the flashes often happened when the eye already started to follow the

pursuit target, leaving more than half of the display width behind the direction of gaze

at the time of the flashes).

The monkeys were rewarded for maintaining their gaze within 3 deg from the

pursuit target position until trial end (occurring 500 ms from target motion onset). This

virtual gaze window for monitoring monkey behavior was relatively relaxed because

we found that the flash could momentarily "attract" gaze position (with both smooth

velocity and saccadic effects; see Results), and also because the monkeys were

highly trained on accurate gaze alignment on the pursuit target anyway. Across

multiple sessions, we collected 3624 trials from monkey M and 1996 trials from

monkey A in this task (with the forward and backward flashes being 2.1 deg in front

of or behind the pursuit target). We also collected 3705 trials from monkey M in the

variant with forward and backward flashes being 6 deg in front of or behind the

pursuit target. The four experimental conditions (normal saccade-free pursuit

initiation without a flash, initiation with a full-screen flash, initiation with a forward

flash, or initiation with a backward flash) were randomly interleaved within a session.

The monkeys typically completed ~300 trials of this task per session, with the

remainder of the session dedicated to the other tasks below, or to other

neurophysiology-related tasks not described in this study.

Experiment 2: Saccadic smooth pursuit initiation with transient position error signals

We repeated the same experiment above, except that there was no initial step in

target position before the target motion onset. That is, in the present experiment, the

monkeys steadily fixated the fixation spot at the center of the display, and the spot

then started moving at a constant speed directly from the center. Because there is an

unavoidable latency in reacting to the spot's motion onset, the monkeys had to

perform an initial catch-up saccade as part of initiating their smooth tracking

behavior, and our purpose here was to explore the influence of instantaneous eye

position error signals introduced by flashes on such an initial catch-up saccade. All

other details were identical to those described above for Experiment 1. We collected

1824 trials from monkey M and 1511 trials from monkey A in this task, and the

forward and backward flashes in this experiment were only at a 2.1 deg eccentricity

relative to the pursuit target.

Experiments 3 and 4: Steady-state pursuit or fixation accompanied by a transient

flash stimulus

To compare the effects of transient instantaneous eye position error signals on

smooth pursuit initiation to effects during steady-state pursuit or during steady-state

gaze fixation, we ran two additional versions of the above experiments, which we

pool here because of their conceptual similarity to each other. In the first variant

(Experiment 3), the flashes could appear during steady-state smooth pursuit as

opposed to during smooth pursuit initiation. Trials started with the fixation spot

presented at +/- 7 deg horizontally from display center, to allow for a long period of

sustained smooth pursuit spanning large portion of the horizontal extent of our

display area. After 300-500 ms of fixation, the target started moving towards (and

past) the center of the display at a speed of 12 deg/s. The motion lasted for ~1500-

1800 ms, and at a random time during steady-state smooth pursuit (0-1500 ms from

motion onset; in steps of ~8 ms), a flash could appear as in Experiments 1 and 2

above (forward and backward flashes were at 2.1 deg). We explored the effects of

the flash and its type (forward, backward, or full-screen) on both smooth velocity

effects as well as on catch-up saccades. We collected 5085 trials from monkey M

and 2849 trials from monkey A in this experiment. In the second variant of the

experiment (Experiment 4), the flashes could happen during steady-state fixation,

and we explored the effects of the flashes on microsaccades (Hafed and

Ignashchenkova, 2013; Buonocore et al., 2017). The flashes could happen 75-999

ms after steady fixation of a central fixation spot. We collected 5222 trials from

monkey M and 3160 trials from monkey A in this experiment.

Data analysis

Detecting catch-up saccades and microsaccades

We numerically differentiated our digitized eye position signals to obtained estimates

of eye velocity and eye acceleration. We used velocity and acceleration criteria to

detect microsaccades and catch-up saccades, as described previously (Krauzlis and

Miles, 1996a; Hafed et al., 2009; Chen and Hafed, 2013). Briefly, a radial velocity

threshold identified candidate saccadic events in the eye position traces, and

acceleration thresholds were then used to refine the start and endpoint of a given

movement. For a subset of data, we used a machine learning approach for saccade

detection (Bellet et al., 2018). All detected microsaccades and catch-up saccades

were manually inspected to correct for misses or false detections.

Measuring smooth eye velocity effects

In Experiment 1, our aim was to isolate the influence of transient instantaneous eye position error signals on smooth pursuit initiation. Our task design minimized the occurrence of catch-up saccades. In the analysis, we replaced all saccades and catch-up saccades from the eye trace with Not-a-Number (NaN) labels that extended in duration to +/- 50 ms around each detected saccade. This way, we made sure that smooth pursuit averages were not influenced by high velocity changes due to the rare saccades that may have occurred. This procedure avoided catch-up saccades during initiation, and, equally important, our exclusion of microsaccades also avoided the potential influence of these fixational eye movements (occurring near target motion onset) on visual sensitivity and behavior (Hafed and Krauzlis, 2010; Hafed, 2013; Chen et al., 2015; Hafed et al., 2015; Tian et al., 2016; Chen and Hafed, 2017; Tian et al., 2018). We then averaged the radial eye velocity traces aligned on motion onset and/or flash onset for the different trial types (e.g. no flash, full-screen flash, forward flash, or backward flash). The no flash eye velocity traces acted as our baseline, to which we compared velocities when there was a flash during pursuit initiation.

To perform such comparisons, we first plotted radial eye velocity after motion onset for the no flash trials as well as for either full-screen, forward, or backward flash trials. Plotting the averages side by side (e.g. see Fig. 1A in Results) allowed us to establish that there was indeed an influence of early transient flashes on "open-loop" smooth pursuit initiation. We then plotted the difference in eye velocity between flash and no flash trials. To do this, we subtracted within each session the average radial eye velocity in no flash trials from the radial eye velocity of every flash trial of a given type (e.g. for forward flashes only); we then repeated this subtraction for all

sessions from a given monkey. This allowed us to achieve a population of velocity

"difference" curves isolating the influence of a given flash type (e.g. Fig. 1B in

Results); we estimated variance in this population by calculating the 95% confidence

interval of the difference curves across all trials in all sessions.

To summarize the impact of a given flash type on eye velocity, we identified a

measurement interval (either 50-100 ms after full flash onset or 50-80 ms after

backward/forward flash onset) in the velocity difference curves described above; this

interval was chosen because it was the interval during which the flash exerted the

influence over eye movements that we were particularly interested in investigating

(see Results). For all trials of a given type (e.g. forward flash, backward flash, or full-

screen flash), we averaged the eye velocity difference in this post-flash interval, and

this gave us a population of measurements across trials. We then compared

populations of measurements across the different flash conditions in our trial types

(e.g. forward versus backward flashes) and also for different flash times relative to

smooth pursuit initiation.

To compare the influence of transient instantaneous eye position error signals

on smooth pursuit initiation to their influence during steady-state smooth pursuit, we

repeated the above analyses but for data containing steady-state pursuit (Experiment

3). For all flash times occurring >200 ms after target motion onset (to ensure that

smooth pursuit was already ongoing) and >200 ms before trial end, we plotted radial

eye velocity from -100 ms to 200 ms relative to flash onset. Importantly, we removed

from the eye traces all catch-up saccades (along with flanking +/- 50 ms intervals) to

avoid effects of high velocity changes on our measurements. We averaged trials from

each type (e.g. full-screen flash, forward flash, or backward flash) to demonstrate the

influence of the flash on smooth eye velocity.

Measuring effects on catch-up saccades and microsaccades

We analyzed catch-up saccade times, amplitudes, and peak velocities. Since the absolute majority of catch-up saccades occurred in the direction of ongoing smooth pursuit (e.g. 97% in monkey A and 98% in monkey M from the pursuit initiation trials in Experiment 2; and 90% in monkey A and 97% in monkey M in Experiment 3), we restricted our catch-up saccade analyses to saccades in the direction of pursuit. We summarized catch-up saccade frequency of occurrence, amplitude, and peak velocity with and without flashes, and as a function of forward or backward flashes. We also analyzed the influence of flash time during smooth pursuit initiation.

We sometimes referred to saccade metrics and kinematics in relation to the eye position error that was present at the time of triggering a saccade. Eye position error was defined as the instantaneous retinotopic position of the pursuit target 10 ms before catch-up saccade onset (i.e. instantaneous target position minus eye position).

To compare changes in catch-up saccade frequencies and metrics/kinematics as a function flash occurrence to changes in other types of saccades, we performed similar analyses on microsaccades during steady-state fixation (Experiment 4). In this case, microsaccades could happen in all directions, so we classified them according to flash direction (for the non-full-screen flash trials); that is, we classified microsaccade directions according to their horizontal component as being in the same direction of the flashed square or opposite it, similar to earlier work (Hafed and Ignashchenkova, 2013; Tian et al., 2016; Tian et al., 2018). Our motivation for comparing catch-up saccades to microsaccades is that microsaccades were recently shown to exhibit kinematic changes after transient visual signals similar to larger saccades (Buonocore et al., 2017); thus, microsaccades are a viable reference for

comparing catch-up saccade effects during smooth tracking to saccades without

such smooth tracking (Heinen et al., 2016; Heinen et al., 2018)

To investigate saccade kinematic changes, we followed recent findings that

specific post-flash saccades have a relationship between saccadic peak velocity and

saccadic amplitude that deviates from the expected saccadic main sequence curve

(Guillaume, 2012; Buonocore et al., 2016; Buonocore et al., 2017). To summarize

this behavior in our own data, we computed the ratio between saccade peak velocity

and amplitude. This value was then compared between a baseline interval before

flash onset, or in some cases from no flash trials, and an interval during the

appropriate post-flash period.

Experimental design and statistical analysis

Each experiment underwent within-subject analyses, and we reported summaries

and statistics from each subject individually. We used two male monkeys, as stated

above, to ensure replicability of results while being constrained by the 3R (Refine,

Reduce, Replace) principles for animal research. Combined with congruence of

several key ingredients of our results with several prior studies in monkeys (e.g.

(Krauzlis and Miles, 1996b; Hafed and Ignashchenkova, 2013; Buonocore et al.,

2017) and humans (e.g. (Schutz et al., 2008; Edelman and Xu, 2009; Buonocore et

al., 2016; Braun et al., 2017), we have high confidence that our results from the 2

monkeys are a general property of the smooth pursuit eye movement system.

In each experiment, we randomly interleaved the relevant conditions for

statistical comparisons (e.g. no flash, full-screen flash, forward flash, and backward

flash in equal proportion). We also ensured a minimum of 300 repetitions per

condition for the great majority of our analyses. Such a large number of repetitions

made using parametric statistical comparisons in our analyses well justified. For

some analyses of catch-up saccades or microsaccades during a "saccadic inhibition"

post-flash phase of trials (Reingold and Stampe, 1999, 2000, 2002; Buonocore and

McIntosh, 2008; Hafed and Ignashchenkova, 2013; Buonocore et al., 2017), there

was (by definition of the phenomenon) a relative rarity of saccades to analyze.

However, we ensured statistical robustness even for these observations by

increasing the number of repetitions per condition that we collected overall (see

above), such that there was still statistical confidence in the relatively fewer inhibition-

phase saccades that we were analyzing (we show 95% confidence intervals for all of

our analyses).

Further statistical validation was obtained by performing, for each animal, two-

way ANOVAs (factor "flash time" by factor "flash direction", with the appropriate

number of factor levels depending on the specific design within the experiment) on

the summary statistics described above (e.g. amplitude of saccades and ratio of

velocity over amplitude in a post-flash period). Post hoc comparisons were followed

up with two-sample t-tests. Either simple or two-sample t-tests were also performed

to compare different conditions when the ANOVA was not necessary; alpha levels

were corrected following Bonferroni methods. We performed all of our statistical tests

using standard Matlab (MathWorks, Inc.) functions.

Results

Smooth pursuit initiation is thought to involve an "open-loop" interval, of ~100 ms, during which eye velocity is presumably not yet under visual feedback error control (Lisberger and Westbrook, 1985). However, visual signals can enter into the processing hierarchy of the oculomotor system significantly earlier than 100 ms. For example, transient visual responses in the brainstem can occur with latencies shorter than ~40-50 ms (Rizzolatti et al., 1980; Boehnke and Munoz, 2008; Chen et al., 2018). We thus hypothesized that there exist mechanisms in which even "open-loop" smooth pursuit initiation can be significantly modified by visual inputs, and in a spatially-specific manner. We tested this by introducing a brief (~8 ms) transient visual stimulus onset that effectively added a transient instantaneous eye position error signal to the smooth pursuit system (Materials and Methods). We found that such a signal was highly effective in modifying smooth pursuit initiation. To demonstrate this, we first present cases in which the transient visual onset consisted of a full-screen white flash, and we then show results when spatially-precise position error signals were introduced by smaller transient flashes.

Transient visual signals perturb smooth pursuit initiation velocity even during "open-loop" intervals

Our monkeys initiated smooth pursuit eye velocity without saccades through our use of a Rashbass step-ramp target trajectory (Experiment 1; Materials and Methods) (Rashbass, 1961). After a period of steady-state fixation, the pursuit target stepped in a direction opposite to the motion trajectory and instantaneously moved at constant speed, such that by the time at which smooth pursuit was initiated, there was no need for a catch-up saccade (de Brouwer et al., 2002). At a random time during the initiation phase of smooth pursuit, we presented a full-screen flash. Figure 1A shows

an example of saccade-free smooth pursuit initiation with and without a full-screen

flash from monkey M. The gray curve shows average radial eye velocity aligned on

the onset of target motion across no-flash trials (i.e. control trials), and error bars

denote 95% confidence intervals. As expected from the task design, eye velocity

increased smoothly to approach target velocity, and the onset of smooth velocity

acceleration occurred after a latency of ~100 ms from target motion onset. However,

when a full-screen flash occurred 50 ms after target motion onset (blue curve in Fig.

1A), the initiation of smooth pursuit was transiently disrupted (indicated by the arrow

and label "Dip" in the figure); there was a transient reduction in eye acceleration

starting ~50 ms after flash onset, and almost coincident with the very earliest phase

of smooth pursuit initiation. This effect can be better visualized in Fig. 1B; here, we

subtracted, for every flash trial, the average radial eye velocity of the no-flash

condition from the given trial's radial eye velocity, and we then averaged all velocity

"difference" curves (Materials and Methods). The average velocity difference curve

showed a clear transient dip starting ~50 ms after full-screen flash onset (error bars

denote 95% confidence intervals). Importantly, we ensured that this data did not

contain any microsaccades or catch-up saccades (Materials and Methods). Thus,

early smooth pursuit initiation (so-called "open-loop" pursuit) is reliably modified by

transient visual signals.

The transient change in saccade-free pursuit initiation that we observed after

full-screen flashes was always time locked to flash onset. For example, Fig. 1C, D

shows results similar to those from Fig. 1A, B above (and from the same monkey),

except that the flash now happened 83 ms after target motion onset rather than after

50 ms. A similar transient decrease in eye acceleration during pursuit initiation was

evident, but it now happened slightly later in the initiation phase of smooth pursuit.

Similarly, Fig. 1E, F shows results with flash onset 117 ms after target motion onset,

and Fig. 1G, H shows results with flash onset 150 ms after target motion onset. In all

cases, there was a transient reduction in eye acceleration (marked by the label "Dip"

in each figure panel), such that the eye velocity difference curves (Figs. 1B, D, F, H)

always showed a negative-going transient ~50 ms after flash onset. These transients

are reminiscent of reductions in saccade and microsaccade frequency with short

latency after full-screen visual flashes (Reingold and Stampe, 1999, 2000, 2002;

Buonocore and McIntosh, 2008; Rolfs et al., 2008; Edelman and Xu, 2009; Bompas

and Sumner, 2011; Hafed and Ignashchenkova, 2013; Tian et al., 2018). However, in

our case, the changes to smooth pursuit initiation happened in a more "analog"

fashion for smooth eye velocity as opposed to a more "digital" or "all-or-none" fashion

for saccade frequency.

To demonstrate that the transient changes in smooth eye velocity were time

locked to full-screen flash onset in both of our monkeys, we computed the velocity

difference curves described above (e.g. Fig. 1B), but we did so for all flash times that

we tested, and also after aligning the curves to flash onset as opposed to target

motion onset. The left panel of Fig. 1I shows the results of this analysis for monkey

A, and the right panel shows the results for monkey M. Note that for presentation

purposes, the figure shows curves from every other flash time, but the results were

identical for all flash times: regardless of flash time, its influence on smooth pursuit

initiation appeared in each monkey ~50 ms after flash onset. To statistically verify this

observation, we averaged the value of the curve for each flash trial in the interval 50-

100 ms after flash onset and tested against zero. As expected, for both monkeys,

there was reliable "inhibition", or a decrease in the difference velocity signal, across

flash onset times (results of the statistical tests are reported in the caption of Fig. 1).

The above results indicate that even open-loop smooth pursuit initiation is

modifiable by visual signals. We confirmed that such modification is similar to what

might happen during sustained, steady-state smooth pursuit, as was recently

reported in humans (Kerzel et al., 2010). Specifically, in a subset of our experiments

(Experiment 3; Materials and Methods), we presented the full-screen flash during

sustained steady-state smooth pursuit. We picked an interval of smooth pursuit eye

movements around the time of full-screen flash onset, and we removed any catch-up

saccades present in such interval (Materials and Methods). Like described above, we

also plotted radial eye velocity aligned on flash onset time (Fig. 2; error bars denote

95% confidence intervals). Full-screen flashes had a very similar effect to their effect

during smooth pursuit initiation: steady-state smooth pursuit eye velocity started

being transiently reduced ~50 ms after flash onset. Therefore, our results so far

indicate that transient visual onsets reliably modify even open-loop smooth pursuit

initiation, and in a manner similar to what happens during steady-state pursuit.

Transient instantaneous eye position error signals modify smooth pursuit initiation

velocity in a direction-dependent manner

Given the results above, might it be possible that transient spatially-specific position

error signals can also modify open-loop smooth pursuit initiation, and in a spatially-

specific manner? To answer this question, we turned back to our saccade-free

smooth pursuit initiation experiments of Fig. 1. We analyzed trials in which we

presented, for only one display frame (~8 ms), a small white square of 1 deg

dimension either 2.1 deg ahead of or 2.1 deg behind the smooth pursuit target. The

square could appear at different times relative to target motion onset; therefore, the

flash "in front" of the pursuit target could sometimes happen either "behind" or "in

front of" instantaneous gaze position depending on flash timing. For example, if

target motion was rightward and a "forward" flash (i.e. 2.1 deg to the right of the

pursuit target) happened ~50 ms after motion onset, then the flash was actually still

to the left of instantaneous gaze position because the target motion had not yet

crossed gaze position (with a step of 4 deg leftward followed by 20.4 deg/s rightward

motion, the target would still be leftward relative to gaze at ~50 ms). On the other

hand, if the same flash happened ~150 ms after motion onset, then the flash was

already ahead of instantaneous gaze position. According to our hypothesis of a

spatially-specific position error signal introduced by the flash, this means that both

"forward" and "backward" flashes relative to pursuit target location should transiently

reduce smooth pursuit acceleration if the flash happened ~50 ms after target motion

onset, but "forward" flashes should increase (rather than decrease) smooth pursuit

acceleration if the flashes happened ~150 ms after target motion onset.

This is indeed what we found. In Fig. 3A, B, we plotted data in a format similar

to Fig. 1A, B above, but only for "forward" flashes happening 50 ms after motion

onset (from monkey M); similarly, in Fig. 3C, D, we plotted the same data for

"backward" flashes happening 50 ms after motion onset (from the same monkey). In

both cases, transient reductions in eye acceleration happened. On the other hand,

when the flash happened 150 ms after motion onset, the transient stimulus increased

eye acceleration for forward flashes (Fig. 3E, F), while it reduced it for backward

flashes (Fig. 3G, H). This is because for forward flashes relative to target location, a

flash occurring 150 ms after target motion onset was also "forward" with respect to

instantaneous gaze position.

We summarized the observations above across both monkeys in Fig. 4A, B.

For each monkey, we measured the value of the velocity difference curve between

flash and no-flash trials (e.g. Fig. 1B, D, F, H and Fig. 3B, D, F, H). We did so by

averaging the value of the curve for each flash trial in the interval 50-80 ms after flash

onset; this gave us a population of measurements for any given flash time and flash

location, and we first performed statistics by estimating 95% confidence intervals on

such a population. We then plotted the velocity difference for a given flash type (e.g.

forward or backward relative to pursuit target position) as a function of flash onset

time relative to pursuit target motion onset (Fig. 4A, B; error bars denote 95%

confidence intervals). The pink curve for each monkey shows the effect of a forward

flash (relative to pursuit target position) on smooth eye velocity, and the green curve

shows the effect of a backward flash. Clearly, there were effects of both forward and

backward flashes on even open-loop pursuit initiation. Specifically, and consistent

with Fig. 3, both monkeys showed reliable influences of spatially-localized flashes on

smooth pursuit initiation, with a transition from inhibition (negative y-axis values) to

enhancement (positive y-axis values) in the forward flash conditions (relative to no

flash pursuit initiation) according to the observations of Fig. 3. We further validated

this result by performing a two-way ANOVA (flash time by flash location) for each of

the monkeys. As expected, we found a significant interaction between the two factors

of flash time and flash location (monkey A: $F(14, 753) = 4.1$, $p = 6.4414*10^{-7}$; monkey

M: $F(14, 1693) = 3.47$, $p = 1.3060*10^{-5}$), supporting the hypothesis that for longer

delays, the two flashes were differentiating inhibition from enhancement of eye

acceleration. In particular, starting from flashes happening at around 100 ms after

motion onset (monkey A: 116 ms; monkey M: 92 ms), forward flashes led to a

smaller inhibition that turned into an enhancement when compared to backward ones

(Fig. 3). Therefore, open-loop smooth pursuit initiation is actually modifiable, and it

reliably reflects the influence of transient instantaneous position error signals

introduced by spatially localized flashes.

To show that enhancement of smooth pursuit initiation acceleration can still

happen even for flashes as early as 50 ms after target motion onset, we repeated the

experiment of Fig. 3 one more time, but this time, the forward or backward flashes

happened 6 deg in front of or behind the pursuit target position. This means that even

if the flash happened ~50 ms after target motion onset, the flash in front of the pursuit

target was also in front of instantaneous gaze position, unlike in Fig. 3. In this case,

summary plots like those shown in Fig. 4A, B should show enhancement for forward

flashes even for early flashes; this was indeed the case (Fig. 4C). In this case, a two-

way ANOVA (flash location by flash time) showed a predominant main effect of the

factor flash location (monkey M: $F(1, 1707) = 121.03$, $p = 3.0190*10^{-27}$), statistically

supporting the interpretation that forward flashes induced enhancement while

backward flashes led to inhibition. We also observed a main effect of flash time

(monkey M: $F(14, 1707) = 2.53$, $p = 0.0014$) suggesting that later flashes had a lower

inhibitory component (in Fig. 4C, it is possible to appreciate this effect as a positive

slope across time for both flash directions) and a small interaction between the two

factors (monkey M: $F(14, 1707) = 1.8$, $p = 0.0341$). Therefore, even the inhibitory

influences seen for early forward flashes in Fig. 3A, B above were not a result of a

generalized inhibitory effect of flashes on eye movements, as sometimes suggested

from the saccadic inhibition literature with similar flashes (Reingold and Stampe,

1999, 2000, 2002; Buonocore and McIntosh, 2008). Instead, the flash was

introducing a spatially-defined geometric addition to the input of the oculomotor

system, as is also the case with the saccadic system (Hafed and Ignashchenkova,

2013; Buonocore et al., 2017). What is most intriguing is that such a spatially-specific

input was able to modify even so-called open-loop smooth pursuit initiation.

Finally, we also analyzed the effects of spatially localized flashes during

sustained pursuit. In the limit of steady-state smooth pursuit, a forward flash was also

effective in increasing saccade-free smooth pursuit eye velocity, and a backward

flash was effective in reducing smooth eye velocity (Fig. 5A, B; also see Fig. 2 for an

analog of this result with full-screen flashes). Therefore, transient instantaneous eye

position error signals have spatially-specific effects on both open-loop and sustained

smooth pursuit eye movements, and the results of Fig. 1 above are not merely a

generalized inhibitory effect of flashes on eye movements.

Catch-up saccades are influenced by transient instantaneous position error signals

similarly to smooth pursuit eye velocity

We explored the analogy between smooth pursuit initiation effects and saccade

effects alluded to above further. Specifically, we recently hypothesized that at the

time of flash-induced action potentials in oculomotor circuits, like the superior

colliculus (SC), a saccade that is triggered would have an amplitude reflecting the

addition of visually-induced action potentials to the already-preprogrammed saccade-

related action potentials (Buonocore et al., 2017). Thus, saccades would violate the

expected monotonic relationship (Zuber et al., 1965; Bahill et al., 1975) between

saccade amplitude and peak velocity, because these saccades would integrate

visually-driven action potentials with already-programmed motor-related action

potentials into the total eye displacement. Our smooth pursuit effects above (Figs. 1-

5) are consistent with this mechanism, and further support for this would emerge if

one were to analyze the trials in which catch-up saccades did occur after flash onset

in our smooth pursuit experiments (as opposed to only saccade-free trials in Figs. 1-

5). Therefore, during steady-state smooth pursuit and the 1x1 deg flashes

(Experiment 3; Materials and Methods), we took only trials in which a catch-up

saccade was triggered 50-150 ms after flash onset (Fig. 6A, G). Such a saccade

putatively happened at the same time at which visually-induced action potentials from

flashes exerted an influence over the oculomotor system in Figs. 1-5.

We aligned all catch-up saccades together and classified them as having a

direction either congruent or incongruent with "forward" or "backward" flashes relative

to the pursuit target in Experiment 3. For comparison, we also analyzed catch-up

saccades on no-flash control trials (using a virtual timing of fictive flashes that did not

occur in reality). Figure 6B, C shows eye position (Fig. 6B) and eye velocity (Fig. 6C)

for catch-up saccades from monkey A congruent with the flash location; the gray

curves show catch-up saccades during steady-state smooth pursuit with no flashes,

and all eye position curves were aligned to eye position at catch-up saccade onset

before averaging. Similarly, Fig. 6D shows eye position and Fig. 6E eye velocity for

catch-up saccades incongruent with flash location, and gray curves show no-flash

catch-up saccades. Error bars in all cases denote 95% confidence intervals. As can

be seen, for forward flashes, catch-up saccades triggered 50-150 ms after flash

onset were larger than those without a flash (two-sample t-tests: $t = 6.992$; $df = 1707$;

$p < 0.0001$) (Fig. 6B). Peak eye velocity seemed to also increase correspondingly for

these movements (Fig. 6C). For backward flashes (Fig. 6D), we observed a small

reduction in amplitude, as it might have been expected (Buonocore et al., 2016;

2017) and also a small reduction in peak velocity. In Figure 6F, we plotted the ratio

between peak velocity and saccadic amplitude as an index of main sequence

violation. Conceptually, if the peak velocity is not appropriately scaled to the

larger/smaller saccades generated during the post-flash interval, the ratio between

velocity and amplitude should lead to values bigger/smaller when compared to the

ratio computed when no flash was presented (Buonocore et al., 2017). In monkey A,

we observed that the peak velocity was in fact scaling with saccadic amplitude when

backward flashes were presented. On the other hand, there was a trend for the ratio

to be smaller than baseline for forward flashes (two-sample t-tests: $t = 1.571$; $df =$

1707; $p = 0.1163$) (Fig. 6F), suggesting the idea that main sequence violation can

reliably happen during the post-flash period even with smooth pursuit. Similar results

in amplitude modulation were observed for monkey M, but with a stronger and more

significant trend for the saccades in violating the main sequence. (Fig. 6G-L). All the

statistics comparing catch-up saccades triggered in the no-flash conditions against

catch-up saccades triggered when either forward or backward flashes were

presented are reported in the caption of Fig. 6.

We also confirmed that during smooth pursuit initiation, there was again an

effect on saccades and not just on smooth eye velocity. In Experiment 2 (Materials

and Methods), target motion started directly from fixation, requiring an initiation

saccade during smooth pursuit initiation. We analyzed the kinematics of such an

initiation saccade (first saccade occurring after motion onset and with latency 30-80

ms from forward or backward flash onset) and compared them to kinematics of

saccades when no flash was presented. We observed reliable modulations of

saccade kinematics similar to the ones that we observed during steady state pursuit

(Fig. 7). In the top row of Fig. 7, saccades were larger than baseline after a forward

flash for both monkey A (Fig. 7A pink bar) and monkey M (Fig. 7B) (Monkey A, two-

sample t-tests: $t = 9.113$; $df = 255$; $p < 0.0001$; monkey M, two-sample t-tests: $t =$

8.411; $df = 335$; $p < 0.0001$). Backward flashes reliably reduced saccade amplitudes

relative to baseline in monkey A (two-sample t-tests: $t = -5.278$; $df = 217$; $p < 0.0001$)

and monkey M (two-sample t-tests: $t = -8.205$; $df = 283$; $p < 0.0001$). To explore

kinematic violations like in the case of catch-up saccades during sustained pursuit

(Fig. 6), we plotted the ratio of peak velocity and amplitude for post-flash saccades

(Fig. 7C, D). This ratio was different from baseline behavior in both monkeys. In

particular, in monkey A, the increase in amplitude after forward flashes was not

accompanied by a corresponding increase in peak velocity, resulting in a smaller

ratio than baseline ($t = 2.459$; $df = 255$; $p = 0.0146$), while no significant violation was

found for backward flashes. In monkey M, the violation after backward flashes

occurred because saccade amplitude decreased without an appropriate scaling down

of the peak velocity ($t = -6.053$; $df = 283$; $= p < 0.0001$).

Taken together, our results so far indicate that the sudden onset of a visual

transient can interfere with the spatial parameters of the motor plan by addition of

visual spikes, and even potentially making saccade amplitude relatively larger for a

given peak velocity in the case of forward flashes, and the opposite pattern for

backward flashes (Buonocore et al., 2016; Buonocore et al., 2017).

The temporal structure of catch-up saccades is reset by transient visual onsets

similarly to the temporal structure of microsaccades during gaze fixation

Finally, we explored the effects of localized visual flashes during stable fixation. The

stimuli that we used were identical to the ones of the previous experiments with the

only difference being that the fixation target was not moving in the present

experiments. In these conditions, we replicated the previous observation that flash

onset strongly reduced the probability to make a microsaccade in an interval of 50 to

150 ms after flash onset for both monkeys (Fig. 8A, B), a phenomenon known as

(micro)saccadic inhibition (Engbert and Kliegl, 2003; Rolfs et al., 2008; Hafed et al.,

2011; Hafed and Ignashchenkova, 2013). More interestingly, during such an

inhibitory period (very similar to the one observed for catch-up saccades; e.g. Fig.

6A, G), the kinematics of microsaccades were also modulated in a way similar to

what we observed for catch-up saccades generated during smooth pursuit initiation.

In particular, for both monkeys, flashes congruent with microsaccade direction were

pulling the amplitude of eye movements higher, but no effect was observed for

backward flashes (Fig. 8C, D). We validated this observation by performing a two-

way ANOVA (flash time by flash location) on microsaccade amplitude for each of the

monkeys. As expected, we found a significant interaction between the two factors

(monkey A: $F(1, 159) = 6.43$, $p = 0.0122$; monkey M: $F(1, 251) = 5$, $p = 0.0263$).

Follow-up analysis confirmed that flashes congruent with microsaccade direction

were indeed leading to the generation of larger microsaccades during the inhibitory period (monkey A, two-sample t-tests: t = -5.53; df = 72; p < 0.0001; monkey M, two-sample t-tests: t = -3.113; df = 131; p = 0.0023). We then tested if the microsaccades launched during the inhibition interval (50-150 ms after flash onset; Fig. 8A, B) were also violating the main sequence. As predicted from previous research on the topic (Guillaume, 2012; Buonocore et al., 2016; Buonocore et al., 2017), and from the results reported above, a pattern of violation was revealed again in both animals. In particular, we observed a general reduction in the velocity over amplitude ratio for both animals (Fig. 8E, F). A two-way ANOVA (flash time by flash location) highlighted a significant main effect of flash time (monkey A: $F(1, 159) = 6.41$, $p = 0.0123$; monkey M: $F(1, 251) = 4.03$, $p = 0.0457$) with no interaction between factors, supporting the hypothesis that the peak velocity was not appropriately scaling with the modulations in amplitude caused by flash onset (Buonocore et al., 2017).

After confirming similarities between the spatiotemporal parameters of catch-up saccades produced during smooth pursuit and microsaccades generated during stable fixation, we turned to analyze the temporal structure of catch-up saccades and microsaccades when no flashes were presented. As a first analysis, we compared inter-saccadic intervals (Fig. 9A, B) and saccade rates (Fig. 9C, D) in the two tasks. In Fig. 9C, D, it is possible to appreciate that both animals produced eye movements with a similar rate, independently of the task being sustained pursuit (orange curve) or stable fixation (violet curve). Moreover, the inter-saccadic interval (i.e. the timing between consecutive movements) was also mainly overlapping between the two tasks, and it was peaking around 200-250 ms in both animals (Fig. 9A, B). Taken together, these results support the hypothesis that the temporal structure of microsaccades and catch-up saccades is similar .

The third observation in our comparison of catch-up saccades to

microsaccades regards instead the eye position error (i.e. the Euclidian distance

between the current fixation location and the eye position 10 ms before producing a

saccade). In Fig. 9E, F, we could observe clearly that there was a substantial

difference in the position error between sustained pursuit (orange) and stable fixation

(violet). In particular, both monkeys had a higher position error during sustained

pursuit, as might be expected. This finding supports the hypothesis that the urge to

make a saccade (i.e. when to trigger an eye movement) was relatively independent

of the eye position error (i.e. where the eyes are relative to the target/fixation) within

the ranges of position error that were present in our study. One might therefore argue

that, up to a certain threshold, an internal clock governing the triggering of

(micro)saccades is fairly constant and unaltered by the ongoing velocity of the eyes

(Tian et al., 2018).

Discussion

We injected brief, perceptually subtle visual flashes around the time of smooth

pursuit initiation, and we observed robust modulations in eye acceleration. These

modulations were spatially-specific, depending on flash location relative to

instantaneous gaze position, and they were therefore not reflecting a generalized

inhibitory effect similar to freezing behavior in prey animals. Instead, eye acceleration

during pursuit initiation was equally easy to increase or decrease, even for flashes

appearing as early as ~50 ms after target motion onset (i.e. even before the onset of

smooth tracking itself). We also observed that when smooth pursuit was initiated with

the aid of catch-up saccades, these saccades were affected by flashes similarly to

how microsaccades (Buonocore et al., 2017) and larger saccades (Buonocore et al.,

2016) are affected, and also similarly to how the smooth eye movements themselves

were modulated. Our results demonstrate that so-called "open-loop" smooth pursuit

is more nuanced than assumed, and they motivate neurophysiological investigations

of rapid sensory and motor processing in the oculomotor system.

Beyond saccadic inhibition

Our experiments were motivated by related research investigating the impacts of

brief flashes on saccades (Reingold and Stampe, 1999; Buonocore and McIntosh,

2008). Initially, such flashes were discovered to transiently inhibit saccades (Reingold

and Stampe, 1999) and microsaccades (Engbert and Kliegl, 2003; Rolfs et al., 2008;

Hafed et al., 2011). However, later research demonstrated that the flashes also have

directionally-specific influences (Hafed and Clark, 2002; Engbert and Kliegl, 2003;

Hafed and Ignashchenkova, 2013; Tian et al., 2016; Tian et al., 2018), as well as

effects on saccade kinematics (Buonocore et al., 2016; Buonocore et al., 2017). For

example, microsaccades occurring 50-100 ms after flash onset and directionally-

congruent with its location are not inhibited as strongly as opposite microsaccades

(Hafed and Ignashchenkova, 2013; Tian et al., 2016; Tian et al., 2018), and

sometimes even enhanced (as in Fig. 6G for monkey M catch-up saccades).

Moreover, congruent microsaccades are enlarged, whereas opposite microsaccades

are smaller, as if to reflect averaging of the saccade-vector command with "visual"

action potentials induced by flash onset (Hafed and Ignashchenkova, 2013; Tian et

al., 2016; Buonocore et al., 2017).

We observed the same kinematic alterations in catch-up saccades as in

microsaccades/saccades (e.g. Figs. 6-8) even though catch-up saccades rode on

high eye velocity. Catch-up saccades were also inhibited after flash onset, and with

directionally-specific patterns (e.g. Fig. 6A, G). In fact, we ran our monkeys on a

fixation task in the same sessions, and replicated our recent findings (Hafed and

Ignashchenkova, 2013; Tian et al., 2016; Buonocore et al., 2017). Our results on

catch-up saccades therefore support recent statements that these saccades behave

similarly to microsaccades (Heinen et al., 2016; Heinen et al., 2018). We now extend

this idea further to include the effects of transient flashes on the saccadic system.

Most intriguingly, even when no catch-up saccades occurred, we still observed

direct correlates of the saccadic phenomena described above on smooth eye

movements. Smooth pursuit initiation was enhanced or inhibited depending on flash

location, and sustained pursuit was also affected. The most interesting added

advantage of performing the experiments on smooth pursuit rather than saccades

was that we could focus on open-loop saccade-free pursuit initiation, demonstrating

that it is equally malleable as sustained pursuit. We specifically found that open-loop

pursuit is easily modifiable by visual signals injecting an instantaneous transient eye

position error. Thus, the term "saccadic inhibition" does not encompass the entire

space of impacts of brief flashes on the oculomotor system.

In our experiments, the catch-up saccades of monkey A, in particular, after

flash onset were not strongly reduced in amplitude for backward flashes, but they

were significantly larger after forward flashes (Fig. 6B, D). We think that observations

like these may be interpreted as reflecting the fact that smooth pursuit was ongoing.

For example, backward flashes reduced smooth eye velocity itself in addition to

influencing saccades (Figs. 2, 5). Therefore, there was a conflict between the

increased eye position error due to slower eye velocity (supporting a bigger catch-up

saccade) and the backward visually-induced action potentials of the flash (supporting

a smaller catch-up saccade). Given the importance of eye position error in smooth

pursuit (de Brouwer et al., 2002), it could be the case that the reduction in smooth

velocity compensated for effects of the flash on catch-up saccade amplitude. By a

similar vein, because monkey A had slightly lower eye velocity than monkey M in

tracking (Fig. 2), there was a bigger likelihood that there was positive eye position

error at the time of the flash onset. This condition, of a plan to make a corrective eye

movement coinciding with a congruent visual flash in the same direction, is known in

microsaccades to have maximal effects on the movement being planned (Tian et al.,

2016; Tian et al., 2018). That is why monkey A had a larger effect of forward flashes

on catch-up saccade amplitude (Fig. 6B).

Smooth pursuit mechanisms

We also observed smooth velocity effects that we think reflect the constant interplay

between target position/motion and the instantaneous ongoing eye behavior. For

example, we found that smooth velocity effects after forward flash onset had a robust

post-enhancement suppression phase. For example, for forward flashes in

Experiments 1 and 3, smooth pursuit acceleration or sustained velocity increased

(what we termed enhancement in Results) and then decreased later (resulting in a

later dip in the velocity difference curves; e.g. Fig. 5). Such a subsequent decrease

could reflect the simple fact that the enhancement causes gaze to overpass the

pursuit target, resulting in negative eye position error. A brief deceleration after the

enhancement thus brings pursuit back on target.

Therefore, our smooth velocity effects relate, in a sense, to studies of

distractor effects on pursuit (Spering et al., 2006). We think that we complement

these observations and highlight the relevance of position error for influencing

smooth eye velocity. Such an influence even happens during fixation with smooth

ocular drifts (Martins et al., 1985; Kowler and Blaser, 1995). In all, this suggests

interesting future neurophysiology experiments that not only focus on the

specification of target motion signals for pursuit, but that investigate, at the level of

the brainstem and other cerebellar and cortical areas, the aggregate interplay

between visual and motor signals, with and without transient flashes. One prediction

of our work is that SC recordings should reveal simultaneity of flash-induced visual

action potentials with alterations in smooth eye velocity, but it would be interesting to

know whether other areas down- or upstream of SC play a more critical role.

What triggers catch-up saccades?

An additional interesting observation from our fixation experiments is that ongoing

microsaccade rate and inter-saccadic intervals during sustained fixation were

qualitatively similar to those during sustained pursuit, despite different position errors.

Even flash-induced modulations in microsaccade or catch-up saccade rate were

similar to each other (i.e. there was rate inhibition followed by rate enhancement with

similar dependencies on saccade direction relative to flash location; Figs. 6, 8). This

raises an intriguing question on what triggers catch-up saccades. It is known that

position error and retinal slip (velocity error) are important for triggering these

saccades (de Brouwer et al., 2002), but there is also tolerance for these errors such

that some combinations of position and velocity errors go briefly uncorrected. This

could explain why a "catch-up saccade rate signature" after flash onset looks

qualitatively identical to the so-called "microsaccade rate signature" after flash onset.

Time since the previous catch-up saccade seems to also matter, and could

additionally explain the above tolerance for position and velocity errors. Indeed, there

seems to be temporal structure to when catch-up saccades occur during smooth

pursuit, similar to that for microsaccades (Fig. 9). Thus, the effect of flashes on the

catch-up saccade rate signature during pursuit (Fig. 6A, G) could be explained by the

same phase-resetting mechanisms associated with microsaccadic temporal structure

(Hafed and Ignashchenkova, 2013).

Microsaccades are also sensitive to eye position error (Guerrasio et al., 2010;

Ko et al., 2010; Tian et al., 2016), extending the similarities to catch-up saccades

mentioned earlier. In the end, a combination of time since the last saccade and

oculomotor errors must matter a great deal for what triggers either catch-up or

microsaccades (Tian et al., 2018).

Implications for experiments

Finally, experiencing our flashes in pilot human experiments revealed a huge

contrast between their perceptual appearance to the observer and their impacts on

the oculomotor system. These brief stimuli (even full flashes) seem, at face value, to

be too transient and inconsequential to affect eye movement behavior (or even

perceptual processing). Yet, they do, and in a most robust manner (e.g. Fig. 1). This

means that certain experimental manipulations might seem so subtle to task

designers but fundamentally systematic for the brain processes being invoked by the

tasks. This can influence experimental result interpretations. For example, there is

interest in probing perceptual state during smooth pursuit initiation by presenting brief

stimuli around the time of eye movements (Spering et al., 2006; Schutz et al., 2008;

Braun et al., 2017). Because our results suggest that pursuit is indeed altered by

brief stimuli, even during initiation, ideal stimuli for these kinds of experiments would

be ones that would not alter retinal image statistics if smooth pursuit were to be

transiently slowed down or accelerated. Otherwise, experimental results may reflect

differences in retinal image statistics when pursuit is altered, rather than more

interesting perceptual and cognitive factors. A second similar example is the

employment of brief stimulus flashes as attentional cues during fixation and

assuming that microsaccades are irrelevant, an assumption that we now know to be

too simplistic (Hafed et al., 2015).

References

Bahill AT, Clark MR, Stark L (1975) The main sequence, a tool for studying human eye movements. Mathematical Biosciences 24:191-204.

Bellet ME, Bellet J, Nienborg H, Hafed ZM, Berens P (2018) Human-level saccade detection performance using deep neural networks. bioRxiv doi: 10.1101/359018.

Boch R, Fischer B, Ramsperger E (1984) Express-saccades of the monkey: Reaction times versus intensity, size, duration, and eccentricity of their targets. Experimental brain research Experimentelle Hirnforschung Expérimentation cérébrale 55.

Boehnke SE, Munoz DP (2008) On the importance of the transient visual response in the superior colliculus. Curr Opin Neurobiol 18:544-551.

Bompas A, Sumner P (2011) Saccadic Inhibition Reveals the Timing of Automatic and Voluntary Signals in the Human Brain. Journal of Neuroscience 31:12501-12512.

Brainard DH (1997) The Psychophysics Toolbox. Spatial vision 10:433-436.

Braun DI, Schutz AC, Gegenfurtner KR (2017) Visual sensitivity for luminance and chromatic stimuli during the execution of smooth pursuit and saccadic eye movements. Vision Res 136:57-69.

Buonocore A, McIntosh RD (2008) Saccadic inhibition underlies the remote distractor effect. Experimental brain research Experimentelle Hirnforschung Expérimentation cérébrale 191:117-122.

Buonocore A, McIntosh RD, Melcher D (2016) Beyond the point of no return: effects of visual distractors on saccade amplitude and velocity. Journal of Neurophysiology 115:752-762.

Buonocore A, Chen C-Y, Tian X, Idrees S, Münch TA, Hafed ZM (2017) Alteration of the microsaccadic velocity-amplitude main sequence relationship after visual transients: implications for models of saccade control. Journal of Neurophysiology 117:1894-1910.

Chen C-Y, Hafed ZM (2017) A Neural Locus for Spatial-Frequency Specific Saccadic Suppression in Visual-Motor Neurons of the Primate Superior Colliculus. Journal of Neurophysiology:jn.00911.02016-00973.

Chen C-Y, Sonnenberg L, Weller S, Witschel T, Hafed ZM (2018) Spatial frequency sensitivity in macaque midbrain. Nature Communications 9:2852.

Chen CY, Hafed ZM (2013) Postmicrosaccadic enhancement of slow eye movements. The Journal of neuroscience : the official journal of the Society for Neuroscience 33:5375-5386.

Chen CY, Ignashchenkova A, Thier P, Hafed ZM (2015) Neuronal Response Gain Enhancement prior to Microsaccades. Curr Biol 25:2065-2074.

Corneil BD (2004) The Superior Colliculus: New Approaches for Studying Sensorimotor
Integration. Methods & New Frontiers in Neuroscience.Edited byWilliam C Hall
andAdonis Moschovakis. Boca Raton (Florida): CRC Press. $119.95. xvi + 324 p + 4 pl;
ill.; index. ISBN: 0–8493–0097–5. 2004. The Quarterly Review of Biology 79:457-457.

Corneil BD, Munoz DP (2014) Overt Responses during Covert Orienting. Neuron 82:1230-
1243.

Corneil BD, Cheng JC, Goonetilleke SC (2013) Dynamic and opposing adjustment of
movement cancellation and generation in an oculomotor countermanding task. J
Neurosci 33:9975-9984.

de Brouwer S, Yuksel D, Blohm G, Missal M, Lefevre P (2002) What triggers catch-up
saccades during visual tracking? Journal of neurophysiology 87:1646-1650.

Edelman JA, Xu KZ (2009) Inhibition of Voluntary Saccadic Eye Movement Commands by
Abrupt Visual Onsets. Journal of Neurophysiology 101:1222-1234.

Engbert R, Kliegl R (2003) Microsaccades uncover the orientation of covert attention. Vision
Research 43:1035-1045.

Everling S, Paré M, Dorris MC, Munoz DP (1998) Comparison of the discharge characteristics
of brain stem omnipause neurons and superior colliculus fixation neurons in monkey:
Implications for control of fixation and saccade behavior. Journal of Neurophysiology
79:511-528.

Evinger C, Kaneko CR (1982) Activity of omnipause neurons in alert cats during saccadic eye
movements and visual stimuli. Journal of Neurophysiology.

Fischer B, Boch R (1983) Saccadic eye movements after extremely short reaction times in the
monkey. Brain Research 260:21-26.

Fischer B, Ramsperger E (1984) Human express saccades: extremely short reaction times of
goal directed eye movements. Experimental brain research Experimentelle
Hirnforschung Expérimentation cérébrale 57:191-195.

Fuchs AF, Robinson DA (1966) A method for measuring horizontal and vertical eye
movement chronically in the monkey. J Appl Physiol 21:1068-1070.

Guerrasio L, Quinet J, Büttner U, Goffart L (2010) Fastigial Oculomotor Region and the
Control of Foveation During Fixation. Journal of Neurophysiology 103:1988-2001.

Guillaume A (2012) Saccadic inhibition is accompanied by large and complex amplitude
modulations when induced by visual backward masking. Journal of Vision 12:5-5.

Hafed ZM (2013) Alteration of visual perception prior to microsaccades. Neuron 77:775-786.

Hafed ZM, Clark JJ (2002) Microsaccades as an overt measure of covert attention shifts.
Vision Research 42:2533-2545.

Hafed ZM, Krauzlis RJ (2010) Microsaccadic suppression of visual bursts in the primate
superior colliculus. J Neurosci 30:9542-9547.

Hafed ZM, Ignashchenkova A (2013) On the Dissociation between Microsaccade Rate and
Direction after Peripheral Cues: Microsaccadic Inhibition Revisited. Journal of
Neuroscience 33:16220-16235.

Hafed ZM, Goffart L, Krauzlis RJ (2009) A neural mechanism for microsaccade generation in
the primate superior colliculus. Science 323:940-943.

Hafed ZM, Lovejoy LP, Krauzlis RJ (2011) Modulation of Microsaccades in Monkey during a
Covert Visual Attention Task. Journal of Neuroscience 31:15219-15230.

Hafed ZM, Chen C-Y, Tian X (2015) Vision, Perception, and Attention through the Lens of
Microsaccades: Mechanisms and Implications. Frontiers in Systems Neuroscience
9:779.

Heinen SJ, Potapchuk E, Watamaniuk SNJ (2016) A foveal target increases catch-up saccade
frequency during smooth pursuit. Journal of Neurophysiology 115:1220-1227.

Heinen SJ, Badler JB, Watamaniuk SNJ (2018) Choosing a foveal goal recruits the saccadic
system during smooth pursuit. Journal of Neurophysiology 120:489-496.

Judge SJ, Richmond BJ, Chu FC (1980) Implantation of magnetic search coils for
measurement of eye position: an improved method. Vision Res 20:535-538.

Kerzel D, Born S, Souto D (2010) Inhibition of steady-state smooth pursuit and catch-up
saccades by abrupt visual and auditory onsets. J Neurophysiol 104:2573-2585.

Kleiner M, Brainard D, Pelli DG (2007) What's new in Psychtoolbox-3? (Abstract). Perception
36.

Ko H-k, Poletti M, Rucci M (2010) Microsaccades precisely relocate gaze in a high visual
acuity task. Nature Neuroscience 13:1549-1553.

Kowler E, Blaser E (1995) The accuracy and precision of saccades to small and large targets.
Vision Research 35:1741-1754.

Krauzlis RJ (2004) Recasting the Smooth Pursuit Eye Movement System. Journal of
Neurophysiology 91:591-603.

Krauzlis RJ, Miles FA (1996a) Release of fixation for pursuit and saccades in humans:
evidence for shared inputs acting on different neural substrates. J Neurophysiol
76:2822-2833.

Krauzlis RJ, Miles FA (1996b) Decreases in the latency of smooth pursuit and saccadic eye
movements produced by the "gap paradigm" in the monkey. Vision Res 36:1973-
1985.

Lisberger SG, Westbrook LE (1985) Properties of visual inputs that initiate horizontal smooth
pursuit eye movements in monkeys. J Neurosci 5:1662-1673.

Martins AJ, Kowler E, Palmer C (1985) Smooth pursuit of small-amplitude sinusoidal motion.
Journal of the Optical Society of America A 2:234.

Missal M, Keller EL (2002) Common inhibitory mechanism for saccades and smooth-pursuit
eye movements. Journal of Neurophysiology.

Pelli DG (1997) The VideoToolbox software for visual psychophysics: transforming numbers
into movies. Spatial vision 10:437-442.

Rashbass C (1961) The relationship between saccadic and smooth tracking eye movements. J
Physiol 159:326-338.

Reingold EM, Stampe DM (1999) Saccadic Inhibition in Complex Visual Tasks. In, pp 249-255.
Boston, MA: Springer US.

Reingold EM, Stampe DM (2000) Saccadic Inhibition and Gaze Contingent Research
Paradigms. In, pp 119-145: Elsevier.

Reingold EM, Stampe DM (2002) Saccadic Inhibition in Voluntary and Reflexive Saccades.
Journal of cognitive neuroscience 14:371-388.

Rizzolatti G, Buchtel HA, Camarda R, Scandolara C (1980) Neurons with complex visual
properties in the superior colliculus of the macaque monkey. Experimental brain
research Experimentelle Hirnforschung Expérimentation cérébrale 38.

Rolfs M, Kliegl R, Engbert R (2008) Toward a model of microsaccade generation: The case of
microsaccadic inhibition. Journal of Vision 8:5-5.

Schmolesky MT, Wang Y, HANES DP, Thompson KG, Leutgeb S, SCHALL JD, Leventhal AG
(1998) Signal timing across the macaque visual system. Journal of Neurophysiology
79:3272-3278.

Schutz AC, Braun DI, Kerzel D, Gegenfurtner KR (2008) Improved visual sensitivity during
smooth pursuit eye movements. Nat Neurosci 11:1211-1216.

Spering M, Gegenfurtner KR, Kerzel D (2006) Distractor interference during smooth pursuit
eye movements. Journal of Experimental Psychology: Human Perception and
Performance 32:1136-1154.

Tian X, Yoshida M, Hafed ZM (2016) A Microsaccadic Account of Attentional Capture and
Inhibition of Return in Posner Cueing. Frontiers in Systems Neuroscience 10:428.

Tian X, Yoshida M, Hafed ZM (2018) Dynamics of fixational eye position and microsaccades
during spatial cueing: the case of express microsaccades. Journal of Neurophysiology
119:1962-1980.

Wurtz RH, Mohler CW (1976) Organization of monkey superior colliculus: enhanced visual
response of superficial layer cells. Journal of Neurophysiology 39:745-765.

Zuber BL, Stark L, Cook G (1965) Microsaccades and the velocity-amplitude relationship for
saccadic eye movements. Science 150:1459-1460.

Figure legends

Figure 1 Transient disruption of open-loop smooth pursuit initiation. (A) We plotted radial eye velocity from monkey M in Experiment 1 when no flashes were presented (gray) and when full flashes were presented 50 ms after target motion onset (light blue). There was a transient dip in eye acceleration relative to the gray curve starting ~50 ms after flash onset. (B) Difference in eye velocity between flash and no-flash trials from A. The flash-induced dip in the difference curve is clearly visible. (C, D) Similar to A, B but for flashes occurring 83 ms after target motion onset. In this case, the transient disruption in open-loop pursuit initiation was slightly delayed relative to A, B because of the slight delay in flash onset time. (E, F) Same for flashes 117 ms after target motion onset. (G, H) Same for flashes after 150 ms from target motion onset. (I) In all cases, and for both monkeys, the disruption was always aligned on flash onset. We plotted eye velocity difference curves as in B, but now aligning the data on flash onset. The different colors denote different flash onset times relative to target motion onset. Regardless of flash time, the velocity difference curves always showed a dip ~50 ms after flash onset. A series of one sample t-tests against zero showed p<0.05 (Bonferroni corrected) for all flash times for both monkeys (with the exception of flash time 83.33 in monkey A). Error bars in all panels denote 95% confidence intervals.

Figure 2 Transient disruption of sustained smooth pursuit. (A) We repeated analyses similar to Fig. 1 but during sustained smooth pursuit from Experiment 3. In monkey A, we plotted radial eye velocity with no flashes were presented (gray) or after full flash onset (light blue). There was a transient reduction on smooth eye velocity ~50 ms after flash onset. (B) Same for monkey M. Error bars denote 95% confidence intervals. Note that monkey A had lower sustained pursuit eye velocity than monkey

1010 M (compare gray curves in both panels). This monkey relied more on catch-up

1011 saccades, as will be evidenced by the catch-up saccade rates discussed later (Fig 9.

1012 C, D).

1013

Figure 3 Directionally-specific modulation of open-loop smooth pursuit initiation by

injected transient eye position error signals. (A, B) Example velocity (A) and velocity

difference (B) traces from monkey M in the same format as in Fig. 1A, B. A flash was

presented 2.1 deg ahead of the pursuit target only 50 ms after target motion onset.

Because pursuit had not yet started at flash onset and because of the geometry of

flash location, the flash was still behind gaze position when it occurred (see text).

This caused a transient dip as in Fig. 1A, B. (C, D) Similar analyses for a flash behind

the pursuit target, again causing a transient dip as in A, B. (E, F) When the forward

flash happened 150 ms after flash onset, it was now ahead of instantaneous gaze

position, and now caused enhancement rather than inhibition of pursuit initiation. (G,

H) A backward flash at the same time caused inhibition. Therefore, pursuit

disruptions are not generalized inhibitory effects, but they reflect spatially-specific

influences of flash position relative to current gaze position. Error bars denote 95%

confidence intervals.

1028

Figure 4 Spatially-specific transient modulations of open-loop smooth pursuit

initiation depending on flash location. (A, B) For forward and backward flashes 2.1

deg ahead of or behind the pursuit target, we confirmed the observations in Fig. 3 in

both monkeys. Very early flashes caused inhibition and later ones caused

enhancement in smooth pursuit velocity/acceleration. For each flash time, we

measured the eye velocity difference in an appropriate interval after flash onset

(Materials and Methods), and we plotted this measurement as a function of flash

onset time relative to target motion onset. For reference, smooth pursuit initiation

typically happens at ~100 ms, meaning that the measurements shown all illustrate

modifications of open-loop smooth pursuit initiation. (C) When the flash was at 6 deg

instead of 2.1 deg, enhancement occurred even for the very earliest flashes,

supporting the conclusion that the early inhibition in A, B was not a generalized

inhibitory effect, but that it depended on instantaneous flash location relative to

instantaneous gaze location. Error bars denote 95% confidence intervals.

1043

Figure 5 Spatially-specific modulations of sustained smooth pursuit eye movements.

(A, b) In both monkeys, we repeated analyses similar to Fig. 2, but for localized

flashes ahead of or behind the pursuit target (by 2.1 deg). The gray curve shows

sustained pursuit velocity without flashes. The colored curves show velocity

modulations with short latency after flash onset. Forward flashes caused enhanced

smooth eye velocity followed by reduction. Backward flashes primarily caused only

reduction in eye velocity. Note that the reduction in eye velocity after the

enhancement in the forward flash conditions may reflect that the enhancement

caused pursuit to overpass the target. Subsequent slowing of the eye could return

gaze to its normal position relative to the target when no flashes take place. Error

bars denote 95% confidence intervals.

1055

Figure 6 Modulations in catch-up saccades during sustained smooth pursuit by flash

onsets. (A) Catch-up saccade rate as a function of time from flash onset for forward

and backward flashes. A characteristic inhibition followed by rebound was obvious,

and it was very similar to microsaccadic modulations during fixation (e.g. see Fig. 8A,

B). Note that at the onset of inhibition, the forward flash condition was less effective

in inhibiting saccades as the backward flash condition (also see G for an even

stronger effect in monkey M). These few saccades that did occur, were often

dramatically altered by the flash. (B) For example, aligning eye position on catch-up

saccade onset for these specific post-flash saccades reveals that they were larger

(pink) than normal catch-up saccades (gray) when a forward flash was presented.

(C) Their radial eye velocity was also higher than when no flash occurred. (D, E)

Backward flashes caused minimal change in saccade amplitude. (F) When we

checked saccade kinematics for the specific post-flash saccades occurring near the

onset of the inhibition phase in A, we found that the larger saccades in B violated the

main sequence curve of peak velocity versus amplitude for saccades: relative to

baseline (gray curve), their peak velocity did not scale as much as the scaling of

amplitude. This is consistent with microsaccadic effects in the literature (also see Fig.

8C, D). (G-L) Similar analyses for monkey M. Also in this case, forward flashes

affected catch-up saccade amplitude (two-sample t-tests: $t = 3.701$; $df = 1742$; $p =$

0.0002) (H, J), and backward flashes made catch-up saccades violate the main

sequence curve (two-sample t-tests: $t = 1.884$; $df = 1683$; $p = 0.0598$)

(green in L). Error bars denote 95% confidence intervals.

1078

Figure 7 Alterations of catch-up saccade kinematics in pursuit initiation with

saccades (Experiment 2). (A, B) Catch-up saccade amplitudes for post-flash

saccades were larger than in no flash trials (gray curve) when a forward flash was

presented but smaller when a backward flash was presented. (C, D) Similar to Fig. 6,

the changes in saccade amplitudes were often associated with altered kinematics

(Buonocore et al., 2016). For example, monkey A had lower peak velocity than

expected for forward flashes, similar to Fig. 6, and monkey M had higher peak

velocity than expected for backward flashes. Thus, kinematic alterations occur during

sustained pursuit (Fig. 6) and also during pursuit initiation (this figure). Error bars

denote 95% confidence intervals.

1089

Figure 8 Analyses similar to Figs. 6-7 for microsaccades during gaze fixation. (A, B)

Microsaccade rate was robustly affected by flash onset, as expected, and very

similarly to the catch-up saccade effects during sustained pursuit. (C-F)

Microsaccade amplitude and kinematic effects were also altered, but in an even

stronger manner than during smooth pursuit. These analyses all replicate our

previous study (Buonocore et al., 2017). Error bars denote 95% confidence intervals.

1096

Figure 9 Similar temporal structure of catch-up saccades to microsaccades despite

significantly larger eye position errros during pursuit. (A, B) We plotted inter-saccadic

intervals during pursuit or fixation. The distributions looked very similar in both tasks,

although monkey A often had very rapid back-to-back microsaccades with 100 ms

inter-saccadic interval. (C, D) Microsaccade and catch-up saccade rates were also

comparable in general. Monkey A made more catch-up saccades because of the

lower pursuit eye velocity (Fig. 2). (E, F) Despite similarities in inter-saccadic

intervals between the tasks, pursuit was always associated with larger eye position

errors at saccade onset than fixation. Error bars denote 95% confidence intervals.

1106

1107

1108   Figure 1

1109

1110

1111

1112

1113 Figure 2

1114

1115

1116

1117

 Figure 3

1123   Figure 4

1124

1125

1126

1127

1128   Figure 5

1129

1130

1131

1132

Figure 6

1138   Figure 7

1139

1140

1141

1142

1143　　Figure 8

1144

1145

1146

1147

1148

1149

 Figure 9