1

Life-cycle mediated effects of urbanization on parasite communities in the estuarine fish, *Fundulus heteroclitus*

James M. Alfieri[1] and Tavis K. Anderson[2]

[1] Ecology and Evolutionary Biology, Texas A&M University, College Station, TX, United States
of America
[2] Virus and Prion Research Unit, National Animal Disease Center, USDA - ARS, Ames, IA,
United States of America

* Corresponding author
E-mail: JamesMAlfieri@gmail.com

Abstract

This study examined the relationship between urbanization and parasite community structure in the estuarine fish, *Fundulus heteroclitus*. We measured landscape and physicochemical factors associated with urbanization at 6 sites from 4 collection periods. Concurrently, we quantified the metazoan parasite community in *F. heteroclitus* collected at those sites, with 105 fish studied per site during the 4 collection periods. Parasite community composition differed between sites. Variation in the prevalence and intensity of infection of two indirect life-cycle parasites, *Lasiocotus minutus* and *Glossocercus caribaensis*, were the primary parasite species that determined this pattern. Sediment potassium and aquatic osmium were the most important physicochemical factors in structuring parasite communities, and habitat dominance was the most important landscape factor. Our data supports the hypothesis that urbanization, acting at both landscape and physicochemical scales, can have a significant impact on parasite community structure. This, however, varied by parasite life history: there was little effect of urbanization on the prevalence and intensity of direct life-cycle parasites, but significant variation was dedicated for indirect life-cycle parasites. This study demonstrates how anthropogenically driven landscape change influences fine-scale parasite population dynamics.

Introduction

Urbanization significantly alters estuarine environments through landscape composition changes, habitat fragmentation, and contamination (Paul and Meyer, 2001; Lowe and Peterson, 2014; Lerberg et al. 2000). These anthropogenic modifications can affect the diversity and population dynamics of free-living animals and their associated parasite communities (Lotze et al., 2006; Hechinger et al. 2007). Individual parasite species may be affected by urbanization

through direct exposure to the abiotic environment (e.g. contaminants and altered

physicochemical parameters) (Anderson and Sukhdeo, 2010; Blanar et al. 2011) or through

changes in host population dynamics (Anderson and Sukhdeo, 2013a). Parasite species co-

occurring in a host species – the parasite community – simultaneously use the same host

resource, but each parasite species may have a different life history strategy. Notably, indirect

life-cycle parasites have life stages that are exposed to the external environment and are

dependent upon trophic interactions; whereas direct life-cycle parasite transmission largely does

not involve trophic interactions. Consequently, urbanization may differentially affect parasite

species but data that associates urbanization with parasite communities and life-history strategy

are sparse (but see Calegaro-Marques 2014 and Blanar et al. 2016).

Parasite life-cycles can be broadly characterized as either indirect or direct. Indirect life-

cycle parasites require different species of hosts to complete their life-cycle and are largely

dependent on trophic interactions which occur more frequently in locally stable communities

(Anderson and Sukhdeo, 2013a). This essential interaction has been used to demonstrate how

parasites can be indicators of free-living diversity (Hechinger et al., 2007), and as such, it has

been argued that a healthy ecosystem is one that is rich in parasites (Hudson et al., 2006). In

urbanized estuaries, host population dynamics may be less stable, resulting in reduced diversity

or less predictable host dynamics, causing a concomitant decrease in the richness and abundance

of indirect life-cycle parasites (Anderson and Sukhdeo, 2013b). On the other hand, direct life-

cycle parasites require only one species of host to complete their life-cycle and are largely

dependent on host population densities, as abundant hosts increase the probability of a parasite

encountering a suitable environment to colonize (Arneberg et al., 1998). A feature of urbanized

estuaries is habitat fragmentation leading to small patch sizes, which can result in increased host

densities, causing a concomitant increase in the richness and abundance of direct life-cycle

parasites (Wang and Moskovits, 2001; Hussain et al. 2013).

Many of the studies that examine the interaction between host and parasite in urbanized

ecosystems have been conducted with human pathogens (Bradley and Altizer, 2007), consider a

single host and a single parasite (Johnson et al., 2009), or a single contaminant and landscape

factor. Further, linking the regional process of urbanization to local transmission and diversity of

parasite communities has yet to reveal consistent patterns (Schotthoefer et al., 2011; Chapman et

al., 2015; Blanar et al., 2016; Johnson et al., 2016). One potential explanation for the absence of

consistent results is that urbanization differentially affects parasite species: the size and direction

of effect is contingent upon parasite life history. In this study, we assessed the effect of multiple

landscape composition and physicochemical factors on 24 parasite communities of the salt marsh

fish, *Fundulus heteroclitus*. The parasite community in *F. heteroclitus* includes indirect and

direct life-cycle species, and the impact of urbanization on the structure parasite communities

should be mediated through different life-cycle strategies. We predicted that urbanized sites

would have low diversity and prevalence of complex life-cycle parasites; but direct life-cycle

parasites would increase in diversity and abundance.

Materials and methods

Study site description

We examined 6 sites along the coastline of Georgia, USA, that reflected variation in

urbanization, measured through % impervious surface within a 250 m radius around each site

(Fig 1). We used this metric to select sites because impervious surface is a strong predictor of

different effects of urbanization, such as landscape composition changes, habitat fragmentation,

and contaminants (Paul and Meyer, 2001; Lerberg et al. 2000). The 250 m radius around each

site was chosen because this is similar to the home range size of the focal host organism,

Fundulus heteroclitus (Lotrich, 1975; Teo and Able, 2003). The 250 m radius criteria was used

as an objective measure to select sites representing a range of urbanization and its effects,

however, for the landscape composition analyses, we used a larger spatial scale at the watershed

level that is described below.

94   Quantifying the landscape and physicochemical environment

We defined the boundaries of the landscape to the lowest level watershed provided by the United

States Geological Survey (USGS) Watershed Boundary Dataset (WBD): the hydrological unit

code (HUC) 12-digit boundary (Lotrich, 1975). We used the 2011 National Land Cover Dataset

(NLCD) and National Oceanic and Atmospheric Administration's (NOAA) Coastal Change

Analysis Program (C-CAP) for the regional land cover analysis. Regional landscape composition

and change were measured using ArcGIS (ArcGIS Version 10.3, Environmental Systems

Research Institute, Redlands, California). We also calculated a series of regional landscape

metrics: patch density, largest patch index, edge density, mean salt marsh size, % salt marsh, %

developed land, and total area using FRAGSTATS software (FRAGSTATS, version 4.2,

University of Massachusetts, Amherst, Massachusetts). Patch density is an index that expresses

the number of continuous habitat types, or patches, per unit area, and is a metric for landscape

heterogeneity. Largest patch index is the size of the largest habitat patch relative to the size of the

landscape, and is a measure of landscape dominance. Edge density is an index of the amount of

habitat edges per unit area, and the contrast between developed land and salt marsh was weighted

at 0.5 to evaluate salt marsh habitat fragmentation by urban structures or landscapes. Mean salt

marsh size is an index of the size of a salt marsh habitat patch, with larger patches weighted in

order to decrease the sensitivity of the mean to small habitat patch sizes. The % salt marsh and

developed land is the percentage of the landscape comprised of those categories, and is a metric

for habitat size and dominance. Total area refers to the area of the watershed. In addition, we

used the National Oceanic and Atmospheric Administration's (NOAA) Coastal Change Analysis

Program (C-CAP) to evaluate % landscape change within the HUC-12 digit watersheds for the

period of time from 1996 – 2010.

We collected information on the physicochemical environment from each site at flow and

ebb tide in March, June, September 2015 and February 2016. Flow and ebb tide measurements

were averaged for each site and time point for analysis. We measured pH, temperature, salinity,

and conductivity using a handheld multiparamenter instrument (Yellow Springs Instruments

Professional Plus; Yellow Springs Instruments, Yellow Springs, Ohio). In addition, we

quantified the trace metal composition, nutrient concentrations, and chlorophyll *a* concentration

during these dates. To extract chlorophyll *a*, we followed a standard acidification method (EPA

445.0), whereby a volume of 100 ml was filtered using Whatman GF/F filters (47 mm; 0.7 μm

nominal pore size). Phytoplankton pigments were then extracted from the filters in 8 ml of 90 %

acetone at 0 ℃ for 24 hours. Fluorescence was measured using a Trilogy fluorometer (Turner

Designs, San Jose, California). For the nutrient and trace metal quantification, 2 samples from

each collection time and site were sent to the University of Georgia Laboratory for

Environmental Analysis (Athens, Georgia). These samples were filtered using a microwave-

assisted nitric acid method (EPA 3015), and total nitrogen, total phosphorus, and 72 metals were

analyzed with inductively coupled plasma mass spectrometry (ELAN 9000; PerkinElmer,

Waltham, Massachusetts). In addition to water analyses, we quantified the trace metal

composition of sediment by collecting 3 - 8 cm core samples from randomized locations within

each site in September 2015. These samples were sent to the University of Georgia Laboratory

for Environmental Analysis (Athens, Georgia), and were prepared using a microwave-assisted

nitric acid method (EPA 3052).

Quantifying the parasite community in *Fundulus heteroclitus*

We used the parasite community of a focal species, the common marsh killifish *Fundulus*

heteroclitus, to assess the impact of human modification of the environment on parasite

establishment and richness. The host community (benthic invertebrates, birds, fishes) transmits at

least 22 species of helminth parasites to the common killifish (Harris and Vogelbein 2006). At

each timepoint, 5 baited minnow traps were randomly placed at every site, during the flood tide

and collected during the ebb tide. We combined all 5 traps, and randomly subsampled to 30

killifish to obtain a representative population of each site and date combination. The fish were

separately housed in the animal facility at Georgia Southern University and necropsied within 1

week (IACUC #I14013).

Each fish was euthanized in a buffered 300 mg L^{-1} solution of tricaine methanesulfonate

(MS-222) until cessation of opercula movement followed by sex determination and

measurements of total length and weight. The MS-222 solution, all external surfaces, gills, and

opercula were examined for ectoparasites and helminths. The viscera, gonads, spleen, liver,

heart, intestine, and bladder were removed and examined for helminths. All helminths from the

first collection were heat fixed, stored in 70% ethanol, stained in acetocarmine, and mounted in

Permount. Parasites from all collections were identified with keys and primary literature (Khalil

et al., 1994; Harris and Vogelbein, 2006; Bray et al., 2008; Anderson et al., 2009).

Statistical Analyses

156 To visualize the variation in parasite component community structure, we performed non-

157 metric multi-dimensional scaling (NMDS). In the NMDS, average abundances of parasite

158 species in each component community were calculated and square root transformed to control

159 for the effect of high abundance parasite taxa. A resemblance matrix was constructed using

160 Bray-Curtis dissimilarity distance, and stress was calculated using Kruskal's stress formula 1. To

161 determine whether there were significant differences in parasite component community structure,

162 we performed a permutational multivariate ANOVA (PerMANOVA). For this analysis, we used

163 site as a factor and performed 999 permutations. Additionally, we performed a 2-way crossed

164 similarity percentage (SIMPER) analysis to identify the parasite species that contributed to 5%

165 or greater of the variation in parasite community structure between sites (PRIMER-E LTD.,

166 Plymouth, U.K.).

167  To determine the role of landscape composition and physicochemical factors in

168 determining parasite community composition, we used a mulitivariate regression tree using R

169 packages rpart and mvpart (R Core Development Team, 2016; De'ath 2004; Therneau et al.

170 2019). Regression tree analysis is beneficial for determining the effects of urbanization on a

171 response variable because the method does not assume independence between samples, and

172 makes no *a priori* assumptions on the relationship between the response and predictor variables.

173 Prior to this analysis, we conducted a variable reduction process to simplify the 48 water and 65

174 sediment trace metal and nutrient concentration data (JMP Pro 12.0.0, SAS Institute Inc). To do

175 this, we generated a correlation matrix of element concentrations, placed all variables in one

176 cluster, and then conducted binary splits based on the magnitude of the second eigenvalues.

177 Variables were iteratively reassigned to clusters, and cluster membership was finalized when the

178 squared correlation to the first principal component was highest. In each cluster, the variable that

179 explained the most variation was used in the subsequent regression tree analysis. In the

180 regression tree analysis, the multivariate response variable was the average abundances of the

181 parasite species and the predictor variables were the landscape metrics and representative

182 physicochemical factors.

Results

Parasite community in *Fundulus heteroclitus*

Eight taxa of metazoan parasites were identified from 630 fish, representing 24 different

parasite communities. The direct life-cycle parasites included the branchiuran *Argulus funduli*,

the copepod *Ergasilus funduli*, the leech *Myzobdella lugubris,* and the monogeneans *Swingleus*

ancistrus and *Fundulotrema prolongis*. The indirect life-cycle parasites included the digenean

Lasiotocus minutus, the cestode *Glossocercus caribaensis*, and the nematode genera

Contracaecum sp. These 8 taxa of parasites infected more than 70% of the killifish examined

(Tables 1 and 2).

Table 1. Prevalence (%) and mean intensity (SE) of the 3 complex life cycle parasite taxa

infecting the salt marsh fish, *Fundulus heteroclitus*. The sites include: Fort Pulaski (FP),

Cockspur Island (CI), Tybee (TB), and Skidaway Island (SK) in the Savannah River Estuary;

Shellman Bluff (SB) as a representative of the Sapelo Sound; and St. Marys (SM) as part of the

St. Marys Estuary. The sites are presented from top to bottom to reflect increasing levels of

impervious surface.

	Lasciotocus minutus		*Glossocercus caribaensis*		*Contracaecum* sp.	
	Prevalence	Intensity	Prevalence	Intensity	Prevalence	Intensity

SK	26.7	14.0 ± 2.9	44.8	4.0 ± 0.6	1.0	1.0 ± 0.0
SB	24.8	24.6 ± 4.4	39.0	3.9 ± 0.5	7.6	1.0 ± 0.0
FP	22.9	9.5 ± 1.7	448	4.2 ± 0.5	13.3	1.0 ± 0.0
CI	2.9	7.3 ± 2.4	35.2	3.4 ± 0.6	5.7	1.0 ± 0.0
SM	1.9	4.0 ± 3.0	91.4	6.0 ± 0.5	6.7	1.0 ± 0.0
TB	0	-	30.5	2.2 ± 0.3	3.8	1.0 ± 0.0

Table 2. Prevalence and mean intensity (SE) of the 5 direct life cycle parasite taxa infecting the salt marsh fish, *Fundulus heteroclitus*. The sites include: Fort Pulaski (FP), Cockspur Island (CI), Tybee (TB), and Skidaway Island (SK) in the Savannah River Estuary; Shellman Bluff (SB) as a representative of the Sapelo Sound; and St. Marys (SM) as part of the St. Marys Estuary. The sites are presented from top to bottom to reflect increasing levels of impervious surface.

Myzobdella lugubris	*Swingleus ancistrus*	*Fundulotrema prolongis*	*Ergasilus funduli*	*Argulus funduli*

	Prevalence	Intensity	Prevalence	Intensity	Prevalence	Intensity	Prevalence	Intensity	Prevalence	Intensity
SK	0	-	40.0	2.8 ± 0.4	29.5	4.6 ± 1.0	24.8	3.4 ± 0.7	15.2	1.4 ± 0.2
SB	1.9	1.0 ± 0	36.2	3.1 ± 0.3	20.0	4.2 ± 0.9	19.0	3.8 ± 0.7	20.0	1.4 ± 0.2
FP	9.5	1.2 ± 0.2	38.1	4.5 ± 0.7	29.5	6.0 ± 1.1	14.3	3.6 ± 0.7	24.8	1.4 ± 0.1
CI	0	-	8.6	2.0 ± 0.5	16.2	3.6 ± 0.8	20.0	3.9 ± 1.1	7.6	1.1 ± 0.2
SM	0	-	13.3	2.7 ± 0.5	20.0	3.6 ± 0.9	22.9	3.4 ± 0.9	12.4	1.2 ± 0.2
TB	0	-	16.2	1.4 ± 0.3	19.0	3.5 ± 0.5	11.4	3.1 ± 0.5	10.5	1.3 ± 0.1

Assessed with NMDS and PerMANOVA, the parasite community composition differed

between sites ($P < 0.001$, Pseudo $F_{5,12} = 6.11$; Fig 2). SIMPER analysis indicated that the

observed differences in parasite community structure were driven by indirect life-cycle parasites:

the digenean *Lasciotocus minutus*, and the cestode *Glossocercus caribaensis*. This analysis

demonstrated that direct life-cycle parasites did not contribute to the variation in parasite

community structure.

Landscape and physicochemical predictors of parasite community

in *Fundulus heteroclitus*

The cluster analysis variable reduction process on the water and sediment trace metal and

nutrient concentrations resulted in 16 clusters, explaining 83.3% of the variance. We constructed

a multivariate regression tree that designated each of the 24 parasite community observations (6

sites, 4 collection periods) to terminal leaf nodes: these terminal leaf nodes categorize the factors,

where the factors are all the collected landscape and physicochemical variables collected from

the 6 sites across 4 collection periods that predict the observed parasite community structure. In

multivariate regression tree analysis, variable importance is determined as the reduction in the

loss function attributed to each variable at each split summed across the full tree. The optimum

tree size contained 3 leaves, and the primary variable that was most important in determining the

parasite community structure was the potassium concentrations in the sediment, with largest

patch index (LPI) and aquatic osmium concentrations as secondary predictor variables (CV Error

= 0.5608). Parasite communities that were dominated by the indirect life-cycle cestode

Glossocercus caribaensis were characterized by sediment potassium levels with a mean of 4063

+/- 270 mg/L, and an LPI of less than 22. Parasite communities that were dominated by the

indirect life-cycle digenean *Lasiotocus minutus* were characterized by sediment potassium levels

with a mean of 4706 +/- 468 mg/L and an osmium concentration of greater than 0.01 mg/L.

Discussion

The goal of this study was to determine the effects of urbanization, mediated through

alterations in landscape composition and physicochemical factors, in structuring parasite

communities. We determined that parasite community composition differed between sites with

different levels of urbanization. Variation in the prevalence and intensity of infection by indirect

life-cycle parasites, the digenean *Lasiotocus minutus* and the cestode *Glossocercus caribaensis*

were the primary drivers for the differences in parasite communities. Although the prevalence of

direct life-cycle parasites differed among sites, this difference did not contribute to the overall

variation in community composition. Additionally, we found that the landscape composition and

physicochemical factors that best predicted the parasite communities were potassium sediment

concentration, largest patch index, and aquatic osmium concentration. These data support the

prediction that in the common marsh killifish, parasite community composition is explained by

metrics associated with human modification of the environment, and parasite responses to

urbanization are dependent upon life history.

Our results indicated that sediment potassium, water osmium, and largest patch index

were the most important factors in determining the structure of parasite communities. Although

potassium is not necessarily a contaminant, the concentrations we observed have been reported

to significantly affect estuarine communities (Ajitthamol et al. 2016). The upstream watersheds

of the high potassium sites are dominated by cotton and soy agriculture, which are especially

dependent on potassium fertilizers. The other major contaminant we found to structure parasite

communities is the rare element osmium. The concentrations that we observed suggest extreme

point source pollution (Turekian et al. 2007). Taken together, these data suggest that our sites

may have contamination that impacts the survival of benthic organisms, which may subsequently

alter trophic interactions and parasite transmission that relies upon those interactions. The major

landscape factor in determining parasite community composition, LPI, is a simple measure of

landscape "dominance." LPI typically decreases as distance from urban centers increase (Nong

et al. 2018). We found an LPI value of 22 to best predict parasite community structure, and this

value was also associated with lower sediment and aquatic contaminant levels. This metric

suggests that urbanization is a major driver of parasite community variation, whereby areas with

lower levels of urbanization likely support more diverse and stable community of organisms that

allow the completion of indirect parasite life-cycles.

Previous studies have shown a relationship between urbanization and parasite community

variation. Blanar et al. (2016) demonstrated a correlation between parasite community structure

and local contaminants and land use within 5 km. In this study, the relative abundance of the

indirect life-cycle digenean was positively associated with an increase in crude oil contaminants,

whereas the opposite pattern was observed for the directly transmitted monogenean parasites.

This observation follows results of an earlier meta-analysis, where parasite life history (>1

obligatory host in life-cycle vs. 1 obligatory host in life-cycle) and habitat (external vs. internal)

informed whether aquatic pollution had a significant effect on parasite population dynamics

(Blanar et al., 2009). Additionally, Calegaro-Marques and Amato (2014) revealed an association

between an urban-rural gradient and life-cycle variation in parasite communities of Rufous-

bellied thrushes. Our results concur with these studies, and support the proposition that

urbanization disproportionately affects indirect life-cycle parasites, which is then reflected in the

parasite community structure.

We found that sites within urbanized watersheds contained a greater prevalence and

intensity of the autogenic (whole life-cycle is aquatic) parasite *Lasiotocus minutus,* a digenean

that uses the focal fish species, *Fundulus heteroclitus,* as the definitive host, molluscs as the first

intermediate host, and fish or crustaceans as the second intermediate host (Stunkard, 1980). This

pattern was unexpected, as this parasite and associated hosts are continuously exposed to the

external aquatic environment, and would be exposed to any contaminants or deleterious

physicochemical parameters. However, the most important physicochemical parameter in

structuring the parasite communities, sediment potassium, may alter trophic structures in a

favorable way to the host communities. For example, potassium has been shown to increase

phytoplankton abundance in estuaries. This increase in basal resource would disproportionately

affect benthic grazers, such as the 1st intermediate hosts for *L. minutus*. The second complex

life-cycle parasite, the cestode *G. caribaensis*, is allogenic (life-cycle is aquatic and terrestrial),

and did not follow the same pattern as *L. minutus*, and was more dominant in sites associated

with less impacts of urbanization. This allogenic cestode has fish-eating birds as definitive hosts

and infects a number of fish intermediate hosts (Scholz et al., 2004). This parasite may appear to

be sensitive to urbanization because its definitive host is vagile and can select habitat outside of

urbanized salt marshes. Consequently, the difference in prevalence across sites may be due to the

effect of the abiotic conditions on parasite life stages external to the host, or through effects on

critical aquatic life-cycle hosts, such as the mollusc and bird (Anderson and Sukhdeo, 2010).

Future research should include population size assessments of intermediate and definitive hosts

to better disentangle the direct effects of urbanization (through exposure to the environment) and

indirect effects (through population dynamics of the hosts).

Coastal urbanization and anthropogenic disturbance are rapidly altering free-living and

parasite community dynamics. Our results suggest that urbanization and its effects are important

in structuring parasite communities of the common marsh killifish, complex life-cycle parasites

are more affected than direct life-cycle parasites, and the degree to which individual parasite

species respond to urbanization is determined by their specific life history strategy. Further, our

data suggest that urbanization likely affects interactions of free-living hosts, and their

environment which is then detected as variation in the parasite community.

Acknowledgements

We acknowledge the advice and support provided by Dr. Clark Alexander and Mike Robinson at

the Applied Coastal Research Laboratory (Georgia Southern University), and Dr. Sayed Hassad

at the Laboratory of Environmental Analysis (University of Georgia). Field assistance was

provided by Emily Dodd, Sarah Dunn, Maria St. Jean, and Jackson Tomlinson. Comments on an

earlier draft by Drs. Risa Cohen, James Roberts, and Oscar Pung improved this manuscript. This

research was funded in part by a Research Assistantship from the Institute of Coastal Plain

Science (Georgia Southern University) to JMA. TKA was supported by the Office of Research

Services and Sponsored Programs at Georgia Southern University (GSU) and by an appointment

to the USDA-ARS Research Participation Program administered by the Oak Ridge Institute for

Science and Education (ORISE) through an interagency agreement between the U.S. Department

of Energy (DOE) and USDA under contract number DE-AC05-06OR23100. The funders had no

role in study design, data collection and interpretation, or the decision to submit the work for

publication. Mention of trade names or commercial products in this article is solely for the

purpose of providing specific information and does not imply recommendation or endorsement

by the USDA. USDA is an equal opportunity provider and employer.

References

Paul, M. J., and J. L. Meyer. 2001. Streams in the urban landscape. Annual Review of Ecology

and Systematics 32:333-365.

Lowe, M. R., and M. S. Peterson. 2014. Effects of Coastal Urbanization on Salt-Marsh Faunal

Assemblages in the Northern Gulf of Mexico. Marine and Coastal Fisheries 6:89-107.

Lerberg, S. B., A. F. Holland, and D. M. Sanger. 2000. Responses of tidal creek macrobenthic

communities to the effects of watershed development. Estuaries 23:838-853.

Lotze, H. K., H. S. Lenihan, B. J. Bourque, R. H. Bradbury, R. G. Cooke, M. C. Kay, S. M.
Kidwell, M. X. Kirby, C. H. Peterson, and J. B. C. Jackson. 2006. Depletion, degradation,
and recovery potential of estuaries and coastal seas. Science 312:1806-1809.

Hechinger, R. F., K. D. Lafferty, T. C. Huspeni, A. J. Brooks, and A. M. Kuris. 2007. Can
parasites be indicators of free-living diversity? Relationships between species richness
and the abundance of larval trematodes and of local benthos and fishes. Oecologia
151:82-92.

Anderson, T. K., and M. V. K. Sukhdeo. 2010. Abiotic versus biotic hierarchies in the assembly
of parasite populations. Parasitology 137:743-754.

Blanar, C. A., D. J. Marcogliese, and C. M. Couillard. 2011. Natural and anthropogenic factors
shape metazoan parasite community structure in mummichog (Fundulus heteroclitus)
from two estuaries in New Brunswick, Canada. Folia Parasitologica 58:240-248.

Anderson, T. K., and M. V. K. Sukhdeo. 2013a. Qualitative community stability determines
parasite establishment and richness in estuarine marshes. PeerJ 1:e92; DOI
10.7717/peerj.92.

Calegaro-Marques C, Amato SB (2014) Urbanization Breaks Up Host-Parasite Interactions: A
Case Study on Parasite Community Ecology of Rufous-Bellied Thrushes (Turdus
rufiventris) along a Rural-Urban Gradient. PLoS ONE 9(7): e103144.
https://doi.org/10.1371/journal.pone.0103144

Blanar, C. A., M. Hewitt, M. McMaster, J. Kirk, Z. Wang, W. Norwood, and D. Marcogliese.
2016. Parasite community similiarity in Athabasca River trout-perch (Percopsis
omiscomaycus) varies with local-scale land use and sediment hydrocarbons, but not
distance or linear gradients. Parasitology Research 115:3853-3866.

Hudson, P. J., A. P. Dobson, and K. D. Lafferty. 2006. Is a healthy ecosystem one that is rich in

parasites? Trends in Ecology & Evolution 21:381-385.

Anderson, T. K., and M. V. K. Sukhdeo. 2013b. The relationship between community species

richness and the richness of that parasite community in Fundulus heteroclitus. Journal of

Parasitology 99:391-396.

Arneberg, P., A. Skorping, B. Grenfell, and A. F. Read. 1998. Host densities as determinants of

abundance in parasite communities. Proceedings of the Royal Society B-Biological

Sciences 265:1283-1289.

Wang, Y. Q., and D. K. Moskovits. 2001. Tracking fragmentation of natural communities and

changes in land cover: Applications of landsat data for conservation in an urban

landscape (Chicago wilderness). Conservation Biology 15:835-843.

Bradley, C. A., and S. Altizer. 2007. Urbanization and the ecology of wildlife diseases. Trends in

Ecology & Evolution 22:95-102.

Johnson, D. S., J. W. Fleeger, and L. A. Deegan. 2009. Large-scale manipulations reveal that

top-down and bottom-up controls interact to alter habitat utilization by saltmarsh fauna.

Marine Ecology Progress Series 377:33-41.

Lotrich, V. A. 1975. Summer home range and movements of Fundulus heteroclitus (Pisces-

Cyprinodontidae) in a tidal creek. Ecology 56:191-198.

Teo, S. L. H., and K. W. Able. 2003. Habitat use and movement of the mummichog (Fundulus

heteroclitus) in a restored salt marsh. Estuaries 26:720-730.

Courtney E. Harris and Wolfgang K. Vogelbein. "Parasite of Mummichogs, Fundulus

heteroclitus, from the York River, Virginia, U.S.A., with a Checklist of Parasites of Atlantic

Coast Fundulus Spp," Comparative Parasitology 73(1), 72-110, (1 January 2006).

Khalil, L.F., Jones, A. & Bray, R.A. 1994. Order Cyclophyllidea (diagnosis and key to families).

Keys to the cestode parasites of vertebrates. CAB International, Wallingford, U.K." 305-307.

R Core Team (2019). R: A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria. URL

https://www.R-project.org/.

Ajithamol, A., Venkadesh, B., Michael Babu, M. and Saraswathy, S. 2016. "Impacts of nutrients

nitrogen, phosphorus, potassium and their role in the sediment of Manakudy estuary, Tamilnadu,

S. India", International Journal of Current Research, 8, (04), 29248-29251.

Nong DH, Lepczyk CA, Miura T, Fox JM (2018) Quantifying urban growth patterns in Hanoi

using landscape expansion modes and time series spatial metrics. PLoS ONE 13(5):

e0196940. https://doi.org/10.1371/journal.pone.0196940

Stunkard, H. W. 1980. Stages in the life-cycle of the digenetic trematode, Lasiotocus minutus

(Manter, 1931) Thomas, 1959. Biological Bulletin 159:497-498.

Scholz, T., R. A. Bray, R. Kuchta, and R. Repova. 2004. Larvae of gryporhynchid cestodes

(Cyclophyllidea) from fish: a review. Folia Parasitologica 51:131-152.

Fig 1. Map of the 6 salt marsh study sites in coastal Georgia, USA, including 2011 land cover

data. The sites include Fort Pulaski (FP), Cockspur Island (CI), Tybee (TB), and Skidaway

Island (SK) in the Savannah River Estuary; Shellman Bluff (SB) as a representative of the

Sapelo Sound; and St. Marys (SM) as part of the St. Marys Estuary. Red circles represent a 250

m buffer around the study site. For the purpose of this work, land cover data was reclassified

from the original 21 categories to 10 (Salt Marsh, Developed, Freshwater Wetland, Open Water,

Forest, Agriculture, Bare Land, and Grassland): 2 land cover categories were not present in the

study area (Freshwater Aquatic Bed and Unconsolidated Shore).

Fig 2. Non-metric multidimensional scaling plot of complex and direct life cycle parasite

component communities of the salt marsh fish, *Fundulus heteroclitus*, from 6 sites in coastal

Georgia, USA. Distances are based on Bray-Curtis dissimilarities of the square root transformed

mean abundance of 8 parasite taxa of *Fundulus heteroclitus*. The sites include: Fort Pulaski (FP)

(closed triangles), Cockspur Island (CI) (open triangles), Tybee (TB) (open squares), and

Skidaway Island (SK) (open circles) in the Savannah River Estuary; Shellman Bluff (SB) (closed

circles) as a representative of the Sapelo Sound; and St. Marys (SM) (closed squares) as part of

the St. Marys Estuary.