LOCAL AND REGIONAL URBANIZATION: CORRELATES OF PARASITE

COMMUNITY COMPOSITION AND RICHNESS IN THE SALT MARSH FISH,

FUNDULUS HETEROCLITUS

James M. Alfieri[1,*] and Tavis K. Anderson[1,*]

[1]Department of Biology, Georgia Southern University, P.O. Box 8042-1, Statesboro, Georgia

30460.

Corresponding author:

James M. Alfieri, Graduate Program in Ecology & Evolutionary Biology, Texas A&M

University, TAMU MS 2475, College Station, TX 77843. E-mail: jamesmalfieri@gmail.com

*Present address: James M. Alfieri, Graduate Program in Ecology & Evolutionary Biology,

Texas A&M University, TAMU MS 2475, College Station, TX 77843; Tavis K. Anderson, Virus

and Prion Research Unit, National Animal Disease Center, USDA-ARS, 1920 Dayton Ave,

Ames, Iowa 50010.

ABSTRACT: This study examined the relationship between regional urbanization and the local

transmission and establishment of parasites in the salt marsh fish, *Fundulus heteroclitus*. To

determine the relative importance of abiotic and biotic factors across local and regional scales, we

quantified the parasite community in *Fundulus heteroclitus* collected from 6 sites selected to

reflect a gradient of urbanization (determined by variation in impervious surface). A total of 630

fish was studied, with 105 collected per site in 4 collection periods from 2015-16: concurrently,

we measured local abiotic parameters and regional landscape descriptors. Trace metal

concentrations differed between sites, but the relationship between contaminant concentration and

urbanization was not consistent. Algal biomass, determined through measurement of chlorophyll

a, was significantly higher in urbanized sites, but there was not a corresponding increase in total

nitrogen or total phosphorus concentrations. Parasite component communities were different

between sites, with the pattern primarily associated with the presence or absence of a complex life-

cycle digenean, *Lasiotocus minutus*. The most consistent predictors of parasite infracommunity

richness within *F. heteroclitus* were host size, % regional salt marsh, and regional patch density.

Our data demonstrate that measures of regional urbanization do not have a consistent effect on the

local abiotic environment and that parasite infracommunity richness is best explained through the

consideration of regional abiotic and local biotic factors. This study demonstrates how

anthropogenically driven landscape change influences fine-scale parasite transmission and parasite

species richness.

INTRODUCTION

In areas associated with high levels of urbanization, there is significant alteration of estuarine habitat by fragmentation resulting in a mosaic of habitat sizes and shapes, and variation in the abiotic environment (Lowe and Peterson, 2014). This anthropogenic habitat modification has resulted in positive, negative, or neutral changes in the diversity and population dynamics of free-living communities (Lotze et al., 2006; Aronson et al., 2014). For parasites, transmission from one host to another involves life stages external to the host along with life stages forming the durable within-host association. Consequently, the effect of urbanization on parasite populations and communities may then be twofold; either directly through changes to, and variation in, the abiotic environment (e.g., availability of contaminants and altered physicochemical parameters, e.g., Anderson and Sukhdeo, 2010; Blanar et al., 2011; Pietrock and Marcogliese, 2003; Morley et al., 2003); or indirectly through modification of host population dynamics (e.g., Hechinger et al., 2007; Anderson and Sukhdeo, 2013a).

The mechanism by which regional urbanization affects local abiotic and biotic conditions, and potentially the transmission and establishment of parasites is likely mediated by the construction of impervious surfaces (Paul and Meyer, 2001). Impervious surfaces increase the volume and rate of terrestrial runoff following rain events, leading to more frequent introductions and higher concentrations of contaminants and freshwater into estuarine habitats, with a range of effects on free-living organisms (Sanger et al., 1999). For example, a regional landscape analysis by Lerberg et al. (2000) found that the watersheds that contained greater than 50% impervious surfaces were associated with higher concentrations of metals (Cu, Cr, Pb, Zn, Cd, and Hg) resulting in a decrease in the abundance and richness of crustaceans and bivalves. Hutchins et al. (2014) found a positive correlation between the amount of impervious surfaces in a regional

landscape and the concentration of nutrients in the water column, which can lead to declines in

fish abundance through habitat degradation, toxic algal blooms, and altered algal communities

(Johnson et al., 2009). In a study contrasting urban and undeveloped estuaries, Krebs et al. (2014)

demonstrated that urban estuaries had lower, variable salinities, resulting in altered fish community

structure. Additional studies have demonstrated how thermal effluent associated with industrial

discharge in estuaries alters fish growth rate and the fish community (Jones et al., 1996), and how

human development may alter pH, dissolved organic matter, and oxygen concentration affecting

foraminiferal faunas (Hayward et al., 2004). A common theme among these studies is that the

regional construction of impervious surfaces alters the local abiotic environment via increases in

contaminant amounts and variation in physicochemical parameters, generally resulting in

significant changes to the free-living community.

The free-living community, and changes in its diversity and dynamics due to urbanization,

is likely to affect the diversity and richness of parasites for two reasons. Firstly, the host is

exploited as an energy source and critical habitat, and over evolutionary time this has resulted in

parasite life strategies that are integrally coupled with host species dynamics. This essential

interaction has been used to demonstrate how parasites can be indicators of free-living diversity

(Hechinger et al., 2007), and it has been argued that a healthy ecosystem is one that is rich in

parasites (Hudson et al., 2006). Further, complex life cycle parasites are largely dependent upon

trophic interactions, which occur more frequently in locally stable communities (Anderson and

Sukhdeo, 2013a). In increasingly urbanized salt marshes, host population dynamics may be less

stable, resulting in reduced diversity or less predictable host dynamics, causing a concomitant

decrease in the richness of parasites (Anderson and Sukhdeo, 2013b). Concurrently, a feature of

increasingly urbanized salt marshes is habitat fragmentation, resulting in higher densities of hosts

in habitat patches (Wang and Moskovits, 2001). Higher host densities may lead to increased

abundance of directly transmitted parasites, as the probability of a parasite transmission stage

encountering a suitable host is increased (Arneberg et al., 1998).

Many of the studies that examine the interaction between host and parasite in urbanized

ecosystems have been conducted with human pathogens (Bradley and Altizer, 2007), on a local

spatial scale (Anderson and Sukhdeo, 2010; Blanar et al., 2011), or consider a single host and a

single parasite (Johnson et al., 2009). Further, linking the regional process of urbanization to local

transmission and diversity of parasite communities has yet to reveal consistent patterns

(Schotthoefer et al., 2011; Chapman et al., 2015; Blanar et al., 2016; Johnson et al., 2016). One

potential explanation for the absence of consistent results is that urbanization differentially affects

parasite species: the size and direction of effect is contingent upon parasite life history. For

example, Blanar et al. (2016) demonstrated a correlation between parasite community structure

and local contaminants and land use within 5 km. In this study, the relative abundance of the

complex life-cycle digenean was positively associated with an increase in crude oil contaminants,

whereas the opposite pattern was observed for the directly transmitted monogeneans. This

observation follows results of an earlier meta-analysis, where parasite life history (>1 obligatory

host in life cycle vs. 1 obligatory host in life cycle) and habitat (external vs. internal) informed

whether aquatic pollution had a significant effect on parasite population dynamics (Blanar et al.,

2009).

In this study, we assessed the effect of regional urbanization and the local abiotic

environment on the parasite community of the salt marsh fish, *Fundulus heteroclitus,* within 6 salt

marshes. We asked whether local factors, such as host size and the immediate abiotic environment,

or if regional landscape factors, such as land-use change and composition, were more important in

structuring the observed parasite community. We predicted that urban salt marshes would be

characterized by a high degree of regional habitat fragmentation and poor local abiotic conditions

(e.g., reduced water quality with nutrient enrichment and high contaminant concentrations).

Consequently, urbanized sites would have low diversity and prevalence of complex life cycle

parasites; but direct life cycle parasites would increase in diversity and abundance.

MATERIALS AND METHODS

Study site description

We examined three estuaries along the coastline of Georgia, USA: the Savannah River

Estuary, Sapelo Sound, and the St. Marys Estuary. The human population in the coastal counties

of Georgia increased by 62% between 1970 and 2000, and is expected to increase 51.2% over the

next 15 years (Ross and Barringer, 2006). The Savannah River Estuary experienced heavy

industrial growth during the first 60 years of the 1900s, resulting in high concentrations of

contaminants such as polychlorinated biphenyls, chlorinated pesticides, polycyclic aromatic

hydrocarbons (Alexander et al., 1999). In contrast, the landscape surrounding Sapelo Sound is

comprised mostly of agriculture and forested land, and there is relatively little urban development.

The landscape surrounding the St. Marys Estuary represents an intermediate between Savannah

River and Sapelo Sound; it is comprised of urban development, agriculture, and forest and there is

evidence of regular anthropogenic impact (Morrison et al., 2008).

From 2015-2016, we surveyed 6 salt marsh sites from within the three estuary systems that

reflected a gradient in urbanization, measured through % impervious surface within a 250 m radius

around each site (Figure 1). In the Savannah River Estuary, sites included Fort Pulaski (FP)

(32°1'9''N, 80°54'6''W), Cockspur Island (CI) (32°2'27''N, 80°51'1''W), Tybee (TB)

(32°0'23''N, 80°51'6''W), and Skidaway Island (SK) (31°58'32''N, 81°1'56''W). As a

representative site of the Sapelo Sound, Shellman Bluff (SB) (31°64'50''N, 81°22'3''W) was

included. St. Marys (SM) (30°43'19''N, 81°32'34''W) was chosen as it is representative of the St.

Marys Estuary.

Quantifying the regional and local environment

We defined the boundaries of the regional environment to the lowest level watershed

provided by the United States Geological Survey (USGS) Watershed Boundary Dataset (WBD):

the hydrological unit code (HUC) 12-digit boundary (Lotrich, 1975). We used the 2011 National

Land Cover Dataset (NLCD) and National Oceanic and Atmospheric Administration's (NOAA)

Coastal Change Analysis Program (C-CAP) for the regional land cover analysis. Regional

landscape composition and change were measured using ArcGIS (ArcGIS Version 10.3,

Environmental Systems Research Institute, Redlands, California). We also calculated a series of

regional landscape metrics: patch density, largest patch index, edge density, mean salt marsh size,

% salt marsh, % developed land, and total area using FRAGSTATS software (FRAGSTATS,

version 4.2, University of Massachusetts, Amherst, Massachusetts). Patch density is an index that

expresses the number of patches per unit area, and is a metric for landscape heterogeneity. Largest

patch index is the size of the largest habitat patch relative to the size of the landscape, and is a

measure of landscape dominance. Edge density is an index of the amount of habitat edges per unit

area, and the contrast between developed land and salt marsh was weighted at 0.5 to evaluate salt

marsh habitat fragmentation by urban structures or landscapes. Mean salt marsh size is an index

of the size of a salt marsh habitat patch, with larger patches weighted in order to decrease the

sensitivity of the mean to small habitat patch sizes. The % salt marsh and developed land is the

percentage of the landscape comprised of those categories, and is a metric for habitat size and

dominance. Total area refers to the area of the watershed. In addition, we used the National

Oceanic and Atmospheric Administration's (NOAA) Coastal Change Analysis Program (C-CAP)

to evaluate % landscape change within the HUC-12 digit watersheds for the period of time from

1996 – 2010.

We collected information on the local abiotic parameters from each site at flow and ebb

tide in March, June, September 2015 and February 2016. We measured pH, temperature, salinity,

and conductivity using a handheld multiparamenter instrument (Yellow Springs Instruments

Professional Plus; Yellow Springs Instruments, Yellow Springs, Ohio). In addition, we quantified

the trace metal composition, nutrient concentrations, and chlorophyll *a* concentrations during these

dates. To extract chlorophyll *a*, we followed a standard acidification method (EPA 445.0), whereby

a volume of 100 ml was filtered using Whatman GF/F filters (47 mm; 0.7 μm nominal pore size).

Phytoplankton pigments were then extracted from the filters in 8 ml of 90 % acetone at 0 °C for

24 hours. Fluorescence was measured using a Trilogy fluorometer (Turner Designs, San Jose,

California). For the nutrient and trace metal quantification, 2 samples from each collection time

and site were sent to the University of Georgia Laboratory for Environmental Analysis (Athens,

Georgia). These samples were filtered using a microwave-assisted nitric acid method (EPA 3015),

and total nitrogen (TN), total phosphorus (TP), and 72 metals were analyzed with inductively

coupled plasma mass spectrometry (ELAN 9000; PerkinElmer, Waltham, Massachusetts). In

addition to water analyses, we quantified the trace metal composition of sediment by collecting 3

- 8 cm core samples from randomized locations within each site in September 2015. These samples

were sent to the University of Georgia Laboratory for Environmental Analysis (Athens, Georgia),

and were prepared using a microwave-assisted nitric acid method (EPA 3052).

Quantifying the parasite community in Fundulus heteroclitus

We used the parasite community of a focal species, the common marsh killifish *Fundulus*

heteroclitus, to assess the impact of human modification of the environment on parasite

establishment and richness. The host community (benthic invertebrates, birds, fishes) transmits at

least 11 species of helminth parasites to the common killifish (Anderson and Sukhdeo, 2010; 2011;

2013b). In addition, the home range estimates of killifish are in concordance with our measure of

local urbanization (% impervious surface within a 250 m radius of each site) (Lotrich, 1975; Teo

and Able, 2003). At each site, killifish were collected using 5 minnow traps baited with dry dog

food. Traps were placed during the flood tide and collected during the ebb tide in March 2015,

with subsequent collections in June 2015, September 2015, and February 2016. We randomly

subsampled each collection to 30 killifish from each site on each collection date. The fish were

housed in the animal facility at Georgia Southern University and necropsied within 1 week

(IACUC #I14013).

Each fish was euthanized in a buffered 300 mg L^{-1} solution of tricaine methanesulfonate

(MS-222) until cessation of opercula movement followed by sex determination and measurements

of total length and weight. The MS-222 solution, all external surfaces, gills, and opercula were

examined for ectoparasites and helminths. The viscera, gonads, spleen, liver, heart, intestine, and

bladder were removed and examined for helminths. All helminths from the first collection were

heat fixed, stored in 70% ethanol, stained in acetocarmine, and mounted in Permount. Parasites

from all collections were identified with keys and primary literature (Khalil et al., 1994; Harris

and Vogelbein, 2006; Bray et al., 2008; Anderson et al., 2009).

Statistical analyses

To determine site-level differences in water pH, salinity, conductivity, water and sediment

trace metal concentration, we performed a series of one-way analysis of variance (ANOVA) tests,

using collection date as a blocking factor. All trace metal and nutrient concentration data were log

transformed to meet the assumptions of normality and homogeneity of variances. If a significant

difference was found ($P < 0.05$), a post-hoc multiple pairwise comparison using Tukey's honest

significance test was performed. To test for differences in parasite component community species

richness between sites, we conducted a Kruskal-Wallis one-way ANOVA (our data did not meet

the parametric assumptions of normality and equal variances). We followed this analysis with a

post-hoc multiple pairwise comparison using the Steel-Dwass test (JMP Pro 12.0.0, SAS Institute

Inc., Cary, NC).

To visualize the variation in parasite component community structure, we performed non-

metric multi-dimensional scaling (NMDS). In the NMDS, average abundances of parasite species

in each component community were calculated and square root transformed to control for the effect

of high abundance parasite taxa. A resemblance matrix was constructed using Bray-Curtis

dissimilarity distance, and stress was calculated using Kruskal's stress formula 1. To determine

whether there were significant differences in parasite component community structure, we

performed a permutational multivariate ANOVA (PerMANOVA). For this analysis, we used site

as a factor and performed 999 permutations. As an overall significant difference was found ($P <$

0.05), we performed post-hoc pairwise PerMANOVA tests to identify differences among the factor

levels, and used the Bonferroni correction to preserve the type I error rate. In addition to the post-

hoc pairwise PerMANOVA tests, we performed a two-way crossed similarity percentage

(SIMPER) analysis to identify the parasite species that contributed to 5% or greater of the variation

in component community structure between sites (PRIMER-E LTD., Plymouth, UK).

To determine the role of local and regional factors in determining parasite infracommunity

richness, we used fully cross-validated regression trees and random forests with 1000 trees using

R (R Core Development Team, 2016). Prior to this analysis, we conducted a variable reduction

process to simplify the 48 water and 65 sediment trace metal and nutrient concentration data (JMP

Pro 12.0.0, SAS Institute Inc., Cary, NC). This analysis generated a correlation matrix of the log

transformed concentration data, placed all variables in one cluster, and then conducted binary splits

based on the magnitude of the second eigenvalues. Variables were iteratively reassigned to

clusters, and cluster membership was finalized when the squared correlation to the first principal

component was highest. In each cluster, the variable that explained the most variation was used in

the subsequent regression tree and random forest analysis. In the regression tree and random forest

analyses the response variable was parasite species richness and the predictor variables were host

characteristics such as size and sex, local abiotic measures determined through the PCA variable

reduction process, and the complete set of regional abiotic metrics. To avoid overfitting, we used

a cost-complexity algorithm to prune the regression tree: this algorithm removed auxiliary nodes

that would result in no significant loss in the mean square error (De'ath and Fabricius, 2000). To

validate the structure of the generated regression tree, we used random forest methods to generate

predictions on the weighted average of response value based on 1000 regression trees (Liaw and

Wiener, 2014).

RESULTS

Abiotic variation in regional and local environments

The salt marsh sites were selected on a gradient of local urbanization measured by the

percent of impervious surface in a 250 m buffer (Figure 1: Table 1). This measure of local

urbanization was not linearly correlated with the regional metric of urbanization (% developed

land in the watershed: Table 1). Skidaway Island (SK), which was characterized by the least

amount of impervious surface in the 250 m buffer at 5%, contained 20.60% developed land in the

watershed. Tybee Island (TB), which was characterized by the greatest amount of impervious

surface in a 250 m buffer at 66%, contained 6.17% developed land in the watershed. Shellman

Bluff (SB), which was characterized by 7% impervious surface in a 250 m buffer, contained the

least percentage of developed land in the watershed at 3.77%. Fort Pulaski (FP) was characterized

by 35% impervious surface in a 250 m buffer, but contained the greatest percentage of developed

land in the watershed at 24.06%.

Water column trace metal concentration differed across sites for 10 metals, but there was

no pattern of increasing metal concentration corresponding with increasing impervious surface or

regional urbanization (Table 2). For all metals there was variation due to the collection date

blocking factor, however seasonal patterns could not be determined given the single year of

sampling. Similarly, sediment trace metal concentration differed across sites for 13 metals, but

there was no linear pattern associated with measures of impervious surface or regional urbanization

(Table 3). No statistically significant differences were detected for water column TN, TP, pH,

temperature, or salinity. Water column chlorophyll *a* concentration differed across sites, with

increases in concentration corresponding with increasing impervious surface ($P < 0.0001$, $F_{5,66} =$

9.6438: Table 4).

The parasite community in Fundulus heteroclitus

Eight taxa of metazoan parasites were identified from the 630 fish. The direct life cycle

parasites included the branchiuran *Argulus funduli*, the copepod *Ergasilus funduli*, the leech

Myzobdella lugubris, and the monogeneans *Swingleus ancistrus* and *Fundulotrema prolongis*. The

complex life cycle parasites included the digenean *Lasiotocus minutus*, the metacestode

Glossocercus caribaensis, and the nematode *Contracaecum* sp.. These 8 taxa of parasites infected

more than 70% of the killifish examined and the prevalence for: *A. funduli* was 15.08%; *E. funduli*

was 18.73%; *M. lugubris* was 1.90%; *S. ancistrus* was 25.40%; *F. prolongis* was 22.38%; *L.*

minutus was 13.17%; *G. caribaensis* was 47.62%; and *Contraececum* sp. was 6.34% (Tables 5 and

6).

There was a difference in mean infracommunity species richness between sites ($P < 0.0001$,

$H_5 = 50.73$: Figure 2). Post-hoc tests revealed no difference between sites TB and CI, and both of

these sites had lower infracommunity richness than sites SK, SB, FP, and SM. Assessed with

NMDS and PerMANOVA, the parasite component communities were separate, and there was a

global difference in parasite component communities between sites ($P < 0.001$, Pseudo $F_{5,12} =$

6.11) (Figure 3). Pairwise PerMANOVA contrasts revealed no significant differences between

sites, however this was due to a conservative corrected alpha level of 0.003. Parasite communities

were generally similar between sites characterized by similar % impervious surface. Site SK

differed from sites CI ($P = 0.031$, Pseudo $F_{1,6} = 5.97$), SM ($P = 0.026$, Pseudo $F_{1,6} = 13.12$), and

TB ($P = 0.027$, Pseudo $F_{1,6} = 10.07$). Site SB differed from sites CI ($P = 0.032$, Pseudo $F_{1,6} = 5.99$),

SM ($P = 0.030$, Pseudo $F_{1,6} = 15.70$), and TB ($P = 0.032$, Pseudo $F_{1,6} = 13.59$). Site FP differed

from sites CI ($P = 0.027$, Pseudo $F_{1,6} = 4.21$), SM ($P = 0.04$, Pseudo $F_{1,6} = 7.53$), and TB ($P =$

0.031, Pseudo $F_{1,6} = 8.25$). SIMPER analysis indicated that the observed differences in parasite

community structure were driven by complex life-cycle parasites: the digenean *Lasciotocus*

minutus and the cestode *Glossocercus caribaensis* (Table 5). The sites containing higher local %

impervious surface were characterized by a lower prevalence and intensity of the digenean

Lasciotocus minutus. There was no difference in prevalence or intensity of the cestode

Glossocercus caribaensis prevalence or intensity among sites SK, SB, FP, CI, and TB – however,

site SM differed, with significantly higher prevalence and intensity of infection. Although the

SIMPER analysis did not identify any direct life cycle parasites that contributed to variation in

parasite communities, sites containing higher local % impervious surface generally had lower

prevalence and intensity for the monogeneans *Swingleus ancistrus* and *Fundulotrema prolongis*,

and the crustaceans *Argulus funduli* and *Ergasilus funduli* (Table 6).

Correlates of parasite species richness in Fundulus heteroclitus

The cluster analysis variable reduction process on the water and sediment trace metal and

nutrient concentrations resulted in 7 and 8 clusters, respectively. The 7 clusters in the water data

explained 77.1% of the variance, while the 8 clusters in the sediment concentration data explained

87% of the variance. Subsequently, we selected the variable that explained the highest variation

within each of the 15 total clusters for use in the regression tree and random forest analyses

(Supplemental Table 1).

We constructed regression trees that designated each of the 630 host observations of

infracommunity species richness to terminal leaf nodes: these terminal leaf nodes categorize the

local and regional drivers determining parasite species richness. We followed this regression tree

with a random forest analysis to assess the structure and validity of the generated regression tree.

For the tree using parasite species richness as the response variable, the optimum tree size

contained 5 leaves. In regression tree analysis, variable importance is determined as the reduction

in the loss function attributed to each variable at each split summed across the full tree. The

variables that were the most important in determining parasite species richness were host length,

host weight, regional % salt marsh, and patch density. Fish that had the highest infracommunity

richness were characterized by a total length less than 6.85 cm, weighed less than 11.25 g, and this

was independent of regional factors (mean richness of 3.158, N = 38). Fish that had the lowest

infracommunity richness were characterized by a total length greater than 6.85 cm and were

collected from a site with a regional % salt marsh less than 42.92 (mean parasite species richness

of 0.7333, N = 150). The most important variables for determining parasite species richness in the

random forest models were host length, host weight, regional % salt marsh, largest patch index,

patch density, and mean salt marsh patch size. The local parameters host length, host weight, and

the regional parameters % salt marsh, and patch density were detected as important predictors of

parasite species richness in both approaches.

DISCUSSION

The goal of this study was to determine the role of local and regional factors in structuring

parasite communities. The sites, selected on an urbanization gradient, had significantly different

parasite component community composition and richness. Variation in the prevalence and

intensity of infection by complex life cycle parasites, the digenean *Lasiotocus minutus* and the

cestode *Glossocercus caribaensis*, were the primary cause for the differences in the parasite

communities between sites. Although the prevalence of direct life cycle parasites differed among

sites, this difference did not contribute to the overall variation in community composition. The

best predictors of parasite infracommunity richness included local and regional measures: host

size, % regional salt marsh, and patch density. The data do not support the prediction that regional

urbanization is linearly associated with decreased parasite infracommunity richness; rather, our

results suggest that habitat size, both local and regional, and landscape heterogeneity have

differential effects on structuring the parasite community mediated by parasite life history.

Complex life cycle parasites can be broadly classified as autogenic, where the parasite

species completes its entire life cycle within the confines of an aquatic system, or allogenic where

the parasite uses hosts that are outside the aquatic environment (Esch et al., 1988). The extension

of this concept is that the local abiotic environment will have a greater effect on autogenic

parasites, as the parasites and hosts are continuously exposed to the aquatic environment. We

found that sites with the highest local impervious surface contained a lower prevalence and

intensity of the autogenic parasite *Lasiotocus minutus,* a digenean that uses the focal fish species,

Fundulus heteroclitus, as the definitive host, molluscs as the first intermediate host, and fish or

crustaceans as the second intermediate host (Stunkard, 1980). This pattern was expected, as this

parasite and associated hosts are continuously exposed to the external aquatic environment, and

would be exposed to any contaminants or deleterious physicochemical parameters. The second

complex life cycle parasite, the metacestode of *G. caribaensis*, is allogenic, and did not follow the

same pattern as *L. minutus*, and was prevalent in a site that had a high amount of impervious

surface. This allogenic cestode has fish-eating birds as definitive hosts and infects a number of fish

intermediate hosts (Scholz et al, 2004). This pattern was not expected, as there are still life stages

where the parasite and hosts are exposed to the external aquatic environment. However, it is

plausible to suggest that because the parasite spends less time in the aquatic environment the

parasite has a lesser risk of exposure to contaminants or deleterious physicochemical parameters.

Further, *G. caribaensis* has a vagile definitive host that may select habitat outside of urbanized salt

marshes, lessening the impact of local and regional urbanization. Consequently, the difference in

prevalence across sites may be due to the effect of the abiotic condition on parasite life stages

external to the host, or through effects on critical aquatic life cycle hosts, such as the mollusc

(Anderson and Sukhdeo, 2010).

Local abiotic factors have the potential to adversely impact the environmental stages of

parasites. For example, higher concentrations of nitrogen and phosphorus result in greater

mortality in *Fasciola hepatica* cercariae (Brglez and Wikerhauser, 1968), and experiments have

demonstrated that survival of cercarial *Philophtalmus* sp. is reduced at lower salinities compared

to survival at normal and high salinity (Lei and Poulin 2011). Similarly, Siddall and Des Clers

(1994) demonstrated that exposure to sewage sludge in seawater lowered the viability of cercariae

and miracidia of the digenean, *Zoogonoides viviparus*, due to trace concentrations of heavy metals

such as Hg, Cu, Cd, and Pb. However, our data show that local abiotic parameters (e.g., pH,

temperature, salinity, nitrogen, phosphorus) did not vary significantly across the sites, and the

regression tree and random forest analyses did not associate any local abiotic factor with parasite

species richness. Consequently, the dynamics we observe for the autogenic parasite, the digenean

L. minutus, are more likely to be determined by changes to the host population rather than affects

on life stages exposed to the environment.

Areas with higher levels of impervious surface are likely associated with stressors that

affect host populations. For example, King et al. (2004) demonstrated that the survival of the

marine bivalve *Tellina deltoidalis* could decrease to less than 10% in low salinity waters, which is

a common occurrence in urbanized watersheds following rain events. In a study germane to ours,

Hattenrath et al. (2010) manipulated nutrient concentrations in a marine system and found that a

toxin-producing dinoflagellate, *Alexandrium fundyense*, had a 4-fold increase in abundance and

toxin densities increased 8-fold. The nutrient concentrations in this experiment were parameterized

by field data collected during two harmful algal blooms in 2007 and 2008 that were associated

with large-scale bivalve population declines (Hattenrath et al., 2010). One striking pattern in our

data was an almost linear increase in chlorophyll *a* across the 6 sites, i.e., sites with more

impervious surfaces had a higher concentration of chlorophyll *a* (Table 4). Chlorophyll *a*

concentration is a proxy for phytoplankton abundance, and can be associated with eutrophication

and harmful algal blooms (Anderson et al., 2002). Further, chlorophyll *a* is often elevated in

estuaries nested in a developed landscape (Reed et al., 2016). The chlorophyll *a* values in our study

are not indicative of harmful algal blooms, but the linear increase in concentration suggests that

water quality is impacted. Though we did not measure mollusc abundance, it is plausible to suggest

that the elevated chlorophyll *a* levels and the consequent reduction in water quality affect the

molluscan first intermediate host and subsequently the observed parasite communities.

Through regression tree and random forest analyses, we demonstrated that host size was

the strongest determinant of parasite species richness in *F. heteroclitus*. Generally, larger hosts

contain more area for a parasite to colonize, allowing for a greater number of niches and concurrent

species (Guegan et al., 1992; Morand and Poulin, 1998). This species-area relationship is one of

the more consistent patterns in ecology (Arrhenius, 1921; Macarthur and Wilson, 1963), and

parasitologists have regularly found positive correlations between host size and infracommunity

richness (Rigby and Lawton, 1981; Morand et al., 2000; Behnke et al., 2001; Timi and Poulin,

2003; Kamiya et al., 2014). Our data demonstrate the opposite; that the strongest predictor for a

high infracommunity richness was a host with a smaller body size, which is in accordance with a

growing body of literature that finds no relationship between size and parasite richness (Feliu et

al., 1997; Krasnov et al., 2004; Korallo et al., 2007). Strona and Fattorini (2014) have argued that

ecological processes (e.g., competition and coextinction avoidance) are the more important

determinants of species richness, and the host size-species richness pattern is not a general

phenomenon (Guegan and Kennedy, 1993; Strona and Lafferty, 2013). Further, Anderson and

Sukhdeo (2011, 2013a) found that the predictors of parasite species richness in *F. heteroclitus*

were associated with ecological interactions and trophic generality. We suggest a similar

explanation for our data: smaller *F. heteroclitus* have a wider diet breadth facilitating infection by

complex life cycle parasites, and they also form higher density schools facilitating direct

transmission parasite establishment. This proposition is supported by Thompson (2015) who found

that *F. heteroclitus* displayed differential size-selective foraging where small individuals were

more likely to consume a diverse diet, consequently increasing the opportunity for infection by

multiple trophically transmitted parasite species. In addition, small *F. heteroclitus* individuals in

Georgia are generally restricted to surface pools in intertidal zone habitats resulting in increased

density which would likely increase direct parasite transmission (Kneib, 1987). Surface pools may

also act as focal points for complex life cycle transmission as small individuals in these pools are

vulnerable to predation by bird definitive hosts (Butner and Brattstrom, 1960; Kneib, 1982). Thus,

in this system, the negative relationship between body size and parasite richness is most likely due

to correlates with body size, such as trophic generality and other ecological interactions.

The secondary predictors for parasite infracommunity richness in *F. heteroclitus* were

associated with regional variables; the % regional salt marsh and patch density. Using a *Daphnia*

model system with 16 parasite taxa, Ebert et al. (2001) determined that time since colonization,

and pool volume - a proxy for host population size - were positively correlated with parasite

richness. To explain an increase of within-habitat transmission rates, they invoked principles from

general epidemiological models, arguing that the structure of the system allowed a lower

persistence threshold size for the parasite (Anderson and May, 1978; 1981; Gregory, 1990). A low

threshold allows for more parasite species to overcome that host density threshold and persist

(Dobson et al., 1992). The data from the present study similarly supports these models and the

observations of Ebert et al. (2001): larger contiguous areas of salt marsh patches, i.e. a greater %

salt marsh, likely contain a larger host population size, sustaining greater parasite species richness.

This proposition is further supported by Rudershausen et al. (2016) who determined that *F.*

heteroclitus population size was associated with creeks that had watersheds with high percentages

of contiguous marsh. Thus, the sites with a large % regional salt marsh likely contain larger

population sizes of *F. heteroclitus*, increasing the chances of a parasite encountering a suitable

host and establishing in the host population.

Coastal urbanization and anthropogenic disturbance are rapidly altering free-living and

parasite community dynamics. Our results suggest that regional urbanization may not always

negatively affect parasite species richness. The factors that influence the richness of the observed

parasite community are both local and regional, and the degree to which individual parasite species

respond to urbanization is determined by life-history strategy. Further, our data do not align with

the relatively general pattern relating species richness and host (habitat) size; instead, we suggest

that species richness in this system is determined by the interactions between free-living hosts and

their environment which is then detected as variation in the parasite community.

ACKNOWLEDGEMENTS

We acknowledge the advice and support provided by Dr. Clark Alexander and Mike

Robinson at the Applied Coastal Research Laboratory (Georgia Southern University), and Dr.

Sayed Hassad at the Laboratory of Environmental Analysis (University of Georgia). Field

assistance was provided by Emily Dodd, Sarah Dunn, Maria St. Jean, and Jackson Tomlinson.

Comments on an earlier draft by Drs. Risa Cohen, James Roberts, and Oscar Pung greatly

improved this manuscript. This research was funded in part by a Research Assistantship from the

Institute of Coastal Plain Science (Georgia Southern University) to JMA. TKA was supported by

the Office of Research Services and Sponsored Programs at Georgia Southern University (GSU)

and by an appointment to the USDA-ARS Research Participation Program administered by the

Oak Ridge Institute for Science and Education under contract number DE-AC05-06OR23100. The

funders had no role in study design, data collection and interpretation, or the decision to submit

the work for publication. Mention of trade names or commercial products in this article is solely

for the purpose of providing specific information and does not imply recommendation or

endorsement by the USDA. USDA is an equal opportunity provider and employer.

LITERATURE CITED

Alexander, C., R. Smith, B. Loganathan, J. Ertel, H. L. Windom, and R. F. Lee. 1999. Pollution

history of the Savannah River Estuary and comparisons with Baltic Sea pollution history.

Limnologica **29:** 267 - 273.

Anderson, R. M., and R. M. May. 1978. Regulation and stability of host-parasite population

interactions. 1. Regulatory processes. Journal of Animal Ecology **47:** 219-247.

Anderson, R. M., and R. M. May. 1979. Population biology of infectious diseases. Nature **280:**

361-367.

Anderson, R. M., and R. M. May. 1981. The population dynamics of micro-parasites and their

invertebrate hosts. Philosophical Transactions of the Royal Society B-Biological Sciences

291: 451-524.

Anderson, D. M., P. M. Glibert, and J. M. Burkholder. 2002. Harmful algal blooms and

eutrophication: Nutrient sources, composition, and consequences. Estuaries **25:** 704-726.

Anderson, T. K., and M. V. K. Sukhdeo. 2010. Abiotic versus biotic hierarchies in the assembly

of parasite populations. Parasitology **137:** 743-754.

Anderson, T. K., and M. V. K. Sukhdeo. 2011. Host centrality in food web networks determines

parasite diversity. PLoS ONE **6:** e26798.

Anderson, T. K., and M. V. K. Sukhdeo. 2013a. Qualitative community stability determines

parasite establishment and richness in estuarine marshes. PeerJ **1:** e92.

Anderson, T. K., and M. V. K. Sukhdeo. 2013b. The relationship between community species

richness and the richness of that parasite community in *Fundulus heteroclitus*. Journal of

Parasitology **99:** 391-396.

Angold, P. G., J. P. Sadler, M. O. Hill, A. Pullin, S. Rushton, K. Austin, E. Small, B. Wood, R.

Wadsworth, R. Sanderson, and K. Thompson. 2006. Biodiversity in urban habitat

patches. Science of the Total Environment **360:** 196-204.

Arneberg, P., A. Skorping, B. Grenfell, and A. F. Read. 1998. Host densities as determinants of

abundance in parasite communities. Proceedings of the Royal Society B-Biological

Sciences **265:** 1283-1289.

Aronson, M.F., C. H. Nilon, C.A. Lepczyk, T. S. Parker, P. S. Warren, S. S. Cilliers, M. A.

Goddard, A. K. Hahs, C. Herzog, M. Katti, and F. A. La Sorte. 2016. Hierarchical filters

determine community assembly of urban species pools. Ecology **97:** 2952-2963.

Arrhenius, O. 1921. Species and area. Journal of Ecology **9:** 95-99.

Behnke, J. M., C. J. Barnard, A. Bajer, D. Bray, J. Dinmore, K. Frake, J. Osmond, T. Race, and

E. Sinski. 2001. Variation in the helminth community structure in bank voles

(*Clethrionomys glareolus*) from three comparable localities in the Mazury Lake District

region of Poland. Parasitology **123:** 401-414.

Blanar, C.A., K. R. Munkittrick, J. Houlahan, D. L. MacLatchy, D. J. and Marcogliese. 2009.

Pollution and parasitism in aquatic animals: a meta-analysis of effect size. Aquatic

Toxicology **93:** 18-28.

Blanar, C. A., D. J. Marcogliese, and C. M. Couillard. 2011. Natural and anthropogenic factors

shape metazoan parasite community structure in mummichog (*Fundulus heteroclitus*)

from two estuaries in New Brunswick, Canada. Folia Parasitologica **58:** 240-248.

Blanar, C. A., M. Hewitt, M. McMaster, J. Kirk, Z. Wang, W. Norwood, and D. Marcogliese.

2016. Parasite community similiarity in Athabasca River trout-perch (*Percopsis*

omiscomaycus) varies with local-scale land use and sediment hydrocarbons, but not

distance or linear gradients. Parasitology Research **115:** 3853-3866.

Bradley, C. A., and S. Altizer. 2007. Urbanization and the ecology of wildlife diseases. Trends in

Ecology & Evolution **22:** 95-102.

Brglez, J., and T. Wikerhauser. 1968. On the effect of some fertilizers and copper sulphate upon

the survival of cercariae and metacercariae of *Fasciola hepatica*. Wiadomosci

parazytologiczne **14:** 675-677.

Butner, A., and B. H. Brattstrom. 1960. Local movements in *Menidia* and *Fundulus*. Copeia **2:**

139-141.

Burkholder, J. M., E. J. Noga, C. H. Hobbs, and H. B. Glasgow. 1992. New phantom

dinoflagellate is the causative agent of major estuarine fish kills. Nature **358:** 407-410.

Chapman, J. M., D. J. Marcogliese, C. D. Suski, and S. J. Cooke. 2015. Variation in parasite

communities and health indices of juvenile *Lepomis gibbosus* across a gradient of

watershed land-use and habitat quality. Ecological Indicators **57:** 564-572.

Chen, H. W., W. C. Liu, A. J. Davis, F. Jordan, M. J. Hwang, and K. T. Shao. 2008. Network

position of hosts in food webs and their parasite diversity. Oikos **117:** 1847-1855.

Daszak, P., A. A. Cunningham, and A. D. Hyatt. 2000. Wildlife ecology - Emerging infectious

diseases of wildlife - Threats to biodiversity and human health. Science **287:** 443-449.

De'ath, G., and K. E. Fabricius. 2000. Classification and regression trees: A powerful yet simple

technique for ecological data analysis. Ecology **81:** 3178-3192.

Dobson, A. P., S. V. Pacala, J. D. Roughgarden, E. R. Carper, and E. A. Harris. 1992. The

parasitse of Anolis lizards in the norther Lesser Antilles 1. Patterns of distribution and

abundance. Oecologia **91:** 110-117.

Ebert, D., J. W. Hottinger, and V. I. Pajunen. 2001. Temporal and spatial dynamics of parasite

richness in a Daphnia metapopulation. Ecology **82:** 3417-3434.

Esch, G. W., C. R. Kennedy, A. O. Bush, and J. M. Aho. 1988. Patterns in helminth communities

in freshwater fish in Great Britain - Alternative strategies for colonization. Parasitology

96: 519-532.

Feliu, C., F. Renaud, F. Catzeflis, J. P. Hugot, P. Durand, and S. Morand. 1997. Comparative

analysis of parasite species richness of Iberian rodents. Parasitology **115:** 453-466.

Gray, B. R., D. Bushek, J. W. Drane, and D. Porter. 2009. Associations between land use and

Perkinsus marinus infection of eastern oysters in a high salinity, partially urbanized

estuary. Ecotoxicology **18:** 259-269.

Gregory, R. D. 1990. Parasites and host geographic range as illustrated by waterfowl. Functional

Ecology **4:** 645-654.

Guegan, J. F., A. Lambert, C. Leveque, C. Combes, and L. Euzet. 1992. Can host body size

explain the parasite species richness in tropical freshwater fishes? Oecologia **90:** 197-

204.

Guegan, J. F., and C. R. Kennedy. 1993. Maximum local helminth parasite community richness

in British freshwater fish - A test of the colonization time hypothesis. Parasitology **106:**

91-100.

Hattenrath, T. K., D. M. Anderson, and C. J. Gobler. 2010. The influence of anthropogenic

nitrogen loading and meteorological conditions on the dynamics and toxicity of

Alexandrium fundyense blooms in a New York (USA) estuary. Harmful Algae **9:** 402-

412.

Hayward, B. W., H. R. Grenfell, K. Nicholson, R. Parker, J. Wilmhurst, M. Horrocks, A. Swales,

and A. T. Sabaa. 2004. Foraminiferal record of human impact on intertidal estuarine

environments in New Zealand's largest city. Marine Micropaleontology **53:** 37-66.

Hechinger, R. F., K. D. Lafferty, T. C. Huspeni, A. J. Brooks, and A. M. Kuris. 2007. Can

parasites be indicators of free-living diversity? Relationships between species richness

and the abundance of larval trematodes and of local benthos and fishes. Oecologia **151:**

82-92.

Hudson, P. J., A. P. Dobson, and K. D. Lafferty. 2006. Is a healthy ecosystem one that is rich in

parasites? Trends in Ecology & Evolution **21:** 381-385.

Hutchins, P. R., E. M. Smith, E. T. Koepfler, R. F. Viso, and R. N. Peterson. 2014. Metabolic

responses of estuarine microbial communities to discharge of surface runoff and

groundwater from contrasting landscapes. Estuaries and Coasts **37:** 736-750.

Huxham, M., and D. Raffaelli. 1995. Parasites and food web patterns. Journal of Animal

Ecology **64:** 168-176.

Johnson, D. S., J. W. Fleeger, and L. A. Deegan. 2009. Large-scale manipulations reveal that

top-down and bottom-up controls interact to alter habitat utilization by saltmarsh fauna.

Marine Ecology Progress Series **377:** 33-41.

Johnson, P. T. J., C. L. Wood, M. B. Joseph, D. L. Preston, S. E. Haas, and Y. P. Springer. 2016.

Habitat heterogeneity drives the host-diversity-begets-parasite-diversity relationship:

evidence from experimental and field studies. Ecology Letters **19:** 752-761.

Jones, G. K., J. L. Baker, K. Edyvane, and G. J. Wright. 1996. Nearshore fish community of the

Port river Barker inlet estuary, south Australia 1. Effect of thermal effluent on the fish

community structure, and distribution and growth of economically important fish species.

Marine and Freshwater Research **47:** 785-799.

Kamiya, T., K. O'Dwyer, S. Nakagawa, and R. Poulin. 2014. What determines species richness

of parasitic organisms? A meta-analysis across animal, plant and fungal hosts. Biological

Reviews **89:** 123-134.

King, C. K., M. C. Dowse, S. L. Simpson, and D. F. Jolley. 2004. An assessment of five

Australian polychaetes and bivalves for use in whole-sediment toxicity tests: Toxicity

and accumulation of copper and zinc from water and sediment. Archives of

Environmental Contamination and Toxicology **47:** 314-323.

Kneib, R. T. 1987. Predation risk and use of intertidal habitats by young fishes and shrimp.

Ecology **68:** 379-386.

Kneib, R. T. 1982. The effects of predation by wading birds (Ardeidae) and blue crabs

(Callinectes-Sapidus) on the population size structure of the common mummichog,

Fundulus heteroclitus. Estuarine Coastal and Shelf Science **14:** 159-165.

Korallo, N. P., M. V. Vinarski, B. R. Krasnov, G. I. Shenbrot, D. Mouillot, and R. Poulin. 2007.

Are there general rules governing parasite diversity? Small mammalian hosts and

Gamasid mite assemblages. Diversity and Distributions **13:** 353-360.

Krasnov, B. R., G. I. Shenbrot, I. S. Khokhlova, and A. A. Degen. 2004. Flea species richness

and parameters of host body, host geography and host 'milieu'. Journal of Animal

Ecology **73:** 1121-1128.

Krebs, J. M., C. C. McIvor, and S. S. Bell. 2014. Nekton community structure varies in response

to coastal urbanization near mangrove tidal tributaries. Estuaries and Coasts **37:** 815-831.

Kuhn, M. 2016. caret: Classification and Regression Training. R package.

Lafferty, K. D., S. Allesina, M. Arim, C. J. Briggs, G. De Leo, A. P. Dobson, J. A. Dunne, P. T.

J. Johnson, A. M. Kuris, D. J. Marcogliese, et al. 2008. Parasites in food webs: the

ultimate missing links. Ecology Letters **11:** 533-546.

Lei, F., and R. Poulin. 2011. Effects of salinity on multiplication and transmission of an intertidal

trematode parasite. Marine Biology **158:** 995-1003.

Lerberg, S. B., A. F. Holland, and D. M. Sanger. 2000. Responses of tidal creek macrobenthic

communities to the effects of watershed development. Estuaries **23:** 838-853.

Liaw, A., and M. Wiener. 2015. randomForest: Breiman and Cutler's Random Forests for

Classification and Regression. R package.

Lotrich, V. A. 1975. Summer home range and movements of *Fundulus heteroclitus* (Pisces-

Cyprinodontidae) in a tidal creek. Ecology **56:** 191-198.

Lotze, H. K., H. S. Lenihan, B. J. Bourque, R. H. Bradbury, R. G. Cooke, M. C. Kay, S. M.

Kidwell, M. X. Kirby, C. H. Peterson, and J. B. C. Jackson. 2006. Depletion, degradation,

and recovery potential of estuaries and coastal seas. Science **312:** 1806-1809.

Lowe, M. R., and M. S. Peterson. 2014. Effects of Coastal Urbanization on Salt-Marsh Faunal

Assemblages in the Northern Gulf of Mexico. Marine and Coastal Fisheries **6:** 89-107.

Macarthur, R. H., and E. O. Wilson. 1963. Equilibrium theory of insular zoogeography.

Evolution **17:** 373-387.

Machut, L. S., and K. E. Limburg. 2008. Anguillicola crassus infection in *Anguilla rostrata* from

small tributaries of the Hudson river watershed, New York, USA. Diseases of Aquatic

Organisms **79:** 37-45.

Magle, S. B., P. Reyes, J. Zhu, and K. R. Crooks. 2010. Extirpation, colonization, and habitat

dynamics of a keystone species along an urban gradient. Biological Conservation **143:**

2146-2155.

Morand, S., and R. Poulin. 1998. Density, body mass and parasite species richness of terrestrial

mammals. Evolutionary Ecology **12:** 717-727.

Morand, S., M. S. Hafner, R. D. M. Page, and D. L. Reed. 2000. Comparative body size

relationships in pocket gophers and their chewing lice. Biological Journal of the Linnean

Society **70:** 239-249.

Morley, N.J., S. W. B. Irwin, and J. W. Lewis. 2003. Pollution toxicity to the transmission of

larval digeneans through their molluscan hosts. Parasitology **126:** S5-S26.

Morrison, C. R., D. S. Bachoon, and K. W. Gates. 2008. Quantification of enterococci and

bifidobacteria in Georgia estuaries using conventional and molecular methods. Water

Research **42:** 4001-4009.

Norris, K., and M. R. Evans. 2000. Ecological immunology: life history trade-offs and immune

defense in birds. Behavioral Ecology **11:** 19-26.

Nunn, C. L., S. Altizer, K. E. Jones, and W. Sechrest. 2003. Comparative tests of parasite species

richness in primates. American Naturalist **162:** 597-614.

Paul, M. J., and J. L. Meyer. 2001. Streams in the urban landscape. Annual Review of Ecology

and Systematics **32:** 333-365.

Pietrock, M. and D. J. Marcogliese. 2003. Free-living endohelminth stages: at the mercy of

environmental conditions. Trends in Parasitology **19:** 293-299.

Rigby, C., and J. H. Lawton. 1981. Species - area relationships of arthropods on host plants -

Herbivores on Bracken. Journal of Biogeography **8:** 125-133.

Ross, C. L., and J. Barringer. 2006. Georgia 2030: Population projections for the 10-county

coastal region. Coastal Georgia Regional Development Center.

Rudershausen, P. J., J. A. Buckel, M. A. Dueker, S. J. Poland, and E. Hain. 2016. Comparison of

fish and invertebrate assemblages among variably altered tidal creeks in a coastal

landscape. Marine Ecology Progress Series **544:** 15-35.

Sanger, D. M., A. E. Holland, and G. I. Scott. 1999. Tidal creek and salt marsh sediments in

South Carolina coastal estuaries: I. Distribution of trace metals. Archives of

Environmental Contamination and Toxicology **37:** 445-457.

Scholz, T., R. A. Bray, R. Kuchta, and R. Repova. 2004. Larvae of gryporhynchid cestodes

(Cyclophyllidea) from fish: a review. Folia Parasitologica **51:** 131-152.

Schotthoefer, A. M., J. R. Rohr, R. A. Cole, A. V. Koehler, C. M. Johnson, L. B. Johnson, and

V. R. Beasley. 2011. Effects of wetland vs. landscape variables on parasite communities

of *Rana pipiens*: links to anthropogenic factors. Ecological Applications **21:** 1257-1271.

Siddall, R., and S. Des Clers. 1994. Effect of sewage sludge on the miracidium and cercaria of

Zoogonoides viviparus (Trematoda: Digenea). Helminthologia (Bratislava) **31:** 143-153.

Strona, G., and S. Fattorini. 2014. A few good reasons why species-area relationships do not

work for parasites. Biomed Research International **8:** 271680,

Strona, G., and K. D. Lafferty. 2013. Predicting what helminth parasites a fish species should

have using parasite co-occurrence modeler (PaCo). Journal of Parasitology **99:** 6-10.

Stunkard, H. W. 1980. Stages in the life-cycle of the digenetic trematode, Lasiotocus minutus

(Manter, 1931) Thomas, 1959. Biological Bulletin **159:** 497-498.

Teo, S. L. H., and K. W. Able. 2003. Habitat use and movement of the mummichog (*Fundulus*

heteroclitus) in a restored salt marsh. Estuaries **26:** 720-730.

Therneau, T., B. Atikinson, and B. Ripley. 2015. rpart: Recursive partitioning and regression

trees. version 4.1-10 ed. R package.

Thompson, J. S. 2015. Size-selective foraging of adult mummichogs, *Fundulus heteroclitus*, in

intertidal and subtidal habitats. Estuaries and Coasts **38:** 1535-1544.

Timi, J. T., and R. Poulin. 2003. Parasite community structure within and across host populations

of a marine pelagic fish: How repeatable is it? International Journal for Parasitology **33:**

1353-1362.

Wang, Y. Q., and D. K. Moskovits. 2001. Tracking fragmentation of natural communities and

changes in land cover: Applications of landsat data for conservation in an urban

landscape (Chicago wilderness). Conservation Biology **15:** 835-843.

FIGURE 1. Map of the 6 salt marsh study sites in coastal Georgia, USA, including 2011 land cover

data. The sites include Fort Pulaski (FP), Cockspur Island (CI), Tybee (TB), and Skidaway Island

(SK) in the Savannah River Estuary; Shellman Bluff (SB) as a representative of the Sapelo Sound;

and St. Marys (SM) as part of the St. Marys Estuary. Red circles represent a 250 m buffer around

the study site. For the purpose of this work, land cover data was reclassified from the original 21

categories to 10 (Salt Marsh, Developed, Freshwater Wetland, Open Water, Forest, Agriculture,

Bare Land, and Grassland): 2 land cover categories were not present in the study area (Freshwater

Aquatic Bed and Unconsolidated Shore).

FIGURE 2. Mean infracommunity species richness (± SE) of complex and direct life cycle

parasites within the salt marsh fish, *Fundulus heteroclitus* (N = 630 fish), from 6 sites in coastal

Georgia, USA. The sites include Fort Pulaski (FP), Cockspur Island (CI), Tybee (TB), and

Skidaway Island (SK) in the Savannah River Estuary; Shellman Bluff (SB) as a representative of

the Sapelo Sound; and St. Marys (SM) as part of the St. Marys Estuary. The sites are presented

from left to right reflecting increasing levels of impervious surface.

FIGURE 3. Non-metric multidimensional scaling plot of complex and direct life cycle parasite

component communities of the salt marsh fish, *Fundulus heteroclitus*, from 6 sites in coastal

Georgia, USA. Distances are based on Bray-Curtis dissimilarities of the square root transformed

mean abundance of 8 parasite taxa of *Fundulus heteroclitus*. The sites include: Fort Pulaski (FP)

(closed triangles), Cockspur Island (CI) (open triangles), Tybee (TB) (open squares), and

Skidaway Island (SK) (open circles) in the Savannah River Estuary; Shellman Bluff (SB) (closed

circles) as a representative of the Sapelo Sound; and St. Marys (SM) (closed squares) as part of

the St. Marys Estuary.

Table I. Regional characteristics of six salt marsh sites on coastal Georgia, USA. The sites include: Fort Pulaski (FP), Cockspur Island (CI), Tybee (TB), and Skidaway Island (SK) in the Savannah River Estuary; Shellman Bluff (SB) as a representative of the Sapelo Sound; and St. Marys (SM) as part of the St. Marys Estuary. The variables measured reflect regional (HUC-12 digit watershed) characteristics of landscape heterogeneity and composition. These data were obtained from the National Landcover Dataset, and analyzed in FRAGSTATS and ArcGIS. The units of measure are acres for Total Area, # of land use class patches per watershed area for Patch Density, % of the landscape that the largest land use class patch comprises for Largest Patch Index, the number of edges between different land use classes per watershed area for Edge Density, and mean size of salt marsh patches for Mean Salt Marsh Size.

	SK	SB	FP	CI	SM	TB
% Impervious surface	5	7	35	45	60	66
Total Area	8482	10441	47497	18101	11179	18101
Patch Density	39.6	29.9	24.2	23.9	28.2	23.9
Largest Patch Index	11.7	7.9	19.5	26.9	17.2	26.9
Edge Density	119.8	115.5	93.1	99.1	104.8	99.1
Mean Marsh Size	2.5	3.5	4.1	4.2	3.6	4.2
% Salt Marsh	39.2	25.6	3.2	46.7	33.2	46.7
% Developed	20.6	3.8	24.1	6.2	20.4	6.2
% Area Change to Developed (1996 – 2010)	6.4	1.9	18.5	2.6	7.3	2.6

Table II. Water column trace metal concentrations at six salt marsh sites on coastal Georgia, USA. The sites include: Fort Pulaski (FP), Cockspur Island (CI), Tybee (TB), and Skidaway Island (SK) in the Savannah River Estuary; Shellman Bluff (SB) as a representative of the Sapelo Sound; and St. Marys (SM) as part of the St. Marys Estuary. The sites are presented from left to right reflecting increasing levels of impervious surface. The presented trace metals (μg L^{-1}) represent those that differed across sites.

	SK		SB		FP		CI		SM		TB	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
Fe	994.39	351.45	909.00	356.76	1040.67	405.33	1083.11	424.73	1448.53	578.33	1287.66	414.82
Zn	66.09	25.75	75.23	29.76	59.57	23.49	50.73	19.5	62.22	24.6	91.26	37.25
Br	1.31	0.57	1.15	0.51	1.21	0.51	1.24	0.60	1.65	0.64	1.59	0.69
Sr	0.04	0.02	0.03	0.02	0.04	0.02	0.04	0.02	0.04	0.02	0.05	0.02
Cd	0.98	0.41	0.94	0.37	0.61	0.23	0.54	0.20	1.28	0.53	1.31	0.50
Sn	0.09	0.03	0.18	0.07	0.12	0.06	0.10	0.04	0.18	0.07	0.15	0.07
Ce	5.24	5.24	0.26	0.18	0	0	1.35	1.3	3.69	2.98	9.34	4.45
Pr	0.65	0.61	0.12	0.05	0.07	0.02	0.26	0.16	0.5	0.34	1.21	0.57
Sm	0.04	0.01	0.13	0.04	0.07	0.02	0.13	0.05	0.05	0.03	0.24	0.15
Au	0.96	0.61	9.83	8.18	2.38	1.62	4.84	2.85	3.02	1.49	0.93	0.53

Table III. Sediment trace metal concentrations at six salt marsh sites on coastal Georgia, USA. The sites include: Fort Pulaski (FP), Cockspur Island (CI), Tybee (TB), and Skidaway Island (SK) in the Savannah River Estuary; Shellman Bluff (SB) as a representative of the Sapelo Sound; and St. Marys (SM) as part of the St. Marys Estuary. The sites are presented from left to right reflecting increasing levels of impervious surface. The presented trace metals (mg Kg^{-1}) represent those that were determined to vary significantly between sites.

	SK		SB		FP		CI		SM		TB	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
Li	44.66	3.91	52.99	1.99	40.36	4.09	57.94	3.84	48.94	3.14	40.60	2.65
Na	520.78	29.69	520.01	43.93	404.42	26.07	390.18	11.78	379.78	24.12	294.79	47.17
V	42.27	5.13	54.21	4.52	49.19	3.94	52.20	0.96	40.73	3.49	29.45	4.25
Co	13.74	2.71	9.69	1.1191	6.82	0.55	8.15	0.26	8.05	0.67	6.16	0.42
Cu	32.18	0.83	48.88	18.85	20.15	2.38	21.72	1.96	24.68	2.81	15.24	2.15
Zn	151.82	33.22	146.31	46.33	49.35	6.12	104.05	32.03	69.10	3.16	36.26	6.53
Ga	2.09	0.32	2.54	0.13	2.11	0.14	2.88	0.12	2.48	0.11	1.89	0.15
I	0.01	0	0.01	0	0.02	0	0.01	0	0.01	0	0.01	0
Ba	25.70	2.30	30.60	1.34	23.77	1.30	31.98	1.02	28.05	2.30	21.91	0.75
Nd	20.51	1.07	23.81	1.83	22.79	0.93	21.98	0.18	21.19	0.72	18.51	0.77
Lu	0.04	0	0.07	0	0.05	0	0.06	0.01	0.06	0	0.04	0
Pb	2006.08	936.37	1681.16	1293.47	0	0	0	0	352.15	352.15	0	0
Th	14035.24	917.71	14414.14	314.00	16319.36	1005.79	16876.34	263.17	16883.67	596.68	13906.43	1107.74

Table IV. Mean chlorophyll *a* concentrations (µg L^{-1}) ± SE at six salt marsh sites on coastal Georgia, USA. The sites include: Fort Pulaski (FP), Cockspur Island (CI), Tybee (TB), and Skidaway Island (SK) in the Savannah River Estuary; Shellman Bluff (SB) as a representative of the Sapelo Sound; and St. Marys (SM) as part of the St. Marys Estuary. The sites are presented from left to right reflecting increasing levels of impervious surface. Collections, labeled A through D, occurred in March 2015 (A), June 2015 (B), September 2015 (C), and February 2016 (D).

	SK		SB		FP		CI		SM		TB	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
A	5.45	0.36	6.39	0.66	8.05	1.24	14.15	0.84	11.16	0.92	13.80	0.28
B	8.87	0.60	7.88	1.55	12.36	0.91	18.41	0.57	19.87	1.23	19.13	1.99
C	6.99	0.50	6.24	0.99	8.89	0.81	8.39	3.63	10.46	1.14	10.58	0.21
D	6.29	0.85	7.80	0.78	8.13	0.54	8.47	0.18	8.81	0.80	14.04	0.50

Table V. Prevalence (%) and mean intensity (±SE) of the three complex life cycle parasite taxa infecting the salt marsh fish, *Fundulus heteroclitus*. The sites include: Fort Pulaski (FP), Cockspur Island (CI), Tybee (TB), and Skidaway Island (SK) in the Savannah River Estuary; Shellman Bluff (SB) as a representative of the Sapelo Sound; and St. Marys (SM) as part of the St. Marys Estuary. The sites are presented from top to bottom reflecting increasing levels of impervious surface.

	Lasciotocus minutus		*Glossocercus caribaensis*		*Contracaecum* sp.	
	Prevalence	Intensity	Prevalence	Intensity	Prevalence	Intensity
SK	26.7	14.0 ± 2.9	44.8	4.0 ± 0.6	1.0	1.0 ± 0.0
SB	24.8	24.6 ± 4.4	39.0	3.9 ± 0.5	7.6	1.0 ± 0.0
FP	22.9	9.5 ± 1.7	448	4.2 ± 0.5	13.3	1.0 ± 0.0
CI	2.9	7.3 ± 2.4	35.2	3.4 ± 0.6	5.7	1.0 ± 0.0
SM	1.9	4.0 ± 3.0	91.4	6.0 ± 0.5	6.7	1.0 ± 0.0
TB	0	-	30.5	2.2 ± 0.3	3.8	1.0 ± 0.0

Table VI. Prevalence (%) and mean intensity (±SE) of the 5 direct life cycle parasite taxa infecting the salt marsh fish, *Fundulus heteroclitus*. The sites include: Fort Pulaski (FP), Cockspur Island (CI), Tybee (TB), and Skidaway Island (SK) in the Savannah River Estuary; Shellman Bluff (SB) as a representative of the Sapelo Sound; and St. Marys (SM) as part of the St. Marys Estuary. The sites are presented from top to bottom reflecting increasing levels of impervious surface.

	Myozobdella lugubris		*Swingleus ancistrus*		*Fundulotrema prolongis*		*Ergasilus funduli*		*Argulus funduli*	
	Prevalence	Intensity	Prevalence	Intensity	Prevalence	Intensity	Prevalence	Intensity	Prevalence	Intensity
SK	0	-	40.0	2.8 ± 0.4	29.5	4.6 ± 1.0	24.8	3.4 ± 0.7	15.2	1.4 ± 0.2
SB	1.9	1.0 ± 0	36.2	3.1 ± 0.3	20.0	4.2 ± 0.9	19.0	3.8 ± 0.7	20.0	1.4 ± 0.2
FP	9.5	1.2 ± 0.2	38.1	4.5 ± 0.7	29.5	6.0 ± 1.1	14.3	3.6 ± 0.7	24.8	1.4 ± 0.1
CI	0	-	8.6	2.0 ± 0.5	16.2	3.6 ± 0.8	20.0	3.9 ± 1.1	7.6	1.1 ± 0.2
SM	0	-	13.3	2.7 ± 0.5	20.0	3.6 ± 0.9	22.9	3.4 ± 0.9	12.4	1.2 ± 0.2
TB	0	-	16.2	1.4 ± 0.3	19.0	3.5 ± 0.5	11.4	3.1 ± 0.5	10.5	1.3 ± 0.1