1 **Increased frequency of travel in the presence of cross-immunity may act to**
2 **decrease the chance of a global pandemic**
3

4 **AUTHORS**
5 R.N. Thompson[1,2,3,*], C.P. Thompson[2], O. Pelerman[4], S. Gupta[2], U. Obolski[2,*]
6
7 *Correspondence to: robin.thompson@chch.ox.ac.uk, uri.obolski@zoo.ox.ac.uk
8
AFFILIATIONS
[1]Mathematical Institute, University of Oxford, Andrew Wiles Building, Radcliffe
Observatory Quarter, Woodstock Road, Oxford OX2 6GG, UK
[2]Department of Zoology, University of Oxford, South Parks Road, Oxford OX1 3PS, UK
[3]Christ Church, University of Oxford, St Aldate's, Oxford OX1 1DP, UK
[4]The Chaim Rosenberg School of Jewish Studies, Tel Aviv University, Tel Aviv 69978,
Israel
16

ABSTRACT
The high frequency of modern travel has led to concerns about a devastating pandemic
since a lethal pathogen strain could spread worldwide quickly. Many historical
pandemics have arisen following pathogen evolution to a more virulent form. However,
some pathogen strains invoke immune responses that provide partial cross-immunity
against infection with related strains. Here, we consider a mathematical model of
successive outbreaks of two strains – a low virulence strain outbreak followed by a high
virulence strain outbreak. Under these circumstances, we investigate the impacts of
varying travel rates and cross-immunity on the probability that a major epidemic of the
high virulence strain occurs, and the size of that outbreak. Frequent travel between
subpopulations can lead to widespread immunity to the high virulence strain, driven by
exposure to the low virulence strain. As a result, major epidemics of the high virulence
strain are less likely, and can potentially be smaller, with more connected
subpopulations. Cross-immunity may be a factor contributing to the absence of a global
pandemic as severe as the 1918 influenza pandemic in the century since.

KEYWORDS

major epidemic; antigenic variation; cross-immunity; pathogen diversity; mathematical modelling

1. INTRODUCTION

Outbreaks of infectious disease are responsible for around 14 million deaths annually [1,2]. In recent years, there have been a number of epidemics that have sparked fears that a global pandemic might develop [3]. Outbreaks such as the 2013-16 Ebola epidemic in West Africa did not develop into a global pandemic [4]. Other outbreaks, for example the 2009 H1N1 influenza pandemic, have included cases in countries worldwide [5] but have not been as destructive as initially feared [6]. Historically, however, there have been severe global pandemics. The 1918 'Spanish flu' pandemic killed around 50 million people [7], and the 1958 and 1968 influenza pandemics caused around one million deaths each [8,9]. These outcomes viewed collectively raise a number of questions: how likely is a global pandemic now, and has the lack of a devastating global pandemic in recent years been simply a matter of luck?

A number of factors affect the pandemic potential when a pathogen first appears in a host population. These include the spatial distribution of hosts and the level of mixing between subpopulations [10,11]. When a pathogen enters a population, a high host density within subpopulations and a high contact rate between subpopulations are most likely to represent appropriate conditions for a major epidemic and lead to high epidemic growth rates [12-15]. The modern world increasingly satisfies these conditions, with growing population sizes and large numbers of individuals living in urban centres, as well as the high frequency of worldwide airline travel [16]. Hence, one might expect the probability of a global pandemic, as well as its potential severity, to be at an all-time high.

However, the determinants of the dynamics of infectious disease outbreaks are

numerous and the above assertions ignore an important feature: cross-immunity

obtained from pathogen exposures in previous outbreaks. Viral, bacterial and eukaryotic

pathogens, such as the influenza virus, *Streptococcus pneumoniae* and the malaria

parasite, evolve in an attempt to avoid control by the immune system [17,18] and in

response to interventions such as vaccines [19-21]. Nonetheless, such immune evasion

is not perfect. Hosts that have been infected by a particular strain and acquired

immunity to that strain may also be at least partially immune to infection with related

strains [22-26]. Previous infections with immunologically related strains of a pathogen

can therefore be beneficial to hosts as they might provide protection against future

infections with other potentially more virulent strains.

Cross-immunity is known to affect the dynamics of outbreaks of various pathogens.

Vibrio cholera, for example, exhibits almost complete cross-immunity between strains (a

study by Koelle *et al.* [27] used an estimate that prior infection by a related strain

reduces the infection rate by 95.6%). Human Respiratory Syncytial Virus (hRSV) has

been shown to provide incomplete cross-immunity against infections from the same

virus group [28], and low levels of cross-immunity again hRSV infections from different

virus groups [28]. Furthermore, it provides partial cross-immunity against infections with

human Parainfluenza Virus [29].

Cross-immunity between related pathogens or pathogen strains has also been shown to

impact on pathogen dynamics and the structures of pathogen populations [30,31].

Cross-immunity might also be expected to affect the chance of and threat from a major

epidemic. The 2009 H1N1 pandemic, for example, was not as destructive as feared,

potentially as a result of decreased population susceptibility due to cross-immunity [32-

35]. Here we develop a mathematical model to investigate the impact of cross-immunity

on the chance of a major epidemic. We consider a general system emulating the

dynamics of an outbreak of a pathogen of high virulence in two connected

subpopulations when cross-immunity is present from a previously circulating low

virulence strain. We illustrate the principle that high rates of travel between

subpopulations can decrease the probability of a major epidemic of the high virulence

strain, and that the expected outbreak size can be either increased or decreased when

there is a higher rate of travel between subpopulations. In a connected world, novel

pathogens could spread worldwide through immunologically naïve populations

extremely quickly. However, epidemics of existing pathogens may be less frequent and

potentially smaller due to cross-immunity between pathogen strains – an important, but

previously underappreciated, factor.

102   **2. RESULTS**

Motivated by the spread of pathogens between geographically separate regions, we

considered pathogen transmission in a population consisting of two spatially distinct

subpopulations (Fig 1). Two outbreaks were assumed to occur. First, an outbreak of a

low virulence (LV) strain of the pathogen that has the potential to generate a large

number of cases but that is unlikely to lead to a large number of deaths. Then, an

outbreak of a related high virulence (HV) strain of the pathogen. Individuals infected in

the first outbreak were partially immune against infection in the second outbreak. It is

the probability of a major epidemic of the HV strain and the possible final sizes of the

HV strain outbreak that are our main concerns in this article.

[**Figure 1.** Schematic illustrating how outbreaks are simulated using the two-

subpopulation model. Circles represent susceptible (S_1, S_2), infected (I_1, I_2) and

removed (R_1, R_2) individuals in subpopulations 1 and 2, while the rates of transfer of

individuals between these classes are given on the arrows connecting the circles (see

Methods and Supplementary Material). A major epidemic of the LV strain is assumed to

occur, followed by an outbreak of the HV strain. Individuals infected in the LV strain

epidemic are conferred cross-immunity at a level α against infection with the HV strain.]

<u>Outbreak dynamics in a single subpopulation</u>

We initially considered outbreak dynamics in a single population. When the HV strain

arrived in the population, its effective reproduction number was reduced if individuals

had previously been infected by the LV strain. The higher the cross-immunity (governed

by the level of cross-immunity, $0 \leq \alpha \leq 1$), the lower the effective reproduction number

of the HV strain (Fig 2a). The higher the basic reproduction number of the LV strain,

R_0^{LV}, the lower the expected effective reproduction number of the HV strain. This is

because, when the LV strain had a high basic reproduction number, it generated a large

number of cases, thereby leading to large numbers of individuals with partial cross-

immunity against the HV strain (Fig 2a).

Similarly, when the HV strain arrived in the population, the probability of a major

epidemic (i.e. successful invasion of the host population) decreased as the level of

cross-immunity increased (Fig 2b), as did the final size of major epidemics of the HV

strain (Fig 2c). The expected number of HV strain infections, accounting for the

possibility that outbreaks may fade-out without becoming major epidemics, also

decreased when the level of cross-immunity increased (Fig 2d). Furthermore, these

quantities were also all reduced when the LV strain was more transmissible (see

different lines in Figs 2b-d).

We also considered stochastic simulations of the model (Fig 3), which support the

analytic result that cross-immunity lowers the probability of a major epidemic of the HV

strain in addition to its final size. The results in Figs 3b-c can be compared with Figs 2b-

c: for $\alpha = 0$ and $\alpha = 0.5$, the probability of a major epidemic in Fig 2b when $R_0^{LV} = 2$

takes the values 0.68 and 0.43, respectively – matching the percentage of outbreaks

that are major epidemics in Figs 3b and 3c. Similarly, for $\alpha = 0$ and $\alpha = 0.5$, the total

numbers of infections in Fig 2c when $R_0^{LV} = 2$ are 940 and 693, respectively – which are

also consistent with the major epidemics simulated in Figs 3b-c.

[**Figure 2.** The impact of cross-immunity on outbreaks of the HV strain in a single

population. (a) The effective reproduction number of the HV strain (R_e); (b) The

probability of a major epidemic of the HV strain; (c) The final size of the HV strain

outbreak when a major epidemic occurs ($R_e^{HV}(\infty)$); (d) The expected total number of

individuals infected by the HV strain, allowing for the possibility of a major epidemic or

fade-out without causing a major epidemic. The level of cross-immunity (α) is displayed

on the x-axis, and lines represent different values of the basic reproduction number of

the LV strain, R_0^{LV}. A major epidemic of the LV strain was assumed to occur prior to the

arrival of the HV strain in the population. Outbreaks of the HV strain were seeded with a

single infected individual, with all other individuals susceptible (either immunologically

naïve, S_1^N, or cross-immune following infection with the LV strain, S_1^I). Parameter values:

$N = 1000, \kappa = 1, R_0^{HV} = 3$. For a description of the parameters, see Table S2.]

[**Figure 3.** Cross-immunity leads to fewer and smaller major epidemics. Simulations of

the model in a single population showing: (a) The LV strain major epidemic; (b) The

subsequent HV strain outbreak, if there is no cross-immunity ($\alpha = 0$); (c) As in panel b,

but with cross-immunity ($\alpha = 0.5$). Parameter values:

$N = 1000, \kappa = 1, \mu = 0.143 \text{ day}^{-1}, \beta_{LV} = 2.86 \times 10^{-4} \text{ day}^{-1}$ (so that $R_0^{LV} = 2$), $\beta_{HV} =$

$4.29 \times 10^{-4} \text{ day}^{-1}$ (so that $R_0^{HV} = 3$). For a description of the parameters, see Table S2.

The blue regions represent quantiles of simulations of major epidemics and the white

dotted line represents the median values. In panels b-c, there are also thin light blue

areas at the bottoms of the panels showing outbreaks that fade-out without causing

major epidemics. The light blue area is omitted in panel a, since we only consider major

epidemics of the LV strain. Simulations are run in the same fashion as in the case of

two connected subpopulations but with the rate of travel between subpopulations set to

$\lambda = 0$, and with the outbreaks always seeded with a single infected individual in the first

subpopulation with the remaining $N - 1$ individuals susceptible. In the HV strain

epidemic, each susceptible individual is either immunologically naïve and in the class

S_1^N, or cross-immune following infection with the LV strain and in the class S_1^I.]

<u>Outbreak dynamics in the two-subpopulation model</u>

We then considered outbreak dynamics in two connected subpopulations and assessed

the impact of the extent of travel between subpopulations (governed by the parameter

λ), as well as the within-subpopulation transmissibility of the LV and HV strains

(governed by the parameter κ), on outbreaks of the HV strain. We considered the

effects of these parameters on the probability of a major epidemic of the HV strain (Fig

4a-c), the final size of an outbreak if a major epidemic is assumed to occur (Fig 4d-f),

and the expected final size accounting for the fact that outbreaks can either fade out as

minor outbreaks or take off as major epidemics (Fig 4g-i). We found that increased rates

of travel between subpopulations led to a lower probability of a major epidemic of the

HV strain. This is due to the probability of a major epidemic of the HV strain being

governed by local susceptibility when the HV strain first arrives in the population. Thus,

increased travel between subpopulations increases the probability that the LV strain has

previously caused a major epidemic in the subpopulation where the HV strain arrives

and conferred some immunity against the HV strain in that subpopulation. The

decreased chance of a major epidemic of the HV strain when between-subpopulation

travel was increased was particularly pronounced when the strains driving the LV and

HV epidemics were very transmissible and the level of cross-immunity was high (see

rightmost region of Fig 4c, in which there is a large reduction in the probability of a

major epidemic of the HV strain from the bottom to the top of the figure).

If the between-subpopulation connectivity was high when major epidemics of the HV

strain occurred, those epidemics tended to be more severe than when the between-

subpopulation connectivity was lower (Fig 4d-e). This was not the case at very high

levels of cross-immunity (Fig 4f). However, the expected total number of HV strain

infections, accounting for the possibility that the outbreak either took off and became a

major epidemic or faded out without spreading widely, reduced as the extent of

between-subpopulation travel increased when the level of cross-immunity was high (e.g.

Fig 4h-i) but increased for lower levels of cross-immunity (Fig 4g).

The outcome that the expected final size of major epidemics of the HV strain was

unaffected by between-subpopulation travel when the level of cross-immunity was high

(Fig 4f) can be explained as follows. The LV strain was assumed to always generate a

major epidemic in subpopulation 1. In that subpopulation, the cross-immunity conferred

was so high that a major epidemic of the HV strain could never occur in that

subpopulation. Consequently, the only way that a major epidemic of the HV could occur

in the overall population was if a major epidemic of the LV strain did not occur in

subpopulation 2, and the HV strain outbreak was seeded in subpopulation 2 and went

on to cause a major epidemic. Conditional on this scenario occurring, the size of that

major epidemic would be unaffected by the rate of between-subpopulation travel, as can

be seen in Fig 4f.

[**Figure 4.** The impact of within- and between-subpopulation travel on outbreaks of the

HV strain in the two-subpopulation model. The left, middle and right columns are results

with the level of cross-immunity, α, equal to 0.5, 0.7 and 0.9, respectively. (a)-(c): the

probability of a major epidemic of the HV strain; (d)-(f): the expected final size of major

epidemics of the HV strain; (g)-(i): the expected total number of the HV strain infections,

allowing for the possibility of a major epidemic or fade-out without causing a major

epidemic. A major epidemic of the LV strain is assumed to occur in at least one

subpopulation and is seeded by a single infected individual in subpopulation 1 (see Fig

1 and Supplementary Material). Outbreaks of the HV strain are seeded with a single

infected individual in a subpopulation chosen at random, with all other individuals in

both subpopulations susceptible. Parameter values: $N = 1000$ individuals per

subpopulation, within-subpopulation basic reproduction numbers $R_0^{LV} = 2\kappa, R_0^{HV} = 3\kappa$.

For a description of the parameters, see Table S2.]

We also verified that our results in Fig 4, which were derived numerically for

computational efficiency and based on analytical expressions (see Supplementary

Material), matched the results of stochastic simulations (Fig S1).

246   **3. DISCUSSION**

The large increase in international travel over the last century might be assumed to
have resulted in a high chance of a devastating global pandemic (see e.g. [36]). Here
we have used a general epidemiological model to demonstrate that an important, yet
often overlooked, factor in the dynamics of a newly introduced high-virulence (HV)
pathogen strain is partial immunity driven by exposures to related pathogen strains.
When a HV pathogen strain arrives in a population following an epidemic of a related
but low virulence (LV) strain, the probability of a major epidemic of the HV strain is
decreased. High rates of travel between spatially distinct subpopulations can drive
larger outbreaks of low virulence pathogens, in turn providing higher levels of immunity
if/when a HV strain, which has the potential to cause a devastating epidemic, appears in
the population (Fig 4a-c).
259
Not only did we find that the probability of a major epidemic of the HV strain decreases
when travel between subpopulations increases but the expected final size of the HV
strain outbreak can also be reduced. This was particularly pronounced when the level of
cross-immunity between strains was high (Fig 4i) since lower cross-immunity levels
combined with high travel rates can lead to large epidemics due to increased mixing
between subpopulations (Fig 4g). When between-subpopulation travel was increased,
the reduction in the probability of a major epidemic of the HV strain, and the reduction in
the expected size of the HV strain outbreak, was largely due to cross-immunity reducing
the proportion of outbreaks that proceeded to become major epidemics. When major
epidemics occurred, we found that they were typically larger when there was more
travel between regions (Fig 4d-e), although this was not always the case, particularly
when the level of cross-immunity was very high (Fig 4f).
272
Partial cross-immunity against a highly virulent strain from prior exposure to a less
virulent strain is characteristic of influenza epidemics. For example, it has been
suggested that individuals born before 1890 were protected against the 1918 H1N1
pandemic due to the outbreak in 1889-90 [37] and that individuals infected with multiple
historical seasonal H1N1 influenza strains were protected against the 2009 H1N1
influenza pandemic strain [38]. Potentially related to this, evidence indicates that older

adults are less likely to be severely affected by influenza pandemics than younger

adults, and this could be because of cross-immunity from exposure to related strains

[39]: in the 1977-78 H1N1 epidemic, for example, only adults under 26 years of age

experienced substantial morbidity [40,41]. Not only does cross-immunity protect

individuals from infection, but our results also suggest that cross-immunity might be a

potential explanatory factor as to why there has not been a pandemic as devastating as

the 1918 influenza epidemic in the century since, despite the emergence of a strain

antigenically similar to the 1918 pandemic strain in 2009. Increased travel during this

time period may have led to more widespread cross-immunity against strains of

pandemic potential, due to the global spread of related, less virulent strains. To verify

that travel rates have indeed increased substantially during the 20th century, as a

theoretical exercise we obtained crude estimates of the travel rates from Europe to the

USA, comparing rates during the early 20th century with current travel rates. For the

early 20th century rates, we used registry statistics from USA ports from 1914 to 1924

(see Supplementary Material Section 5). Oceanic travel should approximate overall

trans-Atlantic travel in this time period, as other modes of long-distance travel were rare.

This yielded the approximation $\lambda \approx 3.6 \cdot 10^{-6}$ per day. Conversely, when estimating

current rates of travel from European countries to the USA, approximated using data on

air travel, we obtain much higher values of $\lambda \approx 3.7 \cdot 10^{-4}$ per day. It might be expected

that such a significant rise in travel in this time period might have reduced the risk of a

global pandemic of a pathogen with circulating strains that induce cross-immunity (Fig

4).

Our assertion that higher rates of travel in the presence of cross-immunity may reduce

the impact of an epidemic is supported by evidence that island nations are more

affected during influenza pandemics than those that are connected by land to other

countries. Isolated islands such as the Solomon Islands experienced mortality rates of

in the range 4-10% during the 1918 pandemic [42], whereas more connected regions of

the world that had experienced more recent epidemics saw a mortality rate of 1-3% [43].

Furthermore, with respect to morbidity, we found that out of the 30 countries with the

highest number of cases per capita during the 2009 H1N1 pandemic, 24 were island

nations [44].

While we have chosen to focus on pathogens affecting human populations, we note that

cross-immunity might also impact on the pandemic potential for pathogens of animals

and plants. Cross-immunity between pathogen strains is a feature of pathogens of

animals including *Trypanosoma congolense* [45] and pathogens of plants including

Tobacco mosaic virus and Potato virus X [46]. Moreover, a previous study of the plant

pathogen *Podosphaera plantaginis* found that highly connected populations

experienced less pathogen colonisation and higher pathogen extinction than

populations that were less connected, and attributed this to the evolution of disease

resistance [47]. Furthermore, the effect of cross-immunity on plant or animal disease

outbreaks might also be changing due to alterations in the worldwide movement of

hosts. For example, climate change appears to be modifying the spatial distributions of

animal populations [48] and the plant nursery trade is more active than ever before [49].

We also note that, while we focussed on cross-immunity between different pathogen

strains, a related concept is protection against reinfection with the same (or very similar)

strain – often referred to as homologous interference or superinfection exclusion –

which has been demonstrated for a number of pathogens of plants or animals [50].

We aimed here to develop the simplest model possible characterising the spread of a

pathogen between spatially distinct populations in the presence of cross-immunity. For

assessing the probability of a major epidemic of a specific virulent strain of a pathogen

in a particular host population, the model would need to be extended and adjusted. The

process of mutation of the pathogen from the low virulence strain to the high virulence

strain may have to be modelled explicitly. If the mutation occurs a long period after the

previous major epidemic of the low virulence strain, cross-immunity might be expected

to have waned compared to if the low and high virulence strain epidemics occur in quick

succession. Alternatively, the high virulence strain epidemic might start before the low

virulence strain epidemic has ended. This would require an extension to the

epidemiological model, to permit the low and high virulence strains to co-circulate (for a

preliminary analysis, see Supplementary Information Section 6). The cumulative effects

of a number of past outbreaks of different related strains might also have to be

considered when a high virulence strain enters a host population [25,38]. Different types

of partial cross-immunity could be considered. Here we have assumed that exposure to

the LV strain reduces the probability of infection with the HV strain, whereas for some

infections cross-immunity may instead (or also) act to reduce the severity of disease

[41,51,52]. As an example, epidemiological isolation leading to high mortality rates of

infections may explain the mass mortality in pacific island populations between the 16[th]

and 19[th] centuries [53].

Other extensions to the model might have to be considered for application to specific

epidemics in human, animal or plant populations. For example, to explore the

contrasting effects of cross-immunity (which protects previously infected hosts) and

antibody-dependent enhancement (which would be expected to increase disease

severity in previously infected hosts [54]) for dengue epidemics in human populations, a

detailed model conditioned to the epidemiology of dengue would have to be developed.

In doing this, however, we note that incorporation of additional epidemiological detail in

models is not always useful, and that only necessary features should be included to

allow model outputs to be clearly understood [55]. For all epidemics, control

interventions might have to be included in the model, such as movement bans [56] or

trade restrictions [57] for livestock or commercial plant diseases. These particular

interventions could be included in the model by altering the within-subpopulation or

between-subpopulation rates of travel. Linked to this, behavioural responses of

individuals in the host population might have to be considered even if interventions are

not in place – for example, the awareness of the risk posed by the pathogen might be

higher following the LV strain epidemic, leading to reduced travel during the HV strain

epidemic.

Nonetheless, we have demonstrated the principle that increased global travel might not

necessarily mean that large pandemics are more likely in the present day than

previously. On the contrary, our results demonstrate that there may exist conditions

under which increased travel between subpopulations might reduce the probability and size of major epidemics. We do not contest that an increase in travel may sometimes increase the chance of a major epidemic, as well as its final size. The size of an epidemic of an entirely new pathogen, for example, or a strain that is antigenically distinct from previously circulating strains, is likely to be larger when there is more travel. This is because the pathogen would then be entering a population that is immunologically naïve and additional mixing provides the opportunity for more transmission events. There is therefore a need to test whether or not increased travel leads to smaller or larger epidemics in different theoretical contexts, as well as using data from a range of real-world outbreaks. However, we have focussed on epidemics occurring due to variants of pre-existing pathogens since these have driven a substantial number of past pandemics. Predicting which pathogen is likely to cause the next major pandemic is challenging [58]. Since our results demonstrate how previous outbreaks may have spread cross-immunity against strains related to pre-existing pathogens, our study has led us to support the assertion of the World Health Organisation: the pandemic threat may be greatest from an unknown strain of a known pathogen, or a pathogen that we have not previously encountered, the so-called disease X [59].

4. METHODS

Mathematical model

LV strain outbreak

We first considered an outbreak of a low virulence (LV) strain. Individuals were classified according to whether they were (S)usceptible to the LV strain, (I)nfected, or (R)ecovered and immune to that strain. Within a single subpopulation, the deterministic SIR model is given by

$$\frac{\mathrm{d}S}{\mathrm{d}t} = -\beta_{LV}\kappa IS,$$

$$\frac{\mathrm{d}I}{\mathrm{d}t} = \beta_{LV}\kappa IS - \mu I,$$

$$\frac{\mathrm{d}R}{\mathrm{d}t} = \mu I. \qquad\qquad (1)$$

The parameter κ is a proxy for the frequency of within-subpopulation travel and governs

the magnitude of the contact rate within the subpopulation. This parameter is also used

in the subsequent high virulence (HV) strain outbreak (see below). The LV strain

epidemic was assumed to follow stochastic SIR dynamics, analogous to the system of

equations (1) within each subpopulation, with individuals also travelling between

subpopulations at rate λ per day (see Fig 1 and Supplementary Material).

HV strain outbreak

We then considered a subsequent outbreak of the HV strain in the population, that is

also governed by stochastic SIR dynamics in each subpopulation but extended to

account for cross-immunity.

Individuals infected in the LV strain epidemic were partially protected against the HV

strain epidemic. The extent of immunity against the HV strain conferred by the LV strain

was governed by a parameter $0 \leq \alpha \leq 1$, which changed the probability of successful

infection by a multiplicative factor $1 - \alpha$ if the host had previously been infected by the

LV strain compared to if the host had not previously been infected by the LV strain. The

value $\alpha = 0$ thereby corresponded to no cross-immunity, $\alpha = 1$ corresponded to

complete cross-immunity, and intermediate values of α corresponded to partial cross-

immunity.

Individuals were classified according to whether they were (S_N)usceptible and not

partially cross-immune against the HV strain, (S_I)usceptible and partially cross-immune,

(I)nfected, or (R)ecovered and immune to that strain or dead. The deterministic

analogue in a single subpopulation to the model that we considered to represent the HV
strain outbreak is
430

$$\frac{\mathrm{d}S_N}{\mathrm{d}t} = -\beta_{HV}\kappa IS_N,$$

$$\frac{\mathrm{d}S_I}{\mathrm{d}t} = -\beta_{HV}\kappa(1-\alpha)IS_I,$$

$$\frac{\mathrm{d}I}{\mathrm{d}t} = \beta_{HV}\kappa IS_N + \beta_{HV}\kappa(1-\alpha)IS_I - \mu I,$$

$$\frac{\mathrm{d}R}{\mathrm{d}t} = \mu I. \tag{2}$$

When the HV strain arrived in the subpopulation, initial cases could either lead to a
major epidemic or fade-out as a minor outbreak. The within-subpopulation basic
reproduction numbers of the LV and HV strains were denoted by R_0^{LV} and R_0^{HV},
respectively (see Supplementary Material). In the HV strain outbreak, like in the
previous LV strain outbreak, we also assumed that individuals travelled between
subpopulations at rate λ per day (see Fig 1).

<u>Background theory</u>

Before considering the full model, we derived the probability of a major epidemic and
the expected final size conditional on a major epidemic when the pathogen first arrived
in a subpopulation, ignoring any effects of migration between subpopulations.

LV strain epidemic

When the LV strain, with within-subpopulation basic reproduction number $R_0^{LV} > 1$,
arrived in a single fully susceptible population, the probability of a major epidemic
following was approximately

$$\text{Prob(major epidemic)} \approx 1 - \left(\frac{1}{R_0^{LV}}\right)^{I(0)},$$

where $I(0)$ is the number of infected hosts that were infected initially (see e.g. [60,61]).

If a major epidemic occurred, the expected number of individuals infected over the

course of the epidemic was approximated by the solution, $R^{LV}(\infty)$, of the final size

equation (see equation (S3) in the Supplementary Material),

$$R^{LV}(\infty) = N - N \exp\left(\frac{-R_0^{LV} R^{LV}(\infty)}{N}\right).$$

HV strain epidemic

When the HV strain appeared in a subpopulation, if a major epidemic of the LV strain

had not occurred in that subpopulation, then the probability of a major epidemic was

$$\mathbb{P}\text{(major epidemic)} \approx 1 - \left(\frac{1}{R_0^{HV}}\right)^{I(0)},$$

467   whenever $R_0^{HV} > 1$, where $I(0)$ is the initial number of hosts infected by the HV strain at

468   the start of the HV strain outbreak. The final size if a major epidemic occurred was then

469   given by the solution, $R_0^{HV}(\infty)$, of the final size equation

$$R_0^{HV}(\infty) = N - N \exp\left(\frac{-R_0^{HV} R_0^{HV}(\infty)}{N}\right).$$

However, when the HV strain arrived in the subpopulation, if a major epidemic of the LV

strain had occurred previously, then the population was not fully susceptible due to

partial cross-immunity. The effective reproduction number of the HV strain when it

arrived in the population was then $R_e = \frac{\beta_{HV}\kappa(1-\alpha x)N}{\mu}$, where $x = \frac{R^{LV}(\infty)}{N}$ is the proportion of

individuals infected in the LV strain epidemic (see Supplementary Material Section 2).

The resulting probability of a major epidemic of the HV strain was then

$$\mathbb{P}(\text{major epidemic}) \approx 1 - \left(\frac{1}{R_e}\right)^{I(0)},$$

whenever $R_e > 1$, and if a major epidemic occurred, the final size was approximated by

numerically solving the system of equations (2).

<u>Possible outcomes of LV and HV strain outbreaks in two subpopulations</u>

When we considered the full model, consisting of two subpopulations (denoted 1 and 2),

we assumed without loss of generality that the LV strain arrived in subpopulation 1.

Since we were only interested in the effect of cross-immunity on outbreaks of the HV

strain, we assumed that the LV strain successfully invaded the population. As a result, a

major epidemic of the LV strain occurred in subpopulation 1.

Eight outcomes were then possible (see Table S1). A major epidemic of the LV strain

may or may not occur in subpopulation 2. Then, major epidemics of the HV strain may

or may not occur in each of subpopulations 1 and 2. Calculation of the probabilities of

each of these outcomes, as well as the expected numbers of individuals infected by the

LV and HV strains in each subpopulation, are described in the Supplementary Material.

REFERENCES

1. Taylor LH, Latham SM, Mark EJ. 2001 Risk factors for human disease emergence.
Phil. Trans. R. Soc. B **356**: 983-989.

2. Dye C. 2014 After 2015: infectious diseases in a new era of health and
development. *Phil. Trans. R. Soc. B* **369**, 20130426.

3. Reperant LA, Osterhaus AD. 2017 AIDS, Avian flu, SARS, MERS, Ebola, Zika…
what next? *Vaccine* **35**, 4470-4474.

4. Coltart CE, Lindsey B, Ghinai I, Johnson AM, Heymann DL. 2017 The Ebola
outbreak, 2013–2016: old lessons for new epidemics. *Phil. Trans. R. Soc. B* **372**,
20160297.

5. Girard MP, Tam JS, Assossou OM, Kieny MP. 2010 The 2009 A (H1N1) influenza
virus pandemic: A review. *Vaccine* **28**, 4895-4902.

6. SteelFisher GK, Blendon RJ, Bekheit MM, Lubell K. 2010 The public's response to
the 2009 H1N1 influenza pandemic. *New Eng. J. Med.* **362**, e65.

7. Johnson NP, Mueller J. 2002 Updating the accounts: global mortality of the 1918-
1920 "Spanish" influenza pandemic. *Bull. Hist. Med.* **76**, 105-115.

8. Kilbourne ED. 2006 Influenza Pandemics of the 20th Century. *Emerg. Infect. Dis*. **12**,
9-14.

9. Saunders-Hastings PR, Krewski D. 2016 Reviewing the history of pandemic
influenza: understanding patterns of emergence and transmission. *Pathogens* **5**, 66.

10. Getz WM, Lloyd-Smith JO, Cross PC, Bar-David S, Johnson PL, Porco TC,
Sánchez MS. 2006 "Modeling the invasion and spread of contagious diseases in
heterogeneous populations" in *Disease Evolution: Models, Concepts, and Data
Analyses*, ed. Z Feng, U Dieckmann, SA Levin (DIMACS), 113-144.

11. Engering A, Hogerwerf L, Slingenbergh J. 2013 Pathogen-host-environment
interplay and disease emergence. *Emerg. Microbes Infect.* **2**, e5.

12. Gupta S, Anderson RM, May RM. 1989 Networks of sexual contacts: implications
for the pattern of spread of HIV. *AIDS* **3**, 807-817.

13. Woolhouse ME, Shaw DJ, Matthews L, Liu WC, Mellor DJ, Thomas MR. 2005
Epidemiological implications of the contact network structure for cattle farms and the
20–80 rule. *Biol. Letters* **1**, 350-352.

14. Keeling MJ. 2005 The implications of network structure for epidemic dynamics.
Theor. Pop. Biol. **67**, 1-8.

15. Woolhouse ME. 2011 How to make predictions about future infectious disease risks.
Phil. Trans. Roy. Soc. B **366**, 2045-2054.

16. Mangili A, Gendreau MA. 2005 Transmission of infectious diseases during
commercial air travel. *Lancet* **365**, 989-996.

17. Vossen MT, Westerhout EM, Söderberg-Nauclér C, Wiertz EJ. 2002 Viral immune
evasion: a masterpiece of evolution. *Immunogenetics* **54**, 527-542.

18. Finlay BB, McFadden G. 2006 Anti-immunology: evasion of the host immune
system by bacterial and viral pathogens. *Cell* **124**, 767-782.

19. Smith DJ, Lapedes AS, de Jong JC, Bestebroer TM, Rimmelzwaan GF, Osterhaus
AD, Fouchier RA. 2004 Mapping the antigenic and genetic evolution of influenza virus.
Science **305**, 371-376.

20. Gandon S, Mackinnon MJ, Nee S, Read AF. 2001 Imperfect vaccines and the
evolution of pathogen virulence. *Nature* **414**, 751-756.

21. Obolski U, Lourenço J, Thompson CP, Thompson RN, Gori A, Gupta S. 2018
Vaccination can drive an increase in frequencies of antibiotic resistance among
nonvaccine serotypes of *Streptococcus pneumoniae*. *Proc. Natl. Acad. Sci.* **6**,
201718712.

22. Gog JR, Grenfell BT. 2002 Dynamics and selection of many-strain pathogens. *Proc.*
Natl. Acad. Sci. **99**, 17209-17214.

23. Grenfell BT, Pybus OG, Gog JR, Wood JL, Daly JM, Mumford JA, Holmes EC. 2004
Unifying the epidemiological and evolutionary dynamics of pathogens. *Science* **303**,
327-332.

24. Reich NG, Shrestha S, King AA, Rohani P, Lessler J, Kalayanarooj S, Yoon IK,
Gibbons RV, Burke DS, Cummings DA. 2013 Interactions between serotypes of dengue
highlight epidemiological impact of cross-immunity. *J. R. Soc. Interface* **10**, 20130414.

25. Thompson CP, Lourenço J, Walters AA, Obolski U, Edmans M, Palmer DS, Kooblall
K, Carnell GW, O'Connor D, Bowden TA, *et al*. 2018 A naturally protective epitope of
limited variability as an influenza vaccine target. *Nature Commun.* **9**, 3859.

26. Thompson CP, Obolski U. 2018 Influenza vaccination and the 'diversity paradox'.
Hum. Vaccin. Immunother. **14**, 3005-3009.

27. Koelle K, Pascual M, Yunus M. 2006 Serotype cycles in cholera dynamics. *Proc.*
Roy. Soc. Lond. B **273**, 2879-2886.

28. White LJ, Waris M, Cane PA, Nokes DJ, Medley GF. 2005 The transmission
dynamics of groups A and B human respiratory syncytial virus (hRSV) in England &
Wales and Finland: seasonality and cross-protection. *Epidem. Infect.* **133**, 279-289.

29. Bhattacharyya S, Gesteland PH, Korgenski K, Bjørnstad ON, Adler FR. 2015 Cross-
immunity between strains explains the dynamical pattern of paramyxoviruses. *Proc.*
Natl. Acad. Sci. **112**, 13396-400.

30. Gupta S, Maiden MC, Feavers IM, Nee S, May RM, Anderson RM. 1996 The
maintenance of strain structure in populations of recombining infectious agents. *Nature*
Med. **2**, 437-442.

31. Gupta S, Ferguson N, Anderson R. 1998 Chaos, persistence, and evolution of strain
structure in antigenically diverse infectious agents. *Science* **280**, 912-915.

32. Hancock K, Veguilla V, Lu X, Zhong W, Butler EN, Sun H, Liu F, Dong L, DeVos JR,
Gargiullo PM, *et al*. 2009 Cross-reactive antibody responses to the 2009 pandemic
H1N1 influenza virus. *New Eng. J. Med.* **361**, 1945-1952.

33. Zimmer SM, Crevar CJ, Carter DM, Stark JH, Giles BM, Zimmerman RK, Ostroff
SM, Lee BY, Burke DS, Ross TM. 2010 Seroprevalence following the second wave of
pandemic 2009 H1N1 influenza in Pittsburgh, PA, USA. *PLoS One* **5**, e11601.

34. Ikonen N, Strengell M, Kinnunen L, Österlund P, Pirhonen J, Broman M, Davidkin I,
Ziegler T, Julkunen I. 2010 High frequency of cross-reacting antibodies against 2009
pandemic influenza A(H1N1) virus among the elderly in Finland. *Euro Surveill.* **15**,
19478.

35. Miller MS, Gardner TJ, Krammer F, Aguado LC, Tortorella D, Basler CF, Palese P.
Neutralizing antibodies against previously encountered influenza virus strains increase
over time: A longitudinal analysis. *Sci. Transl. Med.* **5**, 198ra107.

36. Cooper BS, Pitman RJ, Edmunds WJ, Gay NJ. 2006 Delaying the international
spread of pandemic influenza. *PLoS Med.* **3**, e212.

37. Gagnon A, Miller MS, Hallman SA, Bourbeau R, Herring DA, Earn DJ, Madrenas J.
2013 Age-specific mortality during the 1918 influenza pandemic: Unravelling the
mystery of high young adult mortality. *PLoS One* **8**, e69586.

38. Carter DM, Bloom CE, Nascimento EJ, Marques ET, Craigo JK, Cherry JL, Lipman
DJ, Ross TM. 2013 Sequential seasonal H1N1 influenza virus infections protect ferrets
against novel 2009 H1N1 influenza virus. *J. Virol.* **87**, 1400-1410.

39. Luk J, Gross P, Thompson WW. 2001 Observations on mortality during the 1918
influenza pandemic. *Clin. Inf. Dis.* **33**, 1375-1378.

40. Scholtissek C, von Hoyningen V, Rott R. 1978 Genetic relatedness between the
new 1977 epidemic strains (H1N1) of influenza and human influenza strains isolated
between 1947 and 1957 (H1N1). *Virology* **89**, 613-617.

41. Reichert T, Chowell G, McCullers JA. 2012 The age distribution of mortality due to
influenza: pandemic and peri-pandemic. *BMC Med.* **10**, 162.

42. Wilson N, Mansoor O, Lush D, Kiedrzynski T. 2005 Modeling the impact of
pandemic influenza on Pacific Islands. *Emerg. Infect. Dis.* **11**, 347-349.

43. Ansart S, Pelat C, Boelle PY, Carrat F, Flahault A, Valleron AJ. 2009 Mortality
burden of the 1918-1919 influenza pandemic in Europe. *Influenza Other Respir. Viruses*
3, 99-106.

44. Central Intelligence Agency. 2018 The World Factbook. Accessible at
www.webcitation.org.

45. Masumu J, Marcotty T, Geerts S, Vercruysse J, Van den Bossche P. 2009 Cross-
protection between *Trypanosoma congolense* strains of low and high virulence. *Vet.*
Parasitol. **163**, 127-131.

46. Gal-On A, Shiboleth YM. 2006 "Cross protection" in *Natural Resistance*
Mechanisms of Plants to Viruses, ed. G Loebenstein, JP Carr (Dordrecht: Springer),
261–288.

47. Jousimo J, Tack AJM, Ovaskainen O, Mononen T, Susi H, Tollenaere C, Laine A-L.
2014 Ecological and evolutionary effects of fragmentation on infectious disease
dynamics. *Science* **344**, 1289-1293.

48. Walther GR, Post E, Convey P, Menzel A, Parmesan C, Beebee TJ, Fromentin JM,
Hoegh-Guldberg O, Bairlein F. 2002 Ecological responses to recent climate change.
Nature **416**, 389-395.

49. Liebhold AM, Brockerhoff EG, Garrett LJ, Parke JL, Britton KO. 2012 Live plant
imports: the major pathway for forest insect and pathogen invasions of the US. *Frontiers*
Ecol. Environ. **10**, 135-143.

50. Folimonova SY, Robertson CJ, Shilts T, Folimonov AS, Hilf ME, Garnsey SM,
Dawson WO. 2010 Infection with strains of Citrus tristeza virus does not exclude
superinfection by other strains of the virus. *J. Virol.* **84**, 1314-1325.

51. Gupta S, Snow RW, Donnelly CA, Marsh K, Newbold C. 1999 Immunity to non-
cerebral severe malaria is acquired after one or two infections. *Nature Med.* **5**, 340-343.

52. Hellriegel B. 2001 Immunoepidemiology – bridging the gap between immunology
and epidemiology. *Trends Parasitol.* **17**, 102-106.

53. Penman BS, Gupta S, Shanks GD. 2017 Rapid mortality transition of Pacific Islands
in the 19th century. *Epidem. Infect.* **145**, 1-11.

54. Ferguson N, Anderson R, Gupta S. 1999 The effect of antibody-dependent
enhancement on the transmission dynamics and persistence of multiple-strain
pathogens. *Proc. Natl. Acad. Sci.* **96**, 790-794.

55. Thompson RN, Hart WS. 2018 Effect of confusing symptoms and infectiousness on
forecasting and control of Ebola outbreaks. *Clin. Inf. Dis.* **67**, 505-518.

56. Ferguson NM, Donnelly CA, Anderson RM. 2001 The foot-and-mouth epidemic in
Great Britain: pattern of spread and impact of interventions. *Science* **292**, 1155-1160.

57. Thompson RN, Cobb RC, Gilligan CA, Cunniffe NJ. 2016 Management of invading
pathogens should be informed by epidemiology rather than administrative boundaries.
Ecol. Model. **324**, 28-32.

58. Holmes EC, Rambaut A, Andersen KG. 2018 Pandemics: Spend on surveillance,
not prediction. *Nature* **558**, 180-182.

59. World Health Organization. 2018 Annual review of diseases prioritized under the
research and development blueprint. Accessible at
http://www.who.int/emergencies/diseases/2018prioritization-report.pdf?ua=1.

60. Keeling MJ, Rohani P. 2007 *Modeling Infectious Diseases in Humans and Animals*.
Princeton University Press.

61. Thompson RN, Gilligan CA, Cunniffe NJ. 2016 Detecting presymptomatic infection
is necessary to forecast major epidemics in the earliest stages of infectious disease
outbreaks. *PLoS Comp. Biol.* **12**, e1004836.
654

DATA AVAILABILITY

All numerical calculations and stochastic simulations in this manuscript were performed

in Matlab. Accompanying code for the full model can be found at www.github.com/robin-

thompson/PandemicPotentialGivenCrossProtection.

COMPETING INTERESTS

We have no competing interests.

AUTHORS' CONTRIBUTIONS

RNT and SG conceived the research; All authors designed the study; RNT and UO

carried out the research; RNT drafted the manuscript; All authors revised the

manuscript and gave final approval for publication.

ACKNOWLEDGEMENTS

Thanks to members of the Department of Zoology and Mathematical Institute in Oxford

for helpful discussions about this project.

FUNDING

RNT was funded by a Junior Research Fellowship from Christ Church, Oxford. UO was

supported by an EMBO postdoctoral fellowship. CPT was supported by the European

Union's Seventh Framework Programme (FP7/2007-2013)/European Research Council

(614725-PATHPHYLODYN).

(a)

Total number infected ($I + R$) by time t

10 30 50 70 90

Time (t days)

Without cross-immunity ($\alpha = 0$)

With cross-immunity ($\alpha = 0.5$)

(b)

Total number infected ($I + R$) by time t

Percentage of outbreaks that are major epidemics: 67.6%

Percentage of outbreaks that fade out without major epidemic: 32.4%

Time (t days)

(c)

Total number infected ($I + R$) by time t

Percentage of outbreaks that are major epidemics: 43.2%

Percentage of outbreaks that fade out without major epidemic: 56.8%

Time (t days)