Letter

Experimentally reduced insulin/IGF-1 signalling in adulthood extends

lifespan of parents and improves Darwinian fitness of their offspring

Running title: Lifespan extension improves offspring fitness

Martin I. Lind[1, †, *], Sanjana Ravindran[1, †], Zuzana Sekajova[1], Hanne Carlsson[1, 2],

Andrea Hinas[3] and Alexei A. Maklakov[1, 2, *]

[1] Animal Ecology, Department of Ecology and Genetics, Uppsala University, Norbyvägen

18D, Uppsala 752 36, Sweden

[2] School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich

NR4 7TJ, UK

[3] Department of Cell and Molecular Biology, Uppsala University, PO Box 596, Uppsala 751

24, Sweden

[†] these authors contributed equally to the study

* Authors for correspondence:

Martin I. Lind - martin.i.lind@gmail.com

Alexei A. Maklakov - a.maklakov@uea.ac.uk

Total word count: 5714

Keywords: ageing, antagonistic pleiotropy, functional trade-offs, hyperfunction, IIS

signalling, parental effects, senescence

Abstract

Classical theory maintains that ageing evolves via energy trade-offs between reproduction

and survival leading to accumulation of unrepaired cellular damage with age. In contrast, the

emerging new theory postulates that ageing evolves because of deleterious late-life hyper-

function of reproduction-promoting genes leading to excessive biosynthesis in late-life. The

hyper-function theory uniquely predicts that optimizing nutrient-sensing molecular signalling

in adulthood can simultaneously postpone ageing and increase Darwinian fitness. Here we

show that reducing evolutionarily conserved insulin/IGF-1 nutrient-sensing signalling via

daf-2 RNA interference (RNAi) fulfils this prediction in *Caenorhabditis elegans* nematodes.

Long-lived *daf-2* RNAi parents showed normal fecundity as self-fertilizing hermaphrodites

and improved late-life reproduction when mated to males. Remarkably, the offspring of *daf-2*

RNAi parents had higher Darwinian fitness across three different genotypes. Thus, reduced

nutrient-sensing signalling in adulthood improves both parental longevity and offspring

fitness supporting the emerging view that sub-optimal gene expression in late-life lies at the

heart of ageing.

Impact Statement

Understanding mechanisms underpinning ageing is fundamental to improving quality of life

in an increasingly long-lived society. Recent breakthroughs have challenged the long-

standing paradigm that the energy trade-off between reproduction and somatic maintenance

causes organismal senescence via slow accumulation of unrepaired cellular damage with age.

The emerging new theory of ageing provides a conceptually novel framework by proposing

that ageing is a direct consequence of physiological processes optimized for early-life

function, such as growth and early-life reproduction, that are running 'too high' (i.e. at

hyperfunction) in late adulthood. Contrary to the classic view based on damage accumulation,

the hyperfunction theory proposes that suboptimal gene expression in late-life causes ageing

via excessive biosynthesis. Thus, the hyperfunction theory uniquely predicts that longevity

and Darwinian fitness can be simultaneously increased by reducing unnecessarily high levels

of nutrient-sensing signalling in adulthood. Here we show that reducing evolutionarily

conserved nutrient-sensing signalling pathway fulfils this prediction in *Caenorhabditis*

elegans nematodes. We found that downregulation of the insulin/IGF-1 signalling in adult *C.*

elegans nematodes not only improves longevity but, most intriguingly, increases fitness of

the resulting offspring in the next generation. We found support for increase in offspring

fitness across different genetic backgrounds. Our findings contradict the theoretical

conjecture that energy trade-offs between growth, reproduction and longevity is the universal

cause of senescence and provide strong experimental support for the emerging hyperfunction

theory of ageing.

Introduction

Understanding mechanisms underpinning healthy ageing is fundamental to improving quality of life in an increasingly long-lived society. The long-standing paradigm postulates that energy trade-offs between reproduction and somatic maintenance underlie organismal ageing (Kirkwood 1977, Kirkwood and Austad 2000, Kirkwood 2017). This theory is supported by a large number of studies in different taxa that reported a negative correlation between reproduction and survival (reviewed in Hughes and Reynolds 2005, Partridge et al. 2005, Edward and Chapman 2011, Flatt 2011, Nussey et al. 2013). However, the discoveries of environmental interventions that dramatically increase healthy lifespan in model organisms without the cost of reduced reproduction have challenged the current paradigm and suggested that our understanding of the evolution of ageing is incomplete (Dillin et al. 2002, Kenyon 2005, Antebi 2013, Gems and Partridge 2013, Maklakov and Immler 2016, Flatt and Partridge 2018). Specifically, experimental downregulation of nutrient-sensing insulin/IGF-like (IIS) signalling pathway that governs biosynthesis in response to nutrient availability can achieve increased longevity without a concomitant decrease in reproduction in model organisms (Dillin et al. 2002, Kenyon 2010, Kenyon 2011).

Since cost-free lifespan extension contradicts the traditional view of how ageing evolves, several studies investigated the fitness consequences of reduced IIS signalling (Gems et al. 1998, Walker et al. 2000, Jenkins et al. 2004, Savory et al. 2014, Maklakov et al. 2017). Indeed, mutations that reduce nutrient-sensing signalling during the whole life, as well as environmental interventions aimed at mimicking the mutational effect, often have detrimental pleiotropic effects on key life-history traits, such as development, growth rate, body size and early-life reproduction resulting in reduced Darwinian fitness even if total reproduction is

unaffected (Gems et al. 1998, Briga and Verhulst 2015). The first longevity mutant

discovered in *C. elegans*, *age-1*, is a good example because increased longevity, stress

resistance and late-life reproduction come at a cost of reduced early-life reproduction and

total individual fitness (Maklakov et al. 2017). Moreover, a recent literature survey suggests

that all classic longevity-extending mutations across taxa from worms to flies to mice

detrimentally affect life-history traits resulting in reduced fitness (Briga and Verhulst 2015).

Similarly, experimental evolution studies showed that when longevity and fecundity are

increased simultaneously through selection, the organisms pay the price in slow development

and delayed sexual maturation, again resulting in reduced individual fitness (Lind et al.

2017). These results support the theoretical conjecture that genes with antagonistically

pleiotropic effects between early-life and late-life fitness play an important role in the

evolution of ageing (Williams 1957). However, the mechanisms of antagonistic pleiotropy

(AP) remain elusive. The leading hypothesis, the "disposable soma" theory of ageing (DS)

suggests that ageing results from competitive energy allocation between somatic maintenance

and reproduction (Kirkwood 1977, Kirkwood and Holliday 1979, Kirkwood and Austad

2000). Indeed, increased reproductive performance in early-life correlates with reduced

survival and/or reduced performance in late-life in natural populations (Gustafsson and Part

1990, Boonekamp et al. 2014, Lemaitre et al. 2015) and in laboratory experiments (Rose

1984, Charlesworth 1993, Partridge et al. 1999, but see Chen and Maklakov 2012, Kimber

and Chippindale 2013, Chen and Maklakov 2014, Curtsinger 2019).

However, this hypothesis suffered several setbacks in recent years, with many empirical

studies challenging the importance of energy trade-offs in organismal senescence (reviewed

in Flatt 2011, Kenyon 2011, Antebi 2013, Gems and Partridge 2013, Maklakov and Immler

2016, Flatt and Partridge 2018). Instead, several authors proposed that ageing can result from

molecular signalling networks being optimized for development, growth and early-life

reproduction rather than for late-life reproduction and longevity (Blagosklonny 2010, Kenyon

2010, Antebi 2013, Gems and Partridge 2013, Ezcurra et al. 2018). For example, the

hyperfunction theory maintains that ageing is driven by excessive nutrient-sensing molecular

signalling in adulthood, which results in cellular hypertrophy leading to age-related

pathologies (Blagosklonny 2006, 2010, Ezcurra et al. 2018). These ideas can be traced back

to the original AP theory by George Williams, who suggested that the same physiological

processes that are beneficial for fitness early in life can become detrimental for organismal

fitness with age because of the reduced strength of natural selection on late-life function

(Williams 1957).

Williams's AP theory (Williams 1957) provides the population genetic framework for the

evolution of ageing via two different physiological routes: energy trade-offs (the "disposable

soma" theory of ageing) or functional trade-offs (e.g. "hyperfunction" theory of ageing) (Fig.

1). While both of these physiological theories rely on the same underlying principle, they

make uniquely distinct predictions with respect to reproduction costs of longevity. The

"disposable soma" theory maintains that organismal senescence is caused by slow

accumulation of unrepaired cellular damage with age because insufficient energy resources

are allocated to repair as organisms are maximising fitness rather than longevity (Kirkwood

1977, Kirkwood 2017). Therefore, the "disposable soma" theory predicts that increased

allocation of resources to somatic maintenance will increase longevity at the cost of reduced

resources available to current growth and reproduction. On the contrary, the functional trade-

off theory suggests that longevity is compromised by suboptimal physiology in adulthood

because, as discussed first by Williams (1957), selection is not strong enough to fully

optimize the age-specific expression of an allele whose effects are strongly beneficial in

early-life (e.g. during development) and slightly detrimental in late-life (e.g. during

adulthood). Consequently, the functional trade-off theory predicts that experimental

optimization of physiology in adulthood can increase longevity without any cost to

reproduction, or even simultaneously increase longevity and reproduction.

Because the main cost of longevity appears to be associated with reduced early-life function,

it seems plausible that age-specific modification of gene expression can potentially

circumvent this problem. In their landmark study, Dillin et al. (2002) used age-specific RNA

interference (RNAi) approach to knock down *daf-2* gene expression in *C. elegans* nematodes

across the life cycle of the worms. While early-life feeding with bacteria expressing *daf-2*

double-stranded (ds) RNA resulted in reduced early-life reproduction, there was no

detrimental effect of *daf-2* RNAi in adult worms, which enjoyed two-fold lifespan extension

without any cost to reproduction (Dillin et al. 2002). This study provides the strongest

support to date for the hypothesis that ageing results from molecular nutrient-sensing

signalling that is optimized for early-life function but is suboptimal for late-life function.

Nevertheless, while this study provided a powerful example for the cost-free lifespan

extension, it is possible that certain fitness costs have been overlooked. One possibility is that

fecundity costs become apparent only in mated hermaphrodites. In nature, *C. elegans* live in

populations with small (~0.3%) yet appreciable number of males living among self-fertilising

hermaphrodites with sometimes high levels of outcrossing (Sivasundar and Hey 2005), and

mating, as well as mere presence of male-derived pheromones, has pronounced effects of the

life-history of hermaphrodites (Maures et al. 2013, Shi and Murphy 2014, Aprison and

Ruvinsky 2016). While it is certainly interesting to consider how male-derived effects affect

resource allocation in hermaphrodites, it is unlikely that putative trade-off that is only visible

to selection in such rare circumstances can shape the evolution of ageing in this species.

Perhaps more importantly, it is possible that the fitness of the offspring and, therefore,

Darwinian fitness of the parents are compromised. The trade-off between offspring number

and quality is well known from a number of study systems (Stearns 1992), and is a potential

explanation for the apparent lack of fitness costs in the previous studies (Maklakov and

Immler 2016). Alternatively, longevity and Darwinian fitness can be simultaneously

increased by reducing unnecessarily high levels of nutrient-sensing signalling in adulthood.

To distinguish between these possibilities, we need to understand how reduction in nutrient-

sensing signalling in adulthood affects longevity, offspring number and offspring quality.

Here we show that *daf-2* RNAi in adult *C. elegans* results in increased offspring fitness

across three genetic backgrounds. We discuss these findings in the light of the emerging new

theories of ageing and suggest that they support the hypothesis that functional trade-offs

between early-life fitness and late-life fitness shape the evolution of ageing.

Materials and Methods

Strains

We used the *Caenorhabditis elegans* strains Bristol N2 wild-type (Brenner, Genetics 1974),

as well as the mutants *ppw-1(pk2505)* and *rrf-1(pk1417)*, obtained from Caenorhabditis

Genetics Center (CGC, Missouri, USA).

Maintenance

Before each assay, worms were recovered from freezing and synchronised by bleaching for

two generations to remove any freezing effects. The nematode populations were maintained

at 20°C and 60% relative humidity in an environmental test chamber. For regular

maintenance, the worms were kept on NGM agar supplemented with the antibiotics

streptomycin, kanamycin and nystatin (following Lionaki and Tavernarakis 2013), seeded

with the antibiotic-resistant *E. coli* strain OP50-1 (pUC4K).

Outline of the study

The study was run in three separate experiments. In the first experiment, we investigated

lifespan and reproduction of mated and unmated N2 hermaphrodites reared from sexual

maturity onwards on *daf-2* RNAi or empty vector (EV, control) plates. For logistic reasons,

this experiment was conducted in two blocks for mated worms and one block for unmated

worms. In the second experiment, we investigated the lifespan and egg size of unmated N2,

rrf-1(pk1417) and *ppw-1(pk2505)* hermaphrodites on raised from sexual maturity onwards on

daf-2 RNAi or EV plates. In the third experiment, we collected one egg from each parent at

their second day of adulthood (from *daf-2* RNAi and EV treatments) and investigated the

lifespan and reproduction of these offspring on control plates. Because different experiments

differed in setup time, daily reproduction values (and calculations based upon these, such as

λ_{ind}) are only meaningful for comparison between treatments within each experiment.

RNAi

RNase-III deficient, IPTG-inducible HT115 *Escherichia coli* bacteria with empty plasmid

vector (L4440) was used as control (Timmons et al. 2001) and the same HT115 bacteria with

daf-2 RNAi construct from the Vidal library was used as RNAi treatment. RNAi treatment

started from sexual maturity, and continued until the death of the individual. During the

experiments, worms were maintained on 35 mm NGM agar plates (supplemented with 1 mM

IPTG and 50 µg/ml ampicillin) seeded with 0.1 ml L4440 empty vector control or *daf-2*

bacteria grown in LB supplemented with 50 µg/ml ampicillin for 16-20 hours and seeded

(incubated) on the NGM agar plates again for 24 hours (following. Hinas et al. 2012).

Lifespan Assays

Lifespan assays were set up for all treatment combinations described above. In the lifespan

assays, the individual age-synchronised L4 worms were placed on separate 35 mm plates and

the plates were checked daily to record any instances of death. The surviving worms were

moved to new plates daily until their death. Fertile worms, which showed odd developmental

characteristics and low offspring numbers (<36 offspring), were excluded from the final

analysis (3 mated control worms and 7 mated *daf-2* worms).

Reproduction assays

Offspring production was scored in the reproduction assays using the same worms as those

scored for lifespan, except for the parental N2, *ppw-1* and *rrf-1* worms in the second

experiment, where only lifespan was recorded. Unmated individual hermaphrodites were

moved to new plates daily and scored for offspring produced 2.5 days later. In the "mated"

treatment, two male *C. elegans* (from the initial sample population of N2 strain) were placed

on a plate with a single hermaphrodite for two hours every day to allow time for mating.

Offspring production was scored 2.5 days later, as in the "unmated" treatment.

Egg size assays

Egg size was measured in N2, *ppw-1* and *rrf-1* strains (unmated hermpahrodites) growing on

either *daf-2* RNAi or empty vector (EV) plates. Two days after maturation, worms were

placed individually on new plates and observed continually during five hours for the presence

of newly laid eggs, of which the first two eggs were collected. Eggs were picked immediately

after laying and placed under a Leica M165C microscope set on 12x magnification; photos

were taken using a Lumenera Infinity 2-6C digital microscope camera. Egg size was analysed

from photos using *ImageJ* (https://imagej.nih.gov/ij/). Only eggs laid during gastrulation

stage (the normal developmental stage at egg laying) were included in the analyses.

Statistical analyses

Survival was analysed for each experiment in Cox proportional hazard models in *R 3.3.3*.

Mated (EV: n=72, *daf-2* n=68) and unmated (n=25 per treatment) individuals were analysed

separately, as they were run in different blocks. Unmated individuals were analysed using the

coxph function in the package *survival*, with *daf-2* RNAi treatment as a fixed factor. For

mated individuals, we used the *coxme* package in order to fit block as a random effect, in

addition to the fixed effect of RNAi treatment. In the second experiment (n=25 per

treatment), in addition to RNAi treatment, we also fitted the fixed factor strain (N2, *ppw-1*,

rrf-1) and its interaction with treatment using the *coxph* function in the *survival* package.

Reproduction was analysed as total reproduction as well as rate-sensitive individual fitness

λ_{ind}, which encompasses the timing and number of offspring (Brommer et al. 2002, Lind et al.

2016). λ_{ind} is estimated by solving the Euler-Lotka equation for each individual using the

lambda function in the *popbio* package and is analogous to the intrinsic rate of population

growth (Stearns 1992). For all unmated worms (n=25 per treatment), we estimated the fixed

effect of treatment (*daf-2* RNAi or empty vector). For offspring of the three mutants (n=25

per treatment), we also estimated the fixed effect or strain, using linear models. For the mated

worms (EV: n=72, *daf-2* n=68), we also estimated the random effect of block, in addition to

RNAi treatment. These models were implemented as mixed effect models using the *lme4*

package in *R 3.3.3*, and chi-square tests of fixed effects were performed using the *car*

package. Egg size was analysed in a mixed effect model in *lme4*, treating strain and RNAi

treatment as crossed fixed effects, and parent ID as well as block as random effects. We

obtained the following n: N2 on EV: 56, N2 on *daf-2*: 54, *ppw-1* on EV: 44, *ppw-1* on *daf-2*:

42, *rrf-1* on EV: 59, *rrf-1* on *daf-2*: 42.

Results

First off, we confirmed that *daf-2* RNAi significantly extended the lifespan of unmated N2

wild-type hermaphrodite worms (censoring matricide: $z = -4.94$, df = 1, $p < 0.001$, Fig. 2A;

including matricide as dead: $z = -4.97$, df = 1, $p < 0.001$), as expected from previous studies

(Dillin et al. 2002). In addition, for mated N2, *daf-2* RNAi extended lifespan when matricide

was censored ($z = -2.42$, df = 1, $p = 0.016$, Fig. 2B) but not if matricidal worms were

included as dead ($z = 0.16$, df = 1, $p = 0.87$) because of an increase in matricide in the late

reproducing mated *daf-2* RNAi N2.

We did not find any effect of *daf-2* RNAi on total reproduction (unmated: $F = 0.32$, df = 1, p

= 0.58; mated: $\chi^2 = 1.11$, df = 1, $p = 0.29$) or individual fitness λ_{ind} (unmated: $F = 0.30$, df = 1,

$p = 0.59$; mated: $\chi^2 = 0.43$, df = 1, $p = 0.51$) for neither unmated nor mated N2 (Table 1, Fig.

3). However, *daf-2* RNAi had a positive effect on late (day 5+) reproduction for mated

hermaphrodites ($\chi^2 = 24.76$, df = 1, $p < 0.001$, Fig. 3B).

In a second experiment, using unmated hermaphrodites only, we investigated the effect of *daf-2* RNAi on parent lifespan and offspring lifespan and reproduction across three genetic backgrounds (N2 wild-type and the mutants *ppw-1* and *rrf-1*, that are deficient for germline and somatic RNAi, respectively). Parental treatment with *daf-2* RNAi increased lifespan across all genetic backgrounds, both when matricide was censored (treatment: $\chi^2 = 90.39$, df $= 1$, p <0.001; strain: $\chi^2 = 21.8$, df $= 2$, p <0.001; treatment × strain: $\chi^2 = 10.46$, df $= 2$, p $=$ 0.005, Fig. 2A) and included as dead (treatment: $\chi^2 = 85.25$, df $= 1$, p <0.001; strain: $\chi^2 =$ 20.45, df $= 2$, p <0.001; treatment × strain: $\chi^2 = 9.43$, df $= 2$, p $= 0.009$). In accordance with previously published research (Hibshman et al. 2016), parental *daf-2* RNAi increased egg size (treatment: $\chi^2 = 5.11$, df $= 1$, p $=0.024$; strain: $\chi^2 = 13.89$, df $= 2$, p <0.001; treatment × strain: $\chi^2 = 2.68$, df $= 2$, p $= 0.262$, Fig. 4). However, we found that the effect was most pronounced in N2 wildtype worms, and relatively weak in both somatic and germline *daf-2* knockdown (see Fig. 4), suggesting that *daf-2* knockdown in both somatic and reproductive tissues is required to maximize the effect on egg size.

Parental *daf-2* RNAi treatment did not, however, influence the lifespan of their offspring, neither when matricidal worms were censored (treatment: $\chi^2 = 0.04$, df $= 1$, p $= 0.85$; strain: $\chi^2 = 24.2$, df $= 2$, p <0.001; treatment × strain: $\chi^2 = 0.61$, df $= 2$, p $= 0.74$, Fig. 5A) nor when included as dead (treatment: $\chi^2 = 0.01$, df $= 1$, p $= 0.92$; strain: $\chi^2 = 21.8$, df $= 2$, p <0.001; treatment × strain: $\chi^2 = 0.48$, df $= 2$, p $= 0.79$).

In contrast, parental *daf-2* RNAi treatment significantly increased offspring total reproduction (treatment: F $= 15.9$, df $= 1$, p <0.001; strain: F $= 33.7$, df $= 2$, p <0.001; treatment × strain: F $= 0.09$, df $= 2$, p $= 0.91$, Fig. 5B-C) and individual fitness λ_{ind} (treatment: F $= 11.8$, df $= 1$, p

<0.001; strain: F = 13.1, df = 2, p <0.001; treatment × strain: F = 0.18, df = 2, p = 0.84, Fig.

5D) across all genetic backgrounds. Importantly, there was no correspondence between the

effect of parental *daf-2* RNAi on egg size (see above) and offspring total reproduction /

individual fitness, suggesting that factors beyond the amount of resources in the egg

contribute to increased fitness of offspring of *daf-2* RNAi parents.

Discussion

The "disposable soma" theory of ageing postulates that senescence results from competitive

energy allocation between key life-history traits, such as growth, reproduction and somatic

maintenance (Kirkwood 1977, Kirkwood 2017). This theory predicts that genetic and

environmental manipulations that increase energy allocation to growth and somatic

maintenance will result in detrimental effects to reproduction. This is why the findings by

Dillin et al. (2002), which suggested that adult-only downregulation of insulin/IGF-1 by *daf-2*

RNAi can substantially increase lifespan without any detrimental effect to reproduction, were

subsequently scrutinized in an attempt to find the hidden costs of longevity (Jenkins et al.

2004, Partridge et al. 2005). Nonetheless, both the original findings (Dillin et al. 2002) and

our results here, suggest that adult-only *daf-2* RNAi can more than double longevity without

any negative effect on reproduction. Moreover, when supplied with sperm from males, *daf-2*

RNAi-treated parents have improved fecundity in late-life. However, the key question that we

asked in this study was whether treatments that improve parental performance have positive

or negative effects on their offspring. The trade-off between offspring number and offspring

quality is a well known concept in life-history evolution (Stearns 1992) but is rarely

considered in biogerontological research (reviewed in Maklakov and Immler 2016). Germline

maintenance is costly (Sniegowski et al. 2000, Agrawal and Wang 2008, Maklakov and

Immler 2016, Berger et al. 2017), and increased investment into somatic maintenance can, in

theory, result in increased mutation rate and reduced fitness of progeny.

Alternatively, it is possible that instead of energy trade-offs, the evolution of senescence is

governed by functional trade-offs. Functional trade-offs can occur because the physiological

requirements of a young organism can differ substantially from those of a mature one

(Williams 1957). In his classic 1957 paper, George Williams (Williams 1957) described a

hypothetical example of a mutation that positively affects bone calcification in a developing

young organism but increases calcification of the connective tissues of arteries in a mature

one with detrimental consequences. More recently, it has been suggested that nutrient sensing

IIS/TOR molecular signalling pathways that govern growth and development result in

excessive biosynthesis in late-life leading to different pathologies and increased mortality

(Blagosklonny 2006, 2010, Gems and Partridge 2013, Ezcurra et al. 2018). These proximate

explanations rest on the fundamental assumption that the strength of natural selection

declines with age because of environmental mortality from a range of biotic and abiotic

hazards (e.g. predation, pathogens, competition, starvation) (Williams 1957). Because of such

environmental mortality, optimizing development, growth and instantaneous reproduction

may be more important for organismal fitness that optimizing future survival and

reproduction (Williams 1957, Hamilton 1966). Thus, progressively weakening selection in

adulthood may result in suboptimal levels of IIS/TOR signalling leading to pathology and

senescence (Ezcurra et al. 2018). However, unlike the classic energy trade-off theory, the

functional trade-off hypothesis predicts that it should be possible to modify adult physiology

to improve both longevity and fitness.

Here we found that reduced insulin/IGF-1 signalling in adult worms not only improved

longevity, but also increased reproduction and Darwinian fitness of the resulting offspring in

three different genetic backgrounds. This result contradicts the hypothesis that improved

longevity and postponed ageing of *daf-2* RNAi parents comes at the cost of offspring fitness.

Instead, our findings are in line with the hypothesis that suboptimal levels of nutrient-sensing

signalling in adult life accelerate ageing, curtail lifespan and reduce individual fitness. This

result was not caused by direct inheritance of *daf-2* RNAi, since we did not recover the

lifespan extension effect of *daf-2* knockdown in these offspring. Because *daf-2* is essential for

successful development and growth of a young worm (Dillin et al. 2002), these results

suggest that wildtype *C. elegans* nematodes trade-off improved pre-adult performance for

reduced offspring quality. Such trade-offs are at heart of AP theory, but are usually

interpreted as evidence for energy-based trade-offs. Our results clearly demonstrate that this

is not the case, and that adulthood-only *daf-2* RNAi increases offspring fitness. In summary,

our findings suggest that selection on expression of *daf-2* in adulthood is not sufficiently

strong in nature. We predict that such effects may be very common, and suggest that future

studies should aim to quantify the fitness consequences of experimental manipulation of age-

specific gene expression across a broad range of taxa. Such approach will allow us to

estimate the relative importance of energy trade-offs versus functional trade-offs in the

evolution of ageing across the tree of life.

Because previous research found that both dietary restriction and reduction in insulin-like

signalling by *daf-2* RNAi knockdown increased embryo size in *C. elegans* nematodes

(Hibshman et al. 2016), we replicated these results to test whether increased fitness of adult

progeny results from increased resource allocation to eggs by *daf-2* RNAi mothers. While

daf-2 knockdown increased egg size to a different degree in N2, *ppw-1* and *rrf-1* strains, there

was no correlation between the effect of parental *daf-2* RNAi on egg size and offspring

reproductive performance. We provisionally conclude that increased egg size under reduced

maternal insulin-like signalling can contribute to increased offspring fitness, but it is likely

not the sole source of variation in this trait. Recent work has identified oocyte-specific IIS

targets that are different from soma-specific IIS targets suggesting that IIS signalling

regulates reproduction and longevity through different mechanisms (Templeman et al. 2018).

In the future, it will be interesting to test for individual fitness of offspring produced via

genetic manipulation of oocyte-specific targets of IIS signalling pathway. In recent years,

there has been a vigorous debate regarding whether mechanistic understanding of life-history

trade-offs is necessary to advance life-history theory (Flatt and Heyland 2011, Stearns 2011a,

b). Here we used the mechanistic approach to separate between two conceptually different

evolutionary theories of ageing – energy trade-offs and functional trade-offs – in an empirical

study. We argue that unification between the conceptual approach and the mechanistic

understanding may often prove fruitful in this regard.

Author contributions

MIL and AAM designed the study, with the aid of AH. SR, ZS, MIL and HC collected the

data, MIL analysed the data, MIL and AAM drafted the manuscript. All authors contributed

to the revision of the manuscript.

Acknowledgements

This work was supported by ERC Consolidator Grant 724909 GermlineAgeingSoma to AAM

and Swedish Research Council VR Grant 2016-05195 to MIL. The authors declare no

conflicts of interest.

Data accessibility

402   Upon acceptance, the data will be archived at Dryad.

References

Agrawal, A. F., and A. D. Wang. 2008. Increased transmission of mutations by low-condition
females: Evidence for condition-dependent DNA repair. Plos Biology **6**:389-395.

Antebi, A. 2013. Regulation of longevity by the reproductive system. Experimental
Gerontology **48**:596-602.

Aprison, E. Z., and I. Ruvinsky. 2016. Sexually antagonistic male signals manipulate
germline and soma of *C. elegan*s hermaphrodites. Current Biology **00**:XXX-XXX.

Berger, D., J. Stangberg, K. Grieshop, I. Martinossi-Allibert, and G. Arnqvist. 2017.
Temperature effects on life-history trade-offs, germline maintenance and mutation
rate under simulated climate warming. Proceedings of the Royal Society B-Biological
Sciences **284**.

Blagosklonny, M. V. 2006. Aging and immortality - Quasi-programmed senescence and its
pharmacologic inhibition. Cell Cycle **5**:2087-2102.

Blagosklonny, M. V. 2010. Revisiting the antagonistic pleiotropy theory of aging TOR-
driven program and quasi-program. Cell Cycle **9**:3151-3156.

Boonekamp, J. J., M. Salomons, S. Bouwhuis, C. Dijkstra, and S. Verhulst. 2014.
Reproductive effort accelerates actuarial senescence in wild birds: an experimental
study. Ecology Letters **17**:599-605.

Briga, M., and S. Verhulst. 2015. What can long-lived mutants tell us about mechanisms
causing aging and lifespan variation in natural environments? Experimental
Gerontology **71**:21-26.

Brommer, J. E., J. Merila, and H. Kokko. 2002. Reproductive timing and individual fitness.
Ecology Letters **5**:802-810.

Charlesworth, B. 1993. Evolutionary mechanisms of senescence. Genetica **91**:11-19.

Chen, H.-y., and A. A. Maklakov. 2012. Longer lifespan evolves under high rates of

condition dependent mortality. Current Biology **22**:2140-2143.

Chen, H.-y., and A. A. Maklakov. 2014. Condition dependence of male mortality drives the

evolution of sex differences in longevity. Current Biology **24**:2423-2427.

Curtsinger, J. W. 2019. Fecundity for free? Enhanced oviposition in longevous populations of

Drosophila melanogaster. Biogerontology.

Dillin, A., D. K. Crawford, and C. Kenyon. 2002. Timing requirements for insulin/IGF-1

signaling in *C-elegans*. Science **298**:830-834.

Edward, D. A., and T. Chapman. 2011. Mechanisms underlying reproductive trade-offs:

Costs of reproduction.

Ezcurra, M., A. Benedetto, T. Sornda, A. F. Gilliat, C. Au, Q. Zhang, S. van Schelt, A. L.

Petrache, H. Wang, Y. de la Guardia, S. Bar-Nun, E. Tuler, M. J. Wakelam, and D.

Gems. 2018. *C. elegans* eats its own intestine to make yolk leading to multiple

senescent pathologies. Current Biology **28**:2544-2556.

Flatt, T. 2011. Survival costs of reproduction in *Drosophila*. Experimental Gerontology

46:369-375.

Flatt, T., and A. Heyland. 2011. Why mechanisms are important for life-history theory: A

response by Flatt and Heyland. Pages 375-379 *in* T. Flatt and A. Heyland, editors.

Mechanisms of Life History Evolution: The Genetcis and Physiology of Life History

Traints and Trade-Offs. Oxford University Press, Oxford.

Flatt, T., and L. Partridge. 2018. Horizons in the evolution of aging. BMC Biology **16**:93.

Gems, D., and L. Partridge. 2013. Genetics of longevity in model organisms: debates and

paradigm shifts. Annual Review of Physiology **75**:621-644.

Gems, D., A. J. Sutton, M. L. Sundermeyer, P. S. Albert, K. V. King, M. L. Edgley, P. L.

Larsen, and D. L. Riddle. 1998. Two pleiotropic classes of daf-2 mutation affect

larval arrest, adult behavior, reproduction and longevity in Caenorhabditis elegans.

Genetics **150**:129-155.

Gustafsson, L., and T. Part. 1990. Acceleration of senescence in the collared flycatcher

Ficedula albicollis by reproductive costs. Nature **347**:279-281.

Hamilton, W. D. 1966. The moulding of senescence by natural selection. Journal of

Theoretical Biology **12**:12-45.

Hibshman, J. D., A. Hung, and L. R. Baugh. 2016. Maternal diet and insulin-like signaling

control intergenerational plasticity of progeny size and starvation resistance. PLoS

Genetics **DOI: 10.1371/journal.pgen.1006396**.

Hinas, A., A. J. Wright, and C. P. Hunter. 2012. SID-5 is an endosome-associated protein

required for efficient systemic RNAi in *C. elegans*. Current Biology **22**:1938-1943.

Hughes, K. A., and R. M. Reynolds. 2005. Evolutionary and mechanistic theories of aging.

Annual Review of Entomology **50**:421-445.

Jenkins, N. L., G. McColl, and G. J. Lithgow. 2004. Fitness cost of extended lifespan in

Caenorhabditis elegans. Proceedings of the Royal Society of London Series B-

Biological Sciences **271**:2523-2526.

Kenyon, C. 2005. The plasticity of aging: Insights from long-lived mutants. Cell **120**:449-

460.

Kenyon, C. 2011. The first long-lived mutants: discovery of the insulin/IGF-1 pathway for

ageing. Philosophical Transactions of the Royal Society B-Biological Sciences **366**:9-

16.

Kenyon, C. J. 2010. The genetics of ageing. Nature **464**:504-512.

Kimber, C. M., and A. K. Chippindale. 2013. Mutation, condition and the maintenance of

extended lifespan in *Drosophila*. Current Biology **23**: 2283-2287.

Kirkwood, T. B. L. 1977. Evolution of aging. Nature **270**:301-304.

Kirkwood, T. B. L. 2017. The Disposable Soma Theory of Ageing. Pages 23-39 *in*

Shefferson RP, Jones OR, and S.-G. R, editors. The Evolution of Senescence in the

Tree of Life. Cambridge University Press, Cambridge, United Kingdom.

Kirkwood, T. B. L., and S. N. Austad. 2000. Why do we age? Nature **408**:233-238.

Kirkwood, T. B. L., and R. Holliday. 1979. Evolution of aging and longevity. Proceedings of

the Royal Society of London Series B-Biological Sciences **205**:531-546.

Lemaitre, J.-F., V. Berger, C. Bonenfant, M. Douhard, M. Gamelon, F. Plard, and J.-M.

Gaillard. 2015. Early-late life trade-offs and the evolution of ageing in the wild.

Proceedings of the Royal Society B-Biological Sciences **282**.

Lind, M. I., H. Y. Chen, S. Meurling, A. C. G. Gil, H. Carlsson, M. K. Zwoinska, J.

Andersson, T. Larva, and A. A. Maklakov. 2017. Slow development as an

evolutionary cost of long life. Functional Ecology **31**:1252-1261.

Lind, M. I., M. K. Zwoinska, S. Meurling, H. Carlsson, and A. A. Maklakov. 2016. Sex-

specific trade-offs with growth and fitness following lifespan extension by rapamycin

in an outcrossing nematode, *Caenorhabditis remanei*. Journals of Gerontology Series

a-Biological Sciences and Medical Sciences **71**:882-890

Lionaki, E., and N. Tavernarakis. 2013. High-throughput and longitudinal analysis of aging

and senescent decline in *Caenorhabditis elegans*. Pages 485-500 *in* Lorenzo Galluzzi,

Ilio Vitale, Oliver Kepp, and G. Kroemer, editors. Cell Senescence.

Maklakov, A. A., H. Carlsson, P. Denbaum, M. I. Lind, B. Mautz, A. Hinas, and S. Immler.

2017. Antagonistically pleiotropic allele increases lifespan and late-life reproduction

at the cost of early-life reproduction and individual fitness. Proceedings of the Royal

Society B-Biological Sciences **284**.

Maklakov, A. A., and S. Immler. 2016. The expensive germline and the evolution of ageing.

Current Biology **26**:R577-R586.

Maures, T. J., L. N. Booth, B. A. Benayoun, Y. Izrayelit, F. C. Schroeder, and A. Brunet.

2013. Males shorten the life span of *C. elegans* hermaphrodites via secreted

compounds. Science **1244160**.

Nussey, D. H., H. Froy, J. F. Lemaitre, J. M. Gaillard, and S. N. Austad. 2013. Senescence in

natural populations of animals: Widespread evidence and its implications for bio-

gerontology. Ageing Research Reviews **12**:214-225.

Partridge, L., D. Gems, and D. J. Withers. 2005. Sex and death: What is the connection? Cell

120:461-472.

Partridge, L., N. Prowse, and P. Pignatelli. 1999. Another set of responses and correlated

responses to selection on age at reproduction in *Drosophila melanogaster*.

Proceedings of the Royal Society of London Series B-Biological Sciences **266**:255-

261.

Rose, M. R. 1984. Laboratory evolution of posponed senescence in *Drosophila-*

melanogaster. Evolution **38**:1004-1010.

Savory, F. R., T. G. Benton, V. Varma, I. A. Hope, and S. M. Sait. 2014. Stressful

environments can indirectly select for increased longevity. Ecology and Evolution

4:1176-1185.

Shi, C., and C. T. Murphy. 2014. Mating induces shrinking and death in *Caenorhabditis*

mothers. Science **Express**:1-7.

Sivasundar, A., and J. Hey. 2005. Sampling from natural populations with RNAi reveals high

outcrossing and population structure in *Caenorhabditis elegans*. Current Biology

15:1598-1602.

Sniegowski, P. D., P. J. Gerrish, T. Johnson, and A. Shaver. 2000. The evolution of mutation

rates: separating causes from consequences. Bioessays **22**:1057-1066.

Stearns, S. C. 1992. The Evolution of Life Histories. Oxford University Press, New York.

Stearns, S. C. 2011a. Does impressive progress on understanding mechanisms advance life

history theory? Pages 365-373 *in* T. Flatt and A. Heyland, editors. Mechanisms of

Life History Evolution: The Genetcis and Physiology of Life History Traints and

Trade-Offs. Oxford University Press, Oxford.

Stearns, S. C. 2011b. Reply by Stearns: Mechanisms do not yet force the theory to change.

Page 379 *in* T. Flatt and A. Heyland, editors. Mechanisms of Life History Evolution:

The Genetcis and Physiology of Life History Traints and Trade-Offs. Oxford

University Press, Oxford.

Templeman, N. M., S. J. Luo, R. Kaletsky, C. Shi, J. Ashraf, W. Keyes, and C. T. Murphy.

2018. Insulin signaling regulates oocyte quality maintenance with age via Cathepsin B

activity. Current Biology **28**:753-+.

Timmons, L., D. L. Court, and A. Fire. 2001. Ingestion of bacterially expressed dsRNAs can

produce specific and potent genetic interference in *Caenorhabditis elegans*. Gene

263:103-112.

Walker, D. W., G. McColl, N. L. Jenkins, J. Harris, and G. J. Lithgow. 2000. Natural

selection - evolution of lifespan in *C-elegans*. Nature **405**:296-297.

Williams, G. C. 1957. Pleiotropy, natural selection, and the evolution of senescence.

Evolution **11**:398-411.

Table 1. The effect of daf-2 RNAi on reproduction. Total reproduction and individual

fitness (λ_{ind}) for unmated and mated *C. elegans* N2 wild-type treated with either empty vector

(Control) or *daf-2* RNAi from adulthood onwards. All values expressed as mean ± SE.

	Total reproduction		Fitness (λ_{ind})	
RNAi treatment	unmated	mated	unmated	mated
Control	311.0 ± 7.0	595.7 ± 24.3	4.66 ± 0.03	4.47 ± 0.05
daf-2 RNAi	317.4 ± 8.9	630.5 ± 22.2	4.63 ± 0.05	4.50 ± 0.05

Fig. 1. Theoretical framework. The relationship between population genetic theory of

ageing ("antagonistic pleiotropy", Williams 1957) and physiological theories of ageing based

on either energy trade-offs ("disposable soma" (Kirkwood 1977) or functional trade-offs (e.g.

"hyperfunction" (Blagosklonny 2006)). Note that Williams (1957) was the first to provide an

abstract example of a functional trade-off between early-life and late-life allelic effects on

organismal physiology. Recently, Blagosklonny (2006, 2010) put forward a "hyperfunction"

hypothesis that specifically links suboptimal levels of nutrient-sensing signalling to excessive

biosynthesis (hence, hyperfunction) leading to cellular and organismal senescence.

Fig. 2. The effect of *daf-2* RNAi on lifespan. Survival probability for (**A**) unmated or (**B**)

mated N2 wild-type (purple), *ppw-1* (black) and *rrf-1* (orange) mutants, treated with either

daf-2 RNAi (solid lines, filled symbols) or control empty vector (broken lines, open symbols)

from adulthood onwards.

Fig. 3. The effect of *daf-2* RNAi on reproduction. Daily offspring number for (**A**) unmated

or (**B**) mated N2 wild-type worms, treated with either *daf-2* RNAi (solid lines, filled symbols)

or control empty vector (broken lines, open symbols) from adulthood onwards. Symbols

represent mean ± SE.

Fig. 4. The effect of *daf-2* RNAi on egg size. Egg size of unmated, parental worms. N2 wild-

type (purple), *ppw-1* (black) and *rrf-1* (orange) mutants, treated with either *daf-2* RNAi (solid

 lines, filled symbols) or control empty vector (broken lines, open symbols) from adulthood

 onwards. Symbols represent mean ± SE.

579
Fig. 5. The effect of parental *daf-2* RNAi on offspring survival and reproduction.

Offspring worms, unmated, on control (empty vector) plates from parents exposed to *daf-2*

RNAi or control treatment. (**A**) Survival probability, (**B**) daily offspring number, (**C**) Total

reproduction and (**D**) individual fitness (λ_{ind}) of offspring (on control plates) from parents

either exposed to *daf-2* RNAi (solid lines, filled symbols) or control empty vector (broken

lines, open symbols). The colors reflect N2 wild-type (purple), *ppw-1* (black) and *rrf-1*

(orange) mutants. Symbols represent mean ± SE.