iDEG: a single-subject method utilizing local estimates of dispersion to impute differential expression between two transcriptomes

Qike Li[1,2,4,§], Samir Rachid Zaim[1,2,4, §], Dillon Aberasturi[1,2,4], Joanne Berghout[1-3], Haiquan Li[1,2,4], Francesca Vitali[1,2], Colleen Kenost[1,2,5], Helen Hao Zhang[4,7*], Yves A. Lussier[1-6*]

1. Center for Biomedical Informatics and Biostatistics(CB2)[1]
2. Department of Medicine[2],
3. The Center for Applied Genetics and Genomic Medicine[3],
4. Graduate Interdisciplinary Program in Statistics[4],
5. BIO5 Institute[5], University of Arizona Cancer Center[6],
6. Department of Mathematics[7],
7. The University of Arizona, Tucson, AZ 85721, USA

* Corresponding authors
§ Contributed equally

Email of all authors: hzhang@math.arizona.edu, daberasturi@email.arizona.edu, haiquan@email.arizona.edu, ckenost@email.arizona.edu, hzhang@math.arizona.edu, yves@email.arizona.edu

This manuscript has been submitted to the TBC conference and is not under consideration elsewhere. In addition, all authors have reviewed it.

Abstract

Background: Accurate profiling of gene expression in a single subject has the potential to be a powerful precision medicine tool, useful for unveiling individual disease mechanisms and responses. However, expression analysis tools for RNA-sequencing (RNA-Seq) data require replicate samples to estimate gene-wise data variability and make inferences, which is costly and not easily obtainable in clinical practice. Strategies to implement DEGSeq, DESeq, and edgeR for comparing two conditions without replicates (TCWR) have been proposed without evaluation, while NOISeq-sim was validated in a restricted way using qPCR on 400 transcripts. These methods impose restrictive assumptions in TCWR limiting inferential opportunities.

Methods: We propose a new method that borrows information across different genes from the same individual using a partitioned window to strategically bypass the requirement of replicates per condition. We termed this method "iDEG", which identifies <u>i</u>ndividualized <u>D</u>ifferentially <u>E</u>xpressed <u>G</u>enes in a single subject sampled under two conditions without replicates, i.e., a baseline sample (unaffected tissue) vs. a case sample (tumor). iDEG transforms RNA-Seq data such that, under the null hypothesis, differences of transformed expression counts follow a distribution and variance calculated across a local partition of related transcripts at baseline expression. This transformation enables modeling genes with a two-group mixture model from which the probability of differential expression for each gene is then estimated by an empirical Bayes approach with a local false discovery rate control. To compare the performance of iDEG to other methods applied to TCWR, we conducted simulations assuming a Negative Binomial distribution with varying dispersion parameters and percentages of differentially expressed genes (DEGs).

Results: Our extensive simulation studies demonstrate that iDEG's F_1 accuracy scores better than the other methods at 5%<DEGs<20%, while NOISeq is superior for 30%<DEGs<40% - and comparable in between. DEGseq achieves the largest recall and smallest precision. NOISeq obtains ~90% precision and recall at DEG~40% (the best F_1-scores). iDEG obtained a precision>90% and recall>75% and low false positive rate (<1%) in most conditions.

Conclusion: The partitioned window strategy provides a novel and accurate way to borrow information across genes locally and would probably increase the accuracy of all relevant methods.

Software available at:
http://www.lussiergroup.org/publications/iDEG

Keywords: transcriptome, mRNA expression, RNA-seq, differentially expressed genes, single-subject, N-of-1, RNA-Seq, iDEG.

Background

Precision medicine aims to deliver "the right treatments, at the right time, to the right person" (1). However, clinical research, medicine, and pharmacology need new tools to achieve that goal. The prevailing system of one-size-fits-all drug development has led to the ten top-grossing USA drugs being ineffective for more than 75% of users(2), and these patients typically cannot be identified until after therapeutic failure has occurred. The success of precision medicine hinges on identifying the precise aberrant mechanisms at play during an individual's disease course(3) to optimize treatment based on that individual's biology.

Single-subject RNA sequencing (**RNA-Seq**) analysis considers one patient at a time, with the goal of revealing an individual's altered transcriptomic mechanisms. Relative to traditional cohort-based analyses, a major challenge of single-subject RNA-Seq analysis is the estimation of gene expression variance which is required to identify differentially expressed genes (**DEGs**). In cohort-based methods, gene variance is calculated across a heterogenous set of samples, and the statistical methods employed leverage and rely on those replicates. However, they also emphasize consistent and average responses which may not accurately represent a single patient when the disease is heterogenous or stratified. Alternatively, the variance can be assessed between two conditions in one subject and three replicates. Yet, obtaining sufficient isogenic replicates for one subject to answer more precision questions poses a major difficulty due to (i) limited tissue availability, (ii) the risks associated with invasive tissue-sampling procedures, and (iii) general costs and inefficiencies with the current technology. Even though there is a great body of work for identifying DEGs in RNA-Seq data(4-8) and frameworks for N-of-1 studies either for a single analyte or by pooling gene products in pathways(9-13), to the best of our knowledge, no methods have been designed or validated at the gene level to determine the effect size and statistical significance of a single-subject, single RNA-Seq studies in two conditions without replicates (**TCWR**)(14). Strategies to implement standard RNA-seq analysis methods for comparing TCWR have been proposed without comprehensive evaluation. Typically, these standard methods, usually requiring large cohorts, have been adapted to identify DEGs in TCWR by imposing restrictive assumptions or new parameters, thus limiting inferential opportunities.

Three critical obstacles hinder the analysis of single-subject TCWR studies. These include i) *patient-level inferential capability* in absence of biological replicates, ii) sensitivity to *fold-change inflation* in low-expression genes, and iii) *rigid parametric data assumptions* for variance estimation. We propose a new method that borrows information across different genes from the same individual using a partitioned window to strategically bypass the requirement of replicates per condition. To bridge the knowledge gap and develop a technique that can overcome the current technical limitations in analyzing RNA-Seq data, we developed **iDEG** (**i**ndividualized **D**ifferentially **E**xpressed **G**enes). iDEG is a novel RNA-Seq analysis tool designed to analyze paired transcriptomic samples from a single individual to effectively compare case vs. baseline measurements (e.g., tumor vs. healthy tissue) in the absence of biological replicates. iDEG applies a localized variance-stabilizing transformation to estimate a gene's distribution that borrows information from genes with similar baseline expression. While variance-stabilizing transformation has been previously used to identify DEGs across a large number of subjects or replicates, our approach differs from these since it has been developed to be applied directly on two paired transcriptomes from a single subject by computing the localized dispersion parameters in different windows of genes with similar expression at the baseline.

In this work, we evaluated the performances of iDEG and other four standard approaches applied to single-subject TCWR studies. We also designed simulation studies under several conditions to stratify the

range of applicability of our proposed strategy, which could eventually complement other RNA-seq analyses in TCWR studies. This study demonstrates the utility of variance-stabilizing transformations within subject in absence of replicates in two conditions, which is distinct from previous implementations of variance-stabilizing methods conducted across replicates or subjects.

Methods

The iDEG algorithm: iDEG

The iDEG algorithm (**Figure 1**) is an easy-to-implement, single-command function written in R(15) with a computation speed of one second for identifying a subject's DEGs on 8GB Ram computer. The subsequent sections expand on the main iDEG steps shown in **Figure 1**.

Fig. 1. The iDEG Algorithm. 1) Normalize unequal library sizes if necessary. 2) Partition transcriptome into percentile-based windows using ranked baseline expression. 3) For each window: estimate mean expression, variance, and dispersion parameters. 4) Apply the Variance Stabilizing Transformation (**VST**) for each gene expression count. 5) Calculate the standard normal summary statistic "Z_g" for each gene expression count "g". 6) Determine the identified DEG set "$\hat{\mathcal{G}}$" based on a pre-determined α-cutoff.

Modeling read counts via a re-parameterized Negative Binomial (NB) distribution

We model read counts Y_{gd} as following a re-parameterized negative binomial distribution with mean μ_{gd} and dispersion δ_g. Thus, $Y_{gd} \sim NB\left(\mu_{gd}, \delta_g\right)$ with the following probability mass function, mean, and variance, respectively. Since for any subject, both transcriptomes are sequenced separately, they are treated as conditionally independent, conditional on the subject[*].

$$P\left(y_{gd}|\mu_{gd}\delta_g\right) = \left(1 + \delta_g\mu_{gd}\right)^{-\frac{1}{\delta_g}} \frac{\Gamma\left(y_{gd}+\frac{1}{\delta_g}\right)}{y_{gd}!\,\Gamma\left(\frac{1}{\delta_g}\right)}\left(\frac{\delta_g\mu_{gd}}{1+\delta_g\mu_{gd}}\right)^{y_{gd}}; E\left(Y_{gd}\right) = \mu_{gd},\ Var\left(Y_{gd}\right) = \mu_{gd} + \delta_g\mu_{gd}^2 \text{ (\textbf{Eq.1})}$$

---

[*] A note on notation, since iDEG models each patient's paired transcriptome individually, the subscripts for each subject are omitted since only a subject is handled at a time in any given calculation.

To identify DEGs from a pair of transcriptomes, we must test multiple hypotheses $H_0: \mu_{g1} = \mu_{g2}, g = 1,$ ... , G, where μ_{g1} and μ_{g2} are the theoretical mean expression levels for each gene "g" in sample 1 and sample 2, respectively. We define the DEG set by $\mathcal{G} = \{g: \mu_{g1} \neq \mu_{g2}, g = 1, \dots, G\}$ and its set-theoretic complement of non-differentially expressed genes, or "null gene set" by $\mathcal{G}' = \{1, \dots, G\} \backslash \mathcal{G}$. In presence of replicates, each hypothesis can be tested with a two-sample comparison, using Welch's t-test statistic: $t_g = (\bar{Y}_{g1} - \bar{Y}_{g2})/[(S_{g1}^2/N_{g1} + S_{g2}^2/N_{g2})^{1/2}]$ were $\bar{Y}_{g1}$ and $\bar{Y}_{g2}$, S_{g1} and S_{g2}, and N_{g1} and N_{g2} are each groups' respective sample mean, standard deviation, and size. However, when there is only one observation for Y_{g1} and one for Y_{g2}, neither S_{g1}^2 nor S_{g2}^2 are computable. We thus propose iDEG: an algorithm that transforms Y_{g1} and Y_{g2}, such that a simple function of the transformation allows for modeling all genes with the same distribution. This is done by pooling the genes together and estimating their common variance, hence bypassing the single-subject, single-replicate limitation.

Normalize read counts with unequal library sizes (Fig. 1, Panel 1)

In practice, unequal DNA library sizes may exist; thus, the first step is to normalize library sizes if necessary. We use the quantile-adjusted conditional maximum likelihood (**cXML**) procedure by Robinson and Smyth[16], for normalization, and subsequently, iDEG is applied.

Partition genes into windows to estimate local mean and variance (Fig. 1, Panel 2)

Marioni et al demonstrated the aptness of using expression means to estimate a gene's variance[17]. Therefore, by extension, in iDEG we assume that genes of comparable expression levels are assumed to behave similarly (genes with similar means share similar variances). Thus, after normalization, the next step is to group genes into W non-overlapping windows of similar expression levels to approximate each window's local mean and variance parameters. In the re-parameterized NB distribution, the variance of a given gene, g, is a function of its mean, μg, and dispersion, δg. Thus, the genes are partitioned to obtain their local, window-specific parameters. We define the w^{th} window by the $(w-1)^{th}$ and $(w)^{th}$ percentiles, for $w=1,\dots,W$. $\mathcal{G}_w = \{g: (w-1)^{th} \text{ percentile of } Y_{g1} < Y_{g1} < w^{th} \text{ percentile of } Y_{g1}\}$. To provide robust parameter estimates, we recommend a large positive integer, W, so that each window contains between 150 and 200 genes. However, the final predictions are not overtly sensitive to the choice of W (< 10% difference, data not shown – available upon request).

Compute each window's parameters (Fig. 1, Panel 3)

As seen in (**Eq. 1**), $Var(Y_{gd}) = \mu_{gd} + \delta_g \mu_{gd}^2$. Therefore, when estimating variance locally, we are actually estimating the dispersion parameter δ_g for each gene count. This value is required for the variance-stabilizing transformation (**VST**) h calculation (**Eq. 3; Fig. 1 Panel 4**). However, variance cannot be estimated when only a single observation is available. For RNA-Seq data analysis, one common assumption is that the dispersion δ_g is equal across samples 1 and 2, and that dispersion is a function q of the mean, μ_g[18-20]. Thus, in the absence of replicates, we partition genes into small windows to estimate **the functional mean-dispersion relationship, $\delta_g = q(\mu_g)$**, and hence the variance. We propose a two-step nonparametric procedure to obtain: (I) an initial estimate of δ_g by pooling genes locally; and (ii) a refined estimate of δ_g by estimating $q(\mu_g)$ with a smooth curve-fitting technique. In this approach, all non-differentially expressed (null) genes belonging to the same window $\mathcal{G}_w$ roughly have the same mean μ_w^* and the dispersion value δ_w^*. Thus,

$Y_{\text{gd}} \sim NB\left(\mu_w^*, \delta_w^*\right), \;\; \forall g \in \mathcal{G}_w, \;\; d = 1,2$ where μ_w^* and δ_w^* are the window-specific mean and dispersion values for null genes in $\mathcal{G}_w$, while d specifies if the count comes from sample 1 or 2.

The initial window estimates μ_w^* and σ_w^{2*} as $\hat{\mu}_w^* = Median\left(Y_{g1}\right) \; \forall g \in \mathcal{G}_w$ and $\hat{\sigma}_w^* = Median\left(\left|Y_{g1} - Median(Y_{g1})\right|\right) \; \forall g \in \mathcal{G}_w$, respectively. Since $Var(Y_{gd}) = \sigma_{gd}^2 = \mu_{gd} + \delta_g \mu_{gd}^2$, we estimate δ_w^* with $\hat{\delta}_w^* = (\sigma_w^{2*} - \hat{\mu}_w^*)/\hat{\mu}_w^{2*}$, for all windows $w = 1, \cdots, W$. To further improve the estimate of the dispersion parameter, δ_g, a smoothing spline technique is used to fit a functional mean-dispersion relationship, $\delta_g = q(\mu_g)$, by solving the following optimization problem:

$$\hat{q}_\lambda = \min_{q \in Q} \sum_{w=1}^{W}\{\hat{\delta}_w^* - q\left(\hat{\mu}^*{}_w\right)\}^2 + \lambda \int [q''(t)]^2 \, dt \qquad \textbf{(Eq. 2)}$$

where Q is the second-order Sobolev space on [0, 1] containing q, and λ is a smoothing parameter (selected via generalized cross validation)[21]. After the fitted curve $\hat{q}_\lambda$ is obtained as in **Eq. 2**, the refined estimate of $\hat{\delta}_g$ is computed as $\hat{\delta}_g = \hat{q}_\lambda(\hat{\mu}_w^*), \;\; \forall g \in \mathcal{G}_w, \;\; w = 1, \cdots, W$.

Apply the Variance Stabilizing Transformation ($h(Y_{gd})$) to each gene (Fig.1, Panel 4)

Fig. 2. Variance Stabilizing Transformation (VST). Panel **A** depicts the raw difference $D_g = Y_{g1} - Y_{g2}$ for 20,000 simulated genes (**Methods Simulations**), suggesting that the variance of D_g increases as the mean $\boldsymbol{\mu_{g1}}$ increases; hence, there is no uniform cutoff to differentiate DEGs and null genes. Panel **B** illustrates that, for null genes, VST makes the variance of $D_g^* = h(Y_{g1}) - h(Y_{g2})$ constant regardless of mean $\boldsymbol{\mu_{g1}}$.

After fitting δ_g, we apply the variance stabilizing transformation h to the counts, $Y_{gd} \; \forall g \in \mathcal{G}_w$:

$$h\left(Y_{gd}\right) = \frac{1}{\sqrt{\delta_g}}\sinh^{-1}\sqrt{Y_{gd}\delta_g} + \left(\sqrt{\frac{1}{\delta_g} - 1}\right)\sinh^{-1}\sqrt{\frac{Y_{gd} + \frac{3}{4}}{\frac{1}{\delta_g} - \frac{3}{2}}} \,, \forall g \in \mathcal{G}_w \;\; \textbf{(Eq. 3)}$$

This transformation[22] of the expression Y_{gd} in each window w, results in an approximately constant variance across all windows of the transcriptome (**Fig. 2**), regardless of the expression mean, μ_{gd}. That is $h(Y_{gd}) \stackrel{.}{\sim} N(h(\mu_{gd}), 1)$, where d specifies if the count comes from sample 1 or 2. Therefore, the difference of the two independent normal random variables (e.g., $h\left(Y_{g1}\right) - h(Y_{g2})$) approximately follows a common normal distribution with mean 0 and a constant variance: $Z_g = (1/\sqrt{2})\left[h\left(Y_{g1}\right) - h\left(Y_{g2}\right)\right] \sim N(0,1)$. We suggest replacing $\sqrt{2}$ by a robust estimate of standard deviation (e.g., median absolute deviation)[23]. In most single-subject analyses, the estimated dispersion parameter, $\hat{\delta}_g$, is small, but when $\hat{\delta}_g \geq 2/3$, the VST $h(Y_{gd})$ is not numerically stable. To avoid this issue, we suggest replacing h with h^* [24], where $h^*\left(Y_{gd}\right) = (1/\sqrt{\delta_g})\sinh^{-1}\sqrt{Y_{gd}\delta_g}, \;\; g = 1, \cdots, G; \;\; d = 1,2$. If a negative value of $\hat{\delta}_0$ is obtained, we conservatively set it to zero to assume a larger variance.

Compute the summary statistic for each gene (Fig.1, Panel 5)

In the context of noisy data and large-scale inference, performing individual tests neglects the parallel structure of RNA-Seq data. Moreover, actual data mean and variance may not be close to their theoretical values of 0 and 1 due to various reasons (e.g., correlation across genes, correlation between samples, or failed mathematical assumptions)(25). Therefore, we estimate an empirical null distribution $N(\mu_0, \sigma_0)$ to test these individual hypotheses.

Since differentially and non-differentially expressed genes generally follow different distributions, the probability density function of $Z_g, f(z)$, is naturally modeled by a two-group mixture: $f(z) = \pi_0 f_0(z) + \pi_1 f_1(z)$. Here, f_0 and f_1 are the probability density functions of genes in $\mathcal{G}'$ and in $\mathcal{G}$, while π_0 and $\pi_1 = 1 - \pi_0$ are their respective membership proportions. We assume a normal distribution following previous work from Dean and Raftery(26) that applied a two-group mixture model to identify differentially expressed genes, assuming a normal distribution for the null genes and a uniform distribution for the DEGs. However, we relax their assumptions for the marginal distribution and assume an exponential family. We approximate $f(z)$ using a smooth K-parameter exponential family distribution, $f(z) = \exp\left\{\sum_{k=0}^{K} \beta_k z^k\right\}$, and estimate the parameters $(\beta_0, \beta_1, \cdots, \beta_K)^T$ using Efron's approach(27).

Estimate the local false discovery (locfdr) for each and identify DEGs (Fig. 1, Panel 6)

Finally, to control the false discovery rate (**fdr**), we adopt Efron's idea(28-30) to estimate the local fdr (***locfdr***) using the R package *locfdr* and estimate π_0, $f_0(z)$ by maximum likelihood. Efron et al.(31) have shown *locfdr*'s close connection to the BH false discovery rate procedure(32); therefore, after estimating *locfdr(zg)*, it identifies differentially expressed genes by comparing *locfdr(z_g)* to a pre-specified **α**-cutoff value. The finally set of differentially expressed genes identified by applying the iDEG procedure is denoted by $\hat{\mathcal{G}}$.

Simulations (Fig. 2-3)

To compare the performance of iDEG to existing methods – including edgeR(16), DEGSeq(8), DESeq(19), and NOISeq(6) – extensive numerical studies were conducted assuming that RNA-Seq data follow the NB distribution with a varying dispersion parameter δ_g. Of note, these methods assume the NB distribution for data, which is used in the simulation; except for NOISeq that is nonparametric and DEGseq which assumes a binomial distribution. Baseline (normal tissue) and case (tumor sample) transcriptomes are both simulated and assumed to contain G = 20,000 genes; the library size of one transcriptome is 1.5 times larger than the other one. The single-subject RNA-Seq datasets are simulated with different percentages of DEGs, including DEG percentage = 5%, 10%, 15%, 20%, 25%, 30%, 35%, and 40%. Each experiment is repeated 1000 times. $Y_{g1} \sim \text{NB}(\mu_{g1}, \delta_g)$ and $Y_{g2} \sim \text{NB}(\mu_{g2}, \delta_g)$, where the dispersion parameter δ_g has been set to $\delta_g = 0.005 + 9/(\mu_{g1} + 100)$, per Anders and Huber(19). Gene expression means, μ_{g1}, are sampled from: $P(\mu_{g1}) = (1/500)e^{-(1/500)\times \mu_{g1}}$, $g = 1, \cdots, 20000$. For the case transcriptome, we set $\mu_{g2} = \mu_{g1}$ for $g \in \mathcal{G}'$ and $\mu_{g2} = d^s \mu_{g1}$ for $g \in \mathcal{G}$, where $s = (-1)^b$ and b ~ Bernoulli(0.5) is a random variable. Here, s indicates increasing expression (s = 1) or decreasing expression (s = −1) of a gene in the case transcriptome relative to baseline. Finally, for each gene g, we simulate one observation for Y_{g1} and Y_{g2} respectively and test the hypothesis $\mu_{g1} = \mu_{g2}$. At each iteration, a baseline and a case transcriptome are generated to simulate a distinct RNA-Seq dataset. Methods are assessed by

their Precision, Recall, and FPR, and F_1 score, $F_1 = (2 \cdot (\text{precision} * \text{recall}) / (\text{precision} + \text{recall})$. The average number of identified DEGs is also reported.

Of note, we excluded from the comparison GFOLD(7), a standard approach that can be applied to TCWR studies, as it only ranks genes without providing a measure of significance, thus prohibiting the accurate comparison with the remaining techniques using precision-recall curves or ROC curves. In clinical practice, selecting a rank cutoff for GFOLD would be arbitrary and less useful than a method providing significance as selecting the top 10% of genes arbitrarily assumes there are in fact ~10% of DEGs in the transcriptome.

Results

Table 1 and **Figure 3** depict the accuracies obtained in the simulations. Of note, we have also conducted analyses with a Poisson distribution and showed similar ordering of accuracies between the evaluated methods (data not shown).

Table 1. Performance results of TCWR simulations. At different percentage of DEGs in TCWR simulations, distinct methods obtain the best precision and recall, with iDEG, NOISeq and edgeR producing the best combinations of precision and recall. Of note, edgeR, DEGseq, and DESeq were not designed nor validated for studies without replicates; however, their authors proposed to utilize them in these conditions by defining specific parameters. NOISeq-sim, the NOISeq solution for studies without replicates, obtained the best F1 score (**Fig. 3**), and offers high recall and precision with DEGs=40% i.e. when 8,000 genes are dysregulated among 20,000. On the other hand, iDEG obtains high precision with moderate to high recall in all conditions. EdgeR provides moderate precision with very high recall for DEGs>20%.

Proportions of DEGs seeded	Method	Precision	Recall (TPR)	FP	Predicted DEGs
1,000 out of 20,000 (5% of genes)	**iDEG**	**0.93**	0.65	57	*700*
	edgeR	0.31	0.96	2,090	*3,119*
	NOISeq-sim	0.412	0.89	1140	*2,163*
	DESeq	1.0	0.16	0	*162*
	DEGseq	0.086	**0.99**	10,450	*11,397*
4,000 out of 20,000 (20% of genes)	**iDEG**	**0.97**	0.76	112	*3,136*
	edgeR	0.60	0.97	2,560	*6,405*
	NOISeq-sim	0.747	0.91	1120	*4,897*
	DESeq	Not applicable	0	0	*0*
	DEGseq	0.3	**0.99**	9,280	*13,160*
8,000 out of 20,000 (40% of genes)	**iDEG**	**0.98**	0.80	120	*6,484*
	edgeR	0.77	0.97	2,400	*10,145*
	NOISeq-sim	0.893	0.91	840	*8,177*
	DESeq	Not applicable	0	0	*0*
	DEGseq	0.51	**0.99**	7,560	*15,455*

Fig. 3. Performance results. NOISeq-sim's and iDEG's F_1-scores are more accurate than that of other methods at 5%<DEGs<20% (iDEG) and 30%<DEGs<40% (NOISeq-sim). At DEG=5% and FDR<10%, iDEG provides an interesting compromise between precision and recall, while NOISeq provides a better compromise when the percentage of DEGs is higher than 30%. DESeq offers high precision and low recall while DEGSeq shows the opposite. EdgeR also operates at high recall and precision when DEG<45%. **Panel A-B**. Precision recall curve at 10% FDR for 1,000 and 8,000 seeded DEGs among 20,000 transcripts, respectively. **Panel C.** F_1 scores. Average F_1 scores resulting from 1,000 repeated experiments with vertical bars representing one standard deviation.

Discussion

A summary of the range of optimal operation of iDEG and the other methods are provided in **Table 2**. No single method has emerged as the optimal approach for all conditions. Low expression levels are extremely susceptible to unstable fold-change estimation, as a 5-fold increase from 2 to 10 counts on a dynamic range of 0 to 100,000 should not be treated equivalently to that between 10,000 and 50,000. Standard practice filters out genes with counts below a certain threshold (typically 5 or 10). However, this solution does not address fold change (FC) inflation above the threshold (e.g., FC>2 at 15 counts), nor how to compare distinct FCs at different expression levels. Alternatively, favoring absolute count difference to identify DEGs leads to a systemic bias towards genes with high expression. Conversely, favoring FC results in a systemic bias towards lowly expressed genes. Either of these solutions yields higher false positive rates. For DEGs<30%, the variance stabilization within partitioned windows proposed in iDEG appears to address this dilemma of dealing with fold change inflation by comparing FC values relative to their expression levels, while conventional approaches impose stringent data assumptions that may compromise downstream inferential processes. These restrictions occur either by methods always assuming rather than estimating a distribution and its parameters or by treating case and control observations as replicates to compute the variance in their absence.

Table 2. Operating Characteristics. NOISeq is the only method operating at high precision and recall, but only at DEGs>20%.

Operating Characteristics	Method	5% DEG	10% DEG	20% DEG	40% DEG
Moderate or better Recall (>50%)	iDEG	✔	✔	✔	✔
	edgeR	✔	✔	✔	✔
	NOISeq-sim	✔	✔	✔	✔
	DESeq	✘	✘	✘	✘
	DEGseq	✔	✔	✔	✔
Moderate or better Precision (>50%)	iDEG	✔	✔	✔	✔
	edgeR	✘	✔	✔	✔
	NOISeq-sim	✘	✔	✔	✔
	DESeq	✘	✘	✘	✘
	DEGseq	✘	✘	✘	✔
High Recall (>90%)	iDEG	✘	✘	✘	✘
	edgeR	✔	✔	✔	✔
	NOISeq-sim	✔	✔	✔	✔
	DESeq	✘	✘	✘	✘
	DEGseq	✔	✔	✔	✔
High Precision (>90%)	iDEG	✔	✔	✔	✔
	edgeR	✘	✘	✘	✘
	NOISeq-sim	✘	✘	✘	✔
	DESeq	✔	✘	✘	✘
	DEGseq	✘	✘	✘	✘

Although existing methods can operate in the single-subject, single-sample setting, **Table 1** and **Fig. 3** demonstrate the quality of inference of iDEG as compared to conventional methods. We note that the comparison is not fair, since these techniques were not explicitly designed for single-subject, single-sample studies. We simply conducted the simulations against these methods to demonstrate the need for new approaches to study single-subject transcripts in TCWR conditions. In addition, iDEG relies on widely-accepted parametric assumptions of RNA-Seq data; however, more sensitivity studies should be conducted to evaluate the performance when these assumptions are violated. In addition, a true gold standard to evaluate iDEG and other methods is not as simple as obtaining replicates and running conventional methods as pointed out by recent papers(33, 34). We have also witnessed this difficulty in our previous work when investigating real biological datasets and observed that the definition of a correct gold standard was not possible as the overlap of DEGs between distinct methods was as low as 40% in spite of 30 replicates in isogenic conditions(35). We have thus generated multiple distinct reference standards and developed a comprehensive evaluation of methods to identify DEGs in paired conditions using *biological datasets*, and the results are presented as an independent paper due to its comprehensiveness(35).

Given that the purpose of the companion study served to better understand how DE methods perform in clinical studies and for properly building reference standards in TCWR, it would detract from both studies to combine them into a single, larger study. The companion paper(35) complements this simulation study by providing a clinical and biological validation for iDEG, and furthermore, it demonstrates the strengths and limitations of DE detection techniques in TCWR conditions. The study concludes with our recommendation for an ensemble-type approach to mitigate over-relying on any one technique in absence of a gold standard.

Conclusion

Over the past decade, state-of-the-art techniques in RNA-Seq data analysis have delivered powerful new tools(14) for extending large-scale inference to small-sample settings. The primary goal of iDEG is not to replace these, but rather to expand the scope of RNA-Seq studies into the single-subject, single-replicate realm and provide novel research opportunities and test methods for controlling fold change inflation at low expression ranges. In iDEG, we have shown window portioning to be a novel way of borrowing information across genes locally, with early promising results. Furthermore, this novel research technique for local differential gene expression detection can easily be extended to parametric and non-parametric differential expression tools as a way to enrich and refine existing techniques. In future studies, we envision to i) extend the window partitioning component of iDEG into other techniques, ii) expand them from single-subject analysis to cohort analysis, iii) to locally identify differentially expressed pathways (by incorporating ontologies and knowledge graphs), iv) to apply it to other 'omics measures, and v) to further precision-medicine discoveries in single-subject, single-replicate studies.

Acknowledgements. This work was supported in part by The University of Arizona Health Sciences CB2, the BIO5 Institute, NIH (U01AI122275, CA023074, 1UG3OD023171, 1S10RR029030)

References

1.Kaiser J. Obama Gives East Room Rollout To Precision Medicine Initiative. **Science.** 2015.
2.Schork NJ. Personalized Medicine: Time for One-Person Trials. **Nature**. 2015;520(7549):609-11.

3. Topol EJ. Individualized Medicine From Prewomb To Tomb. **Cell.** 2014;157(1):241-53.

4. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor Package for Differential Expression Analysis of Digital Gene Expression Data. **Bioinformatics**. 2009;26(1):139-40.

5. Anders S, Huber W. Differential expression of RNA-Seq data at the gene level–the DESeq package. Heidelberg, Germany: European Molecular Biology Laboratory (**EMBL**). 2012.

6. Tarazona S, García F, Ferrer A, Dopazo J, Conesa A. NOIseq: a RNA-seq differential expression method robust for sequencing depth biases. **EMBnet journal.** 2011;17(B):18-9.

7. Feng J, Meyer CA, Wang Q, Liu JS, Shirley Liu X, Zhang Y. GFOLD: a generalized fold change for ranking differentially expressed genes from RNA-seq data. **Bioinformatics.** 2012;28(21):2782-8.

8. Wang L, Feng Z, Wang X, Wang X, Zhang X. Degseq: an R Package for Identifying Differentially Expressed Genes From Rna-Seq Data. **Bioinformatics.** 2009;26(1):136-8.

9. Schissler AG, Gardeux V, Li Q, Achour I, Li H, Piegorsch WW, et al. Dynamic changes of RNA-sequencing expression for precision medicine: N-of-1-pathways Mahalanobis distance within pathways of single subjects predicts breast cancer survival. **Bioinformatics**. 2015;31(12):i293-i302.

10. Li Q, Schissler AG, Gardeux V, Achour I, Kenost C, Berghout J, et al. N-of-1-pathways MixEnrich: advancing precision medicine via single-subject analysis in discovering dynamic changes of transcriptomes. **BMC Medical Genomics**. 2017;10(1):27.

11. Li Q, Schissler AG, Gardeux V, Berghout J, Achour I, Kenost C, et al. kMEn: Analyzing noisy and bidirectional transcriptional pathway responses in single subjects. **Journal of biomedical informatics**. 2017;66:32-41.

12. Gardeux V, Achour I, Li J, Maienschein-Cline M, Li H, Pesce L, et al. 'N-of-1-pathways' unveils personal deregulated mechanisms from a single pair of RNA-Seq samples: towards precision medicine. **Journal of the American Medical Informatics Association**. 2014;21(6):1015-25.

13. Zaim SR, Li Q, Schissler AG, Lussier YA. Emergence of pathway-level composite biomarkers from converging gene set signals of heterogeneous transcriptomic responses . **Pac Symp Biocomput.** 2018;23:484-495.

14. Vitali F, Li Q, Schissler AG, Berghout J, Kenost C, Lussier YA. Developing a 'personalome' for precision medicine: emerging methods that compute interpretable effect sizes from single-subject transcriptomes. **Briefings in bioinformatics.** 2017.

15. Team RC. R: A language and environment for statistical computing. 2013.

16. Robinson MD, Smyth GK. Small-Sample Estimation of Negative Binomial Dispersion, With Applications To SAGE Data. **Biostatistics.** 2007;9(2):321-32.

17. Marioni JC, Mason CE, Mane SM, Stephens M, Gilad Y. RNA-seq: an assessment of technical reproducibility and comparison with gene expression arrays. **Genome research.** 2008.

18. Law CW, Chen Y, Shi W, Smyth GK. Voom: Precision Weights Unlock Linear Model Analysis Tools for Rna-Seq Read Counts. **Genome Biology**. 2014;15(2):R29.

19. Anders S, Huber W. Differential expression analysis for sequence count data. **Genome biology.** 2010;11(10):R106.

20. Di Y, Schafer DW, Cumbie JS, Chang JH. The NBP negative binomial model for assessing differential gene expression from RNA-Seq. **Statistical Applications in Genetics and Molecular Biology.** 2011;10(1).

21. Craven P, Wahba G. Smoothing Noisy Data With Spline Functions. **Numerische Mathematik.** 1978;31(4):377-403.

22. Laubscher NF. On Stabilizing the Binomial and Negative Binomial Variances. **Journal of the American Statistical Association**. 1961;56(293):143-50.

23. Hampel FR. The influence curve and its role in robust estimation. **Journal of the American Statistical Association.** 1974;69(346):383-93.

24. Montgomery DC. Design and analysis of experiments: John Wiley & Sons; 2008.

25. Efron B. Large-scale inference: empirical Bayes methods for estimation, testing, and prediction: Cambridge University Press; 2012.

26. Dean N, Raftery AE. Normal uniform mixture differential gene expression detection for cDNA microarrays. **BMC bioinformatics**. 2005;6(1):173.

27. Efron B, Tibshirani R. Using Specially Designed Exponential Families for Density Estimation. **The Annals of Statistics.** 1996;24(6):2431-61.

28. Efron B. Large-Scale Simultaneous Hypothesis Testing. **Journal of the American Statistical Association.** 2004;99(465):96-104.

29. Efron B. Size, Power and False Discovery Rates. **The Annals of Statistics.** 2007;35(4):1351-77.

30. Efron B. Correlation and Large-Scale Simultaneous Significance Testing. **Journal of the American Statistical Association**. 2007;102(477):93-103.

31. Efron B, Tibshirani R, Storey JD, Tusher V. Empirical Bayes Analysis of a Microarray Experiment. **Journal of the American Statistical Association**. 2001;96(456):1151-60.

32. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. **Journal of the royal statistical society Series B** (Methodological). 1995:289-300.

33. Costa-Silva J, Domingues D, Lopes FM. RNA-Seq differential expression analysis: An extended review and a software tool. **PloS one**. 2017;12(12):e0190152.

34. Soneson C, Robinson MD. Bias, robustness and scalability in single-cell differential expression analysis. **Nature methods.** 2018;15(4):255.

35. Rachid Zaim, S.;Kenost, C.; Berghout, J.; Zhang, H.H.; Lussier, Y.A. Evaluating single-subject study methods for personal transcriptomic interpretations to advance precision medicine **BioArxiv.** Deposited Sept. 25th 2018, under consideration at the Transaltional Bioinformatics Conference 2019 for proceedings in BMC Medical Genomics.