Molecular Characterisation of the Synovial Fluid Microbiome in Rheumatoid Arthritis Patients and Healthy Control Subjects

Dargham Bayan Mohsen Hammad[1], Veranja Liyanapathirana[2], and Daniel Paul Tonge[1]

[1]School of Life Sciences, Faculty of Natural Sciences, Keele University, ST5 5BG, UK

[2]Department of Microbiology, Faculty of Medicine, University of Peradeniya, Sri Lanka

Corresponding Author:

Dr Daniel Tonge

School of Life Sciences

Faculty of Natural Sciences

Keele University

ST5 5BG

United Kingdom

Author Names

Dr Daniel P Tonge School of Life Sciences, Faculty of Natural Sciences, Keele

University, ST5 5BG, d.p.tonge@keele.ac.uk 01782 733418

Mr Dargham Bayan Mohsen Hammad School of Life Sciences, Faculty of Natural

Sciences, Keele University, ST5 5BG, d.b.m.hammad@keele.ac.uk 01782 733418

Dr Veranja Liyanapathirana Department of Microbiology, Faculty of Medicine,

University of Peradeniya, Sri Lanka veranjacl@yahoo.com

Keywords

Microbiome, synovial fluid, bacteria, fungi, ITS, 16S rRNA

Abstract

The colonisation of specific body sites in contact with the external environment by microorganisms is both well-described and universally accepted, whereas, the existence of microbial evidence in other "classically sterile" locations including the blood, synovial space, and lungs, is a relatively new concept. Increasingly, a role for the microbiome in disease is being considered, and it is therefore necessary to increase our understanding of these. To date, little data support the existence of a "synovial fluid microbiome".

Methods. The presence and identity of bacterial and fungal DNA in the synovial fluid of rheumatoid arthritis (RA) patients and healthy control subjects was investigated through amplification and sequencing of the bacterial 16S rRNA gene and fungal internal transcribed spacer region 2 respectively. Synovial fluid concentrations of the cytokines IL-6, IL-17A, IL22 and IL-23 were determined by ELISA.

Results. Bacterial 16S rRNA genes were detected in 87.5% RA patients, and all healthy control subjects. At the phylum level, the microbiome was predominated by *Proteobacteria* (Control = 83.5%, RA = 79.3%) and *Firmicutes* (Control = 16.1%, RA = 20.3%), and to a much lesser extent, *Actinobacteria* (Control = 0.2%, RA = 0.3%) and *Bacteroidetes* (Control = 0.1%, RA = 0.1%). Fungal DNA was identified in 75% RA samples, and 88.8% healthy controls. At the phylum level, synovial fluid was

predominated by members of the Basidiomycota (Control = 53.9%, RA = 46.9%) and

Ascomycota (Control = 35.1%, RA = 50.8%) phyla. Statistical analysis revealed key

taxa that were differentially present or abundant dependent on disease status.

Conclusions. This study reports the presence of a synovial fluid microbiome, and

determines that this is modulated by disease status (RA) as are other classical

microbiome niches.

Introduction

The Microbiome

The term "microbiome" describes the entire habitat, including the microorganisms (bacteria, archaea, lower and higher eukaryotes, and viruses), their genomes, and the surrounding environmental conditions. Herein, we utilise this term to denote the existence of a microbial community evidenced through the survey of 16S rRNA genes. In contrast, the term "microbiota" refers to the assemblage of viable microorganisms that comprise these communities [1]. Whilst previous estimates stated that the number of microbial cells present in our microbiota exceeded our own by approximately one order of magnitude, current estimates are more conservative and suggest that the number of bacterial cells in the human body is roughly equal to the number of human cells, representing a mass of approximately 0.2 kg [2]. In either case, the microbiota represents a significant source of non-host biological material. The microbiota undertakes essential biological processes and thus it is unsurprising that a number of disease states are associated with changes in microbiome composition, termed "dysbiosis". Whilst the colonisation of specific body sites in contact with the external environment (such as the gastrointestinal tract, skin and vagina) by microorganisms is both well-described and universally accepted [3], the existence of microorganisms and or microbial DNA in other "classically sterile" locations including the blood [4], synovial space [5], and lungs [6-9], is a relatively new concept.

Rheumatoid Arthritis and Dysbiosis

Rheumatoid arthritis (RA) is a chronic autoimmune disease that causes synovial joint inflammation leading to significant joint pain, swelling, and significant disability with increased morbidity and mortality. RA is estimated to affect 0.5 - 1% of the population worldwide [10]. RA aetiology still remains unclear. Although the autoimmune hypothesis is well established the causes generating this self-directed immune reaction are still poorly understood. Although convention still suggests otherwise, an overwhelming volume of epidemiological and experimental data indicates a microbial origin for RA (reviewed extensively by Pretorius and colleagues [11]) and therefore increasingly, a role for the altered microbiome composition in disease initiation and progression is considered. To date, RA has been associated with dysbiosis of the oral [12, 13], gastrointestinal [12, 14-22], and respiratory [23-25] microbiomes and associated with the presence of specific organisms (most commonly *P. mirabilis*) within the urinary tract [26-29], suggesting that RA results in (or from) pan-microbiome dysbiosis. Studies have further begun to associate RA, or stages thereof, with the presence or absence of specific bacteria, including the expansion of rare bacterial linages [16]. Such findings have the potential to increase our disease understanding, and further suggest that the microbiome may afford a valuable source of novel biomarkers and or novel targets for therapeutic modulation.

The classical assertion that RA lacks a microbial component stems from the generalised inability of clinicians and scientists to observe microbial colonies on synthetic media

when samples from RA patients are subjected to routine culture. Such assessments

crudely characterise organisms as "*alive*" or "*dead*" (or perhaps more accurately,

present or not present) based upon this colony forming ability, yet do not routinely

recognise the potential for the presence of viable-non culturable bacteria. Furthermore,

extensive evidence now provides for the existence of bacterial DNA and RNA in

classically sterile areas leaving the possibility of; (a) microbial translocation from

classical niches (e.g. the gastrointestinal tract) followed by opsonisation leaving only the

nucleic acid, (b) the passage of cell-free microbial nucleic acid through the circulatory

system, (c) microbial translocation followed by a state of dormancy due to unfavourable

environmental conditions.   With relevance to the first two scenarios, it is well-

established that the immune system is able to differentiate self from non-self-nucleic

acid (DNA and RNA) through specific pattern recognition receptions [30], and that

foreign DNA is recognised by TLR9, and RNA is recognised by TLR3 on immune cells,

resulting in the   upregulation of type I interferons and various pro-inflammatory

cytokines [31] (also associated with rheumatoid arthritis) [32]. TLR9 is found in the

endosome of peripheral blood mononuclear cells, such as monocytes, macrophages, T,

B, and NK cells, whereas TLR3 is expressed only in the endosomes of myeloid derived

cells such as dendritic cells and monocytes [33]. With relevance to the latter scenario,

dormant or VBNC bacterial cells may persist within the joints of arthritic patients,

undetected by routine culture, whilst retaining the ability to shed inflammatory agents

such as lipopolysaccharide (LPS) and other antigenic components [11], yet such

samples may present as culture-negative. Indeed, bacterial LPS is able to simulate

many of the common RA-associated cytokines [34].

Evidence of Bacteria in the Synovial Fluid: Origin and Transport

Under normal physiological conditions, one expects the synovial space to be sterile.

Indeed, the presence of viable bacteria results in a diagnosis of septic arthritis, a

condition regarded as a true medical emergency [35]. Nevertheless, numerous studies

report the presence of microorganisms, or microorganism-derived genetic material in

the synovial fluid of RA patients and healthy control subjects [36-42]. Interestingly, these

studies overwhelmingly identify bacteria most commonly found to reside within the oral

cavity, and these findings are accompanied by evidence demonstrating the presence of

these DNAs and or antibodies directed against the originating organism in the blood [36,

37, 43-46]. On the basis of these findings, we suggest that bacteria / bacterial nucleic

acid may reach the synovial space via the blood (as evidenced by their concurrent

presence in both fluids), and further, that the inflammatory environment of the synovial

tissues may facilitate the trapping of these bacterial DNAs, increasing their apparent

concentration in this location [37].

Despite extensive epidemiological and experimental evidence supporting a microbial

cause in RA, mounting evidence linking infection of the oral tissues with RA [23, 47-58],

and data demonstrating the appearance of microbial DNA from distant microbiome

niches in the synovial space (see above), there has been limited study of the bacterial

community in this compartment [5] and no global characterisation of the fungal DNA

present in this location to date. To this end, we investigated the presence of bacterial

(via sequencing of the 16S rRNA gene) and fungal (via sequencing of the ITS2 region)

DNA, in the synovial fluid of human RA patients and healthy control subjects using

cutting-edge next generation sequencing and bioinformatic techniques. Furthermore,

we evaluated any apparent changes in the bacterial or fungal communities detected, in

the context of key markers of synovial inflammation (IL-6, IL-17A, IL-22 and IL-23).

Materials and Methods

Donor Population

This study investigated the presence of bacterial and fungal DNA in synovial fluid samples from sixteen rheumatoid arthritis patients, and nine sex and BMI-matched healthy control (disease-free) individuals who donated a sample of their synovial fluid for research purposes. Synovial fluid samples were aspirated from the knee joint using aseptic technique, transferred to a sterile micro-centrifuge tube, and stored at -80^{o}C prior to further analysis. Samples were procured from Sera Laboratories Limited.

The Independent Investigational Review Board Inc. (BRI-0722) ethically approved sample collection by Sera Laboratories Limited from human donors giving informed written consent. Furthermore, the authors obtained ethical approval from Keele University Ethical Review Panel 3 for the study reported herein. All experiments were performed in accordance with relevant guidelines and regulations.

Measuring synovial fluid cytokines

Markers of inflammation were measured in synovial fluid utilising the Human Magnetic Luminex Screening Assay following the manufacturer's instructions (R&D Systems, Minneapolis, USA). Levels of IL-6, IL-17A, IL-22, and IL-23 were analysed by Luminex kit LXSAHM-04. Prior to analysis, synovial fluid samples were centrifuged at 16,000 x g

for four minutes and diluted in a 1:2 ratio by adding 25 µl of sample to 25 µl of Assay

Buffer prior to analysis.

Microbiome Characterisation

Amplification and sequencing of the bacterial 16S rRNA variable region 4 and fungal

ITS region was used to characterise the bacterial and fungal DNA present in the

synovial fluid samples.

The bacterial 16S rRNA and fungal ITS2 genes were amplified by direct PCR utilising

the oligonucleotide primers detailed in (**Table 1)**. Such approaches lack the need for a

separate DNA extraction phase, often associated with the introduction of contaminants

that impact upon all downstream applications.   A first round of PCR using

oligonucleotide primers 16SV4 F/R and ITS2 F/R was carried out with four microliters of

each synovial fluid sample as template in a final volume of 20µl. Reactions comprised

10µl Phusion blood PCR buffer (Thermofisher), 0.4µl (2 U) Phusion blood DNA

polymerase, 1µl of each primer (10µM), and 3.6µl of nuclease-free water that had been

subject to 15 minutes UV-irradiation. A negative control reaction in which synovial fluid

was substituted with UV-irradiated nuclease-free water was run in parallel with our

experimental samples to confirm that none of the reagents were contaminated by target

DNA. Cycling parameters comprised an initial denaturation step performed at 98°C for 5

minutes, followed by 33 cycles of denaturation (98°C, 10 seconds), annealing (55°C, 5

seconds) and extension (72°C, 15 seconds). A final elongation of 7 minutes at 72°C

completed the reaction. Following gel electrophoresis of 5µl of each resulting PCR

product to confirm successful amplification, the remainder (15µl) was purified of excess

primer and PCR reagents using the QIAquick PCR Purification kit.

In order to add sequencing adapters to facilitate MiSeq library preparation, a second

round of PCR amplification was conducted in a total volume of 50µl comprising 10µl 5X

Platinum Super-Fi Buffer, 1µl 10mM dNTP mixture, 0.5µl Platinum Super-Fi

polymerase, 2.5µl of each XT_tagged primer (16SV4_XT F/R or ITS2_XT F/R), 5µL

from the successful first round PCR reaction and 38.5 µL of UV-irradiated nuclease-free

water. Cycling parameters comprised an initial denaturation step performed at 98°C for

2 minutes, followed by 7 cycles of denaturation (98°C, 10 seconds), and annealing /

extension (72°C, 20 seconds) extension (72°C, 15 seconds). A final elongation of 5

minutes at 72°C completed the reaction. Two controls were run in parallel with this

process and these comprised a standard PCR negative control (in which template was

replaced with UV irradiated molecular biology grade water) and the first round PCR

control which was progressed alongside our experimental samples. All PCR products

and our two control reactions were purified using AMPure XP magnetic beads

(Agencourt) at a ratio of 0.8 beads to sample (v/v), eluted in 20µl of UV-irradiated

nuclease-free water, and visualised by agarose gel electrophoresis and ehidium

bromide staining as above. All PCR negative reactions were subject to high sensitivity

DNA quantification using the Qubit 3.0 hsDNA kit (Invitrogen) to quantitate any DNA in

these samples.

Amplicons were barcoded using the Nextera DNA library kit, multiplexed for efficiency,

and sequenced using the Ilumina MiSeq system with a 250bp paired-end read metric.

Bioinformatic analysis was performed using QIIME implemented as part of the Nephele

16S / ITS paired-end QIIME pipeline using open reference clustering against the SILVA

database for bacteria and the ITS database for fungi at a sequence identity of 99%. All

other parameters remained as default. The statistical significance of differences in the

abundance of individual bacteria and fungi between the RA and control donors was

determined by applying a two-tailed, Mann Whitney test using GraphPad Prism V8.0.

Table1: Oligonucleotide primers used in this study

Primer Name	Primer Sequence (5' – 3')	Length
16SV4_F	GTGCCAGCMGCCGCGGTAA	19
16SV4_R	GGACTACHVGGGTWTCTAAT	20
16SV4_XT_F	TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGGTGCCAGCMGC CGCGGTAA	52
16SV4_XT_R	GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGGACTACHVG GGTWTCTAAT	54
ITS2_F	GCA TCG ATG AAG AAC GCAGC	20
ITS2_R	TCCTCCGCTTATTGATATGC	20
ITS2_XT_F	TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG GCATCGATGAAGAACGCAGC	53
ITS2_XT_R	GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG TCCTCCGCTTATTGATATGC	54

Results

Clinical features of donors and results of 16S rRNA and ITS2 PCR amplification

Synovial fluid samples were obtained from 25 human donors. Sixteen patients were diagnosed with RA, and of these, 8 were male and 8 were female. The RA patients' ages ranged from 52 to 74 years, with a mean of 65.3 years. Nine control synovial fluid samples were obtained from 5 males and 4 females. Their ages ranged from 50 to 68 years, with a mean of 61.0 years. There were no significant differences in the ages of the two cohorts (Unpaired T-test; $P = > 0.05$) **(Table 2)**.

Table 2: Characteristics of the patient population.

Patient ID#	Age	Diagnosis	16S rRNA PCR	ITS2 PCR
1339	61 - 65	RA	+	−
1340	66 - 70	RA	+	+
1341	66 - 70	RA	+	+
1342	66 - 70	RA	+	−
1343	66 - 70	RA	+	−
1344	66 - 70	RA	+	−
1345	56 - 60	RA	+	+

1346	50 - 55	RA	+	+
1347	50 - 55	RA	+	+
1348	66 - 70	RA	+	+
1349	71 - 75	RA	+	+
1350	66 - 70	RA	+	+
BRH1095336	66 - 70	RA	+	+
BRH1095337	66 - 70	RA	−	+
BRH1095338	66 - 70	RA	−	+
BRH1095339	66 - 70	RA	+	+
1351	56 - 60	Healthy	+	+
1352	61 - 65	Healthy	+	−
1353	61 - 65	Healthy	+	+
1354	50 - 55	Healthy	+	+
1355	50 - 55	Healthy	+	+
1356	71 - 75	Healthy	+	+
1357	61 - 65	Healthy	+	+
1358	50 - 55	Healthy	+	+
1359	66 - 70	Healthy	+	+

Utilising PCR amplification, bacterial 16S DNA was detected in the synovial fluid of 14

of out 16 patients with RA (87.5%), and in 9 out of 9 (100%) healthy control subjects.

ITS2 amplification, indicative of the presence of fungi, was detected in 12 of 16 (75%)

RA samples, and 8 out of 9 (89%) healthy control samples. Our various experimental

controls designed to identify target contamination amongst the molecular reagents and

plasticware used consistently failed to produce a visible band following PCR and

agarose gel electrophoresis. Furthermore, these samples were subjected to DNA

quantification using the Qubit 3.0 high-sensitivity DNA kit (Invitrogen) and were found to

be below the limit of detection in all cases.

Assessment of inflammatory markers in synovial fluid

Synovial fluid concentrations of interleukin 6 (IL-6), 17A (IL-17A), 22 (IL-22) and 23 (IL-

23) were measured using the Luminex system as described earlier. Mean interleukin

levels in healthy control and RA patient synovial fluid were significantly different, with

cytokines being present at higher levels in the RA synovial fluid in all cases (Unpaired

T- test) (**Table 3**).

Table 3: Differences in cytokine levels among

Cytokine	Control mean (SD) (pg/mL)	RA mean (SD) (pg/mL)	P value (pg/mL)
IL-6	156.43 (100.30)	1257.80 (905.66)	<0.005

IL-17A	11.68 (3.92)	65.44 (16.51)	<0.005
IL-22	220.14 (85.02)	873.00 (376.46)	<0.005
IL-23	145.64 (64.43)	296.21 (97.07)	<0.005

Characterisation of Bacterial populations via 16S rDNA sequencing of synovial fluid

The presence of bacterial DNA in synovial fluid was evaluated via PCR amplification and sequencing of the bacterial 16S rRNA gene, variable region 4. An average of 75,000 reads were generated for each of the samples; 70,658 reads in the RA samples, and 80,533 reads in the control samples. Although the control samples generated more reads on average, this difference was not statistically significant (Unpaired T test; P = > 0.05), and nevertheless, rarefaction was used prior to differential abundance analysis to control for differences in sequencing depth. Following taxonomic classification using the Nephele platform as described above, our first analytical approach utilised Principal Coordinates Analysis (PCoA) to reduce the complexity of the data obtained, and to visualise any obvious separation between the various experimental samples **(Figure 1).**

Figure 1 - Principal Coordinate Analysis plot generated from a Bray Curtis distance matrix of synovial fluid bacterial community structure for control subjects (red) and

arthritic patients (blue) as determined via amplification and sequencing of the 16S rRNA gene variable region 4 (V4). Proportions of variation explained by the principal coordinates are designated on the axes. PCoA found that the maximal variation were 62.2% (PC1), and 22.9% (PC2). The microbiome of samples that appear in close proximity to each other are more similar in composition. The data points in each pane are sized according to the synovial concentration of IL-17A (A), IL-22 (B), IL-23 (C), and IL-6 (D).

At the phylum level, the synovial fluid microbiome was predominated by *Proteobacteria*

(Control = 83.5%, RA = 79.3%) and *Firmicutes* (Control = 16.1%, RA = 20.3%), and to a

much lesser extent, *Actinobacteria* (Control = 0.2%, RA = 0.3%) and *Bacteroidetes*

(Control = 0.1%, RA = 0.1%). At the genus level, our synovial fluid samples were

predominated by the genus *Pseudomonas* (Control = 80.6%, RA = 66.1%, P = 0.30),

followed by the genus *Enterococcus* (Control = 11.7%, RA = 11.4%, P = 0.95). To a

lesser extent, synovial fluid samples contained members of the *Stenotrophomonas*

(Control = 1.5%, RA = 4.2%, P = 0.97), *Raoultella* (Control = <0.1%, RA = 6.5%, P =

0.051) and *Bacillus* (Control = 0%, RA = 1.2%, P = 0.99) genera, the latter two being

detected almost exclusively in the synovial fluid of our RA patients **(Figure 2)**. Finally,

unidentified taxa of the *Bacillales* order (Control = 4%, RA = 7.4%, P = 0.97) were also

present in relatively high abundance yet could not be further taxonomically classified.

Statistical analysis revealed that whilst the abundance of most bacterial DNA, at

different taxonomic levels, was unaffected by disease status, the abundance of

Raoultella was borderline significant (P = 0.051) and observed as the almost exclusive

presence of this organism's DNA in the synovial fluid of RA patients only (**Figure 3**).

Figure 2 - Relative abundance of genera representing >1% of the total bacterial

population in synovial fluid samples of rheumatoid arthritis (RA, n = 14) and control

(Control, n = 9) samples as determined by amplification and sequencing of the 16S

rRNA gene variable region 4. Data are mean abundance expressed as a percentage of

the total bacterial sequence count.

Figure 3 - Relative abundance of the genus *Raoultella*, identified in the synovial fluid of

healthy donors and RA patients by amplification and sequencing of the 16S rRNA gene

variable region. A borderline significant (p = 0.051) increase in the mean abundance of

Raoultella was identified in RA synovial fluid relative to healthy control synovial fluid.

Data are individual sample abundances expressed as a percentage of the total bacterial

sequence count.

Inflammatory markers and modulated by bacterial microbiome PCoA cluster

Based upon the clustering of samples observed following PCoA (**Figure 1**), we

investigated whether the samples in "cluster 1", had significantly different inflammatory

marker levels to the remaining samples, in addition to different microbial populations.

Interestingly, IL-6 was significantly higher in PCoA cluster 1 compared with the

remaining samples (Two Way ANOVA; P = 0.030) yet all other inflammatory markers

were unaffected by PCoA cluster (P > = 0.05).

Characterisation of fungal populations via ITS2 sequencing of synovial fluid

The presence of fungal DNA in synovial fluid was evaluated via PCR amplification and

sequencing of the fungal ITS2 gene. An average of 50,000 reads were generated for

each of the samples; 52,113 reads in the RA samples, and 49,929 reads in the Control

samples. Although the RA samples generated more reads on average, this difference

was not statistically significant ($P = > 0.05$). Following taxonomic classification using the

Nephele platform as described above, our first approach utilised Principal Coordinates

Analysis (PCoA) to reduce the complexity of the data obtained, and to visualise any

obvious separation between the two experimental groups **(Figure 4).**

Figure 4 - Principal Coordinate Analysis plot generated from a Bray Curtis distance

matrix of synovial fluid fungal community structure for control subjects (red) and arthritic

patients (blue) as determined by amplification and sequencing of the ITS2 gene.

Proportions of variation explained by the principal coordinates are designated on the

axes. PCoA found that the maximal variation were 30.6% (PC1), and 20.1%. The

microbiome of samples that appear in close proximity to each other are more similar in

composition. The data points in each pane are sized according to the synovial

concentration of IL-17A (A), IL-22 (B), IL-23 (C), and IL-6 (D).

At the phylum level, synovial fluid was found to be predominated by members of the

Basidiomycota (Control = 53.9%, RA = 46.9%) and Ascomycota (Control = 35.1%, RA =

50.8%) phyla. At the genus level, synovial fluid samples were predominated by the

genus *Malassezia*, which accounted for 45.3% and 40.9% of the total number of fungal

sequences identified in the Control and RA donors respectively ($P = 0.8$). To a lesser

extent, synovial fluid samples also contained organisms from the genera *Penicillium*

(Control = 0.7%, RA = 15%, P = 0.6), *Candida* (Control = 11.9%, RA = 0.00%, P = 0.4),

Pichia (Control = 8.9%, RA = 0.00%, P = 0.4), *Aspergillus* (Control = 0.2%, RA = 6.2%,

P = 0.14), Cladosporium (Control = 5.1%, RA = 0.1%, P = 0.019), and *Debaryomyces*

(Control = 4.7%, RA = 0.4%, P = 0.7). Furthermore, genus level classification revealed

the presence of specific unclassified organisms belonging to the classes

*Tremellomycete*s (Control = 3%, RA = 2.8%, P = 0.98), and *Agaricomycetes* (Control =

3.2%, RA = 2.2%, P = 0.3), and the orders *Hypocreales* (Control = 0.00%, RA = 26.4%,

P = 0.012) and Malasseziales (Control = 1.2%, RA = 0.1%, P = 0.002) (**Figure 5**).

Statistical analysis revealed that whilst the abundance of most fungal DNA was

unaffected by disease state, select organisms were differentially abundant or present

(**Figure 6**). An unclassified member of the *Hypocreales* order (P = 0.012) and a member

of the genus *Aspergillus* was detected almost exclusively in RA synovial fluid samples

only (P = 0.14 including all data, P = 0.057 excluding samples where <1% of reads

mapped to this taxa). Conversely, a member of the order *Malasseziale*s (P = 0.002),

and the genus *Cladosporium* (P = 0.019) were observed almost exclusively in control

synovial fluid.

Figure 5 - Relative abundance of taxa representing >1% of the total fungal population in

each synovial fluid sample (RA, n = 12) and (Control, n = 9) as determined by

amplification and sequencing of the fungal ITS2 gene. Data are mean abundance

expressed as a percentage of the total fungal sequence count.

Figure 6 - Relative abundance of (A) *Hypocreales*, (B), *Aspergillus*, (C) *Malasseziales*,

and (D) Cladosporium identified in the synovial fluid of RA patients and healthy control

subjects. Data determined by the amplification and sequencing of the ITS2 gene. (A)

The relative abundance of order *Hypocreales* was increased in synovial fluid of RA

patients relative to healthy control synovial fluid. (B) The relative abundance of genus

Aspergillus was increased in RA synovial fluid samples compared with healthy subjects

(borderline significant). (C), and (D) The abundance of the order *Malasseziales* and

genus *Cladosporium* were significantly decreased in the synovial fluid of RA patients in

comparison to the synovial fluid of healthy control subjects. Data are individual

expression values expressed as a percentage of the total fungal read count in each

sample. The statistical significance between groups was determined by unpaired t test

where $P \leq 0.05$.

Inflammatory markers and fungal microbiome PCoA cluster

Based solely upon the clustering of samples observed following PCoA, we investigated

whether samples in PCoA "cluster 1" had significantly different inflammatory marker

profiles, in addition to different microbial populations, than the remaining samples. IL-6

was significantly higher in PCoA cluster 1 compared with the remaining samples (Two

Way ANOVA; $P = 0.016$). Further, the interaction between PCoA cluster and disease

status (control or RA) was also significant ($P = 0.048$). All other inflammatory markers

were unaffected by PCoA cluster ($P > = 0.05$).

Discussion

Mounting evidence suggests pan-microbiome dysbiosis in patients with RA. Moreover, various studies spanning many decades link the development of RA with infections of the oral tissues or urinary system, and report the presence of microorganisms or microorganism-derived genetic material in the synovial fluid; a compartment classically considered to be sterile. To our surprise, only one study has investigated the potential for a "synovial fluid microbiome" using next generation sequencing [5]. To this end, we investigated the presence of bacterial and fungal DNA in the synovial fluid of rheumatoid arthritis (RA) patients and healthy control subjects through amplification and sequencing of the 16S rRNA and ITS2 genes.

Evidence of bacteria

Synovial fluid samples from 25 subjects were analysed in this study and our analyses revealed the presence of a complex synovial microbiome community in health and disease. Bacterial 16S rRNA was detected in fourteen out of sixteen (87.5%) RA patients, and nine out of nine (100%) healthy control subjects. At the phylum level, the synovial fluid microbiome was predominated by Proteobacteria (81.1%) and Firmicutes (18.5%), and to a much lesser extent, Actinobacteria (0.3%) and Bacteroidetes (0.1%). These phyla were also identified as the most abundant in the only other comparable study [5], albeit in slightly different proportions. Whilst the major phyla identified appear

similar to those of the blood microbiome [59], they are present in the synovial fluid at

different levels, suggesting a distinct community structure.

At the genus level, our synovial fluid samples were predominated by DNA of the genus

Pseudomonas and *Enterococcus,* to a lesser extent, unidentified taxa of the *Bacillales*

order and *Stenotrophomonas* genus. The genera *Raoultella* and *Bacillus* were detected

almost exclusively in the synovial fluid of our RA patients **(Figure 2 and 3)**. DNA of

Pseudomonas species, which was the most common genus to be identified in the study

cohort, has previously been identified in the synovial fluid of patients with RA [42, 60].

Whilst the bacterial communities characterised in the disease and control states were

largely consistent, the presence of the genus *Raoultella* was largely restricted to the RA

synovial fluid in our study (present only in 4 of our control samples with read counts <

0.3% yet detected in 6 of our RA patient samples with read abundances in the range 6.1

– 23.5%). Interestingly, when present, *Raoultella* often occupied a large proportion of

the total bacterial read count. This is the first study to describe the presence of DNA

from the genus *Raoultella* in RA synovial fluid samples, however, two major species are

capable of causing infections in man: *Raoultella ornithinolytica* and *Raoultella planticola*,

and have been shown to be capable of reaching the articular joint compartment [61-63].

Moreover, *Raoultella* are characterised as histamine generating bacteria [64], a key pro-

inflammatory molecule found in RA blood, articular cartilage, synovial fluid and synovial

tissue [65]. It is plausible that the presence and or indeed abundance of *Raoultella* DNA

may correlate with specific disease features (for example; disease stage, severity,

current disease activity, treatment regimen), and we therefore plan to explore this

relationship further in the future, particularly with the aim of identifying the origin of the

DNA found.

With regards the numerous reports detailing a link between oral and urinary tract

associated bacteria and RA, we reviewed the results of our taxonomic classification

looking specifically for the presence and abundance of these organisms. Bacteria of the

genus *Porphyromonas* were largely absent with the exception of three synovial fluid

samples (Control = 2, RA =1), which returned a small number of reads mapping to this

genus. These results differ from those of Zhao and colleagues who report the presence

of *Porphyromonas* in all of the samples they sequenced [5], irrespective of disease state

[5].   No reads mapping to the genus *Proteus* were identified in any sample analysed,

however numerous other members of the order Enterobacteriales to which *Proteus*

belongs were identified, including *Raoultella*. It is important to note here that none of our

RA patients reported oral or urinary conditions, and it is therefore plausible that the

above bacteria are present only during active oral or urinary disease. A separate study

is therefore warranted to further investigate these phenomena including RA patients

(without co-morbidities), RA patients with poor oral health, and patients with

periodontitis alone (without RA).

Evidence of fungi

Fungal DNA was identified in 12 out of 16 (75%) RA samples, and 8 out of 9 (88.8%)

healthy controls. Our various experimental controls consistently failed to produce a

visible band following PCR and agarose gel electrophoresis. Furthermore, DNA

quantification using the Qubit 3.0 high-sensitivity DNA kit (Invitrogen) was consistently

below the limit of detection. These indicate that the fungal DNA identified originates

from within the samples rather than from contaminants introduced. At the phylum level,

synovial fluid was predominated by members of the Basidiomycota and Ascomycota

phyla. At the genus level, synovial fluid samples were predominated by the genus

Malassezia, which accounted for 45.3% and 40.9% of the total number of fungal

sequences identified in the Control and RA donors respectively. To a lesser extent,

synovial fluid samples also contained organisms from the genera Penicillium, *Candida,*

Pichia, Aspergillus, Cladosporium, and *Debaryomyces.* Furthermore, genus level

classification revealed the presence of specific unclassified organisms belonging to the

classes *Tremellomycete*s and *Agaricomycetes*, and the orders *Hypocreales* and

Malasseziales (**Figure 6**).

Again we identified key taxa that appeared to be differentially present or abundant

dependent on disease status. The order *Hypocreales* was selectively abundant in RA

patients (7 / 12) and absent from healthy control synovial fluid. Interestingly, when

present, members of the Hypocreales order occupied a large proportion of the total

fungal read count (~ 40%). Evidence for the presence of members of this order have

been previously identified in the healthy human oral cavity [66] and found to be

ubiquitous in human plasma [67], however no reports directly link this order with RA.

Further, the abundance of genus *Aspergillus* was borderline significant; detected more commonly in the synovial fluid of RA patients, and occupying an average of ~ 10.0% of the read population when present. *Aspergillus* has been identified in the healthy human gut [68] and oral cavity [66] in various studies, and is a prevalent opportunistic pathogen of the lungs, particularly affecting immunocompromised individuals, where it causes a range of pulmonary diseases [69]. Furthermore, *Aspergillus* galactomannan antigen level has been shown to elevate in a non-specific manner among patients with RA [70], further hinting towards a possible association.

The presence members of the genus Cladosporium were largely absent in the synovial fluid of all RA patients in comparison to the synovial fluid of healthy control subjects (present in 5 / 8 healthy control patients in comparison with 2 / 12 RA samples). Cladosporium has been shown to colonise the healthy human gut [68] and oral cavity [66], and is encountered commonly in human clinical samples [71], thus evidence of Cladosporium DNA in the healthy patient is perhaps not unexpected. That said, it's absence from the synovial fluid of the majority of RA patients may suggest an altered fungal community in some distant microbial niche, that presents as a reduction in translocated Cladosporium DNA. These findings all warrant further investigation.

In-vivo studies have demonstrated that mycotoxins may play a role in the pathogenesis

of RA [72]. Furthermore, cells such as Th17 that are involved in the immunity against

fungi are also implicated in the pathogenesis of RA [73]. These indicate that fungal

presence in synovial fluid of patients with RA may contribute to pathogenesis. However,

as fungal DNA was also found in healthy controls, further studies are needed, with

quantification of the load of fungal DNA and taxa specific components.

553   Changes in microbiome with synovial fluid cytokine levels

Numerous cytokines have been implicated in the pathogenesis of RA. Levels of IL-17-A,

IL-22, IL23, and IL-6 have been identified to be higher in diseased joints than in healthy

[74, 75], and IL-6 has been identified to play a key role in the pathogenesis of RA. It is

implicated in pathogenesis of local joint disease as well as systemic manifestations [76,

77]. Serum and synovial fluid IL-6 levels correlate with disease severity and activity [78,

79]. We found that IL-6 levels were higher among a sub-cluster of the RA population

defined according to the fungal and bacterial microbiome. As mentioned earlier, Th17

cells are important in the immunity against fungal pathogens, and IL-6 is involved in the

differentiation of lymphocytes to Th17 cells [77]. This could indicate that the differences

in the fungal microbiome within patients with RA might be related to disease severity

through an IL-6 – Th17 mediated pathway.

566   Limitations

Bacterial DNA was identified in 14/16 patients with RA, and in all disease free controls whereas fungal DNA was identified in 12 out of 16 RA samples, and 8 out of 9 healthy controls. These results are consistent with other published works that have previously reported the presence of bacterial DNA in the majority, but not necessarily all, samples investigated [80, 81]. A potential limitation of this study is that we did not elect to sequence of experimental controls, and relied on the absence of a detectable signal following agarose gel electrophoresis and QuBit analysis using the high sensitivity DNA kit. Whilst is plausible that sequencing could have revealed the presence of low abundance target DNA introduced during our experimental procedures we suggest that this (potential) DNA would be (1) present at a much lower concentration (given our experimental samples were clearly visible following gel electrophoresis relative to the absence of any bands for our experimental controls), and (2) consistent given that all samples were processed in parallel. Whilst we acknowledge this potential limitation, it does not detract from our finding of statistically different levels of specific bacteria between the two experimental groups, despite all samples being processed at the same time utilising the same specific batches of reagents.

Conclusions

This is one of the first studies to report the existence of a synovial fluid microbiome, and to determine that this is modulated by disease status (RA) as are other classical

microbiome niches. Our recent work on the blood microbiome [4], and that of others [82-

92], characterised a complex community of organisms (or the genetic material thereof,

more specifically) that likely originate in one of the classical microbiome niches (the gut,

mouth, urogenital tract, skin) and reach the circulation through a process termed

atopobiosis. We hypothesise that the same process may be true of the community we

characterised here in the synovial fluid, with the circulatory system acting as a conduit

between their usual place of habitation and the synovial compartment. Further studies

incorporating pan-microbiome analysis are required to support these data.

Preprint

601 This manuscript has been published as a preprint bioRxiv [93].

References

1. Marchesi, J.R. and J. Ravel, *The vocabulary of microbiome research: a proposal.* Microbiome, 2015. **3**: p. 31.

2. Sender, R., S. Fuchs, and R. Milo, *Revised Estimates for the Number of Human and Bacteria Cells in the Body.* PLoS Biol, 2016. **14**(8): p. e1002533.

3. Markova, N.D., *L-form bacteria cohabitants in human blood: significance for health and diseases.* Discov Med, 2017. **23**(128): p. 305-313.

4. Whittle, E., et al., *Multi-Method Characterisation of the Human Circulating Microbiome.* BioRxiv, 2018.

5. Zhao, Y., et al., *Detection and characterization of bacterial nucleic acids in culture-negative synovial tissue and fluid samples from rheumatoid arthritis or osteoarthritis patients.* Sci Rep, 2018. **8**(1): p. 14305.

6. Mao, Q., et al., *Interplay between the lung microbiome and lung cancer.* Cancer Lett, 2018. **415**: p. 40-48.

7. Moffatt, M.F. and W.O. Cookson, *The lung microbiome in health and disease.* Clin Med (Lond), 2017. **17**(6): p. 525-529.

8. O'Dwyer, D.N., R.P. Dickson, and B.B. Moore, *The Lung Microbiome, Immunity, and the Pathogenesis of Chronic Lung Disease.* J Immunol, 2016. **196**(12): p. 4839-47.

9. Wang, L., et al., *Role of the Lung Microbiome in the Pathogenesis of Chronic Obstructive Pulmonary Disease.* Chin Med J (Engl), 2017. **130**(17): p. 2107-2111.

10. Gibofsky, A., *Overview of epidemiology, pathophysiology, and diagnosis of rheumatoid arthritis.* Am J Manag Care, 2012. **18**(13 Suppl): p. S295-302.

627   11.   Pretorius, E., et al., *Major involvement of bacterial components in rheumatoid arthritis and its*
628   *accompanying oxidative stress, systemic inflammation and hypercoagulability.* Exp Biol Med
629   (Maywood), 2017. **242**(4): p. 355-373.

630   12.   Zhang, X., et al., *The oral and gut microbiomes are perturbed in rheumatoid arthritis and partly*
631   *normalized after treatment.* Nat Med, 2015. **21**(8): p. 895-905.

632   13.   Lopez-Oliva, I., et al., *Dysbiotic Subgingival Microbial Communities in Periodontally Healthy*
633   *Patients With Rheumatoid Arthritis.* Arthritis Rheumatol, 2018. **70**(7): p. 1008-1013.

634   14.   Scher, J.U., et al., *Expansion of intestinal Prevotella copri correlates with enhanced susceptibility*
635   *to arthritis.* Elife, 2013. **2**: p. e01202.

636   15.   Taneja, V., *Arthritis susceptibility and the gut microbiome.* FEBS Lett, 2014. **588**(22): p. 4244-9.

637   16.   Chen, J., et al., *An expansion of rare lineage intestinal microbes characterizes rheumatoid*
638   *arthritis.* Genome Med, 2016. **8**(1): p. 43.

639   17.   Liu, X., et al., *Role of the Gut Microbiome in Modulating Arthritis Progression in Mice.* Sci Rep,
640   2016. **6**: p. 30594.

641   18.   Scher, J.U., D.R. Littman, and S.B. Abramson, *Microbiome in Inflammatory Arthritis and Human*
642   *Rheumatic Diseases.* Arthritis Rheumatol, 2016. **68**(1): p. 35-45.

643   19.   Maeda, Y., et al., *Dysbiosis contributes to arthritis development via activation of autoreactive T*
644   *cells in the intestine.* Arthritis Rheumatol, 2016.

645   20.   Vaahtovuo, J., et al., *Fecal microbiota in early rheumatoid arthritis.* J Rheumatol, 2008. **35**(8): p.
646   1500-5.

647   21.   Liu, X., et al., *Analysis of fecal Lactobacillus community structure in patients with early*
648   *rheumatoid arthritis.* Curr Microbiol, 2013. **67**(2): p. 170-6.

649   22.   Wu, X., et al., *Alterations of Gut Microbiome in Rheumatoid Arthritis.* Osteoarthritis and
650   Cartilage, 2017. **25**: p. S287-S288.

23. Brusca, S.B., S.B. Abramson, and J.U. Scher, *Microbiome and mucosal inflammation as extra-*
articular triggers for rheumatoid arthritis and autoimmunity. Curr Opin Rheumatol, 2014. **26**(1):
p. 101-7.

24. Demoruelle, M.K., et al., *The lung may play a role in the pathogenesis of rheumatoid arthritis.* Int
J Clin Rheumtol, 2014. **9**(3): p. 295-309.

25. Scher, J.U., et al., *The lung microbiota in early rheumatoid arthritis and autoimmunity.*
Microbiome, 2016. **4**(1): p. 60.

26. Ebringer, A. and T. Rashid, *Rheumatoid arthritis is an autoimmune disease triggered by Proteus*
urinary tract infection. Clin Dev Immunol, 2006. **13**(1): p. 41-8.

27. Ebringer, A. and T. Rashid, *Rheumatoid arthritis is caused by a Proteus urinary tract infection.*
APMIS, 2014. **122**(5): p. 363-8.

28. Tani, Y., et al., *Antibodies to Klebsiella, Proteus, and HLA-B27 peptides in Japanese patients with*
ankylosing spondylitis and rheumatoid arthritis. J Rheumatol, 1997. **24**(1): p. 109-14.

29. Wilson, C., et al., *Correlation between anti-Proteus antibodies and isolation rates of P. mirabilis*
in rheumatoid arthritis. Rheumatol Int, 1997. **16**(5): p. 187-9.

30. Chi, H. and R.A. Flavell, *Innate recognition of non-self nucleic acids.* Genome Biol, 2008. **9**(3): p.
211.

31. Atianand, M.K. and K.A. Fitzgerald, *Molecular basis of DNA recognition in the immune system.* J
Immunol, 2013. **190**(5): p. 1911-8.

32. Castañeda-Delgado, J.E., et al., *Type I Interferon Gene Response Is Increased in Early and*
Established Rheumatoid Arthritis and Correlates with Autoantibody Production. Front Immunol,
2017. **8**: p. 285.

33.   Enninga, E.A., et al., *Immune Reactivation by Cell-Free Fetal DNA in Healthy Pregnancies Re-*
Purposed to Target Tumors: Novel Checkpoint Inhibition in Cancer Therapeutics. Front Immunol,
2015. **6**: p. 424.

34.   Rossol, M., et al., *LPS-induced cytokine production in human monocytes and macrophages.* Crit
Rev Immunol, 2011. **31**(5): p. 379-446.

35.   Ross, J.J., *Septic Arthritis of Native Joints.* Infect Dis Clin North Am, 2017. **31**(2): p. 203-218.

36.   Martinez-Martinez, R.E., et al., *Detection of periodontal bacterial DNA in serum and synovial*
fluid in refractory rheumatoid arthritis patients. J Clin Periodontol, 2009. **36**(12): p. 1004-10.

37.   Moen, K., et al., *Synovial inflammation in active rheumatoid arthritis and psoriatic arthritis*
facilitates trapping of a variety of oral bacterial DNAs. Clin Exp Rheumatol, 2006. **24**(6): p. 656-
63.

38.   Ogrendik, M., *Rheumatoid arthritis is linked to oral bacteria: etiological association.* Modern
rheumatology, 2009. **19**(5): p. 453-456.

39.   Reichert, S., et al., *Detection of oral bacterial DNA in synovial fluid.* Journal of clinical
periodontology, 2013. **40**(6): p. 591-598.

40.   Témoin, S., et al., *Identification of oral bacterial DNA in synovial fluid of arthritis patients with*
native and failed prosthetic joints. Journal of clinical rheumatology: practical reports on
rheumatic & musculoskeletal diseases, 2012. **18**(3): p. 117.

41.   Gérard, H.C., et al., *Chromosomal DNA from a variety of bacterial species is present in synovial*
tissue from patients with various forms of arthritis. Arthritis Rheum, 2001. **44**(7): p. 1689-97.

42.   Kempsell, K.E., et al., *Reverse transcriptase-PCR analysis of bacterial rRNA for detection and*
characterization of bacterial species in arthritis synovial tissue. Infect Immun, 2000. **68**(10): p.
6012-26.

43.   Hitchon, C.A., et al., *Antibodies to porphyromonas gingivalis are associated with anticitrullinated protein antibodies in patients with rheumatoid arthritis and their relatives.* J Rheumatol, 2010. **37**(6): p. 1105-12.

44.   Lundberg, K., et al., *Antibodies to citrullinated alpha-enolase peptide 1 are specific for rheumatoid arthritis and cross-react with bacterial enolase.* Arthritis Rheum, 2008. **58**(10): p. 3009-19.

45.   Ogrendik, M., et al., *Serum antibodies to oral anaerobic bacteria in patients with rheumatoid arthritis.* MedGenMed, 2005. **7**(2): p. 2.

46.   Moen, K., et al., *Immunoglobulin G and A antibody responses to Bacteroides forsythus and Prevotella intermedia in sera and synovial fluids of arthritis patients.* Clin Diagn Lab Immunol, 2003. **10**(6): p. 1043-50.

47.   Bartold, P.M., R.I. Marshall, and D.R. Haynes, *Periodontitis and Rheumatoid Arthritis: A Review.* J Periodontol, 2005. **76 Suppl 11S**: p. 2066-2074.

48.   Chen, H.H., et al., *Association between a history of periodontitis and the risk of rheumatoid arthritis: a nationwide, population-based, case-control study.* Ann Rheum Dis, 2013. **72**(7): p. 1206-11.

49.   de Pablo, P., T. Dietrich, and T.E. McAlindon, *Association of periodontal disease and tooth loss with rheumatoid arthritis in the US population.* J Rheumatol, 2008. **35**(1): p. 70-6.

50.   Forner, L., et al., *Incidence of bacteremia after chewing, tooth brushing and scaling in individuals with periodontal inflammation.* J Clin Periodontol, 2006. **33**(6): p. 401-7.

51.   Horliana, A.C., et al., *Dissemination of periodontal pathogens in the bloodstream after periodontal procedures: a systematic review.* PLoS One, 2014. **9**(5): p. e98271.

52.   Koziel, J., P. Mydel, and J. Potempa, *The link between periodontal disease and rheumatoid arthritis: an updated review.* Curr Rheumatol Rep, 2014. **16**(3): p. 408.

53.   Ortiz, P., et al., *Periodontal therapy reduces the severity of active rheumatoid arthritis in patients*

721   *treated with or without tumor necrosis factor inhibitors.* J Periodontol, 2009. **80**(4): p. 535-40.

722   54.   Scher, J.U., et al., *Periodontal disease and the oral microbiota in new-onset rheumatoid arthritis.*

723   Arthritis Rheum, 2012. **64**(10): p. 3083-94.

724   55.   Zhao, X., et al., *Association of Periodontitis with Rheumatoid Arthritis and the Effect of Non-*

725   *Surgical Periodontal Treatment on Disease Activity in Patients with Rheumatoid Arthritis.* Med

726   Sci Monit, 2018. **24**: p. 5802-5810.

727   56.   Li, R., et al., *Rheumatoid arthritis and periodontal disease: What are the similarities and*

728   *differences?* Int J Rheum Dis, 2017. **20**(12): p. 1887-1901.

729   57.   Cheng, Z., et al., *Periodontal disease and periodontal bacteria as triggers for rheumatoid*

730   *arthritis.* Best Pract Res Clin Rheumatol, 2017. **31**(1): p. 19-30.

731   58.   Potempa, J., P. Mydel, and J. Koziel, *The case for periodontitis in the pathogenesis of rheumatoid*

732   *arthritis.* Nat Rev Rheumatol, 2017. **13**(10): p. 606-620.

733   59.   Whittle, E., et al., *Multi-Method Characterization of the Human Circulating Microbiome.*

734   Frontiers in Microbiology, 2019. **9**(3266).

735   60.   Olsen-Bergem, H., et al., *Juvenile idiopathic arthritis and rheumatoid arthritis: bacterial diversity*

736   *in temporomandibular joint synovial fluid in comparison with immunological and clinical*

737   *findings.* Int J Oral Maxillofac Surg, 2016. **45**(3): p. 318-22.

738   61.   Bonnet, E., et al., *Joint infection due to Raoultella planticola: first report.* Infection, 2017. **45**(5):

739   p. 703-704.

740   62.   Levorova, J., et al., *Septic arthritis of the temporomandibular joint caused by rare bacteria*

741   *Raoultella ornithinolytica.* Int J Oral Maxillofac Surg, 2017. **46**(1): p. 111-115.

742   63.   Venus, K., S. Vaithilingam, and I.I. Bogoch, *Septic arthritis of the knee due to Raoultella*

743   *ornithinolytica.* Infection, 2016. **44**(5): p. 691-2.

64.   Kanki, M., et al., *Klebsiella pneumoniae produces no histamine: Raoultella planticola and*

Raoultella ornithinolytica strains are histamine producers. Appl Environ Microbiol, 2002. **68**(7):

p. 3462-6.

65.   Kim, K.W., et al., *Histamine and Histamine H4 Receptor Promotes Osteoclastogenesis in*

Rheumatoid Arthritis. Sci Rep, 2017. **7**(1): p. 1197.

66.   Peters, B.A., et al., *The oral fungal mycobiome: characteristics and relation to periodontitis in a*

pilot study. BMC Microbiol, 2017. **17**(1): p. 157.

67.   Beatty, M., et al., *Small RNAs from plants, bacteria and fungi within the order Hypocreales are*

ubiquitous in human plasma. BMC Genomics, 2014. **15**: p. 933.

68.   Hallen-Adams, H.E. and M.J. Suhr, *Fungi in the healthy human gastrointestinal tract.* Virulence,

2017. **8**(3): p. 352-358.

69.   Singh, H., et al., *Pulmonary Aspergilloma in Rheumatoid Arthritis.* Med J Armed Forces India,

2003. **59**(3): p. 254-6.

70.   Horie, M., et al., *Nonspecific elevation of serum Aspergillus galactomannan antigen levels in*

patients with rheumatoid arthritis. Respir Investig, 2016. **54**(1): p. 44-9.

71.   Sandoval-Denis, M., et al., *Cladosporium Species Recovered from Clinical Samples in the United*

States. J Clin Microbiol, 2015. **53**(9): p. 2990-3000.

72.   Jahreis, S., et al., *Mold metabolites drive rheumatoid arthritis in mice via promotion of IFN-*

gamma- and IL-17-producing T cells. Food Chem Toxicol, 2017. **109**(Pt 1): p. 405-413.

73.   van Hamburg, J.P. and S.W. Tas, *Molecular mechanisms underpinning T helper 17 cell*

heterogeneity and functions in rheumatoid arthritis. J Autoimmun, 2018. **87**: p. 69-81.

74.   Kim, K.-W., et al., *Interleukin-22 promotes osteoclastogenesis in rheumatoid arthritis through*

induction of RANKL in human synovial fibroblasts. Arthritis & Rheumatism, 2012. **64**(4): p. 1015-

1023.

75.   Moura, R.A., et al., *Cytokine profile in serum and synovial fluid of patients with established rheumatoid arthritis.* Annals of the Rheumatic Diseases, 2010. **69**(Suppl 2): p. A51-A51.

76.   Yoshida, Y. and T. Tanaka, *Interleukin 6 and rheumatoid arthritis.* Biomed Res Int, 2014. **2014**: p. 698313.

77.   Srirangan, S. and E.H. Choy, *The role of interleukin 6 in the pathophysiology of rheumatoid arthritis.* Ther Adv Musculoskelet Dis, 2010. **2**(5): p. 247-56.

78.   Rajaei, E., et al., *Evaluating the relationship between serum level of interleukin-6 and rheumatoid arthritis severity and disease activity.* Curr Rheumatol Rev, 2019.

79.   Matsumoto, T., T. Tsurumoto, and H. Shindo, *Interleukin-6 levels in synovial fluids of patients with rheumatoid arthritis correlated with the infiltration of inflammatory cells in synovial membrane.* Rheumatol Int, 2006. **26**(12): p. 1096-100.

80.   Sun, W., et al., *Bacterial diversity in synovial fluids of patients with TMD determined by cloning and sequencing analysis of the 16S ribosomal RNA gene.* Oral Surg Oral Med Oral Pathol Oral Radiol Endod, 2008. **105**(5): p. 566-71.

81.   Siala, M., et al., *Analysis of bacterial DNA in synovial tissue of Tunisian patients with reactive and undifferentiated arthritis by broad-range PCR, cloning and sequencing.* Arthritis Res Ther, 2008. **10**(2): p. R40.

82.   Kell, D.B. and E. Pretorius, *On the translocation of bacteria and their lipopolysaccharides between blood and peripheral locations in chronic, inflammatory diseases: the central roles of LPS and LPS-induced cell death.* Integr Biol (Camb), 2015. **7**(11): p. 1339-77.

83.   Bhattacharyya, M., et al., *The conserved phylogeny of blood microbiome.* Mol Phylogenet Evol, 2017. **109**: p. 404-408.

84.   Mangul, S., et al., *Total RNA Sequencing reveals microbial communities in human blood and disease specific effects.* bioRxiv, 2016.

85.   Païssé, S., et al., *Comprehensive description of blood microbiome from healthy donors assessed*

793   *by 16S targeted metagenomic sequencing.* Transfusion, 2016. **56**(5): p. 1138-47.

794   86.   Potgieter, M., et al., *The dormant blood microbiome in chronic, inflammatory diseases.* FEMS

795   Microbiol Rev, 2015. **39**(4): p. 567-91.

796   87.   Amar, J., et al., *Blood microbiota dysbiosis is associated with the onset of cardiovascular events*

797   *in a large general population: the D.E.S.I.R. study.* PLoS One, 2013. **8**(1): p. e54461.

798   88.   Rajendhran, J., et al., *Contrasting circulating microbiome in cardiovascular disease patients and*

799   *healthy individuals.* Int J Cardiol, 2013. **168**(5): p. 5118-20.

800   89.   Dinakaran, V., et al., *Elevated levels of circulating DNA in cardiovascular disease patients:*

801   *metagenomic profiling of microbiome in the circulation.* PLoS One, 2014. **9**(8): p. e105221.

802   90.   Sze, M.A., et al., *Changes in the bacterial microbiota in gut, blood, and lungs following acute LPS*

803   *instillation into mice lungs.* PLoS One, 2014. **9**(10): p. e111228.

804   91.   Jeon, S.J., et al., *Blood as a route of transmission of uterine pathogens from the gut to the uterus*

805   *in cows.* Microbiome, 2017. **5**(1): p. 109.

806   92.   Martel, J., et al., *Pleomorphic bacteria-like structures in human blood represent non-living*

807   *membrane vesicles and protein particles.* Sci Rep, 2017. **7**(1): p. 10650.

808   93.   Hammad, D.B. and D.P. Tonge, *Molecular Characterisation of the Synovial Fluid Microbiome.*

809   bioRxiv, 2018.

Figure 1 - Principal Coordinate Analysis plot generated from a Bray Curtis distance matrix of synovial fluid bacterial community structure for control subjects (red) and arthritic patients (blue) as determined via amplification and sequencing of the 16S rRNA gene variable region 4 (V4). Proportions of variation explained by the principal coordinates are designated on the axes. PCoA found that the maximal variation were 62.2% (PC1), and 22.9% (PC2). The microbiome of samples that appear in close proximity to each other are more similar in composition. The data points in each pane are sized according to the synovial concentration of IL-17A (A), IL-22 (B), IL-23 (C), and IL-6 (D).

Figure 2 - Relative abundance of genera representing >1% of the total bacterial population in synovial fluid samples of rheumatoid arthritis (RA, n = 14) and control (Control, n = 9) samples as determined by amplification and sequencing of the 16S rRNA gene variable region 4. Data are mean abundance expressed as a percentage of the total bacterial sequence count.

Figure 3 - Relative abundance of the genus *Raoultella*, identified in the synovial fluid of healthy donors and RA patients by amplification and sequencing of the 16S rRNA gene variable region. A borderline significant (p = 0.051) increase in the mean abundance of

Raoultella was identified in RA synovial fluid relative to healthy control synovial fluid.

Data are individual sample abundances expressed as a percentage of the total bacterial

sequence count.

Figure 4 - Principal Coordinate Analysis plot generated from a Bray Curtis distance

matrix of synovial fluid fungal community structure for control subjects (red) and arthritic

patients (blue) as determined by amplification and sequencing of the ITS2 gene.

Proportions of variation explained by the principal coordinates are designated on the

axes. PCoA found that the maximal variation were 30.6% (PC1), and 20.1%. The

microbiome of samples that appear in close proximity to each other are more similar in

composition. The data points in each pane are sized according to the synovial

concentration of IL-17A (A), IL-22 (B), IL-23 (C), and IL-6 (D).

Figure 5 - Relative abundance of taxa representing >1% of the total fungal population in

each synovial fluid sample (RA, n = 12) and (Control, n = 9) as determined by

amplification and sequencing of the fungal ITS2 gene. Data are mean abundance

expressed as a percentage of the total fungal sequence count.

Figure 6 - Relative abundance of (A) *Hypocreales*, (B), *Aspergillus*, (C) *Malasseziales*,

and (D) Cladosporium identified in the synovial fluid of RA patients and healthy control

subjects. Data determined by the amplification and sequencing of the ITS2 gene. (A)

The relative abundance of order *Hypocreales* was increased in synovial fluid of RA

patients relative to healthy control synovial fluid. (B) The relative abundance of genus

Aspergillus was increased in RA synovial fluid samples compared with healthy subjects

(borderline significant). (C), and (D) The abundance of the order *Malasseziales* and

genus *Cladosporium* were significantly decreased in the synovial fluid of RA patients in

comparison to the synovial fluid of healthy control subjects. Data are individual

expression values expressed as a percentage of the total fungal read count in each

sample. The statistical significance between groups was determined by unpaired t test

where $P \leq 0.05$.

	Control	RA
■ g__Bacillus	0.00%	1.20%
■ g__Stenotrophomonas	1.50%	4.20%
■ g__Raoultella	0.10%	6.50%
■ o__Bacillales;Ambiguous_taxa;Ambiguous_taxa	4.00%	7.40%
■ g__Enterococcus	11.70%	11.40%
■ g__Pseudomonas	80.60%	66.10%

	Control	RA
o__Malasseziales_unidentified	1.2%	0.1%
g__Debaryomyces	4.7%	0.4%
g__Cladosporium	5.1%	0.1%
c__Agaricomycetes_unidentified	3.2%	2.2%
c__Tremellomycetes_unidentified	3.0%	2.8%
g__Aspergillus	0.2%	6.2%
g__Pichia	8.9%	0.0%
g__Candida	11.9%	0.0%
g__Penicillium	0.7%	15.0%
o__Hypocreales_other_other	0.0%	26.4%
g__Malassezia	45.3%	40.9%