1

Diatoms diversify and turn over faster in freshwater than marine environments

Abstract

Many clades that span the marine–freshwater boundary are disproportionately more diverse in the younger, shorter-lived, and scarcer freshwater environments than they are in the marine realm. This disparity is thought to be related to differences in diversification rates between marine and freshwater lineages. However, marine and freshwaters are not ecologically homogeneous, so the study of diversification across the salinity divide should also account for other potentially interacting variables. In diatoms, freshwater and substrate-associated (benthic) lineages are several-fold more diverse than their marine and suspended (planktonic) counterparts. These imbalances provide an excellent system to understand whether these variables interact with diversification. Using multistate hidden-state speciation and extinction models we found that freshwater lineages diversify faster than marine lineages regardless of whether they inhabit the plankton or the benthos. Freshwater lineages also had higher turnover rates (speciation + extinction), suggesting that habitat transitions impact speciation and extinction rates jointly. The plankton–benthos contrast was also consistent with state-dependent diversification, but with modest differences in diversification and turnover rates. Asymmetric, and bidirectional transitions rejected hypotheses about the plankton and freshwaters as absorbing, inescapable habitats. Our results further suggest that the high turnover rate of freshwater diatoms is related to high turnover of freshwater systems themselves.

Introduction

Freshwater habitats comprise just 2% of the biosphere, yet in lineages that span the salinity gradient the magnitude of freshwater diversity is often comparable, or surpasses, species richness in the marine realm, which covers 70% of Earth's surface (Mora et al. 2011; Appeltans et al. 2012; Grosberg et al. 2012; Wiens 2015; Miller and Wiens 2017). This is striking given that most of these lineages originated in the oceans, and so—all else being equal—have had more time to accumulate diversity within their ancestral marine environment. Across animals there is a disconnect between species richness and time for accumulation of diversity between marine and freshwaters [e.g., flatworms, rotifers, molluscs, and ray-finned fish (Bloom et al. 2013; Betancur-R et al. 2015; Guinot and Cavin 2015; Wiens 2015; Miller and Wiens 2017)]. Direct tests of asymmetry in diversification have revealed that in amniotes, for example, marine lineages have higher extinction rates (Miller and Wiens 2017; but see Vermeij et al. 2018), whereas marine fish have higher net diversification compared to their freshwater counterparts (Betancur-R et al. 2015). To date, most of the research on this question has focused on animal systems, where there does not appear to be a clear consensus on what is driving disparities in species richness across the marine–freshwater divide (Betancur-R et al. 2015; Guinot and Cavin 2015; Wiens 2015; Miller and Wiens 2017). Microbes, including protists, might provide novel insights into these questions, as they are diverse at both ends of the salinity gradient (Mora et al.

2011; Appeltans et al. 2012) and offer alternative perspectives due to their vastly different
dispersal capabilities, population sizes, and physiological features compared to animals.
Here we focus on diatoms, a species-rich lineage of unicellular eukaryotic algae that play
important roles in the global cycling of carbon and oxygen and form the base of many aquatic
food webs (Armbrust 2009). Although, our knowledge of diatom diversity is far from complete,
a clear pattern of imbalanced species richness across the marine–freshwater divide exists, with
many more species known from freshwaters than the ancestral marine environment (Kociolek
et al. 2017). Most diatom genera are thought to be restricted to either marine or freshwaters
(Round and Sims 1980; Round et al. 1990; Mann 1999), and transitions between these
environments are thought to be challenging, as migrants need to mitigate osmotic stress linked
to changes in cellular water potential, salt stress caused by increased uptake or loss of ions, and
changes in the ratios of intracellular ions (Kirst 1990, 1996). These considerations form the basis
of a long-held view that colonizations of freshwaters by diatoms are infrequent and irreversible
events (e.g., the Rubicon hypothesis of Mann 1999). This hypothesis posits that the modern
freshwater flora has been assembled from a small set of phylogenetically diverse colonizers
(rarity) and that once adapted to freshwaters, these lineages rarely, if ever, cross back to their
ancestral marine habitat (irreversibility), implying that freshwaters are an 'absorbing' state.
Although the freshwater diatom flora is clearly a composite of many lineages that trace back to
independent colonization events (Alverson et al. 2007; Ruck et al. 2016), the frequency of
colonizations and irreversibility aspect of the Rubicon hypothesis have only been studied in
small clades, making it difficult to derive broad conclusions (Beaulieu and O'Meara 2018).
Moreover, studies to date have not explored the interaction between habitat transitions and
rates of diversification, which is essential for tests of character-dependent diversification
(Maddison 2006).
An important aspect that studies of diversification in marine and freshwaters have yet to
consider is that the focal contrast (marine vs. freshwater) is almost certainly not the sole
determinant of diversification dynamics, but rather, observed patterns are likely the product of
many potentially interacting variables related to the habitats and traits of different lineages
(Beaulieu and O'Meara 2016; Caetano et al. 2018). In protists, and in diatoms in particular, the
plankton (suspended growth habit) and benthos (substrate-associated growth habit) represent
another prominent environmental contrast, with an imbalanced species richness similar in
magnitude to the marine–freshwater divide. In diatoms, transitions between planktonic and
benthic habitats appear to be linked to both growth habit (e.g., colonial vs. solitary, Nakov et al.
2015) and mode of sexual reproduction (e.g., oogamous vs. anisogamous, Kooistra et al. 2007,
2009). For example, the numerous, flagellum-propelled male gametes of oogamous species
might facilitate colonization of the plankton, whereas the few, aflagellated gametes of
isogamous lineages might hinder survival and reproduction in dilute planktonic environments
(Kooistra et al. 2007, 2009).
Several additional lines of evidence point to potential interactions between the marine–
freshwater and plankton–benthos divides. First, the age and longevity of most freshwater lakes
supporting planktonic communities are linked to Earth's glaciation cycles, with only a few
dozen long-lived tectonic basins whose formation predates the last interglacial (ca. 130–115

thousand years ago; Mackay et al. 2010; Hampton et al. 2018). The majority of freshwater lakes
are geologically recent and have relatively short lifespans, so freshwater planktonic lineages
might turn over faster relative to the marine planktonic lineages from environments that have
existed continuously since the formation of the oceans. This hypothesis is supported by
observed changes the composition of planktonic diatom communities in sedimentary records
from ancient lakes that closely follow glaciation and drought cycles (Edlund and Stoermer 2000;
Cohen et al. 2007; Mackay et al. 2010; Stone et al. 2011).
Marine and freshwater plankton might also differ in population size and connectivity.
Marine populations have fewer physical barriers to migration and are generally expected to
have broader geographic distributions and higher rates of gene flow (Palumbi 1992, 1994). As a
result, marine planktonic diatoms might experience lower speciation and extinction rates
compared to their freshwater counterparts, which inhabit discontinuous and irregularly
distributed habitats. Still, in diatoms, gene flow can be limited among marine populations due
to isolation by distance (Casteleyn et al. 2010) or ecology (Whittaker and Rynearson 2017),
sometimes even at small spatial scales (Rynearson and Armbrust 2004; Godhe and Härnström
2010; Sjöqvist et al. 2015). These findings suggest that population subdivision might be even
stronger in freshwaters (Vanormelingen et al. 2008; Evans et al. 2009). Taken together, these
observations suggest that observed differences in diversification dynamics across the marine–
freshwater divide are confounded by differences in, for example, lineages inhabiting the marine
plankton vs. freshwater plankton or the marine plankton vs. marine benthos.
Here, we characterize diatom evolution across the marine–freshwater and plankton–
benthos divides using a large, time-calibrated phylogeny representing the known breadth of
extant diatom diversity. To account for variation in rates of diversification and colonizations
across environments, we used a hidden state speciation and extinction modeling framework
(HiSSE; Beaulieu and O'Meara 2016) modified for multiple characters. In doing so, we were able
to jointly estimate rates of colonization and diversification across multiple environments and
account for possible interactions between focal characters (i.e., marine plankton, marine
benthos, freshwater plankton, and freshwater benthos). We used this framework to test two sets
of hypotheses. First, we assessed whether diatom diversification (speciation – extinction) or
turnover (speciation + extinction) is faster in freshwaters due to, for example, differences in
habitat longevity or population size and connectivity. We test our marine–freshwater
hypotheses in two ways: irrespective of whether they have a suspended (plankton) or substrate-
associated (benthos) habit as well as by distinguishing between marine vs. freshwater plankton
and marine vs. freshwater benthos. Second, we tested several hypotheses about the
directionality and reversibility of transitions between habitats, given the traditionally held view
that both plankton and freshwaters are 'absorbing' environments where colonists invade, adapt,
and specialize to the point that emigration is unlikely or impossible. We discuss the
implications of our findings for diatom ancestry, diversification, and phylogenetic diversity
across the ecologically important marine–freshwater and plankton–benthos divides.
123

Materials and Methods
Phylogeny and diversity data
We used a time-calibrated phylogeny of 1,151 diatoms reconstructed from a concatenated
alignment of 11 genes (Nakov et al. 2018)—the largest available phylogenetic tree of diatoms
that covers their known phylogenetic and ecological breadth. We scored 1,132 out of the 1,151
species for two characters: "habitat" (plankton or benthos) and "salinity" (marine or freshwater)
based on taxonomic compilations (Round et al. 1990; Hasle and Syvertsen 1996; Lange-Bertalot
and Krammer 2000; Witkowski et al. 2000; Lobban et al. 2012) and previous work on character
evolution (Alverson et al. 2007; Nakov et al. 2015; Ruck et al. 2016).
Given a total estimated diatom species richness of up to 100,000 species (Mann and
Vanormelingen 2013), this phylogeny, although the most taxonomically inclusive to date, still
falls well short of the full diversity of diatoms. To account for under-sampling in our
diversification analyses, we estimated sampling fractions for each environment, e.g., the fraction
of sampled marine species from the total number of marine species described to date. Data for
salinity (marine–freshwater) were obtained from DiatomBase (Kociolek et al. 2017), which
reports the total estimated number of species per genus and the total estimated number of
marine species per genus, allowing for a straightforward calculation of the number of non-
marine (freshwater) species. Similar data were not available for habitat (plankton vs. benthos),
so we used the primary taxonomic literature (Round et al. 1990; Hasle and Syvertsen 1996;
Lange-Bertalot and Krammer 2000; Witkowski et al. 2000; Lobban et al. 2012) to record the
number of species per genus occurring in the plankton or benthos. Based on these numbers, we
then derived a per-genus fraction of planktonic species and multiplied the total estimated
genus-level diversity by this fraction to obtain an estimate of the number of planktonic and
benthic species per genus. We summed these numbers across all genera represented on our tree
to make diatom-wide estimates of species richness in each environment.
149
Analysis of the binary salinity and habitat characters
Hypotheses about associations between lineage diversification and discretely coded traits or
environments can be tested using state speciation and extinction models (-SSE; Maddison et al.
2007; FitzJohn et al. 2009; Goldberg et al. 2011; Beaulieu and O'Meara 2016). These models
jointly estimate rates of evolution of a focal character (i.e., transitions between states) and rates
of diversification (i.e., speciation and extinction), thereby disentangling whether an imbalanced
ratio of diversity between states reflects differences in diversification or asymmetry in rates of
colonization (Maddison 2006). SSE models come in two flavors, ones that assume the rates of
speciation, extinction, and transition are shared by all lineages occupying a particular state (e.g.,
BiSSE, MuSSE, GeoSSE) and those that relax this assumption through the usage of "hidden
states" (HiSSE, HiGeoSSE). A third set of models, called character-independent models (CID),
allow rate variation independent of the focal characters, and serve as non-trivial null models for
state-dependent diversification (Beaulieu and O'Meara 2016; Caetano et al. 2018).
Initially, for each binary character we fit a total of five models: **(1)** a "trivial null" model
(number of parameters, k=3, Fig. S1) with one parameter for the turnover rate (defined as
speciation + extinction, which measures the rate of any event, birth or death, per unit time) for

both observed states, and one parameter controlling the rate of character transitions (e.g.,
marine <-> freshwater); **(2)** a BiSSE model (k=5, Fig. S1) with separate turnover rates estimated
for lineages living in alternate environments and asymmetric transitions between environments
(e.g., marine -> freshwater ≠ freshwater -> marine); **(3)** a HiSSE model (k=13) with separate
turnover rates for each combination of observed and hidden states and eight transition rates for
changes between states (e.g., marine hidden state A -> marine hidden state B). This model
disallowed simultaneous transitions of the observed and hidden states (e.g., the rate marine
hidden state A -> freshwater hidden state B was set to zero; Fig. S1); **(4)** a CID2 model (k=6)
with two turnover rate parameters similar to the BiSSE model, but not linked to the observed
character states; and **(5)** a CID4 model (k=8) with five diversification parameters matching the
complexity of HiSSE (Fig. S1). In all cases, we kept the extinction fraction (defined as extinction/
speciation, which measures the ratio of death and birth events per unit time) constant across
states.
179
Analysis of a combined habitat + salinity characters
For the combined habitat + salinity character, we extended the HiSSE and CID framework to
handle a four-state case [i.e., marine plankton (MP), marine benthos (MB), freshwater plankton
(FP), and freshwater benthos (FB)]. For this analysis, we again fit a trivial null model that
assumed no variation in diversification as well as a multistate speciation and extinction model
(MuSSE; FitzJohn et al. 2009) that linked diversification rates to the four character states but
assumed no variation within regimes (Fig. S1). We then fit two multistate speciation and
extinction models with hidden traits, MuHiSSE, and seven multistate character-independent
models (MuCID) with increasing numbers of hidden states (MuCID2 through MuCID8; Fig S1).
The most complex MuCID8 model had eight parameters for turnover and one for the extinction
fraction, matching the complexity of the MuHiSSE model but with diversification parameters
not linked to the observed character states (Fig. S1). For both types of models, we allowed
transition rates between environments to differ (e.g., MB -> MP ≠ MP -> MB), and we again
disallowed dual transitions (e.g., MB -> FP = 0). The two MuHiSSE models differed in how they
handled transitions between hidden states. The first model (MuHiSSE reduced) assumed one
rate for all transitions between hidden states (e.g., MB hidden state A -> MB hidden state B = FP
hidden state A -> FB hidden state B), while the second model (MuHiSSE full) allowed separate
transition rates between hidden states (e.g., MB hidden state A -> MB hidden state B ≠ MB
hidden state B -> MB hidden state A; Fig S1). Finally, we assessed two alternative
implementations of the conditioning for survival of crown lineages, including the original
diversitree and HiSSE implementations (FitzJohn et al. 2009; Beaulieu and O'Meara 2016)
(HiSSE setting root.type="madfitz") and the conditioning scheme introduced by Herrera-Alsina
et al. (2019) (HiSSE setting root.type="herr_als").
In all cases, the extinction fraction was fixed across observed and hidden states
regardless of whether we allowed separate estimates of turnover rate for each state.. It is
important to note, however, that although extinction fraction is constrained among states,
converting turnover and extinction fraction into speciation and extinction allows estimation of
separate birth and death rates for each state (Beaulieu and O'Meara 2016). We accounted for

incomplete sampling by incorporating the sampling fractions described above directly into our
diversification analyses (FitzJohn et al. 2009; Beaulieu and O'Meara 2016). This approach
assumed random sampling of each observed character state. For all models, we aimed for 50
maximum likelihood optimizations, each initiated from a different starting point.
212
Diversification and transition hypotheses
After obtaining initial results on the fits of state-dependent and state-independent models, we
tested a second set of state-dependent models that constrained certain parameters to equality or
to zero, effectively removing them from the model. These models were designed to test specific
hypotheses related to asymmetry in rates of species turnover (speciation + extinction) or
transitions between environments. For these models, we constrained or dropped parameters for
each hidden state, permitting rate heterogeneity in the presence of our constraints and
maintaining a level of complexity comparable to MuHiSSE. For example, when testing whether
the plankton in freshwaters is an absorbing, dead-end state from which lineages rarely or never
emigrate, we fixed the plankton-to-benthos transition to zero in both hidden states but still
allowed variation in diversification rates within the freshwater plankton and in the rate of
colonization of the marine plankton from the freshwater plankton.
To assess the association between the plankton—benthos contrast and diversification,
we forced the turnover rate to be the same between: **(1)** the plankton and benthos in marine
environments (MP = MB), **(2)** the plankton and benthos in freshwater environments (FP = FB),
and **(3)** the previous two combined (MP = MB and FP = FB). These three models tested
specifically for differences in diversification dynamics related to suspended vs. substrate-
associated growth habit at either or both ends of the salinity gradient. We also evaluated
whether the marine-freshwater contrast mattered in the context of one type of growth habitat,
but not the other. For example, it could be that benthic lineages differ in turnover between
marine and fresh waters, whereas this distinction is less important in the plankton. This set of
hypotheses included models that forced turnover to be the same between: **(4)** the marine and
freshwater plankton (MP = FP), **(5)** marine and freshwater benthos (MB = FB), and **(6)** the two
previous combined (MP = FP and MB = FB).
We also tested hypotheses related to the directionality of transitions between habitats,
given the historical understanding of freshwaters and the plankton as absorbing states. In
modeling terms, these hypotheses translate to strongly asymmetric transition rates, with the
rates of transition outside of the putatively absorbing state equal to zero. For freshwaters, we set
up models in which: **(7)** the freshwater plankton is absorbing (FP -> MP = 0, but FP -> FB $\neq$ 0),
(8) the freshwater benthos is absorbing (FB -> MB = 0, but FB -> FP $\neq$ 0) and **(9)** freshwaters in
general are absorbing irrespective of growth habit (FP -> MP = 0 *and* FB -> MB = 0). For the
plankton, we had an analogous setup with models in which **(10)** freshwater plankton is
absorbing (FP -> FB = 0, but FP -> MP $\neq$ 0), **(11)** the marine plankton is absorbing (MP -> MB =
0, but MP -> FP $\neq$ 0) and **(12)** plankton is absorbing in both marine and freshwaters (FP -> FB =
0 *and* MP -> MB = 0). In all cases, we disabled only the transitions reflecting the change in state
in the trait of interest while allowing transitions between states of the other trait. For example,
for freshwater plankton, when we tested whether freshwaters were absorbing, we disallowed

the transition to marine plankton and allowed migration to the freshwater benthos, but when
we tested whether the plankton was absorbing, we disallowed the transition to the freshwater
benthos and allowed colonization of the marine plankton.
253
Post-processing of HiSSE and MuHiSSE model fits
In addition to estimating the most likely parameter values for predefined regimes, the models
included in HiSSE also allowed us to reconstruct the probabilities of the hidden states occupied
by each lineage. For example, a lineage might be freshwater, but its hidden state might be less
certain, with some probability of occupying either of the hidden states in the model. In such
cases it is unclear which parameter estimate should be assigned to a particular lineage because
the diversification and transition rates might differ substantially among hidden states. One way
to account for this uncertainty is to average the parameter estimates weighted by the
probabilities of occupying alternative hidden states. For example, if the model has two hidden
marine states with turnover waiting times of 5 and 10 million years respectively, and a species
was reconstructed as occupying marine hidden state *A* with probability of 0.3 and marine
hidden state *B* with probability of 0.7, then the waiting time assigned to it, reflecting the
uncertainty in the ancestral state reconstruction, would be the weighted average 5*0.3 + 10*0.7 =
8.5. The alternative would be to disregard the uncertainty in the reconstruction and assume that
the species in question simply occupies marine hidden state *B* (the more likely state) and assign
to it longer waiting time of 10 million years. This type of averaging of per-lineage diversification
parameters provides a measure of rate variation within regimes, because instead of splitting
species and their rates within a regime into two groups corresponding to the two hidden states,
we can obtain a distribution of weighted averages of the parameters. Even if character-
dependent diversification is supported, it is unlikely that all lineages assigned to a regime
would have diversified at the same rate, and this variation will be reflected in the hidden state
reconstructions. This model-averaging approach also alleviates the subjectivity of choosing
arbitrary thresholds for choosing models, and instead puts the emphasis of interpretation on the
parameter estimates (see Caetano et al., 2018 for further discussion of this point).
We reconstructed marginal ancestral states either from a single best model, when
support measured by the Akaike weight was **>** 99% or from all models supported by ≥ 1%
(HiSSE functions MarginRecon and MarginReconMuHiSSE). A vignette with further details
regarding MuHiSSE is available as part of the hisse R package. The data and scripts used here
are available from a Zenodo repository (DOI: 10.5281/zenodo.2558448). Model summaries and
figures for HiSSE objects can also be made with our R package "utilhisse" (https://github.com/
teofiln/utilhisse) and an interactive web application (https://diatom.shinyapps.io/hisse-web).
We also approximated the likelihood surface around the maximum likelihood estimates
using HiSSE's adaptive sampling routine (HiSSE functions SupportRegion and
SupportRegionMuHiSSE, Beaulieu & O'Meara 2016). This procedure samples the parameter
space around the MLEs and re-estimates the likelihood to find a range of parameter values
whose likelihood falls within 2 log-likelihood units of the best fit. A broad range would indicate
a flatter likelihood surface and less confidence in the parameter estimates. How far the sampler
wanders away from an MLE can be controlled with a scaling argument.

292
Results
Binary characters
Diatom diversity data showed strong disparities in species richness across both the marine–
freshwater and plankton–benthos divides (Mora et al. 2011; Kociolek et al. 2017).
Approximately 70% of the species described to date are freshwater and 83% are benthic. The
estimated numbers of species in marine and freshwater habitats are 3221 and 7349 species
respectively, and of these our phylogeny includes 680 (21%) marine and 452 (6%) freshwater
species. The estimated numbers of species in plankton and benthos are 1800 and 8770 species
respectively, and of these our phylogeny includes 358 (20%) planktonic and 774 (9%) benthic
species.
These imbalanced diversity ratios suggest that diversification rates might be related to
one or both of these characters. Indeed, BiSSE models that linked diversification and character
transition rates to the alternate environments provided a much better fit than models that
assumed one rate across the tree (Table 1). Freshwater and benthic lineages, as expected given
the diversity ratios, had waiting times (inverse of the rate) of speciation + extinction (i.e., net
turnover) that were 3.8 and 1.6 times faster, respectively, than their marine and planktonic
counterparts. Similarly, net diversification (speciation – extinction) was 3.7 times faster in
freshwater than marine lineages, and 1.6 times faster in benthic than planktonic diatoms. The
ratio of extinction to speciation (the extinction fraction) was high in both cases: 0.86 for the
marine–freshwater and 0.87 for the plankton–benthos contrast. According to BiSSE models,
both of these environmental gradients appear to have been important in shaping the diversity
of diatoms.
To assess variation in diversification and transition rates within regimes, we also
evaluated character-dependent (HiSSE) and character-independent (CID) models that included
hidden states, i.e., factors other than our focal characters that might have influenced
diversification. For both characters, we found substantial variation in the rates of diversification
and transitions between regimes, as inferred from the stronger support for models with hidden
states relative to BiSSE models (Table 1). A character-dependent HiSSE model was favored for
both binary traits with Akaike weights of $\omega=0.999$ (Table 1). Generally, freshwater lineages
speciated, but also went extinct, faster than marine lineages, resulting in faster overall species
turnover in freshwater (Table 2), but with variation among hidden states. The difference
between speciation and extinction (net diversification) favored marine lineages in one hidden
state (rate ratio marine:freshwater = 2.8), and freshwater lineages in the other hidden state (rate
ratio freshwater:marine = 13.4; Table 2). For habitat, the differences in waiting times to a net
gain of one species between planktonic and benthic lineages were less pronounced, with
plankton:benthos rate ratios ranging from 1.05 in one hidden state to 1.2 in the other (Table 3).
An alternative way to assess these environmental contrasts is to compare the
distributions of diversification rates after accounting for uncertainty in ancestral
reconstructions. Waiting time to a turnover event [either speciation or extinction in My
calculated as 1/(speciation+extinction)] in marine lineages was on average 11.5 My with a
standard deviation (SD) of 4.2 My, whereas freshwater diatoms had a mean waiting time of 3.9

My and SD = 1.4 My between turnover events (Fig. 1A). In contrast, although a HiSSE model
was favored, the ranges for turnover waiting time in planktonic (4.5 ± 1.8 My) and benthic (3.28
± 1.6 My) species were overlapping (Fig. 1B). The variation in net diversification was
proportional to turnover (Fig. 1A,B) and for both binary characters these tests highlighted broad
rate variation in diversification rates within regimes (Fig. 1A,B).
We tested hypotheses regarding the directionality of transitions between environments
by disabling the transition rates plankton -> benthos and freshwater -> marine (to enforce
freshwaters and planktonic environments to be absorbing). In both cases, these models were
poorly supported, suggesting that transitions across both environmental gradients are
bidirectional ('hisse F absorbing' and hisse P absorbing'; Table 1).
344
Combined salinity + habitat character
One way to further evaluate rate-variation within states is to combine salinity and habitat into a
single four-state character: marine–plankton, marine–benthos, freshwater–plankton, and
freshwater–benthos. With this approach, we can specifically test for differences between, for
example, plankton and benthos, while allowing for the possibility that planktonic lineages in
fresh vs. saline waters also differ in diversification. To assess diversification rate differences
across the four habitat + salinity regimes, we implemented a multistate variant of the HiSSE
model (MuHiSSE), with a separate turnover parameter for each of the four states and two
hidden states within each character combination. To guard against false-positive selection of
these highly parameterized models, we designed and tested a set of multistate character-
independent (MuCID) models with increasing numbers of diversification parameters, up to a
MuCID model with eight hidden traits matching the complexity of the MuHiSSE model [eight
turnover and one extinction fraction that can be transformed into eight speciation and eight
extinction rates (Beaulieu and O'Meara 2016; HiSSE functions ParameterTransform or
ParameterTransformMuHiSSE, Fig. S1).
Data for the combined salinity + habitat states mirrored disparities in species richness
observed for the individual binary traits. Marine plankton accounted for 11% (1155 species),
marine benthos 20% (2066 species), freshwater plankton 6% (645), and freshwater benthos 63%
of all diatom species (6704). Our phylogeny included 288 (14%) marine planktonic species, 380
(18%) marine benthic species, 69 (11%) freshwater planktonic species, and 381 (6%) freshwater
benthic species.
Marine planktonic diatoms generally have much deeper fossil records than benthic
lineages (Sims et al. 2006), and freshwater lineages are generally nested within, and younger
than their marine counterparts (Alverson 2014; Ruck et al. 2016). A superficial look at these data
might suggest that younger lineages are disproportionately more diverse than older ones,
perhaps due to differences in diversification dynamics. We compared a total of 23 models (Table
S1): 11 in the base set comparing character-dependent to character-independent models, six
testing specific hypotheses related to diversification parameters, and another six comparing
hypotheses about transition rates. From these 23 models, four were supported by an Akaike
weight (ω ≥ 1%; Table 4) and were retained for further analyses using model averaging (Table
4). These included a model where the plankton–benthos contrast had no effect (MP = MB and

FP = FB, ω = 84%), the most complex MuHiSSE model without constraints on turnover (ω =
10%), a model that constrained the transition rates such that the marine plankton was absorbing
(MP -> MB = 0, but MP -> FP ≠ 0, ω = 5%), and a model where the freshwater plankton was
absorbing (FP -> MP = 0, but FP -> FB ≠ 0, ω = 1%).
Net turnover and net diversification in freshwater habitats were 3 times faster than in
marine environments after averaging across these four models and over their hidden-state
ancestral reconstructions (Fig. 1C). Lineages with suspended and substrate-associated growth
habits had similar turnover rates in both marine and freshwaters due to the weight of the model
without a plankton–benthos effect. The waiting times to a net gain of one species estimated here
were in general agreement with previous estimates of net diversification in diatoms (Nakov et
al. 2018). For example, the net diversification waiting time for marine planktonic lineages under
the MuHiSSE model was 17 My in one and 2.7 My in the second hidden state, whereas the
waiting time for oogamous nonmotile diatoms (most common in the marine plankton) was
previously estimated as between 16.6 and 100 My (phylogenetic data) or roughly 34.5 My (fossil
data) (Nakov et al. 2018). The relatively high global extinction fraction ($\varepsilon = 0.8$; Table S2),
estimated here from extant species only, was similar to calculations made from Cenozoic
marine fossils ($\varepsilon = 0.75$; Nakov et al. 2018). Analyses using alternative implementations of the
conditioning of survival of crown lineages indicated that the two approaches produce very
similar results (Fig. S1, comparison for the base models).
395
Taxonomic bias
Like any diversification analysis that accounts for unsampled diversity, our results rely on the
assumption that the total (described and undescribed) species richness in our focal regimes is
directly proportional to the diversity described from those habitats to date. However, in
addition to differences in diversification rates, these lopsided diversity ratios could also arise
from unequal taxonomic effort or from nonstandard taxonomic practices (e.g., splitting vs.
lumping) between researchers studying different habitats. It is difficult to assess how these
problems might affect our analyses, or analyses of any group, without some non-arbitrary
method of scaling the diversity data or sampling fractions to account for possible unknown
biases related to, for example, over-splitting (e.g., in freshwater taxa) or lower taxonomic
research effort (e.g., in marine taxa). Nevertheless, we repeated the marine–freshwater analysis
with arbitrarily reduced marine sampling fractions, from the observed value of 21% down to
14%, 10%, 6%, and 2%. Irrespective of how low we set the marine sampling fraction, we found
that HiSSE models were still favored over character-independent models. With smaller
sampling fractions, the diversification and turnover rate estimates for marine lineages
increased, but the signal for faster diversification in freshwaters, at least in one of the hidden
states, was detected in all but the analysis with a marine sampling fraction of 2%. In the latter,
marine lineages diversified faster than freshwater in both hidden states. Overall, these results
indicated that, if the past nearly 250 years of taxonomic work on diatoms is completely biased
and non-representative of diatom diversity as a whole, then some of our inferences might be
affected.

To evaluate potential clade-specific sampling biases, we repeated the analysis of the
entire diatom phylogeny on four subclades with sufficient sampling across the two
environmental contrasts. Although it is not expected that model selection or parameter
estimates from the entire phylogeny should match those inferred from subtrees that differ,
among others, in age, diversity, ecology, life history, and genomic attributes (Beaulieu and
O'Meara 2018; Rabosky 2019), we found that the global result, i.e., faster turnover in freshwater
compared to marine lineages, was qualitatively replicated in most of the subtrees (see
Supplementary Text 1 and Fig. S2 for details).
425
Discussion
One of the most prominent features of aquatic biodiversity is the disparity in species richness
between marine and freshwater habitats. Marine habitats are geologically old, have existed
continuously since the formation of the oceans and amount to >99% of the habitable water on
Earth, whereas present-day freshwater habitats are much younger and have shorter lifespans
determined by glaciation, droughts, or tectonic events. Surprisingly, in lineages with substantial
diversity across the salinity gradient, the younger and derived freshwater clades are often more
species rich than their marine counterparts. This difference in diversity is generally a
consequence of higher diversification rates in freshwater environments relative to marine
habitats. However, freshwater lineages also tend to exhibit higher turnover rates, which might
partly explain the preponderance of diverse clades nearer the present. Pulses of speciation
followed by pulses of extinction are indicative of higher turnover rates. Importantly, differences
in turnover rates suggest that transitions into and out of freshwater environments jointly impact
speciation and extinction rates.
440
Rate variation and hidden states
Our analyses of the marine–freshwater and plankton–benthos contrasts, considered either
independently or in combination, revealed strong support for hidden state models, either
character-dependent (salinity and combined) or character-independent (habitat). This indicates,
as would be expected for a lineage as phylogenetically and ecologically diverse as diatoms, that
the focal characters alone (salinity, habitat or their combination) do not explain all of the
variation in diversification rates across the phylogeny, even after partitioning the data into four
regimes corresponding to the states of the combined habitat+salinity character. This further
underscores the need to account for the multitude of environmental, ecological, and trait factors
that were not explicitly considered here, but whose cumulative effect on diversification might
be comparable to the effects of salinity and habitat. To illustrate this point, we can compare the
differences in diversification between environments to the differences between hidden states
within an environment. Marine planktonic lineages reconstructed in one of the hidden states
had a waiting time to a speciation event of about 17 My. This was similar to marine benthic
lineages in the same hidden state (17 My), but was several times longer than marine planktonic
lineages in the other hidden state (2.7 My; Fig. 2; Table S2). In other words, growing suspended
in the water column or associated with a substrate conferred a smaller difference in
diversification than differences conferred by unobserved factors approximated by the hidden

states (Fig. 2; Table S2). This is not to say that salinity has not played an important role in
diatom diversification, but the opposite—that its effect is strong enough to be detected despite
broad variation related to other, unobserved factors.
It is worth considering what some of these unmeasured factors might be. For example,
differences between lotic and lentic habitats, and within the latter, differences in age, size,
depth, and longevity among lakes of glacial, tectonic, and volcanic origin might be driving
some of the variation within freshwaters. Likewise, the drastic contrast in temperature,
nutrients, and light conditions experienced by diatom communities in coastal areas vs. nutrient-
rich marine polar waters or oligotrophic oceanic gyres could account for some of the hidden
state components within marine lineages. Water temperature in particular appears to be an
important but untested covariate, as it changes with latitude, geography (coastal vs. pelagic),
and water depth (linked to our plankton–benthos contrast), and is related to metabolic rates that
one could easily envision influencing diversification dynamics. For example, Crampton et al.
(2016) found that peaks in species turnover of marine Antarctic diatoms coincided with cooling
of southern latitudes and expansion of ground and sea ice. Five such pulses above the baseline
turnover have been inferred over the past 15 My (Crampton et al. 2016), contributing to
relatively fast turnover within Antarctic diatom communities. Similar cooling-driven
fluctuations in turnover are less likely to have occurred in tropical ocean regions where
temperature has been relatively constant over the past 15 My (Powell and Glazier 2017).
Therefore, it is possible that a portion of the variation captured by the hidden states might also
be related to temperature and latitude. In this regard, diatoms are among the few lineages with
a reverse latitudinal diversity gradient, which is thought to be a product of asymmetric range
expansion toward higher latitudes (Powell and Glazier 2017).
Variation in rates of diversification within regimes might also stem from other traits that
covary with diversity patterns in diatoms. For example, motility of vegetative cells, which
allowed colonization of new types of habitats and fundamentally changed how these diatoms
interact with each other and the environment, was also associated with a faster rate of
diversification (Nakov et al. 2018). A comparison between motile and non-motile lineages
showed that the probability of occupying the second hidden state—where net turnover was
much faster (Fig. 3, Table S2)—was higher in motile lineages, though this result was not
significant (phylogenetic ANOVA, p=0.4). It is important to reiterate that this *post hoc*
exploration of potential hidden variables is not meant to draw conclusions about other factors
not directly tested here, but only as an illustration of the many other variables that may have
interacted with diatom diversification (e.g., Lewitus et al. 2018).
493
Diatom ancestry, absorbing habitats, and phylogenetic diversity
Ancestral state reconstructions supported a marine planktonic environment for the most recent
common ancestor of diatoms, but they quickly colonized the marine benthos (see also Kooistra
et al. 2007). The oldest extant benthic lineages were estimated to be about 180–175 My old (Fig.
3). Colonizations of freshwater did not occur until much later—the oldest extant lineages to gain
a foothold in the freshwater plankton date back to 120–130 Mya, some 60 My after the origin of
diatoms (Fig. 3). Apart from a few isolated cases that have not left a substantial mark on present

day diversity (e.g., *Ellerbeckia*, *Terpsinoe*, and *Pleurosira*), diatoms did not diversify greatly within
the freshwater benthos until after the evolution of anisogamous sexual reproduction and axial
cell symmetry in the pennate diatom lineage (Fig. 3). Marine benthic ancestry was inferred for
the MRCA of the pennate clade, and within it, the clade of actively motile raphid diatoms (Fig.
3). Overall, although our reconstructions supported a planktonic ancestry for diatoms, it is clear
that the marine benthos has been the center stage of diatom evolution, seeding diversity across
a range of environments (e.g., the marine plankton and freshwater benthos) throughout their
history (Fig 3; see also Kooistra et al. 2007).
The permeability of the salinity barrier is a recurring topic in diatom ecology and
evolution. Historically, both in diatoms, and also more broadly across protists (Mann 1999;
Cavalier-Smith 2009; Logares et al. 2009), marine-to-freshwater colonizations have been thought
of as difficult and irreversible transitions, which led to the development of the 'Rubicon'
hypothesis, which states that once a lineage colonizes freshwaters there is no return to the
marine environment (Mann 1999). By simultaneously estimating diversification and transition
rates, our models provided a powerful test of this hypothesis, and by accounting for rate
heterogeneity within regimes we could test whether salinity is a Rubicon in all, or just some,
diatom lineages. Models that disallowed freshwater-to-marine or plankton-to-benthos
transitions as required to directly test these hypotheses received no support (Table 1, 4).
However, our analyses showed that the rates of traversal of either environmental contrasts were
dependent on hidden factors. For example, in some lineages, freshwater-to-marine
colonizations were rare and marine-to-freshwater far more common, whereas in others the
waiting time to recolonization of marine habitats was 13 times shorter than the waiting time of
marine-to-freshwater transitions (Table 2). Models that accounted for the possibility that
marine-to-freshwater transitions differ between planktonic and benthic species, or that
plankton-to-benthos migration differs between marine and freshwater lineages, showed similar
variation (MuHiSSE analysis, Fig. 2, Table S2). Therefore, although the notion of irreversibility
of freshwater or plankton colonizations is not supported for the diatom lineage as a whole, it
nonetheless might apply to a subset of diatom lineages in which recolonization of ancestral
habitats, although still possible, takes far longer (Fig. 2, Table 2, 3).
Comparison of the transition rates can also be informative about the phylogenetic
diversity of communities occupying the extremes of these environmental contrasts. This can be
illustrated by comparing the rates of outflow from the freshwater plankton between the first
and second hidden state. For example, in some lineages escaping the freshwater plankton was a
rare event, but in others, outflow from the freshwater plankton was among the most common
transitions with the shortest retention time (Fig. 2). Such contrasts in transition rates suggest
that for some lineages the freshwater plankton is a transient state, which, combined with the
generally short-lived nature of freshwater planktonic habitats over deep geologic timescales,
suggests an explanation regarding the generally narrow phylogenetic breadth of freshwater
planktonic diatom communities. The few lineages that have diversified substantially in the
freshwater plankton are almost entirely restricted to this habitat and have mostly been sourced
from the marine plankton (e.g., the genus *Aulacoseira* within marine planktonic *Melosirales* or
cyclostephanoid diatoms within the ancestrally marine planktonic *Thalassiosirales*). In turn, for

most of the remaining lineages, colonizations of freshwater plankton were sporadic and did not
lead to substantial species diversification (Fig. 3). This is in contrast to the much more
phylogenetically diverse marine plankton, which has been colonized repeatedly by many
marine benthic lineages (Fig 3; see also Kooistra et al. 2007), harbors a considerable diversity of
araphid (e.g., *Asterionellopsis* and *Thalassiosinema*) and raphid diatoms (e.g., *Fragilariopsis* and
Pseudo-nitzschia; Fig. 3), and as suggested by the rates of outflow from the freshwater plankton,
is also recolonized by freshwater planktonic lineages (Fig. 2, Table S2). These observations,
combined with the non-uniform recruitment to the freshwater plankton—several old and
species-rich marine planktonic lineages have not crossed into freshwaters or remain
disproportionately more diverse in saline waters (e.g., Rhizosoleniales, Coscinodiscales, and
Chaetocerales; Fig. 3)—also suggest that marine planktonic diatom assemblages are
phylogenetically and functionally more diverse than their freshwater counterparts.
It is important to consider here that transitions between marine and freshwaters, or
plankton and benthos, certainly do not occur instantaneously. Rather, the most likely colonizers
of freshwater are marine lineages already capable of growing at a lower salinity (brackish) or a
broader range of salinity (euryhaline), for example species living in estuaries or saline inland
lakes. A more appropriate model for these analyses would therefore account for the presence of
euryhaline or brackish species and potentially constrain the transitions between the salinity
extremes by forcing an intermediate euryhaline or brackish state. In fact, an analysis of a small
clade of predominantly benthic diatoms did support a model of ordered, reversible marine<-
>euryhaline<->freshwater transitions, although, it did not account for diversification (Ruck et
al. 2016). Replicating such an analysis at the level of all diatoms and in a SSE modeling
framework would certainly be a welcome approach. Using geographic SSE models (GeoSSE,
Goldberg et al. 2011; or HiGeoSSE, Caetano et al. 2018) where species with euryhaline ranges
diverge into freshwater and marine populations is also an interesting way to model these data
(see e.g., Betancur-R et al. 2015). However, implementation of such approaches is hampered at
this time due to the lack of data for the range of salinity tolerance across the diatom tree, the
patchy information on diatoms in biogeographic databases, and the resulting uncertainties in
estimation of total diversity and sampling fractions for euryhaline or brackish species.
In sum, the marine–freshwater and plankton-benthos contrasts are unquestionably
important determinants of aquatic diversity. In diatoms, a species-rich lineage of algae
important for global ecology and biogeochemistry, we found that the marine–freshwater
diversity imbalance is maintained by faster turnover and faster net diversification in
freshwaters. This pattern holds for both planktonic and benthic habitats, an environmental
divide that itself has not greatly impacted diatom diversification. More generally, our analyses
showed that global diversity patterns are unlikely to be attributable to a few, broadly defined
regimes. Rather they are the product of many potentially interacting traits and environmental
factors whose individual or aggregate effects might equal or surpass the influence of factors
traditionally considered important. Our analyses help illustrate how consideration of multiple
possibly interacting characters provides a more refined understanding of diversification in old,
widespread, and species-rich clades.
584

References

Alverson, A. J. 2014. Timing marine–freshwater transitions in the diatom order Thalassiosirales.

Paleobiology 40:91–101.

Alverson, A. J., R. K. Jansen, and E. C. Theriot. 2007. Bridging the Rubicon: Phylogenetic

analysis reveals repeated colonizations of marine and fresh waters by thalassiosiroid

diatoms. Mol. Phylogenet. Evol. 45:193–210.

Appeltans, W., S. T. Ahyong, G. Anderson, M. V. Angel, T. Artois, N. Bailly, R. Bamber, A.

Barber, I. Bartsch, A. Berta, M. Błażewicz-Paszkowycz, P. Bock, G. Boxshall, C. B. Boyko, S.

N. Brandão, R. A. Bray, N. L. Bruce, S. D. Cairns, T.-Y. Chan, L. Cheng, A. G. Collins, T.

Cribb, M. Curini-Galletti, F. Dahdouh-Guebas, P. J. F. Davie, M. N. Dawson, O. De Clerck,

W. Decock, S. De Grave, N. J. de Voogd, D. P. Domning, C. C. Emig, C. Erséus, W.

Eschmeyer, K. Fauchald, D. G. Fautin, S. W. Feist, C. H. J. M. Fransen, H. Furuya, O.

Garcia-Alvarez, S. Gerken, D. Gibson, A. Gittenberger, S. Gofas, L. Gómez-Daglio, D. P.

Gordon, M. D. Guiry, F. Hernandez, B. W. Hoeksema, R. R. Hopcroft, D. Jaume, P. Kirk, N.

Koedam, S. Koenemann, J. B. Kolb, R. M. Kristensen, A. Kroh, G. Lambert, D. B. Lazarus, R.

Lemaitre, M. Longshaw, J. Lowry, E. Macpherson, L. P. Madin, C. Mah, G. Mapstone, P. A.

McLaughlin, J. Mees, K. Meland, C. G. Messing, C. E. Mills, T. N. Molodtsova, R. Mooi, B.

Neuhaus, P. K. L. Ng, C. Nielsen, J. Norenburg, D. M. Opresko, M. Osawa, G. Paulay, W.

Perrin, J. F. Pilger, G. C. B. Poore, P. Pugh, G. B. Read, J. D. Reimer, M. Rius, R. M. Rocha, J.

I. Saiz-Salinas, V. Scarabino, B. Schierwater, A. Schmidt-Rhaesa, K. E. Schnabel, M. Schotte,

P. Schuchert, E. Schwabe, H. Segers, C. Self-Sullivan, N. Shenkar, V. Siegel, W. Sterrer, S.

Stöhr, B. Swalla, M. L. Tasker, E. V. Thuesen, T. Timm, M. A. Todaro, X. Turon, S. Tyler, P.

Uetz, J. van der Land, B. Vanhoorne, L. P. van Ofwegen, R. W. M. van Soest, J.

Vanaverbeke, G. Walker-Smith, T. C. Walter, A. Warren, G. C. Williams, S. P. Wilson, and

M. J. Costello. 2012. The magnitude of global marine species diversity. Curr. Biol. 22:2189–

2202.

Armbrust, E. V. 2009. The life of diatoms in the world's oceans. Nature 459:185–192.

Beaulieu, J. M., and B. C. O'Meara. 2018. Can we build it? Yes we can, but should we use it?

Assessing the quality and value of a very large phylogeny of campanulid angiosperms.

Am. J. Bot. 105:417–432.

Beaulieu, J. M., and B. C. O'Meara. 2016. Detecting hidden diversification shifts in models of

trait-dependent speciation and extinction. Syst. Biol. 65:583–601.

Betancur-R, R., G. Ortí, and R. A. Pyron. 2015. Fossil-based comparative analyses reveal ancient

marine ancestry erased by extinction in ray-finned fishes. Ecol. Lett. 18:441–450.

Bloom, D. D., J. T. Weir, K. R. Piller, and N. R. Lovejoy. 2013. Do freshwater fishes diversify

faster than marine fishes? A test using state-dependent diversification analyses and

molecular phylogenetics of new world silversides (Atherinopsidae). Evolution 67:2040–

2057. Wiley Online Library.

Caetano, D. S., B. C. O'Meara, and J. M. Beaulieu. 2018. Hidden state models improve state-

dependent diversification approaches, including biogeographical models. Evolution

72:2308–2324. Wiley Online Library.

Casteleyn, G., F. Leliaert, T. Backeljau, A.-E. Debeer, Y. Kotaki, L. Rhodes, N. Lundholm, K.

Sabbe, and W. Vyverman. 2010. Limits to gene flow in a cosmopolitan marine planktonic

diatom. Proc. Natl. Acad. Sci. U. S. A. 107:12952–12957.

Cavalier-Smith, T. 2009. Megaphylogeny, cell body plans, adaptive zones: causes and timing of

eukaryote basal radiations. J. Eukaryot. Microbiol. 56:26–33.

Cohen, A. S., J. R. Stone, K. R. M. Beuning, L. E. Park, P. N. Reinthal, D. Dettman, C. A. Scholz,

T. C. Johnson, J. W. King, M. R. Talbot, E. T. Brown, and S. J. Ivory. 2007. Ecological

consequences of early Late Pleistocene megadroughts in tropical Africa. Proc. Natl. Acad.

Sci. U. S. A. 104:16422–16427.

Crampton, J. S., R. D. Cody, R. Levy, D. Harwood, R. McKay, and T. R. Naish. 2016. Southern

Ocean phytoplankton turnover in response to stepwise Antarctic cooling over the past 15

million years. Proceedings of the National Academy of Sciences 113:6868–6873.

Edlund, M. B., and E. F. Stoermer. 2000. A 200,000-year, high-resolution record of diatom

productivity and community makeup from Lake Baikal shows high correspondence to the

marine oxygen-isotope record of climate change. Limnol. Oceanogr. 45:948–962. Wiley

Online Library.

Evans, K. M., V. A. Chepurnov, H. J. Sluiman, S. J. Thomas, B. M. Spears, and D. G. Mann. 2009.

Highly differentiated populations of the freshwater diatom Sellaphora capitata suggest

limited dispersal and opportunities for allopatric speciation. Protist 160:386–396.

FitzJohn, R. G., W. P. Maddison, and S. P. Otto. 2009. Estimating trait-dependent speciation and

extinction rates from incompletely resolved phylogenies. Syst. Biol. 58:595–611.

Godhe, A., and K. Härnström. 2010. Linking the planktonic and benthic habitat: genetic

structure of the marine diatom Skeletonema marinoi. Mol. Ecol. 19:4478–4490.

Goldberg, E. E., L. T. Lancaster, and R. H. Ree. 2011. Phylogenetic inference of reciprocal effects

between geographic range evolution and diversification. Syst. Biol. 60:451–465.

Grosberg, R. K., G. J. Vermeij, and P. C. Wainwright. 2012. Biodiversity in water and on land.

Curr. Biol. 22:R900–3.

Guinot, G., and L. Cavin. 2015. Contrasting "Fish" Diversity Dynamics between Marine and

Freshwater Environments. Curr. Biol. 25:2314–2318.

Hampton, S. E., S. McGowan, T. Ozersky, S. G. P. Virdis, T. T. Vu, T. L. Spanbauer, B. M.

Kraemer, G. Swann, A. W. Mackay, S. M. Powers, M. F. Meyer, S. G. Labou, C. M. O'Reilly,

M. DiCarlo, A. W. E. Galloway, and S. C. Fritz. 2018. Recent ecological change in ancient

lakes. Limnol. Oceanogr. 37:17.

Hasle, G. R., and E. E. Syvertsen. 1996. Marine diatoms. Identifying Marine Phytoplankton

(Tomas, CR, ed.), 5-385. Academic Press, San Diego.

Herrera-Alsina, L., P. van Els, and R. S. Etienne. 2019. Detecting the Dependence of

Diversification on Multiple Traits from Phylogenetic Trees and Trait Data. Syst. Biol.

68:317–328.

Kirst, G. O. 1996. Osmotic Adjustment in Phytoplankton and MacroAlgae. Pp. 121–129 *in* R. P.

Kiene, P. T. Visscher, M. D. Keller, and G. O. Kirst, eds. Biological and Environmental

Chemistry of DMSP and Related Sulfonium Compounds. Springer US, Boston, MA.

Kirst, G. O. 1990. Salinity Tolerance of Eukaryotic Marine Algae. Annu. Rev. Plant Physiol.

Plant Mol. Biol. 41:21–53. Annual Reviews.

Kociolek, J. P., K. Balasubramanian, S. Blanco, M. Coste, L. Ector, Y. Liu, M. Kulikovskiy, N.

Lundholm, T. Ludwig, M. Potapova, F. Rimet, K. Sabbe, S. Sala, E. Sar, J. Taylor, B. Van de

Vijver, C. E. Wetzel, D. M. Williams, A. Witkowski, and J. Witkowski. 2017. DiatomBase.

Kooistra, W. H. C. F., G. Forlani, and M. De Stefano. 2009. Adaptations of araphid pennate

diatoms to a planktonic existence. Mar. Ecol. 30:1–15.

Kooistra, W. H. C. F., R. Gersonde, L. K. Medlin, and D. G. Mann. 2007. CHAPTER 11 - The

Origin and Evolution of the Diatoms: Their Adaptation to a Planktonic Existence. Pp. 207–

249 *in* P. G. Falkowski and A. H. Knoll, eds. Evolution of Primary Producers in the Sea.

Academic Press, Burlington.

Lange-Bertalot, H., and K. Krammer. 2000. Diatoms of Europe: Diatoms of the European inland

waters and comparable habitats. ARG Ganter.

Lewitus, E., L. Bittner, S. Malviya, C. Bowler, and H. Morlon. 2018. Clade-specific diversification

dynamics of marine diatoms since the Jurassic. Nature Ecology & Evolution 2:1715. Nature

Publishing Group.

Lobban, C. S., M. Schefter, R. W. Jordan, Y. Arai, A. Sasaki, E. C. Theriot, M. Ashworth, E. C.

Ruck, and C. Pennesi. 2012. Coral-reef diatoms (Bacillariophyta) from Guam: new records

and preliminary checklist, with emphasis on epiphytic species from farmer-fish territories.

Micronesica 43:237–479.

Logares, R., J. Bråte, S. Bertilsson, J. L. Clasen, K. Shalchian-Tabrizi, and K. Rengefors. 2009.

Infrequent marine–freshwater transitions in the microbial world. Trends Microbiol. 17:414–

422.

Mackay, A. W., M. B. Edlund, G. Khursevich, and Others. 2010. Diatoms in ancient lakes. The

Diatoms: Applications for the environmental and earth sciences 2:209–228. Cambridge

University Press: Cambridge.

Maddison, W. P. 2006. Confounding asymmetries in evolutionary diversification and character

change. Evolution 60:1743–1746.

Maddison, W. P., P. E. Midford, and S. P. Otto. 2007. Estimating a binary character's effect on

speciation and extinction. Syst. Biol. 56:701–710.

Mann, D. G. 1999. Crossing the Rubicon: the effectiveness of the marine/freshwater interface as

a barrier to the migration of diatom germplasm. Pp. 1–21 *in* Proceedings of the 14th

International Diatom Symposium, Koenigstein, 1999. Koeltz Scientific Books.

Mann, D. G., and P. Vanormelingen. 2013. An inordinate fondness? The number, distributions,

and origins of diatom species. J. Eukaryot. Microbiol. 60:414–420.

Miller, E. C., and J. J. Wiens. 2017. Extinction and time help drive the marine-terrestrial

biodiversity gradient: Is the ocean a deathtrap? Ecol. Lett. 20:911–921. Wiley Online

Library.

Mora, C., D. P. Tittensor, S. Adl, A. G. B. Simpson, and B. Worm. 2011. How Many Species Are

There on Earth and in the Ocean? PLoS Biol. 9:e1001127. Public Library of Science.

Nakov, T., M. Ashworth, and E. C. Theriot. 2015. Comparative analysis of the interaction

between habitat and growth form in diatoms. ISME J. 9:246–255.

Nakov, T., J. M. Beaulieu, and A. J. Alverson. 2018. Accelerated diversification is related to life

history and locomotion in a hyperdiverse lineage of microbial eukaryotes (Diatoms,

Bacillariophyta). New Phytol. 219:462–473.

Palumbi, S. R. 1994. Genetic Divergence, Reproductive Isolation, and Marine Speciation. Annu.

Rev. Ecol. Syst. 25:547–572. Annual Reviews.

Palumbi, S. R. 1992. Marine speciation on a small planet. Trends Ecol. Evol. 7:114–118. Elsevier.

Powell, M. G., and D. S. Glazier. 2017. Asymmetric geographic range expansion explains the

latitudinal diversity gradients of four major taxa of marine plankton. Paleobiology 43:196–

208. Cambridge University Press.

Rabosky, D. L. 2019. Phylogenies and Diversification Rates: Variance Cannot Be Ignored.

Round, F. E., R. M. Crawford, and D. G. Mann. 1990. Diatoms: Biology and Morphology of the

Genera. Cambridge University Press.

Round, F. E., and P. A. Sims. 1980. The distribution of diatom genera in marine and freshwater

environments and some evolutionary considerations. Pp. 301–320 *in* Proceedings of the

sixth symposium on recent and fossil diatoms. Otto Koeltz Science Publishers Hirschberg,

Germany.

Ruck, E. C., T. Nakov, A. J. Alverson, and E. C. Theriot. 2016. Phylogeny, ecology,

morphological evolution, and reclassification of the diatom orders Surirellales and

Rhopalodiales. Mol. Phylogenet. Evol. 103:155–171.

Rynearson, T. A., and E. V. Armbrust. 2004. Genetic differentiation among populations of the

planktonic marine diatom Ditylum brightwellii (Bacillariophyceae). J. Phycol. 40:34–43.

Wiley Online Library.

Sims, P. A., D. G. Mann, and L. K. Medlin. 2006. Evolution of the diatoms: Insights from fossil,

biological and molecular data. Phycologia 45:361–402.

Sjöqvist, C., A. Godhe, P. R. Jonsson, L. Sundqvist, and A. Kremp. 2015. Local adaptation and

oceanographic connectivity patterns explain genetic differentiation of a marine diatom

across the North Sea--Baltic Sea salinity gradient. Mol. Ecol. 24:2871–2885. Wiley Online

Library.

Stone, J. R., K. S. Westover, and A. S. Cohen. 2011. Late Pleistocene paleohydrography and

diatom paleoecology of the central basin of Lake Malawi, Africa. Palaeogeogr.

Palaeoclimatol. Palaeoecol. 303:51–70.

Vanormelingen, P., V. A. Chepurnov, D. G. Mann, K. Sabbe, and W. Vyverman. 2008. Genetic

divergence and reproductive barriers among morphologically heterogeneous sympatric

clones of Eunotia bilunaris sensu lato (Bacillariophyta). Protist 159:73–90.

Vermeij, G. J., R. K. Grosberg, C. R. Marshall, and R. Motani. 2018. The sea as deathtrap:

comment on a paper by Miller and Wiens. Ecol. Lett. 21:938–939.

Whittaker, K. A., and T. A. Rynearson. 2017. Evidence for environmental and ecological

selection in a microbe with no geographic limits to gene flow. Proc. Natl. Acad. Sci. U. S. A.

114:2651–2656.

Wiens, J. J. 2015. Faster diversification on land than sea helps explain global biodiversity

patterns among habitats and animal phyla. Ecol. Lett. 18:1234–1241.

Witkowski, A., H. Lange-Bertalot, D. Metzeltin, H. Lange-Bertalot, and K. Gantner Verlag. 2000.

Diatom flora of marine coasts I. Iconographia Diatomologica. Annotated Diatom

Micrographs. Diversity-Taxonomy-Identiflcation. Vol. 7. ARG Ganter Verlag KG, Ruggell.

Figures
756
Figure 1. Distributions of tip estimates of net turnover (τ, speciation + extinction) and net
diversification (r, speciation − extinction) averaged across hidden states. (A) binary coding
for marine vs. freshwater (HiSSE model), (B) binary coding for planktonic vs. benthic (CID4
model) and (C) multistate coding for the combination of salinity and habitat [averaged across
four MuHiSSE variants (Table 1)]. The circles and error bars show the mean ± standard
deviation across tips of the phylogeny and the carpet points show the individual rates. The
rates are inverted into waiting times (wt) in millions of years. Abbreviations: M, marine; F,
freshwater; P, plankton; B, benthos.
765

Figure 2. Waiting times (inverse of the rates) in million years (My) for speciation, extinction, and transitions for the habitat + salinity under the favored MuHiSSE model (Table 4). The source environment is denoted with a two-letter abbreviation on the y axis. For speciation (λ) and extinction (μ) the color of the end-point circle matches the source environment. For transition rates (q), the color and abbreviation at the end-point circle denote the destination of colonization. Numbers along the lines are the estimated waiting times. Transitions rates estimated as indistinguishable from zero (and, consequently, infinity waiting times) are shown with dashed lines and open circles. Left and right side panels correspond to the two hidden states. Abbreviations: FB, freshwater--benthos; FP, freshwater--plankton; MB, marine--benthos; MP, marine--plankton.

Figure 3. Ancestral state reconstructions of habitats and tip estimates of net turnover (τ,
speciation + extinction) and net diversification (r, speciation - extinction) rates inverted to
waiting times (wt) in millions of years. The high-turnover and high-diversification
freshwater lineages (shorter waiting times) are predominantly benthic within the raphid
clade with motile vegetative cells (e.g., Suri and Cymb) and predominantly planktonic
within the non-motile centric and araphid grades (e.g., Frag, Thal, and Melo). Abbreviations:
Mya, millions of years ago; Melo, Melosirales; Chae, Chaetocerotales; Eupo, Eupodiscales;
Thal, Thalassiosirales; Frag, fragilaroids; Nitz, Bacillariales; Navi, naviculoids; Pinn,
pinnulariids; Cymb, Cymbellales; Suri, Surirellales; Parm, Parmales; Lept, Leptocylindrales
+ Corethrales; Cosc, Coscinodiscales + Rhizosoleniales.
794

795
796
797

798
Tables
Table 1. Comparison of character-dependent and character-independent models for the
binary coding of salinity and habitat. The models favored by AICc are in bold.
Abbreviations: npd, number of diversification parameters (turnover + extinction fraction);
npt, number of transition parameters; *Δ*, difference in AICc; ω, Akaike weight; F, freshwater;
P, plankton.
805
806
807

Model	lnL	npd	npt	AICc	*Δ*	ω
Salinity (Marine vs. Freshwater)						
hisse	-5325.72	**5**	**8**	10677.77	0	**0.999**
cid4	-5341.02	5	3	10698.16	20.39	<0.001
hisse F absorbing	-5393.10	5	6	10808.44	130.67	<0.001
cid2	-5444.22	3	3	10900.52	222.75	<0.001
bisse	-5506.97	3	2	11023.99	346.22	<0.001
Habitat (Plankton vs. Benthos)						
hisse	-5164.73	**5**	**8**	10355.78	0	**0.999**
cid4 (three rates)	-5189.49	5	3	10395.11	39.33	<0.001
hisse P absorbing	-5189.70	5	6	10401.63	45.85	<0.001
cid2	-5250.83	3	3	10513.74	157.96	<0.001
bisse	-5377.10	3	2	10764.25	408.46	<0.001

808
809

810 **Table 2. Parameter estimates (events per million years) and 2 log-likelihood unit confidence**
811 **interval (in brackets) for the best model (hidden-state speciation and extinction, HiSSE) for**
812 **salinity (marine vs. freshwater). The letters next to character states indicate the hidden states.**
813

	Marine A	Freshwater A	Marine B	Freshwater B
Transition rates				
Marine A	-	0.0150 [0.0141-0.0166]	0.0459 [0.0424-0.0509]	0
Freshwater A	0 [0.0-0.0009]	-	0	0.0187 [0.0142-0.0203]
Marine B	0.0029 [0.0026-0.0033]	0	-	0 [0.0-0.0005]
Freshwater B	0	0.1814 [0.162-0.196]	0.0067 [0.0050-0.0068]	-
Diversification rates				
Speciation	0.168 [0.165-0.183]	0.059 [0.056-0.065]	0.035 [0.035-0.04]	0.472 [0.445-0.52]
Extinction	0.082 [0.08-0.091]	0.029 [0.028-0.033]	0.017 [0.017-0.02]	0.232 [0.221-0.262]
Net turnover (speciation + extinction)	0.25 [0.247-0.274]	0.089 [0.085-0.098]	0.053 [0.053-0.06]	0.704 [0.667-0.772]
Net diversification (speciation - extinction)	0.085 [0.082-0.092]	0.03 [0.027-0.034]	0.018 [0.017-0.02]	0.241 [0.222-0.268]
Extinction fraction (extinction/speciation)	0.49 [0.47-0.519]			

814
815

816 **Table 3. Parameter estimates (events per million years) and 2 log-likelihood unit confidence**
817 **interval (in brackets) for the best model (hidden-state speciation and extinction, HiSSE) for**
818 **habitat (plankton vs. benthos). The letters next to character states indicate the hidden states.**
819

	Plankton A	Benthos A	Plankton B	Benthos B
Transition rates				
Plankton A	-	0.0017 [0.0015-0.0019]	0.0027 [0.0012-0.0051]	0
Benthos A	0.0003 [0.0002-0.0007]	-	0	0.0022 [0.002-0.0023]
Plankton B	0.0955 [0.0842-0.0981]	0	-	0.007 [0.0065-0.0073]
Benthos B	0	0.0497 [0.049-0.0608]	0.0013 [0.0011-0.0016]	-
Diversification rates				
Speciation	0.078 [0.065-0.087]	0.082 [0.075-0.095]	0.581 [0.521-0.608]	0.481 [0.449-0.515]
Extinction	0.064 [0.053-0.072]	0.068 [0.061-0.079]	0.479 [0.418-0.505]	0.396 [0.36-0.423]
Net turnover (speciation + extinction)	0.143 [0.118-0.159]	0.15 [0.136-0.173]	1.06 [0.939-1.112]	0.878 [0.809-0.938]
Net diversification (speciation - extinction)	0.014 [0.011-0.016]	0.015 [0.013-0.019]	0.103 [0.092-0.116]	0.085 [0.081-0.097]
Extinction fraction (extinction/speciation)	0.824 [0.8-0.836]			

820
821
822
823

Table 4. Comparison of character-dependent and independent models for the interaction
between diversification and the combined salinity+habitat states. Shown are the models with
support ≥ 1% used in model-averaging. For the full set of results refer to Supplementary
table S1 Abbreviations: npd, number of diversification parameters (turnover and extinction
fraction); npt, number of transition parameters; Δ, difference in AICc; ω, Akaike weight; ε,
extinction fraction.
830

Model	Parameter setup	lnL	npd	npt	AICc	Δ	ω
No habitat effect	ε=fix; MP = MB *and* FP = FB	-5518.47	7	18	11083.93	0	0.84
MuHiSSE full	ε=fix; no additional constraints	-5516.44	9	18	11088.25	4.32	0.10
MP absorbing plankton	ε=fix; MP -> MB = 0, but MP -> FP ≠ 0	-5519.25	9	17	11089.68	5.75	0.05
FP absorbing freshwater	ε=fix; FP -> MP = 0, but FP -> FB ≠ 0	-5520.47	9	17	11092.12	8.19	0.01

831
832
833

1 **Supplementary information**
2
3 **Clade-specific sampling biases**
4
Clade-specific sampling biases could arise as a result of deeper phylogenetic sampling in
lineages of ecological, biotechnological, or public health interest or due to natural variation in
the distribution of lineages across the focal environmental contrasts. One way to assess potential
biases of this type is to check whether a pattern observed across the entire phylogeny is
replicated at shallower phylogenetic scales. There is no reason to expect that the same models
should be favored or similar parameters should apply for subtrees that differ substantially in
age, ecology, life history, genomic features, and so on (Beaulieu and O'Meara 2018; Rabosky
2019). But it might be informative to evaluate in broad terms whether a global result, i.e.,
turnover in freshwater lineages is faster than in marine, is detectable, at least to some degree, in
younger, smaller, clades.
To address this, we analyzed four subtrees with sufficient representation across both the
marine–freshwater and plankton–benthos contrasts (Fig. S2) and asked whether the model
favored for the all-diatom tree was also favored in the subtrees and whether model-averaged
tip-associated turnover rates estimated for the subtrees recovered a pattern similar to that of the
all-diatom phylogeny. For these smaller clades, the CID4 model was favored over HiSSE in all
but one clade, for both salinity and habitat. However, even though a model of character-
dependent diversification was not universally supported, model-averaged tip rates showed
that, on average, freshwater species had higher turnover than marine in all but one clade. For
habitat, in some clades turnover in the plankton was faster, on average, than in the benthos, but
in others, the two habitats had essentially the same mean rates, consistent with the pattern
observed at the all-diatom scale. For the two largest clades, raphid and multipolar diatoms, we
also repeated the combined habitat+salinity analysis and found that model selection was
replicated for the clade of raphid diatoms but not for the clade of multipolar lineages. Model-
averaged rates again resembled the pattern observed for the whole diatom tree, with turnover
faster in freshwater than marine environments.

Table S1. The full set of multistate models examined for the combined salinity (marine–
freshwater) + habitat (plankton–benthos) character. In all cases the extinction fraction (ε,
speciation/extinction) was held constant across states, while turnover (τ, speciation +
extinction) was free to vary across states or constrained to test specific hypotheses.
34

Model	Parameter setup	lnL	AICc	dAIC	AICw
No habitat effect	MP = MB, FP = FB	-5518.47	11083.93	0	0.84
MuHiSSE relax	No additional constraints	-5516.44	11088.25	4.32	0.1
MP absorbing plankton	MP -> MB = 0, MP -> FP ≠ 0	-5519.25	11089.68	5.75	0.05
FP absorbing freshwater	FP -> MP = 0, FP -> FB ≠ 0	-5520.47	11092.12	8.19	0.01
No salinity effect in plankton	MP = FP, MB ≠ FB	-5524.71	11100.6	16.67	0
FP absorbing plankton	FP -> MP ≠ 0, FP -> FB = 0	-5527.68	11106.53	22.6	0
No habitat effect in freshwater	MP ≠ MB, FP = FB	-5530.05	11111.28	27.35	0
No habitat effect in marine	MP = MB, FP ≠ FB	-5530.4	11111.98	28.05	0
No salinity effect in benthos	MP ≠ FP, MB = FB	-5533.29	11117.75	33.82	0
No salinity effect	MP = FP, MB = FB	-5536.56	11120.11	36.18	0
MuHiSSE	Symmetric transitions between hidden states	-5539.58	11132.43	48.5	0
Plankton absorbing	FP -> FB = 0, MP -> MB = 0	-5542.91	11132.81	48.88	0
CID4 relax	No additional constraints	-5563.63	11161.81	77.88	0
CID5 relax	No additional constraints	-5562.7	11164.08	80.15	0
CID3 relax	No additional constraints	-5566.86	11164.15	80.22	0
CID6 relax	No additional constraints	-5562.69	11168.22	84.29	0
CID2 relax	No additional constraints	-5572.27	11170.86	86.93	0
FB absorbing freshwater	FB -> MB = 0, FB -> FP ≠ 0	-5560.22	11171.61	87.68	0
CID7 relax	No additional constraints	-5562.68	11172.36	88.43	0
CID8 relax	No additional constraints	-5562.63	11176.43	92.5	0
Freshwater absorbing	FB -> MB = 0, FP -> MP = 0	-5594.11	11235.21	151.28	0
MuSSE	No additional constraints	-5701.85	11430.03	346.1	0
Dull	No additional constraints	-5903.94	11828.08	744.15	0

35
36
37

38   **Table S2. Parameter estimates and 2 log likelihood support region (in brackets) for the**
39   **combined salinity + habitat character states [marine–plankton (MP), marine–benthos (MB),**
40   **freshwater–plankton (FP) and freshwater–benthos (FB)]. The favored model (ω = 0.84) was**
41   **hidden state multistate speciation and extinction without a habitat effect on diversification**
42   **(Table 4). The letters next to character states indicate the hidden states.**
43

	MPA	MBA	FPA	FBA	MPB	MBB	FPB	FBB
Diversification rates								
Speciation	0.059 [0.058-0.061]		0.105 [0.098-0.111]		0.376 [0.375-0.385]		0.648 [0.626-0.654]	
Extinction	0.048 [0.046-0.05]		0.085 [0.078-0.09]		0.304 [0.298-0.315]		0.523 [0.498-0.535]	
Net turn	0.107 [0.104-0.11]		0.189 [0.176-0.201]		0.68 [0.674-0.7]		1.17 [1.123-1.189]	
Net div	0.011 [0.012-0.011]		0.02 [0.02-0.02]		0.073 [0.077-0.07]		0.125 [0.128-0.12]	
Extinct frac	0.807 [0.795-0.817]							
Transition rates								
	FB1-to-FP1		0.00111 [0.00109-0.00113]		MP1-to-FP1		0 [0-0.00048]	
	FB1-to-MB1		0.00393 [0.0037-0.00399]		MP1-to-MB1		0 [0-0.00065]	
	FB2-to-FP2		0.00047 [0.00045-9e-04]		MP2-to-FP2		0.005 [0.00493-0.00507]	
	FB2-to-MB2		0.00229 [0.00225-0.0023]		MP2-to-MB2		0.00289 [0.00282-0.0029]	
	FP1-to-FB1		0 [0-9e-04]		MB1-to-FB1		0.00155 [0.00153-0.00158]	
	FP1-to-MP1		0 [0-0.00085]		MB1-to-MP1		0.00078 [0.00069-0.00101]	
	FP2-to-FB2		0.01434 [0.01373-0.01465]		MB2-to-FB2		0.01166 [0.0114-0.01192]	
	FP2-to-MP2		0.00868 [0.00852-0.00891]		MB2-to-MP2		0.0033 [0.00318-0.00332]	
	Hidden1-to-2		0.00305 [0.0029-0.00319]					
	Hidden2-to-1		0.06425 [0.06298-0.06564]					

44
45

Figure S1. Character-dependent and character-independent diversification models for binary (A) and multistate (B) coding of salinity and habitat. The top part (grey) depicts the transition rate matrix with row and column names representing observed character states (M=marine, F=freshwater and their combinations) or hidden (H) states. The letters A-H (e.g., MA, MB, FA, FB) differentiate hidden states within a character state. The bottom part (green), labeled with τ and ε shows the parameters for net turnover (speciation+extinction) and extinction fraction (extinction/speciation), respectively. Each τ and ε parameter is linked to the state denoted in the column name. Note that diversification parameters are associated to character states in SSE models, but not in CID models, where they are linked to hidden states. Numbers inside the table represent the indexes of model parameters. For constrained parameters, the same index occurs in multiple cells (e.g., ε was fixed in all models, so it has one parameter in each table). Models for habitat (plankton–benthos) were the same as the binary models for salinity (A).

A. Binary hidden state speciation and extinction

Null

	M	F
M	–	1
F	1	–

τ	2	2
ε	3	3

BiSSE

	M	F
M	–	2
F	1	–

τ	3	4
ε	5	5

HiSSE

	MA	FA	MB	FB
MA	–	3	5	–
FA	1	–	–	7
MB	2	–	–	8
FB	–	4	6	–

τ	9	10	11	12
ε	13	13	13	13

CID2

	M	F	HA	HB
M	–	2		
F	1	–		
HA			–	3
HB			3	–

τ			4	5
ε			6	6

CID4

	M	F	HA	HB	HC	HD
M	–	2				
F	1	–				
HA			–	3	3	3
HB			3	–	3	3
HC			3	3	–	3
HD			3	3	3	–

τ			4	5	6	7
ε			8	8	8	8

B. Multistate hidden state speciation and extinction

Null

	MP	MB	FP	FB
MP	–	3	5	–
MB	1	–	–	7
FP	2	–	–	8
FB	–	4	6	–

τ	9	9	9	9
ε	10	10	10	10

MuSSE

	MP	MB	FP	FB
MP	–	3	5	–
MB	1	–	–	7
FP	2	–	–	8
FB	–	4	6	–

τ	9	10	11	12
ε	13	13	13	13

MuHiSSE

	MPA	MBA	FPA	FBA	MPB	MBB	FPB	FBB
MPA	–	3	5	–	18	–	–	–
MBA	1	–	–	7	–	18	–	–
FPA	2	–	–	8	–	–	18	–
FBA	–	4	6	–	–	–	–	18
MPB	17	–	–	–	–	11	13	–
MBB	–	17	–	–	9	–	–	15
FPB	–	–	17	–	10	–	–	16
FBB	–	–	–	17	–	12	14	–

τ	19	20	21	22	23	24	25	26
ε	27	27	27	27	27	27	27	27

MuCID2

	MP	MB	FP	FB	HA	HB
MP	–	3	5	–		
MB	1	–	–	7		
FP	2	–	–	8		
FB	–	4	6	–		
HA					–	9
HB					10	–

τ					11	12
ε					13	13

MuCID8

	MP	MB	FP	FB	HA	HB	HC	HD	HE	HF	HG	HH
MP	–	3	5	–								
MB	1	–	–	7								
FP	2	–	–	8								
FB	–	4	6	–								
HA –HH					9	10	11	12	13	14	15	16

τ					17	18	19	20	21	22	23	24
ε					25	25	25	25	25	25	25	25

Figure S2. **Tip estimates of net turnover (speciation + extinction) averaged across models and**
hidden states reconstructions. Shown are the clades of that had at least 40 sampled species
and reasonable representation in marine and freshwaters, as well as plankton and benthos.
In addition to the entire diatoms phylogeny (all diatoms), these clades include the raphid
diatoms (raphid), part of araphid diatoms (araphid part), melosiroid diatoms (melosiroid),
and multi-polar diatoms (mediophytes). Top row: binary coding for marine vs. freshwater;
Middle row: binary coding for planktonic vs. benthic; Bottom row: multi-state coding for
combined habitat+salinity states. Abbreviations: M, marine; F, freshwater; P, plankton; B,
benthos.
70

Figure S3. Distributions of tip-estimates of speciation (λ), extinction (μ), net turnover (τ,
speciation + extinction) and net diversification (*r*, speciation - extinction) averaged across
hidden states under the MuHiSSE model and two alternative implementations of the
conditioning for survival of crown lineages (FitzJohn et al. 2009; Herrera-Alsina et al. 2018).
The circles and horizontal error bars show the mean ± standard deviation across tips of the
phylogeny and the carpet points show the individual rates. The rates are inverted into
waiting times (wt) in millions of years. Abbreviations: M, marine; F, freshwater; P, plankton;
B, benthos.
80

Figure S4. Ancestral state reconstructions of environments and the tip estimates of net
turnover (*τ*, speciation + extinction) **and net diversification** (*r*, speciation - extinction)
following Herrera-Alsina et al.'s (2019) implementation of the conditioning for survival of
crown lineages. Rates were inverted to waiting times (wt) in millions of years. The high-
turnover and high-diversification freshwater lineages (lower waiting times) are
predominantly benthic among motile raphid pennate diatoms (e.g., Suri and Cymb) and
predominantly planktonic within the non-motile centric and araphid grades (e.g., Frag, Thal,
and Melo). Abbreviations: Mya, millions of years ago; Melo, Melosirales; Chae,
Chaetocerotales; Eupo, Eupodiscales; Thal, Thalassiosirales; Frag, fragilaroids; Nitz,
Bacillariales; Navi, naviculoids; Pinn, pinnulariids; Cymb, Cymbellales; Suri, Surirellales;
Parm, Parmales; Lept, Leptocylindrales + Corethrales; Cosc, Coscinodiscales +
Rhizosoleniales.
94

95
96
97
98

References

Beaulieu, J. M., and B. C. O'Meara. 2018. Can we build it? Yes we can, but should we use it?

Assessing the quality and value of a very large phylogeny of campanulid angiosperms.

Am. J. Bot. 105:417–432.

FitzJohn, R. G., W. P. Maddison, and S. P. Otto. 2009. Estimating trait-dependent speciation and

extinction rates from incompletely resolved phylogenies. Syst. Biol. 58:595–611.

Herrera-Alsina, L., P. van Els, and R. S. Etienne. 2018. Detecting the dependence of

diversification on multiple traits from phylogenetic trees and trait data. Syst. Biol. syy057.

Rabosky, D. L. 2019. Phylogenies and Diversification Rates: Variance Cannot Be Ignored.

108