Diversity within the Adenovirus fiber knob hypervariable loops influences primary

receptor interactions.

Alexander T. Baker[1], Alex Greenshields-Watson[2], Lynda Coughlan[3], James A. Davies[1], Hanni

Uusi-Kerttula[1], David K. Cole[2,4], Pierre Rizkallah[2], Alan L. Parker[1]*

[1] Division of Cancer and Genetics, School of Medicine, Cardiff University, Cardiff CF14 4XN,
United Kingdom
[2] Division of Infection and Immunity, School of Medicine, Cardiff University, Cardiff CF14
4XN, United Kingdom
[3] Icahn School of Medicine at Mount Sinai, Department of Microbiology, New York, New
York 10029-6574
[4] Present address: Immunocore Ltd, Abingdon, Oxon, United Kingdom
12
<u>Corresponding author:</u>

*Alan L. Parker
Division of Cancer and Genetics
Henry Wellcome Building
Cardiff University School of Medicine
Heath Park
Cardiff
CF14 4XN
Email: ParkerAL@Cardiff.ac.uk

Keywords: adenovirus, receptor, CD46, CAR, crystal structure

Word count: Abstract: 150 words

25   Main body (excluding methods and figure legends): 4027

Figure count: 8 + 3 Supplementary figures + 1 Supplementary Table

Abstract

Adenovirus based vectors are of increasing relevance for wide ranging therapeutic applications. As

vaccines, vectors derived from species D serotypes 26 and 48 are demonstrating promising efficacy

as protective platforms against infectious diseases. Significant clinical progress has been made, yet

definitive studies underpinning mechanisms of entry, infection, and receptor usage are currently

lacking. Here, we performed structural and biological analysis of the receptor binding fiber-knob

protein of Ad26/48, reporting crystal structures, and modelling putative interactions with two

previously suggested attachment receptors, CD46 and CAR. We provide clear evidence of low affinity

interaction with CAR, with modelling suggesting affinity is attenuated through extended, semi-

flexible loop structures, providing steric hindrance. Conversely, *in silico* and *in vitro* experiments

were unable to provide evidence of interaction between Ad26/48 fiber-knob with CD46. Our findings

provide new insight to the cell-virus interactions of Ad26/48, with important implications for the

design and engineering of optimised Ad-based therapeutics.

 ARTICLE

Adenoviruses (Ads) are increasingly important vectors for wide ranging therapeutic interventions,
from gene delivery and oncolytic agents to platforms for vaccine applications[1-3]. As vaccine vectors,
their use clinically has been popularised by their excellent safety profile coupled with their ability to
induce robust cellular and humoral immunogenicity in humans[4]. Phylogenetically, Ads are diverse
and subdivided across 7 species, A-G[5], based classically on serological cross-reactivity, receptor
usage, haemagglutination properties and, more recently, phylogenetic sequence similarity[6,7].

Most experimental and clinical studies have focussed on the well-studied species C adenovirus, Ad5.
Although potently immunogenic, the efficacy of vaccines based on Ad5 appears hampered by high
seroprevalence rates in humans, and enthusiasm for their use as clinical vaccine platforms has been
dampened by the well-publicised failure of the MERCK STEP vaccine trial. This trial, to evaluate an
Ad5-based HIV vaccine encoding HIV gag/pol/nef antigens, was abandoned due to apparent lack of
efficacy upon 1st term analysis. The study also identified a non-significant trend towards increase HIV
acquisition in a specific high-risk, uncircumcised subset of patients who also had high levels of
baseline pre-existing neutralising antibodies to Ad5[8,9]. As a result, attention has switched from Ad5
based vectors towards the development of alternative adenoviral serotypes with lower rates of pre-
existing immunity. Most notably, vectors under development include those based on species D
serotypes including Ad26, which has entered Phase-III clinical trials as an Ebola vaccine and recently
reported promising immunogenicity in an HIV trial, or chimeric vectors utilising the hexon hyper
variable regions (HVRs) of Ad48 which have undergone Phase-I evaluation as an HIV vaccine[3,10-12].
However, despite extensive clinical advances using these vaccine vectors, we possess very limited
knowledge of their basic biology, particularly with regards to the determinants underpinning their
tropism, mechanisms of cellular entry and receptor usage. Therefore, in this study, we address these
shortcomings through analysis of adenoviral diversity in the context of their receptor binding, fiber
proteins. Whilst Ad are historically divided into seven species, A-G, this may underestimate their
diversity[13-15]. Phylogenetic examination of 56 human Adenovirus fiber proteins from different
species shows deviation from the taxonomy expected based upon whole genome taxonomy, likely
due to recombination events as seen in other Adenoviral proteins[14]. Here, we have sought to
generate high resolution crystallographic structures of the cellular interacting fiber-knob domain of
species D Adenoviruses Ad26 and Ad48, to enable modelling of their putative interactions with
either human Coxsackie and Adenovirus Receptor (CAR) or CD46 (Membrane Cofactor Protein,
MCP), both of which have both been previously suggested as primary receptors for these clinically
significant viral vectors[16,17]. In this study, we employ an integrative workflow utilising X-ray
crystallography, *in silico* modelling, and *in vitro* assays to dissect previous findings suggestive of

interactions of Ad26 and Ad48 with CAR[5,18] and for CD46[16,17,19–21] . Our findings provide shed new

light on the cell-virus interactions of adenovirus and with potential implications for the design and

engineering of optimised Ad-based therapeutics, both for vaccine applications and oncolytic

development, allowing us to minimise off-target or undesirable interactions *in vivo.*

RESULTS

Genetic variability in adenovirus fiber-knob protein

We generated phylogenetic trees of human adenoviruses 1-56 which revealed broader

differentiation in the fiber-knob domain (Fig. 1A) than might be expected based upon taxonomy of

the whole virus (Fig. 1B). In both phylogenetic trees the adenoviruses divide into seven clades

corresponding to the seven adenoviral species, A-G. However, the tree based upon the fiber-knob

domain (Fig. 1A) shows the species D adenoviruses forming a greater number of sub-groups than in

the whole genome tree suggesting greater diversity in the receptors of this species than might be

expected when comparing serotypes. This may be the result of recombination events, as reported

previously for other adenoviral proteins[13,15,22–25]. The opposite is observed in species B adenoviruses,

where simpler groupings are seen when analysing fiber-knob domain alignment than by whole

genome analysis. When divided into sub-species based on whole genome they divide into species B1

and B2, but when the tree is generated based on fiber-knob alone the species B viruses do not divide

into similar groups. The significance of this fiber-knob diversity is unclear, but it has previously been

suggested that the species B1/B2 designation may more closely represent the tissue tropism, than

receptor usage[26,27].

We next calculated the amino acid variability at each position in the aligned adenoviral fiber-knob

sequences, which revealed regions of broad conservation corresponding to β-strands which make up

the main fold of the fiber-knob trimer (Fig. 1C). The positions corresponding to the β-strands of Ad5,

as originally reported by Xia et al[28], are shown by arrows. Comparison shows that the more N-

terminal (A, B, and C) and C-terminal (I and J) β-strands have greater homology across the adenoviral

species than the other sheets. This may relate to the intervening loops between the less tightly

conserved β-strands (D, E, F, G, H) being more apical, a region which is often involved in receptor

interactions[27,29–33].

Structural analysis of Ad26 and Ad48 fiber-knob domain

To investigate diversity of species D adenoviruses, recombinant, 6-His tagged fiber-knob protein

from Ad26 and Ad48 (hereafter referred to as Ad26K and Ad48K) were generated, purified and used

to determine X-ray crystallographic structures of Ad26 (PDB: 6FJN) and Ad48 (PDB: 6FJQ) at

resolutions of 0.97Å and 2.7Å, respectively (Fig. 2A-D). The monomers (Red, Green, Blue) form an

anti-parallel β-barrel, typical of Adenoviral fiber-knob protein, as described by Xia *et al* (PDB 1KNB)[28].

Each monomer interacts with two neighbouring copies to form homotrimer with 3-fold symmetry

(Fig. 2B,D) and a highly stable interface (suppl. 1A). Stability analysis using PISA (Protein Interactions,

Surfaces and Assemblies) software calculates the Ad26 and Ad48 trimers to have >20% lower
interface energy than that of Ad5, indicative of a more stable interaction (Suppl. 1A) between the
monomers of the Ad26 and Ad48 fiber-knobs.

As with the pan species analysis (Fig. 1C), variability of the aligned species D fiber-knobs (Fig. 2E)
confirmed that the β-strands comprising the hydrophobic core of the trimers are highly conserved,
with β-strands demonstrating a high degree of overlap in both spatial position and sequence
variability (suppl. 1B).

Hypervariable loops within the fiber-knob domain are stabilised by inter-loop interactions

Particularly relevant to this study are the DG, GH, HI and IJ loops, linking the indicated strands
corresponding to those in the originally reported Ad5 fiber-knob loops (suppl. 1B)[28]. These loops
have previously been shown to be critical in engagement of CD46 for Ad11, 35, and 21[34–37].
Alignment of these loops with the corresponding loops of Ad5 (species C, CAR interacting), Ad35
(species B, CD46 interacting), Ad11 (species B, CD46 interacting), and Ad37 (species D, CAR and Sialic
acid/GD1a glycan interacting) reveals different topologies in these critical receptor interacting
regions (Fig. 3A)[36–39]. The HI loops of Ad26K and Ad48K are most homologous to those of Ad35K and
Ad37K respectively in terms of amino acid sequence identity (Suppl.1B) and spatial alignment. The
Ad26K DG loop is most homologous to Ad35K but incorporates a three amino acid insertion, while
Ad48K DG loop displays a differing and unique topology. The GH and IJ loops of Ad5, 26, 37, and 48K
demonstrate similar spatial arrangements (though the similarity does not extend to the sequence
identity) but differ from the CD46 utilising Ad11K and Ad35K.

The high-resolution structures obtained allowed us to robustly characterise the loops, seen in the
electron density maps (Fig. 3B, C). To assess loop flexibility and mobility, we assessed the B-factors
(also known as temperature factor), a measure of the confidence in the position of an atom which
can be used to infer flexibility. By assessing the B-Factors, the relative flexibility of the moieties of
interest can be inferred. While the apical domains of some loops demonstrated increased B-Factors
relative to the rest of the molecule, the loops' B-factors are surprisingly low (Fig. 3B, C), suggesting
that they may exhibit limited flexibility.

To investigate whether the different loop conformations were the product of flexibility, or
stabilisation we investigated the inter-loop interactions in the Ad26 and Ad48 structures. This
analysis demonstrated that the GH loop of Ad26K (like those of Ad5, 37 and 48K) does not extend
directly away from the G and H β-sheets, but forms a β-hairpin (Fig. 3A, B) that is maintained by
seven polar contacts within the neighbouring IJ loop which restrict the loops' orientation (Fig. 4A,

Suppl. 2A). Polar contacts were also observed at the apex of several loops, most notably the GH and
CD loops of both fiber-knobs (Fig. 4A, B). The IJ loops form fewer intramolecular polar contacts but
are stabilised by interactions with the adjacent CD and DG loops (Suppl. 2C, D). These interactions
retain the apical residues in a stable conformation, rather than leaving the side chains fully labile.

The B-Factors of the Ad48K DG loop were observed to be polarised normal to the hairpin, with the
outer face of the loop having higher B-Factors compared to the inner face (Fig.4C). This is likely the
result of polar contacts formed between Ser-307, Gln-308, Ala-309, and Leu-304 with Asp-359 and
Gln-357 of the opposing monomer stabilising the conformation of the DG loop. The proline rich
nature of this loop provides further rigidity (Suppl. 3).

Based on this analysis of the inter and intra-loop bonds we suggest that these adenoviral loops may
not be fully flexible variable regions, but organised receptor engagement motifs with carefully
evolved structures. This has direct implications for receptor engagement of these viruses, as the
loops govern previously characterised interactions with CAR and CD46, and are directly involved in
their pathogenicity[26,34,40,41].

Modelling of Ad26K and Ad48K in complex with CAR suggests a low affinity binding interaction

Both CD46 and CAR have been proposed as primary attachment receptors for Ad26 and Ad48[16,17].
Previously generated crystal structures of Ad11K in complex with full length CD46 (PDB: 3O8E), and
Ad37K in complex with CAR-D1 domain (PDB: 2J12) reveal the loops to be essential to receptor
interactions[42]. To investigate the ability of Ad26K and Ad48K to bind these receptors we generated
homology models by alignment of the new Ad26K and Ad48K fiber-knob structures modelled with
the existing fiber-knobs in complex with the receptor of interest and performed energy minimisation
to optimise the conformation to achieve the lowest possible energy interface with which to analyse
the interaction. We performed similar experiments with the well described CAR and CD46 binding
fiber-knob proteins of Ad5 (PDB: 6HCN) and Ad35 (PDB: 2QLK), respectively, as controls.

Modelling of Ad26K and Ad48K in complex with the CAR-D1 domain revealed a region of high
homology with the CAR utilising Ad5 fiber-knob, here after termed the α-interface (Fig. 5A).
Sequence alignment with Ad5K shows that many of the residues previously shown to be critical for
CAR interaction in Ad5K are conserved in Ad26K and Ad48K, including Ser-408, Pro-409, and Tyr-
376[38]. The same is true of residues predicted to interact with CAR directly, such as Lys-417 (number
is for Ad5K). The residues predicted to form direct CAR binding interactions for Ad5K, 26K, and 48K
are pictured in complex with the maximum spatial occupancy of the energy minimised CAR-D1 (Fig.
5C). The high levels of homology are seen to extend to the fold as well as the linear sequence.

Calculation of the binding energies for the modelled complexes predicted a stable α-interface for all

complexes modelled, albeit weaker for the non-CAR utilising Ad35K (Fig.5D) which has low sequence

conservation with Ad5K. However, the interaction is complicated by a second CAR interface, termed

the β-interface (Fig. 6A). The loops forming the β-interface with CAR-D1 differ between Ad5, 26, and

48 fiber-knob (Fig. 6B). The shorter Ad5K DG loop does not clash with the CAR-D1 surface, whereas

the extended Ad26K DG loop forms a partial steric clash, with surface seen to clash with the aligned

CAR-D1, and Ad48K DG loop is seen to form an even larger steric clash. Whilst the longer loop of

Ad26K is expected to be more flexible than that of Ad5K, the Ad48K DG loop is surprisingly stable

due to the characteristics described (Fig.4C).

In vitro IC$_{50}$ evaluation of Ad26K and Ad48K confirms diminished affinity CAR interaction

compared to Ad5K.

Our modelling studies indicate that the longer, inflexible DG loop of Ad48 would be likely sterically

hinder the Ad48K: CAR interaction at the β-interface to a greater extent than the more modest

inhibition of the smaller and more labile loop of Ad26K, which in turn would exhibit more inhibition

of CAR binding than that of Ad5K, where no steric inhibition is observed. Competition inhibition

assays using recombinant fiber-knob protein to inhibit antibody binding to CAR receptor in CHO-CAR

cells (which express CAR, while the parental cell line is established to be non-permissive to Ad

infection) support our observation (Fig. 6C). The IC$_{50}$ (the concentration of protein required to inhibit

50% of antibody binding) of Ad5 is 7.0 ng/10^5 cells, while Ad26K and Ad48K demonstrate IC$_{50}$ values

15.7 and 480 times higher at 0.110µg/10^5 cells and 3.359µg/10^5 cells, respectively, reflecting their

reduced ability to engage CAR due to steric inhibition at the β-interface (Fig. 6C).

Modelling of Ad26K and Ad48K in complex with CD46 indicates minimal interaction

A similar approach was taken to model Ad26K and Ad48K in complex with CD46. Alignments with the

previously published Ad11K-CD46 complex were generated and energy minimised to obtain the

lowest energy state of the complex [35,43]. This interface utilises loops DG, GH, HI and IJ to form a

network of polar interactions with the CD46 Sc1 and Sc2 domains (Fig. 7A)[36].

When Ad35K was modelled in complex with CD46, a network of polar contacts between Ad35 and

CD46 was predicted (Fig. 7B) similar to that observed in the Ad11-CD46 complex crystal structure,

PDB: 3O8E (Fig. 7A)[35]. Previously Ad35K residues Phe-243 , Arg-244, Tyr-260, Arg-279, Ser-282, and

Glu-302 (underlined in Fig.7C) have been implicated as key contact residues for CD46 interaction,

and are conserved in Ad11K (highlighted in blue, Fig.7C)[37]. Our modelling suggests that conversely,

Ad26K and Ad48K are predicted to form very few polar contacts with CD46 with just 2 contacts

predicted for Ad26K (Fig. 7D) and 3 predicted for Ad48K (Fig. 7E). Furthermore, they do not share
any of the critical CD46 binding residues which have been reported previously (underlined, Fig. 7C)
for Ad11K and Ad35K, or any of the predicted interacting residues (blue highlight, Fig. 7C).

We again employed PISA to calculate the binding energy of the various modelled and energy
minimised fiber-knob CD46 complexes (fig.8A). Ad11K, the strongest known CD46 binding
adenovirus[34], was predicted to have the lowest binding energy reflecting its high stability interface,
with Ad35 demonstrating a similar but slightly reduced binding energy. Conversely, Ad26K and
Ad48K are predicted to have lower binding energies, similar to that which may be expected for
random proteins passing in solution, indicating that an interaction between either the Ad26K, or
Ad48K with CD46 is unlikely[44].

While still low compared to the known CD46 interacting Ad11K and Ad35K binding energies, that for
Ad5K was higher than expected for a known non-CD46 interacting adenovirus (Fig. 8A). Inspection of
the model shows that this is due to the close proximity of the large Ad5 HI loop to CD46 (Fig. 8B).
The residues involved in the predicted interaction are not conserved in any known CD46 interface
and suggesting these are random interaction. Furthermore, interaction between the DG loop and
CD46 is integral to known CD46 binding interfaces and is prevented by the Ad5K HI loop laying
between them (Fig. 8B).

IC_{50} evaluation of Ad26K and Ad48K confirms lack of CD46 interaction

Antibody competition inhibition assays in CHO-BC1 cells (CHO cells transduced to express the BC1
isoform of CD46) were used to test the predictions made by modelling (Fig. 8C). These data confirm
that recombinant Ad26K and Ad48K proteins are incapable of inhibiting antibody binding to CD46 at
any concentration tested (up to 2ng/cell), whilst the well-defined CD46 interacting Ad35K
demonstrates strong inhibition, with a calculated IC_{50} of 0.003µg/10^5 cells. Based on these findings it
is highly unlikely that Ad26 or Ad48 are capable of engaging CD46 as a primary receptor.

DISCUSSION

This study reveals the crystal structure of two adenovirus proteins critical to primary receptor engagement, Ad26 and Ad48 fiber-knob, which are important viral vectors currently in human clinical trials[3,11,12]. Despite their advanced development, the field lacks fundamental knowledge regarding the mechanisms of infection for these viral vector platforms. The work we described here provides a combined crystallographic, *in silico,* and *in vitro* approach to investigate adenovirus fiber-knob: receptor interactions with CAR and CD46, two receptors previously proposed to be utilised by these viruses[5,9,17].

Analysis of the phylogenetic relationship between 56 Adenovirus serotypes, both whole genome and fiber-knob domain alignment, confirms diversification into seven adenoviral species[7]. However, generating the phylogenetic tree with fiber-knob sequences, rather than whole genomes, shows additional diversity, not revealed by the traditional taxonomic approach. Adenovirus species D breaks up into several additional sub-clades when focused on the fiber-knob, suggesting greater receptor diversity than might be expected based on the whole virus phylogeny. Similar observations have previously been made in species D hexon and penton[13].

In contrast to species D, the phylogeny of species B adenoviruses, which are known to utilise desmoglein 2 and CD46 as primary receptors, is simplified when focused upon the fiber-knob, indicating less diverse receptor usage[40,45–47]. This simplification in comparison to the whole genomic alignment implies that much of the species diversity must lay in other proteins. The E3 protein, for example, is known to be highly diverse within species B adenoviruses, having previously been exploited in the selection of the oncolytic (cancer killing) virus enadenotucirev, which is currently in clinical trials[13,48,49].

That we see such opposing effects on the species B and D phylogenetic trees when focusing on the fiber-knob, highlights the limitations of simple taxonomic approaches. The current Adenoviral taxonomy is based on antibody neutralisation assays, which are limited by antibodies reliance on surface accessible proteins in the capsid, and does not account for diversity in other viral proteins, as the above suggests for species B. This supports a taxonomic proposal based upon viral genetics rather than antibody neutralisation, as has previously been suggested[13,50].

Many studies on adenovirus neutralisation have focused upon neutralising antibodies (NAbs) which bind to the hexon[51–53]. Following intramuscular vaccination with non-replicating adenoviral vectors, most NAbs are targeted to the hexon; a reflection of its high abundance and surface availability in each viral capsid[51,54]. However, during natural infection, many NAbs target the fiber protein[54],

presumably due to the abundance of fiber produced in the early stages of AdV replication to loosen

cell-cell junctions and facilitate viral spread, prior to lysis and entry of large amounts of whole virus

to the blood stream[55]. For individuals with pre-existing anti-adenovirus immunity derived from

natural infections, anti-fiber NAbs are likely to limit the use of vectors with common adenovirus fiber

proteins by neutralisation of the viral vector prior to its therapeutic effect. Thus, for the

development of vectors to circumvent pre-existing anti-adenovirus immunity for therapeutic use

further exploration of this fiber protein diversity may be beneficial, as well as the on-going studies

using hexon HVR pseudotypes to circumvent anti-hexon immunity[11,56].

Analysis of the adenovirus loops reveals an intricate network of polar interactions which stabilise

their three-dimensional structures. These bonds appear to hold the loops in a conformation which,

in the case of Ad11 and Ad35, facilitates receptor binding. In the Ad26 and Ad48 fiber-knob

structures presented in this study the loops are also held in a stable conformation, though not one

which enables CD46 interaction.

Modelling of Ad26K/Ad48K in complex with CD46 suggested few contacts, and interface energy

calculations using these models predict a weak binding energy. This was confirmed by *in vitro*

competition inhibition assays. These findings appear contradictory with previous studies which

suggest CD46 as the primary receptor for these viruses. However, our findings improves our

knowledge of the cell entry mechanisms of these viruses and the vectors derived from them, and

does not diminish the observed effectiveness of these vaccines.   However, if CD46, a protein

expressed on the surface of all nucleated cells, is not the receptor for these viruses then it is as yet

unknown what the primary tissue tropism determinant is for these clinically significant viruses[57,58].

A similar methodology was applied to the interaction Ad26K/Ad48K with CAR. Inspection of the

modelled complexes indicated a conserved α-interface enabling CAR binding in adenovirus 5, 26, and

48, fiber knobs. However, the structure of the β-interface interaction appears to indicate a novel

mechanism modulating the fiber-knob: CAR affinity. When occupying the intermonomer cleft in the

conformation shown in figure 6B, the DG loops of Ad26 and Ad48 are likely to inhibit CAR binding by

steric hindrance, but if the loops were to shift into a conformation which relieves this clash CAR

binding could occur. Therefore, the ability of these vectors to interact with CAR is a function of the

steric hindrance provided by these loops, sterically hindering the ability of the fiber-knob domain to

engage CAR in a permissive conformation.

The inter-loop contacts described in Figure 4 A, B, and supplemental figure 2, and the normalised B-

factors described in figure 6B will influence the molecular dynamics of the DG-loops. Loops which

can occupy a CAR inhibitory conformation but have fewer stabilising contacts, such as that of Ad26K

(Suppl. 2A, C), should be more permissive to CAR binding. While loops which are less flexible and/or

stabilised in a CAR inhibitory conformation, such as Ad48K (Fig. 6B, Suppl. 2B, D) should result in a

fiber-knob which is less able to bind CAR. This hypothesis fits the competition inhibition studies

shown in Figure 6C, which demonstrate that Ad26K has and apparent ~15x lower affinity for CAR

than Ad5K, and Ad48K has 500x lower affinity.

Species D adenoviruses (Suppl. 3) have a large range of different DG loops. Most sequences have

lengths equal to, or greater than, that of Ad26K, making it plausible that they too could modulate

the fiber-knob interaction with CAR. However, the magnitude of this effect will be dependent on the

individual molecular dynamics of the DG-loops and its interactions with adjacent residues.

Assuming this mechanism of CAR binding regulation is broadly applicable, it may have important

implications for adenoviral vector design. The presence of a high affinity receptor for the virus can

mask the low affinity CAR interaction, creating a "hidden tropism" only observed if the virus is forced

to rely upon it. Expression of CAR on human erythrocytes suggests the potential for sequestration of

virotherapies in the blood[59]. CAR expression in lung epithelial tissues offers another site for potential

off target activity[60,61]. Therefore, many virotherapies previously thought to be non-CAR binding

adenoviruses may in fact demonstrate weak CAR tropism, driving off target infections or resulting in

sequestration of the vector in organs other than that target. This may not be of grave consequence

for non-replicating vectors, such as viral vaccines, but in vectors which rely upon controlled

replication of the virotherapy in the target tissues, such as cancer targeting oncolytic vectors, this

could result in off-target infection, dysregulated expression of therapeutic proteins, and reduced

delivery to the desired point of need.

The work undertaken in this study presents, for the first time, the crystal structures of the fiber-knob

domain protein of adenovirus 26 (PDB: 6FJN), and adenovirus 48 (PDB: 6FJQ) fiber-knob protein. In

addition we report a new crystal structure for adenovirus serotype 5 fiber-knob protein (PDB: 6HCN)

with improved resolution compared to the existing structure (PDB: 1KNB)[28]. We utilise these

structures to investigate the ability of these proteins to interact with the putative receptors, CAR and

CD46, through computational modelling and *in vitro* experimentation. We demonstrate that

adenovirus 26 and 48 fiber-knob domains have a weak ability to bind CAR, and no demonstratable

CD46 interaction, suggesting that these viruses are unlikely to utilise these proteins as a primary

receptor *in vivo*. We suggest that CAR binding is moderated by a previously unreported mechanism

of steric inhibition which may apply to other adenoviruses and demonstrate an *in silico* methodology

capable of rapidly predicting these interactions. These findings enhance our understanding of the

virology of adenovirus infection, and have direct implications for virotherapy vector design, which

often rely upon carefully controlled receptor tropisms to achieve specificity and efficacy[9,18,29,62].

METHODS

Analysis of adenovirus genetic diversity

Genome alignment

Representative whole nucleotide genomes of adenoviral species 1-56 were selected from the
National Center for Biotechnology Information (NCBI), and aligned using the EMBL-EBI Clustal
Omega tool[63–65]. Fiber-knob domain amino acid sequences were derived from the same genome
sequences, defined as the translated nucleotide sequence of the fiber protein (pIV) from the
conserved TLW hinge motif to the protein C-terminus, and aligned in the same manner as the whole
genomes. Phylogenetic relationships were inferred using the maximum likelihood method based
upon the Jukes Cantor model for the whole genome nucleotide analysis[66], and the Poisson
correction model for the fiber-knob amino acid analysis[67], using the MEGA X software[68]. Confidence
was determined by bootstrap analysis (500 replicates)[69] and trees displayed condensed at 70%
confidence (percentage confidence values shown at each node).

Fiber-knob amino acid variability:

Amino acid sequence variability scores were calculated from the Clustal omega aligned amino acid
sequences of the fiber-knob domains of either Adenoviruses 1-56, or only the species D
adenoviruses. Analysis was performed using the protein variability server (PVS), using a consensus
base sequence and the Wu-Kabat method[70,71].

Expression and purification of Recombinant Fiber-Knob protein:

Purification of recombinant Ad26K and Ad48K proteins was performed as previously described[18,72,73].
Briefly, fiber-knob DNA sequences were synthesised commercially by GenScript (Nanjing, China) with
flanking BamHI and HindIII restriction digest sites, in frame, cloned into pQE-30 vector by BamHI,
HindIII double digest and ligation, and the resultant plasmid was transformed into SG13009 *E.coli* by
heat-shock. For expression and purification, E.coli were resuspended in lysis buffer (50mM Tris,
pH8.0, 300mM NaCl, 1% (v/v) NP40, 1mg/ml Lysozyme, 1mM β-mercaptoethanol) and incubated at
room temperature for 30mins. Lysate was clarified by centrifugation at 30,000g for 30mins and
filtered through a 0.22μm syringe filter (Millipore, Abingdon, UK). Clarified lysate was then loaded
onto a 5ml HisTrap FF nickel affinity chromatography column (GE) at 2.0ml/min and washed with 5
column volumes into elution buffer A (50mM Tris [pH8.0], 300mM NaCl, 1mM β-mercaptoethanol).
Protein was eluted by 30min gradient elution from buffer A to B (buffer A + 400mM Imidazole).
Fractions were analysed by reducing SDS-PAGE, and Fiber-knob containing fractions further purified
using a superdex 200 10/300 size exclusion chromatography column (GE) in crystallisation buffer (10

mM Tris [pH 8.0] and 30 mM NaCl). Fractions were analysed by SDS-PAGE and pure fractions

concentrated by centrifugation in Vivaspin 10,000 MWCO (Sartorius, Goettingen, Germany)

proceeding crystallisation.

Competition Inhibition Assays:

Sample Preparation:

Competition inhibition assays were performed as previously described[18]. Briefly, CHO cells

expressing the appropriate receptor (CAR: CHO-CAR, CD46: CHO-BC1), were pre-incubated with

$0.0001\text{-}100\mu g/10^5$ cells of recombinant soluble knob protein. Primary staining was performed with

anti-CD46 antibody (MEM-258) or anti-CAR antibody (RmcB) to complex receptors unbound by fiber-

knob protein, and secondary staining was performed with Alexa-647 labelled goat anti-mouse

F(ab')2 (ThermoFisher) [18,72,73]. All antibodies were used at a concentration of 2μg/ml.

Flow cytometric analysis:

Samples were run in triplicate and analysed by flow cytometry on Attune NxT (ThermoFisher), and

analysed using FlowJo v10 (FlowJo, LLC) by gating on singlets, cell population, and Alexa-647 positive

cells, respectively. Total fluorescence (TF) was used as the measure of inhibition, rather than

percentage of fluorescent cells in the total population, to account for the presence of multiple

receptor copies per cell surface which can enable partial inhibition of antibody binding on the cell

surface, while the cell remains fluorescently labelled. TF was defined as the percentage of Alexa-647

positive cells in the single cell population for each sample and multiplied by the median fluorescent

intensity (MFI) of the Alexa-647 positive single cell population in each sample. Data points are the

mean total fluorescence of three biological replicates with error given as the standard deviation

from the mean. IC_{50} curves were fitted by non-linear regression as previously described, and used to

determine the IC_{50} concentrations[18,72,73].

Crystallisation and structure determination.

Protein samples were purified into crystallisation buffer (10 mM TRIS [pH 8.0] and 30 mM NaCl). The

final protein concentration was approximately 7.5 mg/ml. Commercial crystallisation screen

solutions were dispensed into 96-well plates using an Art-Robbins Instruments Griffon dispensing

robot (Alpha Biotech, Ltd), in sitting-drop vapour-diffusion format. Drops containing 200 nl of screen

solution and 200 nl of protein solution were equilibrated against a reservoir of 60 ⬚l crystallisation

solution. The plates were sealed and incubated at 18°C.

Crystals of Ad5K appeared in PACT *Premier* condition D04 (0.1 M MMT, pH 7.0, 20% PEG 1500),

within 1 to 7 days. Crystals of Ad26K appeared within 1 to 7 days, in PACT *Premier* (Molecular

Dimensions, Suffolk, UK) condition A04 (0.1 M MMT [DL-Malic acid, MES monohydrate, Tris], pH 6.0,

25% PEG 1500. Crystals of Ad48K appeared in PACT *Premier* condition D02 (0.1 M Bis-Tris-propane,

pH 6.5, 20% PEG 3350, 0.2M NaNO3), within 2 weeks. Crystals were cryoprotected with reservoir

solution to which ethylene glycol was added at a final concentration of 25%. Crystals were harvested

in thin plastic loops and stored in liquid nitrogen for transfer to the synchrotron. Data were collected

at Diamond Light Source beamline I04, running at a wavelength of 0.9795Å. During data collection,

crystals were maintained in a cold air stream at 100°K. Dectris Pilatus 6M detectors recorded the

diffraction patterns, which were analysed and reduced with XDS, Xia2[74], DIALS, and Autoproc[75].

Scaling and merging data was completed with Pointless, Aimless and Truncate from the CCP4

package[76]. Structures were solved with PHASER[77], and COOT[78] was used to correct the sequences

and adjust the models. REFMAC5[79] was used to refine the structures and calculate maps. Graphical

representations were prepared with PyMOL[80]. Reflection data and final models were deposited in

the PDB database with accession codes: Ad5K, **6HCN**; Ad26k, **6FJN**; and Ad48k, **6JFQ**.

Energy Minimisation:

Energy Minimisations were performed using the YASARA self-parametrising energy minimisation

algorithm as performed by the YASARA energy minimisation server, and results were visualised in

PyMoll[43,80,81].

Calculation of Interface Energy:

Interface energies were calculated using QT-PISA using biological protein assemblies and excluding

crystallographic interfaces[82]. Values are the mean of the three symmetrical interfaces in each trimer

and error is the standard deviation from the mean, any values above -3.0 kcalmol^{-1} were considered

to be background as shown as a red dashed line on graphs[44].

Sequence visualisations:

Performed using the Clustal Omega multiple sequence alignment algorithm and visualised with

BioEdit[63–65,83].

B-Factor Normalisation:

Comparing order between different structures by comparing individual B-factors can be misleading.

Post-refinement B-factors relate to the Wilson B-factor, which can vary widely between data sets,

even from the same crystal preparation. A valid comparison between different structures can be

achieved by comparing normalised B-factors instead. Normalisation was performed by dividing

individual atomic B-factors by the average B-factor of the whole data set, quantifying the range of

internal flexibility in a structure. This normalised B- factor can then be compared between different

data sets.

Acknowledgments:

428   *The authors would like to thank Diamond Light Source for beamtime (proposal mx14843), and*
429   *the staff of beamline I04 for assistance with data collection. We are also grateful to Elmar Kreiger*
430   *and the team who maintain the YASARA energy minimisation server, as well as Pedro Reche and the*
431   *Immunomedicine Group for maintenance of the Protein Variability Server.*

REFERENCES

1. Combination Adenovirus + Pembrolizumab to Trigger Immune Virus Effects - Full Text View - ClinicalTrials.gov. Available at: https://clinicaltrials.gov/ct2/show/NCT02798406?term=adenovirus&recrs=a&cond=Glioblastoma&rank=2. (Accessed: 23rd September 2017)

2. A Safety and Efficacy Study of ChAdOx1 LS2 and MVA LS2 - Full Text View - ClinicalTrials.gov. Available at: https://clinicaltrials.gov/ct2/show/NCT03203421?term=adenovirus&recrs=a&cond=malaria&rank=2. (Accessed: 23rd September 2017)

3. Long-term Safety Follow-up of Participants Exposed to the Candidate Ebola Vaccines Ad26.ZEBOV and/or MVA-BN-Filo - Full Text View - ClinicalTrials.gov. Available at: https://clinicaltrials.gov/ct2/show/NCT02661464?term=adenovirus&recrs=a&cond=ebola&rank=1. (Accessed: 23rd September 2017)

4. Lee, C. S. *et al.* Adenovirus-mediated gene delivery: Potential applications for gene and cell-based therapies in the new era of personalized medicine. *Genes Dis.* **4,** 43–63 (2017).

5. Abbink, P. *et al.* Comparative Seroprevalence and Immunogenicity of Six Rare Serotype Recombinant Adenovirus Vaccine Vectors from Subgroups B and D. *J. Virol.* **81,** 4654–4663 (2007).

6. Family - Adenoviridae. in *Virus Taxonomy* (eds. King, A. M. Q., Adams, M. J., Carstens, E. B. & Lefkowitz, E. J.) 125–141 (Elsevier, 2012). doi:10.1016/B978-0-12-384684-6.00009-4

7. International Committee on Taxonomy of Viruses. *Virus taxonomy: classification and nomenclature of viruses : sixth report of the International Committee on Taxonomy of Viruses.* (Springer-Verlag, 1995).

8. Mast, T. C. *et al.* International epidemiology of human pre-existing adenovirus (Ad) type-5, type-6, type-26 and type-36 neutralizing antibodies: correlates of high Ad5 titers and implications for potential HIV vaccine trials. *Vaccine* **28,** 950–957 (2010).

9. Uusi-Kerttula, H., Hulin-Curtis, S., Davies, J. & Parker, A. L. Oncolytic Adenovirus: Strategies and Insights for Vector Design and Immuno-Oncolytic Applications. *Viruses* **7,** 6009–6042 (2015).

10. Priddy, F. H. *et al.* Safety and Immunogenicity of a Replication-Incompetent Adenovirus Type 5 HIV-1 Clade B *gag/pol/nef* Vaccine in Healthy Adults. *Clin. Infect. Dis.* **46,** 1769–1781 (2008).

11. Baden, L. R. *et al.* First-in-Human Evaluation of a Hexon Chimeric Adenovirus Vector Expressing HIV-1 Env (IPCAVD 002). *J. Infect. Dis.* **210,** 1052–1061 (2014).

12. Barouch, D. H. *et al.* Evaluation of a mosaic HIV-1 vaccine in a multicentre, randomised, double-blind, placebo-controlled, phase 1/2a clinical trial (APPROACH) and in rhesus monkeys (NHP 13-19). *The Lancet* **392,** 232–243 (2018).

13. Robinson, C. M. *et al.* Molecular evolution of human adenoviruses. *Sci. Rep.* **3,** 1812 (2013).

14. Rux, J. J., Kuser, P. R. & Burnett, R. M. Structural and Phylogenetic Analysis of Adenovirus Hexons by Use of High-Resolution X-Ray Crystallographic, Molecular Modeling, and Sequence-Based Methods. *J. Virol.* **77,** 9553–9566 (2003).

15. Singh, G. *et al.* Overreliance on the Hexon Gene, Leading to Misclassification of Human Adenoviruses. *J. Virol.* **86,** 4693–4695 (2012).

16. Teigler, J. E., Iampietro, M. J. & Barouch, D. H. Vaccination with Adenovirus Serotypes 35, 26, and 48 Elicits Higher Levels of Innate Cytokine Responses than Adenovirus Serotype 5 in Rhesus Monkeys. *J. Virol.* **86,** 9590–9598 (2012).

17. Li, H. *et al.* Adenovirus Serotype 26 Utilizes CD46 as a Primary Cellular Receptor and Only Transiently Activates T Lymphocytes following Vaccination of Rhesus Monkeys. *J. Virol.* **86,** 10862–10865 (2012).

18. Uusi-Kerttula, H. *et al.* Pseudotyped αvβ6 integrin-targeted adenovirus vectors for ovarian cancer therapies. *Oncotarget* **7,** 27926–27937 (2016).

19. Camacho, Z. T., Turner, M. A., Barry, M. A. & Weaver, E. A. CD46-Mediated Transduction of a
Species D Adenovirus Vaccine Improves Mucosal Vaccine Efficacy. *Hum. Gene Ther.* **25,** 364–374
(2014).
20. Weaver, E. A. & Barry, M. A. Low Seroprevalent Species D Adenovirus Vectors as Influenza
Vaccines. *PLoS ONE* **8,** (2013).
21. Geisbert, T. W. *et al.* Recombinant Adenovirus Serotype 26 (Ad26) and Ad35 Vaccine Vectors
Bypass Immunity to Ad5 and Protect Nonhuman Primates against Ebolavirus Challenge. *J. Virol.*
85, 4222–4233 (2011).
22. Wang, Y. *et al.* Phylogenetic evidence for intratypic recombinant events in a novel human
adenovirus C that causes severe acute respiratory infection in children. *Sci. Rep.* **6,** 23014 (2016).
23. Ebner, K., Pinsker, W. & Lion, T. Comparative Sequence Analysis of the Hexon Gene in the Entire
Spectrum of Human Adenovirus Serotypes: Phylogenetic, Taxonomic, and Clinical Implications. *J.*
Virol. **79,** 12635–12642 (2005).
24. Li, P. *et al.* Circulation of HAdV-41 with diverse genome types and recombination in acute
gastroenteritis among children in Shanghai. *Sci. Rep.* **7,** 3548 (2017).
25. Al Qurashi, Y. M. A., Alkhalaf, M. A., Lim, L., Guiver, M. & Cooper, R. J. Sequencing and
phylogenetic analysis of the hexon, fiber, and penton regions of adenoviruses isolated from AIDS
patients. *J. Med. Virol.* **84,** 1157–1165 (2012).
26. Pache, L., Venkataraman, S., Nemerow, G. R. & Reddy, V. S. Conservation of fiber structure and
CD46 usage by subgroup B2 adenoviruses. *Virology* **375,** 573–579 (2008).
27. Russell, W. C. Adenoviruses: update on structure and function. *J. Gen. Virol.* **90,** 1–20 (2009).
28. Xia, D., Henry, L. J., Gerard, R. D. & Deisenhofer, J. Crystal structure of the receptor-binding
domain of adenovirus type 5 fiberprotein at 1.7 Å resolution. *Structure* **2,** 1259–1270 (1994).
29. Baker, A. T., Aguirre-Hernández, C., Halldén, G. & Parker, A. L. Designer Oncolytic Adenovirus:
Coming of Age. *Cancers* **10,** 201 (2018).
30. Amoureux, M.-C. *et al.* Polysialic Acid Neural Cell Adhesion Molecule (PSA-NCAM) is an adverse
prognosis factor in glioblastoma, and regulates olig2 expression in glioma cell lines. *BMC Cancer*
10, 91 (2010).
31. Lenman, A. *et al.* Human Adenovirus 52 Uses Sialic Acid-containing Glycoproteins and the
Coxsackie and Adenovirus Receptor for Binding to Target Cells. *PLOS Pathog.* **11,** e1004657
(2015).
32. Arnberg, N. *et al.* Adenovirus Type 37 Binds to Cell Surface Sialic Acid Through a Charge-
Dependent Interaction. *Virology* **302,** 33–43 (2002).
33. Arnberg, N., Edlund, K., Kidd, A. H. & Wadell, G. Adenovirus Type 37 Uses Sialic Acid as a Cellular
Receptor. *J. Virol.* **74,** 42–48 (2000).
34. Cupelli, K. *et al.* Structure of Adenovirus Type 21 Knob in Complex with CD46 Reveals Key
Differences in Receptor Contacts among Species B Adenoviruses. *J. Virol.* **84,** 3189–3200 (2010).
35. Persson, B. D. *et al.* Structure of the Extracellular Portion of CD46 Provides Insights into Its
Interactions with Complement Proteins and Pathogens. *PLOS Pathog.* **6,** e1001122 (2010).
36. Persson, B. D. *et al.* Adenovirus type 11 binding alters the conformation of its receptor CD46.
Nat. Struct. Mol. Biol. **14,** 164–166 (2007).
37. Wang, H. *et al.* Identification of CD46 binding sites within the adenovirus serotype 35 fiber knob.
J. Virol. **81,** 12785–12792 (2007).
38. Kirby, I. *et al.* Identification of Contact Residues and Definition of the CAR-Binding Site of
Adenovirus Type 5 Fiber Protein. *J. Virol.* **74,** 2804–2813 (2000).
39. Nilsson, E. C. *et al.* The GD1a glycan is a cellular receptor for adenoviruses causing epidemic
keratoconjunctivitis. *Nat. Med.* **17,** 105–109 (2011).
40. Fleischli, C. *et al.* Species B adenovirus serotypes 3, 7, 11 and 35 share similar binding sites on
the membrane cofactor protein CD46 receptor. *J. Gen. Virol.* **88,** 2925–2934 (2007).
41. Ismail, A. M. *et al.* Selection Pressure in the Human Adenovirus Fiber Knob Drives Cell Specificity
in Epidemic Keratoconjunctivitis. *J. Virol.* **90,** 9598–9607 (2016).

42. Seiradake, E., Lortat-Jacob, H., Billet, O., Kremer, E. J. & Cusack, S. Structural and Mutational
Analysis of Human Ad37 and Canine Adenovirus 2 Fiber Heads in Complex with the D1 Domain
of Coxsackie and Adenovirus Receptor. *J. Biol. Chem.* **281,** 33704–33716 (2006).

43. Krieger Elmar, Koraimann Günther & Vriend Gert. Increasing the precision of comparative
models with YASARA NOVA—a self-parameterizing force field. *Proteins Struct. Funct.*
Bioinforma. **47,** 393–402 (2002).

44. Jensen, J. H. Predicting accurate absolute binding energies in aqueous solution: thermodynamic
considerations for electronic structure methods. *Phys. Chem. Chem. Phys.* **17,** 12441–12451
(2015).

45. Vassal-Stermann, E. *et al.* Mapping of Adenovirus of serotype 3 fibre interaction to desmoglein 2
revealed a novel 'non-classical' mechanism of viral receptor engagement. *Sci. Rep.* **8,** (2018).

46. Wang, H. *et al.* Desmoglein 2 is a receptor for adenovirus serotypes 3, 7, 11 and 14. *Nat. Med.*
17, 96–104 (2011).

47. Segerman, A., Arnberg, N., Erikson, A., Lindman, K. & Wadell, G. There are two different species
B adenovirus receptors: sBAR, common to species B1 and B2 adenoviruses, and sB2AR,
exclusively used by species B2 adenoviruses. *J. Virol.* **77,** 1157–1162 (2003).

48. Kuhn, I. *et al.* Directed Evolution Generates a Novel Oncolytic Virus for the Treatment of Colon
Cancer. *PLOS ONE* **3,** e2409 (2008).

49. Illingworth, S. *et al.* Preclinical Safety Studies of Enadenotucirev, a Chimeric Group B Human-
Specific Oncolytic Adenovirus. *Mol. Ther. Oncolytics* **5,** 62–74 (2017).

50. HAdV Working Group, http://hadvwg.gmu.edu/.

51. Bradley, R. R., Lynch, D. M., Iampietro, M. J., Borducchi, E. N. & Barouch, D. H. Adenovirus
serotype 5 neutralizing antibodies target both hexon and fiber following vaccination and natural
infection. *J. Virol.* **86,** 625–629 (2012).

52. Bradley, R. R. *et al.* Adenovirus Serotype 5-Specific Neutralizing Antibodies Target Multiple
Hexon Hypervariable Regions. *J. Virol.* **86,** 1267–1272 (2012).

53. Ma, J. *et al.* Manipulating Adenovirus Hexon Hypervariable Loops Dictates Immune
Neutralisation and Coagulation Factor X-dependent Cell Interaction In Vitro and In Vivo. *PLoS*
Pathog. **11,** (2015).

54. Parker, A. L. *et al.* Effect of neutralizing sera on factor x-mediated adenovirus serotype 5 gene
transfer. *J. Virol.* **83,** 479–483 (2009).

55. Walters, R. W. *et al.* Adenovirus Fiber Disrupts CAR-Mediated Intercellular Adhesion Allowing
Virus Escape. *Cell* **110,** 789–799 (2002).

56. Keele, B. F. *et al.* Adenovirus prime, Env protein boost vaccine protects against neutralization-
resistant SIVsmE660 variants in rhesus monkeys. *Nat. Commun.* **8,** 15740 (2017).

57. Andrews, P. W. *et al.* A human cell-surface antigen defined by a monoclonal antibody and
controlled by a gene on human chromosome 1. *Ann. Hum. Genet.* **49,** 31–39 (1985).

58. Uhlen, M. *et al.* A pathology atlas of the human cancer transcriptome. *Science* **357,** eaan2507
(2017).

59. Seiradake, E. *et al.* The Cell Adhesion Molecule "CAR" and Sialic Acid on Human Erythrocytes
Influence Adenovirus In Vivo Biodistribution. *PLOS Pathog.* **5,** e1000277 (2009).

60. Cohen, C. J. *et al.* The coxsackievirus and adenovirus receptor is a transmembrane component of
the tight junction. *Proc. Natl. Acad. Sci.* **98,** 15191–15196 (2001).

61. Awasthi, V., Meinken, G., Springer, K., Srivastava, S. C. & Freimuth, P. Biodistribution of
Radioiodinated Adenovirus Fiber Protein Knob Domain after Intravenous Injection in Mice. *J.*
Virol. **78,** 6431–6438 (2004).

62. Uusi-Kerttula, H. *et al.* Ad5NULL-A20 - a tropism-modified, αvβ6 integrin-selective oncolytic
adenovirus for epithelial ovarian cancer therapies. *Clin. Cancer Res.* clincanres.1089.2018 (2018).
doi:10.1158/1078-0432.CCR-18-1089

63. Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple sequence alignments
using Clustal Omega. *Mol. Syst. Biol.* **7,** 539–539 (2011).

64. Li, W. *et al.* The EMBL-EBI bioinformatics web and programmatic tools framework. *Nucleic Acids*
Res. **43,** W580-4 (2015).
65. McWilliam, H. *et al.* Analysis Tool Web Services from the EMBL-EBI. *Nucleic Acids Res.* **41,**
W597–W600 (2013).
66. Jukes, T. H. & Cantor, C. R. Evolution of Protein Molecules. in *Mammalian Protein Metabolism*
21–132 (Elsevier, 1969). doi:10.1016/B978-1-4832-3211-9.50009-7
67. Zuckerkandl, E. & Pauling, L. Evolutionary Divergence and Convergence in Proteins. in *Evolving*
Genes and Proteins (eds. Bryson, V. & Vogel, H. J.) 97–166 (Academic Press, 1965).
doi:10.1016/B978-1-4832-2734-4.50017-6
68. Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular Evolutionary Genetics
Analysis across Computing Platforms. *Mol. Biol. Evol.* **35,** 1547–1549 (2018).
69. Felsenstein, J. CONFIDENCE LIMITS ON PHYLOGENIES: AN APPROACH USING THE BOOTSTRAP.
Evol. Int. J. Org. Evol. **39,** 783–791 (1985).
70. Kabat, E. A., Wu, T. T. & Bilofsky, H. Unusual distributions of amino acids in complementarity-
determining (hypervariable) segments of heavy and light chains of immunoglobulins and their
possible roles in specificity of antibody-combining sites. *J. Biol. Chem.* **252,** 6609–6616 (1977).
71. Garcia-Boronat, M., Diez-Rivero, C. M., Reinherz, E. L. & Reche, P. A. PVS: a web server for
protein sequence variability analysis tuned to facilitate conserved epitope discovery. *Nucleic*
Acids Res. **36,** W35–W41 (2008).
72. Coughlan, L. *et al.* Retargeting Adenovirus Serotype 48 Fiber Knob Domain by Peptide
Incorporation. *Hum. Gene Ther.* **25,** 385–394 (2014).
73. Coughlan, L. *et al.* In Vivo Retargeting of Adenovirus Type 5 to αvβ6 Integrin Results in Reduced
Hepatotoxicity and Improved Tumor Uptake following Systemic Delivery. *J. Virol.* **83,** 6416–6428
(2009).
74. Winter, G., Lobley, C. M. C. & Prince, S. M. Decision making in xia2. *Acta Crystallogr. D Biol.*
Crystallogr. **69,** 1260–1273 (2013).
75. Vonrhein, C. *et al.* Data processing and analysis with the autoPROC toolbox. *Acta Crystallogr. D*
Biol. Crystallogr. **67,** 293–302 (2011).
76. Dodson, E. J., Winn, M. & Ralph, A. [32] Collaborative computational project, number 4:
Providing programs for protein crystallography. in *Methods in Enzymology* **277,** 620–633
(Academic Press, 1997).
77. McCoy, A. J. *et al.* Phaser crystallographic software. *J. Appl. Crystallogr.* **40,** 658–674 (2007).
78. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta Crystallogr. D*
Biol. Crystallogr. **60,** 2126–2132 (2004).
79. Murshudov, G. N. *et al.* REFMAC5 for the refinement of macromolecular crystal structures. *Acta*
Crystallogr. D Biol. Crystallogr. **67,** 355–367 (2011).
80. Schrödinger, LLC. The PyMOL Molecular Graphics System, Version 2.0.
81. Krieger, E. *et al.* Improving physical realism, stereochemistry, and side-chain accuracy in
homology modeling: Four approaches that performed well in CASP8. *Proteins Struct. Funct.*
Bioinforma. **77,** 114–122 (2009).
82. Krissinel, E. & Henrick, K. Inference of macromolecular assemblies from crystalline state. *J. Mol.*
Biol. **372,** 774–797 (2007).
83. Hall, T. BioEdit: a user-friendly biological sequence alignment editor and analysis program for
Windows 95/98/NT. *Nucleic Acids Symp. Ser.* **41,** 95–98 (1999).
627
628

 Figure legends

Figure 1: Phylogenetic analysis of adenoviruses mapped by whole genome and fiber-knob domain.
Condensed maximum likelihood trees (percentage confidence shown by numbers next to nodes)
were generated from alignments of fiber-knob domain amino acid sequences of adenoviruses 1-56
(A) or whole genome NT sequences (B). Adenoviruses divide into 7 subspecies, as denoted in the
key, regardless of alignment used, but the species D adenoviruses divide into additional sub-species
when determined by fiber-knob alignment. Numbers next to nodes denote confidence. Wu-Kabat
variability analysis of the Clustal omega aligned fiber-knob domains amino acid sequences of
adenoviruses 1-56 (C) reveals regions of low amino acid variability corresponding to beta-sheets. The
locations of Ad5 β-strands, as described by Xia et al (1995), are aligned to the structure and shown
by arrows, the corresponding positions are coloured in red.

Figure 2: Overview of the Ad26 and Ad48 fiber-knob protein structures (PDB 6FJN and 6FJQ,
respectively). The surface representation of the trimeric Ad26K biological assembly is shown side-on
with the cartoon representation shown for the nearest monomer (A) and the top-down view of the
same Ad26K trimer, as it would appear looking towards the virion, is seen as a cartoon
representation (B), with each monomer coloured in Red, Green, or Blue with the hypervariable loops
extending between the β-strands (Dark blue) coloured as follows: AB-Green, BC – Purple, CD –
Brown, DE – Orange, DG – Red, EG – Pink, GH – Purple, HI – Yellow, IJ – Light Blue. The Ad48K trimer
is shown similarly (C,D). The Wu-Kabat variability plot of the fiber-knob domains of species D
adenoviruses shows regions of low variability (E) with the locations of the Ad48K β-strands shown by
arrows above the graph, and the positions coloured blue, or red when the position is a β-strand in
both Ad26K and Ad48K.

Figure 3: Comparison of HI, DG, GH, and IJ loops of adenoviruses used in this study. The
hypervariable loops of Ad26K (Green) and Ad48K (Cyan) relevant to this study (HI, DG, GH, and IJ)
are shown in the context of the control virus fiber-knob domains, Ad5K (Orange), Ad11K (Yellow),
Ad35K (Pink), and Ad37K (Purple). The electron density achieved in the loops of Ad26K (B) and
Ad48K (C) are shown as mesh. The fitted residues are seen as stick representations, with oxygen and
nitrogen atoms coloured red and blue, respectively and other atoms coloured according to their
relative B-factors with warmer colours indicating higher B-factor values.

Figure 4: Hypervariable loop conformations and contacts residues. The residues comprising the
indicated loops of Ad26K (A) and Ad48K (B) are shown diagrammatically with numbers indicating the
start and end residues of each loop depicted. The network of intraloop polar interactions is shown
by solid lines (one polar bond), and dashed lines (two polar bonds, colour variations are only for ease

of viewing) between interacting residues, similar interloop bonds are also present as visualised in

Suppl. 2. Residues forming part of a helical motif are shaded in blue. The Ad48K DG-loop is seen to

form contacts to the opposing monomer across the inter-monomeric cleft (C). The labelled residues

forming polar contacts (shown as sticks) are coloured by relative B-factor, with warmer colours

indicating higher relative B-factors, as is the cartoon representation of the loop. The opposing Ad48K

monomer is seen as a ribbon representation of the carbon-α chain in cyan and the surface of the

Ad48K trimer seen as a semi-transparent grey surface.

Figure 5: Modelling of the Ad26K and Ad48K interaction with CAR at the α-interface. The α -

interface region is shown by the box on the structural alignment of Ad5K (Orange), 26K (Green), and

48K (Cyan) fiber-knob domain crystal structures in complex with CAR-D1 domain (Grey) as

determined by homology alignment to the previously reported Ad37K CAR-D1 structure (PDB: 2J12)

(A). The aligned amino acid sequence of the investigated fiber-knobs (B) and the predicted α-

interface forming CAR-D1 binding residues are highlighted in blue, with the underlined residues

representing the Ad5K amino acids shown by Kirby *et al* (2000) to be important for CAR interaction.

Conservation of key residues can be seen between Ad5K, Ad26K, and Ad48K fiber-knobs. This

conservation is visualised, with the contact residues comprising the α-interface with Ad5K, 26K, and

48K shown as sticks in complex with the energy minimised CAR-D1 domain (Grey), shown as the

surface of the maximum spatial occupancy of the aligned CAR-D1 monomers from each of the

energy minimised models in complex with Ad5K, 26K, and 48K fiber-knobs (C). (D) Plots the binding

energy of the energy minimised fiber-knob proteins to CAR-D1 complex in the α-interface. Lower

binding energy indicates a more stable interface with the red line depicting 3.0 KcalMol^{-1}, which can

be considered background.

Figure 6: Modelling of the Ad26K and Ad48K interaction with CAR at the β-interface. The β-

interface region is shown by the box on the structural alignment of Ad5K (Orange), 26K (Green), and

48K (Cyan) fiber-knob domain crystal structures in complex with CAR-D1 domain (Grey) as

determined by homology alignment to the previously reported Ad37K CAR-D1 structure (PDB: 2J12)

(A). A dot surface shows the surface of Ad5K, 26K, and 48K DG-loops in the inter-monomer cleft (B).

The boxes denote the maximum B-factor of the corresponding loops, which are shown as putty

representations with thicker regions indicating higher relative B-factors, from which we can infer the

relative stability of the loops. Antibody competition inhibition assay (C) shows the relative inhibitory

ability of the Ad5, 35, 26, and 48, fiber-knob domains in CAR expressing CHO-CAR cells, with the

calculated IC$_{50}$ values shown in boxes. IC$_{50}$ curves are fitted by non-linear regression. Error bars

represent standard deviation of 3 biological replicates.

Figure 7: Modelling of the Ad26K and Ad48K with CD46. Red dashes show contacts between the
energy minimised crystal structure of CD46 SC1 and SC2 domains (grey cartoon) and Ad11K in
complex (PDB 3O8E). The known CD46 interacting fiber-knob, Ad35K (purple), is aligned to the
above crystal structure and energy minimised (B). Amino acid sequence alignment of the tested
fiber-knob proteins (C) shows conservation of residues previously shown by Wang *et al* (2007) to be
key to CD46 binding (underlined) between the known CD46 binding fiber-knobs, Ad11K and Ad35K.
Residues highlighted in blue are predicted to form direct contacts with CD46 in the energy
minimised models. Similar alignments to that performed with Ad35K are shown for Ad26K (Green -
D) and Ad48K (Cyan - E). In all models red dashes indicate polar contacts between the residues
shown as stick representations.

Figure 8: Binding energetics and affinities of Ad26K and Ad48K with CD46. Calculation of the
predicted binding energies for the energy minimised fiber-knob: CD46 models are compared on the
bar chart (A), lower Kcalmol^{-1} values indicate a stronger interaction, the red line at 3.0 KcalMol^{-1},
denotes an interface energy which can be considered negligible (random proteins passing in
solution). The HI loop (red) of the Ad5 fiber-knob (orange) is seen to extend between CD46 (grey)
and the DG loop (B). The antibody competition inhibition assay (C) shows the relative inhibitory
ability of the Ad5, 35, 26, and 48, fiber-knob domains in CD46 expressing CHO-BC1 cells, with the
calculated IC$_{50}$ values shown in boxes. IC$_{50}$ curves are fitted by non-linear regression. Error bars
represent standard deviation of 3 biological replicates.

Figure 1

Figure 2

Ad26 Fiber-Knob Domain (6FJN)

Ad48 Fiber-Knob Domain (6FJQ)

A

C

Trimer Side-on

B

D

Trimer Towards virion

E

Figure 3

	HI	DG	GH	IJ
A Aligned loops used in this study				
B Ad26K (6FJN)				
C Ad48K (6FJQ)				

Figure 4

Figure 5

Figure 6

Figure 7

A Ad11

B Ad35

C
```
          ....|....|....|....|....|....|....|....|....|....|....|....|....|....|....|....|
Ad5K   396 NNDKLTLWTTPAPS-PNCRLNA---EKDAKLTLVLTKCGSQILATVSVLAVKGSLAPISGTVQ------SAHLIIRFDEN
Ad11K  128 DDNINTLWTGVNPTEANCQIMNSSESNDCKLILTLVKTGALVTAFVYVIGVSNNFNMLTTH-----RNINFTAELFFDST
Ad26K  183 KDDRRTLWTTPDTS-PNCKMST---EKDSKLTLTLTKCGSQVLGNVSLLAVTGEYHQMTAT-----TKKDVKISLLFDEN
Ad35K  128 KDSINTLWTGINPP-PNCQIVENTNTNDGKLTLVLKNGGLVNGYVSLVGVSDTVNQMFTQ-----KTANIQLRLYFDSS
Ad48K  177 NNDKLTLWTTPDPS-PNCKIDQ---DKDSKLTFVLTKCGSQILANMSLLVVKGKFSMINNKVNGTDDYKKFTIKLLFDEK

          ....|....|....|....|....|....|....|....|....|....|....|....|....|....|....|....|
Ad5K   466 GVLLNNSFLDPEYWNFRNGDLTEGTAYTNAVGFMPNLSAYPKSHGKT----------AKSNIVS-QVYL---NGDKTKP
Ad11K  193 GNLLTRLSSLKTPLNHKSGQNMATGAITNAKGFMPSTTAYPFNDNSRE----------KENYIYGTCYYTA-SDRTAFP
Ad26K  244 GILLPSSSLSKDYWNYRSDDSIVSQKYNNAVPFMPNLTAYPKPSAQNA----------KNYSRTKIIS-NVYL---GALTYQP
Ad35K  202 GNLLTEESDLKIPLKNKSSTAT-SETVASSKAFMPSTTAYPFNTTTRD----------SENYIHGICYYMTSYDRSLFP
Ad48K  253 GVLLKDSSLDKEYWNYRSNNNNVGSAYEEAVGFMPSTTAYPKPPTPPTNPTTPLEKSQAKNKYVS-NVYL--GGQAGNP

          ....|....|....|....|....|....|....|....|....|....|....|....|....|....|....|...
Ad5K   532 VTLTITLNGTQETGDTTPSAYSMSFSWDWSGHNYINE-----IFATSSYTFSYIAQE-----
Ad11K  261 IDISVMLNRRAIND---ETSYCIRITWSWNTGDAPEVQTSATTLVTSPFTFYYIREDD----
Ad26K  313 VIITIAFNQETENG----CAYSITFTFTWQ-KDYSAQ-----QFDVTSFTFSYLTQENKDKD
Ad35K  270 LNISIMLNSRMISS----NVAYAIQFEWNLNASESPESN--IATLTTSPFFFSYITEDDN-
Ad48K  329 VATTVSFNK--ETG----CTYSITFDFAWN-KTYENV-----QFDSSFLTFSYIAQE-----
```

D Ad26

E Ad48

Figure 8