Global metabolomic characterizations of *Microcystis* spp. highlights clonal diversity in natural bloom-forming populations and expands metabolite structural diversity

Séverine Le Manach[1], Charlotte Duval[1], Arul Marie[1], Chakib Djediat[1], Arnaud Catherine[1], Marc Edery[1], Cécile Bernard[1], Benjamin Marie[1,*]

[1]UMR 7245 MNHN/CNRS Molécules de Communication et Adaptation des Micro-organismes, Muséum National d'Histoire Naturelle, 75005 Paris, France.

slemanach@mnhn.fr; cduval@mnhn.fr; armarie@mnhn.fr; djediat@mnhn.fr; arnocat@mnhn.fr; medery@mnhn.fr; cbernard@mnhn.fr; bmarie@mnhn.fr

* to whom correspondence should be addressed at bmarie@mnhn.fr

Abstract

Cyanobacteria are photosynthetic prokaryotes that are able to synthetize a wild rang of secondary metabolites exhibiting noticeable bioactivity, comprising toxicity. *Microcystis* represents one of the most common cyanobacteria taxa constituting the intensive blooms that arise nowadays in freshwater ecosystems worldwide. They produce numerous cyanotoxins (toxic metabolites), which are potentially harmful to Human health and aquatic organisms. In order to better understand the variations in cyanotoxins production between clones of the *Microcystis*, we investigate the diversity of several strains isolated from the same blooms, from different populations in various geographical area.

Twenty-four clonal strains were compared by genotyping with 16S-ITS fragment sequencing and metabolites chemotyping using LC ESI-qTOF mass spectrometry. While, genotyping can only discriminate between the different species, the global metabolomes reveal clear discriminant molecular profiles between strains. These profiles can be clustered primarily according to their global metabolite content, then to their genotype, and finally to their sampling localities. A global molecular network of all metabolites highlights the production of a wide set of chemically diverse metabolites, comprising only few microcystins, but many aeruginosins, cyanopeptolins and microginins, along with a large set of unknown molecules. They represent the molecular biodiversity that still remain to be investigated and characterized at their structure as well as at their potential bioactivity or toxicity levels.

Keywords: cyanobacteria blooms, secondary metabolites, chemiodiversity, mass spectrometry, aquatic environment.

1. Introduction

The frequency and the intensity of cyanobacteria blooms occurring in continental aquatic ecosystems have increased since last decades, due to climate and anthropogenic changes (Carey *et al.*, 2012; Sukenik *et al.*, 2015; Paerl, 2018). These massive cyanobacteria blooms threaten the functioning of aquatic ecosystems through various processes, including an alteration of the trophic network, a decrease of the light penetration within the water column, the decrease of available dissolved oxygen, and also the production of various secondary metabolites potentially toxic for the organisms living in these ecosystems (Carmichael 2008). Indeed, various cyanobacteria genera are able to synthetize a wild range of secondary metabolites (Welker *et al.*, 2012; Shih *et al.*, 2013), with noticeable bioactivity, comprising high toxicity, and which are potentially harmful to Human health and aquatic organisms (Codd et al., 2005; Pearson *et al.*, 2010). These metabolites are also believed to be noticeably involved in the capability of these organisms to proliferate in various environments (Guljamow *et al.*, 2017).

One of the most pervasive bloom-forming cyanobacteria in worldwide freshwater ecosystems is *Microcystis*, as it is now encountered and nowadays proliferate locally in more than 108 countries and on all continents (Sejnohova and Marsalek, 2012; Harke *et al.*, 2016; Ma *et al.*, 2016). Previous documentations also noted that *Microcystis* would occurred during previous decades in only less than 30 countries (Zurawell *et al.*, 2005), suggesting that *Microcystis* is currently proliferating and dominating phytoplankton communities in a wide range of freshwater ecosystems in both temperate and tropical climates. In temperate systems, this organism overwinters in the benthos and during the summer rises to the epilimnion where it can accumulate to form blooms and scums on the water surface (Harke *et al.*, 2016).

Some important features of *Microcystis*, such as buoyancy regulation, storage strategy at the bottom of water column, phosphate (P) and nitrogen (N) uptake capacities and resistance to zooplankton grazing, favour its worldwide spread (Pearl *et al.*, 2011). Indeed, *Microcystis* presents competitive advantages over other cyanobacteria or microalgae in response to nutrient limitation. In addition, many *Microcystis* strains can produce a multitude of bioactive secondary metabolites, including the potent hepatotoxin microcystins (MCs), and thus persistent blooms pose a risk to those who use impaired water resources for drinking water supplies, recreational activities, and fisheries (Codd 2005). Beyond MCs, other toxic compounds or potential secondary metabolites produced by *Microcystis* have also been reported as chemical warfare against grazing of herbivores (Harke *et al.*, 2017).

So far, eleven gene clusters encoding non-ribosomal peptide synthase (NRPS) and/or polyketide synthase (PKS), and two ribosomal ones predicted to be involved in the biosynthesis of secondary metabolites, were found among ten *Microcystis* genomes (Humbert et *al.*, 2013). Seven of these clusters encode enzymes for the biosynthesis of known metabolites (microcystins, aeruginosins, cyanopeptolins, microginins, anabaenopeptins, cyanobactins and microviridins), whereas the six remaining clusters encode enzymes for the biosynthesis of still unidentified products. However, the relationship between cyanobacterial biomass and metabolite concentrations in the environment appears neither systematic nor linear (Briand *et al.* 2002; Liu *et al.*, 2016). Indeed, the production of metabolites, such as microcystins, by *Microcystis* blooms, depends not only on cyanobacterial biomass, but also on the ratio between potentially producing and non-producing genotypes within the population (Via-Ordorika *et al.* 2004).

Despite significant advances in description of the biosynthetic pathways involved in cyanobacterial metabolite production (Dittmann *et al.*, 2015; Wang *et al.*, 2014), the natural functions and the ecological roles played by these molecules are still not well understood

(Holland *et al.*, 2013; Zak and Kosakowska, 2016). The biosynthesis of cyanobacterial secondary metabolites consumes a great deal of metabolic energy inducing a significant cost for the cell (Briand *et al.*, 2012). However, natural environment are colonized by various clones exhibiting different corteges of metabolites synthesized (Briand *et al.*, 2009). It has been proposed that the environment it self could favour the selection of *Microcystis* clones that present the metabolite composition that is the more adapted to face the local ecological conditions (Welker *et al.*, 2007; Martins *et al.*, 2009; Agha and Quesada 2014).

Recently, the development of modern approach on mass spectrometry data treatment by molecular networking tools gave us new opportunity for the description of the occurrence and the diversity of cyanobacterial metabolites (Yang *et al.*, 2013; Briand *et al.*, 2016a). With the aim of contributing to better understand the variations in metabolite production, such as microcystins (MCs), between clones of cyanobacterial blooms from different localities, we investigate in the present study the clonal diversity of several *Microcystis* strains isolated from different freshwater bloom-forming populations from various geographical area using such innovative approach based on high resolution mass spectrometry analyses.

2. Results

2.1. Morphologic and phylogenetic characterization

In order to elucidate the genetic relation between the 24 *Microcystis* strains we have investigated here (Table 1), we performed a first analysis of 1380-pb 16S fragment indicates that all *Microcystis* morpho-species are grouped in a unique and homogenous group, due to high sequence conservation on this fragments (not shown). Using 16S-23S ITS fragments (above 600-bp long), the phylogenetic analysis shows a clear distinction between *M. aeruginosa* and *M. wessenbergii/viridis* morpho-species (Figure 2). Interestingly, the different strains presenting the MC synthesis gene *mcyA* (indicated in red) appear not clustered on the phylogenetic representation, suggesting that be ability of producing MC would be a character disconnected from the phylogeny of strains.

2.2. Global metabolome analyses

The metabolomic shotgun analyses reveal discriminant metabolic profiles between strains collected from both different or identical sites. Whereas previous works had highlighted the metabolic diversity of some *Microcystis* strains based on few identified metabolites or cyanotoxines (Martins *et al.*, 2009; Welker *et al.*, 2004; Welker *et al.*, 2006), we present here a global picture of the metabolome of each strain. Using HR ESI-Qq-TOF with the 24 *Microcystis* strain extracts, 2051 distinct mass ions in a range of 400–2000 Da were recorded (*i.e.* with a signal to noise ratio in excess of 6, and respective relative peak intensity superior to 5000-count in at least one sample threshold), each strain exhibiting between above 100 and 300 different main metabolites of reliable (>10 000 counts) respective intensity (Figure 3; supplementary figure S1). A hierarchical clustering was performed according to Bray-Curtis indexes calculated between all 24 strains according to the relative intensity of each analytes in each strains. This representation (Figure 3) clearly shows a clustering of all strains producing MCs (*mcyA*+/MC+, in red), on one side, and of other strains not producing MCs (*mcyA*-/MC-, in grey). This clustering shows that some strains from the same environment exhibits very similar metabolite fingerprints (*e.g.* PMC 728.11 and 729.11), when other from the same localities exhibits much more dissimilar metabolite fingerprints (*e.g.* PMC 728.11 and 727.11), being more similar with strains from faraway locations (*e.g.* PMC 729.11 and 816.112). Additional non-metric multidimensional scaling (nMDS) and PERMANOVA analyses based on Bray-Curtis index indicates that the ability to produce MC seems to be the

139 first main driver of the global metabolome of *Microcystis* molecular fingerprinting, the
140 species and the localities representing less explicative parameters (supplementary figure 2).
141
2.3. Metabolite molecular network
A molecular network based on the global fragmentation pattern profile of all observed
metabolites in the 24 strains investigated. The GNPS algorithm automatically compares all
MS/MS spectra by aligning them one by one, grouping identical molecules (presenting
identical mass and fragmentation pattern) and assigning cosine score ranking from 0 to 1 to
each alignment, allowing network reconstruction of the link between each molecule according
to the cosine score links between all molecules with a cosine score significance threshold set
to 0.6. The resulting network constituted of a total of 925 nodes from the 1374 different
analytes which MS/MS data have been obtained (supplementary figure 1), represents a
starting point for the annotation of unidentified metabolites, according to respectively
identified molecules within the different clusters, and to their occurrence in *M. aeruginosa*
and/or *M. wesenbergii/viridis* strains (Figure 4).
154
The *Microcystis* strains produce a wide set of chemically diverse metabolites, which principal
clusters can attemptedly been identified thanks to analytical standards available for some
cyanobacterial secondary metabolite family, or to match with components from libraries
publically available from GNPS platform, such as HMDB, NIST14 or METLIN. Considering
the strains producing these analytes, and above a quarter of them appears to be specific of *M.*
wessenbergii/viridis strains, when more than half are specific of *M. aeruginosa*, the rest being
observed on both species.
The grouping of different analytes in the same molecular cluster is based on similarity of their
fragmentation patterns, each cluster being potentially specific of the structure of chemical
families. Among those larger clusters, we were thus able to annotate some of them, being
constituted by ions of small metabolites, such as di-peptides and small peptides (1 cluster in A
area), of microcystins (3 clusters in C area), of anabaenopeptins (3 clusters in D area), of
aeruginosins (2 clusters in E area), of aerucyclamides (3 clusters in F area), of microginins (2
clusters in H area) and cyanopeptolons (6 clusters in I area), together with various clusters of
unknown components, comprising non-identified ions (for example 2 clusters in B area). Less
than a third of the metabolites observed here could be annotated, thanks to their respective
mass and fragmentation patterns when compare to those of the above 800 metabolite of
freshwater cyanobacteria described so far and listed in supplementary table 1. These un-
identified ions that belong to annotated clusters are then considered as potential new
analogues of their respective molecular family.
175
2.4. Known cyanobacteria secondary metabolite clusters
Microcystins
Microcystins are cyclic heptapeptides that have been firstly described from *M. aeruginosa*.
Above 250 different variants have been described so far (Catherine *et al.*, 2017), 138 being
references in our database for cyanobacterial metabolite (supplementary table 1). They are
characterized by the presence of a non-proteinacous amino acid in position 5 (Adda), two
amino acid derived from Asp and Glu in position 3 and 6, respectively, and 2 very variable
positions (2 and 4), that serve as reference to the name of the variant. Three microcystis
clusters were highlighted according to the presence of 5 standard molecules (Dmet(Asp3)-
MC-LR, MC-LR, MC-YR, MC-LA and MC-HtyR) analyses in parallel of the 24 *Microcystis*
extracts with the same protocol (Figure 5). Other components of these clusters correspond to
ions presenting a match of their respective mass with those of other MC variants previously
described (supplementary table 1), or for ¼ of them to potential new analogues. Observation

of their respective MS/MS spectra shows that they present distinct but similar fragmentation
patterns of other known MC variants.
191

Aeruginosins
Aeruginosins constitute a linear tetrapeptide family that have been firstly described from *M.*
aeruginosa, and that represent above 94 different variants that have been described so far
(Supplementary table 1). Their MS/MS fragmentation patterns are often characterized by the
presence of a Choi fragment (immonium with 140.109 *m/z*) and other recurrent fragments
from Hpla or Pla. Their composition is rather variable and the component of this family
exhibit masses comprised between 430 and 900 Da (Welker and von Döhren 2006). The
molecular network obtained from the 24 *Microcystis* strains exhibits 2 aeruginosin clusters
(Figure 6) that were highlighted by to the presence of 2 standard molecules (aeruginosin 98A
and 98B). Other components of these clusters correspond to ions presenting a mass match
with other variants of aeruginosin previously described (supplementary table 1), or for ½ of
all this compounds to potential new analogues, that aim at being characterized now, in further
dedicated works.
205

Anabaenopeptins
Anabaenopeptins constitutes a very diverse family of cyclic hexapeptides that have been
described from now from *Microcystis*, *Planktothrix*, *Anabaena*, *Aphanizomenon* and *Nostoc*.
Above 75 different variants have been described so far (supplementary table 1). They are
characterized by the presence of a peptide bound between the D-Lys placed in position 2, and
the carboxylic group of the amino acid placed in position 6. Except fro the D-Lys (position 2)
all other positions are variable allowing a large structural diversity of the family which
molecules exhibit masses between 750 and 950 Da (Welker and von Döhren 2006). Three
anabaenopeptin clusters were highlighted here (Figure 7) according to the presence of 4
standard molecules (anabaenopeptin A, B, E and oscyllamide Y). Other components of these
clusters correspond to ions presenting a mass with other variants previously described
(supplementary table 1), or for ½ of all of them to compounds that very likely correspond to
potentially new analogues. All observed anabaenopeptin compounds are from *M. aeruginosa*
strains suggesting that *M. wessenbergii/viridis* strains are not capable of the synthesis of
molecules of this family and may not possess the corresponding *apt* synthetic gene cluster.
221

Cyanopeptolins
Cyanopeptolins belong to a large family of cyclic depsipeptides, that also comprises
micropeptins and aeruginopeptins, representing above 170 variants. Those molecules are
characterized by the presence of the non-proteinaceous amino acid Ahp and by a six-aa long
ring formed by an ester bound between Thr or Pro in position 1 and the carboxylic group of
the N-terminal amino acid (position 6). The lateral chain can exhibit a variable length and is
constituted by one or two amino acid and potentially linked to an aliphatic fatty acid (Welker
and von Döhren 2006). In the molecular network, 2 analytes of the cyanopeptolin clusters
correspond to 2 standard molecules (cyanopeptolin B and D), and various other components
correspond to ions presenting a mass that corresponds to different variants previously
described (supplementary table 1), allowing us to annotate them as cyanopeptolin specific
clusters (Figure 8). Above ½ of the analytes present in these clusters correspond to unknown
compounds constituting potential new analogues. We observe here that all these
cyanopeptolin compounds are from *M. aeruginosa* strains suggesting that *M.*
wessenbergii/viridis strains are not capable of the synthesis of molecules of this family and
may not possess the corresponding *mcn/oci* synthetic gene cluster.
238

Microginins

Microginins are linear pentapeptides (the length of the sequence varying from 4 to 6 amino acids) initially identified from *Microcystis aeruginosa*, then from other species and in other genera such as *Planktothrix* (Welker and von Döhren 2006). These molecules are composed by a characteristic non-proteinaceous amino acid Ahda in N-terminy, the other position bearing variable amino acid structures, comprising Tyr, Pro Hty, Trp, Ala, Ser…. Relatively few microginin variants (less than 40) have been described so far (supplementary table 1). According to the molecular network, above two third of the analytes present in the two microginin clusters correspond to unknown compounds constituting potential new analogues, when six standard molecules could have been retrieved in our analysis (microginin 757, 711, BN578, FR1, FR2 and SD755).

Aerucyclamides/cyanobactins

The name "cyanobactin" has been proposed in order to grouped all cyclic peptides containing proteinogenous amino acids that are post-translationally modified in heterocyclic amino acids and isoprenoid derivatives (Sivonen *et al.*, 2010). It comprises various cyclamides (cyclic peptides of 6 amino acids) that have been identified in freshwater cyanobacteria such as *Microcystis*, *Planktothrix* and *Nostoc*, but also in symbiotic cyanobacteria species. More than 30 variants have been described so far, but more molecules could be related to the family that represent a very large variety of chemical structures (Martins and Vasconcelos 2015). Three different cyanobactin clusters were observed in the molecular network, comprising three aerucyclamide standard molecules (aerucyclamide A, B and C) that were identified by GNPS tool. Above ¾ of the analytes from these clusters representing potential new analogues, that need to be characterized by further dedicated analyses.

2.5. Uncharacterized cyanobacteria metabolite clusters
Potential primary metabolite clusters?

Various other important clusters comprising few tens of analytes, that are mostly present in high amount (higher peak intensity showed by larger circles) in a large set of strains from both *M. aeruginosa* and *M. wessenbergii/viridis* (Figure 4). These compounds present molecular masses comprised between 300 and 500 Da, suggesting that it could correspond to relatively small compounds (supplementary figure 3). We can speculate, that these components could correspond to primary metabolites used for the general metabolism of various strains, however additional effort should be made in order to propose annotations for these ubiquitous molecules present in these specific clusters. Interestingly, the compounds from these latter cluster present high fragmentation patterns, illustrated with various similarity and high cosine score (numerous and tick links between nodes), suggesting they should present very similar structures.

Uncharacterized secondary metabolite clusters?

Interestingly, other clusters correspond to unknown components, some of them being only synthesis by *M. wessenbergii/viridis* as exemplified in supplementary figures 4 and 5. These compounds which molecular mass are relatively high (between 950 and 1300 Da) might correspond to specific secondary metabolite families, belonging to other known family of cyanobacterial metabolite, but poorly characterized, such as microviridins and/or aeruginoguanidins, for which only few variants have been characterized and no standard molecules are available so far. Alternatively, they may correspond to completely new family of metabolites, which existence have been suggested by the observation of orphan NRPS/PKS clusters within various *Microcystis* genomes (Humbert *et al.*, 2013).

3. Discussion

3.1. Phylogeny and biogeography of Microcystis strains

The classification of species in genus *Microcystis* is still under large revision. Traditional morphological criteria including colony form, mucilage structure, cell diameter, the density and organization of cells within the colony, pigment content and life cycles are used for morphospecies recognition (Komarek 2016). Five dominant morphospecies (*M. aeruginosa, M. ichthyoblabe, M. viridis, M. novacekii, M. wesenbergii*) of the genus *Microcystis* were suggested by Ostuka *et al.* (2001). However, with the development of molecular and biochemical markers, some contradictory results of *Microcystis* taxonomy have been found. For instance, 16S rRNA analysis revealed no differences among the morpho-species (Otsuka *et al.*, 2000). In consideration of both morphological and molecular markers, it has been suggested to classify *Microcystis* into three groups: the small cell-size group composed of *M. ichthyoblabe* and *M. flos-aquae*, the middle cell-size group based on *M. aeruginosa* (incl. *M. novacekii*) and the large cell-size group represented by *M. wesenbergii* (Whitton, 2012).

According to your phylogenetic reconstruction based on 16S-16S/23S ITS fragment that is in accordance with previous observation (Otsuka *et al.*, 1999), the studied *Microcystis* strains were roughly divided into two groups; the *M. wesenbergii/viridis* group and the *M. aeruginosa* group, in relation with the relative size of the colonies observed under microscopes. In addition, the *mcyA+* and *mcyA-* strains appears broadly disperse on both group, unspecifically. These observations confirm that the phylogenetic relationship between different strains within the genus of *Microcystis* do not correspond to the possession of *mcy* (Tillett et al., 2001). Therefore, the toxic potential, through the MC synthesis, of each strain should assess regardless of its phylogenetical position.

3.2. Biogeography and genetic diversity of Microcystis strains

Despite the very high similarity of *Microcystis* 16S rRNA sequences (>99.5%), low synteny and large genomic heterogeneity have been retrieved from the investigation of *Microcystis* genomes, deciphering a large cryptic diversity from various strains collected from different sites and continents (Humbert *et al.*, 2013). At an other large geographic scale that in our analysis, genetic comparison of various *Microcystis* strains isolated from different Asian and European lakes based on 16S-ITS fragments does not show local-specific clustering effect indicating that the genetic distance between different genotypes from the same lake can be greater than between strains from very distanced environments (Humbert *et al.*, 2005; Haande *et al.*, 2007). Our data on the molecular heterogeneity observed by metabolic fingerprinting between some strains originating from the same site (for example between PMC 810.12 and PMC 816.12, Champs-sur-Marne, France) also support this hypothesis. The fact that some *Microcystis* genotypes seem to be distributed worldwide was also previously reported for bacterioplankton species (Zwart *et al.*, 1998), suggesting that these organisms may possess peculiar physiological capabilities and/or large dispersal capabilities enabling them to compete successfully in a wide range of freshwater environments.

3.3. MC-producing versus non-MC producing metabolic pattern of Microcystis strains

The metabolome diversity has been also used as molecular characters that can help at the discrimination of various chemotypes (Ivanizevic *et al.*, 2011), in addition to classical genotyping approach that sometimes lack of reliable characters for phylogenetic relationship discrimination. Few analytical methods have been experimented for chemotaxonomic characterization of cyanobacteria, based for example on fatty acid compositions (Gugger *et al.*, 2002), or more recently on ribosomal protein analysed globally by MALDI-TOF in *Microcystis aeruginosa* (Sun *et al.*, 2016). Interestingly, this latter chemotaxonomic approach

was able to group all the MC-producing strains in two distinct clades when the non MC-
producing strains were segregate in three other distinct clades. In a previous work, Martins
and collaborators (2009) have analysed the metabolite diversity of various *Microcytis*
aeruginosa strains originated from Portuguese water supplies using MALDI-TOF MS, and
were able to observe a MC production in almost half of the strains investigated. These
observations also illustrate the fact that MC-producing clones could subsist in various
environments, despite the important energetic cost required for MC gene cluster replication
and the translation of its mega-enzyme complex (Briand *et al.*, 2012). The biological
advantage of producing MCs for some clones still remains an enigma, as the functional role of
MC remaining uncharacterized (Agha and Quesada 2014; Gan *et al.*, 2012).
In a similar manner our molecular fingerprint approach based on global metabolome profiling
using ESI-Qq-TOF discriminate clearly MC-producing strains from others. Taken together,
these observations suggest that shotgun mass spectrometry chemotyping of cyanobacteria
could constitute a promising tool for the characterization of rapid biomarkers aim at the
toxicological assessment of strains isolated from the field. The MC production could
constitute a singular treat, that could constitute on of the key drivers of the global metabolic
diversity of *Microcystis* strains, suggesting that MC could play a keystone function in
cyanobacterial metabolite production. Indeed, it has previously hypothesized according to
metabolomic observation that the character of MCs production could be compensate in strains
not producing MCs by the production of other metabolites, such as aeruginosamines (Martins
et al., 2009) for unknown biological reasons (Briand *et al.*, 2016a; Tonk *et al.*, 2009). Such
secondary metabolic compensatory mechanisms, between and within peptide classes, were
previously suspected for *Microcystis* (Martins *et al.*, 2009) or *Planktothrix* (Tonk *et al.*, 2005)
in response to various growth conditions. However, further investigations with wider
sampling are now required in order to increase the data set that could better help to test such
metabolite functional hypothesis.
365
3.3. Secondary metabolite diversity within known metabolite families
The molecular cluster identified by GNPS approach can be annotated thank to match with
spectral databases or the presence of standard molecules present in the request as additional
samples analysed similarly (Yang *et al.*, 2013). In our hands, the global molecular networks
obtained from the 24 strains MS/MS dataset present various cluster that could have been
annotated accordingly as corresponding to main cyanobacterial metabolite families. We
assumed that the nodes of these clusters which molecular masses is similar to already known
cyanobacterial metabolite very likely correspond to these specific metabolites or alternatively
to isobaric analogues from the same family. In addition, all other nodes from those clusters
that do not correspond to neither standard, nor known analogues, could be considered as new
analogues that may correspond to new putative variants that still have to be investigated.
These observations are in accordance with previous investigation indicating that different
Microcystis strains can produce such various known and unknown secondary metabolites,
according to both genetic or targeted metabolome analyses (Humbert *et al.*, 2013; Martins *et*
al., 2009; Welker *et al.*, 2004; Welker *et al.*, 2006). Surprisingly, rare metabolite family such
as aeruginosamine are not successfully detected in our analysis. Indeed, only few molecules
belonging to this family have been yet described (supp. Table 1) and their MS/MS
fragmentation pattern has not been deeply characterized. In addition, the lack of available
standard molecule and of knowledge on their respective fragmentation patterns makes
aeruginosamines challenging to be simply annotate with our GNPS approach.
386
However, our analyses reveal the large molecular diversity of *Microcystis* metabolites,
according to the various new variants of both cyanobacterial metabolite families that remain

to be characterized. The observation of various uncharacterized cluster also suggest that new
metabolite families are needed to be discovered and described from this taxa, and that further
effort are still required. So far, *Microcystis* represent one of the most studied genera for its
production of various metabolite families. However, the biological functions play by these
molecules remains enigmatic and their overgrowing molecular diversity revealed by global
approach, such as GNPS global metabolomic investigation, constitutes one of the questioning
paradox in the field of evolution and diversity microbiology.
396
3.4. Unknown metabolite families
Although, the strains used in this study were not cultured in stringent axenic conditions, no
noticeable contamination by fungi or heterotroph bacteria could have been detected during the
systematic screening of all strains under light microscope prior to the experiment. In addition,
a previous metabolome analyses in PCC 7806 grown under axenic or non-axenic condition do
not detected any variation in the metabolite produced by the cyanobacteria (Briand *et al.*,
2016b). We assume that the metabolite profiles observed here for the 24 strains are
characteristic of the cyanobacteria them self and that the different metabolite observed in this
study, comprising the unknown metabolite clusters highlighted by the network analysis, are
genuine produced by the cyanobacteria themselves.
The non-annotated cluster observed in our GNPS analysis can potentially correspond to novel
variant of known cyanotoxin or to completely new family of cyanobacteria metabolite.
Indeed, Humbert and co-workers (2013) have shown that the genome of ten *Microcytis* strains
exhibits at least three orphan clusters with specific NRPS/PKS signature that are virtually
synthesizing so far undescribed metabolite family. One could speculate that the unknown
clusters observed with GNPS approach can correspond to such novel metabolite family, and
structural elucidation of an expending number of novel metabolites revealed by molecular
networking are currently been performed on various cyanobacteria (Boudreau *et al.*, 2015).
415
416
4. Conclusions
Innovative approaches based on shotgun metabolomic analyses using high resolution mass
spectrometry, as those performs in this study, seems to provide a large panel of information
on cyanobacteria chemical diversity relevant for evolutive, ecological and toxicological
purposes, and represents an interesting and relatively easy-to-perform alternative to genome
sequencing for metabolite and/or toxic potential descriptions of cyanobacterial strains.
Global molecular network also allows to depict the chemical diversity of the *Microcystis*
metabolome in an interesting manner, with comparison with classical natural product
chemistry approaches described so far, as in our hand above half of the analytes described in
the global molecular network seems to correspond to metabolites belonging to potentially new
variants of known families or even to family members (presenting original fragmentation
patterns) that are still to be described at the structural and toxicological/bioactivity levels.
429
430
5. Material and methods
6.1. Sampling, isolation and cultivation of Microcystis *monoclonal strains*
The study was carried out with 24 monoclonal non-axenic cultures of *Microcystis* spp.
maintained at 25°C in 15-mL vessels with Z8 media in the PMC (Paris Museum Collection)
of living cyanobacteria (http://www.mnhn.fr/fr/collections/ensembles-collections/ressources-
biologiques-cellules-vivantes-cryoconservees/microalgues-cyanobacteries). Larger volume of
all strains was simultaneously cultivated during one month in triplicates in 50 mL
Erlenmeyer's vessels at 25°C using a Z8 medium with a 16 h: 8 h light/dark cycle (60
$\mu mol.m^{-2}.s^{-1}$). All trains were investigated for their MC production by Adda-microcystin

AD4G2 ELISA kit (Abraxis). Cyanobacterial cells were centrifuged (at 4,000 g for 10 min),
then freeze-dried and weighted, and stored at -80°C prior to DNA and metabolite analyses.
442
6.2. DNA-extraction, PCR, sequencing and Phylogenetic analyses
DNA was extracted with Qiagen Kit (Cat N°69506) according to manufacturer's instructions.
Presence and condition of the extracted DNA was confirmed by observing the 260/280-nm
ratio and the absorbance spectra between 200 and 800 nm using a nanodrop
spectrophotometer (Safas, Monaco). PCR reaction was performed with *mcyA* specific primers
developed for *Microcystis* (*mcyA*_S AAAAACCCGCGCCCTTTTAC and *mcyA*_AS
AGGCAGTTGGAGAATCACGG) in order to investigate the presence of this gene in the
different strains. In parallel, the region containing a fragment 16S rRNA end of the 16S-23S
ITS was amplified using primer couples previously described in Gugger and Hoffman (2004)
and Iteman *et al.* (2000), respectively. The amplification was done in a reaction mixture
containing 0.1 µL (100 µM) of each primer, 12.5 µL MyTaq Red Mix polymerase (Bioline®)
and 2 µL (~200 ng) DNA sample. Final volume of the reaction mixture was 25 µL. The PCR
product was sequenced (Genoscreen, France) using the same primers. The partial 16S-ITS
sequences (above 1980-bp long) of all strains were deposited to GenBank (accession numbers
xxxxx-xxxx).
The *Microcystis* 16S-23S ITS gene sequences were compared to a selection of similar (>93%
identity) sequences retrieved from NCBI based on a standard nucleotide BLAST search (basic
local alignment search tool). The sequences were aligned with CodonCode Aligner and non-
homologous regions of the sequence alignment were manually deleted in BioEdit (Version
7.2.5). The phylogeny of the edited, aligned 16S-23S ITS sequences was performed in the
program MEGA (Version 6). The trees based on maximum likelihood were constructed with
1000 bootstrap replicates, with the branch lengths iterated and global rearrangements done.
465
6.3. Metabolome biomass extraction and analysis by mass spectrometry
The 20 mL of biomasses of the 24 *Microcystis* strain cultures were centrifuged (4,000 rpm, 10
min), the culture media discarded, and then freeze-dried. The lyophilized cells were weighted
then sonicated 2 min in acetonitrile/methanol/water (40/40/20) acidified at 0.1% of formic
acid with a constant ratio of 100 µL of solvent for 1 mg of dried biomass, centrifuged at 4°C
(12,000 g; 5 min). Two µL of the supernatant were then analyzed on an UHPLC (Ultimate
3000, ThermoFisher Scientific) coupled with a mass spectrometer (ESI-Qq-TOF Maxis II
ETD, Bruker).
Ultra high performance liquid chromatography (UHPLC) was performed on 2 µL of each of
the metabolite extracts using a Polar Advances II 2.5 pore C_{18} column (Thermo®) at a 300
µL.min^{-1} flow rate with a linear gradient of acetonitrile in 0.1% formic acid (5 to 90% in 21
min). The metabolite contents were analyzed in triplicate for each strain using an electrospray
ionization hybrid quadrupole time-of-flight (ESI-QqTOF) high resolution mass spectrometer
(Maxis II ETD, Bruker) on positive simple MS or on positive autoMSMS mode with
information dependent acquisition (IDA), on the 50-1500 *m/z* rang at 2 Hz or between 2-16
Hz speed, for MS and MS/MS respectively, according to relative intensity of parent ions, in
consecutive cycle times of 2.5 s, with an active exclusion of previously analysed parents. The
data were analyzed with the DataAnalysis 4.4 and MetaboScape 3.0 software for internal
recalibration (<0.5 ppm), molecular feature search and MGF export. Peak lists were generated
from MS/MS spectra between 1 and 15 min, with a filtering noise threshold at 0.1% maximal
intensity and combining various charge states and related isotopic forms. Metabolite
annotation was attempted according to the precise mass of the molecules and their respective
MS/MS fragmentation patterns with regards to an in-house database of above 700

cyanobacteria metabolites and confirmed with few commercially available standard
molecules analysed similarly in our platform.
491
6.4. Data and statistical analysis
Heatmap representation of the global metabolome of the 24 *Microcystis* spp. monoclonal
strains was performed with Gene-E tool (https://software.broadinstitute.org/GENE-E/) using
the relative quantification (pic area) of 2051 molecular features analysed on HR ESI-Qq-TOF
using MetaboScape 3.0 (Bruker) with a >5000 counts and 400-2000 Da threshold,
considering peak presents in at least 3 different trains and in at least 6 consecutive MS scans.
Then, the hierarchical clustering was performed according to Bray-Curtis distance method.
NMDS and PERMANOVA analyses were performed using MicrobiomeAnalyst platform
(http://www.microbiomeanalyst.ca/) in order to investigate the influence of the species, the
sampling localities and of the production of MCs, described as the variables, on the global
metabolite distribution of the global metabolome observed on ESI-Qq-TOF for the 24 strains.
Using the whole MSMS data (converted in mgf format) obtained for the 24 strains taken
together, a molecular network was created using the online workflow at Global Natural
Products Social molecular networking (GNPS) (http://gnps.ucsd.edu) (Wang *et al.*, 2016).
The data were then clustered with MS-Cluster with a parent mass tolerance of 1.0 Da and an
MS/MS fragment ion tolerance of 0.5 Da to create consensus spectra. Consensus spectra that
contained less than one spectrum were discarded. A network was then created where edges
were filtered to have a cosine score above 0.6 and more than five matched peaks. Further
edges between two nodes were kept in the network only if each of the nodes appeared in each
other's respective top 10 most similar nodes. The spectra in the network were then searched
against the GNPS spectral libraries. All matches kept between network spectra and library
spectra were required to have a score above 0.6 and at least five matched peaks. The clustered
spectra of the network were annotated by comparing monoisotopic mass to our in-house
cyanobacteria metabolite databases according to MS and MS/MS fragmentation pattern
matches. Molecular networks were visualized using Cytoscape 3.2.1.
517

6. Acknowledgements
This work was supported by grants from the CNRS Défi ENVIROMICS "Toxcyfish" project
and from the ATM "Les micro-organismes, acteurs clef des écosystèmes" of the MNHN to
Dr. Benjamin Marie. We would like to thank the French minister for the research for their
financial supports Séverine Le Manach. The MS spectra were acquired at the Plateau
technique de Spectrométrie de Masse Bio-organique, UMR 7245 Molécules de
Communication et Adaptation des Micro-organismes, Muséum National d'Histoire Naturelle,
Paris, France.

Author contributions
SLM, ME, AC, CB and BM conceived and designed the experiments; CD isolated all new
strains of the PMC; CD, SLM, AM, CDJ performed the analysis; SLM, CD, AM, SZ and BM
treated the data. All authors wrote and reviewed the manuscript.

Conflict of interest
The authors declare no conflict of interest.
534

535 **7. References**
536 Agha, R., Quesada, A. (2014) Oligopeptides as biomarkers of cyanobacterial subpopulations.
537 Toward an understanding of their biolgical role. *Toxins* 6, 1929-50.
538 Boudreau, P., Monroe, E., Mehrotra, S., Desfor, S., Korabeynikov, A., Sherman, D., Murray,

T., Gerwick, L., Dorrestein, P., Gerwick, W. (2015) Marine cyanobacterium *Moorea producens* JHB through orthogonal natural products workflows. *PLoS ONE* 10:e0133297.

Briand, E., Escoffier, N., Straub, C., Sabart, M., Quiblier, C., Umbert, J-F. (2009) Spatiotemporal changes in the genetic diversity of a bloom-forming *Microcystis aeruginosa* (cyanobacteria) population. *ISME J.* 3, 419-429.

Briand, E., Bomans, M., Quiblier, C., Saleçon, M-J., Humbert, J-F. (2012) Evidence of the cost of the production of Microcystins by *Microcystis aeruginosa* under different light and nitrate environmental conditions. *PLoS ONE* 7:e29981.

Briand, E., Bormans, M., Gugger, M., Dorrestein, PC., Gerwick, W. (2016a) Changes in secondary metabolic profiles of *Microcystis aeruginosa* strains in response to intraspecific interactions. *Environmental Microbiology* 18:384-400.

Briand, E., Humbert, J-F., Tambosco, K., Bormans, M., Gerwick, W; (2016b) Role of bacteria in the prodcution and degradation of *Microcystis* cyanopeptides. *Microbiology Open* 3:343.

Carey, C.C., Ibelings, B.W., Hoffmann, E.P., Hamilton, D.P., Brookes, J.D. (2012) Eco-physiological adaptations that favour freshwater cyanobacteria in a changing climate. *Water Res.* 46, 1394-1407. doi:10.1016/j.watres.2011.12.016

Carmichael, W. (2008) Cyanobacterial Harmful Algal Blooms: State of the Science and Research Needs. *Adv. Exp. Med. Biol.* 619, 831‑53. doi:10.1007/978-0-387-75865-7

Catherine, A. , Bernard, C. , Spoof, L., Bruno, M. (2017). Microcystins and Nodularins. In Handbook of Cyanobacterial Monitoring and Cyanotoxin Analysis (eds J. Meriluoto, L. Spoof and G. A. Codd). doi:10.1002/9781119068761.ch11

Codd, G.A., Morrison, L.F., Metcalf, J.S. (2005) Cyanobacterial toxins: risk management for health protection. *Toxicol. Appl. Pharmacol.* 203, 264-72. doi:10.1016/j.taap.2004.02.01

Dittmann, E., Gugger, M., Sivonen, K., Fewer, D.P. (2015) Natural Product Biosynthetic Diversity and Comparative Genomics of the Cyanobacteria. *Trends Microbiol.* 23, 642-652. doi:10.1016/j.tim.2015.07.008

Gan, N., Xiao, Y., Zhu, L., Wu, Z., Liu, J., Hu, C., Song, L., 2012. The role of microcystins in maintaining colonies of bloom-forming *Microcystis* spp. *Environ. Microbiol.* 14, 730-742. doi:10.1111/j.1462-2920.2011.02624.x

Gugger, M., Hoffmann, L. (2004) Polyphyly of true branching cyanobacteria (Stigonematales). *Int J Syst Evol Microbiol* 54, 349–57.

Gugger, M., Lyra, C., Suominen, I., Tsitko, I., Humbert, J-F., Salkinoja-Salonen, M., Sivonen, K. (2002) Cellular fatty acids as chemotaxonomic makers of the genera *Anabaena*, *Aphanizomenon*, *Microcystis*, *Nostoc* and *Planktothrix*. *Int. J. System. Evol. Microbiol.* 52, 1007-1015.

Guljamow, A., Kreische, M., Ishida, K., Liaimer, A., Altermark, B., Bähr, L., Hertweck, C., Ehwald, R., Dittmann, E. (2017) High-density cultivation of terrestrial *Nostoc* strains leads to reprogramming of secondary metabolome. *Appl Environ Microbiol.* 83 :e01510-17.

Haande, S., Ballot, A., Rohrlack, T., Fastner, J., Wiedner, C., Edvardsen, B. (2007) Diversity of *Microcystis aeruginosa* isolates (Chroococcales, Cyanobacteria) from East-African water bodies. *Arch. Microbiol.* 188, 15-25. doi:10.1007/s00203-007-0219-8

Harke, M.J., Steffen, M.M., Gobler, C.J., Otten, T.G., Wilhelm, S.W., Wood, S.A., Paerl, H.W. (2016) A review of the global ecology, genomics, and biogeography of the toxic cyanobacterium, *Microcystis* spp. *Harmful Algae* 54, 4-20. doi:10.1016/j.hal.2015.12.007

Harke, M.J., Jankowiak, J.G., Morrell, B.K., Gobler, C.J. (2017) Transcriptomic Responses in
the Bloom-Forming Cyanobacterium *Microcystis* Induced during Exposure to
Zooplankton. *Appl Environ Microbiol.* 15;83(5).

Holland, A., Kinnear, S. (2013) Interpreting the possible ecological role(s) of cyanotoxins:
compounds for competitive advantage and/or physiological aide? *Mar. Drugs* 11,
2239-58. doi:10.3390/md11072239

Humbert, J-F., Duris-Latour, D., Le Berre, B., Giraudet, H., Salençon, M.J. (2005) Genetic
diversity in *Microcystis* populations of a French storage reservoir assessed by
sequencing of 16S-23S rRNA intergenic spacer. *Microbiol Ecology* 49, 308-314.

Humbert, J-F., Barbe, V., Latifi, A., Gugger, M., Camteau, A., Coursin, T., Lajus, A.,
Castelli, V., Oztas, S., Samson, G., Longin, C., Medigue, C., Tandeau de Marsac, N.
(2013) A tribute to disorder in the genome of the bloom-forming freshwater
cyanobacterium *Microcystis aeruginosa. PLoS ONE* 8: e70747.

Iteman, I., Rippka, R., Tandeau de Marsac, N., et al. (2000) Comparison of conserved
structural and regulatory domains within divergent 16SrRNA–23S rRNA spacer
sequences of cyanobacteria. *Microbiology* 146:1275–86.

Ivanisevic, J., Thomas, O., Lejeune, C., Chavaldonné, P., Perez, T. (2011) Metabolic
fingerprinting as an indicator of biodiversity: towards understanding inter-specific
relationships among Homoscleromorpha sponges. *Metabolomics* 7: 289-304.

Komárek, J. (2016). A polyphasic approach for the taxonomy of cyanobacteria: principles and
applications. *European Journal of Phycology*, 51(3), 346-353.

Liu, Y., Xu, Y., Wang, Z., Xiao, P., Yu, G., Wang, G., Li, R. (2016) Dominance and
succession of *Microcystis* genotypes and morphotypes in Lake Taihu, a large and
shallow freshwater lake in China. *Environ. Pollut.* 219, 399-408.
doi:10.1016/j.envpol.2016.05.021

Ma, J., Qin, B., Paerl, H.W., Brookes, J.D., Hall, N.S., Shi, K., Zhou, Y., Guo, J., Li, Z., Xu,
H., Wu, T., Long, S. (2016) The persistence of cyanobacterial (*Microcystis* spp.) blooms
throughout winter in Lake Taihu, China. *Limnol. Oceanogr.* 61, 711-722.
doi:10.1002/lno.10246

Martins, J., Saker, M.L., Moreira, C., Welker, M., Fastner, J., Vasconcelos, V.M. (2009)
Peptide diversity in strains of the cyanobacterium *Microcystis aeruginosa* isolated from
Portuguese water supplies. *Appl. Microbiol. Biotechnol.* 82, 951-961.

Martins, J., Vasconcelos, V. (2015) Cyanobactins from cyanobacteria: Current genetic and
chemical state of knowledge. *Mar. Drugs* 13, 6910-6946. doi:10.3390/md13116910

Otsuka, S., Suda, S., Shibata, S., Oyaizu, H., Matsumoto, S., Watanabe, M.M (2001) A
proposal for the unification of five species of the cyanobacterial genus *Microcystis*
Kützing ex Lemmermann 1907 under the rules of the Bacteriological Code. *Int J Syst*
Evol Microbiol. 51(Pt 3), 873-9.

Otsuka, S., Suda, S., Li, R., Matsumoto, S., Watanabe, M.M. (2000) Morphological
variability of colonies of Microcystis morphospecies in culture. *J Gen Appl Microbiol.*
46(1), 39-50.

Otsuka, S., Suda, S., Li, R., Watanabe, M. (1999) Phylogenetic relationships between toxic
and non-toxic strains of the genus *Microcystis* based on 16S to 23S internal transcribed
spacer sequence, *FEMS Microbiol. Lett.* 172 (1),15–21.

Paerl, HW. (2018) Mitigating Toxic Planktonic Cyanobacterial Blooms in Aquatic
Ecosystems Facing Increasing Anthropogenic and Climatic Pressures. *Toxins* 8;10(2).

Paerl, H.W., Hall, N.S., Calandrino, E.S. (2011) Controlling harmful cyanobacterial blooms
in a world experiencing anthropogenic and climatic-induced change. *Sci. Total Environ.*
409, 1739-45. doi:10.1016/j.scitotenv.2011.02.001

Pearson, L., Mihali, T., Moffitt, M., Kellmann, R., Neilan, B. (2010) On the chemistry,

toxicology and genetics of the cyanobacterial toxins, microcystin, nodularin, saxitoxin
and cylindrospermopsin. *Mar. Drugs* 8, 1650-80. doi:10.3390/md8051650
Šejnohová, L., Maršálek, B. (2012) *Microcystis*, in: Ecology of Cyanobacteria II: Their
Diversity in Space and Time. Springer Netherlands, Dordrecht, p. 195-228.
doi:10.1007/978-94-007-3855-3_7
Shih, P.M., Wu, D., Latifi, A., Axen, S.D., Fewer, D.P., Talla, E., Calteau, A., Cai, F.,
Tandeau de Marsac, N., Rippka, R., Herdman, M., Sivonen, K., Coursin, T., Laurent, T.,
Goodwin, L., Nolan, M., Davenport, K.W., Han, C.S., Rubin, E.M., Eisen, J. a, Woyke,
T., Gugger, M., Kerfeld, C. (2013) Improving the coverage of the cyanobacterial phylum
using diversity-driven genome sequencing. *Proc. Natl. Acad. Sci. U. S. A.* 110, 1053-8.
doi:10.1073/pnas.1217107110
Sivonen, K., Leikoski, N., Fewer, D.P., Jokela, J. (2010) Cyanobactins-ribosomal cyclic
peptides produced by cyanobacteria. *Appl. Microbiol. Biotechnol.* 86, 1213-1225.
doi:10.1007/s00253-010-2482-x
Sukenik, A., Quesada, A., Salmaso, N. (2015) Global expansion of toxic and non-toxic
cyanobacteria: effect on ecosystem functioning. Biod*ivers. Conserv.* 4:889-908.
Sun, L-W., Jiang, W-J., Sato, H., Kawachi, M., Lu, X-W. (2016) Rapid classification and
identification of *Microcystis* strains using MALDI-TOF MS and polyphasic analysis.
PLoS ONE 11: e0156275.
Tillett, D., Parker, D. L., & Neilan, B. A. (2001). Detection of toxigenicity by a probe for the
microcystin synthetase A gene (mcyA) of the cyanobacterial genus *Microcystis*:
comparison of toxicities with 16S rRNA and phycocyanin operon (phycocyanin
intergenic spacer) phylogenies. *Applied and environmental microbiology*, *67*(6), 2810-
2818.
Tonk, L., Visser, P.M., Christiansen, G., Dittmann, E., Snelder, E.O.F.M., Wiedner, C., Mur,
L.R., Huisman, J. (2005) The microcystin composition of the cyanobacter-ium
Planktothrix agardhii changes towards a more toxic variant with increasing light
intensity. *Appl. Environ. Microbiol.* 71, 5177–5181.
Tonk, L., Welker, M., Huisman, J., Visser, P.M. (2009) Production of cyanopeptolins,
anabaenopeptins, and microcystins by the harmful cyanobacteria Anabaena 90 and
Microcystis PCC 7806. *Harmful Algae* 8, 219-224. doi:10.1016/j.hal.2008.05.005
Via-Ordorika, L., Fastner, J., Kurmayer, R., Hisbergues, M., Dittmann, E., Komarek, J.,
Erhard, M., Chorus, I. (2004) Distribution of microcystin-producing and non-
microcystin-producing *Microcystis* sp. in European freshwater bodies: detection of
microcystins and microcystin genes in individual colonies. *Syst Appl Microbiol.* 27(5),
592-602.
Wang, H., Fewer, D.P., Holm, L., Rouhiainen, L., Sivonen, K. (2014) Atlas of nonribosomal
peptide and polyketide biosynthetic pathways reveals common occurrence of
nonmodular enzymes. *Proc Natl Acad Sci U S A.* 111(25), 9259-64.
Welker, M., Brunke, M., Preussel, K., Lippert, I., von Döhren, H. (2004) Diversity and
distribution of *Microcystis* (cyanobacteria) oligopeptide chemotypes from natural
communities studies by single-colony mass spectrometry. *Microbiology* 150, 1785-1796.
doi:10.1099/mic.0.26947-0
Welker, M., Von Döhren, H. (2006) Cyanobacterial peptides - Nature's own combinatorial
biosynthesis. *FEMS Microbiol. Rev.* 30, 530-563. doi:10.1111/j.1574-6976.2006.00022.x
Welker, M., Maršálek, B., Šejnohová, L., von Döhren, H. (2006) Detection and identification
of oligopeptides in *Microcystis* (cyanobacteria) colonies: Toward an understanding of
metabolic diversity. *Peptides* 27, 2090-2103. doi:10.1016/j.peptides.2006.03.014
Welker, M., Ejnohová, L., Némethová, D., von Döhren, H., Jarkovsky, J., Marsálek, B.(2007)
Seasonal shifts in chemotype composition of *Microcystis* sp. communities in the pelagial

and the sediment of a shallow reservoir. *Limnol. Oceanogr.* 52, 609-619.
doi:10.4319/lo.2007.52.2.0609
Welker, M., Dittmann, E., Von Döhren, H. (2012) Cyanobacteria as a source of natural
products. *Methods Enzymol.* 517, 23-46. doi:10.1016/B978-0-12-404634-4.00002-4
Whitton, B.A. (2012) Ecology of Cyanobacteria II, Ecology of Cyanobacteria II: Their
Diversity in Space and Time. Springer Netherlands, Dordrecht. doi:10.1007/978-94-007-
3855-3
Yang, J.Y., Sanchez, L.M., Rath, C.M., Liu, X., Boudreau, P.D., Bruns, N., Glukhov, E.,
Wodtke, A., De Felicio, R., Fenner, A., Wong, W.R., Linington, R.G., Zhang, L.,
Debonsi, H.M., Gerwick, W.H., Dorrestein, P.C. (2013) Molecular networking as a
dereplication strategy. J. *Nat. Prod.* 76, 1686-1699. doi:10.1021/np400413s
Zak, A., Kosakowska, A. (2016) Cyanobacterial and microalgal bioactive compounds-the role
of secondary metabolites in allelopathic interactions. *Oceanol. Hydrobiol. Stud.* 45,
131-143. doi:10.1515/ohs-2016-0013
Zurawell, R.W., Chen, H., Burke, J.M., Prepas, E.E. (2005) Hepatotoxic cyanobacteria: a
review of the biological importance of microcystins in freshwater environments. *J.*
Toxicol. Environ. Health. B. Crit. Rev. 8, 1-37. doi:10.1080/10937400590889412
Zwart, G., Hiorns, W.D., Methe, B.A., van Agterveld, M.P., Huismans, R., Nold, S.C., Zehr,
J.P., Laanbroek, H.J. (1998) Nearly identical 16S rRNA sequences recovered from lakes
in North America and Europe indicate the existence of clades of globally distributed
freshwater bacteria. *Syst Appl Microbiol.* 21(4):546-556.
709

Legends of figures

Table 1: List of *Microcystis* spp. strains used in this study, their area of origin, the ELISA MC screening, the *mcyA* gene presence and their respective 16S-ITS sequence accession numbers. Stains isolated from the same sample collected the same day from different French area are indicated with: [a] = Villerest (2008); [b] = Varennes-sur-Seine (2008); [c] = Eure et Loire (2010); [d] = Valence (2011); [e] = Champs –sur-Marne (2012).

Figure 1: Microcystis *spp.* General view of a representative intense *Microcystis* sp. bloom in a recreational pound (Champs-sur-Marne, © B. Marie) (A). Macrograph of *Microcystis* colonies at surface water (© B. Marie) (B). Example of 15-mL vessels containing the monoclonal strains of *Microcystis* spp. maintained in the Paris' Museum Collection (PMC) of cyanobacteria (MNHN, Paris, © C. Duval) (C). Example of micrograph of the isolated monoclonal culture of the *Microcystis aeruginosa*, where scale bare represents 10 μm (© C. Duval) (D). Representative picture of *Microcystis aeruginosa* cell from PMC 156.02 strain (here in division) under transmission electron microscope, where scale bare represents 0.5 μm (© C. Djediat) (E). General structures of various cyanobacterial metabolites belonging to the microcystin, anabaenopeptin, microginin microviridin, aeruginosin and oscillatorin families (F).

Figure 2: Maximum likelihood (ML) tree based on partial 16S-23S ITS sequences. The sequences obtained in this study are indicated in bold. The strains, which exhibit *mcyA* gene, are indicated in red. Other sequences were retrieved from GenBank, accession numbers in brackets. Bootstrap values >60% are shown at the nodes. The scale bar indicates number of nucleotide substitutions per site. The *Microcystis* sp. AICB832 was used as an out-group.

Figure 3: Heatmap representation of the metabolome of the 24 *Microcystis* spp. monoclonal strains analysed using HR ESI-TOF, representing 2051 different analytes (present in at least three strains, with minimal peak intensity > 5000 counts) identified by MetaboScape software. The hierarchical clustering between strains was performed according to Bray-Curtis distance method. Blue and green squares indicate *M. aeruginosa* and *M. wesenbergii/viridis*, respectively. Sampling localities are: C=Champs-sur-Marne; B=Burkina Faso; E=Eure et Loire; Ne=Netherlands; Sc=Scotland; Se=Senegal; So=Souppes-sur-Loin; Var=Varennes-sur-Seine; Val=Valence; Vi=Villerest.

Figure 4: Molecular network generated from MS/MS spectra from the 24 *Microcystis* strains using GNPS tool (all data and results are freely available on the GNPS server at the address http://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=c017414365e84334b38ae75728715552). The nodes of the analytes detected in *M. aruginosa* or *M. wessenbergii/viridis* strains only are indicated in green and blue, respectively, when analytes detected in both species are indicated in orange. Uncharacterized analytes are indicated by circles constituting potential new analogues. Analytes whom individual masses match with known secondary metabotites from cyanobacteria (listed in supplementary table 1) are indicated as specific shapes. Standard molecules analyses similarly are indicated by a heavy black perimeter. Only cluster regrouping at least 2 analytes are represented.

Figure 5: Microcystins clusters highlighted by the GNPS analysis based on the MS/MS CID fragmentation spectra obtained from the 24 *Microcystis* strains. Analytes detected in *M. aeruginosa* or *M. wessenbergii/viridis* strains only are indicated in green and blue, respectively, when analytes detected in both genera are in orange. Analytes whom individual

masses match with known microcystins are indicated as hexagons. Standard molecules analyses similarly are indicated by a heavy black perimeter. Example of MS/MS spectra and chemical structures are shown for (Asp$_3$)-microcystin-LR, microcystin-YR, -LR, -LF, -LA and -HtyR. Notice that $(M+H)^+$ and $(M+2H)^{2+}$ ions may be grouped in distinct clusters.

Figure 6: Aeruginosin cluster highlighted by the GNPS analysis based on the MS/MS CID fragmentation spectra obtained from the 24 *Microcystis* strains. Analytes detected in *M. aeruginosa* or *M. wessenbergii/viridis* strains only are indicated in green and blue, respectively, when analytes detected in both genera are in orange. Analytes whom individual masses match with known aeruginosins are indicated as scares with right corners. Standard molecules analyses similarly are indicated by a heavy black perimeter. Example of MS/MS spectra and chemical structures are shown for aeruginosin 98A and 98B.

Figure 7: Anabaenopeptin clusters highlighted by the GNPS analysis based on the MS/MS CID fragmentation spectra obtained from the 24 *Microcystis* strains. Analytes detected in *M. aeruginosa* strains are indicated in green. Analytes whom individual masses match with known anabaenopeptin are indicated as scares with round corners. Standard molecules are indicated by a heavy black perimeter. Uncharacterized analytes are indicated by circles constituting potential new analogues. Example of MS/MS spectra and chemical structures are shown for anabaenopeptin A, B, F and Oscillamide Y. Notice that $(M+H)^+$ and $(M+2H)^{2+}$ ions may be represented by different nodes grouped in distinct clusters.

Figure 8: Cyanopeptolin clusters highlighted by the GNPS analysis based on the MS/MS CID fragmentation spectra obtained from the 24 *Microcystis* strains. Analytes detected in *M. aeruginosa* strains are indicated in green. Analytes whom individual masses match with known cyanopeptolin are indicated as parallelepiped. Standard molecules are indicated by a heavy black perimeter. Uncharacterized analytes analyses similarly are indicated by circles constituting potential new analogues. Example of MS/MS spectra and chemical structures are shown for cyanopeptolin A and B.

Figure 9: Microginin clusters highlighted by the GNPS analysis based on the MS/MS CID fragmentation spectra obtained from the 24 *Microcystis* strains. Analytes detected in *M. aeruginosa* strains only are indicated in green and blue, when analytes detected in both genera are in orange. Analytes whom individual masses match with known microginin are indicated as 45°-tilted scares. Standard molecules analyses similarly are indicated by a heavy black perimeter. Uncharacterized analytes are indicated by circles constituting potential new analogues. Example of MS/MS spectra and chemical structures are shown for microginin FR1, FR2, 711 and 757.

Figure 10: Aeruclyclamide clusters highlighted by the GNPS analysis based on the MS/MS CID fragmentation spectra obtained from the 24 *Microcystis* strains. Analytes detected in *M. aeruginosa* or *M. wessenbergii/viridis* strains only are indicated in green and blue, respectively, when analytes detected in both genera are in orange. Analytes whom individual masses match with known aerucyclamides are indicated as octogons. Uncharacterized analytes are indicated by circles constituting potential new analogues. Standard molecules analyses similarly are indicated by a heavy black perimeter. Example of MS/MS spectra and chemical structures are shown for aerucyclamide A, B and D.

808 **Supplementary figures**
809

810 **Supplementary Figure 1**: Representation of the analytes from the 24 *Microcystis* strains
811 analysed by MS simple and MS/MS mode, exhibiting the good representativeness of analytes
812 selected for MS/MS analyses. All analysed ions are represented according to their respective
813 RT and *m/z* ratio. For MS/MS data, the size of the circle being representative of their
814 maximum peak intensity.
815

816 **Supplementary Figure 2**: NMDS analysis of global metabolite patterns of the 24 *Microcystis*
817 spp. monoclonal strains analysed using HR ESI-TOF, with PERMANOVA analyses
818 performed on MicrobiomeAnalyst platform with Bray-Curtis index according to the MC
819 production (left) and to the genera (right). "MC production", "species", and "locality" factor
820 present significant impact on the global metabolome.
821

Supplementary Figure 3: Unknown cluster "1" highlighted by the GNPS analysis based on
the MS/MS CID fragmentation spectra obtained from the 24 *Microcystis* strains. This cluster
of uncharacterized molecules that present high fragmentation similarity main correspond to a
new kind of metabolites that still need to be characterized. Analytes detected in *M.*
aeruginosa or *M. wessenbergii/viridis* strains only are indicated in green and blue,
respectively, when analytes detected in both genera are in orange.
828

829 **Supplementary Figure 4**: Unknown cluster "2" highlighted by the GNPS analysis based on
830 the MS/MS CID fragmentation spectra obtained from the 24 *Microcystis* strains. This cluster
831 of uncharacterized molecules that present high fragmentation similarity main correspond to a
832 new kind of metabolites that still need to be characterized. Analytes detected in *M. aruginosa*
833 and *M. wessenbergii/viridis* strains are indicated in orange.
834

835 **Supplementary Figure 5**: Unknown cluster "3-5" highlighted by the GNPS analysis based
836 on the MS/MS CID fragmentation spectra obtained from the 24 *Microcystis* strains. This
837 cluster of uncharacterized molecules that present high fragmentation similarity main
838 correspond to a new kind of metabolites that still need to be characterized. Analytes detected
839 in *M. wessenbergii/viridis* strains only are indicated in blue.
840

841 **Table 1.**

Strain name	Species	Country	Locality/area	MC ELISA detection	*mcyA* PCR detection	16S-ITS Accession number
PCC 7806	*M. aeruginosa*	Netherland	Braakman	+	+	xxxxxx
PCC 7820	*M. aeruginosa*	Scotland	Balgavies	+	+	xxxxxx
PMC 95.02[a]	*M. aeruginosa*	France	Villerest	-	-	xxxxxx
PMC 98.15[a]	*M. aeruginosa*	France	Villerest	-	-	xxxxxx
PMC 155.02	*M. aeruginosa*	Sénégal	Djoudj	-	-	xxxxxx
PMC 156.02	*M. aeruginosa*	Sénégal	Djoudj	-	-	xxxxxx
PMC 241.05	*M. aeruginosa*	Burkina Faso	Ouahigouya	+	+	xxxxxx
PMC 265.06	*M. aeruginosa*	Burkina Faso	Sian	-	-	xxxxxx
PMC 566.08[b]	*M. wesenbergii/viridis*	France	Varennes sur Seine	-	-	xxxxxx
PMC 567.08[b]	*M. wesenbergii/viridis*	France	Varennes sur Seine	-	-	xxxxxx
PMC 570.08	*M. aeruginosa*	France	Souppes sur Loing	-	-	xxxxxx
PMC 671.10[c]	*M. wesenbergii/viridis*	France	Eure et Loire	-	-	xxxxxx
PMC 672.10[c]	*M. wesenbergii/viridis*	France	Eure et Loire	-	-	xxxxxx
PMC 673.10[c]	*M. wesenbergii/viridis*	France	Eure et Loire	-	-	xxxxxx
PMC 674.10[c]	*M. wesenbergii/viridis*	France	Eure et Loire	-	-	xxxxxx
PMC 679.10[c]	*M. aeruginosa*	France	Eure et Loire	+	+	xxxxxx
PMC 727.11[d]	*M. aeruginosa*	France	Valence	-	-	xxxxxx
PMC 728.11[d]	*M. aeruginosa*	France	Valence	+	+	xxxxxx
PMC 729.11[d]	*M. aeruginosa*	France	Valence	+	+	xxxxxx
PMC 730.11[d]	*M. aeruginosa*	France	Valence	-	-	xxxxxx
PMC 807.12[e]	*M. wesenbergii/viridis*	France	Champs sur Marne	+	+	xxxxxx
PMC 810.12[e]	*M. aeruginosa*	France	Champs sur Marne	-	-	xxxxxx
PMC 816.12[e]	*M. aeruginosa*	France	Champs sur Marne	+	+	xxxxxx
PMC 826.12[e]	*M. aeruginosa*	France	Champs sur Marne	-	-	xxxxxx