# α-Synuclein binds extracellular complex N-linked glycans

Melissa Birol[1], Slawomir P. Wojcik[2], Andrew D. Miranker[2] and Elizabeth Rhoades*[2]

[1]Department of Chemistry, University of Pennsylvania, 231 South 34th Street, Philadelphia, PA 19104

[2]Department of Molecular Biophysics and Biochemistry, Yale University, 266 Whitney Avenue, New Haven CT 06511

**Abstract**

Fibrillar deposits of α-Synuclein (αS) are the hallmark of Parkinson's disease. Recent evidence suggests that disease may propagate through cell-to-cell transmission of toxic forms of αS. Relevant to this mechanism, here we report the discovery that αS binds extracellular N-linked glycans. We find that αS binds to proteoliposomes derived from neuronal-lineage cells and that removal of N-linked glycans both decreases binding and alters the conformation of bound αS. Cellular internalization of both monomer and fibrillar αS is specific to neuronal-lineage cells and is reduced dramatically upon cleavage of extracellular N-linked glycans. Our results suggest that interactions with extracellular glycans may play an important role in cellular uptake of αS relevant both to normal proteostasis as well as to transmission of pathogenic αS species.

**Introduction**

The pathologies of Parkinson's disease (PD) and related synucleinopathies are characterized by the accumulation of aggregates of the neuronal protein α-Synuclein (αS) (Goedert, 2001). The prevailing hypothesis is that toxicity is mediated by prefibrillar oligomers of αS. Emerging evidence suggests that cell-to-cell transmission of toxic αS species underlies disease progression (Guo and Lee, 2014), although molecular details of this process, including the identity of relevant extracellular inter-action partners, are lacking.

αS is a small, soluble protein that is intrinsically disordered in the cytoplasm (Theillet et al., 2016). It binds peripherally to anionic lipid bilayers through its N-terminal domain, which becomes α-helical upon binding (Dikiy and Eliezer, 2014). The localization of αS to nerve terminals (George et al., 1995, Diao et al., 2013) and to cellular lipid raft domains (Fortin et al., 2004) suggests there are components or properties of cellular membranes which may be important for αS binding and function which may not be fully reproduced by simple lipid mixtures. Indeed,

specific components of the extracellular membrane, including proteins (Mao et al., 2016) and

proteoglycans (Ihse et al., 2017), have been identified as having roles in the uptake of pathogenic

αS species. These observations motivated our interest in better understanding αS interactions with

complex cellular membranes.

**Results**

The lipid and protein composition of giant plasma membrane vesicles (GPMVs) closely

resembles the cell plasma membrane (Bauer et al., 2009). GPMVs were harvested from SH-SY5Y

and HEK cells and incubated with monomeric N-terminally acetylated αS labeled with Alexa 488

(αS-AL488) (SI) (Trexler and Rhoades, 2012). We used the N-terminally acetylated form of the

protein as this modification is ubiquitously present on human-derived protein, both in healthy

persons and those with PD (Anderson et al., 2006). αS forms large, bright assemblies on the

exterior of the SH-SY5Y GMPVs and causes their clustering (Figures 1A, C – Figure supplement

1A), while it binds only weakly to HEK GPMVs (Figures 1B, C – Figure supplement 1B). This

observation suggests a specific extracellular membrane component found in SH-SY5Y cells and

absent in HEK cells strongly enhances αS binding.

Both proteins and lipids in the outer plasma membrane are heavily modified by

glycosylation (Moremen et al., 2012) and extracellular glycosoaminoglycans (GAGs) have been

implicated in the cellular uptake of fibrillar forms of αS (Holmes et al., 2013). Carbohydrates were

selectively removed from the extracellular surface of the SH-SY5Y GPMVs by incubation with

endoglycosidases. Treatment with peptide-N-glycosidase F (PNGaseF), which cleaves N-linked

glycans results in a loss of inhomogenous binding and the bright αS assemblies (Figure 1 – Figure

supplement 1C, D), as well as significantly decreasing the amount of bound αS (Figure 1C). By

comparison, bright αS assemblies are still observed after treatment with Heparinase I/III, which

removes GAGs, or endoglycosidase H (EndoH), which cleaves high mannose N-linked

carbohydrates (Figures 1D, E – Figure supplement 1E). Uniform binding of Alexa 488 labeled

concanavalin A (conA-AL488), a lectin which binds to α-mannose structures, to SH-SY5Y

GPMVs demonstrates that glycoproteins are distributed throughout the GPMV bilayer in the

absence of αS (Figure 1 - Figure supplement 1C); the addition of αS results in clustering of conA-

stained proteins (Figure 1 - Figure supplement 1F, G). αS appears to induce assembly through

binding to and cross-linking multiple membrane glycoproteins, likely observed because GPMVs

lack an intact cytoskeleton which would restrict large-scale rearrangement of plasma membrane

proteins.

Figure 1. (A) Images of SH-SY5Y GPMVs incubated with 100 nM αS-AL488 and unlabeled αS (concentrations

indicated). (B) As in (A) but with HEK GPMVs. (C) Binding of 80 nM αS-AL488 to GPMVs quantified by FCS as

loss of protein from solution. The number of molecules (N) in control wells lacking GPMVs is indicated by the red-

dashed line. (D) Representative images of SH-SY5Y GPMVs incubated with 100 nM αS-AL488 and 80 μM unlabeled

αS after pretreatment of GPMVs with the indicated endoglycosidase. (E) Binding of αS-AL488 to Heparinase- and

EndoH-treated SH-SY5Y GPMVs by FCS, as in Figure 1C. Binding +/- PNGaseF from Figure 1C shown for

comparison. The data are presented as mean±SD, n=3. Scale bars=20 μm.

The following figure supplement is available for Figure 1:

Figure supplement 1. αS binding to and clustering of SH-SY5Y GPMVs is dependent on N-linked glycans

Intramolecular FRET measurements were made of αS labeled at residues 9 and 72,
encompassing much of membrane-binding domain of αS. Mean energy transfer efficiencies ($ET_{eff}$)
of αS bound to SH-SY5Y GPMVs, without and with PNGaseF treatment, are 0.43±0.08 and
0.19±0.06, respectively (Figure 2). This lower $ET_{eff}$ is comparable to that of the elongated helix
previously observed by single molecule FRET for αS upon binding to synthetic lipid bilayers
(Trexler and Rhoades, 2009). αS assumes multiple conformations on even simple synthetic lipid
vesicles (Robotta et al., 2011); the relatively broad FRET distributions observed here may reflect
a similar degree of conformational heterogeneity. In the absence of complex N-linked glycans, αS
binds to GPMVs through interactions with the lipid bilayer resulting in a predominantly extended
conformation. In the presence of N-linked glycans, αS binding is enhanced and requires a
conformation distinct from the extended, membrane-bound α-helix.

103 Figure 2 Intramolecular FLIM-FRET measurements of αS bound to untreated or PNGaseF-treated SH-SY5Y GPMVs.
104 Left: Representative image of GPMV, colored by donor fluorophore lifetime shown in the sacle bar. Histogram of
105 $ET_{eff}$ calculated from the images, with Gaussian fits shown. The pixels used to calculate the histograms are indicated
106 by dashed lines on the GPMV images.

Figure 3 (A) Diffusion time of αS-AL488 (80 nM) as a function of PNGaseF-derived glycan concentration. (B) Comparison of αS-AL488 binding to carbohydrates (30 μg) cleaved by the indicated endoglycosidases. C. (left) Percent increase in peak intensity for selected residues strongly enhanced by glycan binding and (right) cross-sections from $^{15}$N-$^{1}$H HSQC spectra along the nitrogen dimension at 8.15 ppm for αS alone (gray) and in the presence of PNGaseF glycans (orange). For A, B and C, data are presented as mean±S.D., n=3. *P<0.01, **P<0.001 and ***P<0.0001 by the Student's T-Test).

The following figure supplement is available for Figure 3:

Figure supplement 2. αS binding to isolated glycans.

To identify whether αS binding to glycans requires either the associated glycoproteins or a lipid bilayer, binding of αS to glycans in solution was measured by fluorescence correlation spectroscopy (FCS). SH-SY5Y cells were treated with each of the three endoglycosidases above, and the cleaved carbohydrates were isolated. The carbohydrates were titrated into αS-AL488 and the autocorrelation curves were fit to extract the diffusion time and average number of fluorescent molecules, N (Figure 3 - Figure supplement 2A). The diffusion time, which reflects the hydrodynamic size of the diffusing species, of αS increases more than 25% with increasing concentrations of PNGaseF-cleaved glycans, comparable to conA-AL488 (Figures 3A, B – Figure

supplement 2B). In contrast to GPMV images, there is no evidence of glycan-mediated assembly

of αS in solution (Figure 1 - Figure supplement 2C) which may reflect a difference in αS binding

to disperse glycans in solution compared to the relatively high density of glycans on the GPMV

surface (Figure 1A). The addition of carbohydrates obtained from EndoH or Heparinase treatment

of SH-SY5Y cells or PNGaseF treatment of HEK cells results in minimal changes in the diffusion

time of αS by FCS (Figure 3B – Figure supplement 2D). NMR measurements show non-uniform

glycan dependent changes in αS peak intensity in the presence of PNGaseF-cleaved glycans

(Figures 3C – Figure supplement 2E). By a filtration-based assay, both monomer and pre-formed

fibrillar (PFF) αS are found to pull-down PNGaseF-cleaved glycans from solution (Figure 3 -

Figure supplement S2F). The specific association of αS with SH-SY5Y-derived complex N-linked

glycans both in solution (by FCS, NMR and filtration) and on GPMVs (by imaging and FRET)

led us to probe a possible role in cellular internalization of αS.

SH-SY5Y and HEK cells were incubated with monomer or PFF αS-AL488 and uptake of

protein was monitored (SI). Both forms show time-dependent internalization in SH-SY5Y, but not

HEK, cells (Figure 4 – Figure supplement 3A, B). In SH-SY5Y cells, monomer and PFF αS appear

as puncta, which co-localize with an endosomal marker (Figure 4 – Figure supplement 3A, C) and

uptake is inhibited at 4°C, indicating that active endocytotic pathways is required (Figure 4 - Figure

supplement 3D). Clathrin-dependent endocytosis was assessed in SH-SY5Y and HEK cells using

transferrin. Both cell lines showed rapid uptake of transferrin, indicating that the αS results are not

due to inherent differences in clathrin-dependent endocytosis rates between cell lines (Figure 4 –

Figure supplement 3E).

Carbohydrate groups were selectively removed from the extracellular surface of SH-SY5Y

cells without an impact on clathrin-dependent endocytosis (Figure 4 – Figure supplement 4A, B).

After 12 h incubation with αS, a decrease in the number of intracellular puncta relative to untreated

cells is observed, both for monomer (WT and mutants) and PFF forms, reflecting a significant

decrease in uptake by deglycosylated cells (Figure 4A - Figure supplement 4C, D). Quantification

of the kinetics of depletion of monomer and PFF αS from cell media (Figures 4B, C - Figure

supplement 4D) as well as of the kinetics of internalization (Figure S4D) confirm that αS uptake

is significantly slowed in PNGaseF-treated cells. These observations were validated in

cultured primary hippocampal neurons. Both monomer and PFF αS are readily internalized, and

removal of extracellular N-linked glycans by PNGaseF results in a 10-fold decrease in the amount

of internalized αS (Figure 4D), without causing defects in clathrin-dependent endocytosis (Figure

4 – Figure supplement 4B). Under our measurement conditions, neither untreated nor PNGaseF-

treated cells show evidence of increased toxicity upon incubation with monomer or PFF αS (Figure

4 – Figure supplement 4E).

Figure 4 (A) Uptake of monomer or PFF αS-AL488 by SH-SY5Y cells, -/+ PNGaseF treatment quantified by puncta

analysis. (B) Loss of αS-AL488 loss from extracellular medium of SH-SY5Y cells measured by FCS. (C) As in B,

but for monomer and PFF αS and quantified by PAGE. (D) Uptake of monomer and PFF αS-AL488 by mouse

hippocampal neurons, -/+ PNGaseF treatment quantified by total cellular fluorescence. All αS uptake measurements

with 200 nM monomer αS-AL488 or 200 nM αS$^{PFF}$ (monomer units, 20:1 αS:αS-AL488) ; n=100 cells, 3 independent

experiments, *P<0.01, **P<0.001 and ***P<0.0001 by the Student's T-Test).

The following figure supplement is available for Figure 4:

Figure supplement 3. Endocytosis of αS.

Figure supplement 4. PNGaseF treatment does not disrupt clathrin-mediated endocytosis.

Figure supplement 5. N-terminal acetylation of αS is required for its interaction with GPMVs and purified glycans.

Figure supplement 6. Heparinase and EndoH treatment of SH-SY5Y cells does not disrupt uptake of monomer αS.

## Discussion

The data presented in this manuscript provide compelling evidence that αS interacts

specifically to extracellular complex, N-linked glycans on SH-SY5Y cells and primary neurons.

αS binds to N-linked glycans in solution and on cell-derived proteoliposomes. Uptake of both

monomer and PFF αS by SH-SY5Y cells and primary neurons is dependent upon N-linked glycans.

While N-linked glycans are present on all mammalian cells, the glycome differs significantly, in

the specific sugars and in their distributions, between cell types. The selectivity of αS for N-linked

glycans from SH-SY5Y cells and primary neurons over those from HEK cells can be attributed to

its ability to differentiate between the specific complex sugars. N-linked glycans have not

previously been identified as binding partners of αS, nor as important to its cellular uptake. While

internalization of αS PFFs has garnered significant attention because of the potential role in

spreading of aggregates in disease (Guo and Lee, 2014) much less attention has been given to

monomer αS. However, as αS is secreted even under non-pathogenic conditions (Emmanouilidou

et al., 2010), re-uptake of monomer αS by neurons may be important for normal homeostasis.

Critical to our discovery is our use of N-terminally acetylated αS. Measurements using the

unmodified protein show that it does not bind to complex N-linked glycans on GPMVs or in

solution, nor does deglycosylation have an effect on its uptake by SH-SY5Y cells or primary

neurons (Figures 1, 2, 3 and 4 – Figure supplement 5). In contrast to other protein modifications,

N-terminal acetylation occurs co-translationally and is irreversible (Starheim et al., 2012). It is the

predominant modification found on human αS making N-terminally acetylated αS (Anderson et

al., 2006) of greater physiological significance than protein lacking this modification. For many

proteins, e.g. tropomyosin binding to actin (Hitchcock-DeGregori and Heald, 1987), N-terminal

acetylation is required for recognition of binding partners (Drazic et al., 2016), and our current

work suggests that this may be true for αS and N-linked glycans. While the fact that both monomer

and PFF αS bind to N-linked glycans is initially non-intuitive, high-resolution structures of fibrillar

αS show that its N-terminus remains free and disordered (Tuttle et al., 2016). Thus the acetylated

N-terminus is accessible in both monomer and PFF αS for glycan-binding. Although here we do

not observe a specific role for GAGs in the uptake of monomer, acetylated αS (Figure 4 – Figure

supplement 6), prior studies found that they enhance uptake of unmodified fibrillar αS (Ihse et al.,

2017, Holmes et al., 2013). Presumably, these interactions are primarily electrostatic, driven by

positive residues in the β-sheet core of αS fibers, and thus distinct from the N-terminal driven

binding of complex N-linked glycans.

One consequence of our observation is that it elicits a reconsideration of putative binding

partners for αS. Many of the proteins identified as such contain N-linked glycosylation sites,

including glucocerebrosidase (Yap et al., 2011), Rab3b (Cooper et al., 2006) and LAG3 (Mao et

al., 2016), although neither the glycosylation state of these proteins, nor the N-terminal acetylation

of αS have been explicitly considered in prior studies. For lectins, high affinity binding occurs

when the correct cluster of glycans is present and in the correct orientation. Although to our

knowledge there are no examples of entirely disordered lectins, one advantage would be con-

formational flexibility allowing for recognition and binding of N-linked glycans with non-ideal

orientations. It has been shown that both the length and stiffness of the disordered linkers between

carbo-hydrate binding domains in enzymes which degrade cell plant-wall polysaccharides are

important for orientation of active sites (Rozycki and Cieplak, 2016). While the molecular details

of the interaction between αS and complex, N-linked glycans remain to be determined, the results

presented here identify a novel class of binding partners for αS with potential implications in

normal function and disease transmission.

**Material and Methods**

**αS expression and purification.** αS was expressed in *E. coli* BL21 cells; for N-terminally acetylated αS, BL21 stocks containing the NatB plasmid with orthogonal antibiotic resistance were used. The purification of both acetylated (αS) and unmodified (αS$_{un}$) protein was carried out as previously described [S](Trexler and Rhoades, 2012), with minor modifications. Briefly, two ammonium sulfate cuts were used (0.116 g/mL and 0.244 g/mL) with αS precipitating in the second step. The pellet was resolubilized in Buffer A (25 mM Tris pH 8.0, 20 mM NaCl, 1 mM EDTA) with 1 mM PMSF and dialyzed against Buffer A to remove ammonium sulfate. Dialyzed samples were then loaded to an anion exchange column (GE HiTrap Q HP, 5ml) and eluted with a gradient to 1 M NaCl. αS elutes at approximately 300 mM NaCl. Fractions containing αS were pooled and concentrated using Amicon Ultra concentrators (3000 Da MWCO). Concentrated samples were then loaded to a size exclusion column (GE HiLoad 16/600 Superdex75) and eluted at 0.5 ml/min. Fractions containing αS were again pooled and concentrated, then stored at -80°C. All αS constructs used in this work were checked by MALDI to confirm correct mass and presence of acetylation (Figure supplement 7).

For NMR measurements, 15N-labeled αS was grown in E. coli BL21 stocks containing the NatB plasmid in minimal medium (6 g/L Na$_2$HPO$_4$.7H$_2$O, 3 g/L KH$_2$PO$_4$, 0.5 g/L NaCl, 1mM MgS0$_4$, 300 μM CaCl$_2$, 0.5 g/L $^{15}$NH$_4$Cl) instead of LB medium and purified as described above.

**αS labeling.** αS was site-specifically labeled at a single position by introduction of a cysteine at either residue 9 or residue 130. For labeling reactions, freshly purified αS (typically 200–300 μL of ~200 μM protein) was incubated with 1 mM DTT for 30 min at room temperature to reduce the cysteine. The protein solution was passed over two coupled HiTrap Desalting Columns (GE Life Sciences) to remove DTT and buffer exchanged into 20 mM Tris pH 7.4, 50 mM NaCl, 6 M guanidine hydrochloride (GdmCl). The protein was incubated overnight at 4 °C with stirring with a 4x molar excess of Alexa Fluor 488 (AL488) or Alexa Fluor 594 (AL594) maleimide (Invitrogen). The labeled protein was concentrated and buffer exchanged into 20 mM Tris pH 7.4, 50 mM NaCl using an Amicon Ultra 3K Concentrator (Millipore), with final removal of unreacted dye and remaining GdmCl by passing again over a set of coupled desalting columns equilibrated with 20 mM Tris pH 7.4, 50 mM NaCl.

For dual fluorophore labeling for FRET measurements, cysteines were introduced at
residues 9 and 72. The protein was labeled as described above, with the following modifications.
The protein was first incubated with donor fluorophore AL488 maleimide at a ratio of protein:dye
of 1:0.5 for 2 h at room temperature with stirring. Then, a 4x molar excess of acceptor fluorophore
AL594 maleimide (Invitrogen) was added, and the reaction was continued overnight at 4 °C. The
labeled protein was separated from unreacted dye as described above. αS labeled at these positions
have been extensively studied in our lab; as documented in our previous publications, they do not
perturb αS binding to lipid membranes and serve as excellent reporters of different conformations
of membrane-associated αS(Trexler and Rhoades, 2009, Middleton and Rhoades, 2010).

**Fibril Formation.** αS pre-formed fibrils (PFFs) were prepared as previously described (Volpicelli-
Daley et al., 2014). Briefly, 100 μM αS was mixed with 5 μM αS-AL488 in 20 mM Tris pH 7.4
and 100 mM NaCl. To induce aggregation, this solution was incubated at 37°C for 5 days with
agitation (1500 rpm on an IKA MS3 digital orbital shaker) in parafilm-sealed 1.5 mL Eppendorf
tubes to ensure minimal solvent evaporation. The aggregation reaction was analyzed by Congo
Red absorbance by diluting 10 μl of the aggregation solution in 140 μl 20 μM Congo Red. The
mature fibrils were then pelleted by centrifugation (13,200 rpm for 90 mins at 4°C) and the
supernatant removed. Fibers were resuspended in an equal volume (relative to supernatant) of 20
mM Tris pH 7.4, 100 mM NaCl. Mature fibers were subsequently fragmented on ice using a
sonicator (Diagenode Biorupter™ UCD-300 bath sonicator set to high, 30s sonication followed
by a delay period of 30s, 10 min total) to form PFFs.

**Assessment of fibrillar material** TEM and PAGE were used to characterize fibrillar αS (Figure
supplement 7). For TEM, 10 μL of aggregated protein samples (from both before and after
sonication) were incubated on TEM grids (400-mesh Formvar carbon coated copper, Electron
Microscopy Sciences) for 1–2 minutes. Sample solution was wicked with filter paper and grid was
washed with water to remove any excess material and improve background contrast. Grids were
then incubated with 1% (w/v) aqueous uranyl acetate (UA) solution (10 μL) for 30–60 s. Excess
UA was wicked away with filter paper and grids were air dried. TEM images were collected using
a JOEL JEM 1011 TEM (operating voltage 100 kV) equipped with a charge-coupled device
camera (ORIUS 832. 10W; Gatan). For PAGE, aggregated protein solutions were centrifiuged to
pellet the aggregated material. The upernatant was removed and pellet was resuspended in the

starting volume of buffer. Both supernatant and resuspended pellet (20 µL) were loaded on a 4-

12% polyacrylamide gel. Gels were imaged using a Typhoon FLA7000 gel imager (GE Life

Sciences) using Coomassie stain mode.

**Cell culture.** Human neuroblastoma (SH-SY5Y) and human embryonic kidney 293T (HEK) cells

were grown at 37 °C under a humidified atmosphere of 5% $CO_2$. The SH-SY5Y cells were cultured

in Dulbecco's Modified Eagle's Medium (DMEM) plus 10% fetal bovine serum, 50 U/ml penicillin

and 50 µg/ml streptomycin. The HEK cells were cultured in DMEM supplemented with 10% FBS,

2mM L-glutamine and 100 units/ml penicillin-streptomycin.

Cells were passaged upon reaching ~95% confluence (0.05% Trypsin-EDTA, Life

Technologies), propagated and/or used in experiments. Cells used in experiments were pelleted

and resuspended in fresh media lacking Trypsin-EDTA.

**Primary Neuronal Culture**. Primary neuronal cultures were prepared from E15-E17 embryos of

CD1 mice. All procedures were performed according to the NIH Guide for the Care and Use of

Experimental Animals and were approved by the University of Pennsylvania Institutional Animal

Care and Use Committee (IACUC). Dissociated hippocampal neurons were plated onto sterile,

poly-D-lysine coated on IBIDI chambers at 200,000 cells/coverslip for live cell imaging and were

allowed to mature for 5 days in complete neuronal medium (Neurobasal without phenol red

(Thermo Fisher), 5% B27 supplement (Thermo Fisher)). Medium was partially exchanged every

3-4 days.

**Giant Plasma Membrane Vesicles.** Giant plasma membrane vesicles (GPMVs) are blebs

obtained directly form the cell plasma membrane that contain lipid bilayers and the embedded

membrane proteins, but lack the other biological components of the cell [S](Bauer et al., 2009).

GPMVs were isolated from SH-SY5Y and HEK cells according to established methods [S](Sezgin

et al., 2012)**.** Briefly, cells were plated in 25 $cm^2$ culture flasks and cultured for 48 h**,** washed with

GPMV buffer (10 mM HEPES, 150 mM NaCl, 2 mM $CaCl_2$, pH 7.4) twice and then exposed to

25 mM formaldehyde and 2 mM DTT for 2 h to induce blebbing**.** To reduce the content of DTT,

GPMVs were dialyzed in GPMV buffer prior to use in experiments. GPMVs were also created

using *N*-ethylmaleimide as the blebbing reagent, with comparable results. The phospholipid

content of final material was measured by total phosphate assay.

**Phosphate assay.** Lipid concentrations for GPMV preparations were estimated by measuring total phosphate, assuming that all measured phosphate is from phospholipids, and that all lipids are phospholipids. This is a practical assumption designed to ensure reproducibility.

**Enzymatic cleavage of carbohydrates.** For cleavage of carbohydrates from GPMVs, endoglycosidases were added to the GPMVs in GPMV buffer at concentrations recommended by the manufacturers (PNGaseF - 5,000 units/ml; EndoH - 2500 units; heparinaseI/II - 2500 units) and incubated at 37°C for 6 h. PNGaseF was tagged with a chitin binding domain (Remove-IT PNGaseF, New England Biolabs, MA, USA). For the images shown in this manuscript, the enzyme was removed by incubation of GPMVs with 50 μl of chitin binding magnetic beads. However, control experiments were conducted without removal of PNGaseF and found to be comparable. Cleavage of N-linked glycans by PNGaseF was confirmed by comparing images of GPMVs +/- PNGaseF treatment after incubation with 50 nM concanavalin A (conA), a lectin which binds to binding to α-D-mannose and α-D-glucose moieties, or wheat germ agglutinin, a lectin which binds to N-acetyl-D-glucosamine and sialic acid. A significant decrease in the amount of both proteins is observed in these images (Figure supplement 1).

For cleavage of carbohydrates from cells, cells were first plated for 42 h. After 42 h, media was removed from cells and replaced with FBS-free media complimented with the endoglycosidase (PNGaseF - 5,000 units/ml; EndoH - 2500 units; heparinaseI/II - 2500 units). The cells were incubated at 37°C under a humidified atmosphere of 5% $CO_2$ for an additional 6 h. The media was then removed and replaced with cell growth media prior to the addition of αS. Cleavage of N-linked glycans from cells was confirmed by comparing images +/- PNGAseF treatment after incubation with 50 nM conA-AL488, showing a significant reduction in the amount of conA bound (Figure supplement 4A).

To ensure that PNGaseF is removed from GPMVs or cells after incubation (and therefore does not remain associated with either, blocking potential αS binding sites), we compared the amount of PNGaseF added to either GPMVs or cells with that removed after incubation. PNGaseF (5,000 units/ml) containing a chitin domain was added to chambers containing either GPMVs or cells and incubated at 37°C for 6 h, as for the experiments described above. After incubation, the buffer or media containing PNGaseF was removed incubated with chitin magnetic beads to isolate and concentrate the enzyme. Blank chambers containing only PNGaseF in buffer or media were

subjected to the same treatment. 20 uL of each sample was run on a 4-12% polyacrylamide gels and stained with Coomassie blue. Gels were imaged using a Typhoon FLA7000 gel imager (GE Life Sciences) using Coomassie stain mode. The gels indicate that essentially all of the enzyme is removed (Figure supplement 8).

**Quantification of carbohydrates.** Concentrations of carbohydrates isolated from GPMVs or cells were quantified by using the Total Carbohydrate Quantification Assay Kit (Abcam, MA, USA) following the manufacturer's instructions. Briefly, the carbohydrates are first hydrolyzed to monomer sugar units and then converted to furfural or hydrofurfural. These compounds are converted to chromogens, which can be detected by absorbance at 490 nm. Glucose was used to generate a standard curve for calculated of the total carbohydrate concentration of the samples.

**αS-captured carbohydrate pulldown assay.** Carbohydrates cleaved and isolated from GPMVs (50 µg) were incubated with 100 µM αS in 100 µL 10 mM HEPES, 150 mM NaCl, 2 mM $CaCl_2$, pH 7.4 for 1 h at room temperature. Binding reaction mixes were transferred to Amicon Ultra (3000 Da MWCO) centrifugal concentration devices that had been washed with 500 µL deionized water. The concentrators were centrifuged for 5 min at 4200 rpm and the filtrates collected. The flow-through contains carbohydrates that did not bind αS and thus were not retained in the chamber of the concentrator. The amount of carbohydrate in the flow-through was quantified by the total carbohydrate assay as described above. The fraction of carbohydrate bound and retained by αS ($C_{captured}$) was calculated relative to the starting concentration (50 µg) of the carbohydrate mixture, using;

$$C_{captured} = C_{total} - C_{flow-through} \qquad \text{(eq. 1)}$$

Where $C_{total}$ corresponds to the absorption of the starting stock concentration of carbohydrates used and $C_{flow-through}$ is the absorption of the flow-through.

Samples containing αS with no glycans or conA with 50 µg glycans were used as negative and positive controls, respectively. As expected, no signal at 490 nm was detected for the αS-only sample; the conA results are shown in Figure supplement 2. The absorbance of flow-through at 280 nm was measured, confirming that all protein was retained in the concentrator and could not be detected in the flow-through.

**Cell imaging and analysis.** All cell imaging was carried out by confocal fluorescence microscopy using an Olympus FV3000 scanning system configured on a IX83 inverted microscope platform with a 60x Plan-Apo/1.1-NA water-immersion objective with DIC capability (Olympus, Tokyo, Japan). For all experiments the gain setting for the blue channel was kept constant from sample to sample (excitation 488 nm, emission BP 500-540 nm). For detection of αS-AL594, the green channel was used (excitation 561 nm, emission BP 570-620 nm). Images were obtained in 8-well ibidi chambers (μ-Slide, 8-well glass bottom, ibidi GmbH, Germany) coated with Poly-D-lysine and were seeded with 20000-25000 cells/well. Cells were cultured for 48 h after passage before beginning experiments. For cellular uptake experiments, 200 nM αS-AL488 was incubated with cells for 0 to 24 h before acquiring images. For experiments using deglycosylated cells, cells were pretreated with the tested endoglycosidase for 6 h as described above prior to addition of protein. For colocalization with lysosomes, cells were treated with 75 nM Lysotracker Deep Red (Life Technologies) for 1 h prior to imaging. For all experiments, the gain setting for each channel was kept constant from sample to sample.

Image acquisition and processing were performed with the software accompanying the FV3000 microscope and Image J software (Schneider et al., 2012). For SH-SY5Y cells, internalized αS was quantified either by analysis of the punctate structures in the cells or by the total cellular fluorescence; for primary neurons, internalized αS was quantified by total cellular fluorescence. For total cellular fluorescence, the integrated fluorescence intensity of the cells is calculated and reported. Cellular puncta were analyzed using the Image J particle analysis plug-in. This algorithm detects puncta through a user-defined threshold and counts the number of puncta that meet or exceed the threshold. The threshold was initially defined by manual identification and averaging of a subset of puncta. Colocalization with lysosomes was computed by obtaining a Pearson coeffieient using the ImageJ plugin for co-localization (Coloc_2).

**Endocytosis inhibition.** For inhibition of exocytosis experiments, monomer αS-AL488 (200 nM) or PFFs (200 nM in monomer units, 20:1 αS:αS-AL488) was initially incubated with cells for 30 minutes at 4°C. Control cells were moved to 37°C while the endocytosis inhibited cells were incubated at 4°C for another 4 h before acquiring images. For all experiments, the gain setting for each channel was kept constant from sample to sample. Image acquisition and processing were

achieved using the software accompanying the FV3000 microscope and Image J software
(Schneider et al., 2012).

**GPMV imaging and analysis.** All GPMV images were carried using a PicoQuant MicroTime 200
time-resolved fluorescence system based on an inverted Olympus IX73 microscope (Olympus,
Tokyo, Japan) with a 60x Plan-Apo/1.4-NA water-immersion objective using a 482 nm excitation
laser and a frame size of $512 \times 512$ pixels. Images acquired with this instrument were in lifetime
mode but were integrated to obtain intensity based images comparable to typical confocal images.
This instrument has the advantage of very sensitive avalanche photodiode detectors (SPADs) that
are capable at detecting nM concentrations of protein. Fluorescence intensities were analyzed by
via the lifetime mode (both intensity and FRET images) using SymPhoTime 64 (PicoQuant). The
intensity of images was then adjusted on ImageJ analysis program (Schneider et al., 2012).

For FLIM-FRET experiments, measurements were made of donor-only and donor-acceptor
labeled proteins. SPAD signals were processed with the TimeHarp 300 photon counting board and
analyzed with the SymPhoTime 64 software (PicoQuant) taking into account the instrument
response function to allow consideration of short lifetime components with a high accuracy. FLIM
images were acquired for 180 s, with a pixel integration time of 40 μs per pixel and an average
photon count rate of 10000–30000 counts per second. Regions of interest of the GPMV membrane
were selected from FLIM images and fluorescent lifetimes were obtained from TCSPC decay
curves fitted by an exponential equation using the SymPhoTime 64 software. Fitting of the
fluorescence images was then performed pixel wise with a single exponential model on all pixels
above an intensity threshold of 200 photons. By characterizing donor lifetime in the absence and
presence of acceptor, FRET efficiency ($ET_{eff}$) can be calculated from:

439
$$ET_{eff} = 1 - \frac{\tau_{DA}}{\tau_D} \qquad\qquad (eq.\,2)$$

where $\tau_{DA}$ and $\tau_D$ and are the donor excited state lifetime in the presence and absence of acceptor.
Six FLIM images were recorded for each of three biological repeats per condition. The histograms
shown in Figure 2 represents $ET_{eff}$ values for selected pixels from equatorial sections of the
GPMVs as indicated on the images to their left. The histograms were fit with a Gaussian function
to extract the mean $ET_{eff.}$

Images were obtained in 8-well NUNC chambers (Thermo Scientific, Rochester, NY,

USA) coated with Poly-D-lysine, containing 250 µl of GMPV at 5 µM phospholipid concentration

and 80 nM of αS labeled with Alexa 488. For all experiments using these chambers, the chambers

were passivated by polylysine-conjugated PEG treatment to prevent any nonspecific absorption to

the chamber surfaces (Middleton and Rhoades, 2010).

**Fluorescence correlation spectroscopy (FCS).** Fluorescence correlation spectroscopy (FCS)

measurements were carried out on a lab-built instrument, as described previously (Trexler and

Rhoades, 2009). A 488-nm-wavelength laser was adjusted to 5 µW prior to entering the

microscope. Fluorescence emission was collected through the objective and separated from laser

excitation using a Z488RDC Long-Pass Dichroic and an HQ600/200M Band-Pass Filter

(Chroma). The fluorescence emission was focused into the aperture of a 50-µm-diameter optical

aperture fiber (OzOptics) directly coupled to an avalanche photodiode. A digital correlator

(Flex03LQ-12; Correlator.com) was used to generate the autocorrelation curves. For each

experiment, 30 autocorrelation curves of 10 s each were acquired and averaged together to obtain

statistical variations. These average autocorrelation curves were then fit to a function for single

fluorescent particles undergoing Brownian motion in a 3D Gaussian volume weighted by the

inverse square of the SD:

462
$$G(\tau) = \frac{1}{N} \times \frac{1}{1+\frac{\tau}{\tau_D}} \times \sqrt{\frac{1}{1+\frac{s^2\tau}{\tau_D}}} \qquad\qquad (eq.\,3)$$

where G(τ) is the autocorrelation function for translational diffusion as a function of time τ, N is

the average number of fluorescent molecules in the laser focal volume, and $\tau_D$ is the translational

diffusion time of the particles. The structure factor, s, is the ratio of the radial to axial dimensions

of the focal volume and was obtained from a calibration procedure using Alexa 488 hydrazide dye

solutions (s = 0.2) and fixed for all subsequent fitting. Several datasets of autocorrelation curves

obtained in the presence of carbohydrates were analyzed considering a single component system.

For carbohydrate binding, both the diffusion time, $\tau_D$, (Figures 3A, 3B and Figure supplement 2B,

2D, 5C and 9) and the number of molecules, N, (Figures 1C, 1E, 4B and Figure supplement 2C,

3E, 5B, 5D and 9) were analyzed.

**GPMV binding and cellular uptake by FCS.** FCS was used to monitor αS both binding to

GPMVs and uptake by cells. FCS measurements were made on the instrument described above.

For GPMV binding, experiments were performed in 8-well NUNC chambers (Thermo Scientific,

Rochester, NY, USA) containing 250 µl of GMPV at 5 µM phospholipid concentration. The laser

focal volume was located to a height of 100 µm from the bottom surface of the wells, above all

GMPVs. αS labeled with Alexa 488 (80 nM) was added to wells with GPMVs and the

autocorrelation measurements were taken immediately and after 60 minutes. Each curve was

integrated for 30 s, repeated 10 times and data was analyzed as described above. Binding was

quantified by measuring the change in the number of molecules, N, of fluorescently labeled αS

present in the solution surrounding the GMPVs relative to the starting concentration. There was

no change in the diffusion time of the αS that remained in the cell media over this time course,

evidence that protein is stable and the fluorophore intact (Figure supplement 9).

For cell uptake experiments, cells were plated in phenol red-free medium in 8-well ibidi

chambers (µ-Slide, 8-well glass bottom, ibidi GmbH, Germany) coated with Poly-D-lysine wells

and cultured for 48 h prior to measurements. Control wells contained the same volume of media

without cells; these control for non-specific adsorption of protein to the well surfaces. αS-AL488

(200 nM) was added to the wells at the start of the experiment, and the autocorrelation curves were

taken in the medium well above the cells, at a height of 100 µm from the bottom surface of the

wells. The curves were collected at regular intervals, over a period ranging from 2 to 24 h, and

each autocorrelation curve integrated for 30 s, repeated 10 times. Data were analyzed as described

above. Cellular uptake was assessed by measuring the change in the number of molecules of

fluorescently labeled αS present in the media surrounding the cells relative to the starting

concentration. In between measurements, the chambers were returned to the incubators to maintain

the temperature in the wells at 37°C. As a negative control for both the GPMV binding and cell

uptake studies, GPMVs or cells were incubated with 80 nM eGFP. There was no evidence of eGFP

binding to the GPMVs nor of internalization by SH-SY5Y cells (Figure supplement 10).

**Cellular uptake by PAGE**. As an orthogonal approach to the FCS and imaging approached

described above, uptake of both monomer and PFF αS were quantified by PAGE. SH-SY5Y cells

were incubated with 200 nM αS, as described above for FCS uptake experiments. After incubation

for the desired time (1, 3, 5, 8, 12, or 24 h) the media was removed from cells and stored at 4°C.

After all samples were collected, 20 µl of each sample were run on a 4-12% polyacrylamide gel.

To quantify the amount of protein internalized, an identical set of experiments was carried

out, with modifications as described in the following. For each time point, transferrin-AL488 (100

nM) was added 30 min prior to the due time of the time point to serve as a loading control. At the

desired time points, the cells were detached from the wells by 0.05% Trypsin-EDTA (Life

Technologies), pelleted and lysed in 250 μl RIPA lysis buffer (Thermo Fisher). Cell lysates (20 μl

of stock) were run on PAGE gels. Cell lysates (20 μl) were run on 4-12% polyacrylamide gels.

Gels of both extracellular and internalized αS were imaged using a Typhoon FLA7000 gel

imager (GE Life Sciences) using fluorescent imaging mode to detect αS-AL488 or transferrin-

AL488. Image J was used to quantify the bands.

**Statistical analysis.** Data are expressed as the mean ± SD, and were examined by a one-way

analysis of variance (n = 3). More than three experiments were performed and similar results were

obtained. P values less than 0.05 were considered to be significant.

**Trypan Blue Quenching.** Trypan blue solution 0.4% (Thermo Fisher) and fresh neurobasal

without phenol red, B27, or antibiotic supplementation were equilibrated at 37°C. A 10X dilution

of trypan blue was prepared freshly in the warmed neurobasal media. The trypan blue solution was

then added to cells dropwise and incubated at 37°C for an hour prior to imaging. Trypan Blue

quenching was performed for all imaging experiments performed using PFFs and when monomer

protein was introduced in experiments utilizing primary neurons. Trypan Blue quenching was used

eliminate signal from extracellular material allowing the fluorescence quantification of

intracellular fluorescence signal.

**Nuclear Magnetic Resonance.** $^1$H-$^{15}$N HSQC NMR titrations were carried out at 25°C using

Varian 600 MHz or Agilent 800 MHz spectrometers equipped with room temperature probes. A

uniformly labeled $^{15}$N-αS solution was added to either N-linked glycans obtained by PNGaseF

cleavage from SH-SY5Y cells or commercially available mono- or trisaccharides at the

concentrations indicated in Figures 4 and S2E in GPMV buffer (10 mM HEPES, 150 mM NaCl,

2 mM CaCl2, pH 7.4, 10% $D_2O$; final concentration of $^{15}$N-αS was 350 μM). HSQC spectra were

collected with VnmrJ software using built-in pulse sequence including WATERGATE solvent

suppression and analyzed with Mnova software suite (Mestrelab). Standard parameters for zero-

filling, apodization and baseline correction were applied. $^1$H chemical shifts were referenced using

water resonance and $^{15}N$ chemical shifts were referenced indirectly based on gyromagnetic ratios

of respective nuclei. Previously assigned αS backbone resonances were used(Kang et al., 2011).

Binding to the PNGaseF derived glycans results in non-uniform peak intensity increases

throughout the sequence of αS. For analysis, 10 peaks showing large increases and 10 peaks

showing small or no increases were selected to roughly cover the entire sequence. The same

residues were used for each analyzed dataset. For each set of peaks, the relative magnitude of

increase (as compared to αS in solution without glycans) expressed as a percentage was calculated

and averaged. A difference between two numbers is reported in Figure 3C of the manuscript. The

same calculation was carried out for the mono- and tri-saccharides, which do not bind to αS, shown

in Figure supplement 2E.

**Acknowledgments**

This research was supported by NIH R01 NS079955 and the Michael J. Fox Foundation (to E.R.).

We thank T. Baumgart for use of his confocal microscope and V. M.-Y. Lee for use of her

sonicator. We also thank M. Maronski of the Neuron Cell Core in the The Mahoney Institute for

Neurosciences in the Perlman School of Medicine at the University of Pennsylvania for

hippocampal neuron preparation and R. Jin for providing the vesicle schematic adapted for Figure

2. All data are available from authors by request.

**Competing interests**

The authors declare that no competing interests exist.

**References**

ANDERSON, J. P., WALKER, D. E., GOLDSTEIN, J. M., DE LAAT, R., BANDUCCI, K., CACCAVELLO, R. J., BARBOUR, R., HUANG, J., KLING, K., LEE, M., DIEP, L., KEIM, P. S., SHEN, X., CHATAWAY, T., SCHLOSSMACHER, M. G., SEUBERT, P., SCHENK, D., SINHA, S., GAI, W. P. & CHILCOTE, T. J. 2006. Phosphorylation of Ser-129 is the dominant pathological modification of alpha-synuclein in familial and sporadic Lewy body disease. *J Biol Chem,* 281**,** 29739-52, 10.1074/jbc.M600933200.

BAUER, B., DAVIDSON, M. & ORWAR, O. 2009. Proteomic analysis of plasma membrane vesicles. *Angew Chem Int Ed Engl,* 48**,** 1656-9, 10.1002/anie.200803898.

COOPER, A. A., GITLER, A. D., CASHIKAR, A., HAYNES, C. M., HILL, K. J., BHULLAR, B., LIU, K., XU, K., STRATHEARN, K. E., LIU, F., CAO, S., CALDWELL, K. A., CALDWELL, G. A., MARSISCHKY, G., KOLODNER, R. D., LABAER, J., ROCHET, J. C., BONINI, N. M. & LINDQUIST, S. 2006. Alpha-synuclein blocks ER-Golgi traffic and Rab1 rescues neuron loss in Parkinson's models. *Science,* 313**,** 324-8, 10.1126/science.1129462.

DIAO, J., BURRE, J., VIVONA, S., CIPRIANO, D. J., SHARMA, M., KYOUNG, M., SUDHOF, T. C. & BRUNGER, A. T. 2013. Native alpha-synuclein induces clustering of synaptic-vesicle mimics via binding to phospholipids and synaptobrevin-2/VAMP2. *Elife,* 2**,** e00592, 10.7554/eLife.00592.

DIKIY, I. & ELIEZER, D. 2014. N-terminal acetylation stabilizes N-terminal helicity in lipid- and micelle-bound alpha-synuclein and increases its affinity for physiological membranes. *J Biol Chem,* 289**,** 3652-65, 10.1074/jbc.M113.512459.

DRAZIC, A., MYKLEBUST, L. M., REE, R. & ARNESEN, T. 2016. The world of protein acetylation. *Biochim Biophys Acta,* 1864**,** 1372-401, 10.1016/j.bbapap.2016.06.007.

EMMANOUILIDOU, E., MELACHROINOU, K., ROUMELIOTIS, T., GARBIS, S. D., NTZOUNI, M., MARGARITIS, L. H., STEFANIS, L. & VEKRELLIS, K. 2010. Cell-produced alpha-synuclein is secreted in a calcium-dependent manner by exosomes and impacts neuronal survival. *J Neurosci,* 30**,** 6838-51, 10.1523/JNEUROSCI.5699-09.2010.

FORTIN, D. L., TROYER, M. D., NAKAMURA, K., KUBO, S., ANTHONY, M. D. & EDWARDS, R. H. 2004. Lipid rafts mediate the synaptic localization of alpha-synuclein. *J Neurosci,* 24**,** 6715-23, 10.1523/JNEUROSCI.1594-04.2004.

GEORGE, J. M., JIN, H., WOODS, W. S. & CLAYTON, D. F. 1995. Characterization of a novel protein regulated during the critical period for song learning in the zebra finch. *Neuron,* 15**,** 361-72.

GOEDERT, M. 2001. Alpha-synuclein and neurodegenerative diseases. *Nat Rev Neurosci,* 2**,** 492-501, 10.1038/35081564.

GUO, J. L. & LEE, V. M. 2014. Cell-to-cell transmission of pathogenic proteins in neurodegenerative diseases. *Nat Med,* 20**,** 130-8, 10.1038/nm.3457.

HITCHCOCK-DEGREGORI, S. E. & HEALD, R. W. 1987. Altered actin and troponin binding of amino-terminal variants of chicken striated muscle alpha-tropomyosin expressed in Escherichia coli. *J Biol Chem,* 262**,** 9730-5.

HOLMES, B. B., DEVOS, S. L., KFOURY, N., LI, M., JACKS, R., YANAMANDRA, K., OUIDJA, M. O., BRODSKY, F. M., MARASA, J., BAGCHI, D. P., KOTZBAUER, P. T., MILLER, T. M., PAPY-GARCIA, D. & DIAMOND, M. I. 2013. Heparan sulfate proteoglycans mediate

internalization and propagation of specific proteopathic seeds. *Proc Natl Acad Sci U S A,*
110**,** E3138-47, 10.1073/pnas.1301440110.
IHSE, E., YAMAKADO, H., VAN WIJK, X. M., LAWRENCE, R., ESKO, J. D. & MASLIAH, E. 2017. Cellular
internalization of alpha-synuclein aggregates by cell surface heparan sulfate depends on
aggregate conformation and cell type. *Sci Rep,* 7**,** 9008, 10.1038/s41598-017-08720-5.
KANG, L., WU, K. P., VENDRUSCOLO, M. & BAUM, J. 2011. The A53T mutation is key in defining
the differences in the aggregation kinetics of human and mouse alpha-synuclein. *J Am*
*Chem Soc,* 133**,** 13465-70, 10.1021/ja203979j.
MAO, X., OU, M. T., KARUPPAGOUNDER, S. S., KAM, T. I., YIN, X., XIONG, Y., GE, P., UMANAH, G.
E., BRAHMACHARI, S., SHIN, J. H., KANG, H. C., ZHANG, J., XU, J., CHEN, R., PARK, H.,
ANDRABI, S. A., KANG, S. U., GONCALVES, R. A., LIANG, Y., ZHANG, S., QI, C., LAM, S.,
KEILER, J. A., TYSON, J., KIM, D., PANICKER, N., YUN, S. P., WORKMAN, C. J., VIGNALI, D.
A., DAWSON, V. L., KO, H. S. & DAWSON, T. M. 2016. Pathological alpha-synuclein
transmission initiated by binding lymphocyte-activation gene 3. *Science,* 353,
10.1126/science.aah3374.
MIDDLETON, E. R. & RHOADES, E. 2010. Effects of curvature and composition on alpha-synuclein
binding to lipid vesicles. *Biophys J,* 99**,** 2279-88, 10.1016/j.bpj.2010.07.056.
MOREMEN, K. W., TIEMEYER, M. & NAIRN, A. V. 2012. Vertebrate protein glycosylation: diversity,
synthesis and function. *Nat Rev Mol Cell Biol,* 13**,** 448-62, 10.1038/nrm3383.
ROBOTTA, M., BRAUN, P., VAN ROOIJEN, B., SUBRAMANIAM, V., HUBER, M. & DRESCHER, M.
2011. Direct evidence of coexisting horseshoe and extended helix conformations of
membrane-bound alpha-synuclein. *Chemphyschem,* 12**,** 267-9, 10.1002/cphc.201000815.
ROZYCKI, B. & CIEPLAK, M. 2016. Stiffness of the C-terminal disordered linker affects the
geometry of the active site in endoglucanase Cel8A. *Mol Biosyst,* 12**,** 3589-3599,
10.1039/c6mb00606j.
SCHNEIDER, C. A., RASBAND, W. S. & ELICEIRI, K. W. 2012. NIH Image to ImageJ: 25 years of image
analysis. *Nat Methods,* 9**,** 671-5.
SEZGIN, E., KAISER, H. J., BAUMGART, T., SCHWILLE, P., SIMONS, K. & LEVENTAL, I. 2012.
Elucidating membrane structure and protein behavior using giant plasma membrane
vesicles. *Nat Protoc,* 7**,** 1042-51, 10.1038/nprot.2012.059.
STARHEIM, K. K., GEVAERT, K. & ARNESEN, T. 2012. Protein N-terminal acetyltransferases: when
the start matters. *Trends Biochem Sci,* 37**,** 152-61, 10.1016/j.tibs.2012.02.003.
THEILLET, F. X., BINOLFI, A., BEKEI, B., MARTORANA, A., ROSE, H. M., STUIVER, M., VERZINI, S.,
LORENZ, D., VAN ROSSUM, M., GOLDFARB, D. & SELENKO, P. 2016. Structural disorder of
monomeric alpha-synuclein persists in mammalian cells. *Nature,* 530**,** 45-50,
10.1038/nature16531.
TREXLER, A. J. & RHOADES, E. 2009. Alpha-synuclein binds large unilamellar vesicles as an
extended helix. *Biochemistry,* 48**,** 2304-6, 10.1021/bi900114z.
TREXLER, A. J. & RHOADES, E. 2012. N-Terminal acetylation is critical for forming alpha-helical
oligomer of alpha-synuclein. *Protein Sci,* 21**,** 601-5, 10.1002/pro.2056.
TUTTLE, M. D., COMELLAS, G., NIEUWKOOP, A. J., COVELL, D. J., BERTHOLD, D. A., KLOEPPER, K.
D., COURTNEY, J. M., KIM, J. K., BARCLAY, A. M., KENDALL, A., WAN, W., STUBBS, G.,
SCHWIETERS, C. D., LEE, V. M., GEORGE, J. M. & RIENSTRA, C. M. 2016. Solid-state NMR

structure of a pathogenic fibril of full-length human alpha-synuclein. *Nat Struct Mol Biol,*
23**,** 409-15, 10.1038/nsmb.3194.
VOLPICELLI-DALEY, L. A., LUK, K. C. & LEE, V. M. 2014. Addition of exogenous alpha-synuclein
preformed fibrils to primary neuronal cultures to seed recruitment of endogenous alpha-
synuclein to Lewy body and Lewy neurite-like aggregates. *Nat Protoc,* 9**,** 2135-46,
10.1038/nprot.2014.143.
YAP, T. L., GRUSCHUS, J. M., VELAYATI, A., WESTBROEK, W., GOLDIN, E., MOAVEN, N., SIDRANSKY,
E. & LEE, J. C. 2011. Alpha-synuclein interacts with Glucocerebrosidase providing a
molecular link between Parkinson and Gaucher diseases. *J Biol Chem,* 286**,** 28080-8,
10.1074/jbc.M111.237859.
650